

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/107875/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Dekker, Anne R.J., Verheij, Theo J.M., Broekhuizen, Berna D.L., Butler, Christopher C, Cals, Jochen W.L., Francis, Nicholas , Little, Paul, Sanders, Elisabeth A.M., Yardley, Lucy, Zuithoff, Nicolaas P.A. and Van Der Velden, Alike W. 2018. Effectiveness of GP online training and an information booklet for parents on antibiotic prescribing for children with RTI in primary care: a cluster randomised controlled trial. *Journal of Antimicrobial Chemotherapy* 73 (5) , pp. 1416-1422. 10.1093/jac/dkx542

Publishers page: <https://doi.org/10.1093/jac/dkx542>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1 **Effectiveness of GP online training and an information booklet for parents on antibiotic**
2 **prescribing for children with RTI in primary care: a cluster randomised controlled trial**

3 Anne R.J. DEKKER^{1*}, Theo J.M. VERHEIJ¹, Berna D.L. BROEKHUIZEN¹, Christopher C.
4 BUTLER², Jochen W.L. CALS³, Nick A. FRANCIS⁴, Paul LITTLE⁵, Elisabeth A.M. SANDERS⁶,
5 Lucy YARDLEY⁷, Nicolaas P. A. ZUITHOFF¹, Alike W. VAN DER VELDEN¹

6 Author Affiliations:

7 ¹Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht,
8 Utrecht, The Netherlands; ²Nuffield Department of Primary Care Health Sciences, University of
9 Oxford, Oxford, UK; ³Department of Family Medicine, CAPHRI School for Public Health and
10 Primary Care, Maastricht University, Maastricht, the Netherlands; ⁴Department of Primary Care
11 and Public Health, School of Medicine, Cardiff University, Cardiff, UK; ⁵Primary Care Medical
12 Group, University of Southampton Medical School, Southampton, UK; ⁶Department of Paediatric
13 Immunology, Wilhelmina Children's Hospital, University Medical Center, Utrecht, the
14 Netherlands; ⁷Academic Unit of Psychology, Faculty of Social and Human Sciences, University
15 of Southampton, Southampton, UK.

16

17 *Corresponding author. Tel: +31-88-756-8364, Fax: +31-88-756-8099, E-mail: a.r.j.dekker-
18 8@umcutrecht.nl

19 **Running title:** Improving antibiotic use for children with RTI

20

21 **ABSTRACT**

22 **Objectives:** Antibiotics are too often prescribed in childhood respiratory tract infection (RTI),
23 despite limited effectiveness, potential side-effects, and bacterial resistance. We aimed to
24 reduce antibiotic prescribing for children with RTI by online training for general practitioners
25 (GP) and information for parents.

26 **Methods:** A pragmatic cluster randomised, controlled trial in primary care. The intervention
27 consisted of an online training for GPs and an information booklet for parents. The primary
28 outcome was the antibiotic prescription rate for children presenting with RTI symptoms, as
29 registered by GPs. Secondary outcomes were number of consultations within the same
30 disease episode, consultations for new episodes, hospital referrals and pharmacy dispensed
31 antibiotic courses for children.

32 **Results:** After randomisation, GPs of in total 32 general practices registered 1009
33 consultations. An antibiotic was prescribed in 21% of consultations in the intervention group,
34 compared to 33% in the usual care group, controlled for baseline prescribing (RR 0.65, 95% CI
35 0.46-0.91). The probability of reconsulting during the same RTI episode did not differ
36 significantly between the intervention and control group, nor did the numbers of consultations for
37 new episodes and hospital referrals. In the intervention group antibiotic dispensing was reduced
38 with 32 courses per 1000 children/year, compared to the control group, and adjusted for
39 baseline prescribing (RR 0.78, 95% CI 0.66-0.92). The numbers and proportion of second
40 choice antibiotics did not differ significantly.

41 **Conclusion:** A concise, feasible, online GP training, with an information booklet for parents
42 showed a relevant reduction in antibiotic prescribing for children with RTI.

43 This trial was registered at the Dutch Trial Register (NTR), registration number: NTR4240

44 INTRODUCTION

45 Respiratory tract infections (RTI), including ear-infections, are the most common indication for
46 consulting a general practitioner (GP) during childhood and for prescribing antibiotics.^{1,2} Most
47 RTIs are viral and self-limiting, and many high-income countries have guidelines aiming to
48 restrict the use of antibiotics.³⁻⁵ However, even in a low-prescribing country like the Netherlands,
49 one third of antibiotic prescriptions for children are not congruent with guideline
50 recommendations.⁶ The main drivers of over-prescription are GPs' interpretation of patient or
51 parent expectations, time pressure, diagnostic and prognostic uncertainty and unfamiliarity with
52 recent guidelines.⁷⁻⁹ General practice has a major contribution and responsibility towards
53 antibiotic stewardship, since primary care is a driver of antibiotic resistance.¹⁰⁻¹² Efforts to
54 reduce antibiotic prescribing in primary care have been ongoing for decades, most often
55 focusing on antibiotic use in adults, and consisting of a wide range of strategies.¹³⁻¹⁶ A different
56 approach might be needed for childhood RTI, because of child-specific indications and risk
57 factors, and communication with parents instead of patients themselves.¹⁷ Multifaceted
58 approaches have been shown to be most effective, however, broad implementation of these
59 interventions is rare because of time and costs.^{13,18-20} Online educational programs could be a
60 feasible and cost-effective intervention that could be broadly implemented, updated easily, and
61 ensure a more enduring antibiotic stewardship. Little *et al.* showed that such an intervention
62 was effective in improving antibiotic management of adults with lower RTI.¹⁶ In children, only
63 online instruction on the use of information material was studied in the UK, which was
64 effective.²¹ In our study we aimed to assess the effects of an online training for GPs and an
65 information booklet for parents on antibiotic prescribing for children with RTI in general practice.

66

67 **METHODS**

68 **Trial design**

69 The RAAK (Rational Antibiotic use Kids) study was a pragmatic, cluster randomised, two-arms,
70 controlled trial with measurements before and after the intervention, to allow for adjustment for
71 baseline antibiotic prescribing (baseline audit). GPs within a general practice influence each
72 other and patients within a practice are often managed by different GPs, therefore, the general
73 practice was the unit of randomisation and the unit of analysis to minimize contamination and
74 dilution of the intervention effect. GPs in the control group practised care as usual. We followed
75 the Consolidated Standards of Reporting Trials guidelines, extended for cluster randomised
76 trials.²²

77 **Ethics approval**

78 This trial was exempted by the Ethics Committee of the University Medical Center Utrecht from
79 obtaining parents' or patients' consent (reference number METC 13-237/C). The trial assigned
80 GPs with the aim to improve their prescribing behaviour according to the national practice
81 guidelines. Children were not the subject of the intervention and were treated according to the
82 guidelines.

83 **General practices and participants**

84 For the baseline audit, GPs were asked to register 40 consecutive consultations of children
85 younger than 18 years with signs and symptoms of RTI (nose, ear, throat and/or lower RTI
86 symptoms), presenting at their general practice during the winter season 2013-2014. GPs
87 registered the following anonymous information on consultation report forms: age, duration of
88 symptoms, fever, most prominent symptoms, findings of physical examination, overall illness
89 severity (1= minimally ill, 5=severely ill), the International Classification of Primary Care code for

90 diagnosis, and whether an antibiotic was prescribed, including which one. General practices
91 were excluded if GPs registered less than ten patients in total per general practice, since low
92 numbers could result in poor estimations of the baseline antibiotic prescription rates. After
93 randomisation and implementation of the intervention, this registration of consultations was
94 repeated in the follow-up audit, during the winter season 2014-2015. In addition, parents were
95 invited to fill in a diary for up to two weeks following the index consultation, and give permission
96 to review the child's medical records after six months to collect secondary outcomes.

97 **Intervention**

98 The intervention consisted of online training for GPs and a written information booklet for
99 parents. These were adapted from an intervention for adults that was: a) theory-based: the
100 educational content was designed to promote positive expectations and self-confidence in GPs
101 and patients to manage the infection without antibiotics, b) person-based: the content was
102 developed with extensive feedback from GPs and patients to ensure that it addressed their
103 concerns and was persuasive.²³⁻²⁵ The online training consisted of three parts. The first part was
104 a general background about the relevance of prudent antibiotic use and information about
105 antibiotic-related problems. We presented over-prescription by percentages of prescribed
106 antibiotics, not congruent with guideline recommendations, from a recent Dutch study, to make
107 GPs aware of their responsibility in prudent antibiotic use.⁶ The second part informed about the
108 child-specific parts of the four national RTI guidelines of the Dutch College of GPs⁵, including
109 assessment of disease severity, risk factors, signs and symptoms, when to prescribe antibiotics,
110 and the advised first and second choice antibiotic treatment. This part was summarized in a
111 printable document, which is available as supplementary data at JAC online. The third part
112 focused on training in enhanced communication skills, supported by videos of consultation
113 techniques. The communication skills training was based on the elicit-provide-elicite framework,
114 used in prior antibiotic interventions, adapted to communication with parents.^{7,15,26} In summary,

115 the GP first elicits what the parent's main worries and expectations are. Crucially, the GP
116 actively asks how the parent feels about and what he/she expects from antibiotics. Secondly,
117 the GP provides information relevant to the parents individual understanding and interest,
118 including findings from the medical history and physical examination of the child. Then, the GP
119 elicits the parents interpretation about what has been said and done, to reach mutual agreement
120 and concludes with concrete safety netting, explaining specific signs and symptoms when to
121 reconsult.

122 GPs were invited by email to commence the training. If the training was not started or
123 completed, a weekly reminder email was automatically sent with the request to complete the
124 online training.

125 The booklet contained the following information in text and pictograms: epidemiology of RTI,
126 their predominant viral cause, self-limiting prognosis, rationale to withhold antibiotics, and
127 antibiotic related problems, including bacterial resistance. Additionally, self-management
128 strategies for their child and signs and symptoms when to consult the GP were explained.

129 **Outcomes, sample size, and randomisation**

130 The primary outcome was the antibiotic prescription rate per general practice in the follow-up
131 audit, as documented on the consultation report forms filled in by the GPs.¹⁶ The following
132 secondary outcomes were assessed from the patients' medical records: number of
133 reconsultations during the same disease episode, number of consultations for new RTI
134 episodes and the number of hospital referrals during a follow-up of six months. Total and types
135 of dispensed antibiotic courses for all children under 18 years were collected via the Dutch
136 Foundation for Pharmaceutical Statistics.²⁷ Affiliated pharmacies of the participating general
137 practices (n=68) were asked for permission to collect all dispensed antibiotics that resulted from
138 prescribing by the participating GPs of that practice. Numbers of dispensed systemic antibiotics

139 (ATC-code J01) were collected via an online module for the complete years prior to and after
140 introducing the online training. Total numbers of antibiotics mainly used for RTIs were:
141 tetracyclines (J01AA), amoxicillin (J01CA), pheneticillin (J01CE), amoxicillin/clavulanate
142 (J01CR) and macrolides (J01FA). Amoxicillin (J01CA) and pheneticillin (J01CE) were
143 considered as first choice antibiotics, the others as second choice. The numbers of registered
144 children in the practice for the corresponding year were collected. The median duration of the
145 time being logged-in and the short online evaluation of the GP training were assessed.

146 We calculated that we would need a minimum of 157 consultations per arm, to be able to detect
147 an absolute difference of 15% in prescribing rate (42% and 27%), with 80% power and a 5%
148 significance level. To adjust for clustering of the effect within general practices, we assumed an
149 intra-cluster coefficient of 0.07 and a cluster size of 40, requiring a total of 1171 consultations in
150 both arms.²⁸ In order to achieve this we set out to ask 30 practices to register 40 consultations
151 each. Simple random allocation was performed by a computer generated list on general practice
152 level.

153 **Data analysis**

154 The primary analysis was according to the principle of intention-to-treat and assessed the
155 intervention effect on antibiotic prescribing to children as registered by the GPs in the follow-up
156 audit. We aggregated the data to the cluster level and used a generalized linear model for
157 Poisson distributed count outcomes, controlled for overdispersion.²⁹ We calculated Rate Ratios
158 (RR) with corresponding 95% Confidence Intervals (CI) and adjusted for baseline prescription
159 rates per general practice, as assessed in the year before the intervention. We chose not to
160 adjust for signs/symptoms, or diagnosis, because the interpretation, judgment and use of these
161 variables were part of the educational aspect of the online training.³⁰ The secondary outcomes
162 were also aggregated to the cluster level and analysed similarly as the primary outcome.

163 Pharmacy antibiotic dispensing data were retrieved per practice. The numbers of total
164 dispensed antibiotics were analysed using a generalized linear model and controlled for the
165 numbers of dispensed antibiotics in the year preceding the intervention, and the numbers of
166 children in the practice. Prescription of second choice antibiotics was analysed related to the
167 total number of children and to the total number of dispensed antibiotics and was controlled for
168 baseline prescribing. Analyses were done in SPSS version 21.

169 **RESULTS**

170 **Practice flow**

171 Before randomisation, 38 practices agreed to participate (Figure 1). Preceding the intervention,
172 three practices were excluded, as they did not register any consultation during the baseline
173 audit. Finally, 35 practices were randomised to the control or intervention arm. Three out of 35
174 randomised practices were excluded during the follow-up audit. They had not registered enough
175 consultations, because of sick leave of participating GPs. Therefore, pharmacy data of these
176 practices could neither be obtained reliably. One single-handed GP was excluded for the
177 pharmacy data, since his practice moved during the study period to another part of the city.
178 Practices of the intervention and control group were comparable with respect to their total list
179 size and numbers of listed children (Table 1).

180 **Registration of consultations**

181 During the baseline audit 1009 consultations of children with symptoms of RTI were registered
182 by 75 GPs from 35 general practices (Figure 1). The mean antibiotic prescription rate from this
183 baseline audit was 29.6% (35.7%, SD 4.8 in the control group versus 24.2%, SD 4.3 in the
184 intervention group). The follow-up audit included 1009 consultations in total, 532 from control
185 and 477 from intervention practices. Consultations were comparable between the intervention
186 and control group with respect to childrens' age, duration of illness before consultation, illness
187 severity and presentation with fever (Table 2). Numbers of registered symptoms appeared to be
188 higher in the intervention group as compared to the control group, especially for earache (37.1%
189 versus 29.3%).

190 **Intervention**

191 The training was completed by all 40 GPs of the intervention group. Their median time logged-in
192 was one hour and 18 minutes. Based on GPs' evaluation, the first and second part of the
193 training, with the general background and information of the four guidelines, were valued
194 highest, with a mean score of 4.5 (1=low value, 5=high value); the third part about
195 communication skills scored a mean of 4.2.

196 **Numbers analysed**

197 Analysis of the primary outcome was performed on 475 consultations in the 15 practices
198 allocated to the intervention, and 531 consultations in 17 practices allocated to usual care.
199 Three consultations lacked the primary outcome and were excluded from analyses. In 535
200 (53%) consultations of children, the parent gave permission to anonymously collect secondary
201 outcomes after six months from the child's medical record and was willing to fill in a diary. These
202 consultations showed no relevant differences compared to consultations in which parents were
203 not willing to participate in the study (data not shown). Secondary outcomes of 508 children
204 were available for analyses, 27 cases were lost to follow-up.

205 **Outcomes**

206 In 21.4% of consultations an antibiotic was prescribed in intervention practices, compared to
207 33.2% in the control group. The rate ratio after adjustment for baseline prescription was 0.65
208 (95% CI 0.46-0.91, Table 3). The intra-cluster coefficient was 0.09. The mean number of
209 reconsultations per 100 children within the same disease episode was lower in the intervention
210 group (42), as compared to the control group (64), but did not differ significantly (RR 0.66, Table
211 4). The probability of consultation for new RTI within six months did not differ significantly (RR
212 1.06), nor of hospital referrals (RR 0.66). General practices exposed to the intervention reduced
213 antibiotic dispensing with 32 courses per 1000 children per year, relative to the control group,
214 and based on the full year's pharmacy data (RR 0.78, 95% CI 0.66-0.92, Table 5). Adjusted for

215 the year preceding the intervention, the number of dispensed antibiotics was 114 per 1000
216 children in the intervention group and 146 per 1000 children in the control group. The number of
217 dispensed second choice antibiotics in the intervention group was lower (39.9/1000 children) as
218 compared to the control group (49.2/1000 children), however, this difference was not significant.
219 The percentage of second choice antibiotics neither differed between the control and
220 intervention group (34.1%, versus 34.4%).

221 **DISCUSSION**

222 Online training of GPs and information booklets for parents resulted in less antibiotic
223 prescriptions, measured by GPs' registrations of consultations, as well as by data of total yearly
224 antibiotic dispensing to children with RTIs. The intervention did not result in a significant
225 reduction in second choice antibiotics, reconsultations in the same disease episode,
226 consultations for new RTI episodes, or hospital referrals.

227 Outcomes of previous studies vary depending on setting, study population, and type of
228 intervention.^{13,18-20,31,32} Relatively intensive interventions targeting both parents and clinicians are
229 considered to be most effective, and decrease antibiotic prescribing rates by 6-21%.¹⁹ Focusing
230 on GP-parent communication, supported by written information, also showed to be
231 important.^{14,18,19,31-33} In our study, the prescription rates adjusted for baseline prescription
232 differed 11.8%. This effect was striking, particularly as our baseline prescription rates was
233 already low in comparison with other countries. Previous studies often used complex and time
234 consuming interventions, whereas our online training was feasible, concise and without
235 personal (academic) involvement and showed a long-term effect on antibiotic prescribing.
236 Online GP training to reduce antibiotic prescribing for children has not been used yet in primary
237 care, except for one study in the UK.²¹ This study primarily focused on consulting behaviour,
238 using an information booklet endorsed by the GP; the online training was about how to use the
239 booklet and did not include guideline education and background of antibiotic-related problems.²¹

240 **Strengths and Limitations**

241 This cluster randomised controlled trial showed a convincing effect on antibiotic prescribing
242 using GPs' registrations and pharmacy dispensing data during a full year after the intervention.
243 In the context of continuously improving RTI treatment in children, our study aimed to make a
244 simple, concise and feasible intervention, which was valued by GPs and parents.³⁴ The

245 pragmatic study design did not interfere with daily practice and did not require large time
246 investments or organizational adaptations. Our focus on the total childhood population with
247 broad eligibility criteria, and without selection of subgroups, or controlling for patient
248 characteristics, makes our results reliable and generalizable. By measuring both antibiotic
249 prescribing outcomes in the year preceding the intervention, we were able to control for baseline
250 prescribing, making our results more robust, since the number of clusters was not large.^{35,36} Our
251 study also has potential limitations. First, the pharmacy data could include GPs in the
252 intervention group who did not receive the online training, since some GPs who were not
253 involved in the trial, for example temporary locums or GPs in training, prescribed antibiotics on
254 behalf of participating GPs. This may have diluted the real, potentially higher, intervention effect.
255 This change of employees in the participating practices was increasing over time, and
256 prevented us from reliably measuring the intervention effect in the second year. Secondly, our
257 study was not powered to study whether severe complications could occur more frequently due
258 to reduced antibiotic prescriptions, nevertheless there was no evidence suggesting an adverse
259 effect of the intervention. Our intervention taught GPs according to the evidence-based
260 guidelines.⁵ We therefore expect no risk of inducing under-prescription. Another Dutch
261 intervention, aiming to reduce antibiotic prescribing showed that both over- and
262 underprescribing improved.²⁷ And, a substantial reduction in antibiotic prescriptions was shown
263 to be safe in a recent population-based study.³⁷ Finally, there is a non-significant difference in
264 reconsultation in the intervention and control group, with large within group variation. Many
265 parents of registered children were not invited to participate due to time constraints during the
266 consultation and only half of the invited parents were willing to keep a diary and gave
267 permission to assess the medical records of their child.

268 **Conclusion**

269 The intervention was effective in reducing antibiotic prescribing, and was feasible and
270 acceptable.³⁴ Given the minimal training time and the clear impact on antibiotic prescriptions it is
271 likely to be cost-effective. To implement this intervention at a national level some aspects could
272 be further developed, e.g. considering presenting the information booklet electronically,
273 stimulating informal learning activities including self-reflection, and potential linkage to a
274 structural antibiotic stewardship program.^{34,38}

275 This trial was registered at the Dutch Trial Register (NTR), registration number: NTR4240

276 **OTHERS**

277 **Acknowledgements**

278 We thank the GPs and children/parents for their participation in the RAAK trial, and the
279 pharmacists and Foundation for Pharmaceutical Statistics for sharing and facilitating retrieval of
280 the antibiotic dispensing data. Eveline Noteboom is thanked for practical assistance and Dr. F.
281 Grosfeld for expert advice.

282 **Registration**

283 This trial was registered at the the Dutch Trial Register (NTR), registration number: NTR4240.

284 **Funding**

285 This trial was funded by the Netherlands Organization for Health Research and Development
286 (ZonMw, grant number: 2052.00008). This work was conducted independently from the study
287 funder.

288 **Transparency declarations**

289 None to declare.

290 **Author contributions**

291 Alike van der Velden, Theo Verheij, Lidewij Broekhuizen, Christopher Butler, Jochen Cals, Nick
292 Francis, Paul Little and Lucy Yardley conceived and designed the study. Anne Dekker
293 organised the trial and collected all data. Anne Dekker, Alike van der Velden, Theo Verheij and
294 Peter Zuithoff analysed and interpreted the data. Anne Dekker, Alike van der Velden and Theo
295 Verheij wrote the first draft of the manuscript, and all coauthors critically revised the manuscript
296 for intellectual content. All authors approved the final version and agreed to serve as guarantors
297 of the work.

298 **REFERENCES**

- 299 1. Dekker AR, Verheij TJ, van der Velden AW. Antibiotic management of children with infectious
300 diseases in Dutch primary care. *Fam Pract* 2017; **34**: 169-174.
- 301 2. van der Linden MW, van Suijlekom-Smit LWA, Schellevis FG *et al*. Tweede Nationale Studie
302 naar ziekten en verrichtingen in de huisartspraktijk. Het kind in de huisartspraktijk. 2005,
303 Rotterdam/Culemborg: erasmus MC/Twin Design. ISBN: 90-74494-14 -5.
- 304 3. Clinical guideline: Respiratory tract infections (self-limiting): prescribing antibiotics. NICE
305 guideline development group. <https://www.nice.org.uk/guidance/cg69>.
- 306 4. Harris AM, Hicks LA, Qaseem A. High Value Care Task Force of the American College of
307 Physicians and for the Centers for Disease Control and Prevention. Appropriate antibiotic use
308 for acute respiratory tract infection in adults: Advice for high-value care from the american
309 college of physicians and the centers for disease control and prevention. *Ann Intern Med* 2016;
310 **164**: 425-434.
- 311 5. Clinical guidelines: acute cough, acute sore throat, acute otitis media and acute rhinosinusitis.
312 Nederlands Huisartsen Genootschap (Dutch College of GPs). [https://www.nhg.org/nhg-](https://www.nhg.org/nhg-standaarden)
313 [standaarden](https://www.nhg.org/nhg-standaarden)
- 314 6. Dekker AR, Verheij TJ, van der Velden AW. Inappropriate antibiotic prescription for
315 respiratory tract indications: most prominent in adult patients. *Fam Pract* 2015; **32**: 401-7.
- 316 7. Cabral C, Horwood J, Hay AD *et al*. How communication affects prescription decisions in
317 consultations for acute illness in children: A systematic review and meta-ethnography. *BMC*
318 *Fam Pract* 2014; **15**: 63.

- 319 8. Macfarlane J, Holmes W, Macfarlane R *et al.* Influence of patients' expectations on antibiotic
320 management of acute lower respiratory tract illness in general practice: Questionnaire study.
321 *BMJ* 1997; **315**: 1211-1214.
- 322 9. Coenen S, Francis N, Kelly M *et al.* Are patient views about antibiotics related to clinician
323 perceptions, management and outcome? A multi-country study in outpatients with acute cough.
324 *PLoS One* 2013; **8**: e76691.
- 325 10. Costelloe C, Metcalfe C, Lovering A *et al.* Effect of antibiotic prescribing in primary care on
326 antimicrobial resistance in individual patients: Systematic review and meta-analysis. *BMJ* 2010;
327 **340**: c2096.
- 328 11. Goossens H, Ferech M, Vander Stichele R *et al.*, ESAC Project Group. Outpatient antibiotic
329 use in Europe and association with resistance: A cross-national database study. *Lancet* 2005;
330 **365**: 579-587.
- 331 12. Antimicrobial resistance: Global report on surveillance. World Health Organization; 2014.
332 www.who.int/drugresistance/documents/surveillancereport/en/.
- 333 13. Arnold SR, Straus SE. Interventions to improve antibiotic prescribing practices in ambulatory
334 care. *Cochrane Database Syst Rev* 2005; **4**: CD003539.
- 335 14. Butler CC, Simpson SA, Dunstan F *et al.* Effectiveness of multifaceted educational
336 programme to reduce antibiotic dispensing in primary care: Practice based randomised
337 controlled trial. *BMJ* 2012; **344**: d8173.
- 338 15. Cals JW, Butler CC, Hopstaken RM *et al.* Effect of point of care testing for C reactive protein
339 and training in communication skills on antibiotic use in lower respiratory tract infections: cluster
340 randomised trial. *BMJ* 2009; **338**: b1374. doi:10.1136/bmj.b1374.


- 341 16. Little P, Stuart B, Francis N *et al.* Effects of internet-based training on antibiotic prescribing
342 rates for acute respiratory-tract infections: A multinational, cluster, randomised, factorial,
343 controlled trial. *Lancet* 2013; **382**: 1175-1182.
- 344 17. Mangione-Smith R, McGlynn EA, Elliott MN *et al.* The relationship between perceived
345 parental expectations and pediatrician antimicrobial prescribing behavior. *Pediatrics* 1999; **103**:
346 711-718.
- 347 18. Andrews T, Thompson M, Buckley DI *et al.* Interventions to influence consulting and
348 antibiotic use for acute respiratory tract infections in children: A systematic review and meta-
349 analysis. *PLoS One* 2012; **7**: e30334.
- 350 19. Hu Y, Walley J, Chou R, *et al.* Interventions to reduce childhood antibiotic prescribing for
351 upper respiratory infections: Systematic review and meta-analysis. *J Epidemiol Community*
352 *Health* 2016; **70**: 1162–1170.
- 353 20. Vodicka TA, Thompson M, Lucas P *et al.* Reducing antibiotic prescribing for children with
354 respiratory tract infections in primary care: A systematic review. *Br J Gen Pract* 2013; **63**: e445-
355 e454.
- 356 21. Francis NA, Butler CC, Hood K *et al.* Effect of using an interactive booklet about childhood
357 respiratory tract infections in primary care consultations on reconsulting and antibiotic
358 prescribing: A cluster randomised controlled trial. *BMJ* 2009; **339**: b2885.
- 359 22. Campbell MK, Piaggio G, Elbourne DR *et al.* Consort 2010 statement: Extension to cluster
360 randomised trials. *BMJ* 2012; **345**: e5661.

- 362 23. Yardley L, Douglas E, Anthierens S *et al.* Evaluation of a web-based intervention to reduce
363 antibiotic prescribing for LRTI in six European countries: Quantitative process analysis of the
364 GRACE/INTRO randomised controlled trial. *Implement Sci* 2013; **8**: 134-5908-8-134. doi:
365 10.1186/1748-5908-8-134.
- 366 24. Yardley L, Morrison L, Bradbury K *et al.* The person-based approach to intervention
367 development: Application to digital health-related behavior change interventions. *J Med Internet*
368 *Res* 2015; **17**: e30.
- 369 25. Anthierens S, Tonkin-Crine S, Douglas E *et al.* General practitioners' views on the
370 acceptability and applicability of a web-based intervention to reduce antibiotic prescribing for
371 acute cough in multiple european countries: A qualitative study prior to a randomised trial. *BMC*
372 *Fam Pract* 2012; **13**: 101.
- 373 26. Rollnick S, Kinnersley P, Butler C. Context-bound communication skills training:
374 Development of a new method. *Med Educ* 2002; **36**: 377-383.
- 375 27. van der Velden AW, Kuyvenhoven MM, Verheij TJ. Improving antibiotic prescribing quality
376 by an intervention embedded in the primary care practice accreditation: The ARTI4 randomized
377 trial. *J Antimicrob Chemother* 2016; **71**: 257-263.
- 378 28. Killip S, Mahfoud Z, Pearce K. What is an intracluster correlation coefficient? Crucial
379 concepts for primary care researchers. *Ann Fam Med* 2004; **2**: 204-208.
- 380 29. Faraway JJ. Extending the linear model with R. Generalized linear, mixed effects and
381 nonparametric regression models. Boca Raton: Chapman & Hall/CRC, 2016; 399.

- 382 30. van Duijn HJ, Kuyvenhoven MM, Tiebosch HM *et al.* Diagnostic labelling as determinant of
383 antibiotic prescribing for acute respiratory tract episodes in general practice. *BMC Fam Pract*
384 2007; **8**: 55.
- 385 31. Boonacker CW, Hoes AW, Dikhoff MJ *et al.* Interventions in health care professionals to
386 improve treatment in children with upper respiratory tract infections. *Int J Pediatr*
387 *Otorhinolaryngol* 2010; **74**: 1113-1121.
- 388 32. O'Sullivan JW, Harvey RT, Glasziou PP *et al.* Written information for patients (or parents of
389 child patients) to reduce the use of antibiotics for acute upper respiratory tract infections in
390 primary care. *Cochrane Database Syst Rev* 2016; **11**: CD011360.
- 391 33. de Bont EG, Alink M, Falkenberg FC *et al.* Patient information leaflets to reduce antibiotic
392 use and reconsultation rates in general practice: A systematic review. *BMJ Open* 2015; **5**:
393 e007612.
- 394 34. Spaan NR, Dekker ARJ, van der Velden AW *et al.* Informal and formal learning of general
395 practitioners. *Journal of Workplace Learning* 2016; **28**: 378-391.
- 396 35. Donner A, Klar N. Design and analysis of cluster randomization trials in health
397 research. London: Arnold, 2000; p.178.
- 398 36. Blair PS, Turnbull S, Ingram J *et al.* Feasibility cluster randomised controlled trial of a within-
399 consultation intervention to reduce antibiotic prescribing for children presenting to primary care
400 with acute respiratory tract infection and cough. *BMJ Open* 2017; **7**: e014506.
- 401 37. Gulliford MC, Moore MV, Little P *et al.* Safety of reduced antibiotic prescribing for self
402 limiting respiratory tract infections in primary care: Cohort study using electronic health records.
403 *BMJ* 2016; **354**: i3410.

404 38. Spoelman WA, Bonten TN, de Waal MW *et al.* Effect of an evidence-based website on
405 healthcare usage: An interrupted time-series study. *BMJ Open* 2016; **6**: e013166.

406 **Figure 1 Trial profile, practice flow**



424 **Table 1 Characteristics of general practices allocated to the intervention and control**
425 **group**

	Intervention (n=15)	Control (n=17)
Median list size total (IQR)	2980 (2491-4850)	3275 (2589-3589)
Median list size children <18 years (IQR)	604 (518-999)	664 (421-810)
Participating GPs	40	35
Male/female GPs	46%/54%	43%/57%
Mean age GP (SD)	46 (11)	45.3 (9.5)

426 IQR= interquartile range

427 **Table 2 Characteristics of consultations of the follow-up audit after allocation to the**
 428 **intervention or control group**

	Intervention (n=477)	Control (n=532)
Mean age, years (SD)	4.7 (4.4)	4.4 (4.1)
Median duration of illness before consultation, days (IQR)	5 (3-14)	5 (3-10)
Mean GPs' perception of illness severity, 1 = not ill, 5 = severely ill (SD)	1.6 (0.8)	1.9 (1.0)
Fever (%)	257 (53.9)	278 (52.3)
Earache (%)	177 (37.1)	156 (29.3)
Runny nose (%)	387 (81.1)	375 (70.5)
Sore throat (%)	128 (26.8)	121 (22.7)
Cough (%)	358 (75.1)	381 (71.6)

429 IQR= interquartile range

430 **Table 3 Effectiveness of the intervention on antibiotic prescription rates**

	Intervention	Control	RR (95% CI)
Crude antibiotic prescription rate (95% CI)	20% (95/475) (15.4-26)	36.9% (196/531) (30.8-44.3)	0.54 (0.4-0.74)*
Adjusted antibiotic prescription rate** (95% CI)	21.4% (16.6-27.6)	33.2% (27-40.8)	0.65 (0.46-0.91)*

431 Data were retrieved from GP-registered consultations. *P<0.05. **Adjusted for baseline

432 prescription.

433 **Table 4 Effectiveness of the intervention on reconsultation, consultations for new RTI**
 434 **episodes and hospital referrals**

	Intervention (n=311)	Control (n=197)	RR (95% CI)
Absolute number of reconsultations	132	126	
Mean number of reconsultations/100 children (95% CI)	42 (29-63)	64 (43-96)	0.66 (0.38-1.16)
Absolute number of new RTI consultations	252	150	
Mean number of new RTI consultations/100 children (95% CI)	81 (64-103)	76 (56-104)	1.06 (0.72-1.58)
Absolute number of hospital referrals	24	23	
Mean number of hospital referrals/100 children (95% CI)	8 (5-13)	12 (7-20)	0.66 (0.31-1.40)

435 Data were retrieved from the child's medical registries.

436 **Table 5 Effectiveness of the intervention on total and second choice yearly dispensed**
 437 **antibiotics**

		Intervention	Control	RR (95% CI)
Total antibiotics/1000 children/year (95% CI)	Crude	110 (89.1-136)	161 (137-189)	0.68 (0.52-0.89)*
	Adjusted**	114 (100-129)	146 (132-162)	0.78 (0.66-0.92)*
Number of second choice antibiotics/1000 children/year (95% CI)	Crude	39.3 (29.1-53.1)	54.8 (43.3-69.4)	0.72 (0.49-1.05)
	Adjusted**	39.9 (32.6-48.7)	49.2 (41.7-58.1)	0.81 (0.63-1.05)
Percentage of second choice antibiotics/total antibiotics (95% CI)	Crude	35.7% (29-44)	34% (28.9-40)	1.05 (0.81-1.37)
	Adjusted**	34.1% (29.6-39.3)	34.4% (30.8-38.3)	0.99 (0.83-1.19)

438 Data were retrieved from a full year's pharmacy dispensing data. *P<0.05. ** Adjusted for
 439 baseline prescription.