

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/109355/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Howes, Joanna-Marie, Pugh, Nicholas, Hamaia, Samir W, Jung, Stephanie M., Knauper, Vera, Malcor, Jean-Daniel and Farndale, Richard W 2018. MMP-13 binds to platelet receptors α IIb β 3 and GPVI and impairs aggregation and thrombus formation. *Research and Practice in Thrombosis and Haemostasis* 2 (2) , pp. 370-379. 10.1002/rth2.12088 file

Publishers page: <https://doi.org/10.1002/rth2.12088> <<https://doi.org/10.1002/rth2.12088>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



MMP-13 binds to platelet receptors α IIb β 3 and GPVI and impairs aggregation and thrombus formation.

Joanna-Marie Howes^{1*}, Nicholas Pugh², Samir W. Hamaia¹, Stephanie M. Jung¹, Vera Knäuper³, Jean-Daniel Malcor¹, Richard W. Farndale¹.

¹Department of Biochemistry, University of Cambridge, Downing Site, Cambridge, CB2 1QW, UK;

²Department of Biomedical and Forensic Sciences, Anglia Ruskin University, Cambridge, CB1 1PT;

³Cardiff University Dental School, Dental Drive, CF14 4XY, Cardiff, UK.

Running title: MMP-13 binds to α IIb β 3 and GPVI

*To whom correspondence should be addressed:

Joanna-Marie Howes, Department of Biochemistry, University of Cambridge, Downing Site, Cambridge, CB2 1QW, UK. Phone: +44(0)1223333643; Fax: +44(0)1223333345; E-mail: jmh206@cam.ac.uk

This study was supported by Programme grants from British Heart Foundation, RG/09/003/27122 and RG/15/4/31268 to RWF. Work in V. Knäuper's laboratory is supported by Tenovus.org.

Keywords: Integrin α IIb β 3 (α IIb β 3), GPVI collagen receptor, Matrix Metalloproteinase-13, platelets, thrombosis.

ESSENTIALS

- **MMP-13 has the potential to influence platelet function and thrombus formation directly**
- **We sought to elucidate whether MMP-13 is able to bind to specific platelet receptors**
- **MMP-13 is able to bind to platelet α IIb β 3 (α IIb β 3) and glycoprotein (GP)VI**
- **These interactions are sufficient to inhibit platelet aggregation and thrombus formation**

BACKGROUND. Acute thrombotic syndromes lead to atherosclerotic plaque rupture with subsequent thrombus formation, myocardial infarction and stroke. Following rupture, flowing blood is exposed to plaque components, including collagen, which triggers platelet activation and aggregation. However, plaque rupture releases other components into the surrounding vessel which have the potential to influence platelet function and thrombus formation. **OBJECTIVES.** Here we sought to elucidate whether matrix metalloproteinase-13 (MMP-13), a collagenolytic metalloproteinase up-regulated in atherothrombotic and inflammatory conditions, affects platelet aggregation and thrombus formation. **RESULTS.** We demonstrate that MMP-13 is able to bind to platelet receptors α IIb β 3 (α IIb β 3) and platelet glycoprotein (GP)VI. The interactions between MMP-13, GPVI and α IIb β 3 are sufficient to significantly inhibit washed platelet aggregation and decrease thrombus formation on fibrillar collagen. **CONCLUSIONS.** Our data demonstrate a role for MMP-13 in the inhibition of both platelet aggregation and thrombus formation in whole flowing blood, and may provide new avenues of research into the mechanisms underlying the subtle role of MMP-13 in atherothrombotic pathologies.

Platelet-extracellular matrix and platelet-platelet adhesions are central to the formation of thrombi. MMP-13 is up-regulated in inflammation, and is elevated in the atherosclerotic plaque, contributing to its vulnerability [1]. It is also implicated in the progression and remodelling of cerebral tissue in stroke [2]. Plaque rupture releases MMP-13 into the local environment where it has direct access to plasma proteins, blood cells and platelets. Following injury to the blood vessel wall, specific platelet receptors mediate platelet-collagen and platelet-platelet interactions. GPIIb α binds to immobilised von Willebrand factor (VWF) in the vessel wall, initiating platelet capture [3], and glycoprotein (GP)VI binds directly to collagen and activates platelets. Integrin α 2 β 1 stabilises the early stages of the platelet-collagen interaction, and integrin α IIb β 3 supports platelet-platelet interactions mediated by fibrinogen and VWF [4-7].

MMP-2 and -9 have previously been shown to bind to platelet receptors and/or to modulate platelet function [8-13]. Here, we hypothesised that MMP-13 may also interact directly with platelet receptors GPVI, integrin $\alpha 2\beta 1$, or the platelet adhesive integrin $\alpha \text{IIb}\beta 3$ to modulate platelet adhesion, aggregation and thrombus formation. Our work identifies potential roles for MMP-13 in modulating the recruitment or activation of platelets in thrombotic pathologies.

METHODS

MMP-13 expression, purification and activation

ProMMP-13 and its MMP-13 catalytic (CAT, 249-451) domain were expressed, purified, activated and dialysed as previously described [14-16]. The structurally homologous but catalytically inactive ProMMP-13(E204A) was a kind gift from Dr. R. Visse (Kennedy Institute of Rheumatology Division, Imperial College London, London, UK). GST-Hemopexin (HPX) domain (264-471) was expressed in *E. coli* using the pGEX-2T expression vector, the forward primer TCCGCGTGGATCCCTCTATGGTCCAGGAGATGAA and the reverse primer GCAA-ATTCCATTTTGTGGTGTGAAGAATTCAT, which contain BamHI and EcoRI restriction sites respectively, as previously described [16].

Washed platelet preparation and platelet adhesion assays

Plates were coated with 10 $\mu\text{g/ml}$ MMP-13 variants in Tris buffered saline (TBS) for 1 h at 24 °C. Plates were then blocked with 5 % BSA in TBS for 20 minutes at 24 °C and washed with TBS prior to the addition of washed platelets. Platelets were purified and adhesion assays conducted as previously described [17, 18]. Glanzmann thrombasthenic blood was kindly provided by Prof M. Makris, Royal Hallamshire Hospital, Sheffield, UK. GR144053 (4-[4-(4-aminoiminomethyl)-1-piperazinyl]-1-piperidineacetic acid hydrochloride trihydrate) was purchased from Calbiochem, Nottingham, UK. The $\alpha 2\beta 1$ integrin-binding peptide GFOGER

(GPC[GPP]5-GFOGER-(GPP)5-GPC) and GPVI-binding peptide CRP-(XL) (GCO-[GPO]10-GCOG; cross-linked where appropriate) and the inert GPP10 (GPC-[GPP]10-GPC) were generated as previously described [7] along with the anti-GPVI scFvs 10B12 and 1C3 and the non-GPVI-binding scFv 2D4 [19-23] which were a kind gift from Dr P. Smethurst. Human fibrinogen type I was purchased from Sigma, UK. Anti- $\alpha 2\beta 1$ antibody 6F1 was a kind gift from Prof B. Coller (Mount Sinai Hospital, New York, USA). RGDS (Arg-Gly-Asp-Ser) and cyclic RGD (H-Cys-Arg-Gly-Asp-Phe-Pro-Ala-Ser-Ser-Cys-OH) were purchased from Bachem, Weil am Rhein, Germany. The fibrinogen-derived peptide, Lys-Gln-Ala-Gly-Asp-Val (KQAGDV), was purchased from Innovagen, Sweden. Inhibitory antibodies/compounds were used at 10 μ M (6F1, 10B12, 1C3, 2D4, cRGD, GR144053, KQAGDV) or 100 μ M (fibrinogen and RGDS).

Flow cytometry

In activation experiments, whole blood diluted 1:4 with HEPES buffered saline (HBS) was mixed for 10 min at 24 °C with an equal volume of 10 μ g/ml mouse anti P-Selectin (Abcam, Cambs, UK) and the following agonists 2 mM proMMP-13(E204A) or MMP-13, 100 μ g/ml CRP-XL, 100 μ g/ml HORM® equine collagen I fibers (Takeda, Linz, Austria), thrombin activating peptide (TRAP; 500 μ M; Sigma, UK) calcium ionophore A23187 (100 μ M; Sigma UK) or HBS (negative control) added. Alexa 488 conjugated anti-mouse (30 μ g/ml final concentration; Jackson Immuno Research, Ely, UK) was then added and after 10 min at 24 °C the volume was made up to 500 μ l with isotonic solution. After 30 min fluorescence was measured using an Accuri C6 flow cytometer (BD Biosciences, Oxford, UK). In inhibition experiments, whole blood was pre-incubated with proMMP-13(E204A), GR144053 (20 μ M) or 10B12 (10 μ g/ml) for 20 minutes prior to the addition of CRP-XL.

Solid phase adhesion assays.

Recombinant human α IIb β 3 and GPVI monomer were obtained from R&D Systems (Abingdon, Oxford, UK). Recombinant extracellular domain of GPVI (GPVI_{ex}, comprising D1D2 (amino acids 1–214; 42 kDa) fused with the Fc domain of human IgG (GPVI-Fc2, 150 kDa) was prepared as previously described [24].

HB 96-well plates (Nunc, Langensfeld, Germany) were coated with recombinant GPVI monomer or dimer (10 μ g/ml in Phosphate-Buffered Saline (PBS)) for 1 h at 24 °C. All further incubations were performed at room temperature for 1 h unless otherwise stated. The wells were washed three times with adhesion buffer (1 mg/ml BSA in PBS containing 0.1 % (v/v) Tween-20) between each incubation step. The wells were then blocked with 50 mg/ml BSA in TBS prior to the addition of MMP-13 at a concentration of 83 nM (unless otherwise stated) for 1 h at 24 °C in adhesion buffer. Rabbit anti-MMP-13, raised against MMP-13 hinge region (Abcam, Cambridge, UK), and goat anti-rabbit HRP (Dako, Stockport, UK) were added at a dilution of 1:2000 in adhesion buffer prior to the addition of a TMB substrate system (Sigma, UK) and the plates read at 450 nm.

Aggregometry

Washed platelet aggregation was performed using a Chrono-Log turbidimetric aggregometer (Labmedics, Abingdon on Thames, UK). 250 μ l aliquots of platelets, 2×10^8 /ml in calcium-free Tyrodes buffer (CFT), were pre-incubated for 1 h with 80 nM MMP-13 or vehicle control prior to the addition of receptor agonists in a maximum volume of 5 μ l. Thrombin, calcium ionophore A23187 (San Diego, CA, USA), bovine collagen I fibers (Ethicon Corp, Somerville,

NJ, USA), HORM® and CRP–XL were prepared and employed to activate platelets as previously described [25, 26]. Aggregations were allowed to proceed for 5 min.

Cleavage of platelet receptors and their substrates by MMP-13

Recombinant human (rh)GPVI, purified α IIB β 3 (100 μ g/ml, R&D Systems) and human fibrinogen type I (1 mg/ml) were incubated with MMP-13 or MMP-13(E204A) (8 μ M final concentration) for 2 h at 37 °C. An equal volume of Tris buffer was used as a negative control. Reducing sample buffer was then added to the mixture in preparation for electrophoresis and Western blotting.

***In vitro* sheddase activity assays**

Dialysed MMP-13 at a final concentration of 130 nM was incubated with washed platelets for 60 min at 37 °C. Positive controls for shedding included thrombin (1 U/ml, Sigma, UK) combined with fibrous type I collagen (1 mg/ml), the calcium ionophore A23187 (1 μ g/ml). The platelets were then pelleted at 1500 g for 1 min. The supernatants were aspirated and centrifuged again to ensure platelet depletion. This new supernatant was retained for analysis. Where indicated, platelet lysate was resuspended in reducing sample buffer.

Electrophoresis and Western blotting

Protein samples in reducing sample buffer were boiled for 5 min and applied to 4-12 % NuPage® Gels and separated by electrophoresis using the Xcell SureLock™ system (Invitrogen) under reducing conditions. Proteins were then transferred on to nitrocellulose membrane (Millipore, Bedford, UK) at 40 V overnight at 4 °C using a Mini Protean II system (Bio-Rad, Hemel Hempstead, UK). Following transfer, the PVDF was blocked (5 % non-fat dry powdered milk, 0.1 % Tween 20 in TBS) for 1 h and primary antibody was then added

(1:1000 dilution) and incubated for 2 h at room temperature. Anti-human GPVI was a kind gift from Dr P. Smethurst, and anti- $\beta 3$ was obtained from Abcam, Cambridge, UK. Following washes with TBST, the membrane was incubated with HRP conjugated secondary antibody (1:10000 dilution/TBST) for 1 h at 24 °C. The PVDF was developed using a chemiluminescent substrate (GE Healthcare, Amersham, Bucks, UK).

Whole blood perfusion experiments

Whole blood was pre-incubated with either carrier (TBS) or 80 nM MMP-13 for 1 h prior to perfusion over 10 μ g/ml type I fibrous collagen as previously described [17, 25]. Where indicated, slides were coated with MMP-13(E204A) alone as a (negative) control.

RESULTS

Adhesion assays were performed in the presence of 2 mM EDTA or Mg^{2+} to ablate or support integrin-mediated adhesion. Platelet adhesion to MMP-13 preparations was significantly reduced, but not abolished, by EDTA, suggesting both integrin-dependent and -independent contributions, whereas EDTA fully abolished binding to the collagen-binding integrin-specific peptide GFOGER (Figure 1A).

Platelet pre-incubation with the α IIB β 3 antagonists, GR144053, cRGD and RGDS, and with anti-GPVI scFv 10B12 and 1C3, all caused a substantial and significant reduction ($p < .01$) in platelet adhesion to proMMP-13 (Figure 1A), with residual adhesion being observed in the presence of EDTA remaining above negative control levels (non-specific substrates). This may indicate co-operative binding to α IIB β 3 and GPVI. Interaction between MMP-13 and integrin α 2 β 1 was less prominent, since blocking antibody 6F1 had just a small effect, and was not studied further. Platelet pre-incubation with the fibrinogen-derived peptide, KQAGDV, had

no effect on platelet adhesion, indicating that MMP-13 binds α IIB β 3 closer to the primary RGD-binding site. Soluble fibrinogen also did not block platelet adhesion to MMP-13, in line with the need for platelet activation for soluble fibrinogen binding to α IIB β 3 to occur, whereas immobilised fibrinogen is already competent to bind. The GPVI-specific scFv, 1C3, does not target the collagen-binding site at the apex of GPVI, unlike 10B12, and was unable to inhibit the adhesion of washed platelets to CRP (Figure 1B). 1C3 binding requires both GPVI Ig domains and is thought to reduce platelet activation by inhibiting receptor clustering; its epitope includes isoleucine 148 [21, 23], located in strand E on the opposite face of D2 to the crystal structure dimerisation interface located in strand G [24]. An indifferent control, the anti-HLA-A2 scFv, 2D4, was inactive in these experiments. In subsequent experiments, only low platelet binding was observed to isolated CAT and HPX domains of MMP-13 in comparison with the intact protein (Figure 1C), indicating that neither domain alone governs the interaction between the MMP and platelets, and supporting the possibility that two sites on MMP-13 co-operate to bind α IIB β 3 and GPVI.

Competition assays in which washed platelets were pre-incubated with increasing amounts of the catalytically-dead MMP-13(E204A) provided further evidence that MMP-13 interacts with both GPVI and α IIB β 3; like GR144053 and 10B12, MMP-13 can compete α IIB β 3 off immobilised fibrinogen and GPVI off CRP (IC₅₀ 150 ng/ml and ~10 ng/ml respectively; Figures 2A(i-iv)). Solid phase binding assays to coated isolated receptors revealed that MMP-13 was able to bind weakly to GPVI monomer, but strongly to the GPVI dimer (Figure 2A(v)). Similar assays of adhesion to recombinant α IIB β 3 revealed some binding of its native ligand, fibrinogen, but little or no binding of MMP-13, regardless of whether Mg²⁺, Mn²⁺ or Ca²⁺ was present, nor could we detect binding of MMP-13 to purified α IIB β 3 (results not shown). Adhesion of α IIB β 3-null Glanzmann platelets to MMP-13, however, was markedly reduced (Figure 2B(i)). Blockade of α IIB β 3 on healthy platelets resulted in the same adhesion level as

seen for α IIB β 3-null platelets. As expected, binding of Glanzmann platelets to fibrinogen was abolished (Figure 2B(ii)) and to CRP was unaffected (Figure 2B(iii)). Our results indicate that, whilst MMP-13 appears able to bind to α IIB β 3 on the platelet surface, recombinant α IIB β 3 used here cannot reproduce this effect.

Whilst it was able to cleave the recombinant α IIB β 3 β -chain and GPVI in solution, as well as fibrinogen α and β chains (Figure 3A), MMP-13 was unable to either cause or mediate shedding of either receptor in situ (Figure 3B).

Pre-incubation of washed platelets for 1 h with 80 nM MMP-13(E204A) significantly reduced platelet aggregation to a series of agonists, and for the mid-range dose of each, analysed using 2-way ANOVA, the inhibitory effect of MMP-13 was significant ($p < .01$). Prominent amongst these stimuli were: CRP-XL, ionophore A23187, and bovine fibrillar collagen I, for which it was easier to establish mid-range doses than for thrombin and equine fibrillar collagen. A summary of results is shown in Figure 4A and representative traces in Figure 4B. MMP-13 does not activate platelets measured by flow cytometry: no change in fluorescence using the anti-P-Selectin antibody was observed following the incubation of whole blood with pro-MMP-13(E204A) or MMP-13, whereas clear expression was seen following treatment with CRP-XL, TRAP, HORM® and ionophore A23187 (Figure 5A). In addition, MMP-13 does not promote the aggregation of washed platelets (Figure 4B(i)). Subsequent flow cytometry experiments revealed that unlike the anti-GVI scFv 10B12, neither proMMP-13(E204A) nor GR144053 (a potent α IIB β 3 antagonist) are able to alter secretion following platelet activation via CRP-XL (Figure 5B). This would suggest that in solution, the polymeric CRP-XL is a more potent ligand than MMP-13, and that the interaction of MMP-13 with α IIB β 3 predominates over that with GPVI.

We investigated the influence of MMP-13 or MMP-13(E204A) on platelet adhesion and activation in flowing blood *in vitro*, using fibrillar collagen I coatings and a shear rate of 1000s⁻¹

¹. Pre-incubation of whole blood with MMP-13 resulted in significantly reduced platelet surface coverage ($p < .05$), mean thrombus height ($p < .01$), and ZV_{50} ($p < .05$), using one-way ANOVA and Holm multiple comparison test; Figure 6A(i-iii). ZV_{50} is the height within a Z-stack at which thrombus volume = 50% and describes the activation state of adhered platelets in flowing human blood [25]. Data obtained using pre-incubations with MMP-13(E204A) reached significance only for mean thrombus height ($p < .05$, Figure 6A(ii)). MMP-13(E204A)-coated slides were not able to support platelet adhesion under flow (Figure 6A(i-iii)). These results indicate that the interaction of MMP-13(E204A) with platelet GPVI and $\alpha IIb\beta 3$ is sufficient to reduce platelet thrombus height. Catalytically active MMP-13, whilst unable to cleave these receptors off the platelet surface, appears more able to inhibit platelet deposition. MMP-13 co-coated with collagen type I did not significantly alter platelet aggregate formation under flow conditions (Figure 6B(i-iii)). Interaction of active MMP-13 with other blood components is not excluded by the present work, and further study is indicated.

DISCUSSION

We have previously shown that degradation by MMP-13 has the potential to modulate platelet adhesion to collagen [17]. MMPs are zymogens; proteolysis is required to expose their catalytic site. Here we show that surprisingly, all forms of MMP-13, pro- and active wild type enzyme as well as their catalytically inactive mutant counterparts, were able to support a high level of platelet adhesion under static conditions. This adhesion was inhibited by the anti-GPVI scFvs 10B12 and 1C3 suggesting that the relatively large MMP-13 occludes the sites of both 10B12 and 1C3 binding on the receptor. MMP-13 was also able to bind strongly to the GPVI dimer. Although GPVI dimerization increases upon platelet activation, dimeric GPVI is also present on resting platelets and is required for their initial interaction with exposed collagen [26].

Crystallography of the proMMP-13 structure in complex with pro-domain peptides revealed a dimeric form as an HPX-mediated dimer like some other metalloproteinases, although in this study [27], MMP-13 was not dimeric in solution. Conceivably, interaction of MMP-13 with platelet surface GPVI dimer may provide a template for dimerization of the MMP. Platelet adhesion to MMP-13 was also inhibited by the anti- α IIB β 3 compound GR144053, and binding of Glanzmanns α IIB β 3-null platelets to MMP-13 was significantly reduced. Following pre-incubation of washed platelets with MMP-13, neither GPVI nor α IIB β 3 was shed from the platelet surface. It would appear, therefore, that whilst able to bind to platelet α IIB β 3 and GPVI, the orientation of MMP-13 on the platelet surface does not allow access of its CAT domain to the cleavage site, which, for other sheddases, resides close to the trans-membrane region and is regulated by membrane structure [28] or substrate phosphorylation [29]. Pre-incubation with MMP-13 did not result in platelet activation or aggregation. Here it is worth noting that MMP-13 has been reported to cleave and thus activate PAR-1 on cardiac cells [30]. This has not been demonstrated on platelets, and may result in platelet activation concomitant with α IIB β 3 inhibition, however in this case the catalytically inactive proMMP-13(E204A) is rendered unable to cleave the PAR-1 receptor.

Coated as a substrate, MMP-13 is independently unable to support platelet adhesion in whole flowing blood, and its co-localisation with collagen does not result in an increase in platelet binding. In solution however, MMP-13 is able to interact with platelet receptors GPVI and α IIB β 3 thereby modulating both platelet aggregation and thrombus formation under flow. Whilst MMP-13 is able to compete with immobilised CRP-XL for occupation of the GPVI receptor, our flow cytometry experiments reveal that the interaction of the MMP with platelets is insufficient to compete with the polymeric solution-phase CRP-XL and so alter platelet secretion. In this respect, it behaves much like the α IIB β 3 antagonist, GR144053, and it would appear therefore that the inhibitory effects of MMP-13 in solution are mediated predominantly

through α I**IIb** β 3. At concentrations comparable to those reached in stroke patient plasma and found to correlate with severity of infarction [31], MMP-13 can interact with both GPVI and α I**IIb** β 3, and can compete with CRP and fibrinogen for occupation of these receptors. MMP-13 is unable to cleave GPVI and α I**IIb** β 3 from the platelet surface however, and appears to exert its effects by direct physical blockade of receptor engagement.

Until now, the role of MMP-13 in atherothrombosis has been considered to be restricted to collagen proteolysis and remodelling, rendering plaque more friable and prone to rupture [1]. However, MMPs are now emerging as important mediators of platelet function [32, 33]. MMPs -1 and -2 are released from activated platelets where they co-localise with integrins at the sites of platelet-platelet interaction [10, 34]. Active MMP-1 and -2 can stimulate platelet function, suggesting receptor engagement and proteolysis [34, 35]. MMPs in atherosclerotic lesions, released from the injured vessel wall itself or from platelets and monocytes, and that can also interact with platelets, are likely to interfere with the progression of plaque rupture, subsequent thrombosis and its associated pathologies including stroke, reperfusion injury and haemorrhagic transformation. Indeed, these processes are associated with an upregulation of MMP activity [2, 31, 36]. In mice, MMP-13 is the key mediator of collagen degradation in atheroma and confers instability onto the vulnerability plaque cap [37-39]. Disruption of the blood brain barrier (BBB) by MMPs is associated with haemorrhagic transformation following ischemic stroke [36, 40, 41], whilst MMPs -9 and -13 are implicated in the early pathology of stroke progression, and plasma MMP-13 levels correlate with lesion volume [2, 31]. In addition, the platelet collagen receptor GPVI has been identified in models of models of reperfusion injury [42], is associated with increased risk of stroke development, and is also seen after ischemic stroke [43].

Here we demonstrate that MMP-13 can exert an anti-thrombotic effect; inhibiting platelet aggregation and thrombus formation in flowing whole blood. It may be that this

metalloproteinase has multiple roles in the pathology of ischaemic stroke; firstly by undermining the stability of the fibrous cap of atheroma and so promoting its rupture, then modulating the BBB to increase bleeding risk, and finally acting on platelets to impair the aggregatory interactions, by antagonising GPVI and $\alpha\text{IIb}\beta\text{3}$ which would normally protect against bleeding. MMP-13 would appear therefore to modulate the architecture around sites of infarction to increase both risk of stroke and its haemorrhagic complications. The effect of MMP-13 will depend upon its local level and the exposure of MMP-13-binding matrix components and warrants further investigation.

FIGURE LEGENDS

Figure 1. Washed platelet adhesion assays. (A) Platelets adherent to 10 $\mu\text{g/ml}$ coated MMP-13 variants in the presence of 2 mM Mg^{2+} (red bars) or 2 mM EDTA (orange bars) where stated. Where appropriate, platelets were pre-incubated with anti-GPVI, $\alpha\text{IIb}\beta 3$ or $\alpha 2\beta 1$ antagonists. BSA and GPP₁₀ were used as Mg^{2+} -independent negative controls. The platelet $\alpha 2\beta 1$ binding-peptide GFOGER was included as an Mg^{2+} dependent positive control. * $p < .05$; ** $p < .01$; † (one-way ANOVA and Holm multiple comparison test) relative to untreated platelets in either the presence of Mg^{2+} or EDTA, as appropriate. (B) Inhibition of platelet adhesion to CRP by anti-GPVI scFvs as described above (C) Platelets adherent to MMP-13(E204A) and MMP-13 CAT and HPX domains. CRP-XL was used as a positive control. ** $p < .01$ (one-way ANOVA and Holm multiple comparison test) relative to adhesion to MMP-13(E204A). Data represent mean $A_{405} \pm \text{S.E.}$ of three experiments.

Figure 2. Competition, GPVI and Glanzmann platelet binding assays. (A) MMP-13(E204A) and either 10B12 or GR144053 were used to obtain IC_{50} values for the inhibition of washed platelet adhesion to 10 $\mu\text{g/ml}$ coated fibrinogen (i, iii) or CRP (ii, iv) respectively. (v) Adhesion of MMP-13(E204A) to recombinant human GPVI monomer and dimer. Plates were coated with 10 $\mu\text{g/ml}$ GPVI or BSA as a negative control. MMP-13(E204A) at a concentration of 83 nM was allowed to adhere for 1 h at room temperature, then detected using an antibody directed at the MMP-13 linker region, as described in Experimental Procedures. Data represent mean $A_{450} \pm \text{S.E.}$ of three experiments. (B) Platelets from a healthy donor (red bars) and from a Glanzmann thrombasthenic individual (orange bars) were allowed to adhere to MMP-13, Fibrinogen and CRP-XL coated plates. Where appropriate, platelets were pre-incubated with anti-GPVI (1C3), or $\alpha\text{IIb}\beta 3$ antagonists as described for Figure 1. Data represent mean $A_{405} \pm \text{S.E.}$ of duplicate readings for one experiment due to the rarity of the Glanzmann donor.

Figure 3. Platelet receptor cleavage and shedding assays. (A) Degradation analysis of recombinant platelet receptors by active and MMP-13(E204A). Recombinant human (rh)GPVI and purified $\alpha\text{IIb}\beta 3$ and fibrinogen type I were incubated with MMP-13 or MMP-13(E204A) for 2 h at 37 °C. An equal volume of Tris buffer was used as a negative control. Samples were subjected to electrophoresis under reducing conditions and Coomassie stained. Images are representative of three experiments. (B) Shedding analysis of platelet receptors. Washed platelets were incubated with the calcium ionophore A23187, a thrombin and collagen type I mixture or MMP-13 for 1 h at 37 °C. The platelets were then pelleted, the supernatant isolated and subjected to SDS-PAGE under reducing conditions and Western blotted. Platelet GPVI and the integrin $\beta 3$ chain were detected using the appropriate antibodies as described in materials and methods. Recombinant human GPVI or $\alpha\text{IIb}\beta 3$ were loaded onto the gels where appropriate as positive controls. Images are representative of three experiments.

Figure 4. Inhibition of platelet aggregation by MMP-13(E204A). Different agonists were added to washed platelets following pre-incubation with 80 nM MMP-13(E204A). The equivalent volume of 0.01 M acetic acid was used as a negative control. Mean donor responses performed in duplicate and repeated three times with different donors are shown in (A), and representative individual traces in response to MMP-13 only, A23187, thrombin, cross-linked collagen related peptide (CRP-XL), HORM® and type I collagen shown in (B).

Figure 5. Activation of platelets in whole blood. Whole blood was mixed with anti P-Selectin and the agonists 2 mM proMMP-13(E204A)/MMP-13, 100 $\mu\text{g/ml}$ CRP-XL, 100 $\mu\text{g/ml}$ HORM® equine collagen I fibers, thrombin activating peptide (TRAP; 500 μM) calcium ionophore A23187 (100 μM) or HBS (negative control) added. Alexa 488 conjugated anti-mouse was then added and after 10 min at 24 °C the volume was made up to 500 μl with isotonic solution. After 30 min fluorescence was measured using an Accuri C6 flow cytometer (BD Biosciences, Oxford, UK). Data represent mean $A_{450} \pm \text{S.E.}$ of three separate donors. ** $p < .005$; (one-way ANOVA and Holm multiple comparison test).

Figure 6. Platelet adhesion and thrombus deposition on fibrillar type I collagen. Untreated whole blood and blood pre-incubated with 80 nM MMP-13 or negative control where stated was drawn through a flow chamber for 5 min over (A) collagen type I fibers or (B) collagen type I fibers co-coated

with MMP-13 using a syringe pump to generate a wall shear rate of 1000s^{-1} , corresponding to arteriolar conditions. Surface coverage (i) mean height (ii) and (iii) ZV_{50} are the mean taken from a minimum of three different donors as measured using confocal microscopy. * $p < .05$; ** $p < .01$; (one-way ANOVA and Holm multiple comparison test) relative to MMP-untreated platelets.

Role and contribution of each author:

Vera Knäuper, Jean-Daniel Malcor and Stephanie Jung provided essential materials, Nicholas Pugh assisted with flow experiments, Richard Farndale designed the research, helped analyse the data and write the manuscript, and Joanna-Marie Howes designed and performed the research, and wrote the manuscript.

The authors would also like to thank Dr. Isuru Induruwa (Stroke Medicine, Department of Clinical Neurosciences, University of Cambridge) for constructive reading of this manuscript.

Disclosure of Conflict of Interest

The authors have no conflict of interest.

REFERENCES

1. Sukhova GK, Schonbeck U, Rabkin E, Schoen FJ, Poole AR, Billingham RC, and Libby P. Evidence for increased collagenolysis by interstitial collagenases-1 and -3 in vulnerable human atheromatous plaques. *Circulation* 1999; **99**; 2503-2509.
2. Ma F, Martinez-San Segundo P, Barcelo V, Morancho A, Gabriel-Salazar M, Giralt D, Montaner J, and Rosell A. Matrix metalloproteinase-13 participates in neuroprotection and neurorepair after cerebral ischemia in mice. *Neurobiol Dis* 2016; **91**; 236-246.
3. Kroll MH, Harris TS, Moake JL, Handin RI, and Schafer AI. von Willebrand factor binding to platelet GpIb initiates signals for platelet activation. *J Clin Invest* 1991; **88**; 1568-1573.
4. Barnes MJ, Knight CG, and Farndale RW. The collagen-platelet interaction. *Curr Opin Hematol* 1998; **5**; 314-320.
5. Bennett JS. Platelet-fibrinogen interactions. *Ann N Y Acad Sci* 2001; **936**; 340-354.
6. Farndale RW. Collagen-induced platelet activation. *Blood Cells Mol Dis* 2006; **36**; 162-165.
7. Morton LF, Hargreaves PG, Farndale RW, Young RD, and Barnes MJ. Integrin alpha 2 beta 1-independent activation of platelets by simple collagen-like peptides: collagen tertiary (triple-helical) and quaternary (polymeric) structures are sufficient alone for alpha 2 beta 1-independent platelet reactivity. *Biochem J* 1995; **306**; 337-344.
8. Momi S, Falcinelli E, Giannini S, Ruggeri L, Cecchetti L, Corazzi T, Libert C, and Gresele P. Loss of matrix metalloproteinase 2 in platelets reduces arterial thrombosis in vivo. *J Exp Med* 2009; **206**; 2365-2379.
9. Santos-Martinez MJ, Medina C, Jurasz P, and Radomski MW. Role of metalloproteinases in platelet function. *Thromb Res* 2008; **121**; 535-542.
10. Sawicki G, Sanders EJ, Salas E, Wozniak M, Rodrigo J, and Radomski MW. Localization and translocation of MMP-2 during aggregation of human platelets. *Thromb Haemost* 1998; **80**; 836-839.
11. Sebastiano M, Momi S, Falcinelli E, Bury L, Hoylaerts MF, and Gresele P. A novel mechanism regulating human platelet activation by MMP-2-mediated PAR1 biased signaling. *Blood* 2017; **129**; 883-895.
12. Sheu JR, Fong TH, Liu CM, Shen MY, Chen TL, Chang Y, Lu MS, and Hsiao G. Expression of matrix metalloproteinase-9 in human platelets: regulation of platelet activation in in vitro and in vivo studies. *Br J Pharmacol* 2004; **143**; 193-201.
13. Zaman AG, Helft G, Worthley SG, and Badimon JJ. The role of plaque rupture and thrombosis in coronary artery disease. *Atherosclerosis* 2000; **149**; 251-266.
14. Chung L, Dinakarbandian D, Yoshida N, Lauer-Fields JL, Fields GB, Visse R, and Nagase H. Collagenase unwinds triple-helical collagen prior to peptide bond hydrolysis. *Embo J* 2004; **23**; 3020-3030.
15. Kennedy AM, Inada M, Krane SM, Christie PT, Harding B, Lopez-Otin C, Sanchez LM, Pannett AA, Dearlove A, Hartley C, et al. MMP13 mutation causes spondyloepimetaphyseal dysplasia, Missouri type (SEMD(MO)). *J Clin Invest* 2005; **115**; 2832-2842.
16. Knauper V, Lopez-Otin C, Smith B, Knight G, and Murphy G. Biochemical characterization of human collagenase-3. *J Biol Chem* 1996; **271**; 1544-1550.
17. Howes JM, Pugh N, Knauper V, and Farndale RW. Modified platelet deposition on matrix metalloproteinase 13 digested collagen I. *J Thromb Haemost* 2015; **13**; 2253-2259.
18. Smethurst PA, Onley DJ, Jarvis GE, O'Connor MN, Knight CG, Herr AB, Ouweland WH, and Farndale RW. Structural basis for the platelet-collagen interaction: the smallest motif within collagen that recognizes and activates platelet Glycoprotein VI contains two glycine-proline-hydroxyproline triplets. *J Biol Chem* 2007; **282**; 1296-1304.

19. Jarvis GE, Raynal N, Langford JP, Onley DJ, Andrews A, Smethurst PA, and Farndale RW. Identification of a major GpVI-binding locus in human type III collagen. *Blood* 2008; **111**; 4986-4996.
20. Konitsiotis AD, Raynal N, Bihan D, Hohenester E, Farndale RW, and Leitinger B. Characterization of high affinity binding motifs for the discoidin domain receptor DDR2 in collagen. *J Biol Chem* 2008; **283**; 6861-6868.
21. O'Connor MN, Smethurst PA, Davies LW, Joutsu-Korhonen L, Onley DJ, Herr AB, Farndale RW, and Ouwehand WH. Selective blockade of glycoprotein VI clustering on collagen helices. *J Biol Chem* 2006; **281**; 33505-33510.
22. Siljander PR, Hamaia S, Peachey AR, Slatter DA, Smethurst PA, Ouwehand WH, Knight CG, and Farndale RW. Integrin activation state determines selectivity for novel recognition sites in fibrillar collagens. *J Biol Chem* 2004; **279**; 47763-47772.
23. Smethurst PA, Joutsu-Korhonen L, O'Connor MN, Wilson E, Jennings NS, Garner SF, Zhang Y, Knight CG, Dafforn TR, Buckle A, et al. Identification of the primary collagen-binding surface on human glycoprotein VI by site-directed mutagenesis and by a blocking phage antibody. *Blood* 2004; **103**; 903-911.
24. Horii K, Kahn ML and Herr AB. Structural basis for platelet collagen responses by the immune-type receptor glycoprotein VI. *Blood* 2006; **108**; 936-942.
25. Pugh N, Simpson AM, Smethurst PA, de Groot PG, Raynal N, and Farndale RW. Synergism between platelet collagen receptors defined using receptor-specific collagen-mimetic peptide substrata in flowing blood. *Blood* 2010; **115**; 5069-5079.
26. Jung SM, Moroi M, Soejima K, Nakagaki T, Miura Y, Berndt MC, Gardiner EE, Howes JM, Pugh N, Bihan D, et al. Constitutive dimerization of glycoprotein VI (GPVI) in resting platelets is essential for binding to collagen and activation in flowing blood. *J Biol Chem* 2012; **287**; 30000-30013.
27. Tochowicz A, Goettig P, Evans R, Visse R, Shitomi Y, Palmisano R, Ito N, Richter K, Maskos K, Franke D et al. The Dimer Interface of the Membrane Type 1 Matrix Metalloproteinase Hemopexin Domain. *J Biol Chem* 2011; **286**; 7587-7600.
28. Sommer A, Bhakdi S, and Reiss K. How membrane asymmetry regulates ADAM17 sheddase function. *Cell Cycle* 2016; **15**; 2995-2996.
29. Parra LM, Hartmann M, Schubach S, Li Y, Herrlich P, and Herrlich A. Distinct Intracellular Domain Substrate Modifications Selectively Regulate Ectodomain Cleavage of NRG1 or CD44. *Mol Cell Biol* 2015; **35**; 3381-3395.
30. Jaffre F, Friedman AE, Hu Z, Mackman N, and Blaxall BC. beta-adrenergic receptor stimulation transactivates protease-activated receptor 1 via matrix metalloproteinase 13 in cardiac cells. *Circulation* 2012; **125**; 2993-3003.
31. Rosell A, Alvarez-Sabin J, Arenillas JF, Rovira A, Delgado P, Fernandez-Cadenas I, Penalba A, Molina CA, and Montaner J. A matrix metalloproteinase protein array reveals a strong relation between MMP-9 and MMP-13 with diffusion-weighted image lesion increase in human stroke. *Stroke* 2005; **36**; 1415-1420.
32. Nanni S, Melandri G, Hanemaaijer R, Cervi V, Tomasi L, Altimari A, Van Lent N, Tricoci P, Bacchi L, and Branzi A. Matrix metalloproteinases in premature coronary atherosclerosis: influence of inhibitors, inflammation, and genetic polymorphisms. *Transl Res* 2007; **149**; 137-144.
33. Trivedi V, Boire A, Tchernychev B, Kaneider NC, Leger AJ, O'Callaghan K, Covic L, and Kuliopulos A. Platelet matrix metalloproteinase-1 mediates thrombogenesis by activating PAR1 at a cryptic ligand site. *Cell* 2009; **137**; 332-343.
34. Galt SW, Lindemann S, Allen L, Medd DJ, Falk JM, McIntyre TM, Prescott SM, Kraiss LW, Zimmerman GA, and Weyrich AS. Outside-in signals delivered by matrix metalloproteinase-1 regulate platelet function. *Circ Res* 2002; **90**; 1093-1099.30.

35. Sawicki G, Salas E, Murat J, Miszta-Lane H, and Radomski MW. Release of gelatinase A during platelet activation mediates aggregation. *Nature* 1997; **386**; 616-619.
36. Chaturvedi M, and Kaczmarek L. MMP-9 inhibition: a therapeutic strategy in ischemic stroke. *Mol Neurobiol* 2014; **49**; 563-573.
37. Quillard T, Araujo HA, Franck G, Tesmenitsky Y, and Libby P. Matrix metalloproteinase-13 predominates over matrix metalloproteinase-8 as the functional interstitial collagenase in mouse atheromata. *Arterioscler Thromb Vasc Biol* 2014; **34**; 1179-1186.
38. Cheng C, Tempel D, van Haperen R, van Damme L, Algur M, Krams R, and de Crom R. Activation of MMP8 and MMP13 by angiotensin II correlates to severe intra-plaque hemorrhages and collagen breakdown in atherosclerotic lesions with a vulnerable phenotype. *Atherosclerosis* 2009; **204**; 26-33.
39. Quillard T, Tesmenitsky Y, Croce K, Travers R, Shvartz E, Koskinas KC, Sukhova GK, Aikawa E, Aikawa M, and Libby P. Selective inhibition of matrix metalloproteinase-13 increases collagen content of established mouse atherosclerosis. *Arterioscler Thromb Vasc Biol* 2011; **31**; 2464-2472.
40. Lakhan SE, Kirchgessner A, Tepper D, and Leonard A. Matrix metalloproteinases and blood-brain barrier disruption in acute ischemic stroke. *Front Neurol* 2013; **4**; 32.
41. Jickling GC, Liu D, Stamova B, Ander BP, Zhan X, Lu A, and Sharp FR. Hemorrhagic transformation after ischemic stroke in animals and humans. *J Cereb Blood Flow Metab* 2014; **34**; 185-199.
42. Nieswandt B, Kleinschnitz C, and Stoll G. Ischaemic stroke: a thrombo-inflammatory disease? *J Physiol* 2011; **589**; 4115-4123.
43. Bigalke B, Stellos K, Geisler T, Seizer P, May AE, Melms A, Luft A, and Gawaz M. Expression of platelet glycoprotein VI is associated with transient ischemic attack and stroke. *Eur J Neurol* 2010; **17**; 111-117.