

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/111103/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Rainer, Timothy H. , Leung, L.Y., Chan, C.P.Y., Leung, Y.K., Cheng, N.M., Lai, P.B.S., Cheung, Y.S. and Graham, C.A. 2017. Circulating human leucine-rich  $\alpha$ -2-glycoprotein 1 mRNA and protein levels to detect acute appendicitis in patients with acute abdominal pain. *Clinical Biochemistry* 50 (9) , pp. 485-490.  
10.1016/j.clinbiochem.2017.02.010

Publishers page: <http://dx.doi.org/10.1016/j.clinbiochem.2017.02.01...>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



**1. Full title**

Circulating human leucine-rich  $\alpha$ -2-glycoprotein 1 mRNA and protein levels to detect acute appendicitis in patients with acute abdominal pain

**2. Running head**

Risk-assessment tool for abdominal pain in the Emergency Department

**3: Authors**

Rainer TH<sup>1\*</sup>, Leung LY<sup>1</sup>, Chan CPY<sup>1</sup>, Leung YK<sup>1</sup>, Cheng NM<sup>1</sup>, Lai PBS<sup>2</sup>, Cheung YS<sup>2</sup>, Graham CA<sup>1</sup>,

**4: Name of institution**

<sup>1</sup> Accident and Emergency Medicine Academic Unit, The Chinese University of Hong Kong.

<sup>2</sup> Department of Surgery, The Chinese University of Hong Kong.

**5. Corresponding author**

\* Correspondence: Professor T H Rainer, Director, Accident and Emergency Medicine Academic Unit, Chinese University of Hong Kong, 2/F, Main Clinical Block and Trauma Centre, Prince of Wales Hospital, Shatin, New Territories, Hong Kong.

E-mail address: [thrainer@cuhk.edu.hk](mailto:thrainer@cuhk.edu.hk) (T.H. Rainer)

Telephone: +852 26321034

Fax: +852 26481469

**6. Key words:** abdominal pain; acute abdomen; acute appendicitis; diagnosis; mRNA

**7. Previous presentation**

**1.** Rainer TH, Leung LY, Chan CPY, Cheng NM, Lai PBS, Cheung YS, Graham CA. Add-On LRG1 Tests For Improving The Prediction Of Acute Appendicitis In Emergency Department Patients With Acute Abdominal Pain: Prospective Observational Study  
At: 16<sup>th</sup> International Conference of Emergency Medicine. Organized by the African College of Emergency Medicine. Cape Town. 18 – 21 April 2016

**2.** Leung LY, Rainer TH, Chan CPY, Leung YK, Lai PBS, Cheung YS, Graham CA. Circulating leucine-rich  $\alpha$ -2-glycoprotein 1 to detect acute appendicitis in patients with acute abdominal pain. *Conference Programme and Abstracts 2014*; C278: 178

At: 15<sup>th</sup> International Conference of Emergency Medicine. Organized by the Hong Kong College of Emergency Medicine. Hong Kong. 11 – 14 June 2014

**8. List of abbreviations:** AA: acute appendicitis; CT-computed tomography; LRG1: Leucine-rich-2-glycoprotein;.

**9. Human gene**

Glyceraldehyde 3-phosphate dehydrogenase: GAPDH

Leucine-rich alpha-2-glycoprotein 1: LRG1

**Word count abstract: 248**

**Word count manuscript: 3663**

## Abstract

**Background:** Elevated levels of circulating plasma and urine leucine-rich-2-glycoprotein-1 (LRG1) protein has been found in patients with acute appendicitis (AA) and may be useful for diagnosis. This study aimed to investigate whether combined tests including circulating LRG1 mRNA levels improves the early diagnosis of AA.

**Methods:** Between December 2011 and October 2012, a prospective study was conducted on patients aged 18 years or older presenting to the ED with acute abdominal pain (<7 days of symptom onset). Levels of whole blood LRG1 mRNA levels and plasma LRG1 protein taken from these patients within 24 hours of arrival (mean 12.4h) were analyzed. The primary outcome was AA.

**Results:** Eighty-four patients (40 (47.6%) with AA and 44 (52.4%) without AA; mean age 35 years; 41.6% males) were recruited. Median whole blood LRG1 mRNA and plasma LRG1 levels were higher in AA patients than in non-AA. Of 40 AA patients, 13 (32.5%) were diagnosed as complicated AA, and had median LRG1 mRNA level higher than in patients with complicated AA. In ROC analysis of LRG1 mRNA (normalized to GAPDH), LRG1 protein and Alvarado score for discriminating AA and non-AA, the areas under the curve (AUC) were 0.723, 0.742 and 0.805 respectively. The combination of normalized LRG1 mRNA, LRG1 protein and Alvarado score demonstrated the largest AUC (0.845). **Conclusion:** A combination of modified whole blood LRG1 mRNA levels, serum LRG1 protein and Alvarado score at the ED may be useful to diagnose simple and complicated AA from other causes of abdominal pain..

## **Introduction**

Acute appendicitis (AA) is a common life-threatening abdominal emergency, which in 2010, claimed 34800 deaths worldwide [1]. In the USA simple AA accounts for an average of 1.8 days in hospital, perforated AA accounts for 5.2 hospital days [2] and the incidence is increasing [3]. AA is commonly diagnosed based on clinical history, physical examination, simple laboratory tests and imaging [4-7], including the Alvarado score) [8, 9], white cell count or C-reactive protein [10], urinary 5-hydroxyindoleacetic acid (5-HIAA) [11]), and ultrasonography , CT and MRI [12-14] (see Appendix for a summary). The gold standard is CT imaging but this is not always available, and its radiation carries some cancer risk especially in the young. This has led some to search for alternative pathways for accurately diagnosing AA [7].

Circulating biomarkers have the potential to improve the diagnostic accuracy of AA in cases where utilizing CT or MRI would be inappropriate, delayed or unavailable. Leucine-rich  $\alpha$ -2-glycoprotein-1 (LRG1) belongs to the leucine-rich repeat (LRR) family of proteins, many of which are involved in protein-protein interaction, signal transduction, and cell adhesion [15]. The biological function of LRG1 is unclear, but recently studies have demonstrated that LRG1 is expressed during granulocyte differentiation [16] and required for pathological angiogenesis [17]. Using a proteomic approach, LRG1 has recently been identified as a specific marker of AA [18,19]. High expressions of LRG1 protein have been found in the inflamed appendices of patients with AA, and increases in its level have been observed in urine and plasma of children with AA [18-20]. Its concentration correlated with histological severity of appendicitis [19-20]. A new diagnostic marker of specific LRG1 peptides using selected ion monitoring mass spectrometry has been developed, and superior diagnostic performance (AUC: 0.98) has been

demonstrated in the urine of children with AA. However, over 24 hours are required to detect urine LRG1 using this method, which limits its application in emergency settings. A commercial ELISA for LRG1 has also become available, and with shorter processing times than mass spectrometry. However, immunoassay interference resulted in inadequate performance for clinical use [19-20].

Although previous studies have studied changes in protein levels, such levels are dependent upon the upstream expression of LRG1 mRNA which encodes LRG1 protein. Nucleic acids are well regarded as early markers of acute illness and injury [21-36], and we have previously demonstrated a potential clinical role for plasma DNA as a prognostic marker in patients with acute abdominal pain [21].

In adult patients aged over 18 years presenting to an emergency department with acute abdominal pain, what is the add-on diagnostic and risk-stratification value of circulating levels of LRG1 and LRG1 mRNA in patients with AA? We hypothesise that there are significant differences in levels of circulating LRG1 and LRG1 mRNA between patients with AA, and those patients without AA, and that there will be a positive correlation between circulating levels and the severity of appendicitis. Thus the aims of this study were (1) to investigate the diagnostic value of plasma LRG1 and whole blood LRG1 mRNA level in patients with suspected AA, and (2) to elucidate the correlation between whole blood LRG1 mRNA and histological severity of appendicitis, and (3) to investigate early temporal relationships in circulating LRG1 and LRG1 mRNA in patients with acute abdominal pain. This may enable the development of novel protein

and mRNA-based blood markers or combinations to improve the diagnostic accuracy of simple and complicated AA.

## **Materials and methods**

### **Subjects and data collections**

Approval was obtained from Institutional Review Board of the Chinese University of Hong Kong to conduct this prospective study (CREC 2015.710). Written consent was obtained either from the patient or a relative in all cases.

Eligible patients included those aged 18 years and above, presenting to the Emergency Department of the Prince of Wales Hospital, Hong Kong, with abdominal pain of less than 7 days duration. Thirty-one healthy volunteers matched for mean age and sex were recruited. Final diagnosis was determined by the presence or absence of appendicitis on gross and histologic examination.

### **Inclusion and exclusion criteria**

Patients aged 18 or above presenting to the ED with acute abdominal pain of likely surgical cause within 7 days of symptom onset were recruited. Patients were excluded if they were below 18 years of age, lack of consent, pregnant, had external blunt or penetrating trauma (due to an external force associated with a motor vehicle crash, fall or assault etc.), had known non-surgical causes for abdominal pain such as diabetic ketoacidosis, urinary tract infection, gastro-esophageal reflux, or indigestion (dyspepsia), had chronic medical conditions (e.g. inflammatory bowel disease, cancer, sickle cell anemia), or were taking chronic anti-inflammatory medications.

138

139 **Definition**

140 *Acute abdominal pain* was defined as pain occurring within 7 days of onset and in an area  
141 extending below the lower ribs, above the inguinal line and between the mid-axillary lines.

142 *Acute appendicitis (AA)* was defined as the presence of transmural inflammation of appendix or  
143 the presence of pus in the lumen of the appendix [23].

144 *Acute appendicitis like syndrome (AALS)* is usually characterized by clinical symptoms and  
145 physical examination. Clinical symptoms were classified as typical and atypical. Typical  
146 appendicitis usually included abdominal pain beginning in the region of the umbilicus for several  
147 hours, associated with anorexia, nausea or vomiting. The pain was then localized in the right  
148 lower quadrant, where tenderness developed. Atypical appendicitis lacked this typical  
149 progression and may include pain in the right lower quadrant as an initial symptom. Atypical  
150 appendicitis often requires ultrasound scan and/or CT scan to assist diagnosis.

151 The Alvarado score is also used for AA diagnosis [24]. The score has 6 clinical items (based on  
152 clinical symptoms and physical examination) and 2 laboratory measurements with a total of 10  
153 points. A score below 5 is strongly against a diagnosis of appendicitis, while a score of 7 or more  
154 is strongly predictive of acute appendicitis.

155 *Healthy controls* were defined as age- and sex-matched volunteers with no history of recent  
156 acute illness within 3 months, chronic illness, smoking or medication.

157 *Histologic severity of appendicitis* was classified as having no inflammatory features (normal),  
158 foci of neutrophilic infiltration in the wall or mucosa (focal), scattered transmural infiltration  
159 (mild), dense transmural infiltration with tissue distortion (moderate), or dense transmural  
160 infiltration with tissue necrosis or wall perforation (severe) [8].



161

162 **Data collection and measurable parameters**

163 Using a standardized protocol, an English- and Cantonese-speaking research assistant collected  
164 demographic and previous medical data including age, sex, symptom onset time, time between  
165 sample collection and operative care, medical history (e.g. abdominal pain, seizures,  
166 hypertension, diabetes mellitus, ischaemic heart disease, atrial fibrillation, hyperlipidaemia,  
167 smoking etc.) and current medication.

168

169 **Preparation of plasma and RNA extraction**

170 A 10 ml venous blood was taken by standard venipuncture and collected into EDTA-tubes.  
171 Whole blood was collected and stored in TrizolLS (Invitrogen) at -80°C for further analysis.  
172 Plasma was collected after centrifugation and stored at -80°C for further analysis of LRG1  
173 protein level. Total RNA was extracted from 400 ul whole blood and has been previously  
174 described [25].

175

176 **One-step RT-qPCR for LRG1 mRNA and GAPDH mRNA**

177 One-step real-time RT-qPCR was used for measuring the LRG1 mRNA concentrations in the  
178 whole blood RNA samples, based on previously reported methods [25]. The RT-qPCR assay for  
179 LRG1 was developed and optimized. The calibration curve for LRG1 mRNA quantification was  
180 prepared by assaying serial dilutions of HPLC-purified single-stranded synthetic DNA  
181 oligonucleotides (Sigma) specifying a 77-bp LRG1 amplicon, with concentrations ranging from  
182  $1 \times 10^7$  copies to  $1 \times 10^1$  copies. The amplification primers were LRG1F  
183 (5'- ACTGCAACCCGCTTAACA -3') and LRG1R (5'- TCCCAAAGTGCTGGGATTAC -3'),

and the dual-labeled fluorescent probe was LRG1P [5'-(FAM) AATAATCCTGCCTTTGGCCGGGT (TAMRA)- 3', where FAM is 6-carboxyfluorescein and TAMRA is 6-carboxytetramethylrhodamine]. For normalization, reference gene, glyceraldehyde 3-phosphate dehydrogenase (GAPDH) mRNA was measured and the assay for GAPDH has also been well established and described [25]. The concentration of LRG1 and GAPDH mRNA in the whole blood sample of patients and healthy controls were measured in duplicate.

#### **ELISA for LRG1 protein analysis**

Plasma LRG1 was quantified by human LRG assay (IBL, Fujioka, Japan) according to manufacturer's protocols. All samples and reagents were brought to room temperature 30 minutes before use. The level of LRG1 protein in plasma of healthy controls and patients were measured in duplicate.

#### **Outcome measures**

The primary outcome was the presence or absence of AA. The secondary outcome was the severity of appendicitis.

#### **Statistical analysis**

Descriptive statistics and data comparison tests (chi-squared, Fisher exact, Mann-Whitney, Kruskal-Wallis tests), Receiver Operating Characteristic (ROC) analysis, logistic regression, as well as diagnostic strength were carried out using MedCalc12.3 software (MedCalc Software bvba).

## **Results**

### **Baseline characteristics**

Between 14<sup>th</sup> December 2011 and 21<sup>st</sup> October 2012, 84 patients (40 (47.6%) with AA and 44 (52.4%) without AA; median age 35 years; 41.6% males) presenting to the emergency department with acute abdominal pain of less than seven days duration were recruited. The characteristics of the 84 patients are shown in Table 1. Thirty-one healthy controls, matched for mean age and sex were also recruited (median age 32 years, 48.4% male).

### **Whole Blood LRG1 mRNA and plasma LRG1 in AA diagnosis**

Table 2 shows the differentiating features between patients with and patients without AA. Alvarado score and haematemesis were the only discriminating clinical features. Median concentrations of whole blood LRG1 mRNA were significantly different between patients with and patients without AA ( $1.3$  v  $2.2 \times 10^5$  copies/ $\mu$ l blood;  $p=0.0134$ ). Median whole blood LRG mRNA normalized to GAPDH was also significantly different between patients with and patients without AA ( $205$  v  $371$  copies/pg GAPDH;  $p=0.0004$ ). In addition, median plasma LRG1 protein was higher in AA patients than non-AA patients ( $54$  vs  $26$  mg/l;  $p<0.0001$ ).

Figure 1A shows the increase in median LRG1 mRNA concentrations from healthy controls through non-AA, simple AA to complicated AA (Kruskal-Wallis  $p<0.0001$ ). Figure 1B shows the increase in median LRG mRNA concentrations normalized to GAPDH from healthy controls through non-AA, simple AA to complicated AA (Kruskal-Wallis  $P=0.0013$ ). There are significant dose-response increases with increasing severity. Figure 1C shows the increase in median plasma LRG1 concentrations from healthy controls through non-AA, simple AA to

complicated AA (Kruskal-Wallis  $P < 0.0001$ ). There are significant dose-response increases with increasing severity.

Figures 2 shows the receiver operating characteristic (ROC) curves for LRG1 mRNA concentrations, non-normalized and normalized to GAPDH, plasma LRG1 concentrations, and combination of LRG1 mRNA and protein concentrations in patients with non- versus AA. The area under the curve (AUC) of LRG1 mRNA increased from 0.657 to 0.723 after normalization to GAPDH. When compared to LRG1 mRNA, the plasma LRG1 produced a larger AUC (0.742 vs 0.657). The combination of LRG1 mRNA and plasma LRG1 demonstrated larger AUC (0.743) (Table 3). In Table 3, combination of LRG1 mRNA (normalized to GAPDH) and protein demonstrated the larger AUC (0.781). Combination of LRG1 mRNA (normalized to GAPDH), protein and Alvarado score demonstrated the largest AUC (0.845).

Table 3 shows the add on effect and accuracy of whole blood combinations of LRG1 mRNA, LRG1/GAPDH mRNA, plasma LRG1 protein, and Alvarado score for detecting acute appendicitis. The optimal cut off values for LRG1 mRNA and plasma LRG1 in diagnosis of AA were  $2.0 \times 10^5$  copies/ $\mu$ l whole blood (sensitivity: 57.5%; specificity: 72.7%) and 31 mg/l (sensitivity: 77.5%; specificity: 68.2%), respectively. The sensitivity of LRG1 mRNA increased to 95% after being normalized to GAPDH (cut off: 188 copies/pg GAPDH). Combination of LRG1/GAPDH mRNA and protein showed the highest sensitivity, which was 97.5%.

Supplemental Figure 1 shows the receiver operating characteristic (ROC) curves for LRG1 mRNA concentrations, non-normalized and normalized to GAPDH, plasma LRG1

concentrations, and combination of LRG1 mRNA and protein concentrations in patients with simple versus complicated AA. The area under the curve (AUC) of LRG1 mRNA, normalized to GAPDH, protein were 0.694, 0.651 and 0.632 respectively. However, combination of LRG1 mRNA and plasma LRG1 did not improve the diagnostic value (AUC: 0.634) in differentiating complicated AA from simple AA. Diagnostic accuracy of whole blood combinations of LRG1 mRNA, LRG1/GAPDH mRNA, plasma LRG1 protein, and Alvarado score for discriminating simple and complicated AA shows on Supplemental Table 1. In differentiation between simple AA and complicated AA, LRG1 mRNA normalized to GAPDH displayed 100% sensitivity and 33.3% specificity. The sensitivity of LRG1 mRNA and mRNA combined with plasma LRG1 were the same, which was 84.6%, whereas the specificity (63% vs 51.9%) and diagnostic value (AUC: 0.694 vs 0.634) of LRG1 mRNA alone were higher.

Factors including LRG1 mRNA, LRG1/GAPDH, LRG1 protein and Alvarado were subjected to multivariate logistic regression. The logistic regression model for discriminating of acute appendicitis and complicated AA are shown in Table 4. Results show that whole blood LRG1/GAPDH mRNA level and Alvarado score are independent predictors of AA. Whole blood LRG1 mRNA and plasma LRG1 protein predict complicated AA.

## **Discussion**

This study shows that normalized and non-normalized whole blood LRG1 mRNA concentrations measured in patients with acute abdominal pain may be used to differentiate patients with acute appendicitis from other causes of acute abdomen, and that the highest levels occur in patients

with complicated gangrenous appendicitis or appendiceal abscess. These findings raise the possibility of LRG1 mRNA as a diagnostic marker.

The diagnosis of acute appendicitis presents a diagnostic challenge to clinicians even when ultrasound and CT are available. Current laboratory diagnostic markers represent a general acute-phase reactant response that is not specific for acute appendicitis [36,37].

The previous discovery that LRG1 protein was elevated in diseased appendices, and also elevated the blood and urine of children with acute appendicitis, even in the presence of negative imaging, raised the possibility of a novel diagnostic marker [7]. Further studies showed that the commercially available LRG1 ELISA was subject to an immunoassay interference effect [8].

Cellular and circulating proteins are downstream biomarkers in pathological processes and as such may represent a late feature in disease processes. Patients presenting with acute conditions require rapid cellular processes to ‘switch on’ which in turn introduces a delay before biological abnormalities may appear. It is likely that upstream changes in such processes produce molecular changes earlier in acute diseases and may be more useful as early diagnostic and prognostic markers in disease. With this rationale we evaluated changes in LRG mRNA concentrations, the transcript for LRG protein, as a potential marker. The performance of LRG1 mRNA for the detection of AA was moderate but nevertheless showed a promising dose-response effect. In addition, present study demonstrated that combination of whole blood LRG1 normalized to GAPDH, and its plasma protein level and Alvarado score improve the diagnostic

accuracy to acute appendicitis, suggesting that LRG1 would have add on effect on Alvarado score in detecting acute appendicitis.

The use of a blood based LRG1 mRNA to enhance current clinical decision rules may improve the accuracy of diagnosing acute appendicitis. An inexpensive but accurate immunoassay could replace the use of advanced imaging and complex RT-qPCR in patients with equivocal clinical presentations. LRG1 mRNA is likely to be elevated in clinical scenarios involving bacterial infections and so its use should be guided by a reasonable clinical suspicion of appendicitis.

This study is preclinical phase study and limited by the time required for RT-qPCR. Nevertheless, appropriate commercialisation would allow the possibility of a point of care test. The study did had a single gold standard for a single condition – acute appendicitis – but it would be important to evaluate the response of LRG1 mRNA in patients with other causes of abdominal pain. Furthermore, LRG1 mRNA offers add on effect on Alvarado score on detecting acute appendicitis. We had to select out samples for study as we had limited funding but ideally all samples from consecutive patients would be analyzed. We have not performed any comparison with Alvarado score, imaging, or other acute phase proteins so it is unclear whether LRG1 mRNA offers any advantage over these markers.

## **Conclusion**

In conclusion, this study shows that both whole blood LRG1 mRNA and plasma LRG1 concentrations are elevated in patients with acute appendicitis and may have a role as a diagnostic marker. A combination of modified whole blood LRG1 mRNA levels, serum LRG1

321 protein and Alvarado score at the ED may be useful to diagnose AA from other causes of  
322 abdominal pain.

323

324



325 **Table 1** Characteristics of 84 patients presenting to hospital with acute abdominal pain and  
 326 suspected acute appendicitis

327

Characteristics	Value
Age	35[16] 18-66
Sex (male,%)	35 (41.7)
Day from symptom onset (day)	2 [3] 1-7
Time of blood collection from arrival of emergency department (h)	11.8 [10.9] 1.7-23.9
Alvarado	6 [3] 2-10
<b><i>Symptoms (no. of patients, %)</i></b>	
Nausea/vomiting	38 (45.2)
Haematemesis	0 (0)
Diarrhoea	19 (22.6)
Fresh blood in stool	1 (1.2)
Melaena	1 (1.2)
Abdominal distension	40 (47.6)
Poor appetite	53 (63.1)
Heartburn/Indigestion	7 (8.3)
Change bowel habit	25 (29.8)
Jaundice	0 (0)
Dysuria/urinary frequency	12 (14.3)

Syncope/dizziness	29 (34.5)
Fever	32 (38.1)
Virginal discharge	3 (3.6)
<b><i>Pain feature</i></b>	
Tenderness RLQ	84 (100)
Rebound tenderness	35 (41.6)
Migratory RLQ pain	42 (50)
<b><i>Whole blood parameters</i></b>	
LRG1 mRNA level (x10 <sup>5</sup> copies/μl blood)	1.5 [1.8] 0.24-13.01
LRG1 mRNA level (copies/pg GAPDH)	300 [288] 78-3818
GAPDH (pg/μl blood)	567 [390] 6-1856
Plasma LRG1 protein (mg/l)	39 [40] 4-114
<b><i>Type of AA (N = 40)</i></b>	
Simple AA	27 (67.5%)
Complicated AA	13 (32.5%)

---

328 All continuous data are expressed as medians [interquartile range] and the whole range.

329 Numbers may not sum up to 100 because of rounding, multiple factors or absent data

330

331

**Table 2** Comparisons of factors for discriminating acute appendicitis (AA) and non-AA in 84 patients with abdominal pain and suspected AA

Characteristics	Non AA (N=44)	AA (N=40)	<i>p</i> -value
Age	36 [14] 18-58	33 [17] 19-66	0.6412
Sex (male,%)	14 (31.8)	21 (52.5)	0.0764
Day from symptom onset (day)	2 [3] 1-7	2 [0] 1-7	0.5424
Time of blood collection from arrival of emergency department (h)	10.8 [12.2] 1.9-23.9	11.8 [9.5] 1.7-22.9	0.5841
Alvarado	5 [3] 2-8	7 [2] 4-10	<0.0001
<b><i>Symptoms (no. of patients, %)</i></b>			
Nausea/vomiting	20 (45.5)	18 (45)	1.0000
Haematemesis	0 (0)	0 (0)	0.1013
Diarrhoea	16 (36. 4)	3 (7.5)	0.0151
Fresh blood in stool	1 (2.3)	0 (0)	1.0000
Melaena	1 (2.3)	0 (0)	1.0000
Abdominal distension	22 (50)	18 (45)	0.3453
Poor appetite	26 (59.1)	27 (67.5)	0.5000
Heartburn/Indigestion	5 (11.4)	2 (5)	0.6955
Change bowel habit	13 (29.5)	12 (30)	0.3163

Jaundice	0 (0)	0 (0)	0.1013
Dysuria/urinary frequency	9 (20.5)	3 (7.5)	0.3411
Syncope/dizziness	21 (47.7)	8 (20)	0.0894
Fever	13 (29.5)	19 (47.5)	0.1132
Virginal discharge	3 (6.8)	0 (0)	0.2770
<b><i>Pain feature</i></b>			
Tenderness RLQ	44 (100)	40 (100)	0.7434
Rebound tenderness	17 (38.6)	18 (45)	0.6588
Migratory RLQ pain	15 (34.1)	27 (67.5)	0.2446
<b><i>Whole blood parameters</i></b>			
LRG1 mRNA level (x10 <sup>5</sup> copies/μl blood)	1.3 [1.5] 0.3-4.3	2.2 [2.3] 0.2-13	0.0134
LRG1 mRNA level (copies/pg GAPDH)	205 [217] 78-568	371 [232] 149-3818	0.0004
GAPDH (pg/μl blood)	563 [279] 89-1855	591 [615] 6-1503	0.4572
Plasma LRG1 level (mg/l)	26 [38] 4-99	54 [40] 55-114	<0.0001

335

336 All continuous data are expressed as medians [interquartile range] and the whole range.

337 Categorical variables are given as values (percentages).

338 *P* values were derived using the Mann–Whitney test or Fisher exact test as appropriate.

339

340

**Table 3** Add on effect and accuracy (95% CI) of whole blood combinations of LRG1 mRNA, LRG1/GAPDH, plasma LRG1 protein, and Alvarado score for detecting acute appendicitis

	Optimal cut-off	AUC	Improvement in C score*	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)
Alvarado	>5	0.805 (0.714-0.895)	-	92.5 (79.6-98.4)	56.8 (41.0-71.7)	66.1 (52.2-78.2)	89.3 (71.8-97.7)
LRG1 mRNA (x10 <sup>5</sup> copies/ul)	>2.0	0.657 (0.538-0.775)	-	57.5 (40.9-73.0)	72.7 (57.2-85.0)	65.7 (47.8-80.9)	65.3 (50.4-78.3)
LRG1/GAPDH mRNA	>188	0.723 (0.614-0.832)	0.066 (10%)	95.0 (83.1-99.4)	47.7 (32.5-63.3)	62.3 (49.0-74.4)	91.3 (72.0-99.0)
LRG1 protein (mg/l)	>31	0.742 (0.635-0.849)	0.085 (13%)	77.5 (61.6-89.2)	68.2 (52.4-81.4)	68.9 (53.4-81.8)	76.9 (60.7-88.9)
LRG1 mRNA + plasma LRG1 protein	>12.4	0.743 (0.636-0.850)	0.086 (13%)	77.5 (61.5-89.2)	68.2 (61.5-89.2)	68.9 (53.4-81.8)	76.9 (60.7-88.9)
LRG1/GAPDH mRNA + LRG1 protein	>1.7	0.781 (0.663-0.879)	0.124 (19%)	97.5 (86.8-99.9)	50 (34.6-65.4)	63.9 (50.6-75.4)	95.7 (78.1-99.9)
Alvarado + LRG1/GAPDH mRNA +LRG1 protein	>5.6	0.845 (0.764-0.925)	0.188 (29%)	87.5 (73.2-95.8)	65.9 (50.1-79.5)	70 (53.4-58.8)	85.3 (68.9-95.1)

\* from LRG1 mRNA

**Table 4** Logistic regression model of factors for discriminating acute appendicitis (AA) and complicated AA

Factor	Before stepwise		After stepwise	
	Adjusted Odds ratio	P value	Adjusted Odds ratio	P value
	(95%CI)		(95%CI)	
<i>AA vs non AA</i>				
Whole blood LRG1 mRNA *	0.83 (0.23-2.97)	0.7746		
Whole blood LRG1/GAPDH mRNA *	18.76 (3.27-107.62)	0.0010	16.50 (3.10-87.71)	0.0010
Plasma LRG1 protein *	2.70 (0.74-9.81)	0.1322		
Alvarado	2.00 (1.22-3.13)	0.0052	2.22 (1.46-3.37)	0.0002
<i>Complicated AA vs simple AA</i>				
Whole blood LRG1 mRNA *	7.26 (0.78-66.86)	0.0814	9.72 (1.60-59.12)	0.0136
Whole blood LRG1/GAPDH mRNA *	2.27 x 10 <sup>6</sup>	0.9940		
Plasma LRG1 protein *	6.94 (0.85-56.65)	0.0705	5.93 (1.11-31.60)	0.0371
Alvarado	0.81 (0.36-1.85)	0.6220		


\* Optimal cut-off

**Figures**

Figure 1. Box-plot of median level of (A) whole blood LRG1 mRNA (B) LRG1/GAPDH mRNA and (C) plasma LRG1 protein of healthy controls (HC), non acute appendicitis patients (nonAA), patients with simple AA and patients with complicated AA.


Figure 2 Receiver operating characteristic (ROC) curves of whole blood LRG1 mRNA, LRG1/GAPDH mRNA, plasma LRG1 concentrations, combination of LRG1 mRNA and protein concentrations, combination of LRG1/GAPDH mRNA and LRG1 protein concentrations, and combination of Alvarado (Alv), LRG1/GAPDH mRNA, LRG1 protein concentrations in patients with AA versus non-AA.

**Figure 1A**




370

371 **Figure 1B**



372


373 **Figure 1C**



374



**Figure 2**



## **References**

1. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2095-128.
2. Healthcare Cost and Utilization Project (HCUP). Trends in Rates of Perforated Appendix, 2001–2010. <http://hcup-us.ahrq.gov/reports/statbriefs/sb159.jsp> (Accessed September 2014).
3. Buckius MT, McGrath B, Monk J, Grim R, Bell T, Ahuja V. Changing epidemiology of acute appendicitis in the United States: study period 1993-2008. *J Surg Res* 2012;175:185-90.
4. Smink DS, Finkelstein JA, Garcia Pen˜a BM, Shannon MW, Taylor GA, Fishman SJ. Diagnosis of acute appendicitis in children using a clinical practice guideline. *J Pediatr Surg* 2004;39:458.
5. Kharbanda AB, Taylor GA, Fishman SJ, Bachur RG. A clinical decision rule to identify children at low risk for appendicitis. *Pediatrics* 2005;116:709.
6. Mikaelsson C, Arnbjörnsson E. The value of C-reactive protein (CRP) determinations in patients with suspected acute appendicitis. *Ann Chir Gynaecol* 1984;73:281.
7. Polites SF, Mohamed MI, Habermann EB, Homme JL, Anderson JL, Moir CR, et al. A simple algorithm reduces computed tomography use in the diagnosis of appendicitis in children. *Surgery* 2014;156:448-54.
8. Alvarado A. A practical score for the early diagnosis of acute appendicitis. *Ann Emerg Med* 1986;15:557-64.
9. Memon ZA, Irfan S, Fatima K, Iqbal MS, Sami W. Acute appendicitis: Diagnostic accuracy of alvarado scoring system. *Asian J Surg* 2013;36:144-9.
10. Panagiotopoulou IG, Parasharv D, Lin R, Antonowicz S, Wells AD, Bajwa FM, et al. The

410 diagnostic value of white cell count, C-reactive protein and bilirubin in acute appendicitis and its  
411 complications. *Ann R Coll Surg Engl* 2013;95:215-21.

412 11. Hernandez R, Jain A, Rosiere L, Henderson SO. A prospective clinical trial evaluating  
413 urinary 5-hydroxyindoleacetic acid levels in the diagnosis of acute appendicitis. *Am J Emerg*  
414 *Med* 2008; 26:282-6.

415 12. van Randen A, Laméris W, van Es HW, van Heesewijk HP, van Ramshorst B, Ten Hove W,  
416 et al. A comparison of the accuracy of ultrasound and computed tomography in common  
417 diagnoses causing acute abdominal pain. *Eur Radiol* 2011;21:1535-45.

418 13. Medscape. Appendicitis Imaging. <http://emedicine.medscape.com/article/363818-overview>  
419 (Accessed July 2015).

420 14. Old JL, Dusing RW, Yap W, Dirks J. Imaging for suspected appendicitis. *Am Fam Physician*  
421 2005;71:71-8.

422 15. Kobe B, Kajava AV. The leucine-rich repeat as a protein recognition motif. *Curr. Opin Struct*  
423 *Biol* 2001;11:725–32.

424 16. O'Donnell LC, Druhan LJ, Avalos BR. Molecular characterization and expression analysis of  
425 leucine-rich alpha2-glycoprotein, a novel marker of granulocytic differentiation. *J Leukoc Biol*  
426 2002;72:478-85.

427 17. Wang X, Abraham S, McKenzie JA, Jeffs N, Swire M, Tripathi VB, et al. LRG1 promotes  
428 angiogenesis by modulating endothelial TGF- $\beta$  signalling. *Nature* 2013;499:306-11.

429 18. Kentsis A, Lin YY, Kurek K, Calicchio M, Wang YY, Monigatti F, et al. Discovery and  
430 validation of urine markers of acute pediatric appendicitis using high-accuracy mass  
431 spectrometry. *Ann Emerg Med* 2010;55:62-70.

432 19. Kentsis A, Ahmed S, Kurek K, Brennan E, Bradwin G, Steen H, et al. Detection and

433 diagnostic value of urine leucine-rich  $\alpha$ -2-glycoprotein in children with suspected acute  
 434 appendicitis. *Ann Emerg Med* 2012; 60:78-83.

435 20. Kharbanda AB, Rai AJ, Cosme Y, Liu K, Dayan PS. Novel serum and urine markers for  
 436 pediatric appendicitis. *Acad Emerg Med* 2012;19:56-62.

437 21. Rainer TH, Chan AKC, Lee LLY, Yim VWT, Lam NYL, Yeung SW, et al. Use of plasma  
 438 DNA to predict mortality and need for intensive care in patients with abdominal pain. *Clin Chim*  
 439 *Acta* 2008;398:13-7.

440 22. Chan WYR, Graham CA, Rainer TH, Lam NYL, Chiu RWK, Chik KW, et al. Use of a bone  
 441 marrow transplantation model system to demonstrate the haematopoietic origin of plasma S100B  
 442 mRNA. *Clin Chem* 2007;53:1874-6.

443 23. Rainer TH, Wong KS, Lam W, Lam NY, Graham CA, Lo YM. Comparison of plasma beta-  
 444 globin DNA and S-100 protein concentrations in acute stroke. *Clin Chim Acta* 2007;376:190-6.

445 24. Rainer TH, Lam NYL, Man CY, Chiu RWK, Woo KS, Lo YMD. Plasma beta-globin DNA  
 446 as a prognostic marker in chest pain patients. *Clin Chim Acta* 2006;368:110-3.

447 25. Lam NY, Rainer TH, Wong LK, Lam W, Lo YM. Plasma DNA as a prognostic marker for  
 448 stroke patients with negative neuroimaging within the first 24 h of symptom onset. *Resuscitation*  
 449 2006;68:71-8.

450 26. Rainer TH, Lam NYL. Circulating Nucleic Acids and Critical Illness. In: YMD Lo, RWK  
 451 Chiu, PJ Johnson Eds. *Circulating Nucleic Acids in Plasma or Serum II*. *Ann N Y Acad Sci*  
 452 2006;1075:271-7.

453 27. Chiu RW, Rainer TH, Lo YM. Circulating nucleic acid analysis: diagnostic applications for  
 454 acute pathologies. *Acta Neurochi Suppl* 2005;95:471-4.

455 28. Lam NYL, Rainer TH, Chiu RWK, Joynt GM, Lo YMD. Plasma mitochondrial DNA

456 concentration after trauma. Clin Chem 2004;50:213-6.

457 29. Rainer TH, Lam NYL, Tsui NBY, Ng EKO, Chiu RWK, Joynt GM, et al. Effects of filtration  
 458 on glyceraldehyde-3-phosphate dehydrogenase mRNA in the plasma of trauma patients. Clin  
 459 Chem 2004;50:206-8.

460 30. Rainer TH, Wong LKS, Lam W, Yuen E, Lam NYL, Metreweli C, et al. Prognostic use of  
 461 circulating plasma nucleic acid concentrations in patients with acute stroke. Clin Chem  
 462 2003;49:562-9.

463 31. Lam NY, Rainer TH, Chan LY, Joynt GM, Lo YM. Time course of early and late changes in  
 464 plasma DNA in trauma patients. Clin Chem 2003;49:1286-91.

465 32. Rainer TH. Plasma DNA, prediction and post-traumatic complications. Clin Chim Acta  
 466 2001;313:81-5.

467 33. Lo YMD, Rainer TH, Chan LYS, Hjelm NM, Cocks RA. Plasma DNA as a prognostic  
 468 marker in trauma patients. Clin Chem 2000;46:319-23.

469 34. Marudanayagam R, Williams G, Rees B. Review of the pathological results of 2660  
 470 appendicectomy specimens. J Gastroenterol 2006;41:745-9.

471 35. Ng EKO, Tsui NBY, Lam NYL, Chiu RWK, Yu SCH, Wong CSC, et al. Presence of  
 472 filterable and nonfilterable mRNA in the plasma of cancer patients and healthy individuals. Clin  
 473 Chem 2002;48: 1212-17.

474 36. Leung LY, Chan CP, Leung YK, Jiang HL, Abrigo JM, Wang de F, et al. Comparison of  
 475 miR-124-3p and miR-16 for early diagnosis of hemorrhagic and ischemic stroke. Clin Chim Acta  
 476 2014;433:139-44.

477 37. Kao LS, Tsao KJ. Serum markers in acute appendicitis. J Surg Res 2010;164:69-71.

478 38. Wang LT, Prentiss KA, Simon JZ, Doody DP, Ryan DP.. The use of white blood cell count

479 and left shift in the diagnosis of appendicitis in children. *Pediatr Emerg Care* 2007;23:69-76.

480

481 **Appendix 1**

482

	Sensitivity	Specificity	Accuracy
Plain x-ray	50%	50%	
US (Inexperienced)	75%	86%	80%
ED Physicians	80%	84%	
Alvarado <6	94%	80%	90%
US (Experienced)	90%	100%	96%
CT scan	96 – 100%	95 – 97%	96 – 98%
MRI	100%	94%	

483

484

485

486 **Supplemental Table 1** Accuracy (95% CI) of whole blood LRG1 mRNA, whole blood LRG1  
487 /GAPDH mRNA, plasma LRG1 protein, Alvarado, combination of whole blood LRG1 mRNA  
488 and plasma LRG1 protein, combination of whole blood LRG1/GAPDH mRNA and plasma  
489 LRG1 protein Alvarado, and combination of Alvarado, whole blood LRG1/GAPDH mRNA and  
490 plasma LRG1 protein for discriminating complicated acute appendicitis

	Optimal cut-off	AUC	Sensitivity (%)	Specificity (%)	PPV (%)	NPV (%)	PLR (%)	NLR (%)
Alvarado	>6	0.655 (0.482-0.828)	84.6 (54.6-98.1)	40.7 (22.4-61.2)	40.7 (22.4-61.2)	84.6 (54.6-98.1)	1.43 (0.97-2.11)	0.38 (0.1-1.46)
LRG1 mRNA (x10 <sup>5</sup> copies/ul)	>2.1	0.694 (0.518-0.870)	84.6 (54.6-98.1)	63.0 (42.4-80.6)	52.4 (29.8-74.3)	89.5 (66.7-98.7)	2.28 (1.33-3.93)	0.24 (0.07-0.90)
LRG1 /GAPDH mRNA	>261	0.651 (0.477-0.825)	100 (75.3-100)	33.3 (16.5-54.0)	41.9 (24.6-60.9)	100 (66.4-100)	1.50 (1.15-1.96)	0.00
LRG1 protein (mg/l)	>55	0.632 (0.436-0.829)	76.9 (46.2-95.0)	63.0 (42.4-80.6)	50.0 (27.2-72.8)	85.0 (62.1-96.8)	2.08 (1.17-3.69)	0.37 (0.13-1.03)
Whole blood LRG1 mRNA + plasma LRG1 protein	>0.9	0.634 (0.453-0.814)	84.6 (54.6-98.1)	51.9 (31.9-71.3)	45.8 (25.6-67.2)	87.5 (61.7-98.5)	1.76 (1.12-2.77)	0.30 (0.08-1.12)
Whole blood LRG1/GAPDH mRNA + plasma LRG1 protein	>0.7	0.652 (0.453-0.852)	84.6 (54.6-98.1)	59.3 (38.8-77.6)	50 (28.2-71.8)	88.9 (65.3-98.6)	2.08 (1.25-3.46)	0.72 (0.56-0.85)




Alvarado +	>3.1	0.685	92.3	44.4	44.4	92.3	1.66	0.68
Whole blood		(0.521-0.850)	(64.0-	(25.5-64.7)	(25.5-64.7)	(64.0-99.8)	(1.15-2.41)	(0.52-0.82)
LRG1/GAPDH			99.8)					
mRNA + plasma								
LRG1 protein								

---


491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509

**Supplemental Figure 1** Receiver operating characteristic (ROC) curves for whole blood LRG1 mRNA concentrations, whole blood LRG1 mRNA normalized to GAPDH, plasma LRG1 concentrations, combination of LRG1 mRNA and protein concentrations, combination of LRG1 mRNA normalized to GAPDH and protein concentrations, and combination of Alvarado (Alv), LRG1 mRNA normalized to GAPDH, protein concentration in patients simple versus complicated AA




**Supplemental Figure 2** Temporal changes in (A) LRG1 mRNA, (B) LRG1 mRNA normalized to GAPDH, (C) and plasma LRG1 from symptom onset (days) to blood sampling for the AA and non-AA groups. Data is presented as the mean  $\pm$  SEM. Significant difference in LRG1 mRNA or plasma LRG1 was found between AA and non-AA patient with  $P < 0.05$  by using t-test (\*).

Supplemental Figure 2A



Supplemental Figure 2B



Supplemental Figure 2C

