

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/114247/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

McCully, Michelle L., Kouzeli, Ariadni and Moser, Bernhard 2018. Peripheral tissue chemokines: homeostatic control of immune surveillance T cells. *Trends in Immunology* 39 (9) , pp. 734-747. 10.1016/j.it.2018.06.003

Publishers page: <http://dx.doi.org/10.1016/j.it.2018.06.003>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



# Peripheral tissue chemokines: homeostatic control of immune surveillance T cells

Michelle L. McCully<sup>1</sup>, Ariadni Kouzeli<sup>1</sup> and Bernhard Moser<sup>1,\*</sup>

5

*<sup>1</sup>Department of Infection, Immunology & Biochemistry, School of Medicine, Cardiff University, Heath Park, Cardiff CF14 4XN, United Kingdom*

\* Corresponding author: Bernhard Moser ([moserb@cardiff.ac.uk](mailto:moserb@cardiff.ac.uk))

10 Cellular immunity is governed by a complex network of migratory cues that enable appropriate immune cell responses in a timely and spatially controlled fashion. This review focuses on the chemokines and their receptors regulating the steady state localization of immune cells within healthy peripheral tissues. Steady-state immune cell traffic is not well understood but is thought to involve constitutive (homeostatic) chemokines. The recent discovery of tissue-resident memory T (TRM) cells illustrates  
15 our need for understanding how chemokines control immune cell mobilization and/or retention. These studies will be critical to unravel novel pathways for preserving tissue function (aging) and preventing tissue disease (vaccination).

20

## Tissue immune surveillance, the last frontier in immunology

---

For many reasons that include restricted access to healthy human tissue and sample sizes, there exists a formidable gap in our knowledge about how immune surveillance cells are recruited and retained in healthy tissues. This situation is remarkable for two reasons: 1) the significant complexity of immune surveillance cells present in healthy tissues/organ, notably those constantly exposed to environmental microbes and toxins and, 2) the extended duration this system needs to operate, which can exceed 80 years in humans. Peripheral tissue immune surveillance compartments are composed of both cells of the innate and adaptive immune systems and the diversity in their composition reflects the special need for immune surveillance tailored to distinct body locations.

As, thankfully, most individuals are healthy most of the time, the maintenance of tissue immune surveillance is a long-lasting process whose duration vastly exceeds that of rare episodes of infection and inflammation. Obviously, chemokines and their receptors expressed on target cells in collaboration with adhesion receptors and their ligands are intimately involved in both processes. The reader interested in general aspects of the chemokine system and its involvement in host defence is referred to excellent reviews published in recent years [1-3]. The vast majority of the literature deals with the inflammatory side of the chemokine network. Homeostatic immune cell processes are also controlled by chemokines whose expression is constitutive and tissue-specific. Again, our current understanding of the control of immune cell traffic by chemokines in primary (bone marrow, thymus) and secondary lymphoid tissues (LNs, spleen, Peyers Patches) is well advanced [3]. In clear contrast, our understanding of immune cells localization in peripheral tissues and healthy organs is still very limited. Exciting new findings revealed the importance of the tissue environment on shaping local immune cells, as best examined for tissue macrophages [4-6], and some types of tissue-resident memory T (TRM) cells [7-10]. In this article, we will discuss recent advances in our understanding of how chemokines and their receptors help orchestrate T cell compartments in healthy peripheral tissues with the view that this knowledge may help us understand the steady-state immune processes underpinning tissue health.

## Migration control of circulating T cells

Antigen-experienced T cells in peripheral blood include lymph node-homing central memory T ( $T_{CM}$ ) cells and memory precursor T ( $T_{MP}$ ) cells as well as effector memory T ( $T_{EM}$ ) cells that are largely excluded from secondary lymphoid tissues [11-13]. Naïve,  $T_{CM}$  and  $T_{MP}$  cells, express CCR7 as well as CD62L (E-selectin) for binding to GlyCAM-1 on high endothelial cells (HEV), which enables their sequential entrance into tissue-draining lymph nodes in response to CCL19 and CCL21 under steady-state conditions (reviewed in [14]).  $T_{EM}$  cells lack CCR7 but express combinations of receptors for inflammatory chemokines guiding them to peripheral sites during episodes of inflammation. As such,  $T_{EM}$  cell recruitment is transient and ceases to occur with resolution of inflammation.

In line with the underlying paradigm linking immune function with cell migration properties, the chemokine system defines distinct subsets among circulating memory T cell subsets. This is best exemplified by T helper cell subsets whose functional properties (cytokine secretion profile) correlate with the combinatorial cell surface expression of chemokine receptors, as expertly discussed in recent review articles [2, 15]. Under steady-state conditions, circulating  $T_{CM}$  and  $T_{EM}$  cells are largely excluded from healthy peripheral tissues and organs and, as such, do not appear to contribute to the peripheral immune surveillance system.

During immune responses, pathogen-specific T cells are activated in relevant tissue-draining LNs where they expand and differentiate into effector T cells before being released into the blood stream. The type of chemokine receptors present on effector T cells matches their functional demand at the site of infection/inflammation.

Interestingly, early work in mice has demonstrated that effector T cells also gain access, in reduced numbers, to healthy peripheral tissues throughout the body [16, 17]. This process may help to prevent the spreading of infections to secondary sites and, in addition, may contribute to long-lived peripheral memory T cell compartments in unaffected (healthy) tissues. Further, the gene expression signature of  $T_{MP}$  cells, the survivors of effector T cell contraction phases, includes the lymphoid tissue chemokine receptors CXCR4, CCR7 as well as the inflammatory chemokine receptors CXCR3 and CXCR6 that may potentially guide these cells to inflammatory sites [18, 19]. Collectively, memory T cells in peripheral blood do not gain access

under steady-state conditions to healthy tissues, indicating that their study does not reveal the characteristics typically associated with memory T cells present in healthy peripheral tissues.

## 5 **Control of T cell positioning by homeostatic chemokines**

It is becoming increasingly clear that T cells present in healthy (steady-state) peripheral tissues are not simply a mirror image of circulating T cells. This fundamental understanding raises many exciting questions regarding the mechanisms governing their generation as well as their tissue-specific localization and retention. Recent findings emphasise the importance of target tissues themselves in this process [15, 20]. A point in case is tissue macrophages, many of which are derived from “primitive” myeloid progenitor cells generated in the yolk sac or foetal liver (reviewed in [4, 6, 21, 22]). Following recruitment into diverse tissues, these progenitors develop into specialized macrophages, such as microglia, Kupffer cells, peritoneal macrophages or Langerhans cells.

Obviously, (but with the notable exception of murine dendritic epidermal T cells [DETCs], see below), peripheral tissue memory T cells are not derived from embryonic precursors but instead reflect past immune activation events involving effector T cells and, potentially, TMP cells [16, 17]. Tissue-homing properties are imprinted in part by soluble tissue-derived migration cues that effector T cells are exposed to in draining LNs. Again, and in contrast to inflammatory diseases marked by an array of inducible (inflammatory) chemokines, the functions of constitutively produced chemokines within distinct healthy peripheral tissues are not well understood. In this section, we will describe chemokine receptors and ligands that have been associated with controlling the location of peripheral tissue memory T cells (Table 1). For clarity, we will refer to the entire population of T cells present in healthy non-lymphoid tissues at any given time as peripheral immune surveillance T (TPS) cells. TPS cells include TRM cells as well as migratory memory T cells, NKT cells and  $\gamma\delta$  T cells [23]. Please note that numerous chemokines have dual functions, i.e. are present in healthy tissues yet become upregulated in response to inflammatory stimuli.

**CXCR4/ACKR3** – CXCR4 is the most widely expressed chemokine receptor and binds the single chemokine CXCL12 that is constitutively produced in lymphoid organs and peripheral tissues (reviewed in [24]). It also binds gp120 envelope protein of HIV-1 and acts as HIV-1 co-receptor, together with CD4, for CXCR4-tropic (X4) HIV-1 particles (reviewed in [25]). Besides controlling the positioning of bone marrow and thymic progenitor cells during steady-state immune cell development (reviewed in [3, 26]), this chemokine receptor system participates in embryonic tissue development and wound healing (reviewed in [26, 27]). Recent work has revealed that ACKR3, an atypical chemokine receptor, fulfils an essential role in controlling homeostatic immune (and tissue) cell traffic through its ability to internalize and degrade extracellular CXCL12 (reviewed in [28]). Thus, homeostatic immune/tissue cell traffic is finely tuned by two opposing receptors; cell recruitment and/or retention of CXCR4<sup>+</sup> cells by CXCL12 and inhibition of these CXCR4<sup>+</sup> cell migration processes by ACKR3-mediated CXCL12 degradation. A role for CXCR4/ACKR3 in controlling TPS cell traffic has not been reported.

**CXCR6** – CXCR6 is found on activated and memory lymphocytes, including  $\alpha\beta$  T cells,  $\gamma\delta$  T cells, NK T cells, NK cells and B cells and is also present on TPS cells. Similar to CD8<sup>+</sup> TRM cells, CXCR6<sup>+</sup> memory T cells in human and mouse liver express CD69 and the transcription factor Hobit, that causes tissue retention by preventing the expression of the tissue-emigrant receptors CCR7 and S1P1R [29, 30]. TRM-like CXCR6<sup>+</sup> NK cells have also been detected in human liver [31, 32]. The relative contribution of soluble *versus* membrane-bound CXCL16 to the tissue residency of liver T cells and NK/NK T cells is not known although it is tempting to speculate that hepatocyte/cholangiocyte-associated CXCL16 plays a particular role in this process. Recently, CXCR6 was shown to support retention of memory T cells in mouse skin, supporting the notion that membrane-bound CXCL16 on local tissue cells may facilitate adhesive interactions with local CXCR6<sup>+</sup> T cells [33]. Still, the CXCL16-CXCR6 axis in skin may be more relevant to inflammatory diseases since CXCL16 levels are low in healthy skin but become strongly upregulated during inflammation [34, 35].

**CCR6** – CCR6 is broadly expressed on memory T cells, including Th17 cells, Treg cells,  $\gamma\delta$  T cells, and NK T cells, NK cells, B cells, DCs/LCs and neutrophils (reviewed in [24]). CCL20, the only selective ligand for CCR6, was reported to be present in

normal skin and mucosal epithelia as well as perivascular cells, suggesting a role in homeostatic immune cell traffic [36, 37]. However, it needs to be pointed out that CCL20 is prominently upregulated in inflammatory diseases of the skin and intestine, in agreement with the selective expression of CCR6 on distinct T helper cell subsets, especially Th17 cells [15]. IL-17-producing  $\gamma\delta$  T cells ( $\gamma\delta$ T17) also express CCR6 (together with CCR2) and are widely distributed in mouse mucosa/cutaneous tissues where they contribute to IL-17-driven inflammatory diseases [38]. CCR6 governs the retention of long-term memory  $\gamma\delta$  T cells in healthy tissues as well as LNs [39-41], linking this chemokine receptor expression with tissue immune surveillance by a special subset ( $V\gamma 4^+$   $\gamma\delta$ T17) of  $\gamma\delta$  T cells. It is not clear whether CCR6 controls the same process in human mucosa/cutaneous tissues where  $\gamma\delta$  T cells are far less numerous than in mice.

**CCR9** – CCR9 is selective for only one chemokine, CCL25, whose expression is prominent in intestinal epithelia already in the steady-state but is further enhanced under inflammatory conditions (reviewed in [42]). CCR9 is abundant on small intestinal T cells, including intraepithelial and lamina propria  $\alpha\beta$  and  $\gamma\delta$  T cells, as well as IgA<sup>+</sup> plasmablasts, plasmacytoid DCs and intestinal/hepatic macrophages, which is in clear contrast to peripheral blood where CCR9<sup>+</sup> leukocytes are rare. The CCL25-CCR9 axis contributes to the recruitment of  $\alpha 4\beta 7^+$   $\alpha\beta$  and  $\gamma\delta$  T cells to the small intestine, although T cell recruitment to this region does not solely depend on this chemokine system (reviewed in [43]). Despite the importance of the vast T cell compartment exceeding  $10^{11}$  cells in the human intestinal mucosa, it is currently not clear how (if at all) CCR9 contributes to the immune surveillance traffic at this particular location.

**CCR10** – CCR10 plays a dual role in the control of mucosa/cutaneous lymphocyte traffic where it is found on subsets of  $\alpha\beta$  and  $\gamma\delta$  T cells, Treg cells, innate lymphoid cells (ILCs) and IgA<sup>+</sup> plasmablasts [44, 45]. The tissue selectivity is provided by the two chemokines, CCL27 and CCL28 that specifically bind to CCR10. CCL27 is produced in the skin by basal keratinocytes and perivascular cells in the dermis, is highly elevated in inflammatory skin diseases but is absent in intestinal tissues [46-48]. In clear contrast, CCL28 is produced by mucosal epithelial cells in the colon and secretory organs (salivary and mammary glands) where it orchestrated IgA<sup>+</sup>

plasmablasts traffic [45, 49-52]. In addition, CCL28 was also demonstrated to control memory T cells traffic in the nasal mucosa where it is constitutively expressed by vascular endothelial cells [53]. Intriguingly, in this study CCL28 was proposed to act *via* CCR3, the second receptor for CCL28.

- 5 The CCL27-CCR10 axis plays a redundant role in the recruitment of effector T cells to the inflamed skin [54]. Still, constitutive expression of CCL27 in human and mouse skin also suggests an involvement in the traffic control of TPS cells under steady-state conditions. In addition, resident CCR10<sup>+</sup> CD8<sup>+</sup> T cells in murine skin were shown to enhance the survival of local Treg cells and, *vice versa*, local CCR10<sup>+</sup> Treg  
10 cells affected the number of local conventional  $\alpha\beta$  T cells in the absence of skin inflammation [55, 56]. A similar role has been attributed to ILCs, a major cellular constituent of murine TPS cells in body-lining tissues, whose localization to healthy skin was also CCR10 dependent [57]. However, despite CCL27 expression, CCR10<sup>+</sup> cells are absent in healthy human skin, denoting a species difference regarding the  
15 role played by the CCL27-CCR10 axis in controlling the local TPS cell compartment.

**CX3CR1** –Similar to CXCL16 (see above), CX3CL1, the ligand for CX3CR1, is a chemoattractant in its soluble form whereas membrane-bound CX3CL1 mediates adhesion of CX3CR1<sup>+</sup> cells (reviewed in [24]). Besides distinct tissue macrophages, such as microglia, CX3CR1 is present on NK cells and T cells as well as monocytes,  
20 where its variation of expression distinguishes two major monocyte subsets. A function in tissue homeostasis is inferred from CX3CR1<sup>+</sup> DCs during microbial surveillance in intestinal tissue and from the effect of CX3CR1<sup>+</sup> microglia on neuronal plasticity and synaptic pruning [58, 59]. Of note, the CX3CL1-CX3CR1 axis has recently been implicated in tissue immune homeostasis. Similar to blood monocytes,  
25 the level of CX3CR1 expression in murine CD8<sup>+</sup> T cells defines three subsets, TCM and TEM cells that either lack or express high level of CX3CR1, and a memory T cell subset with intermediate CX3CR1 expression [60]. CX3CR1<sup>int</sup> T cells are long-lived, recirculate through peripheral tissues (thus differing from TRM cells, which are sessile) and give rise to effector and TEM cells upon reinfection. Conversely, another  
30 group reported that LNs contain a sessile memory T cell subset characterized by CX3CR1 (and CD62L) expression with proposed pathogen surveillance function [61]. It is not clear at present whether the CX3CL1-CX3CR1 axis plays any role in the generation and/or maintenance of these novel memory T cells subsets.



**Emerging chemokine systems with a role in tissue homeostasis** – GPR15, a G-protein coupled receptor induced on skin and colon immune cells by local TGF- $\beta$ 1 [62], was recently shown to facilitate the localization of fetal V $\gamma$ 3<sup>+</sup> (alternatively called V $\gamma$ 5<sup>+</sup>)  $\gamma\delta$  T cells in the embryonic epidermis of mice [63]. Fetal V $\gamma$ 3<sup>+</sup>  $\gamma\delta$  T cells are the precursors of dendritic epidermal T cells (DETCs) whose stress surveillance function in mouse skin is well described (reviewed in [64]). The  $\gamma\delta$  T cells equivalent to mouse DETCs do not exist in humans. GPR15 is also involved in the intestinal immune homeostasis in mice (but not humans) *via* its regulation of the local Treg cell compartment [65, 66]. The two identified ligands, a thrombomodulin-derived protein fragment [67] and *C10ORF99*-encoded protein GPR15L [68], do not mediate chemotaxis of GPR15<sup>+</sup> cells, suggesting that a migration-unrelated mechanism underlies the GPR15 controlled processes in tissue immune homeostasis.

CXCL14 and CXCL17 are highly expressed at the steady-state in a broad range of epithelial tissues and organs, suggesting important contributions to tissue immune homeostasis. CXCL14 is broadly expressed in body-lining tissues and internal organs (reviewed in [69]). A large body of literature describes the pro- or anti-tumour effect of CXCL14, which was linked to its function in angiogenesis and tumour immune surveillance. Similar to CXCL12, CXCL14 is involved in tissue development and glucose metabolism that may explain, in part, the striking survival defects observed in CXCL14-KO mice [70-72]. It is a chemoattractant for myeloid cells, notably blood monocytes, and also synergizes with CXCL12 in CXCR4-mediated chemokine responses [73]. Slow progress in CXCL14 research is largely due to the fact that its receptor has remained elusive.

CXCL17 is prominently expressed in the tongue, airways (trachea, lung) and stomach as well as in psoriatic (but not healthy) skin and was shown to chemoattract myeloid cells (macrophages, monocyte-derived suppressor cells) [74, 75]. CXCL17 was recently reported to be a selective chemokine ligand for GPR35 (subsequently called CXCR8), although others could not confirm this finding [76, 77]. GPR35 has long been known for its selectivity for the tryptophan derivative kynurenic acid (and related compounds), linking endogenous and/or microbial amino acid metabolism with intestinal disease (reviewed in [78]). The correlation between homeostatic CXCL17 production and GPR35-linked intestinal disease requires further investigations.

## CCR8 designates tissue-resident memory T cell in healthy human skin

A brief history of CCR8-related chemokine research is summarized in Box 1. The following discussion is restricted to human  $\alpha\beta$  T cells, although similar mechanisms may underlie the generation/maintenance of other numerically minor CCR8<sup>+</sup> lymphocyte subsets ( $\gamma\delta$  T cells, NKT cells, NK cells). CCR8 is found on approximately half of all human skin CD4<sup>+</sup> and CD8<sup>+</sup> T cells [79]. Considering the enormous number of T cells present in human skin [80], we estimate that  $>10^{10}$  cells express CCR8, which is equivalent to all (naïve and memory) T cells present in peripheral blood. In clear contrast, CCR8 expression is rare (approx. 5%) among peripheral blood T cells and is absent on any other type of blood leukocytes, such as  $\gamma\delta$  T cells, B cells, NK cells, monocytes and neutrophils. All peripheral blood CCR8<sup>+</sup> T cells are memory cells belonging to either TCM or TEM subsets and also co-express cutaneous lymphocyte-associated antigen (CLA), indicating that they may be on their way to the skin.

The presence of two memory T cells subsets of equal size in healthy human skin that are distinguished by CCR8 expression raises the question as of how they relate to each other. The hallmark of TRM cells are receptors associated with their tissue residency (CD69<sup>+</sup>, CD103<sup>+/-</sup>) and transcription factors (Hobit<sup>hi</sup>, Blimp<sup>hi</sup>, Eomes<sup>low</sup>, T-bet<sup>low</sup>, KLF2<sup>low</sup>) controlling their tissue location and residency as well as their long term survival [7-10, 29]. Of interest, Runx3 was recently reported to be a master regulator of the transcription factor landscape underpinning the generation of TRM cells [81]. Human skin CCR8<sup>+</sup> T cells bear many of the hallmarks ascribed to mouse TRM cells [7-10], including CD69/CD103 expression, proliferation responses to steady-state growth factors IL-7 and IL-15 and lower expression of Eomes and T-bet. In contrast, skin T cells lacking CCR8 were more variable in phenotypic and functional markers, expressed inhibitory receptors, including PD-1, showed poor proliferative responses *ex vivo* and had substantially higher levels of transcripts for effector molecules (perforin, granzymes, CXCR3, etc.). Underscoring these striking differences, the two major skin populations distinguished by CCR8 also recognized different antigens as demonstrated by TCR V $\beta$  clonotype analyses [79].

How human skin CCR8<sup>+</sup> T cells relate to their CCR8<sup>+</sup> counterparts in blood is not known. Similar to the recently described “migratory” memory T cells in human and mouse skin [82, 83] and “circulating” memory T cells in peripheral blood of mice [60], it is possible that the human blood CCR8<sup>+</sup> compartment contains a subset of

5 recirculating memory cells with homing preferences for healthy skin. Peripheral Treg cells seed mouse skin very early in life [84]. Although a recent report indicated human skin Treg cells were non-migratory [85], studies in mice have demonstrated their continuous emigration from the skin in the steady state [86]. Human skin Treg cells uniformly express CCR8 and, likewise, human peripheral blood contains CCR8<sup>+</sup>  
 10 Treg cells that co-express the skin-homing marker CLA and account for between one third and one half of all circulating FoxP3<sup>+</sup> Treg cells [79, 87]. Interestingly, human CCR8<sup>+</sup> Treg cells have also been found in skin-unrelated tumours, such as breast, lung and colorectal tumours, and their presence has been associated with a poorer prognosis [88, 89]. One group reported that triggering of CCR8 signalling enhanced  
 15 the suppressive function of Treg cells [90]; however, a similar pro-Treg cell effect was not found by another group [88]. Since alternatively activated M2 macrophages frequently found in tumour tissue and Treg cells themselves produce CCL1, it is possible that the CCR8/CCL1 axis has been co-opted by the tumour to promote local immune suppression [91, 92]. How tumour and skin Treg cells relate to each other is  
 20 not known but CCR8 expression may indicate shared localisation properties.

Collectively, we propose that CCR8 may serve one or several roles in the localisation of CCR8<sup>+</sup> TRM cells in human skin as discussed in Figure 1. Our understanding of CCR8 expression on mouse lymphocytes is rudimentary. In support of our model, CCR8 mRNA was detected late during anti-viral immune responses in mouse skin  
 25 CD8<sup>+</sup> TRM cells [93]. Also, lack of CCR8 did not prevent T cell recruitment to skin infection, suggesting that in mice CCR8 may be more important in steady-state immune surveillance processes [33]. A role for CCR8 in effector T cell traffic was also recently reported in a mouse model of atopic dermatitis [94].

### 30 **The critical role of the tissue microenvironment in TPS generation**

It is becoming increasingly clear that the tissue itself regulates the localization and residency of immune cells in peripheral tissues. For instance, homing to distinct

tissue sites, especially those with restricted access (i.e. skin, intestinal tract) requires specific migration cues to be imprinted in tissue draining lymph nodes and not elsewhere (reviewed in [95, 96]). Each tissue produces a characteristic set of soluble factors under steady-state conditions that are necessary to induce changes in gene expression. Human skin epidermis releases factors that induce CCR8 (together with CLA) in TCR-activated (but not resting) naïve  $\alpha\beta$  T cells during *ex vivo* culture [97]; and these factors were subsequently determined to be keratinocyte-derived PGE<sub>2</sub> and 1,25-dihydroxyvitamin D<sub>3</sub>, the active metabolite of vitamin D [98] (Figure 2). In addition, vitamin D also plays a role in the generation of CCR10<sup>+</sup> T cells for migration to epidermal layers, but instead of PGE<sub>2</sub> this requires the presence of IL-12 [99]. Likewise, in the gut, the local environment licenses resident CD103<sup>+</sup> dendritic cells with the ability to produce the vitamin A metabolite all-trans retinoic acid (atRA; reviewed in [100]). atRA in turn imprints a gut-tropism, defined by expression of CCR9 and integrin  $\alpha 4\beta 7$ , in responding T cells [101], IgA-secreting B cells [102] as well as ILCs [103].

Interestingly, vitamin metabolites have also been shown to affect the transcriptional programme in tissue macrophages. In the peritoneum, locally produced atRA was shown to regulate the anatomical localization of peritoneal macrophages by inducing of GATA6 and downstream transcriptional changes [104, 105]. Whether vitamin A (in the gut) or vitamin D (in the skin) induce similar transcriptional programmes in T<sub>H</sub>17 cells has not yet been explored. Still, it is worth noting that Treg cell generation is influenced by the presence of both vitamin A and D metabolites acting either directly [106] or *via* DCs [107, 108].

Vitamin A and D induce signal transduction by binding to specific nuclear receptors, RAR (retinoic acid receptor) and VDR (vitamin D receptor), respectively, which form heterodimers with the RXR (retinoic X receptor) nuclear receptor family members. In adipose tissue, the differentiation of Treg cells requires activation of PPAR $\gamma$  (peroxisome proliferator-activated receptor gamma), a nuclear receptor that also pairs with RXR to induce transcription. More importantly, PPAR $\gamma$  signalling was recently shown to increase survival of mouse and human skin CD8<sup>+</sup> T<sub>H</sub>1 cells, linking nuclear receptor signalling with homeostasis of T<sub>H</sub>1 cells [109]. Similar to vitamins A and D, the ligands for PPAR $\gamma$  are present in the tissues and, together with other local mediators like PGE<sub>2</sub>, IL-15 or TGF $\beta$ , may significantly alter the phenotype and

function of T cells that enter the tissue [98, 110, 111]. The relative contribution of each nuclear receptor will likely depend on the site of T cell differentiation as availability of the functional vitamin metabolites (i.e. atRA in gut, 1,25-dihydroxyvitamin D3 in skin) varies across tissue sites. Together, *via* sensing of diverse tissue metabolites, these data suggest that RXR-containing heterodimeric nuclear receptors play an important role in the localization and functional specialisation of TPS cells and other types of tissue immune surveillance cells.

### Concluding Remarks

Tissue health is the prevailing physiological state of our body whereas tissue disease is commonly short-lived (e.g. acute infections) and only rarely life-threatening and/or unremitting (chronic inflammation, cancer). Tissue health is governed by homeostatic chemokines with tissue-specific expression profiles that retain immune surveillance cells, including TRM cells and migratory TPS cells, which express the corresponding chemokine receptors and adhesion molecules. The peripheral tissue immune cell compartments are vast in terms of complexity and cellularity as best exemplified by the superficial skin or mucosal epithelia in humans. At these locations, homeostatic chemokines can function in different ways as discussed in Figure 1 for CCR8 in human skin TRM cells. In essence, tissue-specific homeostatic chemokines could act on effector T cells by retaining these cells within the tissue following resolution of the infection or they could specifically act on memory precursor T cells that have survived the retraction phase of an immune response. Alternatively, homeostatic chemokines could contribute to the longevity of TPS cells by controlling their sequestration to distinct niches within non-lymphoid tissues rich in survival factors, as reported for virus-specific TRM cells in the genital mucosa [112]. Finally, since memory T cells continuously survey the local tissue environment [113, 114], homeostatic chemokines could enable co-localization of TPS cells with local antigen-presenting cells (APCs) as a means of mounting immediate responses to pathogens that local TPS cells have been trained to recognize.

Many questions remain to be addressed (see Outstanding Questions) as we are only beginning to unravel which chemokine networks are at play in distinct tissues and how they affect the resulting local immune compartment under steady-state

conditions. For instance, we do know that TRM persist for many years/decades in tissues, but we don't know whether local tissue-specific homeostatic chemokines are important for their long-term localization and/or function. We also need to determine whether there is any redundancy in these roles. For example, CXCR6 is expressed  
5 by TRM in many tissues, excluding a role in tissue-specific immune cell localization, yet it may contribute to immune surveillance of micro-infections in response to unregulated CCL20. In fact, some peripheral tissue chemokines fulfill a dual function: homeostatic immune cell localization under steady-state conditions and control of immune cell traffic during brief episodes of local infections.

- 10 It is clear that distinct tissue factors, especially metabolites such as vitamins and lipid derivatives, act to imprint tissue residency. In human skin,  $\alpha\beta$  T cells are not the only lymphocyte subset expressing CCR8. In fact, minor fractions of TPS cells constituting skin-resident  $\gamma\delta$  T cells, NKT cells and ILCs, also express CCR8. This finding illustrates that factors governing tissue residence do not discriminate between  
15 different subsets of TPS cells within the same tissue.

- Finally, as for inflammatory chemokines, homeostatic chemokine networks may be targeted for vaccine purposes to promote long-lived, tissue specific protection. Not only could the presence of receptors for peripheral tissue chemokines on TRM cells (such as CCR8 on cutaneous TRM cells) serve as a marker for successful  
20 vaccination regimens, it may be possible to promote T cell recruitment and survival by modulating their tissue expression. Clearly, future investigations of peripheral tissue chemokines will help us to identify novel pathways for interventions that control cellular processes underpinning tissue health and, in effect, long life.

## Box 1. Summary of CCR8-related research

---

- 5      ● *History of CCL1 and CCR8* – CCL1, the first identified CC chemokine, was discovered in the late 1980's in humans [115] and mice [116] and was originally reported to be a chemoattractant for blood monocytes [117]. CCR8, the receptor for CCL1, was discovered in 1997 and, in agreement with the original description of CCL1 activity [117], CCR8 transcripts were found in human blood monocytes [118]. The second group who cloned CCR8 was unable to detect CCR8 mRNA in human blood monocytes and did not find monocyte migration in response to CCL1 [119]. Identification of murine CCR8 was reported in 1998 [120].
- 10      ● *Additional ligands for CCR8* – Murine (but not human) CCL8 controls inflammatory T cell recruitment in mouse model of atopic dermatitis [94]; human CCL18 was reported to be a lower potency ligand for human CCR8 with overproduction in inflammatory skin lesions [121].
- 15      ● *CCR8-KO mice* – Early reports were inconclusive regarding the role of CCR8 in inflammation [122-124] but more recent data revealed that CCR8 modulates Treg cell and myeloid DC function in allergic airway disease [125] as well as Th2 cell recruitment in a mouse model of atopic dermatitis [94].
- 20      ● *DC generation/localization* – Role of CCR8 in DC/LC function was inferred from studies showing a positive effect on DC differentiation during *in vitro* reverse transendothelial migration of human monocytes [126] yet showing a inhibitory (and possibly indirect) effect on skin DC emigration and LC repopulation in mouse models of skin inflammation and stress [127, 128].
- 25      ● *T cell traffic* – Based on earlier work and studies with CCR8-KO mice, Th2 cells, notably IL-5 producing Th2 and Treg cells are targets for human CCL1 (and mouse CCL8) [87, 94, 129-137]. In addition, based on novel CCR8-specific antibodies, it is now known that human CCR8<sup>+</sup> T cells include 50% of all TPS cells present in healthy human skin, including TRM cells, (V $\delta$ 2<sup>neg</sup>)  $\gamma\delta$  T cells, NKT cells and NK cells [79, 97, 138, 139]. In human blood, CCR8 is found on

30      minor (5%) subsets of TCM and TEM cells and very few (V $\delta$ 2<sup>neg</sup>)  $\gamma\delta$  T cells [79,

139, 140]. In addition, CCR8 is present on all skin Treg cells and on all blood Treg cells that also co-express CLA.

- *Expression of CCL1* – At steady-state, CCL1 is produced by human skin LCs and dermal perivascular cells [134, 139]. Under stimulatory conditions, and in agreement with original findings [116], CCL1 is a product of activated T cells, including human TRM cells [139] and Treg cells [90, 91] in addition to activated mast cells [135, 141, 142], macrophages [92, 134, 143, 144] and vascular/lymphatic endothelial cells [126, 145].
- *Viral CCR8 ligands* – Molluscum contagiosum virus-encoded MC148 [146] and Herpes virus HHV8-encoded vMIP-II [147] are two CCR8 antagonists. Of note, molluscum contagiosum virus is a human skin-tropic poxvirus and its early secreted MC148 protein is a highly selective CCR8, conceptually linking viral infection with evasion of human skin CCR8<sup>+</sup> TRM and other skin-specific immune surveillance cells.



(Figure legends)

**Figure 1. Hypothesis: CCR8<sup>+</sup> TRM cells contribute to tissue health by prevent excessive tissue inflammation.** *Primary infection:* Microbes that have penetrated

the skin are processed by local APCs, including epidermal LCs and dermal DCs. In  
 5 case of first-time exposure, microbe-specific memory T cells are not present at the  
 site of pathogen entry, thus enabling the initiation of a local inflammatory cascade  
 through the interaction of microbial antigens with TLRs on macrophages and tissue  
 cells. This in turn leads to rapid secretion of inflammatory chemokines as well as  
 maturation of LN-homing of CCR7<sup>+</sup> DCs/LCs. CCR8 is expressed on T cells within  
 10 the draining LNs as a consequence of TCR signalling induced during interaction with  
 microbial antigen-presenting DCs/LCs in combination with skin-derived tissue factors  
 (vitamin D, PGE<sub>2</sub>). Besides CCR8, newly generated effector T cells express receptor  
 for inflammatory chemokines (CXCR3, CCR2, CCR5, etc.) as well as skin-homing  
 adhesion molecules (CLA, integrins) guiding the effector T cells selectively to the site  
 15 of skin inflammation. CCR8 on effector T cells is probably not necessary in the  
 recruitment phase of effector T cells. *Tissue recovery:* Following the resolution of  
 infection and inflammation, surviving memory precursor T (TMP) cells give rise to long-  
 lived CCR8<sup>+</sup> TRM cells that co-localize with cells constitutively producing CCL1 in the  
 epidermis or in perivascular niches of the dermis. Some TMP cells may acquire  
 20 CCR7, which enables their tissue exit and potential survival as circulating TCM cells.  
 Whether local tissue factors (vitamin D, PGE<sub>2</sub>) are sufficient for maintaining CCR8  
 expression on skin TRM cells or whether TRM cells need to recirculate through  
 draining LNs in order to maintain CCR8 expression is not clear. *Immune surveillance:*  
 CCR8<sup>+</sup> TRM cells survey healthy skin by continuously interacting with local APCs  
 25 (epidermal LCs, dermal DCs) in search of their cognate antigen. In case of re-  
 infection, microbe-specific TRM cells become activated leading to two separate  
 events. First, activated CCR8<sup>+</sup> TRM cells immediately release CCL1 enabling an  
 amplification of the anti-microbial response by mobilizing additional CCR8<sup>+</sup> TRM cells.  
 Second, activated CCR8<sup>+</sup> TRM cells turn into effector cells capable of neutralizing the  
 30 infectious particles. This “memory” response quickly deals with micro-infections  
 circumventing the need for an inflammatory response involving a new wave of  
 circulating effector T cells. Such quick action of local TRM cells may indeed hold the  
 key for maintaining tissue health and longevity. Finally, since all skin Treg cells

express CCR8, auto-reactive effector T cells may be inhibited by Treg cells following their mobilization in response to CCR8<sup>+</sup> TRM cell-secreted CCL1.

**Figure 2. Tissue environments control local memory T cell compartments.**

- 5 Expression of the skin-homing markers CCR8 and cutaneous lymphocyte-associated antigen (CLA) on memory T cells is controlled by TCR stimulation in the presence of the CCR8-inducing factors 1,25(OH)<sub>2</sub>D<sub>3</sub> (calcitriol) and prostaglandin E<sub>2</sub> (PGE<sub>2</sub>). In addition to dietary supplementation, sunshine (UVB irradiation) in the epidermis of sun-exposed skin converts cholesterol into vitamin D<sub>3</sub>, which is further processed by
- 10 intracellular hydroxylases into bioactive 1,25(OH)<sub>2</sub>D<sub>3</sub>. PGE<sub>2</sub> is continuously formed in the upper layers of the epidermis where end-stage keratinocytes undergo apoptosis. 1,25(OH)<sub>2</sub>D<sub>3</sub> and PGE<sub>2</sub> act on T cells during their engagement with DCs either in the skin or in skin-draining lymph nodes by means of engaging the nuclear receptor VDR-RXR and cell surface receptor EP4, respectively. The gut-homing markers
- 15 CCR9 and integrin  $\alpha$ 4 $\beta$ 7 on intestinal memory T cells are controlled by food-derived vitamin A metabolites. The subset of CD103<sup>+</sup> DCs in the intestine is capable of converting the inactive metabolite retinol through intracellular reduction and oxidation reactions into bioactive all-trans retinoic acid (atRA) that acts on the nuclear receptor RAR-RXR. Induction of CCR9 and  $\alpha$ 4 $\beta$ 7 expression occurs during engagement of T
- 20 cells with CD103<sup>+</sup> DCs in the mesenteric lymph nodes, and the development of T cell subsets such as TH17 cells and Tregs is further controlled by additional cytokines (IL-6, TGF- $\beta$ , etc.)

## Glossary

**Naive T cells (T<sub>N</sub> cells):** Antigen-inexperienced circulating CD4<sup>+</sup> and CD8<sup>+</sup> αβ T cells that are excluded from healthy and inflamed peripheral tissues. Expression of CCR7 and CD62L enables naïve T cells to continuously enter secondary lymphoid tissues (e.g LNs) *via* high-endothelial venules. In the T-zone of LNs naïve T cells make multiple contacts with DCs, screen the repertoire of presented antigenic peptide-MHC complexes and, in case of cognate interaction, proliferate and develop into effector T cells.

**Effector T cells (T<sub>E</sub> cells):** Effector T cells are produced in response to microbial challenges, are quickly recruited to sites of inflammation in response to locally produced inflammatory cytokines and provide immediate defensive functions during local re-challenge. Effector T cells are short lived (KLRG1<sup>+</sup>, CD127<sup>-</sup>) and disappear during the immune response contraction phase.

**Helper T cells (T<sub>H</sub> cells):** Circulating effector or memory CD4<sup>+</sup> αβ T cells distinguished by the profile of cytokines they secrete upon re-stimulation. TH subsets include TH1, TH2, TH9, TH17, TH22, etc. and express characteristic combinations of chemokine receptors that enable their selective recruitment to inflammatory sites where the distinct cytokines are needed.

**Follicular B helper T cells (T<sub>FH</sub> cells):** Circulating memory CD4<sup>+</sup> αβ T cells specialised in providing B cell help to T cell-dependent antigens during the germinal centre reaction in the B cell compartments of LNs. T<sub>FH</sub> cells express CXCR5, which guides them to the B cells at the follicular-medullary junctions where CXCL13 is being produced.

**Regulatory T cells (T<sub>reg</sub> cells):** CD4<sup>+</sup> αβ T cells specialised in dampening T cell responses through engagement of inhibitory receptors (e.g. CTLA-4), secretion of anti-inflammatory stimuli (IL-10, IL-35, nucleotides) or depletion (CD25) of IL-2. Treg cells are distinguished by the transcription factor FoxP3 and high levels of CD25, which are generally absent in resting conventional T cells.

**Central memory T cells (T<sub>CM</sub> cells):** Circulating memory CD4<sup>+</sup> and CD8<sup>+</sup> TCM cells continuously enter secondary lymphoid tissues (e.g LNs) *via* high-endothelial venules

and survey local dendritic cells for the presence of cognate antigen. TCM cells express CCR7 and CD62L enabling their entry and localisation within LNs.

**Effector memory T cells (TEM cells):** Circulating memory CD4<sup>+</sup> and CD8<sup>+</sup> TEM cells that do not express CCR7 and CD62L and, therefore, are excluded from secondary lymphoid tissues. TEM cells are also excluded from healthy peripheral tissues but express variable combinations of receptors for inflammatory chemokines guiding them to site of inflammation where the corresponding chemokines are produced.

**Memory precursor T cells (TMP cells):** Alternatively called stem-cell memory T (TSCM) cells and memory precursor effector T (TMPEC) cells). TMP cells share a KLRG1<sup>low</sup>T-bet<sup>low</sup>IL-7Ra<sup>hi</sup> phenotype, survive contraction phase of adaptive immune responses, and give rise to long-lived circulating and possibly tissue-resident memory T cells.

**Peripheral tissue immune surveillance T cells (TPS cells):** The combination of CD3<sup>+</sup> T cells present in healthy peripheral tissues, including tissue resident memory and circulating  $\alpha\beta$  T cells,  $\gamma\delta$  T cells and NK T cells.

**Resident memory T cells (TRM cells):** CD4<sup>+</sup> and CD8<sup>+</sup> T cells with long-term residence in healthy tissues forming part of the local immune surveillance system. TRM cells often express CD69  $\pm$  CD103, are generated during ongoing adaptive immune responses and are in disequilibrium with circulating memory T cells.

## (References)

1. Zlotnik, A. and Yoshie, O. (2012) The Chemokine Superfamily Revisited. *Immunity* 36 (5), 705-716.
2. Griffith, J.W. et al. (2014) Chemokines and Chemokine Receptors: Positioning Cells for Host Defense and Immunity. *Annual Review of Immunology*, Vol 32 32, 659-702.
- 5 3. Schulz, O. et al. (2016) Chemokines and Chemokine Receptors in Lymphoid Tissue Dynamics. *Annu Rev Immunol* 34, 203-42.
4. Davies, L.C. et al. (2013) Tissue-resident macrophages. *Nature Immunology* 14 (10), 986-995.
5. Epelman, S. et al. (2014) Origin and Functions of Tissue Macrophages. *Immunity* 41 (1), 21-35.
6. Amit, I. et al. (2016) The role of the local environment and epigenetics in shaping macrophage identity and their effect on tissue homeostasis. *Nat Immunol* 17 (1), 18-25.
- 10 7. Mueller, S.N. et al. (2013) Memory T cell subsets, migration patterns, and tissue residence. *Annu Rev Immunol* 31, 137-61.
8. Schenkel, J.M. and Masopust, D. (2014) Tissue-resident memory T cells. *Immunity* 41 (6), 886-97.
9. Park, C.O. and Kupper, T.S. (2015) The emerging role of resident memory T cells in protective immunity and inflammatory disease. *Nat Med* 21 (7), 688-97.
- 15 10. Mueller, S.N. and Mackay, L.K. (2016) Tissue-resident memory T cells: local specialists in immune defence. *Nat Rev Immunol* 16 (2), 79-89.
11. Sallusto, F. et al. (1999) Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. *Nature* 401 (6754), 708-712.
- 20 12. Gattinoni, L. et al. (2011) A human memory T cell subset with stem cell-like properties. *Nat. Med.* 17 (10), 1290-1297.
13. Joshi, N.S. et al. (2007) Inflammation directs memory precursor and short-lived effector CD8(+) T cell fates via the graded expression of T-bet transcription factor. *Immunity*. 27 (2), 281-295.
14. Sallusto, F. and Lanzavecchia, A. (2000) Understanding dendritic cell and T-lymphocyte traffic through the analysis of chemokine receptor expression. *Immunol.Rev.* 177, 134-140.
- 25 15. Sallusto, F. (2016) Heterogeneity of Human CD4(+) T Cells Against Microbes. *Annu Rev Immunol* 34, 317-34.
16. Reinhardt, R.L. et al. (2001) Visualizing the generation of memory CD4 T cells in the whole body. *Nature* 410 (6824), 101-105.
- 30 17. Masopust, D. et al. (2004) Activated primary and memory CD8 T cells migrate to nonlymphoid tissues regardless of site of activation or tissue of origin. *J Immunol* 172 (8), 4875-82.
18. Yu, B. et al. (2017) Epigenetic landscapes reveal transcription factors that regulate CD8(+) T cell differentiation. *Nat Immunol* 18 (5), 573-582.
19. Pace, L. et al. (2018) The epigenetic control of stemness in CD8(+) T cell fate commitment. *Science* 359 (6372), 177-186.
- 35 20. Moser, B. et al. (2004) Chemokines: multiple levels of leukocyte migration control. *Trends Immunol.* 25 (2), 75-84.
21. Ginhoux, F. et al. (2016) New insights into the multidimensional concept of macrophage ontogeny, activation and function. *Nat Immunol* 17 (1), 34-40.
- 40 22. Okabe, Y. and Medzhitov, R. (2016) Tissue biology perspective on macrophages. *Nat Immunol* 17 (1), 9-17.
23. McCully, M.L. and Moser, B. (2011) The human cutaneous chemokine system. *Frontiers Chemoattractants* 2, 1-15.
24. Bachelier, F. et al. (2014) International Union of Basic and Clinical Pharmacology. [corrected]. LXXXIX. Update on the extended family of chemokine receptors and introducing a new nomenclature for atypical chemokine receptors. *Pharmacol Rev* 66 (1), 1-79.
- 45 25. Loetscher, P. et al. (2000) Chemokines and their receptors in lymphocyte traffic and HIV infection. *Adv.Immunol.* 74, 127-180.
26. Nagasawa, T. (2014) CXC chemokine ligand 12 (CXCL12) and its receptor CXCR4. *Journal of Molecular Medicine-Jmm* 92 (5), 433-439.
- 50

27. Luo, Q. et al. (2016) Role of Stromal-Derived Factor-1 in Mesenchymal Stem Cell Paracrine-Mediated Tissue Repair. *Curr Stem Cell Res Ther* 11 (7), 585-92.
28. Graham, G.J. et al. (2012) The biochemistry and biology of the atypical chemokine receptors. *Immunol Lett* 145 (1-2), 30-8.
- 5 29. Mackay, L.K. and Kallies, A. (2017) Transcriptional Regulation of Tissue-Resident Lymphocytes. *Trends Immunol* 38 (2), 94-103.
30. Stelma, F. et al. (2017) Human intrahepatic CD69 + CD8+ T cells have a tissue resident memory T cell phenotype with reduced cytolytic capacity. *Sci Rep* 7 (1), 6172.
31. Stegmann, K.A. et al. (2016) CXCR6 marks a novel subset of T-bet(lo)Eomes(hi) natural killer cells  
10 residing in human liver. *Sci Rep* 6, 26157.
32. Lunemann, S. et al. (2017) Hobit expression by a subset of human liver-resident CD56(bright) Natural Killer cells. *Sci Rep* 7 (1), 6676.
33. Zaid, A. et al. (2017) Chemokine Receptor-Dependent Control of Skin Tissue-Resident Memory T Cell Formation. *J Immunol* 199 (7), 2451-2459.
- 15 34. Scholz, F. et al. (2007) Constitutive expression and regulated release of the transmembrane chemokine CXCL16 in human and murine skin. *J Invest Dermatol* 127 (6), 1444-55.
35. Gunther, C. et al. (2012) CXCL16 and CXCR6 are upregulated in psoriasis and mediate cutaneous recruitment of human CD8+ T cells. *J Invest Dermatol* 132 (3 Pt 1), 626-34.
36. Homey, B. et al. (2000) Up-regulation of macrophage inflammatory protein-3 alpha/CCL20 and CC  
20 chemokine receptor 6 in psoriasis. *J.Immunol.* 164 (12), 6621-6632.
37. Kwon, J.H. et al. (2002) Colonic epithelial cells are a major site of macrophage inflammatory protein 3alpha (MIP-3alpha) production in normal colon and inflammatory bowel disease. *Gut* 51 (6), 818-826.
38. Cai, Y. et al. (2011) Pivotal role of dermal IL-17-producing gammadelta T cells in skin  
25 inflammation. *Immunity* 35 (4), 596-610.
39. Hartwig, T. et al. (2015) Dermal IL-17-producing gammadelta T cells establish long-lived memory in the skin. *Eur J Immunol* 45 (11), 3022-33.
40. Ramirez-Valle, F. et al. (2015) Inflammation induces dermal Vgamma4+ gammadeltaT17 memory-like cells that travel to distant skin and accelerate secondary IL-17-driven responses. *Proc Natl Acad Sci U S A* 112 (26), 8046-51.
- 30 41. Zhang, Y. et al. (2016) Migratory and adhesive cues controlling innate-like lymphocyte surveillance of the pathogen-exposed surface of the lymph node. *Elife* 5.
42. Mowat, A.M. and Agace, W.W. (2014) Regional specialization within the intestinal immune system. *Nat Rev Immunol* 14 (10), 667-85.
- 35 43. Marsal, J. and Agace, W.W. (2012) Targeting T-cell migration in inflammatory bowel disease. *J Intern Med* 272 (5), 411-29.
44. Homey, B. et al. (2000) Cutting edge: the orphan chemokine receptor G protein-coupled receptor-2 (GPR-2, CCR10) binds the skin-associated chemokine CCL27 (CTACK/ALP/ILC). *J.Immunol.* 164 (7), 3465-3470.
- 40 45. Wang, W. et al. (2000) Identification of a novel chemokine (CCL28), which binds CCR10 (GPR2). *J.Biol.Chem.* 275 (29), 22313-22323.
46. Ishikawa-Mochizuki, I. et al. (1999) Molecular cloning of a novel CC chemokine, interleukin-11 receptor  $\alpha$ -locus chemokine (ILC), which is located on chromosome 9p13 and a potential homologue of a CC chemokine encoded by molluscum contagiosum virus. *FEBS Lett.* 460 (3), 544-548.
- 45 47. Morales, J. et al. (1999) CTACK, a skin-associated chemokine that preferentially attracts skin-homing memory T cells. *Proc.Natl.Acad.Sci.USA* 96 (25), 14470-14475.
48. Homey, B. et al. (2002) CCL27-CCR10 interactions regulate T cell-mediated skin inflammation. *Nat.Med.* 8 (2), 157-165.
49. Pan, J. et al. (2000) A novel chemokine ligand for CCR10 and CCR3 expressed by epithelial cells in mucosal tissues. *J.Immunol.* 165 (6), 2943-2949.
- 50 50. Hernandez-Ruiz, M. and Zlotnik, A. (2017) Mucosal Chemokines. *J Interferon Cytokine Res* 37 (2), 62-70.

51. Matsuo, K. et al. (2018) CCL28-Deficient Mice Have Reduced IgA Antibody-Secreting Cells and an Altered Microbiota in the Colon. *J Immunol* 200 (2), 800-809.
52. Morteau, O. et al. (2008) An indispensable role for the chemokine receptor CCR10 in IgA antibody-secreting cell accumulation. *J Immunol* 181 (9), 6309-15.
- 5 53. Danilova, E. et al. (2015) A role for CCL28-CCR3 in T-cell homing to the human upper airway mucosa. *Mucosal Immunol* 8 (1), 107-14.
54. Reiss, Y. et al. (2001) CC chemokine receptor (CCR)4 and the CCR10 ligand cutaneous T cell-attracting chemokine (CTACK) in lymphocyte trafficking to inflamed skin. *J.Exp.Med.* 194 (10), 1541-1547.
- 10 55. Fu, Y. et al. (2016) Cutting Edge: Skin CCR10+ CD8+ T Cells Support Resident Regulatory T Cells through the B7.2/Receptor Axis To Regulate Local Immune Homeostasis and Response. *J Immunol* 196 (12), 4859-64.
56. Xia, M. et al. (2014) CCR10 regulates balanced maintenance and function of resident regulatory and effector T cells to promote immune homeostasis in the skin. *J Allergy Clin Immunol* 134 (3), 634-644 e10.
- 15 57. Yang, J. et al. (2016) Selective programming of CCR10(+) innate lymphoid cells in skin-draining lymph nodes for cutaneous homeostatic regulation. *Nat Immunol* 17 (1), 48-56.
58. Regoli, M. et al. (2017) The Multifaceted Personality of Intestinal CX3CR1(+) Macrophages. *Trends Immunol* 38 (12), 879-887.
- 20 59. Limatola, C. and Ransohoff, R.M. (2014) Modulating neurotoxicity through CX3CL1/CX3CR1 signaling. *Front Cell Neurosci* 8, 229.
60. Gerlach, C. et al. (2016) The Chemokine Receptor CX3CR1 Defines Three Antigen-Experienced CD8 T Cell Subsets with Distinct Roles in Immune Surveillance and Homeostasis. *Immunity* 45 (6), 1270-1284.
- 25 61. Bottcher, J.P. et al. (2015) Functional classification of memory CD8(+) T cells by CX3CR1 expression. *Nat Commun* 6, 8306.
62. Konkel, J.E. et al. (2017) Transforming Growth Factor-beta Signaling in Regulatory T Cells Controls T Helper-17 Cells and Tissue-Specific Immune Responses. *Immunity* 46 (4), 660-674.
63. Lahl, K. et al. (2014) Orphan chemoattractant receptor GPR15 mediates dendritic epidermal T-cell recruitment to the skin. *Eur J Immunol* 44 (9), 2577-81.
- 30 64. Vantourout, P. and Hayday, A. (2013) Six-of-the-best: unique contributions of gammadelta T cells to immunology. *Nat.Rev.Immunol.* 13 (2), 88-100.
65. Nguyen, L.P. et al. (2015) Role and species-specific expression of colon T cell homing receptor GPR15 in colitis. *Nat Immunol* 16 (2), 207-213.
- 35 66. Kim, S.V. et al. (2013) GPR15-mediated homing controls immune homeostasis in the large intestine mucosa. *Science* 340 (6139), 1456-9.
67. Pan, B. et al. (2017) The fifth epidermal growth factor like region of thrombomodulin alleviates LPS-induced sepsis through interacting with GPR15. *Thromb Haemost* 117 (3), 570-579.
68. Suply, T. et al. (2017) A natural ligand for the orphan receptor GPR15 modulates lymphocyte recruitment to epithelia. *Sci Signal* 10 (496).
- 40 69. Lu, J. et al. (2016) CXCL14 as an emerging immune and inflammatory modulator. *J Inflamm (Lond)* 13, 1.
70. Meuter, S. et al. (2007) Murine CXCL14 is dispensable for dendritic cell function and localization within peripheral tissues. *Mol.Cell Biol.* 27 (3), 983-992.
- 45 71. Nara, N. et al. (2007) Disruption of CXC motif chemokine ligand-14 in mice ameliorates obesity-induced insulin resistance. *J.Biol.Chem.* 282 (42), 30794-30803.
72. Tanegashima, K. et al. (2010) CXCL14 deficiency in mice attenuates obesity and inhibits feeding behavior in a novel environment. *PLoS.ONE.* 5 (4), e10321.
73. Collins, P.J. et al. (2017) Epithelial chemokine CXCL14 synergizes with CXCL12 via allosteric modulation of CXCR4. *FASEB J* 31 (7), 3084-3097.
- 50 74. Burkhardt, A.M. et al. (2014) CXCL17 is a major chemotactic factor for lung macrophages. *J Immunol* 193 (3), 1468-74.

75. Oka, T. et al. (2017) CXCL17 Attenuates Imiquimod-Induced Psoriasis-like Skin Inflammation by Recruiting Myeloid-Derived Suppressor Cells and Regulatory T Cells. *J Immunol* 198 (10), 3897-3908.
76. Maravillas-Montero, J.L. et al. (2015) Cutting edge: GPR35/CXCR8 is the receptor of the mucosal chemokine CXCL17. *J Immunol* 194 (1), 29-33.
- 5 77. Park, S.J. et al. (2018) GPR35 mediates Iodoxamide-induced migration inhibitory response but not CXCL17-induced migration stimulatory response in THP-1 cells; is GPR35 a receptor for CXCL17? *Br J Pharmacol* 175 (1), 154-161.
78. Tan, J.K. et al. (2017) Metabolite-Sensing G Protein-Coupled Receptors-Facilitators of Diet-Related Immune Regulation. *Annu Rev Immunol* 35, 371-402.
- 10 79. McCully, M.L. et al. (2018) CCR8 Expression Defines Tissue-Resident Memory T Cells in Human Skin. *J Immunol* 200 (5), 1639-1650.
80. Clark, R.A. et al. (2006) The vast majority of CLA<sup>+</sup> T cells are resident in normal skin. *J.Immunol.* 176 (7), 4431-4439.
- 15 81. Milner, J.J. et al. (2017) Runx3 programs CD8(+) T cell residency in non-lymphoid tissues and tumours. *Nature* 552 (7684), 253-257.
82. Watanabe, R. et al. (2015) Human skin is protected by four functionally and phenotypically discrete populations of resident and recirculating memory T cells. *Sci Transl Med* 7 (279), 279ra39.
83. Bromley, S.K. et al. (2013) Recirculating memory T cells are a unique subset of CD4<sup>+</sup> T cells with a distinct phenotype and migratory pattern. *J Immunol* 190 (3), 970-6.
- 20 84. Scharschmidt, T.C. et al. (2015) A Wave of Regulatory T Cells into Neonatal Skin Mediates Tolerance to Commensal Microbes. *Immunity* 43 (5), 1011-21.
85. Sanchez Rodriguez, R. et al. (2014) Memory regulatory T cells reside in human skin. *J Clin Invest* 124 (3), 1027-36.
86. Tomura, M. et al. (2010) Activated regulatory T cells are the major T cell type emigrating from the skin during a cutaneous immune response in mice. *J Clin Invest* 120 (3), 883-93.
- 25 87. Colantonio, L. et al. (2002) Skin-homing CLA<sup>+</sup> T cells and regulatory CD25<sup>+</sup> T cells represent major subsets of human peripheral blood memory T cells migrating in response to CCL1/I-309. *Eur.J.Immunol.* 32 (12), 3506-3514.
88. Plitas, G. et al. (2016) Regulatory T Cells Exhibit Distinct Features in Human Breast Cancer. *Immunity* 45 (5), 1122-1134.
- 30 89. De Simone, M. et al. (2016) Transcriptional Landscape of Human Tissue Lymphocytes Unveils Uniqueness of Tumor-Infiltrating T Regulatory Cells. *Immunity* 45 (5), 1135-1147.
90. Barsheshet, Y. et al. (2017) CCR8+FOXP3<sup>+</sup> Treg cells as master drivers of immune regulation. *Proc Natl Acad Sci U S A* 114 (23), 6086-6091.
- 35 91. Hoelzinger, D.B. et al. (2010) Blockade of CCL1 inhibits T regulatory cell suppressive function enhancing tumor immunity without affecting T effector responses. *J Immunol* 184 (12), 6833-42.
92. Asai, A. et al. (2012) CCL1 released from M2b macrophages is essentially required for the maintenance of their properties. *J Leukoc Biol* 92 (4), 859-67.
93. Mackay, L.K. et al. (2013) The developmental pathway for CD103(+)CD8<sup>+</sup> tissue-resident memory T cells of skin. *Nat Immunol* 14 (12), 1294-301.
- 40 94. Islam, S.A. et al. (2011) Mouse CCL8, a CCR8 agonist, promotes atopic dermatitis by recruiting IL-5<sup>+</sup> T(H)2 cells. *Nat.Immunol.* 12 (2), 167-177.
95. Mora, J.R. and Von Andrian, U.H. (2006) T-cell homing specificity and plasticity: new concepts and future challenges. *Trends Immunol.* 27 (5), 235-243.
- 45 96. Sigmundsdottir, H. and Butcher, E.C. (2008) Environmental cues, dendritic cells and the programming of tissue-selective lymphocyte trafficking. *Nat.Immunol.* 9 (9), 981-987.
97. McCully, M.L. et al. (2012) Epidermis instructs skin homing receptor expression in human T cells. *Blood* 120 (23), 4591-4598.
98. McCully, M.L. et al. (2015) Skin Metabolites Define a New Paradigm in the Localization of Skin Tropic Memory T Cells. *J Immunol* 195 (1), 96-104.
- 50 99. Sigmundsdottir, H. et al. (2007) DCs metabolize sunlight-induced vitamin D3 to 'program' T cell attraction to the epidermal chemokine CCL27. *Nat.Immunol.* 8 (3), 285-293.



100. Agace, W.W. and Persson, E.K. (2012) How vitamin A metabolizing dendritic cells are generated in the gut mucosa. *Trends Immunol* 33 (1), 42-8.
101. Iwata, M. et al. (2004) Retinoic acid imprints gut-homing specificity on T cells. *Immunity*. 21 (4), 527-538.
- 5 102. Mora, J.R. et al. (2006) Generation of gut-homing IgA-secreting B cells by intestinal dendritic cells. *Science* 314 (5802), 1157-60.
103. Kim, M.H. et al. (2015) Retinoic Acid Differentially Regulates the Migration of Innate Lymphoid Cell Subsets to the Gut. *Immunity* 43 (1), 107-19.
104. Rosas, M. et al. (2014) The Transcription Factor Gata6 Links Tissue Macrophage Phenotype and Proliferative Renewal. *Science* 344 (6184), 645-648.
- 10 105. Okabe, Y. and Medzhitov, R. (2014) Tissue-specific signals control reversible program of localization and functional polarization of macrophages. *Cell* 157 (4), 832-44.
106. Urry, Z. et al. (2012) The role of 1 $\alpha$ ,25-dihydroxyvitamin D3 and cytokines in the promotion of distinct Foxp3+ and IL-10+ CD4+ T cells. *Eur J Immunol* 42 (10), 2697-708.
- 15 107. Bakdash, G. et al. (2015) Retinoic acid primes human dendritic cells to induce gut-homing, IL-10-producing regulatory T cells. *Mucosal Immunol* 8 (2), 265-78.
108. van der Aar, A.M. et al. (2011) Vitamin D3 targets epidermal and dermal dendritic cells for induction of distinct regulatory T cells. *J Allergy Clin Immunol* 127 (6), 1532-40 e7.
109. Pan, Y. et al. (2017) Survival of tissue-resident memory T cells requires exogenous lipid uptake and metabolism. *Nature* 543 (7644), 252-256.
- 20 110. Casey, K.A. et al. (2012) Antigen-independent differentiation and maintenance of effector-like resident memory T cells in tissues. *J.Immunol.* 188 (10), 4866-4875.
111. Mackay, L.K. et al. (2015) T-box Transcription Factors Combine with the Cytokines TGF-beta and IL-15 to Control Tissue-Resident Memory T Cell Fate. *Immunity* 43 (6), 1101-11.
- 25 112. Iijima, N. and Iwasaki, A. (2014) T cell memory. A local macrophage chemokine network sustains protective tissue-resident memory CD4 T cells. *Science* 346 (6205), 93-8.
113. Ariotti, S. et al. (2012) Tissue-resident memory CD8+ T cells continuously patrol skin epithelia to quickly recognize local antigen. *Proc.Natl.Acad.Sci.U.S.A* 109 (48), 19739-19744.
114. Beura, L.K. et al. (2018) Intravital mucosal imaging of CD8(+) resident memory T cells shows tissue-autonomous recall responses that amplify secondary memory. *Nat Immunol* 19 (2), 173-182.
- 30 115. Miller, M.D. et al. (1989) A novel polypeptide secreted by activated human T lymphocytes. *J.Immunol.* 143, 2907-2916.
116. Burd, P.R. et al. (1987) Cloning and characterization of a novel T cell activation gene. *J.Immunol.* 139, 3126-3131.
- 35 117. Miller, M.D. and Krangel, M.S. (1992) The human cytokine I-309 is a monocyte chemoattractant. *Proc.Natl.Acad.Sci.USA* 89, 2950-2954.
118. Tiffany, H.L. et al. (1997) Identification of CCR8: A human monocyte and thymus receptor for the CC chemokine I-309. *J.Exp.Med.* 186, 165-170.
119. Roos, R.S. et al. (1997) Identification of CCR8, the receptor for the human CC chemokine I-309. *J.Biol.Chem.* 272, 17251-17254.
- 40 120. Goya, I. et al. (1998) Identification of CCR8 as the specific receptor for the human  $\beta$ -chemokine I-309: Cloning and molecular characterization of murine CCR8 as the receptor for TCA-3. *J.Immunol.* 160, 1975-1981.
121. Islam, S.A. et al. (2013) Identification of human CCR8 as a CCL18 receptor. *J.Exp.Med.* 210 (10), 1889-1898.
- 45 122. Goya, I. et al. (2003) Absence of CCR8 does not impair the response to ovalbumin-induced allergic airway disease. *J.Immunol.* 170 (4), 2138-2146.
123. Chung, C.D. et al. (2003) CCR8 is not essential for the development of inflammation in a mouse model of allergic airway disease. *J.Immunol.* 170 (1), 581-587.
- 50 124. Chensue, S.W. et al. (2001) Aberrant in vivo T helper type 2 cell response and impaired eosinophil recruitment in CC chemokine receptor 8 knockout mice. *J.Exp.Med.* 193 (5), 573-584.

125. Buckland, K.F. et al. (2007) Remission of chronic fungal asthma in the absence of CCR8. *J Allergy Clin Immunol* 119 (4), 997-1004.
126. Qu, C. et al. (2004) Role of CCR8 and other chemokine pathways in the migration of monocyte-derived dendritic cells to lymph nodes. *J.Exp.Med.* 200 (10), 1231-1241.
- 5 127. Yabe, R. et al. (2015) CCR8 regulates contact hypersensitivity by restricting cutaneous dendritic cell migration to the draining lymph nodes. *Int Immunol* 27 (4), 169-81.
128. Nagao, K. et al. (2012) Stress-induced production of chemokines by hair follicles regulates the trafficking of dendritic cells in skin. *Nat Immunol* 13 (8), 744-52.
129. Annunziato, F. et al. (1998) Molecules associated with human Th1 or Th2 cells. *Eur.Cytokine Netw.* 9 Suppl. 3, 12-16.
- 10 130. D'Ambrosio, D. et al. (1998) Selective up-regulation of chemokine receptors CCR4 and CCR8 upon activation of polarized human type 2 Th cells. *J.Immunol.* 161, 5111-5115.
131. Zingoni, A. et al. (1998) The chemokine receptor CCR8 is preferentially expressed in Th2 but not Th1 cells. *J.Immunol.* 161, 547-551.
- 15 132. Panina-Bordignon, P. et al. (2001) The C-C chemokine receptors CCR4 and CCR8 identify airway T cells of allergen-challenged atopic asthmatics. *J.Clin.Invest* 107 (11), 1357-1364.
133. Sebastiani, S. et al. (2001) Chemokine receptor expression and function in CD4+ T lymphocytes with regulatory activity. *J.Immunol.* 166 (2), 996-1002.
134. Gombert, M. et al. (2005) CCL1-CCR8 interactions: an axis mediating the recruitment of T cells and Langerhans-type dendritic cells to sites of atopic skin inflammation. *J.Immunol.* 174 (8), 5082-5091.
- 20 135. Gonzalo, J.A. et al. (2007) Coordinated involvement of mast cells and T cells in allergic mucosal inflammation: critical role of the CC chemokine ligand 1:CCR8 axis. *J.Immunol.* 179 (3), 1740-1750.
136. Soler, D. et al. (2006) CCR8 expression identifies CD4 memory T cells enriched for FOXP3+ regulatory and Th2 effector lymphocytes. *J.Immunol.* 177 (10), 6940-6951.
- 25 137. Coghill, J.M. et al. (2013) CC chemokine receptor 8 potentiates donor Treg survival and is critical for the prevention of murine graft-versus-host disease. *Blood* 122 (5), 825-36.
138. Ebert, L.M. et al. (2006) Homing and function of human skin gammadelta T cells and NK cells: relevance for tumor surveillance. *J.Immunol.* 176 (7), 4331-4336.
- 30 139. Schaerli, P. et al. (2004) A Skin-selective Homing Mechanism for Human Immune Surveillance T Cells. *J.Exp.Med.* 199 (9), 1265-1275.
140. Mutalithas, K. et al. (2010) Expression of CCR8 is increased in asthma. *Clin.Exp.Allergy* 40 (8), 1175-1185.
141. Nakajima, T. et al. (2002) Marked increase in CC chemokine gene expression in both human and mouse mast cell transcriptomes following Fcepsilon receptor I cross-linking: an interspecies comparison. *Blood* 100 (12), 3861-3868.
- 35 142. Gilcrest, H. et al. (2003) Human cord blood-derived mast cells synthesize and release I-309 in response to IgE. *Life Sci* 73 (20), 2571-81.
143. N'Diaye, M. et al. (2006) Aryl hydrocarbon receptor- and calcium-dependent induction of the chemokine CCL1 by the environmental contaminant benzo[a]pyrene. *J Biol Chem* 281 (29), 19906-15.
- 40 144. Hoshino, A. et al. (2007) Inhibition of CCL1-CCR8 interaction prevents aggregation of macrophages and development of peritoneal adhesions. *J Immunol* 178 (8), 5296-304.
145. Das, S. et al. (2013) Tumor cell entry into the lymph node is controlled by CCL1 chemokine expressed by lymph node lymphatic sinuses. *J Exp Med* 210 (8), 1509-28.
- 45 146. Lüttichau, H.R. et al. (2000) A highly selective CC chemokine receptor (CCR)8 antagonist encoded by the poxvirus mollusum contagiosum. *J.Exp.Med.* 191 (1), 171-179.
147. Kledal, T.N. et al. (1997) A broad-spectrum chemokine antagonist encoded by Kaposi's sarcoma-associated herpesvirus. *Science* 277, 1656-1659.

## Highlights

---

Chemokines orchestrate immune cell trafficking, and thus the site of chemokine production can define the local immune cell composition.

5 Healthy peripheral tissues host a highly complex network of immune surveillance cells, including macrophages, antigen-presenting cells and functionally diverse subsets of T cells and innate lymphoid cells. Homeostatic chemokines responsible for their recruitment, tissue retention and/or immune effector function have remained ill-defined.

10 Peripheral tissue chemokines orchestrate the function of the local immune surveillance system and, therefore, are essential contributors to tissue health and longevity.

In human skin, CCR8 and its ligand CCL1 are thought to play a critical role in the migration and maintenance of resident immune surveillance cells.

## Outstanding Questions

---


- Are homeostatic chemokine systems, such as CCL1-CCR8 in human skin, necessary for TRM recruitment, localization and/or retention, survival or memory responses to local antigens?
- 5      - What is the role of tissue non-specific chemokine receptors present on TRM cells? Do they synergize with receptors for homeostatic chemokines or are they mainly involved in controlling immune cell mobilization in response to local infections?
- How do tissue factors, including vitamins, imprint tissue residency / survival?
- 10      Do they also work on other lymphocyte populations (e.g. ILCs,  $\gamma\delta$  T cells)?
- Since numerous memory  $\alpha\beta$  T cells in human skin are not tissue-resident (i.e. do not express tissue residence markers CD69  $\pm$  CD103), what chemokine networks regulate their recirculation?
- Can peripheral tissue chemokine networks be manipulated to promote tissue-specific immunity, e.g. Treg cell recruitment in the case of autoimmune
- 15      diseases or localization and/or maintenance of memory T cells in response to vaccination?

**Table 1.** Chemokine receptors involved in peripheral tissue T cell traffic


Chemokine receptor	Cellular expression	Ligands	Function in tissue lymphocyte traffic	Refs
CXCR4 (ACKR3)	Haematopoietic and non-haematopoietic stem cells, thymocytes, leukocytes, tissue cells	CXCL12	CXCR4 is a chemokine receptor whereas ACKR3 binds and internalizes CXCL12, thereby regulating extracellular CXCL12 concentrations. The CXCL12-CXCR4/ACKR3 axis controls bone marrow stem cell and progenitor localization, thymocyte development, plasma B cell homing as well as tissue (brain, heart, vasculature) development and repair. CXCR4 also functions as co-receptor for HIV-1	[3, 25, 26, 28]
CXCR6	$\alpha\beta$ T, $\gamma\delta$ T, NKT, NK, plasma cells	CXCL16	CXCL16 is either membrane-bound (cell-cell adhesion) or soluble (chemotaxis). CXCR6 associated with memory T cell and /NK cell compartments in liver and skin.	[30-33]
CCR6	$\alpha\beta$ T, $\gamma\delta$ T, Treg, ILCs, NKT, NK, B, immature DCs	CCL20	CCL20 is present at low levels in healthy skin but is massively upregulated during inflammation. Besides inflammatory involvement, CCR6 contributes to the maintenance of memory $\gamma\delta$ T17 cells in healthy skin and draining LNs.	[39-41]
CCR8	Human skin lymphocytes ( $\alpha\beta$ T, $\gamma\delta$ T, NKT, NK). Mouse thymocytes, Th2 and Treg	CCL1 CCL8 (mouse) CCL18 (human)	CCL1, the primary ligand for CCR8, is constitutively expressed in human skin and rapidly released by activated skin TRM cells. Murine CCL8 and human CCL18 are alternative ligands for mouse and human CCR8, respectively. Murine CCR8 is primarily a T cell-associated chemokine receptor with expression on skin-homing Th2 and Treg cells. CCR8 on human skin T cells is discussed in the main text.	[94, 121, 138, 139]
CCR9	Thymocytes, intestinal ( $\alpha\beta$ T, $\gamma\delta$ T)	CCL25	CCL25 is primarily produced in small intestinal tissues. CCR9 is present almost exclusively on intestinal $\alpha\beta$ T cells, $\gamma\delta$ T cells and plasmablasts but its function in intestinal T cell traffic, including T cell recruitment and maintenance of memory T cell compartment appears to be redundant.	[43]
CCR10	Effector T, B, ILCs, melanocytes	CCL27 CCL28	CCL27 and CCL28 share mutually exclusive expression pattern, CCL27 being present in the skin whereas CCL28 is being expressed in colon and secretory organs. CCR10 expressing CD8 <sup>+</sup> T cells and Treg cells support survival of T cells as well as CCR10 <sup>+</sup> ILCs. CCR10 on melanocytes/melanoma cells may control their skin retention.	[55-57]

CX3CR1	Monos, LCs, mΦ, NK, αβ T, γδ T, platelets, neurons	CX3CL16	<p>CX3CL1 exists in two forms, membrane-bound (cell adhesion) or secreted (chemotaxis).</p> <p>Expression of CX3CR1 on macrophages (microglia, LCs) and intestinal DCs was shown to support local immune surveillance functions. Graded surface levels of CX3CR1 were used to distinguish effector/memory T cells subsets, including long-lived recirculating and LN-sessile memory T cells.</p>	[60, 61]
--------	--	---------	--	----------

## Primary Infection



## Tissue Recovery



## Immune Surveillance

