

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/115758/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Best, Damian, Avenell, Alison and Bhattacharya, Siladitya 2017. How effective are weight-loss interventions for improving fertility in women and men who are overweight or obese? A systematic review and meta-analysis of the evidence. *Human Reproduction Update* 23 (6) , pp. 681-705. 10.1093/humupd/dmx027

Publishers page: <http://dx.doi.org/10.1093/humupd/dmx027>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1 **How effective are weight loss interventions for improving fertility in women and men**
2 **who are overweight or obese? A systematic review and meta-analysis of the evidence.**

3 **Authors:** Damian Best¹, Alison Avenell², Siladitya Bhattacharya¹

4 ¹Aberdeen Fertility Centre, University of Aberdeen, Aberdeen, United Kingdom, AB25 2ZL;

5

6 ²Health Services Research Unit, University of Aberdeen Foresterhill, Aberdeen, United Kingdom,

7 AB25 2ZD

8

9 Corresponding author Damian Best: dajbest1@gmail.com

10 Obstetrics & Gynaecology, Faculty of Medical Sciences, University of the West Indies Queen

11 Elizabeth Hospital, St. Michael, Barbados, BB11155.

12 **Running title:** Do weight loss interventions improve fertility?

13 **AUTHOR:** there is a limit of 50 characters for the running title. [Is this alternative acceptable?](#)

14

15

Style Definition: TOC 1: Font: Not Bold, Do not check spelling or grammar

Commented [BDA1]: Yes

16 Table of Contents

17 **AUTHOR:** the table of contents does not appear to agree with the text. Only first (e.g.
18 **Introduction, Methods**) and second level headings (e.g. *Criteria for considering studies for*
19 *this review*) can be included in the Table of Contents (although third etc. level headings are
20 allowed in the main text). The styles used for headings in the main text should be consistent
21 and only second level headers included here. Please would you recheck and edit both the
22 main text and Table of Contents accordingly? Thank you.

23 Abstract

24 Introduction

25 Methods

26 Criteria for considering studies for this review

27 Search methods for identification of studies

28 Data collection and analysis

29 Assessment of risk of bias in individual studies

30 Summary measures

31 Synthesis of results

32 Assessment of heterogeneity

33 Sensitivity analyses

34 Results

35 Outcome of the search

36 Description of populations

37 Description of interventions

38 Quality of studies

39 Studies involving female infertility participants

40 Comparisons with metformin

41 Barriers to weight loss in infertile women

Formatted: Font: (Default) Times New Roman, 12 pt, Bold, Check spelling and grammar

Formatted: Heading 2

Formatted: Font: (Default) Times New Roman, 12 pt, Check spelling and grammar

Formatted: Font: (Default) Times New Roman, 12 pt, Bold, Check spelling and grammar

Formatted: Font: (Default) Times New Roman, 12 pt, Check spelling and grammar

Formatted: Font: (Default) Times New Roman, 12 pt, Bold, Font color: Auto, Check spelling and grammar

Formatted: Font: (Default) Times New Roman, 12 pt, Bold, Check spelling and grammar

Formatted: Font: (Default) Times New Roman, 12 pt, Check spelling and grammar

Formatted: Font: (Default) Times New Roman, 12 pt, Check spelling and grammar

Formatted: Check spelling and grammar

Formatted: Font: (Default) Times New Roman, 12 pt, Check spelling and grammar

Commented [BDA2]: Adjustments have been made

Formatted: Font: (Default) Times New Roman, 12 pt, Check spelling and grammar

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Check spelling and grammar

Formatted: Normal, Indent: Left: 0.39 cm, Tab stops: Not at 15.9 cm

42 Studies involving male infertility participants

43 Discussion

44 Conclusions

45

46 **Abstract**

47

48 **BACKGROUND**

49 The prevalence of obesity is increasing worldwide, with a corresponding increase in overweight
50 and obese patients referred with infertility. This systematic review aimed to determine whether
51 non-surgical weight reduction strategies result in an improvement in reproductive parameters
52 affected by obesity, e.g. delayed time to pregnancy, oligozoospermia and azoospermia. No prior
53 reviews have examined this within the general fertility population, or in both sexes.

54

55 **OBJECTIVE AND RATIONALE**

56 Our objective was to answer the question: "In overweight and obese women, men and couples
57 seeking fertility treatment, what non-surgical weight loss interventions have been used, and
58 how effective are they at weight loss and improving reproductive outcomes?"

59

60 **SEARCH METHODS**

61 An electronic search of MEDLINE, EMBASE and the Cochrane Library was performed for studies
62 published between [AUTHOR: correct?] January 1966 and March 2016. Text word and MESH
63 search terms used related to infertility, weight, and barriers to weight loss. Inclusion criteria
64 were: an intervention to change lifestyle evaluated in any study design in participants of either
65 gender with an unfulfilled desire to conceive. Studies were excluded if they included
66 participants not attempting pregnancy, with illnesses that might cause weight fluctuations, or
67 studies evaluating bariatric surgery. Two reviewers performed data extraction and quality

Commented [BDA3]: Yes

68 assessment using the Cochrane Risk of Bias Tool for randomised trials, and a ratified checklist
69 (ReBIP) for non-randomised studies.

70

71 **OUTCOMES**

72 A total of 40 studies were included, of which 14 were RCTs. Primary outcomes were pregnancy,
73 live birth rate and weight change. In women, reduced calorie diets and exercise interventions
74 were more likely than control interventions to result in pregnancy [Risk Ratio 1.59, 95% CI
75 (1.01, 2.50)], and interventions resulted in weight loss and ovulation improvement, where
76 reported. Miscarriage rates were not reduced by any intervention.

77

78 **WIDER IMPLICATIONS**

79 Overweight and obese persons seeking fertility should be educated on the detrimental effects of
80 fatness and the benefits of weight reduction, including improvement in pregnancy rates. A
81 combination of a reduced calorie diet, by reducing fat and refined carbohydrate intake, and
82 increased aerobic exercise should form the basis of programmes designed for such individuals.
83 A lack of randomised studies in men and couples, and studies evaluating barriers to undertaking
84 weight loss in infertile populations is evident, and future research should examine these issues
85 further.

86

87 **Key words:** infertility, obesity, weight loss, systematic review, diet, exercise

88 **Introduction**

89

90 Overweight is defined by the World Health Organization (WHO) (World Health Organization,
91 2016) as a BMI ≥ 25 kg/m², and obesity as ≥ 30 kg/m². The prevalence of obesity is increasing
92 worldwide (World Health Organization, 2016), with more than 600 million obese adults,
93 including 15% of women, in 2014 - double the prevalence reported three decades earlier.

94 Infertility is defined by the failure to achieve a clinical pregnancy after 12 months or more of
95 regular unprotected sexual intercourse (Zegers-Hochschild *et al.*, 2009; National Institute for
96 Health and Clinical Excellence, 2013). Using this definition, infertility affects about 1 in 7
97 couples in the UK, which is similar to an estimate of prevalence in the USA (Thoma *et al.*, 2013).

98 An increasingly overweight and obese population has led to a greater proportion of individuals
99 with high BMI being evaluated in infertility settings (Vahratian and Smith, 2009).

100

101 BMI has been associated with delayed conception in women in a dose-dependent fashion
102 (Gesink Law *et al.*, 2007; Wise *et al.*, 2013; Wise *et al.*, 2010). It is thought to produce
103 anovulation (Zain and Norman, 2008; Klenov and Jungheim, 2014), which results in menstrual
104 irregularity (Hartz *et al.*, 1979; Zain and Norman, 2008); impairment of oocyte development
105 (Klenov and Jungheim, 2014) and quality (Carrell *et al.*, 2001; Metwally *et al.*, 2007); and it also
106 has direct effects on the endometrium (Bellver *et al.*, 2011). With ovulation induction (OI)
107 treatment, higher doses of medications (Dickey *et al.*; Balen *et al.*, 2006a) and a longer period of
108 stimulation (Balen *et al.*, 2006b) are required with a higher BMI. With IVF, the chances of
109 achieving a pregnancy decrease with each unit increase in BMI (Ferlitsch *et al.*, 2004), and it is
110 thought that implantation rates are lower in obese women (Bellver *et al.*, 2010; Styne-Gross *et al.*
111 *et al.*, 2005). An increased risk of miscarriage following IVF has also been reported (Maheshwari
112 *et al.*, 2007).

113

114 Obesity may affect male fertility in a dose-dependent manner (Sallmen *et al.*, 2006). Postulated
115 mechanisms include increased testicular temperature with prolonged sitting (Hammoud *et al.*,
116 2012), and increased oestrogen production in fat with disruption of the hypothalamo-pituitary-
117 gonadal axis (Schneider *et al.*, 1979; Shukla *et al.*, 2014). Some studies have suggested reduced
118 sperm concentration and motility (Hammiche *et al.*, 2012; Hakonsen *et al.*, 2011; Sermondade
119 *et al.*, 2013), and lower clinical pregnancy rates with IVF (Keltz *et al.*, 2010; Umul *et al.*, 2015)
120 with increasing male adiposity. Where both partners are obese, greater difficulty achieving
121 pregnancy is expected (Ramlau-Hansen *et al.*, 2007).

122

123 Patients attending for fertility treatments are often advised to optimize their weights to
124 improve outcomes, and in some cases to access assistance with funding (National Institute for
125 Health and Clinical Excellence, 2013; Farquhar and Gillett, 2006; Infertility Network UK, 2015)
126 or to reduce the risks of obstetric complications (The Royal Australian and New Zealand College
127 of Obstetricians and Gynaecologists, 2011).

128

129 The purpose of this systematic review was to investigate first whether weight loss interventions
130 for infertile patients achieve their goal in reducing weight, and second whether they result in
131 improved fertility outcomes. A prior systematic review of 11 studies in women (Sim K.A. *et al.*,
132 2014) focussed on assisted reproduction only and used a much more limited search for studies
133 up until 2014. Our review used an extensive search until 2016 and presents data from 40
134 studies with women and men with infertility, not just patients requiring assisted reproduction.

135

136 **Methods**

137

138 This systematic review was undertaken according to a pre-specified protocol.

139

140 ***Criteria for considering studies for this review***

141

142 *Types of studies*

143 We included intervention studies in overweight or obese participants with any design and any
144 duration of follow-up.

145

146 *Types of participants*

147 Participants could be of either sex. They were required to have both infertility, defined by an
148 unfulfilled desire to conceive of any duration or attendance for infertility investigations or
149 treatment, and a prescribed period of attempt at weight loss. A BMI ≥ 25 kg/m² was used to
150 define overweight, and ≥ 30 kg/m² to define obesity (World Health Organization, 2016);
151 participants were required to be at least overweight. Morbid obesity was defined by BMI ≥ 40
152 kg/m². Studies were excluded if participants were not actively attempting pregnancy, or had an
153 illness or eating disorder that might result in weight fluctuation.

154

155 *Types of interventions*

156 Any intervention for weight loss allowed inclusion, for example prompt for weight reduction,
157 dietary modification, exercise, psychological or behavioural counselling, or the drug orlistat.
158 Bariatric surgery was not included. Alternative or control interventions were included where
159 evaluated in included studies.

160

161 *Types of outcome measures*

162 Primary outcomes were weight change, achievement of pregnancy, and live birth, the latter
163 defined as the complete expulsion or extraction from its mother of a product of fertilisation,
164 which, after such separation, breathes or shows any other evidence of life such as heart beat,
165 umbilical cord pulsation, or definite movement of voluntary muscles, irrespective of whether
166 the umbilical cord has been cut or the placenta is attached (National Institute for Health and
167 Clinical Excellence, 2013).

168
169 Secondary outcomes were BMI change; waist circumference change (North American
170 Association for the Study of Obesity *et al.*, 2000); miscarriage, defined as a loss of pregnancy
171 before 23+6 weeks of gestation (Anonymous, 2012); ovulation improvement, with anovulation
172 defined by suboptimal rise in LH, or a below-threshold serum progesterone level taken in the
173 mid-luteal phase of the menstrual cycle (National Institute for Health and Clinical Excellence,
174 2013); improvement in menstrual regularity, with irregular cycles defined as outside the range
175 of 26-36 days (National Institute for Health and Clinical Excellence, 2013); and time to
176 conception.

178 ***Search methods for identification of studies***

179
180 The dates searched were 1 January 1966 to 19 March 2016. No restriction was placed on
181 language of origin, ~~or publication status~~ and studies could be included regardless of whether
182 they had been published **(AUTHOR: please would you clarify what you mean by 'status' here?)**.

183 The electronic databases used were MEDLINE, EMBASE and the Cochrane Library. Additional
184 studies were identified through review of the references of the retrieved papers, contact with
185 study authors, and online searching of Google and Google Scholar (Google Inc., Mountain View,
186 CA, USA). The search strategy used for MEDLINE and EMBASE is given in the **Supplementary**
187 **Data (AUTHOR: we tend not to include Appendices, therefore these data have been divided and**
188 **renamed as supplementary files.)**.

Commented [BDA4]: Reworded "publication status for clarity"

Formatted: Font: Bold

Commented [BDA5]: Thank you

189

190 **Data collection and analysis**

191

192 *Selection of studies*

193 The titles and abstracts obtained were screened by one author (DB). To verify the process of
194 selection, the titles and abstracts for the year 2000 were independently screened by DB and AA,
195 with full agreement with respect to papers to be included or excluded. Thus full text versions of
196 all relevant papers were retrieved by DB for further scrutiny by both reviewers. Any
197 disagreement between the two as to which studies to include was resolved by discussion.

198

199 *Data collection process*

200 Data were extracted by one reviewer (DB), and checked by a second (AA). Discrepancies were
201 resolved by discussion between the reviewers, and a third reviewer (SB) was available for
202 referral as necessary. Study authors were contacted as required for missing data or clarification.

203

204 For each included study, the information collected included study design; methods; location,
205 setting and time period; information about the participants (demographics, infertility diagnosis,
206 BMI, eligibility criteria); drop-outs; the types of interventions assessed and their descriptions;
207 and outcomes, including pregnancy, live births, weight or BMI change, miscarriage, ovulation
208 and menstrual change. For men, semen analysis parameters were captured. (AUTHOR: please
209 would you re-state those outcomes here, for clarity? Thank you.). Any data on barriers to weight
210 loss were also captured. Data on costs and cost-effectiveness were sought but not found.

211

212 *Assessment of risk of bias in individual studies*

213 RCTs were assessed for their methodological quality using the Cochrane Risk of Bias Tool
214 (Higgins *et al.*, 2011).

215

Formatted: Font: Bold

Commented [BDA6]: Done

216 Non-randomised studies were assessed using a checklist developed for the Review Body for
217 Interventional Procedures (ReBIP) (64) (AUTHOR: please would you use the correct format
218 citation?). The ReBIP is an independent review body that carries out systematic reviews for the
219 National Institute for Health and Care Excellence's Interventional Procedures Programme. The
220 checklist was adapted from several sources (Verhagen *et al.*, 1998; Khan *et al.*, 2001; Downs
221 and Black, 1998), and has been used in many systematic reviews. It includes assessment of
222 sample selection; clarity of inclusion and exclusion criteria; baseline comparability of
223 participants; consecutive selection of patients; prospective data collection; clarity of
224 intervention descriptions; experience of person administering the intervention;
225 appropriateness of staff, place and facilities where participants were treated; consideration of
226 important incomes on clinical effectiveness, cost-effectiveness, or learning curves; use of
227 objective outcome measures; blinding of outcome assessment; adequacy of duration of follow-
228 up to detect important effects on outcomes of interest; information provision on dropouts, and
229 their similarity to completers; identification of important prognostic factors; and adjustment of
230 analyses for confounding factors.

231

232 *Summary measures*

233 Results were summarised descriptively. In the outcome summary tables, numbered rankings
234 are provided in an effort to demonstrate which study interventions reported more favourable
235 outcomes, with a lower number indicating a better outcome. Meta-analysis was undertaken
236 where possible. Risk ratios (RR) and mean differences with 95% CI between groups are
237 reported.

238

239 *Synthesis of results*

240 Data were imported into Review Manager Version 5.3.5 (The Cochrane Collaboration, Oxford,
241 UK) for quantitative synthesis. A random effects, rather than a fixed effect model, was used for

Formatted: Font: Bold

Commented [BDA7]: Citation not required. Sources mentioned below

242 meta-analysis, owing to unavoidable population and intervention heterogeneity, as anticipated
243 in interventions for weight loss.

244

245 *Assessment of heterogeneity*

246 Statistical heterogeneity was investigated by visual inspection of forest plots, the Q-test (with
247 $p < 0.1$ implying statistical heterogeneity) and by examining the I^2 statistic (Higgins and
248 Thompson, 2002), where a result $> 50\%$ was taken to indicate substantial statistical
249 heterogeneity.

250

251 *Sensitivity analyses*

252 Sensitivity analysis was planned to compare randomised trials judged to be at low risk of bias
253 for allocation concealment or randomisation with those judged to be at higher risk of bias;
254 however, the small numbers of trials precluded this.

255 **Results**

256

257 ***Outcome of the search***

258

259 Details of the selection process for studies is summarised in the PRISMA flow diagram (Figure

260 1) **(AUTHOR: the quality of this figure still seems poor. If possible, please can you send to me a**

261 **higher quality file?** Seventeen study authors were contacted for clarification or further

262 information; ten provided helpful further information.

263

264 Full details of all the studies and interventions are provided in Supplementary Table SI. There
265 were 14 RCTs, six non-randomised studies with comparison groups (NRCTs), and twenty cohort
266 studies with interventions. Ten studies took place in Australia, seven in the USA, three each in
267 Italy and the Netherlands, and two in Denmark. One each came from Belgium, Brazil, Canada,
268 Egypt, France, Germany, India, Iran, Jordan, Pakistan, Russia, Saudi Arabia, Slovakia, Turkey and
269 the UK. Three studies were multi-centre.

270

271 Enrolment varied from eight participants (Faure *et al.*, 2014) to 577 (Mutsaerts *et al.*, 2016).

272 Attrition rates ranged from 10.6% (Mutsaerts *et al.*, 2016) to 66.7% (Turner-McGrievy *et al.*,
273 2014).

274

275 ***Description of populations***

276

277 Mean age of study groups varied from 25 years (Lazurova *et al.*, 2004) to 35.4 years (Chavarro
278 *et al.*, 2012). Ethnicity was reported in six papers, with between 55% (Turner-McGrievy *et al.*,
279 2014) and 87.5% white (Mutsaerts *et al.*, 2016).

280

Formatted: Font: Bold

Commented [BDA8]: I have re-entered the data into the online prisma diagram generation tool, and will attach all file types so that the best quality file can be used

281 Duration of infertility, where reported, ranged from a mean of 19.5 months (Palomba *et al.*,
282 2008) to 11 years (Aliyeva *et al.*, 1993). Some studies addressed women with any infertility
283 history. Others centred on women undergoing fertility treatments or planning to start
284 immediately following the intervention. Some were exclusively catered to persons with
285 polycystic ovary syndrome (PCOS), with the syndrome defined by Rotterdam 2004 consensus
286 criteria (ESHRE and Group, ASRM-Sponsored PCOS Consensus Workshop, 2004) in ten cases,
287 National Institutes of Health (NIH) criteria (Zawadzki and Dunaif, 1992) in one case, ultrasound
288 appearance of the ovaries in two cases, and not specified in five cases. One study used PCOS as
289 an exclusion criterion (Moran *et al.*, 2003).

290

291 In eight studies, patients were anovulatory, and in seven studies patients had irregular
292 menstrual cycles. Three papers were centred on patients who were resistant to clomiphene
293 citrate for ovulation induction.

294

295 The lowest mean BMI was 24.5 kg/m² (Chavarro *et al.*, 2012), and the highest was 44 kg/m²
296 (Mahoney, 2014; Hakonsen *et al.*, 2011).

297

298 Two studies were centred on men. One had participants with sperm DNA fragmentation who
299 were overweight or obese (Faure *et al.*, 2014), while the other examined obese men only
300 (Hakonsen *et al.*, 2011). One study examined a weight loss intervention in couples (Homan *et*
301 *al.*, 2012).

302

303 ***Description of interventions***

304

305 Table 1 provides brief details of the types of interventions tested. In general, interventions were
306 poorly described, with insufficient detail to allow their replication. Where described, most
307 reducing diets followed healthy eating principles by reducing fat and refined carbohydrate

308 intake. Where available, studies described calorie intakes ranging from 1000 to 2000kcal/d
309 (depending on body size), apart from three studies that tested very low calorie diets as part of
310 their reducing diets (Tsagereli 2006, Kiddy 1992, van Dam 2004) and one that tested a very low
311 carbohydrate ketogenic diet (Mavroupolos 2005). If physical activity advice or programmes
312 were incorporated into interventions, they were generally for aerobic exercise. There was no
313 clear evidence that any particular regime was more effective for weight loss, providing there
314 was a weight reduction diet prescribed.

315
316 Five of the 16 interventions incorporating both a dietary and physical activity component
317 provided some part of the intervention in a group setting. Only two interventions were
318 described involving couples. One study (Chavarro *et al.*, 2012) tracked participants' weights
319 after administering a lifestyle questionnaire. While this is not strictly a weight-loss intervention,
320 we have included it with the other cohort studies, because there is evidence that weight loss
321 occurs with enrolment in studies tracking weight, even with minimal intervention (Waters *et al.*,
322 2012; Johns *et al.*, 2016).

323

324

325 **Quality of studies**

326

327 The quality of the studies included was variable. For full details of risk of bias assessment for ~~all~~
328 the randomised studies please see Supplementary Table SII. ~~The quality of the studies included~~
329 ~~was variable.~~ Three of fourteen RCTs had four or five out of seven domains with low risk of bias,
330 e.g. allocation concealment, incomplete outcome data, but the Lifestyle trial (Mutsaerts *et al.*,
331 2016) stood out as the one with best quality. All studies were of high risk of bias with respect to
332 blinding of participants to treatment allocation, and all had low risk of bias in at least two
333 domains. Five of 26 non-randomised studies, having eight to nine of 14 domains with low risk of

334 bias, were judged to be good quality. [Full details of the risk of bias assessment for the non-](#)
335 [randomised studies are given in Supplementary Table SIII.](#)

336

337 ***Studies involving female infertility participants***

338

339 *Pregnancy*

340 Aggregated data from six RCTs show that women who received a combination of a reducing diet

341 and exercise had higher pregnancy rates than those who underwent standard **care** **(AUTHOR: is**
342 **there a word missing? Please would you recheck the sense here? Thank you.)**, 238/434 (54.8%)

343 versus 221/443 (49.9%); RR 1.59, 95% CI (1.01, 2.50) (Figure 2). The trial by Mutsaerts *et al*

344 (2016)) contributed significantly to the heterogeneity, $I^2=71%$ and its exclusion in a post-hoc

345 sensitivity analysis reduced I^2 to 0%. Although the numbers of participants in other RCTs were

346 too few for any meaningful conclusions to be drawn, a very small trial was unable to

347 demonstrate any advantage of diet alone, while another suggested that orlistat might be

348 beneficial. The results of an NRCT (Clark *et al*, 1998) support the result of the meta-analysis

349 favouring diet and exercise. The study by Koning *et al* (2015) did not show any advantage

350 associated with motivational interviewing.

Formatted: Font: Bold

Commented [BDA9]: Added the word 'care'

351 Table 2 details the reproductive outcomes for all included studies. Diet and exercise studies
352 without control populations achieved rates that were comparable to those included in the meta-
353 analysis.

354

355 *Live births*

356 Data from five RCTs (Figure 3) did not favour diet and exercise over standard care in terms of
357 live birth rates which were 48.9% (195/399) versus 46.8% (190/406) respectively; RR 1.54,
358 95% CI (0.93, 2.56). Mutsaerts et al (Mutsaerts *et al.*, 2016) again contributed significantly to
359 the heterogeneity, $I^2=69\%$. Its exclusion in a post-hoc sensitivity analysis reduced I^2 to 0%, and
360 resulted in diet and exercise having a statistical advantage, RR 1.86, 95% CI (1.25, 2.77). The
361 NRCT (Clark *et al.*, 1995; Clark *et al.*, 1998) also favoured the intervention arm, 45/67 (67.2%)
362 versus no treatment (0/20); RR 28.10, 95% CI (1.81, 436.85). Motivational interviewing was
363 also not associated with an increased live birth rate.

364

365 Table 2 details live birth rates for all included studies. Women in studies on diet and exercise
366 without control groups had lower success rates than those included in the meta-analysis.
367 Khaskheli et al (Khaskheli *et al.*, 2013), who did not provide details of their intervention,
368 reported a 37.6% (32/85) livebirth rate, while Crosignani et al, (Crosignani *et al.*, 2003), who
369 administered a 1200 kcal/day diet along with aerobic exercise, achieved a 24.2% (8/33) rate.
370 The numbers of participants in the other studies included in Table 2 were too few for any
371 meaningful conclusions to be drawn.

372 *Spontaneous and IVF conceptions*

373 Pooled data from RCTs (Figures 4 and 5) show that, reducing diets and exercise were not
374 associated with a higher chance of spontaneous conception [28.3% (86/304) versus 15.9%
375 (50/315); RR 2.20, 95% CI (0.98, 4.93)].

376

377 *Miscarriages*

378 A combination of a reducing diet and exercise was associated with a pooled miscarriage rate of
379 13.4 per hundred women (48/357) or 23.3% of pregnancies (48/206) (Figures 6 and 7). The
380 control arm had a pooled miscarriage rate of 8.7 per hundred women (31/355) or 15.6% of
381 pregnancies (31/199). This shows no benefit of diet and exercise [combined RR 0.96, 95% CI
382 (0.89, 1.04) per woman and RR 0.91, 95% CI (0.91, 0.82, 1.01) per pregnancy]. Miscarriage rates
383 in the other studies on diet and exercise vary widely (Table 2).

384

385 *Ovulation*

386 One RCT (Palomba *et al.*, 2010) found a 37.5% (12/32) ovulation improvement with a
387 combination of reducing diet and exercise, while its control group had a 9.3% (3/32)
388 improvement rate; RR 4.00 95% CI (1.25, 12.84) (Figure 8). An NRCT, also using diet and
389 exercise, ~~that~~ had the highest rate of improvement of all included studies in its intervention
390 group. [AUTHOR: again, does this sentence read correctly? Please would you recheck?], 89.6%
391 (60/67), while none of the participants in the control group saw improvement; RR 37.37, 95%
392 CI (2.41, 578.65).

393

394 One RCT (Kumar and Arora, 2014) compared the use of 120mg of orlistat twice daily and a
395 lifestyle modification programme with use of the lifestyle programme alone. The orlistat group
396 had a 50% (15/30) rate of ovulation, while the control group had a 6.7% (2/30) rate; RR 7.50,
397 95% CI (1.88, 29.99).

398

Formatted: Font: Not Bold, Not Italic

Commented [BDA10]: Removed the word 'that' before had

Formatted: Font: Bold

399 One RCT (Legro *et al.*, 2015) compared the proportions of total clomiphene treatment cycles
400 that were ovulatory in each group. The group whose treatment was preceded by diet and
401 exercise advice and the oral contraceptive pill (OCP) had the best ovulation rate, 67.1% (94/140),
402 while the lifestyle alone group and control (OCP alone) group had rates of 60.3% (82/136) and
403 46.1% (71/154) respectively.

404
405 Another RCT (Thomson *et al.*, 2008) randomised its participants, who had PCOS, to receive an
406 energy-restricted high-protein diet alone (DO), combined with aerobic exercise (DA), or
407 combined with both aerobic and resistance exercise (DC). DO saw a 50% (6/12) improvement
408 in ovulation, DA a 50% (3/6) improvement, and DC a 42.8% (3/7) improvement.

409
410 Table 2 shows the ovulation rates in cohort studies of diet without controls. One study using a
411 very low calorie diet (van Dam *et al.*, 2004) had a relatively good ovulation improvement rate,
412 60% (9/15). Other studies had comparatively low rates, including one using a hypocaloric 1434
413 kcal/day diet (Thomson *et al.*, 2009) which achieved a 19.2% (10/52) rate, and another
414 (Turner-McGrievy *et al.*, 2014) which saw no improvement using either a low-calorie diet
415 [LCD] (**AUTHOR: please define LCD**) or a vegan diet. One study compared a hypocaloric high
416 protein diet to structured exercise training (SET) (Palomba *et al.*, 2008), and achieved a 25%
417 (5/20) improvement rate with diet, and a 65% (13/20) improvement with exercise.

418
419 A study comparing ovulation improvement in women receiving acupuncture to those receiving
420 an unspecified diet (Aliyeva *et al.*, 1993) had a 55% (11/20) improvement with the former, and
421 46.7% (28/60) improvement in the latter; RR 1.18, 95% CI (0.73, 1.90).

422
423 *Menstrual irregularity*

424 In clomiphene citrate resistant PCOS women, a single trial (Palomba *et al.*, 2010), showed that
425 hypocaloric diet and exercise was significantly more likely to result in menstrual improvement,

Commented [BDA11]: Placed full nomenclature "low calorie diet" ahead of the abbreviation LCD

Formatted: Font: Bold

426 than observation only: 34.4% (11/32) versus 9.3% (3/32); RR 3.67, 95% CI (1.13, 11.92)
427 (Figure 9). Other studies without control groups also suggest that diet and exercise were
428 associated with improvement in menstrual regularity (Table 2). There were no RCTs evaluating
429 the role of dietary interventions alone or motivational interviewing in regulating menstrual
430 cyclicity. Data from observational and often uncontrolled studies of other interventions show
431 variable levels of success (Table 2).

432

433 *Oocyte retrieval*

434 Data from a single RCT (Figure 10) suggest that a low glycaemic index diet (Becker *et al.*, 2015)
435 could improve oocyte yield in IVF. The mean (SD) number of oocytes in women in the
436 intervention arm was 7.75 (5.39), compared to 4.18 (SD 3.01) in controls; mean difference -
437 3.57, 95% CI (-6.87, -0.27).

438

439 *Time to conception*

440 The Lifestyle trial (Mutsaerts *et al.*, 2016) reported that the median time to conception (TTC)
441 resulting in a term live birth was 7.2 months (interquartile range (IQR) 2.6, 12.0) in the
442 intervention group, who underwent a multidisciplinary programme based on diet and exercise,
443 versus 5.2 (IQR 2.4, 10.1) in the control group, who were allowed fertility treatment from the
444 start (P=0.06). An RCT comparing energy-restricted low protein and high protein diet groups
445 who both received exercise instruction (Moran *et al.*, 2003) reported a 4-5 week TTC regardless
446 of group. A cohort study which advised women to lose 10% of their body weight (Kort *et al.*,
447 2013; Kort *et al.*, 2014) found that those who met the target had a mean TTC of 227 days, while
448 those who had not had a mean of 231 days.

449

450 *Weight loss*

451 Aggregated data from RCTs suggest that a combination of a reducing diet and exercise resulted
452 in greater weight loss than that achieved in control groups (Figure 11); mean difference -3.98 kg

Formatted: Font: Not Bold, Not Italic

453 (95% CI -4.85, -3.12). Diet alone was also shown to be more effective in a small RCT; mean
454 difference -5.23kg (95% CI -7.42, -3.04) (Becker *et al.*, 2015).

455

456 In non-randomised trials, Clark et al (Clark *et al.*, 1998) reported significantly greater weight
457 loss from diet and exercise; mean difference -9.00kg (95% CI -10.88, -7.12). Two non-
458 randomised trials testing motivational interviewing also found greater weight loss; mean
459 difference -3.21kg (95% CI -5.93, -0.49kg). Changes in weight, BMI and waist circumference for
460 other studies are provided in Table 3. Generally results are not inconsistent with studies with
461 either randomised or non-randomised control groups.

462

463 ***Comparisons with metformin***

464

465 *Pregnancy*

466 Three small RCTs evaluated lifestyle interventions versus metformin (Figure 12). None was able
467 to show any superiority of metformin over the alternative intervention, but it is difficult to be
468 confident about these outcomes owing to the small sample sizes involved.

469

470 *Ovulation*

471 Two small RCT of diet or drug treatment (Figure 13) were unable to show any benefit over
472 metformin. Details of other studies are in Table 2.

473

474 *Menstrual irregularity*

475 Metformin did not appear to improve menstrual regularity in a number of small trials
476 comparing it to diet and exercise with or without drug treatment, diet alone or acarbose (Figure
477 14). Details of other studies are in Table 2.

478

479 *Weight loss*

480 Weight, BMI and waist circumference changes for studies comparing metformin with lifestyle
481 interventions are provided in Table 3.

482 **AUTHOR: please would you mention Supplementary Table SIII (Quality assessment for non-**
483 **randomised studies) at the appropriate place in the Results section? (Please renumber the**
484 **Supplementary Tables if order in text requires it.) Thank you.**

Formatted: Font: Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Commented [BDA12]: Lines 328-330

485 **Summary of the key findings in infertile women**

486

487 In overweight and obese infertile women, weight loss is associated with improved chances of
488 becoming pregnant, and possibly spontaneous conception pre-IVF. Ovulation and menstrual
489 irregularity are also aided. There appears to be no significant difference between weight loss
490 interventions and controls with respect to rates of miscarriage and IVF conceptions. The few
491 studies that included metformin showed no significant difference between weight loss
492 interventions and the drug for achieving pregnancy or improvement in ovulation status;
493 similarly there was no difference between the two groups in menstrual improvement.

494

495 **Barriers to weight loss in infertile women**

496

497 Very little has been reported regarding the perceived barriers to weight loss in overweight and
498 obese infertile women. Two of the included studies attempted in some way to evaluate these.
499 In the trial by Thomson et al (Thomson *et al.*, 2008), 104 participants were allocated to receive
500 an energy restricted diet only (DO), the same diet combined with five sessions of aerobic
501 walking and jogging exercise per week (DA), or the diet combined with aerobic and resistance
502 training exercises (DC), where resistance exercise replaced two of the aerobic sessions (110)

503 **(AUTHOR: please would you use the correct format for the citation? Thank you.)** Forty-three of
504 the participants in the trial (DO=13, DA=11 and DC=19) completed a validated Exercise
505 Benefits/Barriers Scale at weeks 0, 10 and 20 (Thomson *et al.*, 2016). This was to determine
506 their perceptions of the benefits and barriers to participation in exercise. The scale included 43

Commented [BDA13]: Second citation removed as wasn't required.

Formatted: Font: Bold

507 items with statements related to ideas about exercise and a four-point Likert scale, with 4 =
508 strongly agree, 3 = agree, 2 = disagree and 1 = strongly disagree. At baseline, the statements
509 with the highest agreement were: “exercise tires me” [mean (SD), 2.86 (0.71)]; “exercise is hard
510 work for me [2.79 (0.67)], and “I am fatigued by exercise” [2.60 (0.62)]. Those with the lowest
511 agreement were: “my family members do not encourage me to exercise” [1.56 (0.59)], “my
512 spouse does not encourage exercising” [1.63 (0.70)], and “I think people in exercise clothes look
513 funny” [1.72 (0.59)].

514
515 Barrier scores were reduced by week 10 in this study, regardless of treatment allocation, with
516 no further reduction by week 20 ($P \leq 0.001$). Time expenditure and physical exertion as
517 perceived barriers saw a significant reduction ($P \leq 0.003$), while family discouragement saw no
518 change ($P = 0.6$).

519
520 Participants in the study by Galletly et al (Galletly *et al.*, 1996), in addition to completing a 24-
521 week programme with weekly meetings comprising 1 hour of exercise and 1 hour of dietary and
522 psychiatric counselling, completed the Rosenberg Self-Esteem Scale and the General Hospital
523 Anxiety and Depression Scale at the start and end of the study. At baseline, those who dropped
524 out before the end of intervention exhibited higher anxiety ($p < 0.02$) and depression scores
525 ($p < 0.069$), and lower self-esteem scores ($p < 0.056$) than those who completed. Completion was
526 associated with a significant improvement in self-esteem ($p < 0.0001$) and depression ($p < 0.006$),
527 and a non-significant reduction in anxiety scores.

528
529 Based on what little evidence is available, overweight infertile women appear most deterred
530 from exercise by the perception that it causes fatigue and is hard work. These perceptions, as
531 well as depression, seem to decrease with continuation of an exercise programme. Very few
532 women seem to be discouraged from exercise by family members. Those women who
533 discontinue exercise would appear to be more anxious and/or depressed at the outset.

534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560

Studies involving male infertility participants

Achievement of pregnancy, live birth rate, and Improvement in sperm DNA integrity

One cohort study (Faure *et al.*, 2014) administered a nutritionist-led personalised dietary programme of an unreported nature coupled with exercise, aimed at reducing intra-abdominal fat over a 3-8 month period, to eight men who had significant abdominal fat and at least 25% sperm DNA fragmentation and unexplained infertility (Table 4). The partners of all eight achieved pregnancy, three naturally and five through IUI, and all went on to have live births. Six who allowed their semen samples to be analysed both before and after the intervention all had an improvement in the degree of sperm DNA fragmentation.

AUTHOR: could the following sections be edited to avoid repeating most of the data that are shown in Table 4, as we try to minimise such repetition in the results section? Thank you,

Improvement in semen analysis parameters

A cohort of 44 men with BMI 33-61 kg/m² underwent a programme based on a healthy diet and daily exercise (Hakonsen *et al.*, 2011). Semen analyses before and afterward were compared in 27 men, and the results were as shown in Table 4.

The group with 3.5-12.1% weight loss (Group 1, n=9) saw a mean change in sperm concentration of -11 million/ml (95% CI -49, +27). The group with 12.2-17.1% (Group 2, n=9) loss had a mean change of +19 million/ml (95% CI -23, 61). The group with 17.2-25.4% loss (Group 3, n=9) had a mean change of +17 million/ml (95% CI -24, 58).

Group 1 saw a mean change in progressive sperm motility of -2%, 95% CI (-15,11). Group 2 had a mean change +4%, 95% CI (-10, 18). Group 3 had a mean change +11%, 95% CI (-3, 25).

Formatted: Font: Bold
Formatted: Normal, Line spacing: single
Formatted: Font: Bold

561 ~~Group 1 saw a mean change in normal sperm morphology of 0%, 95% CI (-2, 4). Group 2 had a~~
562 ~~mean change of +1%, 95% CI (-3, 4). Group 3 had a mean change of +4%, 95% CI (1,7).~~

563

564 *Change in weight*

565 ~~The cohort study by Hakonsen et al (Hakonsen et al., 2011) achieved median weight change~~
566 ~~22kg, range -4 to 39 kg. The cohort study by Faure et al (Faure et al., 2014) achieved a mean~~
567 ~~weight reduction of 4.40 (SD 5.0 kg). Table 4 also summarises the weight loss achieved in both~~
568 ~~studies.~~

569

570 *Summary*

571 The relative paucity of studies in overweight and obese men seeking fertility has made it
572 difficult to draw firm conclusions regarding the benefits of weight loss. However, improvements
573 in sperm concentrations, motility and normal morphology have been suggested, though these
574 have not been statistically significant. Improvement in sperm DNA integrity has also been
575 suggested, and this might be linked to an improved live birth rate.

576

Formatted: Normal

577 **Discussion**

578

579 **Key findings**

580

581 The present systematic review and meta-analysis found that the weight loss interventions,
582 particularly diet and exercise, improved pregnancy rates and ovulatory status. A trend toward
583 more natural pregnancies, but not IVF pregnancies, occurred. Miscarriage rates were unaffected
584 by weight loss interventions and, as reports of higher oocyte yield and improved menstrual
585 regularity are based on single trials, further research is needed before firm conclusions can be
586 made. A lack of RCTs in men and couples, and of studies evaluating barriers to undertaking
587 weight loss in overweight and obese infertile populations, is evident. The quality of the studies
588 we included was variable, but the one with the lowest risk of bias was the Lifestyle trial
589 (Mutsaerts *et al.*, 2016), whose intervention also ranked highly for both pregnancy and live
590 birth rates.

591

592 **Women**

593

594 *Pregnancies and live births*

595 Weight loss interventions, particularly those with reducing diets and exercise, were more likely
596 than controls to result in pregnancy. Live births were reported by ~~only a few~~ relatively fewer
597 studies. **[AUTHOR: correct?]**, usually involving women undergoing fertility treatments such as
598 OI or IVF. Data from studies predating the Lifestyle trial in 2016 (Mutsaerts *et al.*, 2016)
599 suggested that diet and exercise combinations were superior to standard care, but inclusion of
600 this trial changed this. Those in the intervention arm underwent a 6-month programme based
601 on diet and exercise, prior to undergoing 18 months of fertility treatments. Those in the control
602 arm commenced fertility treatments immediately. The primary outcome was vaginal birth of a
603 healthy singleton of at least 37 weeks' gestation within 24 months after randomisation. The trial

Commented [BDA14]: Reworded.

Formatted: Font: Bold

604 was well conducted, with low risk of bias. This was the largest study in the review, and it carried
605 substantial weight during meta-analysis. The results deviated from the trend exhibited by the
606 preceding trials, resulting in marked increase in heterogeneity. It is possible that the results
607 might have been different if the control group underwent 6 months of no intervention
608 observation prior to 18 months of treatment. The trial found that significantly more women in
609 the intervention group had ongoing pregnancies from natural (AUTHOR: the term natural' is
610 usually preferred throughout but I leave the final decision to you.) conception than the control
611 group, consistent with our other findings.

Formatted: Font: Bold

Commented [BDA15]: I'll accept your suggestion

612
613 Studies reporting the highest pregnancy and live birth rates were not necessarily those with
614 highest weight loss. Clark et al, had a high live birth rate of 67.2% in the diet and exercise arm of
615 their NRCT, while also achieving one of the highest weight changes, mean -10.2kg (SD 4.3)
616 (Clark *et al.*, 1995; Clark *et al.*, 1998). On the other hand, the intervention arm of the Mutsaerts
617 et al (2016) RCT reported a live birth rate of 53.2%, but only achieved a mean weight change of
618 -4.4 kg (SD 5.8).

619
620 *Miscarriages*

621 Studies with some of the lowest rates of weight loss reported fewer miscarriages in their
622 intervention arms, but the data are insufficient to draw conclusions (Tables 2 and 3).

623
624 *Ovulation*

625 Weight loss interventions, regardless of their nature, were significantly more likely than control
626 interventions to achieve ovulation in anovulatory women. Improvements were not necessarily
627 greatest in those studies achieving the greatest weight loss.

628

629 *Menstrual irregularity*

630 A single RCT suggested that diet and exercise were significantly more likely to improve
631 menstrual cycles than a control intervention (Palomba *et al.*, 2010). High rates of menstrual
632 improvement were seen in studies with good weight or BMI reduction. Qublan et al (2007), for
633 example, with a 1200-1400 kcal diet in their RCT, had a 61.9% menstrual improvement and
634 achieved one of the better BMI reductions, with a mean change of -4.8 kg/m².

635

636 *Spontaneous and IVF conceptions*

637 Studies included women expected to proceed to in-vitro treatments after a period of weight
638 loss. Many women were able to conceive without further assistance through weight loss alone.
639 All interventions showed an improved rate of spontaneous conception in comparison to a
640 control. There was however no significant difference between interventions and controls when
641 it came to IVF conceptions. Similar spontaneous conception rates were seen regardless of type
642 of intervention and/or amount of weight lost.

643

644 *Change in weight, BMI and waist circumference*

645 No class of intervention was consistently better in achieving reduction in weight compared to
646 others. Trials with longer interventions and follow-up sometimes suffered from higher rates of
647 study drop-out, for example 20.4% in the 12-month Sim et al RCT (Sim *et al.*, 2014a) and 10.6%
648 in the 6-month Mutsaerts et al RCT (Mutsaerts *et al.*, 2016), both of which mainly had diet and
649 exercise. Others had no dropout: Karlsen et al after 7-8 months (Karlsen *et al.*, 2013) and
650 Koning et al after 6 months (Koning, 2015), and both included motivational interviewing. This
651 suggests that enhancing motivation plays a key role in compliance. Programmes with strict
652 dietary and exercise modifications may be more likely to see discontinuation. The study with
653 the most weight lost, the cohort study by Mavropoulos et al (Mavropoulos *et al.*, 2005), utilised

654 a very restrictive diet for 6 months, and had very low recruitment (n=11) and a high rate of
655 drop-out (58.5%).

656

657 *Metformin*

658 The meta-analyses showed that weight loss interventions have a non-significant advantage over
659 metformin with respect to achievement of pregnancy or improvement of ovulation status. There
660 was also no significant difference in menstrual regularity improvement. In light of these
661 findings, and the gastro-intestinal side-effects common with metformin (Kumar and Arora,
662 2014; Qublan *et al.*, 2007), lifestyle interventions should remain the first line therapy for
663 improvement in ovulation and menstruation.

664

665 *Interventions in infertile men*

666

667 There was very little evidence available to judge the effectiveness of weight reduction strategies
668 in men, and this is an area ripe for future research. The two studies both utilised a diet and
669 exercise combination, with neither providing adequate details of the intervention.

670

671 *Types of diet*

672

673 As expected, the dietary interventions used were based on caloric restriction, and were usually
674 consistent with weight reduction advice from national guidelines (National Clinical Guideline
675 Centre, 2014; Dietz *et al.*, 2015). It is hard to say, based on the data obtained, whether any
676 particular degree of restriction was superior for achievement of weight loss, particularly given
677 the frequent pairing of diet with exercise. Some of the RCTs used individualized diets, tailoring
678 caloric deficit to baseline weight (Becker *et al.*, 2015; Duval *et al.*, 2015b). Others mentioned a
679 hypocaloric range within which dietary intakes would fall (Kumar and Arora, 2014; Qublan *et*
680 *al.*, 2007; Thomson *et al.*, 2008; Turner-McGrievy *et al.*, 2014), generally between 1200 and

681 1800 kcal/day. Dieticians were used to assist with dietary advice in a few cases (Duval *et al.*,
682 2015b; Turner-McGrievy *et al.*, 2014; Thomson *et al.*, 2008; Mutsaerts *et al.*, 2016). The RCT
683 with the lowest recruitment and highest discontinuation rate used a vegan diet in one of its
684 study arms (Turner-McGrievy *et al.*, 2014), suggesting poor acceptance.

685

686 ***Types of exercise***

687

688 The majority of exercise interventions sought to increase weekly aerobic activity in participants
689 in order to increase caloric expenditure. The most frequent feature seen in the RCTs was an
690 increase in the number of steps or amount of walking by participants (Duval *et al.*, 2015b;
691 Karimzadeh and Javedani, 2010; Legro *et al.*, 2015; Mutsaerts *et al.*, 2016; Sim *et al.*, 2014a;
692 Thomson *et al.*, 2008). Fewer RCTs (**AUTHOR: correct, or studies?**) included strength or
693 resistance training (Karimzadeh and Javedani, 2010; Moran *et al.*, 2011b; Thomson *et al.*, 2008),
694 and this was always in addition to the basic aerobic approach. Thomson *et al.* found that
695 addition of resistance training did not result in a significantly different weight loss in their RCT
696 of diet alone, diet with aerobic exercise, or diet with aerobic and resistance exercise (Thomson
697 *et al.*, 2008). Physical activity interventions were in some cases individualised (Duval *et al.*,
698 2015a; Moran *et al.*, 2011a) and in others in the form of group sessions (Moran *et al.*, 2003),
699 with occasional use of a kinesiologist (Duval *et al.*, 2015a) or physiotherapist (Kumar and
700 Arora, 2014).

701

702 ***Types of weight loss medication***

703

704 The use of weight loss drugs is contraindicated during pregnancy (Department of Veterans
705 Affairs, 2014), and the studies using them aimed to reduce weight prior to attempting
706 pregnancy. One trial (Legro *et al.*, 2015) recommended barrier contraception during its
707 intervention. Oral medications used in the studies included orlistat, a lipase inhibitor;

Commented [BDA16]: Correct.

Formatted: Font: Bold

708 sibutramine, a selective serotonin and norepinephrine reuptake inhibitor; and acarbose, an
709 alpha-glucosidase inhibitor shown to induce modest weight loss, though not suitable for weight
710 maintenance (Hauner *et al.*, 2001). Of these, sibutramine, which has been withdrawn in Europe
711 and the USA but is still available on the internet, has been shown in a large study to have a risk
712 of cardiovascular defects in unborn infants (Källén, 2014), while the same study showed no risk
713 of birth defects from orlistat use. The safety of acarbose in pregnancy is not established. Orlistat
714 was shown to be superior to a control with respect to achievement of pregnancy and ovulation
715 in a single study (Kumar and Arora, 2014). Until further evidence is available, lifestyle
716 interventions should still be considered the first line therapy, with drug use reserved for
717 monitored trials.

718

719 ***Strengths and limitations of the review***

720

721 This review has added to the scope of the systematic review by Sim *et al.* (2014b) by going
722 beyond examining overweight and obese women undergoing fertility treatment to encompass
723 individuals of both genders, and couples, from a variety of infertility circumstances undergoing
724 non-surgical weight loss programmes. A wide search strategy meant that we were thus able to
725 capture 40 studies, in comparison to the 11 in that review (Sim *et al.*, 2014b). Author contact,
726 initiated with 17 study authors, was another strength, as this allowed unreported information
727 to be gained, and clarification of unclear information. Nevertheless, we cannot exclude
728 publication bias, where studies with less positive outcomes remain unreported.

729

730 ***Clinical recommendations***

731

732 Overweight and obese persons seeking fertility should be educated on the effects of being
733 overweight or obese on the ability to achieve pregnancy, and the benefits of weight reduction,
734 including improvement in pregnancy rates, and a reduced need for OI and IVF.

735
736 A combination of a reduced calorie diet, which is not overly restrictive, and aerobic exercise,
737 intensified gradually, should form the basis of programmes designed for such individuals. As
738 compliance is key to success, coached sessions of achievable frequency, e.g. weekly, for up to 6
739 months, should be considered. Motivational interviewing techniques might also be useful. An
740 advantage of fertility care is that couples are catered to, and dual enrolment may result in better
741 adherence, as partners tend to motivate each other.

742

743 **Research recommendations**

744

745 Future research would ideally be in the form of large multi-centre RCTs including both women
746 and men, ideally couples, attending fertility clinics and having at least one partner with a BMI
747 within the overweight or obese range. Based on the data from RCTs of diet and exercise
748 examined here, we estimate that a trial would require over 454 participants to detect a
749 difference of 15% in pregnancy rates (90% power, $p < 0.05$, control pregnancy rate 50%,
750 intervention rate 65%). This figure does not take account of losses to follow-up, and would need
751 to be considerably inflated, depending on drop-out rates in feasibility and pilot work.

752

753 A reasonable control intervention would be the administration of dietary and exercise advice
754 only-, to be undertaken for a brief period ~~only~~ **(AUTHOR: correct?)**, to reduce persons allocated
755 to this arm feeling as though their fertility treatments were being delayed with no intervention.
756 Similar durations employed between intervention and control groups prior to starting fertility
757 treatments would allow for fair comparisons with respect to weight loss and reproductive
758 outcomes.

759

760 A long duration of follow-up, spanning decades, might answer further questions, including
761 whether obstetric problems related to weight such as gestational diabetes or caesarean rates

Commented [BDA17]: Rearranged location of the word 'only'

Formatted: Font: Bold

762 might be reduced in imminent or future pregnancies. The permanency of lifestyle changes made
763 while attempting fertility might also be established, as well as their impact on family members,
764 and the maintenance of weight loss in the long term. ~~The role of epigenetic changes might also~~
765 ~~be explored. (AUTHOR: although important, and you are discussing possible future research~~
766 ~~here, this subject has not been mentioned so far and seems a little unrelated to the overall~~
767 ~~approach here. It is also unclear what particular studies you would perform, therefore I suggest~~
768 ~~removing it, which would not detract from the discussion. If you agree, please edit accordingly?~~
769 ~~Thank you.)~~

Formatted: Font: Bold

Commented [BDA18]: Removed reference to epigenetic studies.

771 **Conclusions**

772
773 Non-surgical weight reduction strategies in the infertile have been shown to improve
774 reproductive outcomes in both men and women. Diet, exercise, a combination of the two,
775 weight-loss medication and motivational interviewing have all been efficacious in reducing
776 obesity. In women, weight loss from diet and exercise is associated with improved chances of
777 becoming pregnant, with a trend toward improved live birth rate. Ovulation and menstrual
778 irregularity are also aided. At present there appears to be no significant difference between
779 weight loss interventions and control interventions with respect to rates of miscarriage and IVF
780 conceptions. Well-designed RCTs should also shed light on the effect of weight loss on numbers
781 of oocytes retrieved and other parameters in IVF cycles, and on TTC, as well as the role of
782 interventions in couples. In men, improvements in sperm concentrations, motility and normal
783 morphology have been suggested, as well as in sperm DNA integrity, but further high quality
784 research is needed to confirm these findings and demonstrate improvement in live birth rates in
785 their partners.

786 **Acknowledgements**

787 Mrs Cynthia Fraser provided useful assistance in the formulation of a search strategy. Authors
788 responding to queries included Gabrielle Turner McGrievy, Ann Van Oers, Kamilla Karlsen, Kyra
789 Sim, Amany Salama, Rebecca Thomson, Cherrie Galletly, and Hanno Pijl.

790 **Authors' roles**

791 Dr Damian Best was involved in design and conduct of the review, data analysis, drafting the
792 manuscript and critical discussion. Professor Alison Avenell was involved in the design and
793 conduct of the review, supervised data analysis, checked data extraction, and was involved in
794 manuscript revision and critical discussion. Professor Siladitya Bhattacharya was involved in
795 the design of the review and contributed to manuscript revision and critical discussion.

796 **Funding**

797 No specific funding was sought. All authors were employed by the University of Aberdeen in a
798 research capacity. The Health Services Research Unit is core funded by the Chief Scientist Office
799 of the Scottish Government Health and Social Care Directorate.

800

801 **Conflict of interest**

802 None declared.

803

804 **References**

805 Anonymous . National Collaborating Centre for Women's and Children's Health (UK). Ectopic
806 Pregnancy and Miscarriage. 2012. RCOG.

807 Aliyeva EA, Fanchenko ND, Parshutin NP, Gasparov AS, Pshenichnikova TI. The effect of a
808 decrease in body weight in patients with the polycystic ovary syndrome. *Akush Ginekol* 1993;**3**; 33-
809 36.

810 Balen A, Dresner M, Scott E, Drife J. Should obese women with polycystic ovary syndrome receive
811 treatment for infertility? *BMJ* 2006a;**332**; 434-435.

812 Balen A, Platteau P, Andersen A, Devroey P, Sorensen P, Helmggaard L, Arce C. The influence of
813 body weight on response to ovulation induction with gonadotrophins in 335 women with World
814 Health Organization group II anovulatory infertility. *BJOG* 2006b;**10**; 1195-1202.

815 Becker GF, Passos EP, Moulin CC. Short-term effects of a hypocaloric diet with low glycemic index
816 and low glycemic load on body adiposity, metabolic variables, ghrelin, leptin, and pregnancy rate in
817 overweight and obese infertile women: a randomized controlled trial. *Am J Clin Nutr* 2015;**6**; 1365-
818 1372.

819 Bellver J, Ayllon Y, Ferrando M, Melo M, Goyri E, Pellicer A, Remohi J, Mesequeur M. Female
820 obesity impairs in vitro fertilization outcome without affecting embryo quality. *Fertil Steril* 2010;**93**;
821 447-454.

822 Bellver J, Martinez-Conejero JA, Labarta E, Alama P, Melo A M, Remohi J, Pellicer A, Horcajadas A
823 J. Endometrial gene expression in the window of implantation is altered in obese women especially in
824 association with polycystic ovary syndrome. *Fertil Steril* 2011;**95**; 2335-2341.

825 Carrell DT, Jones KP, Peterson CM, Aoki V, Emery R B, Campbell R B. Body mass index is
826 inversely related to intrafollicular hCG concentrations, embryo quality and IVF outcome. *Reprod*
827 *Biomed Online* 2001;**2**; 109-111.

828 Chavarro J, Ehrlich S, Colaci D, Wright D, Toth T, Petrozza J, Hauser R. Body mass index and short
829 term weight change in relation to treatment outcomes in women undergoing assisted reproduction.
830 *Fertil Steril* 2012;**1**; 109-116.

831 Clark AM, Thornley B, Tomlinson L, Galletley C, Norman RJ. Weight loss in obese infertile women
832 results in improvement in reproductive outcome for all forms of fertility treatment. *Hum Reprod*
833 1998;**6**; 1502-1505.

834 Clark A, Ledger W, Galletley C, Tomlinson L, Blaney F, Wang X, Norman R. Weight loss results in
835 significant improvement in pregnancy and ovulation rates in anovulatory obese women. *Hum Reprod*
836 1995;**10**; 2705-2712.

837 Crosignani PG, Colombo M, Vegetti W, Somigliana E, Gessati A, Ragni G. Overweight and obese
838 anovulatory patients with polycystic ovaries: parallel improvements in anthropometric indices,
839 ovarian physiology and fertility rate induced by diet. *Hum Reprod* 2003;**9**; 1928-1932.

840 Department of Veterans Affairs. VA/DoD clinical practice guideline for screening and management
841 of overweight and obesity. 2014.

842 Dickey RP, Taylor SN, Curole DN, Rye H P, Lu Y P, Pyrzak R. Relationship of clomiphene dose and
843 patient weight to successful treatment. *Hum Reprod* 1997;**12**; 449-453.

844 Dietz WH, Baur LA, Hall K, Puhl RM, Taveras EM, Uauy R, Kopelman P. Management of obesity:
845 improvement of health-care training and systems for prevention and care. *Lancet* 2015;**9986**; 2521-
846 2533.

847 Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological
848 quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol
849 Community Health* 1998;**6**; 377-384.

850 Duval K, Belan M, JeanDenis F, Baillargeon J. An interdisciplinary lifestyle intervention improves
851 clinically relevant fertility outcomes in obese infertile women-preliminary results. *Fertil Steril*
852 2015a;**3 SUPPL. 1**; e97.

853 Duval K, Langlois M, Carranza-Mamane B, Pesant M, Hivert M, Poder TG, Lavoie HB, Ainmelk Y,
854 St-Cyr Tribble D, Laredo S et al. The Obesity-Fertility Protocol: a randomized controlled trial
855 assessing clinical outcomes and costs of a transferable interdisciplinary lifestyle intervention, before
856 and during pregnancy, in obese infertile women. *BMC Obesity* 2015b;**2**; 47.

857 ESHRE TR, Group, ASRM-Sponsored PCOS Consensus Workshop. Revised 2003 consensus on
858 diagnostic criteria and long-term health risks related to polycystic ovary syndrome. *Fertil Steril*
859 2004;**1**; 19-25.

860 Farquhar C, Gillett W. Prioritising for fertility treatments?should a high BMI exclude treatment?
861 *BJOG* 2006;**10**; 1107-1109.

862 Faure C, Dupont C, Baraibar MA, Ladouce R, Cedrin-Durnerin I, Wolf JP, Lévy R. In subfertile
863 couple, abdominal fat loss in men is associated with improvement of sperm quality and pregnancy: a
864 case-series. *PLoS ONE* 2014;**2**; e86300.

865 Ferlitsch K, Sator O M, Gruber M D. Body mass index, follicle stimulating hormone and their
866 predictive value in in vitro fertilization. *J Assist Reprod Genet* 2004;**21**; 431-436.

867 Galletly C, Clark A, Tomlinson L, Blaney F. A group program for obese, infertile women: Weight
868 loss and improved psychological health. *J Psychosom Obstet Gynaecol* 1996;**2**; 125-128.

869 Gesink Law DC, Macle hose RF, Longnecker MP. Obesity and time to pregnancy. *Hum Reprod*
870 2006;**22**; 414-420.

871 Hakonsen L, Thulstrup A, Aggerholm A, Olsen J, Bonde J, Andersen C, Bungum M, Ernst E, Hansen
872 M, Ernst E et al. Does weight loss improve semen quality and reproductive hormones? Results from a
873 cohort of severely obese men. *Reprod Health* 2011;**1**; 24.

874 Hammiche F, Laven JSE, Twigt JM, Boellaard WPA, Steegers EAP, SteegersTheunissen RP. Body
875 mass index and central adiposity are associated with sperm quality in men of subfertile couples. *Hum
876 Reprod* 2012;**8**; 2365-2372.

877 Hammoud A, Meikle A, Reis L, Gibson M, Peterson C, Carrell D. Obesity and Male Infertility: A
878 Practical Approach. *Semin Reprod Med* 2012;**6**; 486-495.

879 Hartz A, Barboriak P, Wong A, Katayama K, Rimm A. The association of obesity with infertility and
880 related menstrual abnormalities in women. *Int J Obes* 1979;**1**; 57-73.

881 Hauner H, Petzinna D, Sommerauer B, Toplak H. Effect of acarbose on weight maintenance after
882 dietary weight loss in obese subjects. *Diabetes Obes Metab* 2001;**6**; 423-427.

- 883 Higgins JP, Altman DG, Gotzsche PC, Juni P, Moher D, Oxman AD, Savovic J, Schulz KF, Weeks L,
884 Sterne JA et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*
885 2011;**343**; d5928.
- 886 Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;**11**; 1539-
887 1558.
- 888 Homan G, Litt J, Norman RJ. The FAST study: Fertility ASsessment and advice Targeting lifestyle
889 choices and behaviours: A pilot study. *Hum Reprod* 2012;**8**; 2396-2404.
- 890 Infertility Network UK. NHS Funding for Fertility Treatment in Scotland. 2015: **October**.
- 891 Johns DJ, Hartmann-Boyce J, Jebb SA, Aveyard P, on behalf of the Behavioural Weight Management
892 Review Group. Weight change among people randomized to minimal intervention control groups in
893 weight loss trials. 2016;**4**; 772-780.
- 894 Källén BA. Antiobesity drugs in early pregnancy and congenital malformations in the offspring.
895 2014;**6**; e571-e576.
- 896 Karimzadeh MA, Javedani M. An assessment of lifestyle modification versus medical treatment with
897 clomiphene citrate, metformin, and clomiphene citrate-metformin in patients with polycystic ovary
898 syndrome. *Fertil Steril* 2010;**1**; 216-220.
- 899 Karlsen K, Humaidan P, Sorensen LH, Alsbjerg B, Ravn P. Motivational interviewing: A part of the
900 weight loss program for overweight and obese women prior to fertility treatment. *Gynecol Endocrinol*
901 2013;**9**; 839-842.
- 902 Keltz J, Zapantis A, Jindal S, Lieman H, Santoro N, Polotsky A. Overweight Men: Clinical pregnancy
903 after ART is decreased in IVF but not in ICSI cycles. *J Assist Reprod Genet* 2010;**9-10**; 539-544.
- 904 Khan KS, Ter Riet G, Glanville J, Sowden AJ, Kleijnen J. Undertaking systematic reviews of research
905 on effectiveness: CRD's guidance for carrying out or commissioning reviews. 2001. NHS Centre for
906 Reviews and Dissemination.
- 907 Khaskheli M, Baloch S, Baloch AS. Infertility and weight reduction: Influence and outcome. *J*
908 *College Phys Surg Pak* 2013;**23**; 798-801.
- 909 Klenov V, Jungheim E. Obesity and reproductive function: a review of the evidence. *Curr Opin*
910 *Obstet Gynecol* 2014;**26**; 455-460.
- 911 Koning AMH. Fertility treatment in obese women. *Thesis* 2015:.
- 912 Kort J, Kim S, Lathi R. A Cohort Study to Evaluate the Impact of Weight Loss on Fertility Outcomes
913 in an Overweight/obese Population. *Fertil Steril* 2013;**3**; S4.
- 914 Kort J, Winget C, Kim S, Lathi R. A retrospective cohort study to evaluate the impact of meaningful
915 weight loss on fertility outcomes in an overweight population with infertility. *Fertil Steril* 2014;**5**;
916 1400-1403.
- 917 Kumar P, Arora S. Orlistat in polycystic ovarian syndrome reduces weight with improvement in lipid
918 profile and pregnancy rates. *J Hum Reprod Sci* 2014;**4**; 255-261.

- 919 Lazurova I, Dravecka I, Kraus V, Petrovicova J. Metformin versus sibutramine in the treatment of
920 hyperinsulinemia in chronically anovulating women. *Bratisl Lek Listy* 2004;**5-6**; 207-210.
- 921 Legro RS, Dodson WC, KrisEtherton PM, Kunselman AR, Stetter CM, Williams NI, Gnatuk CL,
922 Estes SJ, Fleming J, Allison KC et al. Randomized controlled trial of preconception interventions in
923 infertile women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 2015;**11**; 4048-4058.
- 924 Maheshwari A, Stofberg L, Bhattacharya S. Effect of overweight and obesity on assisted reproductive
925 technology - a systematic review. *Hum Reprod Update* 2007;**5**; 433-444.
- 926 Mahoney D. Lifestyle modification intervention among infertile overweight and obese women with
927 polycystic ovary syndrome. *J Am Assoc Nurse Pract* 2014;**6**; 301-308.
- 928 Mavropoulos JC, Yancy WS, Hepburn J, Westman EC. The effects of a low-carbohydrate, ketogenic
929 diet on the polycystic ovary syndrome: A pilot study. *Nutr Metab* 2005;**1**; 35.
- 930 Metwally M, Cutting R, Tipton A, Skull J, Ledger L W, Li C T. Effect of increased body mass index
931 on oocyte and embryo quality in IVF patients. *Reprod Biomed Online* 2007;**5**; 532-538.
- 932 Moran L, Noakes M, Clifton P, Tomlinson L, Norman R. Dietary composition in restoring
933 reproductive and metabolic physiology in overweight women with polycystic ovary syndrome. *J Clin
934 Endocrinol Metab* 2003;**2**; 812-819.
- 935 Moran LJ, Hutchison SK, Norman RJ, Teede HJ. Lifestyle changes in women with polycystic ovary
936 syndrome. *Cochrane Database Syst Rev* 2011a; 007506.
- 937 Moran L, Tsagareli V, Norman R, Noakes M. Diet and IVF pilot study: Short-term weight loss
938 improves pregnancy rates in overweight/obese women undertaking IVF. *Aust N Z J Obstet Gynaecol*
939 2011b;**51**; 455-459.
- 940 Mutsaerts MAQ, Van Oers AM, Groen H, Burggraaff JM, Kuchenbecker WKH, Perquin DAM, Koks
941 CAM, Van Golde R, Kaaijk EM, Schierbeek JM et al. Randomized trial of a lifestyle program in
942 obese infertile women. *N Engl J Med* 2016;**20**; 1942-1953.
- 943 National Clinical Guideline Centre. Obesity: Identification and management of overweight and
944 obesity in children, young people and adults. Commissioned by the National Institute for Health and
945 Care Excellence. 2014; 1.
- 946 National Institute for Health and Clinical Excellence. CG156 Fertility: Assessment and treatment for
947 people with fertility problems. 2013.
- 948 North American Association for the Study of Obesity, National Heart, Blood Institute, National
949 Institutes of Health (US), NHLBI Obesity Education Initiative. The practical guide: identification,
950 evaluation, and treatment of overweight and obesity in adults. 2000. National Institutes of Health,
951 National Heart, Lung, and Blood Institute, NHLBI Obesity Education Initiative, North American
952 Association for the Study of Obesity.
- 953 Palomba S, Falbo A, Giallauria F, Russo T, Rocca M, Tolino A, Zullo F, Orio F. Six weeks of
954 structured exercise training and hypocaloric diet increases the probability of ovulation after
955 clomiphene citrate in overweight and obese patients with polycystic ovary syndrome: A randomized
956 controlled trial. *Hum Reprod* 2010;**11**; 2783-2791.

- 957 Palomba S, Giallauria F, Falbo A, Russo T, Oppedisano R, Tolino A, Colao A, Vigorito C, Zullo F,
958 Orio F. Structured exercise training programme versus hypocaloric hyperproteic diet in obese
959 polycystic ovary syndrome patients with anovulatory infertility: A 24-week pilot study. *Hum Reprod*
960 2008;**3**; 642-650.
- 961 Qublan HS, Yannakoula EK, Al-Qudad MA, El-Uri FI. Dietary intervention versus metformin to
962 improve the reproductive outcome in women with polycystic ovary syndrome. A prospective
963 comparative study. *Saudi Med J* 2007;**11**; 1694-1698.
- 964 Ramlau-Hansen H, Thulstrup M, Nohr A, Bonde P, Sørensen I, Olsen J. Subfecundity in overweight
965 and obese couples. *Hum Reprod* 2007;**6**; 1634-1637.
- 966 Sallmen M, Sandler DP, Hoppin JA, Blair A, Baird DD. Reduced fertility among overweight and
967 obese men. *Epidemiology* 2006;**5**; 520-523.
- 968 Schneider G, Kirschner A M, Merkwitz R, Ertel H N. Increased estrogen production in obese men. *J*
969 *Clin Endocrinol Metab* 1979;**4**; 633-638.
- 970 Sermondade N, Faure C, Fezeu L, Shayeb G A, Bonde P J, Jensen T, Van Wely M, Cao J, Martini C
971 A, Eskandar M et al. BMI in relation to sperm count: an updated systematic review and collaborative
972 meta-analysis. *Hum Reprod Update* 2013;**3**; 221-231.
- 973 Shukla K, Chambial S, Dwivedi S, Misra S, Sharma P. Recent scenario of obesity and male fertility.
974 *Andrology* 2014;**6**; 809-818.
- 975 Sim KA, Dezarnaulds GM, Denyer GS, Skilton MR, Caterson ID. Weight loss improves reproductive
976 outcomes in obese women undergoing fertility treatment: A randomized controlled trial. *Clin Obes*
977 2014a;**2**; 61-68.
- 978 Sim KA, Partridge SR, Sainsbury A. Does weight loss in overweight or obese women improve
979 fertility treatment outcomes? A systematic review. *Obes Rev* 2014b;**10**; 839-850.
- 980 Styne-Gross A, Elkind-Hirsch K, Scott Jr T R. Obesity does not impact implantation rates or
981 pregnancy outcome in women attempting conception through oocyte donation. *Fertil Steril* 2005;**83**;
982 1629-1634.
- 983 The Royal Australian and New Zealand College of Obstetricians and Gynaecologists. Ovarian
984 stimulation in assisted reproduction. 2011; 1.
- 985 Thoma ME, McLain AC, Louis JF, King RB, Trumble AC, Sundaram R, Louis GMB. Prevalence of
986 infertility in the United States as estimated by the current duration approach and a traditional
987 constructed approach. *Fertil Steril* 2013;**5**; 1324-1331. e1.
- 988 Thomson RL, Buckley JD, Brinkworth GD. Perceived exercise barriers are reduced and benefits are
989 improved with lifestyle modification in overweight and obese women with polycystic ovary
990 syndrome: a randomised controlled trial. *BMC Womens Health* 2016;**1**; 1.
- 991 Thomson RL, Buckley JD, Moran LJ, Noakes M, Clifton PM, Norman RJ, Brinkworth GD. The
992 effect of weight loss on anti-Mullerian hormone levels in overweight and obese women with
993 polycystic ovary syndrome and reproductive impairment. *Hum Reprod* 2009;**8**; 1976-1981.
- 994 Thomson RL, Buckley JD, Noakes M, Clifton PM, Norman RJ, Brinkworth GD. The effect of a
995 hypocaloric diet with and without exercise training on body composition, cardiometabolic risk profile,

- 996 and reproductive function in overweight and obese women with polycystic ovary syndrome. *J Clin*
997 *Endocrinol Metab* 2008;**9**; 3373-3380.
- 998 Turner-McGrievy GM, Davidson CR, Wingard EE, Billings DL. Low glycemic index vegan or low-
999 calorie weight loss diets for women with polycystic ovary syndrome: A randomized controlled
1000 feasibility study. *Nutr Res* 2014;**6**; 552-558.
- 1001 Umul M, Kose S, Bilen E, Altuncu AG, Oksay T, Guney M. Effect of increasing paternal body mass
1002 index on pregnancy and live birth rates in couples undergoing intracytoplasmic sperm injection.
1003 *Andrologia* 2015;**3**; 360-364.
- 1004 Vahratian A, Smith YR. Should access to fertility-related services be conditional on body mass index?
1005 *Hum Reprod* 2009;**7**; 1532-1537.
- 1006 van Dam EWCM, Roelfsema F, Veldhuis JD, Hogendoorn S, Westenberg J, Helmerhorst FM, Frölich
1007 M, Krans HMJ, Meinders AE, Pijl H. Retention of estradiol negative feedback relationship to LH
1008 predicts ovulation in response to caloric restriction and weight loss in obese patients with polycystic
1009 ovary syndrome. *Am J Physiol Endocrinol Metab* 2004;**4**; E615-E620.
- 1010 Verhagen AP, de Vet HC, de Bie RA, Kessels AG, Boers M, Bouter LM, Knipschild PG. The Delphi
1011 list: a criteria list for quality assessment of randomized clinical trials for conducting systematic
1012 reviews developed by Delphi consensus. *J Clin Epidemiol* 1998;**12**; 1235-1241.
- 1013 Waters L, George AS, Chey T, Bauman A. Weight change in control group participants in
1014 behavioural weight loss interventions: a systematic review and meta-regression study. 2012;**1**; 120.
- 1015 Wise L, Palmer J, Rosenberg L. Body size and time-to-pregnancy in black women. *Hum Reprod*
1016 2013;**10**; 2856-2864.
- 1017 Wise L, Rothman K, Mikkelsen E, Sorensen H, Riis A, Hatch E. An internet-based prospective study
1018 of body size and time-to-pregnancy. *Hum Reprod* 2010;**1**; 253-264.
- 1019 World Health Organization. Obesity and Overweight fact sheet no 311. 2016: **April 17**.
- 1020 Zain M, Norman R. Impact of obesity on female fertility and fertility treatment. *Womens Health*
1021 *(Lond)* 2008;**2**; 183-194.
- 1022 Zawadzki JK, Dunaif A. Diagnostic criteria for polycystic ovary syndrome: towards a rational
1023 approach. In Dunaif A, Givens JR, Haseltine FP and Merriam GR (eds) Polycystic ovary syndrome.
1024 1992. Blackwell Scientific Publications, Boston, pp. 377-384.
- 1025 Zegers-Hochschild F, Adamson GD, de Mouzon J, Ishihara O, Mansour R, Nygren K, Sullivan E, van
1026 der Poel S, International Committee for Monitoring Assisted Reproductive Technology, World Health
1027 Organization. The International Committee for Monitoring Assisted Reproductive Technology
1028 (ICMART) and the World Health Organization (WHO) Revised Glossary on ART Terminology,
1029 2009. *Hum Reprod* 2009;**11**; 2683-2687.
- 1030
- 1031
- 1032

1033 **Figure legends**

1034 **AUTHOR: I have files for a Figure 1 (PRISMA chart in tiff format) plus Figures 2-14 in eps**
1035 **format. The legends below are for only 13 figures. Please would you recheck the figure**
1036 **numbering/titles?**

1037 Figure 1. PRISMA flow diagram illustrating the process of selection of studies for inclusion in the
1038 systematic review. **AUTHOR: please would you provide a higher quality image for Production?**

1039
1040 Figure 2. **AUTHOR: it may be helpful to also state in all titles that these are Forest plots.** Forest
1041 plot - Pregnancies associated with diet, diet and exercise, orlistat and motivational interviewing
1042 versus control.

1043

1044 Figure 3. Forest plot - Intervention versus control: live births.

1045 Figure 4. Forest plot - Intervention versus control: spontaneous conceptions.

1046 Figure 5. Forest plot - Intervention versus control: IVF conceptions.

1047 Figure 6. Forest plot - Intervention versus control: miscarriages for all participants enrolled.

1048 Figure 7. Forest plot - Intervention versus control: miscarriages for women who became
1049 pregnant.

1050 Figure 8. Forest plot - Intervention versus control: ovulation improvement.

1051 Figure 9. Forest plot - Intervention versus control: improvement of menstrual irregularity.

1052 Figure 10. Forest plot - Intervention versus control: oocytes retrieved per female participant.

1053 Figure 11. Forest plot - Intervention versus control: change in weight (kg).

1054 Figure 12. Forest plot - Intervention versus metformin: women who became pregnant.

1055 Figure 13. Forest plot - Intervention versus metformin: improvement in ovulation.

1056 Figure 14. Forest plot - Intervention versus metformin: improvement in menstrual irregularity
1057 per irregular menstruating female participant

1058

1059 **AUTHOR: we tend not to include Appendices, therefore these data have been**
1060 **divided and renamed as supplementary files. The quality of the file for the**

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Commented [BDA19]: Ok

Formatted: Font: Not Bold

Commented [BDA20]: As before - attached all available formats from generation tool so that best image can be selected

Formatted: Font: Bold

Commented [BDA21]: Good idea

Formatted: Font: Bold

Formatted: Indent: Left: 1.27 cm, No bullets or numbering

1061
1062
1063
1064
1065

Supplementary Data (search strategy) is not high enough for Production, therefore please would you send to me a high quality file, eps or tiff format, and preferably at 300dpi? Word and Powerpoint are acceptable if not pixelated or blurry at 400%. Thank you.

1066

1067

Commented [BDA22]: I can sent the original .txt file, which should solve the problems with the image

1068

1069

AUTHOR: for accuracy, please could you include *et al* in the references in the tables below, where required? Thank you.

Formatted: Font: Not Bold

Formatted: Font: Not Bold, Italic

Formatted: Font: Not Bold

Formatted: Font: 12 pt, Bold

Table 1. Types of interventions evaluated in studies of weight loss interventions and fertility in women and men who are overweight or obese.

Reducing diet/healthy eating and exercise programme or advice	
Duval Duval <i>et al</i> 2015 G	RCT
Karimzadeh Karimzadeh <i>et al</i> 2010	RCT
Moran Moran <i>et al</i> 2011	RCT
Mutsaerts Mutsaerts <i>et al</i> 2016	RCT
Palomba Palomba <i>et al</i> 2010 G	RCT
Sim Sim <i>et al</i> 2014 G	RCT
Clark Clark <i>et al</i> 1998 G	Non-RCT
Crosignani Crosignani <i>et al</i> 2003	Cohort
De Frène 2015	Cohort
Faure Faure <i>et al</i> 2014	Cohort
Galletly Galletly <i>et al</i> 1996a G	Cohort
Galletly Galletly <i>et al</i> 1996b G	Cohort
Hakonsen Hakonsen <i>et al</i> 2011	Cohort
Hollman Hollman <i>et al</i> 1996	Cohort
Khaskheli Khaskheli <i>et al</i> 2013	Cohort
Mahoney Mahoney <i>et al</i> 2014	Cohort
Miller Miller <i>et al</i> 2008 G	Cohort
Salama Salama <i>et al</i> 2015	Cohort
Reducing diet/healthy eating, weight loss drugs, and exercise programme or advice	
Legro Legro <i>et al</i> 2015	RCT
Lazurova Lazurova <i>et al</i> 2004	Non-RCT
Kort Kort <i>et al</i> 2014	Cohort
Reducing diet/health eating, or type of diet	

<u>Becker</u> <u>Becker et al</u> 2015	RCT
<u>Moran</u> <u>Moran et al</u> 2003	RCT
<u>Qublan</u> <u>Qublan et al</u> 2007	RCT
<u>Turner-McGrievy</u> <u>Turner-McGrievy et al</u> 2014	RCT
<u>Aliyeva</u> <u>Aliyeva et al</u> 1993	Non-RCT
<u>Palomba</u> <u>Palomba et al</u> 2008 G	Non-RCT
<u>Awartani</u> <u>Awartani et al</u> 2012	Cohort
<u>Kiddy</u> <u>Kiddy et al</u> 1992	Cohort
<u>Mavropoulos</u> <u>Mavropoulos et al</u> 2005 G	Cohort
<u>Thomson</u> <u>Thomson et al</u> 2009	Cohort
<u>Tsagareli</u> <u>Tsagareli et al</u> 2006	Cohort
<u>Van Dam</u> <u>Van Dam et al</u> 2004	Cohort
Behavioural therapy, including motivational interviewing	
<u>Karlsen</u> <u>Karlsen et al</u> 2013	Non-RCT
<u>Koning</u> <u>Koning et al</u> 2015 C	Non-RCT
<u>Homan</u> <u>Homan et al</u> 2012	Cohort
Exercise programme	
<u>Thomson</u> <u>Thomson et al</u> 2008 G	RCT
<u>Palomba</u> <u>Palomba et al</u> 2008	Non-RCT
Orlistat	
<u>Kumar</u> <u>Kumar et al</u> 2014	RCT
Metformin	
<u>Karimzadeh</u> <u>Karimzadeh et al</u> 2010	RCT
<u>Kumar</u> <u>Kumar et al</u> 2014	RCT
<u>Qublan</u> <u>Qublan et al</u> 2007	RCT
<u>Sonmez</u> <u>Sonmez et al</u> 2005	RCT
<u>Lazurova</u> <u>Lazurova et al</u> 2004	Non-RCT
Acarbose	
<u>Sonmez</u> <u>Sonmez et al</u> 2005	RCT
Detailed lifestyle questionnaire	

Study ID	Intervention arm	Pregnancies per participant			% Live births per participant		% Miscarriages per participant (fewer = better)		% Miscarriages per pregnancy (fewer = better)		% Improved ovulation per anovulatory participant		% Improved menstrual irregularity / irregular menses per participant		Time to conception per participant (less = better)	
		N	Rank	%	Rank	%	Rank	%	Rank	%	Rank	%	Rank	%	Rank	%
Palomba Palomba et al 2008	Exercise	20	14	35.0			7	5.0	5	14.3	5	65.0				
Homan Homan et al 2012	Motivational interviewing	35	15	34.8			5	4.3	3	12.5						
Awartani Awartani et al 2012	Weight reduction advised	90	16	34.4												
Qublan Qublan et al 2007	Diet	24	17	33.3			4	4.2	3	12.5	10	50.0	11	61.9		
Legro Legro et al 2015	Diet & exercise	50	18	32.0	9	26.0	9	6.0	10	18.8	6	60.3				
Crosignani Crosignani et al 2003	Diet & exercise	33	19	30.3	10	24.2					8	55.6	6	66.7		
Aliyeva Aliyeva et al 1993	Diet	60	20	30.0							14	46.7	8	63.3		
Kiddy Kiddy et al 1992	Very low calorie diet/Low calorie diet	24	21	29.2	14	16.7	10	8.3	13	28.6			20	29.4		
Hollman Hollman et al 1996	Diet & exercise	35	22	28.6	12	22.9	8	5.7	11	20.0			3	80.0		
Legro Legro et al 2015	Diet, exercise, oral contraceptive pill	50	23	28.0	11	24.0	3	4.0	5	14.3	4	67.1				
Qublan Qublan et al 2007	Metformin	22	24	27.3			6	4.5	7	16.7	15	45.5	12	61.1		
Kumar Kumar et al 2014	Metformin	30	25	26.7							20	23.3				
Becker Becker et al 2015	Diet	14	26	21.4	13	21.4										
Karimzadeh Karimzadeh et al 2010	Diet & exercise	75	26	20.0									6	66.7		
Lazurova Lazurova et al 2004	Metformin	30	28	20.0									4	70.0		
Miller Miller et al 2008	Diet & exercise	12	29	16.7									1	83.3		
Karimzadeh Karimzadeh et al 2010	Metformin & clomiphene	88	30	14.8									10	62.5		
DeFrene 2015	Diet & exercise	23	31	13.0												
Karimzadeh Karimzadeh et al 2010	Metformin	90	32	12.2									13	55.6		
Salama Salama et al 2015	Mediterranean diet & exercise	58	33	12.1									9	62.8		

Study ID	Intervention arm	N	Pregnancies per participant		% Live births per participant		% Miscarriages per participant (fewer = better)		% Miscarriages per pregnancy (fewer = better)		% Improved ovulation per anovulatory participant		% Improved menstrual irregularity / irregular menses per participant		Time to conception per participant (less = better)	
			Rank	%	Rank	%	Rank	%	Rank	%	Rank	%	Rank	%	Rank	%
Turner-McGrievy Turner-McGrievy <i>et al</i> 2014	Low calorie diet	9	34	11.1							22	0.0				
Palomba Palomba <i>et al</i> 2008	Diet	20	35	10.0			1	0.0	1	0.0	19	25.0				
Galletly Galletly <i>et al</i> 1996(a)	Exercise & advice	96	36	9.4												
Moran Moran <i>et al</i> 2003	High protein diet	14	37	8.7									21	26.1	1	4-5 weeks
Lazurova Lazurova <i>et al</i> 2004	Sibutramine, diet, exercise	15	38	6.7									14	53.3		
Thomson Thomson <i>et al</i> 2008	Diet & aerobic exercise	31	39	6.3							10	50.0	17	42.9		
Moran Moran <i>et al</i> 2003	Low protein diet	14	40	4.3									22	22.7	1	4-5 weeks
Thomson Thomson <i>et al</i> 2008	Diet	30	41	3.3							10	50.0	24	21.4		
Palomba Palomba <i>et al</i> 2010	Diet & exercise, Clomiphene	32	42	3.1							18	37.5	19	34.4		
Thomson Thomson <i>et al</i> 2008	Diet & aerobic & resistance exercise	33	43	3.0							16	42.8	16	44.4		
Turner McGrievy 2014	Vegan diet	9	44	0							22	0.0				
Sonmez Sonmez <i>et al</i> 2005	Acarbose	15									2	86.7	5	69.2		
Sonmez Sonmez <i>et al</i> 2005	Metformin	15									3	80.0	2	80.0		
Van Dam Van Dam <i>et al</i> 2004	Very low calorie diet	15									6	60.0				
Thomson Thomson <i>et al</i> 2009	Diet	52									20	19.2	18	42.3		
Palomba Palomba <i>et al</i> 2010	Diet & exercise	32									21	12.5	25	12.5		
Mahoney Mahoney <i>et al</i> 2014	Motivational interviewing, diet, exercise												23	22.2		

Table 3. Weight reduction outcomes for women (lower numbers for ranking indicate better outcomes).

Study ID	Intervention Arm	N	Mean Weight change (kg) Per Participant		Mean BMI change (kg/m ²) Per Participant		Mean Waist Circumference change (cm) Per Participant	
			Rank	Change	Rank	Change	Rank	Change
Mavropoulos et al 2005	Low calorie ketogenic diet	5	1	-12.34 ± 6.37	8	-4.0		
Hollman et al 1996	Diet & exercise	35	2	-10.2 ± 7.9	10	-3.4 (median)		
Clark et al 1998	Diet & exercise	67	3	-10.2 ± 4.3	9	-3.7 ± 1.6		
Thomson et al 2008	Diet & aerobic exercise	31	4	-10.1 ± 5.6			1	-11.7 ± 6.1
Karlsen et al 2013	Motivational interviewing	110	5	-9.3	11	-3.3		
Thomson et al 2009	Diet	52	6	-9.0 ± 0.8 (SEM)			4	-10.4 ± 0.9 (SEM)
Thomson et al 2008	Diet	30	7	-8.6 ± 6.0			3	-10.8 ± 7.1
Thomson et al 2008	Diet & aerobic & resistance exercise	3	8	-8.6 ± 5.2			2	-11.0 ± 6.3
Moran et al 2003	High protein diet	14	9	-8.5 ± 1.1				
Kumar et al 2014	Orlistat	30	10	-7.81 ± 0.66%	4	-8.12 ± 6.71%		
Kumar et al 2014	Metformin	30	11	-7.78 ± 0.57% (SEM)	3	-8.40 ± 0.65% (SEM)	14	-2.68 ± 0.16% (SEM)
Kiddy et al 1992	Very low calorie diet/low calorie diet	24	12	-6.9 ± 6.7				
Moran et al 2003	Low protein diet	14	13	-6.9 ± 0.8 (SEM)				
Sim et al 2014	Diet & exercise	26	14	-6.6 ± 4.6	14	-2.4 ± 1.6	6	-8.7 ± 5.6
Salama et al 2015	Mediterranean diet & exercise	58	15	-6.3 ± 20.8	16	-2.37 ± 7.28	7	-6.45 ± 14.13
Legro et al 2015	Lifestyle	50	16	-6.2, 95% CI (-7.3, -5.3)			8	-6.3, 95% CI (-9.2, -3.4)
Galletly et al 1996(b)	Diet and exercise	58	17	-6.2 ± 4.5	13	-2.4 ± 1.7		
Legro et al 2015	Diet, exercise, oral contraceptive pill	50	18	-6.1, 95% CI (-7.0, -5.2)			9	-6.2, 95% CI (-9.1, -3.3)
Tsagareli et al 2006	Very low calorie diet	6	19	-5.6; Range -8.2 to -5.3	17	-2.1; Range -1.8 to -3.2	5	-10.0; Range -7 to -14
Galletly et al 1996(a)	Diet & exercise	96	20	-5.2 ± 5.1				
Becker et al 2015	Diet	14	21	-4.51 ± 0.83 (SEM)			15	-1.31 ± 10.5
Palomba et al 2010	Diet & exercise, Clomiphene	32	22	-4.42 ± 9.22	12	-2.64 ± 4.26	11	-4.92 ± 5.87

Study ID	Intervention Arm	N	Mean Weight change (kg) Per Participant		Mean BMI change (kg/m ²) Per Participant		Mean Waist Circumference change (cm) Per Participant	
			Rank	Change	Rank	Change	Rank	Change
Mutsaerts Mutsaerts et al 2016	Diet & exercise	280	23	-4.4 ± 5.8	21	-1.3 (Median) IQR (-2.5, -0.07)		
Koning Koning et al 2015	Motivational interviewing	102	24	-4.3 ± 0.8 (SEM)	20	-1.4 ± 0.3 (SEM)		
Palomba Palomba et al 2010	Diet & exercise	32	25	-4.21 ± 8.56	15	-2.39 ± 3.54	12	-3.84 ± 6.35
Moran Moran et al 2011	Diet & exercise	18	26	-3.8 ± 3.0	19	-1.4 ± 1.1	10	-5.3 ± 4.6
Mahoney Mahoney et al 2014	Motivational interviewing, diet, exercise	9	27	-3.18 ± 2.27	24	-0.7 ± 7.31		
Turner-McGrievy Turner-McGrievy et al 2014	Vegan diet	9	28	-2.1 ± 3.5				
Turner-McGrievy Turner-McGrievy et al 2014	Low calorie diet	9	29	-0.4 ± 0.9				
Chavarro Chavarro et al 2012	Questionnaire	170	30	&0.3 (Median) IQR (-0.1, 1.8)				
Kort Kort et al 2014	Diet & exercise	52			1	-13.89% ± 3.71% in 17 with 10% weight loss -3.79% ± 3.98% in 35 with <10% weight loss		
Khaskheli Khaskheli et al 2013	Diet & exercise	85			2	-9.6 ± 1.23		
Lazurova Lazurova et al 2004	Sibutramine, diet, exercise	15			5	-4.6 ± 4.2 (SEM)		
Qublan Qublan et al 2007	Diet	24			6	-4.8		
Qublan Qublan et al 2007	Metformin	22			7	-4.1		
Miller Miller et al 2008	Diet & exercise	12			18	-2.06 ± 0.51 (SEM)		
Sonmez Sonmez et al 2005	Acarbose	15			22	-1.1 ± 2.78		
Lazurova Lazurova et al 2004	Metformin	30			23	-0.85 ± 0.38 (SEM)		
Sonmez Sonmez et al 2005	Metformin	15			25	-0.3 ± 2.47		
Homan Homan et al 2012	Motivational interviewing	35		"47% of the overweight participants had a modest loss of between 1 and 5 kg (not statistically significant)"				-3.8 in ½ of those attending follow-up and not pregnant
Duval Duval et al 2015	Diet & exercise	24		10/24 had ≥ 5% weight loss				12/24 had ≥ 5cm reduction (compared to 11/31 in control group)

Study ID	Intervention Arm	Mean Weight change (kg) Per Participant			Mean BMI change (kg/m ²) Per Participant		Mean Waist Circumference change (cm) Per Participant	
		N	Rank	Change (compared to 11/31 in control group)	Rank	Change	Rank	Change
Karimzadeh Karimzadeh et al 2010	Diet & exercise	75						Significantly lower waist circumference (P=0.001) in the lifestyle group than in the clomiphene, metformin, and clomiphene/metformin groups
Aliyeva Aliyeva et al 1993	Diet	60		“30-40% decrease in body weight in 2-4 months”				
Aliyeva Aliyeva et al 1993	Acupuncture	20		“Slower rate and extent of weight reduction: 7-9 months”				
Awartani Awartani et al 2012	Weight reduction advised	90				23/90 reduced their BMI below 35; 65/90 did not		
Crosignani Crosignani et al 2003	Diet & exercise	33						Those with 5% weight loss had mean waist circumference 94 cm, SD 9 Those with 10% weight loss had mean 86, SD 7
Van Dam Van Dam et al 2004	Very low calorie diet	15		14/15 loss at least 10% of body weight				
Palomba Palomba et al 2008	Exercise	20		Ovulatory (n=13): Mean -5.6%, SD 1.6% Anovulatory (n=7): Mean -2.0%, SD 0.2%		Ovulatory (n=13): Mean -10.0%, SD 3.7% Anovulatory (n=7): Mean -1.4%, SD 2.1%		Ovulatory (n=13): Mean -9.6%, SD 2.1% Anovulatory (n=7): Mean -2.5%, -SD 2.9%
Palomba Palomba et al 2008	Diet	20		Ovulatory (n=5): Mean -10.5%, SD 4.1% Anovulatory (n=15): Mean -2.3%, SD 3.1%		Ovulatory (n=5): Mean -15.4%, SD 3.9% Anovulatory (n = 15): Mean -4.0%, SD 5.2%		Ovulatory (n=5): Mean -9.4%, SD 2.5% Anovulatory (n=15): Mean -2.8%, SD 3.1%

Table 4. Reproductive and weight outcomes for infertile men.

	Faure <i>Faure et al</i> 2014 N=8	Hakonsen <i>Hakonsen et al</i> 2011 N=27 of 44 at follow-up
Clinical pregnancy rate	8/8	
Live birth rate	8/8	
Improvement in sperm DNA integrity	6/6 consenting to semen analysis upon completion	
Change in sperm concentration (million/ml)		Group 1 (n=9) Mean -11 95% CI -49, 27
Group 1: Weight loss 3.5-12.1%		Group 2 (n=9) Mean +19 95% CI -23, 61
Group 2: Weight loss 12.2-17.1%		Group 3 (n=9) Mean +17 95% CI -24, 58
Group 3: Weight loss 17.2-25.4%		
Change in sperm motility (% motile)		Group 1 (n=8) Mean -2 95% CI -15, 11
		Group 2 (n=9) Mean +4 95% CI -10, 18
		Group 3 (n=9) Mean +11 95% CI -3, 25
Change in normal sperm morphology (% normal forms)		Group 1 (n=9) Mean 0 95% CI -2, 4
		Group 2 (n=9) Mean +1 95% CI -3, 4
		Group 3 (n=9) Mean +4 95% CI 1, 7
Change in weight (kg)	Mean -4.40 SD 5.00	Median -22 Range -4 to -39
Change in BMI (kg/m ²)	Mean -1.22 SD 1.38	
Change in Waist circumference (cm)	Mean -8.50 SD 7.89	