

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/117189/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Anderson, James, Jathoul, Amit and Syed, Aisha 2019. Synthesis and bioluminescence of electronically modified and rotationally restricted colour-shifting infraluciferins. *Tetrahedron* 75 (3) , pp. 347-356.
[10.1016/j.tet.2018.11.061](https://doi.org/10.1016/j.tet.2018.11.061)

Publishers page: <https://doi.org/10.1016/j.tet.2018.11.061>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Synthesis and bioluminescence of electronically modified and rotationally restricted colour-shifting infraluciferins

James C. Anderson, Amit P. Jathoul and Aisha J. Syed

Department of Chemistry, University College London, 20 Gordon Street, London, WC1H 0AJ, UK.

and


School of Biosciences, University of Cardiff, Sir Martin Evans Building, Museum Avenue, Cardiff, CF10 3AX, UK

Supplementary Information


Copies of ^1H and ^{13}C NMR spectra

Graphs of emission kinetics of enzymes with different compounds acquired in the PIO


¹H – NMR: 4-((tert-butyldimethylsilyl)oxy)-2-nitroaniline (15)




¹H – NMR: 6-((tert-butyldimethylsilyl)oxy)-1H-benzo[d]imidazole-2-carbonitrile (16)




¹³C – NMR: 6-((tert-butyldimethylsilyl)oxy)-1H-benzo[d]imidazole-2-carbonitrile (**16**)




¹H – NMR: 6-hydroxy-1H-benzo[d]imidazole-2-carbonitrile (**17**)




¹H – NMR: 2-(6-hydroxy-1*H*-benzo[*d*]imidazol-2-yl)-4,5-dihydrothiazole-4-carboxylic acid (**10**)




¹H – NMR: 8-hydroxybenzo[4,5]imidazo[1,2-*a*]pyridine-3-carbonitrile (**20**)




¹³C-NMR: 8-hydroxybenzo[4,5]imidazo[1,2-*a*]pyridine-3-carbonitrile (20**)**




¹H – NMR: 2-(8-hydroxybenzo[4,5]imidazo[1,2-*a*]pyridin-3-yl)-4,5-dihydrothiazole-4-carboxylic acid (12**)**




¹³C-NMR: 2-(8-hydroxybenzo[4,5]imidazo[1,2-*a*]pyridin-3-yl)-4,5-dihydrothiazole-4-carboxylic acid (**12**)




¹H – NMR: 2-methyl-6-((triisopropylsilyl)oxy)-1*H*-benzo[*d*]imidazole (**22**)




¹³C-NMR: 2-methyl-6-((triisopropylsilyl)oxy)-1*H*-benzo[*d*]imidazole (**22**)




¹H – NMR: 2-methyl-6-((triisopropylsilyl)oxy)-1-((2- (trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazole (**23a**)




¹³C-NMR: 2-methyl-6-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazole (**23a**)




¹H – NMR: 2-methyl-5-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazole (**23b**)




¹³C-NMR: 2-methyl-5-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazole (**23b**)




¹H – NMR: 6-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazole-2-carbaldehyde (**24a**)




¹³C-NMR: 6-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazole-2-carbaldehyde (**24a**)




¹H – NMR: 5-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl) ethoxy) methyl)-1*H*-benzo[*d*]imidazole-2-carbaldehyde (**24b**)




¹³C-NMR: 5-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl) ethoxy) methyl)-1*H*-benzo[*d*]imidazole-2-carbaldehyde (**24b**)




¹H – NMR: methyl (2-(diethoxyphosphoryl)ethanethioyl)serinate (**25**)




¹H – NMR: methyl (*E*)-(3-(6-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)prop-2-enethioyl)serinate (**26a**)




¹³C-NMR: methyl (*E*)-(3-(6-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)prop-2-enethioyl)serinate (**26a**)




¹H – NMR: methyl(*E*)-(3-(5-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)prop-2-enethioyl)serinate (**26b**)




¹³C-NMR: methyl(*E*)-(3-(5-((triisopropylsilyl)oxy)-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)prop-2-enethioyl)serinate (**26b**)




¹H – NMR: methyl (*E*)-(3-(6-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)prop-2-enethioyl)serinate (**27a**)




¹³C-NMR: methyl (*E*)-(3-(6-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)prop-2-enethioyl)serinate (**27a**)




¹H – NMR: methyl (*E*)-(3-(5-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)prop-2-enethioyl)serinate (**27b**)




¹³C-NMR: methyl (*E*)-(3-(5-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)prop-2-enethioyl)serinate (**27b**)




¹H – NMR: methyl (*E*)-2-(2-(6-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylate (**28a**)




¹³C-NMR: methyl (*E*)-2-(2-(6-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylate (**28a**)




¹H – NMR: methyl (*E*)-2-(2-(5-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylate (**28b**)




¹³C-NMR: methyl (*E*)-2-(2-(5-hydroxy-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylate (**28b**)




¹H – NMR: methyl (*E*)-2-(2-(6-hydroxy-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylate




¹³C-NMR: methyl (*E*)-2-(2-(6-hydroxy-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylate




HMBC spectrum: methyl (*E*)-2-(2-(6-hydroxy-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylate




¹H – NMR: (*E*)-2-(2-(6-hydroxy-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylic acid (**13**)




¹³C-NMR: (E)-2-(2-(6-hydroxy-1*H*-benzo[*d*]imidazol-2-yl)vinyl)-4,5-dihydrothiazole-4-carboxylic acid (**13**)




Emission kinetics of enzymes with different compounds acquired in the PIO
SI Figure:
A




B



C



D



E

