

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/118609/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Barbieri, Mirko and Heard, Charles M. 2019. Isolation of punicalagin from Punica granatum rind extract using mass-directed semi-preparative ESI-AP single quadrupole LC-MS. *Journal of Pharmaceutical and Biomedical Analysis* 166 , pp. 90-94. 10.1016/j.jpba.2018.12.033

Publishers page: <http://dx.doi.org/10.1016/j.jpba.2018.12.033>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Isolation of punicalagin from *Punica granatum* rind extract using mass-directed semi-preparative ESI-AP single quadrupole LC-MS

Mirko Barbieri, Charles M. Heard ^{a,*}

^a School of Pharmacy & Pharmaceutical Sciences, Cardiff University, Cardiff CF10 3NB, Wales, United Kingdom

^{**}Email address: heard@cardiff.ac.uk Tel: +44 2920 875819

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

Abstract

We are living in a era of alarming increases in microbial resistance to currently available antibiotics, and there is a growing need for new pharmaceutical products to treat infectious diseases. The pomegranate is an edible fruit that has virucidal and antimicrobial activities which is primarily attributable to the high concentration of hydrolysable tannins. Punicalagin, a high molecular weight tannin (1084.7), accounts for approximately 70% of the total and is concentrated in the fruit exocarp (rind). It is the focus of much research, although it is prohibitively expensive to purchase which presents an obstacle to further exploitation and development. Here we describe a method for the isolation of punicalagin from pomegranate rind extract and total pomegranate tannins using an Agilent preparative mass-directed LC-MS single quadrupole ESI-API system, where the ionization was set as negative and the mass signal acquired in Single Ion Monitoring. Thanks to the automatic fraction collector, the method can be used in automatic mode and is capable of producing punicalagin with >95% purity.

Keywords: Punica Granatum, Pomegranate Rind Extract, TPT, Punicalagin, LC-MS

1. Introduction

We are currently living in an era of alarming and unprecedented increases in microbial resistance to our current arsenal of antibiotic drugs, and no new drugs have come onto the market in 30 years or so. New products are thus urgently needed [1, 2]. In addition, there is a clear predilection by patients for medicines that are of natural origin compared to synthetic drugs [3, 4].

The fruit of *Punica granatum* L. (pomegranate) has been used as a healthy and healing fruit for many years. It has long been used in Middle Eastern and Chinese medicine for its antioxidant, anti-inflammatory [5] bactericidal [6] and virucidal [7] properties. The phytochemicals constituents of pomegranates include many chemical species including sugars, flavonoids, anthocyanin, tannins, [8] and can be obtained from most parts of the fruit but the exocarp, or rind, has been shown to contain the highest concentration of polyphenols. A major class of compounds is the hydrolysable ellagitannins of which punicalagin (2,3-(S)-hexahydroxydiphenoyl-4,6-(S,S)-gallagyl-D-glucose) constitutes up to 70% [Figure 1]. It is a large molecule (MW 1084.71) and is comprised of 3 moieties: gallagic acid and ellagic acid linked via a glucose unit. The molecule naturally adopts one of two anomeric forms, α and β , depending on whether the glucose is in the boat or chair conformation, and the two exist in a 1:2 ratio under standard conditions.

Ongoing research shows that punicalagin is primarily responsible for antimicrobial, anti-inflammatory and virucidal properties of PRE. The microbicidal activities of PRE and punicalagin have been shown to undergo significant potentiation when co-administered with metal ions, eg Zn (II) [9, 10]. Punicalagin therefore has potential as a therapeutic agent as outlined; however, it is very expensive to purchase (eg £217 for 10 milligrams from one major supplier) which presents a barrier to further development, as substantial quantities are required to carry out the necessary pre-clinical drug development work.

PRE is typically prepared from pomegranate rinds by extraction into boiling water [9] followed by centrifugation and filtration. The non-tannin components of PRE can be removed by column chromatography to yield concentrated total pomegranate tannins (TPT), for example using Amberlite XAD-16 resin. Punicalagin may then be isolated by column chromatography using a Sephadex LH-20 column which entails a complex gradient elution [11] or by countercurrent chromatography [12]. The methods typically used to isolate punicalagin thus require numerous time-consuming laboratory steps and lengthy sample preparation.

Given the need for new antimicrobial molecules has arisen over recent years, fast and reproducible methods to isolate significant amounts of antimicrobial molecules has the potential to expedite the development of new therapeutic products. Mass-directed purification is a type of preparative LCMS in which the instrument may be set to collect column eluate upon detecting a pre-selected mass or mass range, once the chromatographic conditions have been established. These sophisticated instruments are available from a number of manufacturers including Waters and Shimadzu, although we have in our laboratory an Agilent 1260 Infinity Automated LC/MS Purification System [13]. In the current work we describe a method for the isolation of high-purity punicalagin using this system.

81 2. Materials and methods

82

83 2.1 Materials

84

85 Amberlite XAD-16 (Alfa Aesar), LCMS-grade water and methanol were purchased from Fisher Scientific,
86 Loughborough, UK. Pomegranates, of Spanish origin, were purchased from a local store.

87

88

89

90 2.2 Preparation of Pomegranate Rind Extract (PRE)

91

92 The rinds from five pomegranates were cut in small pieces of approx. 2cm and blended in deionized water,
93 25% w/v, before being boiled for 10 min and transferred to 50mL tubes and centrifuged 7 times at 5840rpm
94 for 30 min on each occasion. The supernatant was then vacuum filtered through 0.45µm nylon membranes
95 (Whatman, UK) to yield a clear dark yellow-orange solution which was then freeze-dried and stored at -20°C
96 until required [14].

97

98

99 2.3 Preparation of Total Pomegranate Tannins (TPT) from PRE

100

101 Based on a previously reported method [11] a glass column was slurry-packed with 50g of Amberlite XAD-
102 16 resin in degassed and deionized water. This was then washed with methanol, followed by deionized
103 water, and then was left overnight to equilibrate. A 2cm layer of sand was added to the top and the bottom.
104 Freeze-dried PRE was reconstituted in deionized water at a concentration of 200 mg/mL and loaded onto the
105 column. Firstly, the water-soluble components were washed out using a copious amount of water, then the
106 TPT fraction was eluted after washing with 70mL of methanol and collected. This solution was then rotary
107 evaporated using a Buchi instrument with the temperature set at 40°C and the vacuum at 50mbar. The TPT
108 product was stored at 2-4 °C until further use. Analysis of the tannin-free fraction (section 2.4) confirmed that
109 no tannins were present (data not shown).

110

111 2.4 Analytical HPLC analysis of TPT

112

113 The TPT was analysed using an Agilent 1100 system fitted with a Kinetex C18 150 x 4.6mm 5µm 100Å RP
114 column (Phenomenex, Macclesfield, UK). Based on a previous method [9] a gradient mobile phase was used
115 composed of: solvent A H₂O + 0.1%TFA and solvent B: MeOH + 0.1%TFA. The gradient timetable
116 started with 5%B, 0-3 min 10%B, 3-15 min 20%B, 15-20 min 40%B, 20-25 min 60%B, 25-27 min 5%B, 27-

117 35 min 5%B. A 1 mg/mL sample in deionized H₂O was prepared, the injection volume was 20 µL and the
118 detection was performed at a wavelength of 258 nm with a UV detector.

119

120

121 2.5. Semi-preparative Isolation of Punicalagin using an LC-MS system

122

123 The isolation was performed using a modular semi-preparative LC-MS ESI-AP single quadrupole system with
124 automatic fraction collector (Agilent Technologies, Stockport, UK). Waters X-Select CSH C18 OBD 5µm
125 130Å columns were used: firstly a 50 x 10mm scout column which was used to set up a reliable extraction
126 method, then a 250 x 19mm semi-preparative column. The solvents used were H₂O (channel A) and MeOH
127 (channel B); no mobile phase acidification was used. The separation was performed in Single Ion Monitoring
128 (SIM) negative ionization mode, with a capillary voltage of 4000V, drying gas temperature of 350°C
129 and gas nebulizer pressure of 32 psi [15]. The threshold of the automatic fraction collector was set at 2000.

130

131 The gradient was scaled from the analytical method to be reproducible with these columns. Firstly, the scout
132 column method was converted from the analytical one, and a run was performed to check the reliability of the
133 separation. The sample was 100mg/mL in H₂O and 300µL were injected. Further runs were carried out to
134 control the consistency of the separations. The gradient used with this column started at t₀ with 5%B, 0-0.38
135 min 10%B, 0.38-1.80 min 20%B, 1.80-2.39 min 40%B, 2.39-2.98 min 60%B, 2.98-3.22 min 5%B, 3.22-4.16
136 min 5%B. Although even with this column was possible to isolate punicalagin, only low amounts were
137 collected, so the method evolved to the big semi-preparative column. The gradient for this column started
138 with 5%B, 0-7.77 min 10%B, 7.77-33.36 min 20%B, 33.36-44.02 min 40%B, 44.02-54.69 min 60%B, 54.69-
139 58.95 min 5%B, 58.95-76.01 min 5%B. This sample load was 225mg in 1.5mL H₂O. The flow rate for both
140 the columns was 20mL/min. Both manual and automatic collection were performed. As already stated above,
141 the threshold for the automatic collection was set above 2000. The samples obtained were then dried to yield
142 the purified punicalagin; firstly, the methanolic fraction was evaporated using a Buchi rotary evaporator (50
143 mbar, 50 °C), then the remaining solvent (aqueous) was removed by freeze-drying.

144

145

146 3. Results and discussions

147

148 As can be seen in the analytical HPLC chromatogram [Figure 2], the two anomers of punicalagin were well
149 separated in the preparative LCMS method described. The two main peaks α and β punicalagin eluted with a
150 different concentration of solvent B and always tend to an approximate ratio of 1:2 ratio. The peak at the end
151 can be identified as ellagic acid, one of the major building blocks of punicalagin. The analysis of the TPT
152 was performed to prove the presence of punicalagin, so it was possible to proceed with the purification.
153 Because of the differences in the stationary phase between the preparative and the analytical columns, the

chromatogram of the preparative HPLC doesn't appear well separated, but thanks to the mass spectrometric collection triggering, it was still possible to isolate highly pure punicalagin. Figure 2 shows the plot of the run with the small scout column. The vertical lines show the windows in which the punicalagin (alpha the first and beta the second) was collected; the ratio of the two isomers was found also in the mass chromatogram.

The analytical resolution of punicalagin utilises acidification, typically by trifluoroacetic acid (TFA) [9]. In LCMS the preferred acidifying reagent is formic acid as, compared to TFA, it tends not to leave behind residual molecules. However, we found that when the punicalagin fractions were evaporated to dryness, it was discovered that nearly all the punicalagin had degraded. This was attributed to the increasing acidity that accompanied drying as the acid was also increasing in concentration. Consequently, it was decided to omit altogether the formic acid to ensure greater stability of the punicalain, albeit at the expense of superior chromatographic resolution.

The semi-preparative column report [Figure 3] shows a different HPLC chromatogram and a similar SIM chromatogram, probably due to the different sample loading in the two columns: in the small scout column it was closer to the optimal loading. In this case, even though both the little signals in the mass spectrum were slightly above 2000, the only fraction collected was the high peak at 22 min. For completeness, each fraction collected was re-run through the preparative LC-MS [Figure 4] to check the purity of the punicalagin, and all the signals detected were approximately the same: the differences represented the different grade of purity (average of >95% pure). These tests were performed after 1 hour.

The purity of the collected punicalagin product was determined by analytical HPLC [9]. Figure 5 shows that 95% of the integrated areas (compared to the total areas) was accounted for by the punicalagin anomer peaks. Interestingly, we attempted to collect individual anomers separately, by programming the instrument to collect at times corresponding to α and β punicalagin accordingly - it would be of interest to determine if the anomers give rise to differing biological (eg anti-microbial, anti-inflammatory) activities. However, even though collections were seemingly successful using the acid-free elution, it was noted that each 'purified' anomer reverted to the 1:2 ratio after drying, with β -punicalagin is the more abundant anomer at neutral pH [16].

4. Conclusions

A rapid, reliable and automatable method to isolate punicalagin from TPT was developed using the 1260 Infinity Automated LC/MS purification system. This facilitating the development work in order to bring to market new punicalagin-based antimicrobial products at a time of great need whilst avoiding large expenses to purchase it from suppliers. Work is ongoing to purify punicalagin directly from PRE, thus avoiding the prior concentration of TPT through Amberlite. However, the TPT separation step is not onerous, and does provide a material that is freely soluble in methanol, useful for easy injection into the LCMS.

191 **Acknowledgments**

192

193 We are very grateful for the support of Wellcome Trust ISSF and the Erasmus+ student exchange program.

194

195

196

197 **References**

198

199

200 [1] E. Crouch, L. Dickes, A. Kahle. Review on antibiotic resistance. *Adv Pharmacoepidemiol Drug Saf*
201 2015, 4:3.

202

203 [2] V.K.Sharma, N. Johnson, L. Cizmas, T.J. McDonald, H. Kim. A review of the influence of treatment
204 strategies on antibiotic resistant bacteria and antibiotic resistance genes. *Chemosphere*
205 150 (2016) 702-714.

206

207 [3] J.A. Astin. Why patients use alternative medicine - results of a national study. *JAMA* 279 (1998) 1546-
208 1553.

209

210

211 [4] M.V. Seeman, N. Seeman, J.A. Cho. Worldwide preferences for natural remedies for “nervousness”
212 and common colds. *Journal of Psychiatry and Brain Science*; 3 (2018) 1-11.

213

214 [5] E.P. Lansky, R.A. Newman, *Punica granatum* (pomegranate) and its potential for prevention and
215 treatment of inflammation and cancer, *J. Ethnopharmacol.* 109 (2007) 177–206.
216 doi:10.1016/j.jep.2006.09.006.

217

218 [6] G. Millo, Antibacterial inhibitory effects of *Punica granatum* gel on cariogenic bacteria : an in vitro
219 Study, (2017) 152–157.

220

221 [7] J. Lu, K. Ding, Q. Yuan, Determination of punicalagin isomers in pomegranate husk,
222 *Chromatographia.* 68 (2008) 303–306. doi:10.1365/s10337-008-0699-y.

223

224 [8] K. Subashini, Review of Phytochemical Screening for Pomegranate Peel Extract.pdf, (n.d.).

225 doi:10.4010/2016.772.

226

227 [9] Houston DMJ, Bugert JJ, Denyer SP, Heard CM. Potentiated virucidal activity of pomegranate rind
228 extract (PRE) and punicalagin against Herpes simplex virus (HSV) when co-administered with zinc
229 (II) ions, and antiviral activity of PRE against HSV and aciclovir-resistant HSV. PlosOne, 0179291.

230

231 [10] D.M.J. Houston, J. Bugert, S.P. Denyer, C.M. Heard, Anti-inflammatory activity of Punica granatum
232 L. (Pomegranate) rind extracts applied topically to ex vivo skin, Eur. J. Pharm. Biopharm. 112 (2017)
233 30–37. doi:10.1016/j.ejpb.2016.11.014.

234

235 [11] N. Seeram, R. Lee, M. Hardy, D. Heber, Rapid large scale purification of ellagitannins from
236 pomegranate husk, a by-product of the commercial juice industry, Sep. Purif. Technol. 41 (2005) 49–
237 55. doi:10.1016/j.seppur.2004.04.003.

238

239 [12] J. Lu, Y. Wei, Q. Yuan, Preparative separation of punicalagin from pomegranate husk by high-speed
240 countercurrent chromatography, J. Chromatogr. B. 857 (2007) 175–179.
241 doi:10.1016/j.jchromb.2007.06.038.

242

243 [13] [https://www.agilent.com/en/products/liquid-chromatography/infinitylab-lc-purification-](https://www.agilent.com/en/products/liquid-chromatography/infinitylab-lc-purification-solutions/infinitylab-preparative-scale-lc-purification-solutions/1260-infinity-automated-lc-ms-purification-system)
244 [solutions/infinitylab-preparative-scale-lc-purification-solutions/1260-infinity-automated-lc-ms-](https://www.agilent.com/en/products/liquid-chromatography/infinitylab-lc-purification-solutions/infinitylab-preparative-scale-lc-purification-solutions/1260-infinity-automated-lc-ms-purification-system)
245 [purification-system](https://www.agilent.com/en/products/liquid-chromatography/infinitylab-lc-purification-solutions/infinitylab-preparative-scale-lc-purification-solutions/1260-infinity-automated-lc-ms-purification-system) (accessed 1 November 2018)

246

247 [14] D.M.J. Houston, B. Robins, J.J. Bugert, S.P. Denyer, C.M. Heard, In vitro permeation and biological
248 activity of punicalagin and zinc (II) across skin and mucous membranes prone to Herpes simplex
249 virus infection, Eur. J. Pharm. Sci. 96 (2017) 99–106. doi:10.1016/j.ejps.2016.08.013.

250

251 [15] V. Brighenti, S.F. Groothuis, F.P. Prencipe, R. Amir, S. Benvenuti, F. Pellati, Metabolite
252 fingerprinting of Punica granatum L. (pomegranate) polyphenols by means of high-performance
253 liquid chromatography with diode array and electrospray ionization-mass spectrometry detection, J.
254 Chromatogr. A. 1480 (2017) 20–31. doi:10.1016/j.chroma.2016.12.017.

255

256 [16] M. Kraszni, A. Marosi, C.K. Larive, NMR assignments and the acid–base characterization of the

257 pomegranate ellagitannin punicalagin in the acidic pH-range. Anal Bioanal Chem 405 (2013) 5807–
258 5816.

259

260

261

262

263

264

265 Figures:

- 266 1. Chemical structure of punicalagin
- 267 2. HPLC chromatogram of PRE
- 268 3. LC-MS report of TPT using the Scout column
- 269 4. LC-MS report of TPT using the Semi-preparative column
- 270 5. LC-MS report of collected pure punicalagin