

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/118822/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Wadey, Rebecca M., Connolly, Katherine D., Mathew, Donna, Walters, Gareth, Rees, D. Aled and James, Philip E. 2019. Inflammatory adipocyte-derived extracellular vesicles promote leukocyte attachment to vascular endothelial cells. *Atherosclerosis* 283 , pp. 19-27. 10.1016/j.atherosclerosis.2019.01.013

Publishers page: <https://doi.org/10.1016/j.atherosclerosis.2019.01....>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Inflammatory adipocyte-derived extracellular vesicles promote leukocyte attachment to vascular endothelial cells

Authors: Rebecca M. Wadey^a, Katherine D. Connolly^a, Donna Mathew^a, Gareth Walters^a, D. Aled Rees^b, Philip E. James^a.

Affiliations: ^aCardiff School of Sport and Health Sciences, Cardiff Metropolitan University, 200 Western Avenue, Cardiff, CF5 2YB, UK; ^bNeuroscience and Mental Health Research Institute, Cardiff University, Hadyn Ellis Building, Maindy Road, Cardiff, CF24 4HQ, UK.

Email Addresses:

Rebecca M. Wadey: wadeyrm@hotmail.co.uk

Katherine D. Connolly: kdconnolly@cardiffmet.ac.uk

Donna Mathew: domathew@cardiffmet.ac.uk

Gareth Walters: gwalters@cardiffmet.ac.uk

D. Aled Rees: reesda@cardiff.ac.uk

Philip E. James: pjames@cardiffmet.ac.uk

Address for Correspondence:

Professor Philip E. James
Cardiff School of Sport and Health Sciences
Cardiff Metropolitan University
200 Western Avenue
Cardiff
CF5 2YB, UK
E-mail: pjames@cardiffmet.ac.uk
Tel: +44 (0) 2920 417 129

Abstract

Background and Aims: Obesity is associated with an increased risk of cardiovascular disease, but the mechanisms involved are not completely understood. In obesity, the adipocyte microenvironment is characterised by both hypoxia and inflammation. Therefore, we sought to determine whether extracellular vesicles (EVs) derived from adipocytes in this setting might be involved in mediating cardiovascular disease, specifically by promoting leukocyte attachment to vascular endothelial cells.

Methods: Mature 3T3-L1 adipocytes were incubated for 24 hours under control, TNF- α (30 ng/mL), hypoxia (1% O₂), or TNF- α +hypoxia (30 ng/mL, 1% O₂) conditions. EVs were isolated by differential ultracentrifugation and analysed by nanoparticle tracking analysis. Primary human umbilical vein endothelial cells (HUVECs) were treated with EVs for 6 hours before being lysed for Western blotting to investigate changes in adhesion molecule production, or for use in leukocyte attachment assays.

Results: EVs from adipocytes treated with TNF- α and TNF- α +hypoxia increased vascular cell adhesion molecule (VCAM-1) production in HUVECs compared to basal level (4.2 ± 0.6 and 3.8 ± 0.3 -fold increase, respectively ($p < 0.05$)), an effect that was inhibited by an anti-TNF- α neutralising antibody. Production of other adhesion molecules (E-selectin, P-selectin, platelet endothelial cell adhesion molecule and VE-Cadherin) were unchanged. Pre-incubating HUVECs with TNF- α +hypoxia EVs significantly increased leukocyte attachment compared to basal level (3.0 ± 0.4 -fold increase ($p < 0.05$)).

Conclusions: Inflammatory adipocyte EVs induce VCAM-1 production in vascular endothelial cells, accompanied by enhanced leukocyte attachment. Preventing adipocyte derived EV-induced VCAM-1 upregulation may offer a novel therapeutic target in the prevention of obesity-driven cardiovascular disease.

Word Count: 245/250

Introduction

Obesity is a risk factor for the development of cardiovascular disease (CVD)¹ but the complex mechanisms linking the two together are still to be fully elucidated. Obesity combined with metabolic syndrome (an umbrella term encompassing insulin resistance, dyslipidaemia, vascular endothelial dysfunction, hypertension and a hypercoagulable state), further increases the risk of CVD^{2, 3}. Whilst a complete understanding of the mechanisms that trigger adipose tissue to become metabolically dysregulated in obesity are not fully understood, evidence suggests that hypoxia and inflammation play a role.

Obese adipose tissue contains localised regions of hypoxia⁴⁻⁷ thought to arise due to increased cell number and cell size beyond the distance oxygen can diffuse from capillaries. Free fatty acids and cytokines produced by hypoxic adipocytes attract and activate macrophages and in doing so, induce local tissue inflammation⁸. Under such conditions, physiological adipokine production is dysregulated. For example, release of pro-inflammatory tumour necrosis factor (TNF)- α is increased⁹ whilst anti-inflammatory adiponectin is decreased¹⁰. The elevation of TNF- α and decrease of adiponectin in plasma are individually associated with increased leukocyte attachment to vascular endothelial cells via increased production of vascular endothelial cell adhesion molecules^{11, 12}. Subsequent phagocytosis of oxidised low-density lipoproteins within vessel walls results in lipid-laden foam cell formation¹³ and later, atherosclerotic plaque development.

Like most cells, adipocytes release extracellular vesicles (EVs)^{14, 15}. EVs are membrane-bound structures that can convey biological information from a cell of origin to a recipient cell to achieve a target effect. Based on diameter and biogenesis they are categorised into three broad groupings: exosomes (~80-100 nm; released from multivesicular bodies upon plasma membrane fusion), microvesicles (~200-1000 nm; shed directly from the plasma membrane) and apoptotic bodies (~1-5 μ m; released as blebs during apoptosis). *In vitro* data show that adipocytes release EVs containing adipokines¹⁶⁻¹⁹. In obesity, adipocytes reside in a hypoxic and inflammatory environment, therefore the content and function of adipocyte-derived EVs in this environment may be different to adipocyte-derived EVs from lean adipose tissue. Adipocyte EVs circulating in obesity may contribute to vascular endothelial dysfunction which later leads to atherosclerotic plaque formation. We sought to test this hypothesis *in vitro* using a leukocyte attachment assay with vascular endothelial cells isolated from umbilical cords and EVs derived from adipocytes cultured in conditions representative of lean and obese (hypoxic and inflammatory) adipose tissue.

Materials and Methods

Ethical approval for this study was granted by the NHS Health Research Authority (ethics committee reference: 14/NW/1459), and Cardiff Metropolitan University's Research Ethics Committee.

Adipocyte Culture and EV Isolation

3T3-L1 adipocytes were cultured for 14 days as described by Connolly *et al*¹⁶. For all EV-isolation experiments, serum-containing culture medium was replaced with serum-free medium for 24 hours. During this time, adipocytes were exposed to one of four treatments representative of control (95 % air / 5 % CO₂), inflamed (30 ng/mL TNF- α (13473019, ThermoFisher, USA)), hypoxic (1 % O₂), and inflamed and hypoxic (30 ng/mL TNF- α , 1 % O₂) adipose tissue. Cell media were centrifuged at 4 °C at 1,000 *g* (5 minutes), 15,000 *g* (15 minutes), and finally 100,000 *g* (1 hour) to pellet EVs. EVs were resuspended in 1 \times PBS and quantified by nanoparticle tracking analysis (NTA; a technique that tracks nanoparticles in suspension using a laser, and based on Brownian motion, determines particle size and concentration). NTA was performed using a NanoSight LM10 with a 642 nm laser (Malvern Instruments Ltd, Malvern, UK), using software version 2.3, screen gain 4 and camera level 10. Five, one-minute videos were recorded per sample with analysis screen gain 10 and detection threshold 4. Temperature ranged from 20 to 23.5 °C.

HUVEC Isolation and Culture

Human umbilical cords were obtained following elective Caesarean sections. Saline was used to flush blood out of the umbilical vein, before one end was clamped. Collagenase type 1A (1 mg/mL; C5894, Sigma) in Medium 199 (M199; 31150, ThermoFisher) at 37 °C, was syringed into the vein until the cord became taut and then the end of the cord was also clamped. After 15 minutes, one clamp was released allowing the vascular endothelial cell suspension to be collected into a Falcon tube. Copious growth medium (M199 containing 10 % (v/v) foetal bovine serum (10500-064, ThermoFisher), human epidermal growth factor (1 ng/mL; 13453029, ThermoFisher), hydrocortisone (1 μ g/mL; H0888, Sigma), gentamycin (35 μ g/mL; G1272, Sigma) and amphotericin (0.5 μ g/mL; A2942, Sigma)) was added to terminate the enzymatic digestion. Cells were centrifuged (300 *g*, 5 minutes, 4 °C), resuspended in growth medium, and plated in 96-well plates for leukocyte attachment assays or 6-well plates for protein assays. Plates were pre-coated with 1 % (w/v) bovine skin gelatin (G9391, Sigma) in 1 \times PBS. HUVECs were given 2 hours to adhere, before the medium was aspirated and replaced. HUVECs reached confluency within 4-7 days, were never passaged, and were used for experiments within 7 days.

Leukocyte Isolation

Blood (10 mL) was obtained from healthy volunteers by venepuncture and transferred into a universal container (UC) containing Heparin (100 μ L, 5,000 I.U/mL,

Wockhardt, India). Dextran (2.5 mL, 6 % (w/v), Sigma) dissolved in 1× balanced salt solution (BSS; 0.13 M NaCl, 2.6 mM KCl, 8.0 mM Na₂HPO₄, 1.83 mM KH₂PO₄, pH 7.4) was added and mixed by a single inversion. Blood was transferred into a fresh UC and allowed to fractionate for 1 hour. The buffy coat layer (~1.5 mL) was transferred into a fresh UC. Cells were collected by centrifugation (300 g, 2 minutes, room temperature) and the pellet resuspended in sterile H₂O to burst any contaminating erythrocyte membranes. After 10 seconds the UC was filled with BSS and the leukocytes were pelleted as above. The pellet was resuspended in Krebs-BSA (0.1 % (w/v) bovine serum albumin (BSA) in 1× Krebs (1.2 M NaCl, 0.48 M KCl, 0.12 M KH₂PO₄, 0.12 M MgSO₄, 0.13 M CaCl₂, 2.5 M HEPES, pH 7.4)) that had been passed through a 0.22 µm pore sterile-filter (Merck Millipore, USA) for sterilisation and remove possible contaminating serum extracellular vesicles. Cells were incubated on ice, in darkness, with CellTrace™ (1:1000; C34851, Invitrogen). After 10 minutes, leukocytes were centrifuged, resuspended in Krebs-BSA and left to settle on ice for 30 minutes. Prior to the attachment assay, the leukocyte suspension was diluted 1:10 with Krebs-BSA pre-warmed to 37 °C.

Leukocyte Attachment Assay

Serum-free M199 (150 µL) containing $1.2 \pm 0.4 \times 10^{10}$ (Control), $2.5 \pm 0.4 \times 10^{10}$ (TNF-α), $8.6 \pm 0.8 \times 10^{10}$ (Hypoxia) and $1.2 \pm 0.3 \times 10^{11}$ (TNF-α & Hypoxia) adipocyte EV/mL (mean ± SEM; N=3, n=9), was added to individual wells of a 96-well plate containing a confluent monolayer of HUVECs. Wells were also used to determine basal leukocyte attachment (no EVs; negative control) or 100 ng/mL TNF-α to increase HUVEC VCAM-1 production (no EVs; positive control). After 6 hours, EVs were removed with three Krebs washes before 150 µL of the fluorescently-labelled leukocyte suspension was added to all wells. Plates were incubated for 30 minutes before non-adherent cells were removed with three Krebs washes. Three images from around the centre of each well were captured using an inverted fluorescence microscope. The percentage of total image area covered by leukocytes was determined using Image J software (1.49v; National Institutes of Health, USA). Incubation timings were based on published protocols²⁰. Three separate sets of adipocyte-derived EVs were used in this experiment and each set of EVs was tested on three HUVEC samples (N=3, n=9).

Effect of Adipocyte EVs on HUVEC Adhesion Molecule Production

Serum-free M199 (1.5 mL) containing $1.2 \pm 0.4 \times 10^{10}$ (Control), $2.5 \pm 0.4 \times 10^{10}$ (TNF-α), $8.6 \pm 0.8 \times 10^{10}$ (Hypoxia) and $1.2 \pm 0.3 \times 10^{11}$ (TNF-α & Hypoxia) adipocyte EV/mL (mean ± SEM, N=3, n=3), was added to individual wells of a 6-well plate of HUVECs for 6 hours. Another well was used for the determination of basal protein production (no EVs; negative control). EVs were removed with three 1× PBS washes before HUVECs were lysed using radioimmunoprecipitation assay buffer (Invitrogen) containing protease inhibitors (Roche, Switzerland). Lysates were centrifuged (12,000 g, 10 minutes, 4 °C) and supernatants frozen until required. The protein concentration of each sample was determined using a NanoDrop spectrophotometer

1 and samples analysed by Western blotting. Lysates were mixed with sample buffer
2 and reducing agent (Invitrogen) and heated (95 °C, 10 minutes). Ten µg of total
3 protein from each sample was loaded onto 4-12 % Bis-Tris gels (Novex™,
4 ThermoFisher) and SDS-PAGE performed (180 V, 1 hour). Resolved proteins were
5 transferred to polyvinylidene difluoride membranes (40 V, 75 minutes). Membranes
6 were blocked using 5 % (w/v) non-fat dried milk in 1× Tris-buffered saline containing
7 0.05 % Tween-20 (TBST) (1 hour). Primary antibodies directed at proteins of interest
8 (vascular cell adhesion molecule (VCAM-1; ab134047), E-selectin (ab18981), P-
9 selectin (ab59738), platelet endothelial cell adhesion molecule (PECAM; ab28364),
10 vascular endothelial (VE)-cadherin (ab33168), endothelial nitric oxide synthase
11 (eNOS; ab76198) (all Abcam, Cambridge, UK)) were diluted 1:500 in blocking buffer
12 and incubated with membranes overnight. Membranes were then incubated with
13 horseradish peroxidase (HRP)-conjugated anti-rabbit IgG (1:1000; 10794347,
14 ThermoFisher) (2 hours). Protein bands developed on enhanced chemiluminescence
15 film (10607665, Fisher Scientific). Membranes were re-probed for β-actin (1:2000;
16 4970S, Cell Signalling Technology (CST), Netherlands) to confirm equal loading.
17 Band densitometry was performed using Image J. Three sets of adipocyte-derived
18 EVs were used in this experiment with each set tested on one HUVEC sample (N=3,
19 n=3).

20 **Effect of Adipocyte EVs on the Plasma Membrane Expression of HUVEC** 21 **Adhesion Molecules**

22 HUVECs were treated with 3T3-L1 EVs from each condition (control, TNF-α,
23 hypoxia, TNF-α+hypoxia) as described above, then washed twice and detached from
24 the culture plate using 200 µL 1× PBS. HUVECs were collected by centrifugation
25 (300 g, 5 minutes) and then re-suspended in ice-cold FACS Buffer (1% BSA (v/v) in
26 1× PBS). Flow cytometry was used to assess the surface adhesion molecule profile
27 of HUVECs. Antibodies used for flow cytometric analysis were obtained from
28 Biolegend® (BioLegend, San Diego, CA, USA). They include; phycoerythrin (PE)
29 anti-human CD144, allophycocyanin (APC) anti-human VCAM-1, Alexafluor 647 anti-
30 human PECAM-1, PE anti human E-Selectin, and PECy7 anti-human ICAM-1. Cells
31 were incubated with antibodies for 30 minutes at 4 °C in darkness before being
32 analysed on an FC500 MPL flow cytometer (Beckman Coulter equipped with 488 nm
33 and 633 nm lasers) and data captured on MPX Cytometer List Mode Data
34 Acquisition and Analysis Software (version 2.2). Acquisition was terminated upon
35 recording 10,000 events, and cells gated based on their forward scatter and side
36 scatter characteristics. Fluorescence minus one (FMO) stains were used to set the
37 positive gates for each antibody (See supplemental information and supplementary
38 Figure 3 for gating strategy). Three separate sets of adipocyte- derived EVs were
39 used in this experiment and each set of EVs was tested on one individual HUVEC
40 sample (N=3, n=3). Mean Fluorescence Intensity (MFI) was used to compare
41 surface marker expression between treatments.

TNF- α Neutralisation Assay

A TNF- α neutralising antibody was used to determine whether EVs derived from adipocytes treated with TNF- α , mediated VCAM-1 upregulation in HUVECs. To neutralise any pre-existing TNF- α , HUVECs and adipocyte-derived EVs were incubated separately in serum-free culture medium containing 0, 3, 10, 30, 100 or 300 ng/mL TNF- α neutralising antibody. After 2 hours, HUVEC media were removed and the EV-containing media were added to the corresponding wells. After 6 hours, HUVECs were lysed and VCAM-1 levels assessed by Western blotting, as previously described. Four sets of adipocyte-derived EVs were used in this experiment with each set tested on one HUVEC sample (N=4, n=4).

Analysis of 3T3-L1 Cell and EV Proteins

Adipocyte cell and EV lysates were assessed by Western blotting as previously described. Ten μ g of total protein was loaded per lane for cell lysates, and 5 μ g of total protein for EV lysates. Primary antibodies directed at fatty acid binding protein-4 (FABP4; 3544S, CST), adiponectin (2789S, CST), peroxisome proliferator-activated receptor gamma (PPAR γ ; 2443S, CST) and perilipin (9349S, Cell Signalling Technology), were used at a 1:500 dilution, and HRP-conjugated anti-rabbit IgG was used at a 1:1000 dilution. Three sets of adipocyte cell and EV lysates were used (N=3, n=3).

Statistical Analyses

Data are presented as mean or mode \pm SEM. A one-way ANOVA with Tuckey's Multiple Comparison Test was used to analyse differences. Data were analysed using GraphPad Prism (version 6; GraphPad Software Inc., USA) and *p*-values <0.05 were considered significant.

Results

Effect of Adipocyte EVs on Leukocyte-to-Endothelial Cell Attachment

HUVECs pre-treated with TNF α +hypoxia-derived adipocyte EVs, increased leukocyte attachment to the same extent as the positive control (Figure 1A). These increases were significant when compared to those observed for HUVECs not treated with EVs and HUVECs pre-treated with control adipocyte EVs (Figure 1A). Leukocyte attachment was also greater following treatment with TNF α +hypoxia-derived EVs compared to hypoxia-derived adipocyte EVs. Pre-treating HUVECs with TNF- α -derived adipocyte EVs also increased leukocyte attachment compared to HUVECs not treated with EVs. No other differences were observed in leukocyte attachment to HUVECs after pre-treatment with adipocyte EVs. Using size-exclusion chromatography, we confirm that TNF- α is associated with EVs; there is no free TNF- α (Supplementary Figure 1).

Effect of Adipocyte EVs on HUVEC Protein Production

Following treatment of HUVECs with adipocyte EVs, Western blotting was used to examine the production of proteins involved in leukocyte attachment (Figure 2). VCAM-1 production was increased in HUVECs treated with TNF- α EVs and TNF- α +hypoxia EVs compared to that of untreated HUVECs. No change was observed with control EVs nor hypoxia EVs (Figure 2A). The production of other adhesion proteins including E-selectin (Figure 2B), P-selectin (Figure 2C) and PECAM (Figure 2D) were unaffected by treatment with adipocyte EVs. In addition, no effect on the production levels of the vascular endothelial cell marker proteins VE-Cadherin (Figure 2E) and eNOS (Figure 2F) were identified.

Effect of Adipocyte EVs on the Plasma Membrane Expression of HUVEC Adhesion Molecules

Flow cytometry confirmed the expression of VCAM-1, E-selectin, PECAM-1 and ICAM-1 on HUVEC plasma membranes (Figure 3). However, there were no statistically significant differences in the mean fluorescence intensity for each protein between any of the treatment groups. TNF- α treatment (no EVs) of HUVECs was used as a positive control in all experiments and showed positive expression of adhesion markers, other than for PECAM-1 which showed high expression at baseline and following all treatment conditions.

TNF- α Neutralisation Prevents VCAM-1 Upregulation in HUVEC

To determine if the upregulation of VCAM-1 in HUVECs seen following treatment with TNF- α and TNF- α +hypoxia adipocyte EVs (Figure 2) is due to EV-associated TNF- α , a TNF- α neutralising antibody was used. Compared to untreated HUVECs, TNF- α +hypoxia EVs increased VCAM-1 production (Figure 4). Compared to the level of VCAM-1 production achieved following treatment with TNF- α +hypoxia EVs, co-treatment with 100 ng/mL and 300 ng/mL neutralising antibody completely inhibited VCAM-1 upregulation (Figure 4).

Adipokine Production in Adipocytes and Adipocyte-Derived EVs

No effect on the production of FABP4 in cell lysates due to any treatment was identified (Figure 5A). FABP4 in was increased in EVs lysates of cells treated with TNF- α regardless of normoxia or hypoxia (Figure 5B). Adiponectin production was decreased in cells treated with TNF- α +hypoxia compared to that of control cells, but individually, TNF- α and hypoxia had no effect (Figure 5C). All treatments appeared to decrease adiponectin in EV lysates compared to that of control cells (Figure 5D). PPAR γ production in cell lysates decreased in response to hypoxia and TNF- α +hypoxia (Figure 5E). Conversely, hypoxia- and TNF- α +hypoxia-derived EVs contained more PPAR γ (Figure 5F). Compared to that of control cells, perilipin production decreased in response to hypoxia and TNF- α +hypoxia (Figure 5G). No differences in the production of perilipin in EVs were identified (Figure 5H).

Effect of Inflammatory and Hypoxic Stimuli on Adipocyte EV Yield and Size.

Compared to control adipocytes, treatment with TNF- α , hypoxia and TNF- α +hypoxia all increased the number of EVs produced per cell (Figure 6A). Treatment with TNF- α , hypoxia, or TNF- α +hypoxia all decreased EV size compared to control adipocytes (Figure 6B).

Discussion

Increased adiposity has long been recognised as a risk factor for cardiovascular disease development¹. In obesity, adipose tissue inflammation is associated with vascular inflammation²¹ and one of the primary mediators of this process is the inflammatory cytokine, TNF- α . Whilst TNF- α is secreted by adipocytes themselves, levels in obese adipose tissue are predominantly raised by activated macrophages²². TNF- α desensitizes adipocytes to insulin²³, but distally and in terms of effects on vascular endothelial cells, plasma TNF- α levels positively correlate with adhesion molecule production, disrupted eNOS activity and oxidative stress²⁴. We are the first to show that EVs derived from adipocytes residing within a hypoxic and inflammatory, TNF- α -containing environment (modelling the pathophysiological inflammatory nature of obese adipose tissue *in vivo*), are implicated in the onset of vascular disease by increasing vascular endothelial cell adhesion molecule production and thereby promoting leukocyte attachment. In addition, we show that inflammatory and hypoxic stimuli affect the content of adipocyte-derived EVs, as well as their yield and size.

Leukocyte-to-endothelial cell attachment was increased following treatment of HUVECs with TNF- α and TNF- α +hypoxia derived adipocyte EVs when compared to untreated HUVECs. However, when compared to the level of leukocyte attachment following treatment of HUVECs with control EVs, attachment was only increased following treatment with TNF- α +hypoxia EVs. This suggests that a factor(s) conferred by hypoxia, can elicit a functional effect on vascular endothelial cells that exacerbates leukocyte adhesion in the presence of inflammation, and this is worthy of future investigation. The data do however, indicate that inflammatory EVs, regardless of whether they were produced under normoxic or hypoxic conditions, prime endothelial cells for subsequent leukocyte attachment.

Whilst flow cytometry confirmed the expression of the adhesion molecules VCAM-1, E-selectin, PECAM-1 and ICAM-1 on the surface of HUVEC plasma membranes, no differences between EV treatment groups were observed. Although not significant VCAM-1 surface expression did, however, show a trend to increase in cells treated with TNF- α and TNF- α +hypoxia derived adipocyte EVs and mirrored the increase in VCAM-1 detected by Western blotting. We hypothesize that in order to detect subtle changes in surface marker expression between treatment groups will likely require a very high number of HUVEC/experimental replicates and it is acknowledged further studies will be required in order to investigate differences in surface expression.

The fact that inflammatory adipocyte EVs increase HUVEC VCAM-1 production, does however, offer a mechanism through which leukocyte-to-endothelial cell attachment is achieved. TNFR1 receptors have been shown to induce VCAM-1 mRNA and protein via a pathway mediated by NF- κ B but not ERK, p38MAPK or JNK kinase²⁵. Future experiments aim to determine if the TNF- α delivered to HUVECs by

3T3-L1 EVs activates TNFR1 receptors and mediates VCAM-1 upregulation via a similar mechanism. As TNF- α neutralisation prevents VCAM-1 upregulation, we hypothesise that this would also prevent the increase in leukocyte-to-endothelial cell attachment following incubation of HUVEC with TNF- α and TNF- α +hypoxia derived adipocyte EVs. *In vivo*, VCAM-1 is involved in the firm attachment of leukocytes to endothelial cells prior to their transmigration through the vessel wall^{26, 27}, and whilst enhanced VCAM-1 production accompanied by leukocyte attachment is known to contribute to the progression of atherosclerosis^{28, 29}, we can now suggest that this is at least partly mediated by EVs derived from hypoxic and inflamed adipocytes.

In this study, EVs were not processed further to remove possible contaminating “free” TNF- α . However, we do provide evidence that TNF- α co-elutes with CD63 and FABP-4 following size exclusion chromatography of EV samples and that no “free” TNF- α elutes in later fractions. This indicates that TNF- α within EV samples is EV-associated (Supplementary Figure 1). In addition, it is important to note that the leukocytes for the leukocyte-to-endothelial attachment assay were isolated from heparin-anticoagulated blood. Heparin may activate platelets³⁰ causing them to expose P-selectin. P-selectin can consequentially bind to P-selectin glycoprotein on leukocytes, and this may have affected leukocyte attachment³¹. However, this effect would be equal across all experimental conditions, and as such, the results observed in this study can be attributed to 3T3-L1 EVs. It is also important to note that once isolated from blood, the leukocytes were resuspended in 0.22 μ m sterile-filtered Krebs-BSA. As such, there is a chance that bovine serum EVs smaller than 0.22 μ m were present in this buffer and that they too could have affected leukocyte adhesion. Likewise, this effect would have been equal across all conditions and consequentially the levels of leukocyte attachment observed across all experimental conditions would remain in proportion to each other.

Our results add to the growing body of literature confirming that adipocytes not only release EVs^{15, 16, 32}, but that the information conveyed through their content has a functional effect on other cell types. As such, we investigated the effect of inflammatory and hypoxic stimuli on the production of several proteins (FABP4, adiponectin, PPAR γ and perilipin) linked to cardiovascular disease in both adipocyte cell and EV lysates. We hypothesise that differences in the content of these proteins within EVs derived from inflammatory and/or hypoxic adipocytes may confer mechanistic influences on leukocyte attachment to vascular endothelial cells. Future studies will seek to identify and elucidate such mechanisms.

Plasma levels of FABP4, a fatty-acid chaperone protein³³, have been shown to increase in obesity³⁴ and are associated with vascular endothelial cell dysfunction³⁵. We observed no change in the FABP4 content of adipocyte cell lysates in response to inflammatory or hypoxic stimuli. This result supports a previous study whereby FABP4 production in adipocyte cell lysates was also shown not to change in response to hypoxia³⁶. What is interesting, however, is that the authors of this study

1 did identify an increased level of FABP4 in adipocyte culture media in response to
2 hypoxia. In our study, hypoxia did not increase the FABP4 content of EVs suggesting
3 that perhaps FABP4 is released by adipocytes in an EV-independent mechanism in
4 this setting. However, we did observe an increase in FABP4 in the lysates of EVs
5 derived from TNF- α treated adipocytes (regardless of normoxia or hypoxia). As such,
6 future experiments will aim to elucidate the influence of FABP4 within inflamed
7 adipocyte-derived EVs on vascular endothelial dysfunction.

8 Adiponectin is abundant within the circulation of healthy people and exerts regulatory
9 insulin-sensitising and anti-inflammatory effects^{37,38}. Plasma adiponectin levels
10 negatively correlate with degree of obesity³⁹ and therefore also correlate with
11 increased obesity-related co-morbidities including insulin resistance, type II diabetes
12 and cardiovascular disease⁴⁰. We found that a combined inflammatory and hypoxic
13 stimulus decreased adiponectin in adipocyte cell lysates. EV-associated adiponectin
14 also appeared to decrease compared to control in response to all treatments, but
15 particularly hypoxia. Using the same adipocyte cell model, Chen *et al.*, showed that
16 hypoxia prevents soluble adiponectin secretion⁴⁰, complementing our finding. Our
17 data add to this by suggesting that at least a proportion of this reduction in
18 adiponectin secretion under hypoxic conditions is because it is not being packaged
19 into EVs. Reduced adiponectin within adipocyte EVs may contribute to the decrease
20 in plasma adiponectin during obesity and thereby may contribute to the development
21 of cardiovascular disease.

22 PPAR γ is a nuclear receptor regulating the transcription of genes central to fatty acid
23 and energy metabolism⁴¹. We found that hypoxia decreased PPAR γ production in
24 adipocyte cell lysates. Down-regulation of intracellular PPAR γ in response to hypoxia
25 is well documented^{16, 42} and is mediated by inhibition of its transcription by hypoxia
26 inducible factor-1 (HIF-1 α). Interestingly, in our study we also observed a
27 simultaneous increase of PPAR γ within the lysates of EVs derived from hypoxic
28 adipocytes, confirming that adipocyte EVs are selectively packaged depending on
29 environmental cues. We hypothesize that HIF-1 α may also be implicated in the
30 selective packaging of PPAR γ into EVs destined for release, as a further means to
31 reduce intracellular PPAR γ levels. Overall, this finding highlights that EVs do not
32 necessarily possess the same content as their cell-of-origin, but rather that their
33 content reflects the cell-of-origin's current state.

34 Perilipin is a protein localised to the surface of lipid droplets and is involved in
35 lipolysis⁴³. In our study, hypoxia decreased perilipin production within adipocyte cell
36 lysates when normalised to total protein. Wang *et al.*,⁴⁴ previously showed no
37 difference in total perilipin production per individual adipocyte sampled from lean and
38 obese adipose tissue but when normalised to total protein or fat cell surface area,
39 perilipin levels were also significantly lower in obese samples. We propose that
40 hypoxia restricts perilipin production in adipose tissue in obesity *in vivo*. In addition,
41 whilst perilipin levels are increased in the circulation in obese mice and humans¹⁵

1 this may be because it is released in to the circulation from adipocytes as a free
2 protein rather than being associated with EVs. This hypothesis fits with our results as
3 we identified no difference in the perilipin content of adipocyte EVs in response to
4 inflammatory and hypoxic stimuli.

5 In terms of specific EV character, our group has previously shown that EVs isolated
6 from adipocytes express the vesicular proteins CD9, CD63, Alix, tumour
7 susceptibility gene (TSG101)¹⁶, and now show that they also exhibit characteristic
8 cup-shape morphology by electron microscopy (Supplementary Figure 1). Here we
9 show the yield and size of adipocyte-derived EVs can be modulated independently
10 by external stimuli. TNF- α , hypoxia and TNF- α +hypoxia increased the yield of
11 EVs/cell whilst decreasing their size. These physical changes may themselves also
12 confer distinct functional effects. However, it is important to note that the size data
13 presented is based solely on raw NTA values; no refractive index⁴⁵ nor mathematical
14 modelling⁴⁶ was applied.

15 In summary, we provide evidence that adipocytes residing in a hypoxic and
16 inflammatory environment produce EVs capable of inducing VCAM-1 production in
17 vascular endothelial cells, and that this effect promotes leukocyte attachment. We
18 also show that both inflammatory and hypoxic stimuli not only influence the
19 adipokine content of adipocytes and their EVs, but also effect EV yield and size.
20 Future studies will further explore the functional impact inflammatory and hypoxic
21 adipocyte EVs have on cardiovascular disease. Indeed, the functional effects of
22 adipocyte EVs in the setting of obesity, are likely to be not solely limited to vascular
23 endothelial cells. We anticipate that a better understanding of the mechanisms
24 through which adipocyte EVs negatively impact the vascular endothelium will open
25 the door for the development of novel therapies for preventing and treating obesity-
26 driven cardiovascular disease.

Conflict of Interest

The authors declare no conflict of interest.

Financial Support

This work was supported by the British Heart Foundation (grant reference: PG/14/51/30686).

Author contributions

Rebecca M. Wadey performed the experiments. Katherine D. Connolly performed pilot experiments and manuscript editing. Donna Mathew and Gareth Walters performed the flow cytometry experiments. D. Aled Rees and Philip E. James were the Principal Investigators leading the project.

Acknowledgments

We would like to thank the clinical staff at University Hospital Wales for assisting with the collection of umbilical cords, the volunteers who donated blood samples, and Dr Justyna Wiczak and Mrs Margaret Munnery for performing phlebotomy. We also thank Dr Errin Johnson (EM facility, Sir William Dunn School of Pathology, University of Oxford) for performing the transmission electron microscopy.

Figure Legends

Figure 1 – Leukocyte attachment to HUVECs pre-treated with and without adipocyte-derived EVs. (A) Ratio of percentage area covered by leukocytes to a confluent monolayer of HUVECs pre-treated with (C) control, (T) TNF- α , (H) hypoxia, and (TH) TNF- α +hypoxia-derived adipocyte EVs relative to that of (B; Basal) HUVECs not treated with EVs (negative control). Direct treatment with TNF- α acted as a positive control (mean \pm SEM; N=3, n=9; *** $p<0.001$). (B) Representative fluorescent images used to determine the percentage area of HUVECs covered by leukocytes. White dots are leukocytes stained with CellTraceTM. Scale bar = 200 μ m.

Figure 2 – Expression of adhesion and endothelial marker proteins in HUVECs following treatment with adipocyte EVs. Densitometry graphs and representative Western blots showing the ratio of (A) VCAM-1 (110 kDa), (B) E-Selectin (80 kDa), (C) P-Selectin (140 kDa), (D) PECAM (130 kDa), (E) VE-Cadherin (115 kDa) and (F) eNOS (140 kDa) expression in HUVECs following treatment with (C) control, (T) TNF- α , (H) hypoxia and (TH) TNF- α +hypoxia derived adipocyte EVs, relative to that of (B; basal) untreated HUVECs (mean \pm SEM; N=3, n=3; *** $p<0.001$). β -actin confirms equal loading (bottom blots). Ten μ g total protein per lane.

Figure 3 - Surface expression of HUVEC adhesion proteins following treatment with adipocyte EVs. Mean fluorescence intensity graphs showing the ratio of (A) VCAM-1 (-APC), (B) E-Selectin (-PE), (C) PECAM-1 (-AlexoFluor) and (D) ICAM-1 (-PE/Cy7) expressed on the plasma membrane of HUVECs following treatment with (C) control, (T) TNF- α , (H) hypoxia and (TH) TNF- α +hypoxia derived adipocyte EVs, relative to that of (B; basal) untreated HUVECs. Direct treatment with TNF- α was used as a positive control (mean \pm SEM; N=3, n=3 for all except ICAM-1 where N=2, n=2).

Figure 4 – TNF- α neutralisation inhibits endothelial VCAM upregulation following treatment with TNF- α +hypoxia adipocyte EVs. (A) Western blotting densitometry of VCAM-1 expression in control HUVEC lysates, and lysates of HUVECs treated with TNF- α +hypoxia EVs (denoted by +) in the presence of 0, 3, 10, 30, 100 and 300 ng/ml TNF- α neutralising antibody (mean \pm SEM; N = 4, n = 4; ** $p < 0.01$). (B) Representative VCAM-1 (110 kDa; top) and β -actin (40 kDa; bottom) blots. Twenty μ g total protein per lane.

Figure 5. FABP4, adiponectin, PPAR γ and perilipin expression in adipocyte and adipocyte-derived EV lysates. Densitometry graphs and representative Western blots for FABP4 and adiponectin in cell and EV lysates following (C) control, (T) TNF- α , (H) hypoxia and (TH) TNF- α +hypoxia treatment of adipocytes. FABP4 expression in (A) cell lysates (mean \pm SEM; N=6, n=6) and (B) EV lysates (mean \pm SEM; N=4, n=4) (15 kDa). Adiponectin expression in (C) cell lysates (mean \pm SEM; N=5, n=5; * $p < 0.05$) and (D) EV lysates (mean \pm SEM; N=2, n=2) (30 kDa). PPAR γ expression in (E) cell lysates (mean \pm SEM; N=6, n=6; *** $p < 0.001$) and (F) EV lysates (mean \pm SEM; N=3, n=3; ** $p < 0.01$) (53 and 57 kDa). Perilipin expression in (G) cell lysates (mean \pm SEM; N=5, n=5; *** $p < 0.001$) and (H) EV lysates (mean \pm SEM; N=3, n=3) (56 kDa). Twenty μ g total protein per lane of cell lysates, and 5 μ g total protein per lane of EV lysates.

Figure 6. Effect of inflammatory and hypoxic stimuli on adipocyte EV yield and size. The effect of (C) control, (T) TNF- α , (H) hypoxia and (TH) TNF- α +hypoxia treatments on (A) EV yield per cell (mean \pm SEM; n=3, n=9; *** $p < 0.001$) and (B) EV size (mode \pm SEM; n=3, n=9; *** $p < 0.001$).

References

1. Hubert HB, Feinleib M, McNamara PM, Castelli WP. Obesity as an independent risk factor for cardiovascular disease: A 26-year follow-up of participants in the framingham heart study. *Circulation*. 1983;67:968-977
2. Ortega FB, Lee DC, Katzmarzyk PT, Ruiz JR, Sui X, Church TS, Blair SN. The intriguing metabolically healthy but obese phenotype: Cardiovascular prognosis and role of fitness. *Eur Heart J*. 2013;34:389-397
3. Kaur J. A comprehensive review on metabolic syndrome. *Cardiol Res Pract*. 2014;2014:943162
4. Ye J, Gao Z, Yin J, He Q. Hypoxia is a potential risk factor for chronic inflammation and adiponectin reduction in adipose tissue of ob/ob and dietary obese mice. *Am J Physiol Endocrinol Metab*. 2007;293:E1118-1128
5. Ye J. Emerging role of adipose tissue hypoxia in obesity and insulin resistance. *Int J Obes (Lond)*. 2009;33:54-66
6. Pasarica M, Rood J, Ravussin E, Schwarz JM, Smith SR, Redman LM. Reduced oxygenation in human obese adipose tissue is associated with impaired insulin suppression of lipolysis. *J Clin Endocrinol Metab*. 2010;95:4052-4055
7. Hosogai N, Fukuhara A, Oshima K, Miyata Y, Tanaka S, Segawa K, Furukawa S, Tochino Y, Komuro R, Matsuda M, Shimomura I. Adipose tissue hypoxia in obesity and its impact on adipocytokine dysregulation. *Diabetes*. 2007;56:901-911
8. Trayhurn P, Wang B, Wood IS. Hypoxia and the endocrine and signalling role of white adipose tissue. *Arch Physiol Biochem*. 2008;114:267-276
9. Ouchi N, Parker JL, Lugus JJ, Walsh K. Adipokines in inflammation and metabolic disease. *Nat Rev Immunol*. 2011;11:85-97
10. Bulló M, Salas-Salvadó J, García-Lorda P. Adiponectin expression and adipose tissue lipolytic activity in lean and obese women. *Obes Surg*. 2005;15:382-386
11. Ouchi N, Kihara S, Arita Y, Maeda K, Kuriyama H, Okamoto Y, Hotta K, Nishida M, Takahashi M, Nakamura T, Yamashita S, Funahashi T, Matsuzawa Y. Novel modulator for endothelial adhesion molecules: Adipocyte-derived plasma protein adiponectin. *Circulation*. 1999;100:2473-2476
12. Ouedraogo R, Gong Y, Berzins B, Wu X, Mahadev K, Hough K, Chan L, Goldstein BJ, Scalia R. Adiponectin deficiency increases leukocyte-endothelium interactions via upregulation of endothelial cell adhesion molecules in vivo. *J Clin Invest*. 2007;117:1718-1726

- 1 13. Bakhai A. Adipokines--targeting a root cause of cardiometabolic risk. *QJM*.
2 2008;101:767-776
- 3 14. Robbins PD, Morelli AE. Regulation of immune responses by extracellular
4 vesicles. *Nat Rev Immunol*. 2014;14:195-208
- 5 15. Eguchi A, Lazic M, Armando AM, Phillips SA, Katebian R, Maraka S,
6 Quehenberger O, Sears DD, Feldstein AE. Circulating adipocyte-derived
7 extracellular vesicles are novel markers of metabolic stress. *J Mol Med (Berl)*.
8 2016;94:1241-1253
- 9 16. Connolly KD, Guschina IA, Yeung V, Clayton A, Draman MS, Von Ruhland C,
10 Ludgate M, James PE, Rees DA. Characterisation of adipocyte-derived
11 extracellular vesicles released pre- and post-adipogenesis. *J Extracell*
12 *Vesicles*. 2015;4:29159
- 13 17. Aoki N, Jin-no S, Nakagawa Y, Asai N, Arakawa E, Tamura N, Tamura T,
14 Matsuda T. Identification and characterization of microvesicles secreted by
15 3t3-l1 adipocytes: Redox- and hormone-dependent induction of milk fat
16 globule-epidermal growth factor 8-associated microvesicles. *Endocrinology*.
17 2007;148:3850-3862
- 18 18. Kralisch S, Ebert T, Lossner U, Jessnitzer B, Stumvoll M, Fasshauer M.
19 Adipocyte fatty acid-binding protein is released from adipocytes by a non-
20 conventional mechanism. *Int J Obes (Lond)*. 2014;38:1251-1254
- 21 19. Kranendonk ME, Visseren FL, van Balkom BW, Nolte-'t Hoen EN, van
22 Herwaarden JA, de Jager W, Schipper HS, Brenkman AB, Verhaar MC,
23 Wauben MH, Kalkhoven E. Human adipocyte extracellular vesicles in
24 reciprocal signaling between adipocytes and macrophages. *Obesity (Silver*
25 *Spring)*. 2014;22:1296-1308
- 26 20. Lever R, Rose MJ, McKenzie EA, Page CP. Heparanase induces
27 inflammatory cell recruitment in vivo by promoting adhesion to vascular
28 endothelium. *Am J Physiol Cell Physiol*. 2014;306:C1184-1190
- 29 21. Hagita S, Osaka M, Shimokado K, Yoshida M. Adipose inflammation initiates
30 recruitment of leukocytes to mouse femoral artery: Role of adipo-vascular axis
31 in chronic inflammation. *PLoS One*. 2011;6:e19871
- 32 22. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW.
33 Obesity is associated with macrophage accumulation in adipose tissue. *J Clin*
34 *Invest*. 2003;112:1796-1808
- 35 23. Hotamisligil GS, Shargill NS, Spiegelman BM. Adipose expression of tumor
36 necrosis factor- α : Direct role in obesity-linked insulin resistance. *Science*.
37 1993;259:87-91
- 38 24. Zhang H, Zhang J, Ungvari Z, Zhang C. Resveratrol improves endothelial
39 function: Role of $\text{tnf}\{\alpha\}$ and vascular oxidative stress. *Arterioscler Thromb*
40 *Vasc Biol*. 2009;29:1164-1171

- 1 25. Zhou Z, Connell MC, MacEwan JD. TNFR1-induced NF- κ B, but not ERK,
2 p38MAPK or JNK activation, mediates TNF-induced ICAM-1 and VCAM-1
3 expression on endothelial cells. *Cell Sig.* 2007;19: 1238-1248
- 4 26. Granger D, Senchenkova E. *Inflammation and the microcirculation*. San
5 Rafael, California: Morgan & Claypool Life Sciences; 1 edition; 2010.
- 6 27. Robinson LA, Tu L, Steeber DA, Preis O, Platt JL, Tedder TF. The Role of
7 Adhesion Molecules in Human Leukocyte Attachment to Porcine Vascular
8 Endothelium: Implications for Xenotransplantation. *J Immunol.*
9 1990;161:6931-6938.
- 10 28. Onat D, Brillon D, Colombo PC, Schmidt AM. Human vascular endothelial
11 cells: A model system for studying vascular inflammation in diabetes and
12 atherosclerosis. *Curr Diab Rep.* 2011;11:193-202
- 13 29. Cybulsky MI, Iiyama K, Li H, Zhu S, Chen M, Iiyama M, Davis V, Gutierrez-
14 Ramos JC, Connelly PW, Milstone DS. A major role for VCAM-1, but not
15 ICAM-1, in early atherosclerosis. *The Journal of Clinical Investigation.*
16 2001;107:1255–1262.
- 17 30. Durcin M, Fleury A, Taillebois E, Hilairet G, Krupova Z, Henry C, Truchet S,
18 Trötz Müller M, Köfeler H, Mabilieu G, Hue O, Andriantsitohaina R, Martin P,
19 Le Lay S. Characterisation of adipocyte-derived extracellular vesicle subtypes
20 identifies distinct protein and lipid signatures for large and small extracellular
21 vesicles. *J Extracell Vesicles.* 2017;6:1305677
- 22 31. Gao C, Boylan B, Fang J, Wilcox DA, Newman DK, Newman PJ. Heparin
23 promotes platelet responsiveness by potentiating α IIb β 3-mediated outside-in
24 signaling. *Blood.* 2011;117:4946-4952
- 25 32. Falati S, Liu Q, Gross P, Merrill-Skoloff G, Chou J, Vandendries E, Celi A,
26 Croce K, Furie BC, Furie B. Accumulation of Tissue Factor into Developing
27 Thrombi In Vivo Is Dependent upon Microparticle P-Selectin Glycoprotein
28 Ligand 1 and Platelet P-Selectin. *JEM.* 2003;197:1585-1598.
- 29 33. Furuhashi M, Hotamisligil GS. Fatty acid-binding proteins: Role in metabolic
30 diseases and potential as drug targets. *Nat Rev Drug Discov.* 2008;7:489-503
- 31 34. Xu A, Wang Y, Xu JY, Stejskal D, Tam S, Zhang J, Wat NM, Wong WK, Lam
32 KS. Adipocyte fatty acid-binding protein is a plasma biomarker closely
33 associated with obesity and metabolic syndrome. *Clin Chem.* 2006;52:405-
34 413
- 35 35. Aragonès G, Saavedra P, Heras M, Cabré A, Girona J, Masana L. Fatty acid-
36 binding protein 4 impairs the insulin-dependent nitric oxide pathway in
37 vascular endothelial cells. *Cardiovasc Diabetol.* 2012;11:72
- 38 36. Wu LE, Samocha-Bonet D, Whitworth PT, Fazakerley DJ, Turner N, Biden TJ,
39 James DE, Cantley J. Identification of fatty acid binding protein 4 as an
40 adipokine that regulates insulin secretion during obesity. *Mol Metab.*
41 2014;3:465-473

- 1 37. Makki K, Froguel P, Wolowczuk I. Adipose tissue in obesity-related
2 inflammation and insulin resistance: Cells, cytokines, and chemokines. *ISRN*
3 *Inflamm*. 2013;2013:139239
- 4 38. Arita Y, Kihara S, Ouchi N, Takahashi M, Maeda K, Miyagawa J, Hotta K,
5 Shimomura I, Nakamura T, Miyaoka K, Kuriyama H, Nishida M, Yamashita S,
6 Okubo K, Matsubara K, Muraguchi M, Ohmoto Y, Funahashi T, Matsuzawa Y.
7 Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity.
8 *Biochem Biophys Res Commun*. 1999;257:79-83
- 9 39. Aprahamian TR, Sam F. Adiponectin in cardiovascular inflammation and
10 obesity. *Int J Inflamm*. 2011;2011:376909
- 11 40. Chen B, Lam KS, Wang Y, Wu D, Lam MC, Shen J, Wong L, Hoo RL, Zhang
12 J, Xu A. Hypoxia dysregulates the production of adiponectin and plasminogen
13 activator inhibitor-1 independent of reactive oxygen species in adipocytes.
14 *Biochem Biophys Res Commun*. 2006;341:549-556
- 15 41. Ivanova EA, Parolari A, Myasoedova V, Melnichenko AA, Bobryshev YV,
16 Orekhov AN. Peroxisome proliferator-activated receptor (ppar) gamma in
17 cardiovascular disorders and cardiovascular surgery. *J Cardiol*. 2015;66:271-
18 278
- 19 42. Yun Z, Maecker HL, Johnson RS, Giaccia AJ. Inhibition of ppar gamma 2
20 gene expression by the hif-1-regulated gene dec1/stra13: A mechanism for
21 regulation of adipogenesis by hypoxia. *Dev Cell*. 2002;2:331-341
- 22 43. Kern PA, Di Gregorio G, Lu T, Rassouli N, Ranganathan G. Perilipin
23 expression in human adipose tissue is elevated with obesity. *J Clin Endocrinol*
24 *Metab*. 2004;89:1352-1358
- 25 44. Wang Y, Sullivan S, Trujillo M, Lee MJ, Schneider SH, Brolin RE, Kang YH,
26 Werber Y, Greenberg AS, Fried SK. Perilipin expression in human adipose
27 tissues: Effects of severe obesity, gender, and depot. *Obes Res*.
28 2003;11:930-936
- 29 45. Gardiner C, Shaw M, Hole P, Smith J, Tannetta D, Redman CW, Sargent IL.
30 Measurement of refractive index by nanoparticle tracking analysis reveals
31 heterogeneity in extracellular vesicles. *J Extracell Vesicles*. 2014;24:25361
- 32 46. van der Pol E, Coumans FAW, Grootemaat AE, Gardiner C, Sargent IL,
33 Harrison P, Sturk A, van Leeuwen TG, Nieuwland R. Particle size distribution
34 of exosomes and microvesicles determined by transmission electron
35 microscopy, flow cytometry, nanoparticle tracking analysis, and resistive pulse
36 sensing. *JTH*. 2014;12:1182-1192

1 **Key Words**

2 Adipocyte, adipokine, obesity, endothelial dysfunction, vesicles.

Materials and Methods

Size Exclusion Chromatography

EVs from 3T3-L1 adipocytes treated with 30 ng/mL TNF- α for 24-hours were collected by differential ultracentrifugation; cell medium was centrifuged at 4 °C at 1,000 *g* (5 minutes), 15,000 *g* (15 minutes), and finally 100,000 *g* (1 hour) to pellet EVs. EVs were re-suspended in 1.0 mL of 0.22 μ m sterile-filtered 1X PBS which was then loaded onto a size-exclusion chromatography column (Exo-spinTM midi columns; Cell Guidance Systems; EX04-20). Thirty 0.5 mL 1X PBS fractions were collected and then Nanoparticle Tracking Analysis, Nanodrop protein concentration and Western blotting assays performed. Nanoparticle Tracking Analysis and Western Blotting were performed as described in the manuscript. The primary antibodies FABP4 (3544S, CST), CD63 (sc-15363, SantaCruz) and TNF- α (6945; CST) were used at a 1:500 dilution. (N=1, n=1). Protein concentration was determined using a NanoDrop 1000 Spectrophotometer (Thermo Scientific).

Transmission Electron Microscopy

3T3-L1 EVs were resuspended in 1X 0.22 μ m-filtered BPS and then fixed with an equal volume of 4 % (v/v) paraformaldehyde and kept in at 4°C until processing for Transmission Electron Microscopy (TEM). Briefly, EVs (10 μ l) were adsorbed onto glow discharged carbon formvar 200 mesh copper grids for 2 minutes, Grids were then blotted using filter paper, stained for 10 seconds with 2% (w/v) uranyl acetate before surplus stain was removed and grids were air dried. Grids were imaged using a FEI Tecnai 12 TEM at 120 kV fitted with a Gatan OneView CMOS camera. (N=1, n=1).

Flow Cytometry Gating Strategy

Acquisition was terminated upon recording 10,000 events and events gated based on their forward scatter and side scatter characteristics, as we have described in detail previously for HUVEC ¹.

Results

Size Exclusion Chromatography

A peak in particle concentration is observed between fractions 6-8 (Supplementary Figure 1A). Protein content shows low levels across the 30 fractions (Supplementary Figure 1B). The protein content is highest in fractions 7 and 8. TNF- α appears in fractions 6-8 which also contain CD63 and FABP4 and does not appear in later fractions where soluble protein would elute (Supplementary Figure 1C). This provides evidence that a small level of TNF- α is strongly associated with EVs. This also confirms that the majority of soluble TNF- α added to the initial cell culture is removed during washing/centrifugation steps

TEM Electron Microscopy








3T3-L1 EVs display typical cup-shape morphology (Supplementary Figure 2).







Flow Cytometry Gating Strategy





Unstained HUVEC and single positive CD144-PE fluorescence (using FL-2 channel) or single positive adhesion marker (using FL-4 channel) were analysed to ensure compensation. Double stained CD144 (PE) +ve and VCAM-1 (APC) +ve was used to confirm HUVEC phenotype (Supplementary Figure 3). Cytometric gating for cell adhesion molecule profile. A FSC versus SSC region was set to exclude cell aggregates and debris. Representative dot plots showing fluorescence-minus-one for unstained cells, CD144+ cells, VCAM-1 + cells, and CD144+ and VCAM-1 + cells.



References














Burnley-Hall N, Willis G, Davis J, Rees DA, James PE. Nitrite-derived nitric oxide reduces hypoxia-inducible factor 1 α -mediated extracellular vesicle production by endothelial cells. *Nitric Oxide* 2017;63:1-12.



A**B**



A**B****C****D**

A**B**






1

'Inflammatory' and 'Inflammatory & Hypoxic' adipocytes release TNF- α -containing extracellular vesicles (EVs)




Image Key:



Adipocyte EVs



Leukocytes



Endothelial Cells



Erythrocytes




VCAM-1 protein

2

Following entry of these EVs into the circulation, there is an increase in vascular endothelial cell VCAM-1 production


3

This is followed by increased leukocyte attachment




N.b. Image not to scale.

C



B



C

