Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/120912/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Cardoso, Diogo N., Ferreira, Nuno G. C. ORCID: <https://orcid.org/0000-0002-3749-3836>, Tourinho, Paula S., Santos, Cátia, Soares, Amadeu M. V. M. and Loureiro, Susana 2019. Mercury accumulation from food decreases collembolans' growth. *Science of the Total Environment* 668 , pp. 25-31. 10.1016/j.scitotenv.2019.02.392 file

Publishers page: <http://dx.doi.org/10.1016/j.scitotenv.2019.02.392>
<<http://dx.doi.org/10.1016/j.scitotenv.2019.02.392>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.

Mercury accumulation from food decreases collembolans' growth

Diogo N. Cardoso¹, Nuno G. C. Ferreira¹, Paula S. Tourinho¹, Cátia Santos^{1,2}, Amadeu M.V.M. Soares¹ and Susana Loureiro¹

¹Department of Biology & Centre for Environmental and Marine Studies (CESAM), University of Aveiro, 3810-193 Aveiro, Portugal.

²Department of Biology, Terrestrial Ecology Unit, Ghent University, K.L. Ledeganckstraat 35, 9000, Ghent, Belgium

Corresponding author:

Diogo N. Cardoso

Address: Department of Biology & Centre for Environmental and Marine Studies (CESAM), University of Aveiro, 3810-193 Aveiro, Portugal.

E-mail: dfilipe@ua.pt

Tel: +351 234 370779

*Graphical Abstract

growth

bioaccumulation

Highlights

- Exposure to mercury through food reduces the growth of collembolans;
- Mercury uptake in the organisms through a dietary exposure increases Hg inside organisms;
- A possible maximum sub lethal concentration of 0.07 ng Hg/g for collembolans was found;
- Food avoidance may occur in the presence of contaminated food at high Hg levels;
- At higher mercury levels, collembolans presented a higher elimination rate.

Mercury accumulation from food decreases collembolans’ growth

Diogo N. Cardoso¹, Nuno G. C. Ferreira¹, Paula S. Tourinho¹, Cátia Santos^{1,2}, Amadeu M.V.M. Soares¹ and Susana Loureiro¹

¹Department of Biology & Centre for Environmental and Marine Studies (CESAM), University of Aveiro, 3810-193 Aveiro, Portugal.

²Department of Biology, Terrestrial Ecology Unit, Ghent University, K.L. Ledeganckstraat 35, 9000, Ghent, Belgium

Corresponding author:

Diogo N. Cardoso

Address: Department of Biology & Centre for Environmental and Marine Studies (CESAM), University of Aveiro, 3810-193 Aveiro, Portugal.

E-mail: dfilipe@ua.pt

Tel: +351 234 370779

Abstract

In the terrestrial environment mercury (Hg) is redistributed and transformed into different inorganic and metal - organic species that are deposited into soils. In the present study, the effects of contaminated food with environmentally relevant concentrations of Hg in the form of HgCl_2 on the soil-dwelling collembolan *Folsomia candida* were assessed. Changes in growth rate and Hg bioaccumulation levels were observed at different concentrations of Hg in food complementing data on the effects of Hg on reproduction and survival using standardized protocols. Collembolan growth was recorded every two days, and a Von Bertalanffy's growth curve was derived along with the growth rate. Collembolan growth was dependent on the Hg food concentration. Also, the final length of animals was affected by the presence of Hg in food, with differences in all treatments comparing to non-exposed organisms. Toxicokinetic patterns from Hg exposure in food at different concentrations were not significantly different from each other in the uptake, but differences were found in the depuration phase. Combining the two approaches, collembolans seems to invest their effort in the depuration process, neglecting their efforts in other vital processes, such as growth. Also, metal contaminated food avoidance possibly occurred, a behavior already reported in the literature, thus decreasing their feeding and contaminant intake. Therefore, growth tests can act an important asset to fill the gaps of bioaccumulation tests and reproductive assays, towards a mechanistic understanding. Changes in growth rate, even at low and environmentally relevant concentrations, could be a warning signal when occurring in species with key roles in ecosystems. Also, this study highlights the importance of these complementary tests for a better and complete approach to risk assessment studies.

Key-words: Mercury, contaminated food, bioaccumulation, Von Bertalanffy's growth curve

Capsule

This study is a step forward on the understanding of mercury exposure effects to soil organisms, using growth and bioaccumulation in time as endpoints.

1 - Introduction

Mercury (Hg) pollution is a worldwide problem, posing a serious threat to ecosystems and consequently to humans, due to its ability to biomagnify along food chains (Boening, 2000). Being a class B metal (Nieboer and Richardson, 1980), Hg has high affinity for reduced sulphur atoms such as those from proteins and peptides containing thiol, which could lead to a disruption of the tertiary structure of proteins, necessary for optimal function and condition of organisms (Valko et al., 2005). Furthermore, Hg can also affect organisms at a cellular level, depleting cellular antioxidation systems, which can lead to the production of reactive oxygen species inducing oxidative stress damage - (e.g., peroxidation of membrane lipids) - (Stohs and Bagchi, 1995). Since Hg is a widely-distributed contaminant, it is transported by air and deposited in areas far from emission sources (Pacyna et al., 2009), on the top-soil layer, and a consequent redistribution and further uptake by vegetation may occur (Miller et al., 2005). Besides that, Hg can be retained in soils for a period of 500 to 1000 years (Pendias et al., 2011), and suffer transformations into organic forms (most commonly methylmercury), through microbial activity, thus becoming more toxic to organisms (Wang et al., 2003). Organic and inorganic forms of Hg will eventually enter food webs and bioaccumulate and biomagnify in soil invertebrates (Pedrini-Martha et al., 2012; Wiener et al., 2006).

Despite the growing concern about the potential adverse effects of elevated mercury concentrations in the environment, the toxicity data available for soil invertebrates is scarce. Some examples are studies in earthworms (Abbasi and Soni, 1983; Beyer et al., 1985; Buch et al., 2017; Fischer and Koszorus, 1992; Mahbub, et al., 2017.), millipedes (Buch et al., 2018),

collembolans and enchytraeids (Buch et al., 2016; Lock and Janssen, 2001). Since those studies were conducted in different substrates, some toxicity differences were observed, as soil characteristics have a significant influence on the bioavailability and toxicity of metals. High pH values and organic matter content could lead to an increase in chemical sorption and consequently a decrease in bioavailability and toxicity (Lock and Janssen, 2001; Sandifer and Hopkin, 1996). The interactions of Hg with temperature were also reported, reducing the cold tolerance of earthworms and collembolans in the presence of Hg, with a dominant synergistic pattern between these two stressors (Bindesbøl et al., 2009; Holmstrup et al., 2008)

Current recommended ecotoxicological tests with the springtail *Folsomia candida* provide useful data regarding their reproduction output and survival, upon a 28-day exposure (ISO, 1999; ISO, 2014). Xenobiotics exposure to collembola is mainly carried out and assessed through the interstitial soil pore water (Fountain and Hopkin, 2005), but the evaluation of other routes of exposure are essential in risk assessment studies. Contaminated food exposure can, therefore, complement the information provided by the already existing standardized tests, facilitating, for example, the observation of growth, and providing additional information on xenobiotics' effects on organisms (Roex et al., 2003). It is widely known that the primary dietary component of most springtails is fungi (Van Straalen and Van Meerendonk, 1987), and most fungal species accumulate metals (e.g., mercury) in their hyphae (Bengtsson et al., 1983), being food a pertinent source of contaminants. Also, using growth as an endpoint is of utmost importance for the equilibrium and sustainability of any species, as a feature closely related to the organism reproduction (Fountain and Hopkin, 2001). In addition, the exposure of xenobiotics through food can also provide new insight on toxicity effects to soil invertebrates as their central role in the ecosystem is related to decomposition processes (Crouau and Moïa, 2006). In an organism, growth is dependent upon food quality availability and the metabolism efficiency, which can be affected by the presence of xenobiotics/contaminants, being,

therefore, a relevant endpoint to assess environmental hazard (Crommentuijn et al., 1993; Folker-Hansen et al., 1996).

Considering the above mentioned, this study aimed at assessing the effects of Hg on *Folsomia candida* growth and to infer if Hg bioaccumulation patterns were related to the observed growth patterns. For that, organisms were exposed to Hg-contaminated food at different relevant concentrations (1 to 4 ppm) for a 28 days' exposure period. During this period collembolan growth was recorded every two days, and a Von Bertalanffy growth curve derived along with the correspondent growth rate. At the same time, a similar experimental setup was conducted to derive bioaccumulation patterns and toxicokinetic parameters (uptake and elimination rates) also at different exposure concentrations. These two experiments complemented each other to understand the mechanisms of toxicity that Hg can induce in *F. candida*.

2 – Material and methods

2.1 - Test species and test chemical

Collembolans from the species *F. candida* were obtained from synchronized laboratory cultures maintained at the University of Aveiro, Portugal. Cultures are kept in plastic boxes lined with a mixture of plaster of Paris and activated charcoal in a ratio of 9:1 (ISO, 2014), in the dark, under a constant temperature regime (20 ± 2 °C). Once a week, granulated dry yeast was added as a food source (Fermipan, Setúbal, Portugal).

The same yeast provided in cultures was contaminated with mercury (II) chloride (HgCl₂ - CAS no: 7487-94-7) purchased from Merck Millipore (99.5 % purity).

2.2 – Growth test with *Folsomia candida*

The growth test was run in two phases: 1) an exposure phase, where organisms were exposed to contaminated yeast for 28 days and 2) a post-exposure phase, where organisms were moved to clean vials with clean food for 14 days. The test started with 10 days old collembolans that were exposed to Hg via food, provided as contaminated yeast with 1, 2, 3 and 4 ppm of Hg. The chosen concentrations are environmental relevant since concentrations ranging from 0.003 to 4.6 mg/kg of Hg have been reported in soils worldwide (E. Steinnes, 1997). Also, the mean content of Hg in soils was estimated to be 1.1 mg/kg, with an average background content in different soil types ranging from 0.58 to 1.8 mg/kg (Xu et al., 2015). Each replicate included one organism in a cylindrical plastic pot (30 cm³), on a moist substrate of 9:1 (w/w) plaster of Paris:charcoal mixture. Granulated dry yeast was previously spiked with the respective Hg solutions and dried by a lyophilization process. Afterward, it was supplied *ad libitum* during the exposure phase and being replaced every week or in the presence of fungi. A total of 10 replicates were used per control, Hg treatments and sampling time. Digital photographs were taken every two days for 42 days, and the collembolan's length was recorded from the end of the posterior abdominal segment to the anterior margin of the head, using the Image J analysis software program (Schneider et al. 2012). Calibration run by using a small millimeter paper placed near the measured organism.

Considering that collembolan finished the test with 52 days old, the appearance and number of new-born juveniles were also recorded.

2.3 – Bioaccumulation test with *Folsomia candida*

Running at the same time as the growth test, 18 replicates with 20 organisms each per Hg concentration were also exposed to the same Hg concentrations following similarly the procedures described in section 2.2. Briefly, organisms were exposed to contaminated food (1, 2, 3 and 4ppm Hg) for 28 days, followed by a non-contaminated yeast exposure for an extra 14

days' period. Organisms were sampled after 7, 14, 21, 28 days during the exposure period and at day 35 and 42 (belonging to the post-exposure test), sacrificing 3 replicates in each sampling time for bioaccumulation analysis (3 replicates used in each one of the 6 sampling times, a total of 18 replicates initially performed). Samples were immediately frozen at -80 °C for following chemical analysis. Before the measurements of total Hg inside the organisms were dried by a lyophilization process.

2.4 - Mercury analysis

Total mercury (THg) concentration in yeast (food source) and collembolans was quantified by atomic absorption spectrophotometry with thermal decomposition using an Advanced Mercury Analyser (AMA) LECO 254 (Costley et al., 2000). The analytical procedure was adapted as described by Cabecinhas et al. (2015) and Vieira et al. (2014): drying time of 60 s, decomposition time of 150 s and waiting time of 45 s. The detection limit of the procedure was 0.02 ng Hg, and first and second working range of analysis (automatically switched) varied between 0.05 – 40 ng Hg and 40-600 ng Hg, respectively.

Mercury determinations were all done in triplicate and blank run in parallel before and after each sample to ensure the equipment was clean from any internal Hg contamination. All blank measurements were run, at least, in triplicate until values obtained were consistently inferior to the equipment detection limit (0.02 ng of Hg); additional blanks were, always performed at the beginning and end of the day.

Accuracy and precision of the analytical method were assessed by the analysis of certified reference materials (NRC TORT-2 Lobster hepatopancreas, and NRC DOLT-3 Dogfish liver) in parallel with samples (Pereira et al., 2008). The average recovery of TORT-2 and DOLT-3 were 101 and 104%, respectively.

2.5 - Data analysis

Growth was described by the Von Bertalanffy expression:

$$L_t = L_{\infty} (1 - e^{-k(t-t_0)}) \quad (1)$$

Where L_t = length at time t (mm), L_{∞} = the asymptotic length (mm), k = growth rate and t_0 = the hypothetical negative time estimated from hatching date for an individual with length 0 (days) (Bertalanffy, 1960). The theoretical growth curve was fitted to the average length of individuals per replicate and the parameters L_{∞} and k were estimated. For comparisons between the effects of Hg treatments and the negative control, slopes of the regression and growth rate values (k) were compared by a generalized likelihood ratio test. Differences in collembolans' length exposed to different Hg treatments, in different days of exposure were analyzed using a one-way ANOVA, followed by post hoc Dunnett's test ($p < 0.05$), comparing to the correspondent negative control (in time). To infer on the possible effects of Hg exposure in collembolan's reproduction output, the day at which newborn juveniles were first observed in each test box were compared between treatments and the control by using a Kruskal-Wallis test followed by Dunn's *post hoc* test. Differences in the weight of collembolans at different sampling times were derived by a one-way ANOVA, followed by post hoc Dunnett's test ($\alpha=0.05$). A two-way ANOVA followed by a Tukey Test ($\alpha=0.05$) was also used to observe statistical differences in the weight of collembolans in the different concentrations of Hg, at different sampling times, also looking at the interaction between these two factors. The R-squared (r^2) was obtained by dividing the sum of squares of each factor and of their interaction by the total sums of squares of the two-way ANOVA assessing the percentage of variance accounted for each factor in the ANOVAs.

For the bioaccumulation test, the toxicokinetics of Hg was described by applying a one-compartment model. Uptake and elimination equations were fitted simultaneously. In the

model, body concentration of Hg was used, and the initial Hg concentration in the organisms ($Q_{(0)}$) was considered to be zero.

For the uptake phase, the following equation was used:

$$Q_{(t)} = \frac{k_1}{k_2} \cdot C_e \cdot (1 - e^{-k_2 \cdot t}) \quad (2)$$

Where $Q_{(t)}$ - Hg body concentrations at t days (ng/mg); k_1 – uptake rate constant ($\text{kg}_{\text{food}}/\text{kg}_{\text{org}} \cdot \text{day}^{-1}$); k_2 – elimination rate constant (day^{-1}); C_e – Hg exposure concentration (ng/mg); and t - time (days).

For the elimination phase, the metal body burden was described as:

$$Q_{(t)} = \frac{k_1}{k_2} \cdot C_e \cdot (e^{-k_2 \cdot (t-t_c)} - e^{-k_2 \cdot t}) \quad (3)$$

where t_c - time the animals are transferred to clean medium (days).

Differences in k_1 and k_2 values between concentrations were tested by a Generalized Likelihood Ratio test (GLR test). The assimilation rate (a) was calculated as $k_1 \cdot C_e$, and the bioaccumulation factor (BAF) was calculated as k_1/k_2 . Also, $\text{BAF}_{\text{org/food}}$ was also calculated as the Concentration of Hg in organisms/concentration of Hg in food although no equilibrium was reached (plateau) during the uptake phase. Hg half-lives (DT_{50}) in the collembolans were calculated as $\ln 2/k_2$. All calculations were performed using SPSS (version 20).

3 – Results and discussion

3.1 Hg measurements in food

The measured concentrations in yeast were 0.918, 2.03, 2.87 and 3.9 ppm of Hg for the nominal concentrations chosen of 1, 2, 3, and 4 ppm, respectively. Since differences between nominal and measured values were lower than 10%, nominal concentrations were used throughout the manuscript. However, measured concentrations were used in the data analysis.

3.2 - Growth test with *Folsomia candida*

Collembolan growth was dependent on Hg concentration in food as observed in Figure 1. A Von Bertalanffy growth curve was fitted to the data in all cases, and the values of L_{∞} and K derived (Table 1). For each Hg treatment and control, two curves were settled, and data are presented separately in Table 1-A for data for the exposure phase only (first 28 days of the experiment), and in Table 1-B for the 42-day period, including both exposure and post-exposure phases (28 days of exposure to Hg-contaminated food and 14 days of recovery with non-contaminated food).

Table 1 - Parameters in von Bertalanffy growth curves. Estimates of mean maximum length (L_{∞}) and growth rate (k) with asymptotic confidence intervals (C.I.) A- using growth measurements for 28 days of exposure (uptake); B – using growth measurements for 42 days (28 days of exposure followed by 14 days of recovery).

A

[Hg] in food (ppm)	L_{∞}	C.I. 95%	K	C.I. 95%	r^2
0	3.619	3.460 – 3.778	0.029	0.027 – 0.031	0.981
1	3.05	2.906 – 3.194	0.036	0.033 – 0.039	0.958
2	2.917	2.795 – 3.038	0.037	0.035 – 0.040	0.965
3	2.776	2.673 – 2.880	0.041	0.038 – 0.044	0.960
4	2.238	2.191 – 2.285	0.057	0.055 – 0.060	0.960

B

[Hg] in food (ppm)	L_{∞}	C.I. 95%	K	C.I. 95%	r^2
0	2.862	2.799 – 2.924	0.042	0.04 – 0.044	0.961
1	2.614	2.551 – 2.669	0.047	0.045 – 0.049	0.942
2	2.529	2.482 – 2.576	0.048	0.046 – 0.050	0.955
3	2.2458	2.416 – 2.500	0.051	0.049 – 0.054	0.949
4	2.106	2.083 – 2.130	0.065	0.063 – 0.068	0.949

As expected, collembolans reached their highest values of L_{∞} (maximum length) when exposed to uncontaminated food decreasing its value with the increase of Hg food concentration for the 28 days and 42 days fitting curves, as observed in Figure 1. Also, the pattern was inverse for collembolans' growth rate (k) with values increasing with the increase of Hg in food (28 and 42 days fitting curves). Curves fitted with the 28 days' dataset always showed higher values for L_{∞} and lower values for k when compared to curves fitted with the 42 days' data set. In addition, the statistical analysis showed significant differences in the growth rate within the same treatment when comparing the 28 and 42 days fitting curves (Table SD1; $p < 0.05$). Using the same approach for collembolans exposed to for both 28 or 42 days curves, the growth rate was statistically different for all treatments comparing to the control curve (Table SD1; $p < 0.05$). Due to those differences, an analysis per day of exposure of collembolans' length was performed for the 28 days of exposure. As expected, statistical differences were found regarding growth for the 4-ppm treatment already at day 12 and onward, comparing to the control (Dunnett's test, $p < 0.05$). For collembolans exposed to contaminated food with 3 and 2 ppm, statistical differences were found at day 14 and onward, while for the 1ppm treatment only at day 20 and onward differences were attained, always comparing with the control animals supplied with non-contaminated yeast (Dunnett's test, $p < 0.05$). These significant differences were maintained during the recovery period thus

showing that organisms were not completely recovering from Hg exposure even when clean food was provided (Dunnett's test, $p < 0.05$).

The time of hatched eggs was also evaluated in our study. Analyzing the hatching day of juveniles, a significant delay was observed for organisms exposed to 4 ppm Hg in food ($H_4 = 15.747$; $p = 0.003$). The mean number of days for juveniles to hatch ($\pm$ SD) in control, 1, 2, 3 and 4 ppm was respectively 22.1 (± 0.78); 21.6 (± 0.70); 22.3 (± 0.71), 22.2 (± 0.73) and 23.78 (± 1.30) days. These results are in accordance with the ones found by Fountain and Hopkins (2011) where reproduction (eggs laid in time) was delayed, due to retarded growth at high metal concentrations. This species of collembolan has an unpigmented body and a gut that crosses the organisms' body, and when they avoid food and start eating the black mixture of plaster of Paris and activated charcoal, their gut acquires a visible black color. With the increase of Hg in food the number of organisms showing a black ribbon along the body also increased (personal observation/data not shown) thus supporting a possible avoiding food behavior. Avoidance behavior is well explained by Fountain and Hopkin (2001), extrapolating that the reduced growth of collembolans, when exposed to higher concentrations of metals, could be explained to presumably use of the substrate as an alternative food source to springtails on an attempt to acquire nutrition from the graphite.

In the study of Marigomez et al. (1986), the terrestrial slug *Arion ater* was exposed to concentrations of mercury chloride ranging 10 to 100 ppm, and significant food consumption and growth reductions were observed for concentrations higher than 10 ppm in a dose-related manner. These results are in accordance with ours since an increase in Hg concentrations leads to lower growth. Another study performed by Abassi and Soni (1983) kept adult earthworms (*Octochaetus pattoni*) in cement tanks for 60 days, at the same density as the one found in the wild, in a mixture of soil and animal dung contaminated with mercuric chloride ranging from 0.5 - 5.0 ppm Hg. The calculated LC_{50} was 2.39 ppm after 10 days' decreasing to 0.79 ppm at

the end of the 60 days' exposure period. Differences in the Hg-induced toxicity could be explained by different exposure routes, since earthworms would ingest the contaminated food, but would also be exposed to the contaminated soil by dermal contact and soil ingestion.

Lock and Janssen (2001) studied the effects of Hg in three different representative soil invertebrates: the enchytraeid *Enchytraeus albidus*, the collembolan *Folsomia candida* and the earthworm *Eisenia fetida*. An EC₅₀ of 9.16 ppm was calculated for *E. fetida* (21 days of exposure), while an EC₅₀ of 22 ppm for *E. albidus* (42 days of exposure) and an EC₅₀ of 3.26 ppm for *F. candida* (28 days of exposure) were derived, based on data from reproduction output for all organisms. A similar study conducted by Liu et al. (2010), using only the species *F. candida* found a higher EC₅₀ for reproduction of 9.29 ppm and an EC₅₀ value for the avoidance of 3.88 ppm in the presence of an agricultural fluvoaquic sandy loam soil (Cambisol, 9.0% clay, 21.8% silt, 69.2% sand) spiked with HgCl₂. Also, a more recent study from Buch et al. (2016) assessed the effects of natural and artificial soils contaminated with Hg in two collembolan species (*Folsomia candida* and *Proisotoma minuta*). For *F. candida* exposed to natural soil, an AC₅₀ (avoidance behavior) of 5.44 (CI 4.13–6.75) ppm, an EC₅₀ for reproduction of 3.40 (3.18–3.62) ppm and an LC₅₀ of 6.12 (3.74–8.50) ppm were attained. Despite the apparent differences between both studies (different exposure routes and different endpoints analyzed), the range of concentrations used in our study is lower than the LC₅₀ and AC₅₀, and the highest concentration used similarly to the EC₅₀ derived by Buch et al. (2016). It should also be noticed that in that study *F. candida* was more sensitive to mercury contamination than *P. minuta*. Even knowing that different exposure routes could lead to differences regarding toxicity, our results reveal that ingestion of contaminated food could be a very sensitive endpoint and growth could be a relevant sensitive endpoint to look at.

3.3 - Bioaccumulation test with *Folsomia candida*

Before Hg analysis, 20 collembolans per replicate from each treatment and control were sampled and weighted at each sampling time. The results of the one-way ANOVA in each sampling time revealed that until day 28 (complete uptake period), no statistical differences were found in the weight of collembolans exposed to different Hg concentrations (ANOVA, $p > 0.05$). Collembolans sampled at day 35 (first sampling time in the recovery period) from the 2 and 3 ppm food exposure presented statistically higher weights than those from the control (Dunnett's test, $p < 0.05$). Surprisingly, no statistical differences were observed for the ones exposed to 4 ppm Hg in food compared to the control. Regarding the collembolans used in day 42, statistical differences in their weight were also observed, with collembola from the 4-ppm exposure showing higher weights (Dunnett's test, $p < 0.05$), on a possible attempt to ingest as much food as possible to compensate the lack of quality feeding when previously contaminated food was supplied, as suggested above. Analyzing the results with a two-way ANOVA, as expected, the time of exposure was the factor that explained the majority of the collembola weight differences (two-way ANOVA, $F_{5,88} = 106.007$, $p < 0.05$) with an R^2 of 0.82. Regarding the factor for Hg concentration no statistical effect in collembolans' weight was derived (two-way ANOVA, $F_{4,88} = 1.062$, $p > 0.05$) with an R^2 of 0.014 with a slight interaction with time of exposure (two-way ANOVA, $F_{20,88} = 2.23$, $p < 0.05$; R^2 of 0.07).

The fit of the one-compartment kinetics model and the corresponding uptake and elimination rate constants are shown in Figure 2 and Table 2. For 1 and 3 ppm treatments, Hg body concentrations did not reach equilibrium after the 28 days of exposure. Mean Hg body concentration ($n=3$) at the end of the uptake phase (28 days) was 0.02, 0.03, 0.07 and 0.07 ppm, corresponding to a total body burden of 0.07, 0.13, 0.25 and 0.26 ng of Hg at exposure concentrations of 1, 2, 3 and 4 ppm, respectively. Within this, Hg values in collembola at day 28 of exposure increased with the increase in Hg concentrations in food, with the two highest concentrations of exposure (3 and 4 ppm) producing similar body burden values.

Considering the size of collembolans decreased with increasing Hg concentrations, and their weight did not follow the same pattern, the hypotheses that an extra source of food occurred can be raised. This was somehow supported by observations of the presence of black material (possibly charcoal) in the digestive system of collembolans during the exposure to the highest Hg concentration. The hypothesis of extra ingestion of plaster of Paris and charcoal to compensate for the lack or low quality of the provided food source can be raised and corroborated with these findings. By looking at the kinetics curves patterns, collembolans exposed to 2 and 4 ppm were those deriving data that formed a (close to) plateau in the uptake phase, showing a potential equilibrium between uptake and elimination. This was also shown by the DT_{50} values at 2 and 4 ppm where Hg half-life in collembola was much lower than those from 1 and 3 ppm, revealing a faster elimination of Hg. The same pattern was also seen when comparing BAF values calculated with the kinetic parameters or through the concentrations measured.

Table 2 - Hg uptake and elimination rate constants (k_1 and k_2), bioaccumulation factor (BAF)_{kinetics} and $BAF_{org/food}$ and time that takes for Hg to be reduced by half inside the organisms (DT_{50}) for the accumulation of Hg in *Folsomia candida* exposed for 28 days to Hg spiked food, followed by an elimination phase of 14 days. The 95 % confidence intervals are presented in brackets.

[Hg] in food (ppm)	K_1 ($kg_{food}/kg_{org}/day$)	K_2 (day^{-1})	$BAF_{kinetics}$ (kg_{food}/kg_{org})	$BAF_{[Hg]org/[Hg]food}$ (kg_{food}/kg_{org})	DT_{50} (days)
1	0.0011 (0.001 - 0.002)	0.0307 (-0.001 - 0.062)	0.0359	0.018	22.6
2	0.0020 (0.001 - 0.003)	0.1038 (0.057 - 0.150)	0.0198	0.0165	6.7
3	0.0014 (0.001 - 0.002)	0.0456 (0.031 - 0.061)	0.0312	0.022	15.2
4	0.0016 (0.001 - 0.002)	0.0972 (0.068 - 0.126)	0.0173	0.017	7.1

Uptake rate constants (k_1) were compared using the likelihood-ratio test. The statistical analysis did not show any significant difference between treatments, thus showing that Hg

uptake and accumulation was concentration independent. The same procedure was conducted for the elimination rate constant (k_2) and statistical differences were observed between concentrations of 1 and 2 ppm ($\chi^2_{(1)} = 5.766$ $p < 0.05$), 1 and 4 ppm ($\chi^2_{(1)} = 5.34$ $p < 0.05$), 2 and 3 pp ($\chi^2_{(1)} = 8.10$ $p < 0.05$), and between 3 and 4 ppm ($\chi^2_{(1)} = 13.04$ $p < 0.05$). These differences in k_2 revealed that collembolans are eliminating Hg faster as Hg exposure increases.

Collembolans exposed to the lowest concentration of Hg (1 ppm) in food showed a low Hg uptake (< 0.02 ng/g), which is in accordance with the no effects observed in their growth during the test. At the highest concentrations (3 and 4 ppm) a maximum of approximately 0.07 ng/g was attained, which could be close to the maximum sub-lethal concentrations that collembolans can tolerate. Organisms exposed to 4 ppm had to compensate food intake with extra food available (as mentioned above) as possibly they could not tolerate any further increase of Hg in their body. Although this avoidance behavior towards the contaminated food along with the ingestion of plaster led to an increase in their weight, no energy could be allocated from the plaster towards their growth (expressed as body size).

The comparison of these data with literature is difficult as studies on mercury bioaccumulation in soil-dwelling organisms are scarce, and toxicity studies are based on soil exposure using reproduction as a preferred endpoint. However, as stated before, Fountain and Hopkin found an avoidance behavior to Cd, Cu, Pb, and Zn exposures in food, at high concentrations. Although they did not assess metal bioaccumulation, graphite was found in the digestive tract which supports our data regarding food compensation. In the study of Zhang et al. (2009) earthworms showed to accumulate high levels of Hg from a polluted soil with total and methylmercury concentrations differing between earthworms' species. The comparison between earthworms (or other species) and collembolans should be analyzed carefully since the routes of exposure may be different. Also, the exposure route may be constant or avoidable, like is the case of food, which can be avoided, or soil under a reproduction test

where avoidance is not an available option. Fountain and Hopkin (2001) also discussed the potential higher tolerance of Collembolans regarding metal exposure through diet since they can avoid contaminated food. When in contact with contaminated media they excrete metals by molting, exfoliating the midgut epithelium where the elements are retained as part of a storage detoxification system, in order to decrease metal accumulation (Köhler, 2002).

In soil ecotoxicology, the presence of soil as exposure matrix or as substrate can provide some difficulties when assessing some crucial endpoints, like growth. In the present case study, the dietary exposure was assessed in plaster of Paris/charcoal, providing continuous monitoring during the exposure period, and assessing the effects in time. This approach would not have been possible in the soil, due to the difficulty of spotting each collembola and following its growth (in time).

4 - Conclusions

This study provides a new approach to understand Hg-induced effects in *F. candida*, by combining an ecotoxicological endpoint (growth) and with Hg bioaccumulation following a timeline. For the first time, it is reported the Hg-induced effects on growth and bioaccumulation from a dietary source to the springtail *F. candida*. Even at low and environmental relevant Hg concentrations, collembola growth rate was impaired. Despite a clear difference in the growth of collembolans with increasing Hg concentrations, collembolans did not reach a complete steady state in all concentrations of the uptake phase, being independent of the Hg exposure concentration.

Regarding depuration, after exposure to the higher Hg concentration, collembolans showed an increase in the k_2 , with a faster elimination and decreasing Hg residence time. Also, by comparing collembolans' length and weight at different Hg concentration, along with their bioaccumulation patterns, it could be highlighted a potential food avoidance and exploring a

different food source available (plaster with charcoal). This was perceived by the dark color of collembola exposed to the highest concentrations. It was also highlighted a possible maximum sub-lethal concentration of 0.07 ng Hg/g of collembolan. The present study can be considered as a step forward in the assessment of the potential risks of mercury or other metals in terrestrial environments, also providing a mechanistic approach looking simultaneously at toxicity and bioaccumulation.

5 - Acknowledgments

This work was supported by CESAM (UID/AMB/50017/2013) through FCT/MEC (through national funds) and the co-funding by the FEDER (POCI-01-0145-FEDER-00763), within the PT2020 Partnership Agreement and Compete 2020. Diogo Cardoso was supported by a FCT PhD grant (SFRH/BD/52569/2014) within the Doctoral Programme Biology and Ecology of Global Changes. The authors would like to thank the laboratory support given by Dr. Abel Ferreira.

6- Bibliography

- Abbasi, S. A., Soni, R., 1983. Stress-induced enhancement of reproduction in earthworm *Octochaetus pattoni* exposed to chromium (vi) and mercury (ii)— implications in environmental management. *International Journal of Environmental Studies*. 22, 43-47.
- Bengtsson, G., et al., 1983. Growth Changes Caused by Metal Uptake in a Population of *Onychiurus armatus* (Collembola) Feeding on Metal Polluted Fungi. *Oikos*. 40, 216-225.
- Bertalanffy, L., Principles and theory of growth. *Fundamental Aspects of Normal and Malignant Growth*, ed. WW Nowinski. Amsterdam: Elsevier publ. co.(Abstr.), 1960.
- Beyer, W. N., et al., 1985. Accumulation of methylmercury in the earthworm, *Eisenia foetida*, and its effect on regeneration. *Bulletin of Environmental Contamination and Toxicology*. 35, 157-162.
- Bindesbøl, A.-M., et al., 2009. Impacts of heavy metals, polyaromatic hydrocarbons, and pesticides on freeze tolerance of the earthworm *Dendrobaena octaedra*. *Environmental Toxicology and Chemistry*. 28, 2341-2347.
- Boening, D. W., (2000). Ecological effects, transport, and fate of mercury: a general review. *Chemosphere*. 40, 1335-1351.

439 Buch, A. C., et al., (2016). Ecotoxicity of mercury to *Folsomia candida* and *Proisotoma minuta*
440 (Collembola: Isotomidae) in tropical soils: Baseline for ecological risk assessment .
441 Ecotoxicology and Environmental Safety. 127, 22-29.

442 Buch, A. C., et al. (2017). "Ecotoxicology of mercury in tropical forest soils: Impact on
443 earthworms." Science of The Total Environment 589: 222-231.

444 Buch, A. C., et al. (2018). "Food preference and ecotoxicological tests with millipedes in litter
445 contaminated with mercury." Science of The Total Environment 633: 1173-1182.

446 Cabecinhas, A. S., et al., 2015. Sensitivity of the sea snail *Gibbula umbilicalis* to mercury
447 exposure – Linking endpoints from different biological organization levels.
448 Chemosphere. 119, 490-497.

449 Coelho, J. P., et al., (2005). Macroalgae response to a mercury contamination gradient in a
450 temperate coastal lagoon-Ria de Aveiro. Estuarine, Coastal and Shelf Science, 65, 492–
451 500.

452 Costley, C. T., et al., 2000. Determination of mercury in environmental and biological samples
453 using pyrolysis atomic absorption spectrometry with gold amalgamation. Analytica
454 Chimica Acta. 405, 179-183.

455 Crommentuijn, T., et al., 1993. Influence of Cadmium on Life-History Characteristics of
456 *Folsomia candida* (Willem) in an Artificial Soil Substrate. Ecotoxicology and
457 Environmental Safety. 26, 216-227.

458 Crouau, Y., Moia, C., 2006. The relative sensitivity of growth and reproduction in the springtail,
459 *Folsomia candida*, exposed to xenobiotics in the laboratory: An indicator of soil
460 toxicity. Ecotoxicology and Environmental Safety. 64, 115-121.

461 Diogo, J. B., et al., 2007. Tolerance of genetically characterized *Folsomia candida* strains to
462 phenmedipham exposure. Journal of Soils and Sediments. 7, 388-392.

463 E. Steinnes, "Mercury," in Heavy Metals in Soils, B. J. Alloway, Ed., pp. 245–259, Blackie
464 Academics and Professional Press, London, UK, 2nd edition, 1997.

465 Fischer, E., Koszorus, L., 1992. Sublethal effects, accumulation capacities and elimination rates
466 of As, Hg and Se in the manure worm, *Eisenia fetida* (Oligochaeta, Lumbricidae).
467 Pedobiologia. 36, 172-178.

468 Folker-Hansen, P., et al., 1996. Effect of Dimethoate on Body Growth of Representatives of the
469 Soil Living Mesofauna. Ecotoxicology and Environmental Safety. 33, 207-216.

470 Fountain, M. T., Hopkin, S. P., 2001. Continuous Monitoring of *Folsomia candida* (Insecta:
471 Collembola) in a Metal Exposure Test. Ecotoxicology and Environmental Safety. 48,
472 275-286.

473 Fountain, M. T., Hopkin, S. P., 2005. *FOLSOMIA CANDIDA* (COLLEMBOLA): A "Standard" Soil
474 Arthropod*. Annu. Rev. Entomol. 50, 201-222.

475 Holmstrup, M., et al., 2008. Exposure to mercury reduces cold tolerance in the springtail
476 *Folsomia candida*. Comparative Biochemistry and Physiology C-Toxicology &
477 Pharmacology. 148, 172-177.

478 ISO, 1999. Soil quality – inhibition of reproduction of Collembola (*Folsomia candida*) by soil
479 pollutants., ISO/DIS 22030, Geneva, Switzerland., 1999.

480 ISO, 2014. ISO 11267, Soil quality -- Inhibition of reproduction of Collembola (*Folsomia*
481 *candida*) by soil contaminants. International Organization for Standardization. Geneve.

482 Kabata-Pendias, A., 2011. Trace elements in soils and plants, 4th ed. CRC Press, Taylor and
483 Francis Group, USA, p. 534

484 Köhler, H.-R., 2002. Localization of metals in cells of saprophagous soil arthropods (Isopoda,
485 Diplopoda, Collembola). Microscopy Research and Technique. 56, 393-401.

486 Liu, Y.-R., et al., 2010. Effects of mercury on reproduction, avoidance, and heat shock protein
487 gene expression of the soil springtail *Folsomia candida*. Environmental Toxicology and
488 Chemistry. 29, 654-659.

- Lock, K., Janssen, C., 2001. Ecotoxicity of mercury to *Eisenia fetida*, *Enchytraeus albidus* and *Folsomia candida*. *Biology and Fertility of Soils*. 34, 219-221.
- Mahbub, K. R., et al. (2017). "Mercury toxicity to *Eisenia fetida* in three different soils." *Environmental Science and Pollution Research* 24(2): 1261-1269.
- Marigomez J. A., et al., 1986. Feeding and growth responses to copper, zinc, mercury and lead in the terrestrial gastropod *arionater* (Linne) *Journal of Molluscan Studies*. 52, 68-78.
- Miller, E., et al., 2005. Estimation and Mapping of Wet and Dry Mercury Deposition Across Northeastern North America. *Ecotoxicology*. 14, 53-70.
- Nieboer, E., Richardson, D. H. S., 1980. The replacement of the nondescript term 'heavy metals' by a biologically and chemically significant classification of metal ions. *Environmental Pollution Series B, Chemical and Physical*. 1, 3-26.
- Pacyna, J. M., et al., 2009. Changes of emissions and atmospheric deposition of mercury, lead, and cadmium. *Atmospheric Environment*. 43, 117-127.
- Pedrini-Martha, V., et al., 2012. Patterns of Urban Mercury Contamination Detected by Bioindication With Terrestrial Isopods. *Archives of Environmental Contamination and Toxicology*. 63, 209-219.
- Pereira, E., et al., 2008. Evaluation of an interlaboratory proficiency-testing exercise for total mercury in environmental samples of soils, sediments and fish tissue. *TrAC Trends in Analytical Chemistry*. 27, 959-970.
- Roex, E. W. M., et al., 2003. Acetylcholinesterase inhibition and increased food consumption rate in the zebrafish, *Danio rerio*, after chronic exposure to parathion. *Aquatic Toxicology*. 64, 451-460.
- Sandifer, R.D., Hopkin, S.P., 1996. Effects of pH on the toxicity of cadmium, copper, lead and zinc to *Folsomia candida* Willen, 1902 (Collembola) in a standard laboratory test system. *Chemosphere* 33, 2475-2486
- Schneider, C.A., Rasband, W.S., Eliceiri, K.W. "NIH Image to ImageJ: 25 years of image analysis". *Nature Methods* 9, 671-675, 2012.
- SPSS, 1997. SigmaScan 5. SPSS Inc., Chicago.
- Stohs, S. J., Bagchi, D., 1995. Oxidative mechanisms in the toxicity of metal ions. *Free Radical Biology and Medicine*. 18, 321-336.
- Tipping, E., Poskitt, J.M., Lawlor, A.J., Wadsworth, R.A., Norris, D.A., Hall, J.R., 2011. Mercury in United Kingdom topsoils; concentrations, pools, and critical limit exceedances. *Environ. Pollut.* 159, 3721-3729.
- Valko, M., et al., 2005. Metals, Toxicity and Oxidative Stress. *Current Medicinal Chemistry*. 12, 1161-1208.
- Van Straalen, N. M., Van Meerendonk, J. H., Biological half-lives of lead in *Orchesella cincta* (L.) (Collembola). *Bulletin of Environmental Contamination and Toxicology*. 38, 213-219.
- Vieira, H. C., et al., 2014. Real and Potential Mercury Accumulation in Human Scalp of Adolescents: A Case Study. *Biological Trace Element Research*. 163, 19-27.
- Wang, D., et al., 2003. Accumulation and transformation of atmospheric mercury in soil. *Science of The Total Environment*. 304, 209-214.
- Wiener, J. G., et al., 2006. Mercury in Soils, Lakes, and Fish in Voyageurs National Park (Minnesota): Importance of Atmospheric Deposition and Ecosystem Factors. *Environmental Science & Technology*. 40, 6261-6268.
- Xu, J., et al., 2015. Sources and remediation techniques for mercury contaminated soil. *Environment International*. 74, 42-53.
- Zhang, Z., et al., 2009. Bioaccumulation of Total and Methyl Mercury in Three Earthworm Species (*Drawida sp.*, *Allolobophora sp.*, and *Limnodrilus sp.*). *Bulletin of Environmental Contamination and Toxicology*. 83, 937-942.

Figure
[Click here to download Figure: Figures_Final.pptx](#)

Figure 1 – Body length of *Folsomia candida* recorded during a 28 day exposure to Hg contaminated yeast (uptake phase), followed by a 14 day exposure to clean yeast (elimination phase). The negative control was clean yeast (diamond) and the vertical dashed line separates the uptake phase from the elimination phase. Data is expressed as average $\pm$ standard error.

Figure 2 - Uptake and elimination kinetics of Hg in *Folsomia candida* exposed to nominal concentrations ranging from 1 to 4 ppm in contaminated dry yeast. Uptake and elimination phases lasted for 28 and 14 days, respectively. The vertical dashed line separates the uptake phase from the elimination phase. Lines represent the modeled Hg body concentration, using the one compartment model.

Supplementary material for on-line publication only

[Click here to download Supplementary material for on-line publication only: supplementary material_final.docx](#)