

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/129588/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Thomas, John Rhidian, Robinson, Chloe V, Mrugala, Agata, Ellison, Amy R, Matthews, Emily, Griffiths, Sian W, Consuegra, Sofia and Cable, Jo 2020. Crayfish plague affects juvenile survival and adult behaviour of invasive signal crayfish. *Parasitology* 147 (6), pp. 706-714. 10.1017/S0031182020000165

Publishers page: <http://dx.doi.org/10.1017/S0031182020000165>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Crayfish plague affects juvenile survival and adult behaviour of invasive signal crayfish

Running title: Crayfish plague negatively affects signal crayfish

John Rhidian Thomas^a, Chloe V Robinson^b, Agata Mrugała^c, Amy R Ellison^{a*}, Emily Matthews^a, Siân W Griffiths^a, Sofia Consuegra^b, Jo Cable^a

^aCorresponding author. Thomasjr6@cardiff.ac.uk. School of Biosciences, Cardiff University, Museum Avenue, CF10 3AX, U. K.

^bDepartment of Biosciences, Swansea University, Singleton Park, Swansea, SA2 8PP, U. K.

^cDepartment of Ecology, Faculty of Science, Charles University, Viničná 7, Prague 2 CZ-12844, Czech Republic

*Present address: Bangor University, School of Natural Sciences, Environment Centre Wales, Bangor, LL57 2UW, U. K.

Abstract

The spread of invasive, non-native species is a key threat to biodiversity. Parasites can play a significant role by influencing their invasive host's survival or behaviour, which can subsequently alter invasion dynamics. The North American signal crayfish (*Pacifastacus leniusculus*) is a known carrier of *Aphanomyces astaci*, an oomycete pathogen that is the causative agent of crayfish plague and fatal to European crayfish species, whereas North American species are considered to be largely resistant. There is some evidence, however, that North American species, can also succumb to crayfish plague, though how *A. astaci* affects such 'reservoir hosts' is rarely considered. Here, we tested the impact of *A. astaci* infection on signal crayfish, by assessing juvenile survival and adult behaviour following exposure to *A. astaci* zoospores. Juvenile signal crayfish suffered high mortality 4-weeks post-hatching, but not as older juveniles. Furthermore, adult signal crayfish with high infection levels displayed altered behaviours, being less likely to leave the water, explore terrestrial areas and exhibit escape responses. Overall, we reveal that *A. astaci* infection affects signal crayfish to a much greater extent than previously considered, which may not only have direct consequences for invasions, but could substantially affect commercially harvested signal crayfish stocks worldwide.

Keywords: crayfish plague; *Aphanomyces astaci*; behaviour; invasive species; signal crayfish; reservoir hosts

35 Key findings

- 36 • Juvenile signal crayfish suffered mortality after exposure to *A. astaci*
- 37 • Adult signal crayfish exhibited altered behavioural responses following exposure
- 38 • Susceptibility to *A. astaci* could affect signal crayfish population management

1. Introduction

Parasites have a significant impact on communities and ecosystems by directly affecting host fitness, with subsequent impacts on population dynamics and overall biodiversity (Hudson et al., 2006; Tompkins et al., 2011; Cable et al., 2017). Despite this, parasites are a fundamental component of healthy ecosystems with wide reaching impacts, from influencing the cycle of biogeochemical nutrients to regulating host density and functional traits (Hatcher et al., 2014; Preston et al., 2016). Parasites can also influence their host's behaviour, which can in turn alter the outcome of competitive interactions, reproductive behaviour and dispersal ability (Bakker et al., 1997; Macnab and Barber, 2012; Barber et al., 2017). During invasions by non-native species to new areas, parasites can play a key role facilitating or hindering the successful spread of invaders, while potentially having catastrophic effects on other related native species (Vilcinskas, 2015).

Crayfish are freshwater crustaceans that are commercially harvested in many countries, but can also reach high densities and exert a significant impact on ecosystems, with several species having become widespread, damaging invaders (Holdich et al., 2014; James et al., 2014; Ercoli et al. 2015). For example, in Great Britain, the North American signal crayfish (*Pacifastacus leniusculus*) has become the most common crayfish species, having largely replaced the native white clawed crayfish (*Austropotamobius pallipes*, see Holdich et al., 2014; James et al., 2014). Crayfish are hosts to many parasites and symbionts, including viruses, bacteria, fungi and helminths that can cause chronic, long-term infections (Longshaw et al., 2012; Kozubíková-Balcarová et al., 2013). One such parasite, the oomycete *Aphanomyces astaci*, the causative agent of crayfish plague, is a key threat to crayfish biodiversity worldwide (Svoboda et al., 2017), having eradicated many populations of native European crayfish (Filipová et al., 2013; Kozubíková-Balcarová et al., 2014) and recent evidence suggests it may have also caused a decline in commercially harvested North American crayfish stocks (Edsman

et al., 2015; Jussila et al., 2015). This obligate parasitic oomycete penetrates host tissues (Söderhäll et al., 1978) and produces motile reproductive zoospores (Cerenius and Söderhäll, 1984), which can reach high densities (up to several hundred zoospores per litre) during a crayfish plague outbreak (Strand et al., 2014). An infected individual can release about 2700 zoospores per week (Strand et al., 2012), and this number can be much higher when the crayfish is dying or moulting (Makkonen et al., 2013; Svoboda et al., 2013).

Generally, North American crayfish species which have co-evolved with *Aphanomyces astaci*, are considered to be chronic but largely asymptomatic carriers. They combat *A. astaci* through consistent production of prophenoloxidase, which activates a melanisation cascade resulting in melanisation of hyphae that prevents their invasion into host soft tissues (Cerenius et al., 1988). Most native European crayfish, on the other hand, apparently only produce prophenoloxidase only in response to infection, which is too slow to effectively melanise the hyphae that then spread into host tissues leading to paralysis and death (Cerenius et al., 2003). The Australian yabby (*Cherax destructor*) also suffers high mortality as a result of crayfish plague, though this species shows some resistance to less virulent strains and survives longer when exposed to highly virulent strains compared to highly susceptible species (Mrugała et al., 2016). In infected European crayfish, severe behavioural changes before death include a lack of coordination and paralysis (Gruber et al., 2014), though to what extent carrier crayfish exhibit behavioural changes is largely unknown and this could play a vital role during new invasions and in commercial crayfish farms. Highly infected crayfish, for example, might be less likely to disperse, which would alter invasion success and introduction to new habitats.

Few studies have directly assessed the effect of the *A. astaci* on North American species, although there is some evidence that they can succumb to the disease and display altered behaviour if also stressed by other factors (Cerenius et al., 1988; Aydin et al., 2014; Edsman et al., 2015). Co-infection of *A. astaci* and *Fusarium* spp., for example, results in

Eroded Swimmeret Syndrome (ESS) in signal crayfish, which causes females to carry fewer eggs (Edsman et al., 2015). Mortality of adult signal crayfish has also been observed in experimental settings, though only when crayfish were exposed to very high zoospore numbers (Aydin et al., 2014). Furthermore, vertical transmission of *A. astaci* (from adults to eggs) has been reported (Makkonen et al., 2010), and little is known on how *A. astaci* might affect juvenile North American crayfish.

Here, we addressed two key issues regarding the effects of *A. astaci* on signal crayfish. First, we tested the hypothesis that juvenile signal crayfish would suffer high mortality upon infection by *A. astaci* zoospores, as it has previously been suggested that juvenile crayfish may be more susceptible to infection compared to adults (Mrugała et al., 2016). Additionally, we assessed the effect of *A. astaci* on adult signal crayfish, hypothesising that even if adults may not suffer significant mortality, behavioural changes would be apparent.

2. Materials and Methods

2.1 Signal crayfish trapping

All adult signal crayfish were collected in February and March 2017 using cylindrical traps ('Trappy Traps', Collins Nets Ltd., Dorset, UK) baited with cat food and checked daily (trapping licence: NT/CW081-B-797/3888/02). The crayfish were collected from a population displaying negligible levels of infection (maximum agent level A1) when assessed in 2014 (Derw Farm pond, Powys, Wales, SO 13891 37557; James et al., 2017). A small subset of individuals (n = 3) re-tested by qPCR (see 2.2) before the experiments began in May 2017 all revealed low levels of infection by *A. astaci*, although elevated compared to 2014 (agent level A2/A3). After removal from traps, crayfish were transferred to individual containers with 500 mL of pond water and transported to the Cardiff University Aquarium (holding licence: W C ILFA 002), where they were maintained individually in 20 L aquaria containing a plant pot

refuge, gravel and air supply delivered via an airstone. The crayfish were held at $13\pm1^{\circ}\text{C}$ under a 12 h light: 12 h dark lighting regime and fed a mixture of frozen peas and *Tubifex* bloodworm (Shirley Aquatics, Solihull, West Midlands, U.K.) once every 2 days. A 50% water change was conducted 1 h after feeding to maintain water quality and remove excess food. Crayfish were acclimatised to the laboratory for at least 4 weeks before the experiments began. Four females were carrying eggs, and upon hatching, the offspring were mixed, moved to 120 L communal aquaria and used in the juvenile infection experiment. Only male crayfish were used in the adult behavioural tests; since a relatively low number of females ($n = 6$) were caught and therefore it was not possible to test an equal number of males and females in this experiment.

2.2 *Aphanomyces astaci* culture and quantification

Crayfish in the present study were exposed to a Group B strain (Pec14) of *Aphanomyces astaci* provided by Charles University in Prague. This strain was isolated from dead *Astacus astacus* from a crayfish plague outbreak in the Černý Brook, Czech Republic (Kozubíková-Balcarová et al., 2014) and demonstrated similarly high virulence towards European *A. astacus* (see Becking et al., 2015) as the strains from Group B (PsI) used in other experimental studies (Makkonen et al., 2012; Jussila et al., 2013). The culture was maintained in Petri dishes of RGY agar (Alderman, 1982; Becking et al., 2015; Mrugała et al., 2016) and zoospores were produced according to the methodology of Cerenius et al. (1988). Briefly, 2-4 agar culture plugs ($\sim 2\text{ mm}^2$) were cut from an RGY culture and placed in flasks containing 200 mL of liquid RGY-medium. Multiple replicates were done each time in order to produce a sufficient number of zoospores. These cultures were allowed to grow at 16°C for 2-4 days on a shaker. Once sufficient mycelial growth had occurred, the cultures were washed to induce sporulation and transferred to separate flasks (containing 500 mL of distilled water). The washing was repeated in distilled water 3-4 times over ~ 8 h. Then, the cultures were incubated at $13\pm1^{\circ}\text{C}$ for 24-36

h until motile zoospores were produced. The number of zoospores was quantified using a haemocytometer.

Following both experiments, crayfish were euthanized by freezing at -20°C for 1 h. For juveniles, the whole crayfish was lysed (TissueLyser, Qiagen) and DNA extracted using a Qiagen DNeasy extraction kit (Qiagen). For adult crayfish, a section of tail-fan and soft-abdominal tissue was removed by dissection, lysed (TissueLyser, Qiagen), and both tissues were pooled (~ 20 mg total), and the DNA extracted using the same kits. Infection intensity was estimated based on the number of PCR-forming units (PFU) determined by qPCR using TaqMan MGB probes and expressed using the semi-quantitative levels A0-A7 for adults (as described by Vrålstad et al., 2009); with slight modification of the protocol as in Svoboda et al. (2014). For juveniles, infection intensity was expressed as number of PCR forming units because a direct comparison cannot be made here between juvenile (whole body) and adult (sample body) infection levels.

2.3 Juvenile crayfish infection

Here, we monitored the survival of juvenile signal crayfish that hatched in the laboratory when exposed either to *A. astaci* at 1, 10 or 100 zoospores mL⁻¹ or to a sham treatment (control). All crayfish used in this experiment hatched within a 3-day period in May 2017. The infection was conducted twice in separate experiments, the first time approximately 4 weeks after the crayfish hatched (n = 25 crayfish per zoospore treatment) and the second time after 10 weeks with different crayfish (n = 17 individuals per zoospore treatment). When the experiment began, crayfish were housed individually in 1 L pots containing distilled water with a gravel substrate for 48 h. After this acclimatisation period, the pots were spiked with 1, 10 or 100 zoospores mL⁻¹ (the control treatment was given a 20% water change). After a 24 h infection period, 80% of the water in all pots was changed. The crayfish were fed crushed algae wafers and frozen

Tubifex bloodworm (Shirley Aquatics, Solihull, West Midlands, U.K.) once every 2 days. A 50% water change was conducted 1 h after feeding to maintain water quality. For 14 days, we recorded crayfish deaths and any moults daily. Crayfish and moulted carapaces were stored in ethanol at -20°C until DNA was extracted.

2.4 Adult crayfish behaviour

Male crayfish behaviour was tested in an arena (Fig. 1) consisting of a tank (L 100 cm x H 53 cm x W 48 cm) with access to a terrestrial area (L 120 cm x H 20 cm x W 20 cm). At the start of the experiment, crayfish were divided into two groups: those destined for ‘high infection’ and those to be kept at ‘low infection’ levels. Those destined for the ‘high-infection’ group (n = 15, mean carapace length 52.2 mm, sd = 4.44) were individually exposed to a dose of 1000 zoospores mL⁻¹ in 500 mL of water for 24 h. Simultaneously, the ‘low-infection’ crayfish (n = 17, mean carapace length 53.1 mm, sd = 4.66) were sham-infected by adding the same amount of distilled water instead of spore-containing water. After the 24 h period, all crayfish were returned to their individual tanks, where they were held for 1 week before their behaviour was assessed. Individual crayfish were placed into the behavioural arena (Fig. 1) and left to acclimatise overnight. Then, at 09:00 h the next day, their behaviour was recorded using an infrared CCTV camera (Sentient Pro HDA DVR 8 Channel CCTV, Maplin, Rotherham, UK) for 24 h (09:00 – 21:00 light and 21:00 – 09:00 dark). During video analysis, the time spent engaged in each of the following four behaviours was recorded for each crayfish: actively walking in water, in a refuge, stationary out of the refuge and moving out of water.

Following this, each crayfish was moved to an aquarium (W 30 cm x L 61 cm x D 37 cm) with covered sides and allowed to settle for 30 min before their response to being gently touched on the rostrum for 10 s was tested. Crayfish typically reacted by raising their chelae (an aggressive, threatening response) and/or retreating using a characteristic ‘tail-flip’

response. If a crayfish retreated, the glass rod was immediately moved again to touch the rostrum. This test was repeated three times with 5 min intervals. Whether the crayfish reacted with a ‘tail-flip’ and/or raised its chelae to attack was recorded. These responses were recorded since behavioural changes that affect a crayfish’s ability to retreat or interact with conspecifics may have subsequent effects on competitive ability, resource acquisition, and ultimately, survival.

Following behavioural tests, crayfish were euthanized and *A. astaci* infection levels were quantified as described in section 2.2.

2.5 Statistical analysis

All statistical analyses were performed in R version 3.5.1 (R Core Team, 2018). For the juvenile crayfish experiment, Kaplan-Meier survival analyses were performed using the ‘survival’ package in R (Therneau, 2018) with separate models run for both time points (4- and 10-weeks post-hatching). Both models included spore concentration and carapace length as independent variables. Model selection was based on AIC. It was not possible to statistically assess the effect of moulting on mortality as an insufficient number of moulting events were recorded.

For the adult crayfish, the time spent moving (active), in shelter, stationary (outside of a shelter) or out of the water was quantified over 24 h for each individual. Generalised Additive Models for Location, Scale and Shape (GAMLSS) models (Stasinopoulos et al., 2008) with appropriate distributions (see Table 1) were used to determine whether ‘treatment’ (i.e. high or low infection) or carapace length (mm) influenced the proportion of time crayfish spent moving, in shelter, out of water or stationary. In the GAMLSS with beta inflated or beta zero inflated distributions, the μ parameter refers to the average amount of time spent engaging in a particular behaviour, whilst v relates to the likelihood of a behaviour not occurring

(Stasinopoulos et al., 2008). To assess the response of crayfish to a touch stimulus, threatening or tail flip escape responses were scored separately. The crayfish were tested three times, and it was noted whether they retreated by tail flipping and/or threatened by raising the chelae at least once during the three tests. These data were analysed in binomial models (i.e., threat/no threat, tail flip/no tail flip), using GAMLSS. Treatment group and carapace length were included as independent variables.

3. Results

3.1 Juvenile crayfish infection

At 4-weeks old, zoospore concentration significantly affected survival of juvenile signal crayfish ($z = 5.971$, $p < 0.001$), with almost all crayfish dying in both the 10 and 100 zoospore mL^{-1} treatments after the 14-day experimental period (Fig. 2). Around half of the crayfish died in the 1 zoospore mL^{-1} treatment, whilst 92% of control treatment crayfish survived. Carapace length also had a significant effect on the survival of these crayfish, with larger individuals surviving longer ($z = -4.387$, $p < 0.001$). In contrast, survival of crayfish exposed to the same infection doses at 10 weeks of age was not significantly affected by the zoospore treatment; all crayfish in the control and 1 zoospore mL^{-1} treatments survived, whilst 88% and 82% of those in the 10 and 100 zoospore mL^{-1} treatments survived. All juvenile crayfish that were tested (Fig. 2) were previously infected (as they were descended from infected females), although those that were exposed to zoospores exhibited significantly elevated infection levels (subset tested for *A. astaci* infection using qPCR; Kruskal-Wallis $X^2 = 9.7534$, $df = 3$, $p = 0.021$; Fig. 2).

3.2 Adult crayfish behaviour

All crayfish in the ‘high-infection’ group displayed agent levels A4-A6 (median number of PFU = 23,050; n[A4] = 3; n[A5] = 13; n[A6] = 1), whilst all crayfish from the ‘low-infection’ group remained at very low (n[A2] = 9) to low (n[A3] = 6) infection levels (median number of PFU = 43). As such, for all analyses, crayfish behaviour was compared in terms of high and low infection.

Adults exposed to *A. astaci* zoospores (high-infection: 1000 zoospores mL⁻¹) were significantly less likely to leave the water and spent on average 1.3% (range: 0 - 3.8%) of the 24 h period in the terrestrial arena compared to those in the low infection-group (sham-infected), which spent 3.5% (range: 0.3 - 9.2%) out of water (GAMLSS, ν , LRT = 5.671, $p = 0.017$). In terms of the other behaviours, there was no significant difference between crayfish from both the high and low-infection groups, which spent 31.8 (standard deviation $\pm 9.1\%$) of the time active, 47.2 $\pm 25\%$ stationary outside of a shelter and 18.6 $\pm 26\%$ in a shelter (Table 1; Appendix Table 2).

Crayfish from the high-infection group were also significantly less likely to mount a tail-flip response to tactile stimulation (GAMLSS, μ , LRT = 4.036, $p = 0.045$), where 35% of those in the high-infection group initiated a tail-flip response at least once compared to 75% of those from the low-infection group. Overall, though there was no significant difference between the two treatment groups, larger crayfish were more likely to display a threat response (GAMLSS, μ , LRT = 4.758, $p = 0.029$), spend less time in a shelter (GAMLSS, ν , LRT = 5.514, $p = 0.019$) and more time stationary outside of a shelter (GAMLSS, ν , LRT = 5.730, $p = 0.017$) compared to smaller crayfish.

4. Discussion

Here, we show that *A. astaci* can cause almost total mortality in juvenile signal crayfish at ecologically relevant zoospore densities (Strand et al., 2012, 2014), though larger, older individuals were less affected. Additionally, we show that a high *A. astaci* burden affects the behaviour of adult crayfish, making them almost half as likely to spend time on land and to escape from tactile stimulation compared to less infected individuals. The low infection levels of our control crayfish did not differ from those frequently observed in *P. leniusculus* populations across Europe (Kozubíková et al., 2011; Filipová et al., 2013; Tilmans et al., 2014) and in Japan (Mrugała et al., 2017); although slightly higher infection levels (A2-A5) were reported in the UK (James et al., 2017). Thus, the high infection group in our study represents the outbreak of a highly virulent strain. Whilst signal crayfish are a highly successful invasive species in Europe that continue to spread (Peay et al., 2010; Holdich et al., 2014; James et al., 2014), the negative impacts of crayfish plague reported here, especially in terms of juvenile mortality, could have consequences for commercially harvested stocks by reducing recruitment and possibly resulting in population crashes. This also supports previous studies which have shown that commercially harvested signal crayfish populations can decline when *A. astaci* is present (Jussila et al., 2016). Furthermore, these results add to growing evidence that *A. astaci* could play a more significant role in regulating invasive signal crayfish population dynamics than previously considered, which could play a role in determining invasion success (Jussila et al., 2015).

In North American crayfish species, *A. astaci* can grow within the carapace, though constant host melanisation of new hyphae prevents spore penetration to soft tissues (Unestam & Weiss, 1970; Nyhlén & Unestam, 1975; Cerenius et al., 2003). In the present study, juvenile signal crayfish suffered extensive dose-dependent mortality when exposed to *A. astaci* zoospores around 4-weeks post-hatching. Slightly older (and therefore larger) crayfish, however, avoided this cost. Many juvenile crayfish studied here probably became infected

rapidly after hatching, having acquired an infection from their mothers via horizontal transmission. Older and larger crayfish possibly have a better-developed immune response, capable of efficiently melanising hyphae. It has been suggested that the immune response of juvenile crayfish to *A. astaci* infection is generally reduced compared to adults (Mrugała et al., 2016), which seems to be the case in the present study. In other invertebrates too, younger individuals exhibit lower immune responses, for instance, snails showing greater susceptibility to schistosome parasites (Dikkeboom et al., 1985). It has also been hypothesised, however, that juvenile crayfish could be less affected due to their higher moulting frequency compared to adults (Reynolds, 2002), allowing them to shed the growing hyphae and lower their *A. astaci* burden. Further research comparing the immunological capacity of juvenile and adult crayfish is required to confirm this. By the 10-week time-point, particularly susceptible individuals may have already succumbed to infection and therefore those used in the current experiment would have been more resistant to the pathogen. This appears unlikely though, since high levels of mortality were not observed in the communal holding tanks. Ecologically, the finding that relatively young crayfish hatchlings are highly susceptible to high doses of zoospores could have significant implications for signal crayfish recruitment and survival, especially in lentic environments, where zoospores are less likely to be washed away from the maternal crayfish.

Adult crayfish suffering from higher *A. astaci* infection levels during the present study exhibited a reduced tendency to leave the water. Although crayfish spend little time out of water in general, this finding suggests that populations of invasive signal crayfish with high burdens of *A. astaci* could be less likely to disperse overland to reach new aquatic habitats, a behavioural trait that can contribute to the spread of invasive crayfish (Grey and Jackson, 2012; Holdich et al., 2014; Puky, 2014; Ramalho and Anastácio, 2014). Other invertebrates are less active when infected by parasites, potentially to avoid the associated fitness costs of dispersal. Flat back mud crabs (*Eurypanopeus depressus*) infected with rhizocephalans, for example,

spend more time in shelter and are less active than uninfected crabs (Belgrad and Griffen, 2015), whilst sponge-dwelling snapping shrimp (*Synalpheus elizabethae*) infected by bopyrid isopods show 50% lower activity levels compared to uninfected individuals (McGrew and Hultgren, 2011). In other invertebrates, many studies have shown that parasites can influence dispersal, though these studies focus on direct host manipulation, which does not seem to be the case here as there is no evidence of *A. astaci* actively manipulating the host. In terms of native European crayfish management, a lower tendency of infected individuals to disperse overland might be beneficial, by reducing the transmission of *A. astaci* to new waterbodies.

Highly infected crayfish were also less likely to respond to tactile stimulation by retreating in a characteristic ‘tail-flip’ manner. This reduced ability to escape could lead to increased predation of highly infected crayfish. *A. astaci* zoospores largely penetrate soft abdominal tissue (Vrålstad et al., 2009), and it is possible that the reduced escape response is directly due to the general pathological effects of the parasite (Unestam & Weiss, 1970). Other parasites, such as *Thelohania contejeani*, also penetrate crayfish tissues, parasitising the muscles and reducing the ability of crayfish to predate and feed (Haddaway et al., 2012). It is also possible that highly infected crayfish exhibit a reduced tendency to move on land to reduce the risk of predation. In the same way, crustaceans become less active and tend to stay in a refuge when moulting, during which they are vulnerable to predators and largely unable to escape (Thomas, 1965; Cromarty et al., 2000).

The crayfish used in the current study were from a population previously considered to be below the detection limit (n = 30 tested by James et al. (2017) exhibited A0-A1 levels). However, given that infection levels A2-A3 were found both among crayfish tested before the experiment began, as well as among those in the group not exposed to zoospores, it is evident that this population has either become infected since 2014, that a previously very low prevalence of *A. astaci* has since increased, or that crayfish present with A2-A3 infection levels

in 2014 were just not trapped by James et al. (2017). Signal crayfish in Europe are generally associated with the Group B strain of *A. astaci* (see Huang, Cerenius and Söderhäll, 1994; Grandjean et al., 2014), which has also been found infecting another Welsh population, approximately 45 miles away from the population studied here (James et al., 2017). Although not confirmed, the crayfish used in the present study were most likely initially infected with a Group B strain and subsequently exposed to another strain from the same group. It is also possible that the tested crayfish were locally adapted to their original *A. astaci* infection (Gruber et al., 2014; Jussila et al., 2015), and the observed behavioural effects resulted from the exposure to the new *A. astaci* strain. As observed by Jussila et al. (2013), even assumed identical *A. astaci* strains may differently affect their crayfish hosts; therefore, the experimental crayfish in the present study likely dealt with multiple infections of closely related *A. astaci* strains. Further research is required, to explicitly compare the behaviour and survival of infected and uninfected signal crayfish, as well as investigate the effects of different *A. astaci* strains (both in single and multiple infections) on the behaviour and survival of infected crayfish.

In summary, we have shown that high levels of *A. astaci* cause severe mortality in young juveniles and affect the behaviour of adult signal crayfish. Mounting evidence suggests that signal crayfish may succumb to *A. astaci* more often than previously considered, which could be having an impact on commercially harvested populations (Aydin et al., 2014; Edsman et al., 2015). The crayfish exposed to zoospores in the present study displayed relatively high plague agent levels of A4-A6 (A7 being the highest level of infection; Vrålstad et al., 2009). A longer period of infection, or higher infection dose, may induce further behavioural responses beyond those reported here, and in some cases even cause mortality as observed by Aydin et al. (2014), where signal crayfish were exposed to 10,000 zoospores mL⁻¹. Female crayfish suffering from ESS carry far fewer fertilised eggs than uninfected females (Edsman et al.,

2015) which, coupled with the high juvenile mortality documented in the present study, could drastically reduce juvenile recruitment and result in population crashes. Similarly, crayfish plague could also have implications for the further spread of signal crayfish by affecting population dynamics, though this species has already successfully colonised large parts of Europe (Holdich et al., 2014) and so the ecological impact may be negligible. Anecdotally, it was assumed that most North American crayfish are infected with *A. astaci*, though molecular methods have demonstrated that it is less prevalent than once thought. In France, for example, just over half of the signal crayfish populations tested were found to be positive for crayfish plague (Filipová et al., 2013), and in the UK the prevalence was 56.5% (James et al., 2017). It is possible, therefore, that the population dynamics of uninfected invasive populations may be affected when infected individuals are translocated and introduced.

Financial Support

This project was funded by Coleg Cymraeg Cenedlaethol (JRT) and the Welsh Government and Higher Education Funding Council for Wales through the Sêr Cymru National Research Network for Low Carbon, Energy and the Environment (NRN-LCEE) AquaWales project.

References

- Alderman, DJ** (1982) *In vitro* testing of fisheries chemotherapeutants. *Journal of Fish Diseases* **5**, 113-123. <https://doi.org/10.1111/j.1365-2761.1982.tb00464.x>
- Aydin, H, Kokko, H, Makkonen, J, Kortet, R, Kukkonen, H and Jussila, J** (2014) The signal crayfish is vulnerable to both the As and the PsI-isolates of the crayfish plague. *Knowledge and Management of Aquatic Ecosystems* **03**. <https://doi.org/10.1051/kmae/2014004>
- Bakker, TCM, Mazzi, D, Zala, S** (1997) Parasite-induced changes in behavior and color make

- 389 *Gammarus pulex* more prone to fish predation. *Ecology* **78**, 1098-1104.
 390 <https://doi.org/10.2307/2265861>
- 391 **Barber, I, Mora, AB, Payne, EM, Weinersmith, KL, Sih, A** (2017) Parasitism, personality
 392 and cognition in fish. *Behavioural Processes* **141**, 205-219.
 393 <https://doi.org/10.1016/j.beproc.2016.11.012>
- 394 **Becking, T, Mrugala, A, Delaunay, C, Svoboda, J, Raimond, M, Viljamaa-Dirks, S,**
 395 **Petrusek, A, Grandjean, F and Braquart-Varnier, C** (2015) Effect of experimental
 396 exposure to differently virulent *Aphanomyces astaci* strains on the immune response of the
 397 noble crayfish *Astacus astacus*. *Journal of Invertebrate Pathology* **132**, 115-124.
 398 <https://doi.org/10.1016/j.jip.2015.08.007>
- 399 **Belgrad, BA, Griffen, BD** (2015) Rhizocephalan infection modifies host food consumption
 400 by reducing host activity levels. *Journal of Experimental Marine Biology and Ecology* **466**,
 401 70-75. <http://dx.doi.org/10.1016/j.jembe.2015.02.011>
- 402 **Cable, J, Barber, I, Boag, B, Ellison, AR, Morgan, ER, Murray, K, Pascoe, EL, Sait, SM,**
 403 **Wilson, AJ and Booth, M** (2017) Global change, parasite transmission and disease control:
 404 lessons from ecology. *Philosophical Transactions of the Royal Society B Biological Sciences*
 405 **372**, 20160088. <https://doi.org/10.1098/rstb.2016.0088>
- 406 **Cerenius, L, Bangyeekhun, E, Keyser, P, Söderhäll, I and Söderhäll, K** (2003) Host
 407 prophenoloxidase expression in freshwater crayfish is linked to increased resistance to crayfish
 408 plague fungus, *Aphanomyces astaci*. *Cellular Microbiology* **5**, 353-357.
- 409 **Cerenius, L and Söderhäll, K** (1984) Repeated zoospore emergence from isolated spore cysts
 410 of *Aphanomyces astaci*. *Experimental Mycology* **8**, 370-377. [https://doi.org/10.1016/0147-](https://doi.org/10.1016/0147-5975(84)90061-6)
 411 [5975\(84\)90061-6](https://doi.org/10.1016/0147-5975(84)90061-6)
- 412 **Cerenius, L, Söderhäll, K, Persson, M and Ajaxon, R** (1987) The crayfish plague fungus
 413 *Aphanomyces astaci* - diagnosis, isolation, and pathobiology. *Freshwater Crayfish* **7**, 131-143.

- 414 **Cromarty, SI, Mello, J and Kass-Simon, G** (2000) Molt-related and size-dependent
 415 differences in the escape response and post-threat behavior of the American lobster, *Homarus*
 416 *americanus*. *Biological Bulletin* **199**, 265-277. <https://doi.org/10.2307/1543183>
- 417 **Dikkeboom, R, Knaap, WPWVD, Meuleman, EA and Sminia, T** (1985) A comparative
 418 study on the internal defence system of juvenile and adult *Lymnaea stagnalis*. *Immunology* **55**,
 419 547-553.
- 420 **Edsman, L, Nyström, P, Sandström, A, Stenberg, M, Kokko, H, Tiitinen, V, Makkonen,**
 421 **J and Jussila, J** (2015) Eroded swimmeret syndrome in female crayfish *Pacifastacus*
 422 *leniusculus* associated with *Aphanomyces astaci* and *Fusarium* spp. infections. *Diseases of*
 423 *Aquatic Organisms* **112**, 219-228. <https://doi.org/10.3354/dao02811>
- 424 **Ercoli, F, Ruukonen, TJ, Koistinen, S, Jones, RI and Hämäläinen, H** (2015) The introduced
 425 signal crayfish and native noble crayfish have different effects on sublittoral macroinvertebrate
 426 assemblages in boreal lakes. *Freshwater Biology* **60**, 1688-1698.
 427 <https://doi.org/10.1111/fwb.12601>
- 428 **Filipová, L, Petrusek, A, Matasová, K, Delaunay, C and Grandjean, F** (2013) Prevalence
 429 of the crayfish plague pathogen *Aphanomyces astaci* in populations of the signal crayfish
 430 *Pacifastacus leniusculus* in France: evaluating the threat to native crayfish. *PLoS One* **8**,
 431 e70157. <https://doi.org/10.1371/journal.pone.0070157>
- 432 **Grandjean, F, Vrålstad, T, Diéguez-Urbeondo, J, Jelić, M, Mangombi, J, Delaunay, C,**
 433 **Filipová, L, Rezinciuc, S, Kozubíková-Balcarová, E, Guyonnet, D, Viljamaa-Dirks, S and**
 434 **Petrusek, A** (2014) Microsatellite markers for direct genotyping of the crayfish plague
 435 pathogen *Aphanomyces astaci* (Oomycetes) from infected host tissues. *Veterinary*
 436 *Microbiology* **170**, 317-324. <https://doi.org/10.1016/j.vetmic.2014.02.020>
- 437 **Grey, J and Jackson, MC** (2012) “Leaves and eats shoots”: direct terrestrial feeding can
 438 supplement invasive red swamp crayfish in times of need. *PLoS One* **7**, e42575.

439 <https://doi.org/10.1371/journal.pone.0042575>

440 **Gruber, C, Kortet, R, Vainikka, A, Hyvärinen, P, Rantala, M.J, Pikkarainen, A, Jussila,**
 441 **J, Makkonen, J, Kokko, H and Hirvonen, H** (2014) Variation in resistance to the invasive
 442 crayfish plague and immune defence in the native noble crayfish. *Annales Zoologici Fennici*
 443 **51**, 371-389. <https://doi.org/10.5735/086.051.0403>

444 **Haddaway, NR, Wilcox, R.H, Heptonstall, REA, Griffiths, HM, Mortimer, RJG,**
 445 **Christmas, M and Dunn, AM** (2012) Predatory functional response and prey choice identify
 446 predation differences between native/invasive and parasitised/unparasitised crayfish. *PLoS*
 447 *One* **7**, e32229. <https://doi.org/10.1371/journal.pone.0032229>

448 **Hatcher, MJ, Dick, JTA and Dunn, AM** (2014) Parasites that change predator or prey
 449 behaviour can have keystone effects on community composition. *Biological Letters* **10**.
 450 <https://doi.org/10.1098/rsbl.2013.0879>

451 **Holdich, DM, James, J, Jackson, C and Peay, S** (2014) The North American signal crayfish,
 452 with particular reference to its success as an invasive species in Great Britain. *Ethology*
 453 *Ecology & Evolution* **26**, 232-262. <https://doi.org/10.1080/03949370.2014.903380>

454 **Huang, TS, Cerenius, L and Söderhäll, K** (1994) Analysis of genetic diversity in the crayfish
 455 plague fungus, *Aphanomyces astaci*, by random amplification of polymorphic DNA.
 456 *Aquaculture* **126**, 1-9. [https://doi.org/10.1016/0044-8486\(94\)90243-7](https://doi.org/10.1016/0044-8486(94)90243-7)

457 **Hudson, PJ, Dobson, AP and Lafferty, KD** (2006) Is a healthy ecosystem one that is rich in
 458 parasites? *Trends Ecology & Evolution* **21**, 381-385. <https://doi.org/10.1016/j.tree.2006.04.007>

459 **James, J, Cable, J and Slater, F** (2014) A.L.I.E.N. databases: Addressing the lack in
 460 establishment of non-natives databases. *Crustaceana* **87**, 1192-1199.
 461 <https://doi.org/10.1163/15685403-00003329>

462 **James, J, Nutbeam-Tuffs, S, Cable, J, Mrugała, A, Viñuela-Rodriguez, N, Petrusek, A**
 463 **and Oidtmann, B** (2017) The prevalence of *Aphanomyces astaci* in invasive signal crayfish

- 464 from the UK and implications for native crayfish conservation. *Parasitology* **144**, 411-418.
 465 <https://doi.org/10.1017/S0031182016002419>
- 466 **Jussila, J, Kokko, H, Kortet, R, Makkonen, J** (2013) *Aphanomyces astaci* Ps1-genotype
 467 isolates from different Finnish signal crayfish stocks show variation in their virulence but still
 468 kill fast. *Knowledge and Management of Aquatic Ecosystems* **411**, 10.
 469 <https://doi.org/10.1051/kmae/2013077>
- 470 **Jussila, J, Vrezec, A, Makkonen, J and Kortet, R** (2015) Invasive crayfish and their invasive
 471 diseases in Europe with the focus on the virulence evolution of the crayfish plague. In Canning-
 472 Clode J. (eds) *Biological Invasions in Changing Ecosystems Vectors, Ecological Impacts,*
 473 *Management and Predictions.* DeGruyter Open Ltd, Warsaw/Berlin, pp. 183-204.
 474 <https://doi.org/10.1515/9783110438666-013>
- 475 **Jussila, J, Tiitinen, V, Edsman, L, Kokko, H, Fotedar, R** (2016) Signal crayfish in Lake
 476 Saimaa could be maladapted to the local conditions due to *Aphanomyces astaci* infection: a
 477 seven-year study. *Freshwater Crayfish* **22**, 53-60. <https://doi.org/10.5869/fc.2016.v22-1.53>
- 478 **Kozubíková-Balcarová, E, Koukol, O, Martín, M.P, Svoboda, J, Petrusek, A and**
 479 **Diéguez-Urbeondo, J** (2013) The diversity of oomycetes on crayfish: morphological vs.
 480 molecular identification of cultures obtained while isolating the crayfish plague pathogen.
 481 *Fungal Biology* **117**, 682-691.
- 482 **Kozubíková-Balcarová, E, Beran, L, Ďuriš, Z, Fischer, D, Horká, Svoboda, J, Petrusek,**
 483 **A** (2015) Status and recovery of indigenous crayfish populations after recent crayfish plague
 484 outbreaks in the Czech Republic. *Ethology Ecology & Evolution* **26**, 299-319.
 485 <https://doi.org/10.1080/03949370.2014.897652>
- 486 **Longshaw, M, Bateman, KS, Stebbing, P, Stentiford, GD and Hockley, FA** (2012) Disease
 487 risks associated with the importation and release of non-native crayfish species into mainland
 488 Britain. *Aquatic Biology* **16**, 1-15. <https://doi.org/10.3354/ab00417>

- 489 **Macnab, V and Barber, I** (2012) Some (worms) like it hot: fish parasites grow faster in
 490 warmer water, and alter host thermal preferences. *Global Change Biology* **18**, 1540-1548.
 491 <https://doi.org/10.1111/j.1365-2486.2011.02595.x>
- 492 **Makkonen, J, Kokko, H, Henttonen, P and Jussila, J** (2010) Crayfish plague (*Aphanomyces*
 493 *astaci*) can be vertically transferred during artificial incubation of crayfish eggs: preliminary
 494 results. *Freshwater Crayfish* **17**, 151-153.
- 495 **Makkonen, J, Strand, DA, Kokko, H, Vrålstad, T and Jussila, J** (2013) Timing and
 496 quantifying *Aphanomyces astaci* sporulation from the noble crayfish suffering from the
 497 crayfish plague. *Veterinary Microbiology* **162**, 750-755.
 498 <https://doi.org/10.1016/j.vetmic.2012.09.027>
- 499 **McGrew, M, Hultgren, KM** (2011) Bopyrid parasite infestation affects activity levels and
 500 morphology of the eusocial snapping shrimp *Synalpheus elizabethae*. *Marine Ecology*
 501 *Progress Series* **43**, 195-204. <https://doi.org/10.3354/meps09123>
- 502 **Mrugała, A, Veselý, L, Petrusek, A, Viljamaa-Dirks, S and Kouba, A** (2016) May *Cherax*
 503 *destructor* contribute to *Aphanomyces astaci* spread in Central Europe? *Aquatic Invasions* **11**,
 504 459-468. <https://doi.org/10.3391/ai.2016.11.4.10>
- 505 **Nyhlén, L and Unestam, T** (1975) Ultrastructure of the penetration of the crayfish integument
 506 by the fungal parasite, *Aphanomyces astaci*, Oomycetes. *Journal of Invertebrate Pathology* **26**,
 507 353-366.
- 508 **Peay, S, Holdich, DM and Brickland, J** (2010) Risk assessments of non-indigenous crayfish
 509 in Great Britain. *Freshwater Crayfish* **17**, 109-122. [https://doi.org/10.1007/s00531-014-1002-](https://doi.org/10.1007/s00531-014-1002-0)
 510 0
- 511 **Preston, DL, Mischler, JA, Townsend, AR and Johnson, PTJ** (2016) Disease ecology meets
 512 ecosystem science. *Ecosystems* **19**, 737-748. <https://doi.org/10.1007/s10021-016-9965-2>
- 513 **Puky, M** (2014) Invasive crayfish on land: *Orconectes limosus* (Rafinesque, 1817) (Decapoda:

- 514 Cambaridae) crossed a terrestrial barrier to move from a side arm into the Danube River at
 515 Szeremle, Hungary. *Acta Zoologica Bulgarica* **66**, 143-146.
- 516 **Ramvalho, RO and Anastácio, PM** (2014) Factors inducing overland movement of invasive
 517 crayfish (*Procambarus clarkii*) in a ricefield habitat. *Hydrobiologia* **746**, 135-146.
 518 <https://doi.org/10.1007/s10750-014-2052-9>
- 519 **Söderhäll, K, Svensson, E and Unestam, T** (1978) Chitinase and protease activities in
 520 germinating zoospore cysts of a parasitic fungus, *Aphanomyces astaci*, oomycetes.
 521 *Mycopathologia* **64**, 9-11.
- 522 **Stasinopoulos, M, Rigby, RA and Akantziliotou, C** (2008) Instructions on how to use the
 523 GAMLSS package in R, second edition. Available at: [http://www.gamlss.com/wp-](http://www.gamlss.com/wp-content/uploads/2013/01/gamlss-manual.pdf)
 524 [content/uploads/2013/01/gamlss-manual.pdf](http://www.gamlss.com/wp-content/uploads/2013/01/gamlss-manual.pdf)
- 525 **Strand, DA, Jussila, J, Johnsen, SI, Viljamaa-Dirks, S, Edsman, L, Wiik-Nielsen, J,**
 526 **Viljugrein, H, Engdahl, F and Vrålstad, T** (2014) Detection of crayfish plague spores in
 527 large freshwater systems. *Journal of Applied Ecology* **51**, 544-553.
 528 <https://doi.org/10.1111/1365-2664.12218>
- 529 **Strand, DA, Jussila, J, Viljamaa-Dirks, S, Kokko, H, Makkonen, J, Holst-Jensen, A,**
 530 **Viljugrein, H and Vrålstad, T** (2012) Monitoring the spore dynamics of *Aphanomyces astaci*
 531 in the ambient water of latent carrier crayfish. *Veterinary Microbiology* **160**, 99-107.
 532 <https://doi.org/10.1016/j.vetmic.2012.05.008>
- 533 **Svoboda, J, Kozubíková-Balcarová, E, Kouba, A, Buřič, M, Kozák, P, Diéguez-**
 534 **Uribeondo, J, Petrusek, A** (2013) Temporal dynamics of spore release of the crayfish plague
 535 pathogen from its natural host, American spiny-cheek crayfish (*Orconectes limosus*), evaluated
 536 by transmission experiments. *Parasitology* **140**, 792-801.
 537 <https://doi.org/10.1017/S0031182012002223>
- 538 **Svoboda, J, Mrugała, A, Kozubíková-Balcarová, E and Petrusek, A** (2017) Hosts and

transmission of the crayfish plague pathogen *Aphanomyces astaci*: a review. *Journal of Fish Diseases* **40**, 127-140. <https://doi.org/10.1111/jfd.12472>

Svoboda, J, Strand, DA, Vrålstad, T, Grandjean, F, Edsman, L, Kozák, P, Kouba, A, Fristad, RF, Koca, SB and Petrusek, A (2014) The crayfish plague pathogen can infect freshwater-inhabiting crabs. *Freshwater Biology* **59**, 918-929. <https://doi.org/10.1111/fw.b.12315>

Terry, M and Therneau, M (2018) Package “survival”. <https://CRAN.R-project.org/package=survival>

Thomas, LR (1965) Moulting behaviour of the western Australian crayfish *Panulirus cygnus* George (Decapoda, Reptantia). *Crustaceana* **11**, 111-113.

Tompkins, DM, Dunn, AM, Smith, MJ and Telfer, S (2011) Wildlife diseases: from individuals to ecosystems. *Journal of Animal Ecology* **80**, 19-38. <https://doi.org/10.1111/j.1365-2656.2010.01742.x>

Unestam, T and Weiss, DW (1970) The host-parasite relationship between freshwater crayfish and the crayfish disease fungus *Aphanomyces astaci*: responses to infection by a susceptible and a resistant species. *Journal of General Microbiology* **60**, 77-90. <https://doi.org/10.1099/00221287-60-1-77>

Vilcinskas, A (2015) Pathogens as biological weapons of invasive species. *PLoS Pathogens* **11**, 1-5. <https://doi.org/10.1371/journal.ppat.1004714>

Vrålstad, T, Knutsen, AK, Tengs, T and Holst-Jensen, A (2009) A quantitative TaqMan® MGB real-time polymerase chain reaction based assay for detection of the causative agent of crayfish plague *Aphanomyces astaci*. *Veterinary Microbiology* **137**, 146-155. <https://doi.org/10.1016/j.vetmic.2008.12.022>

Fig. 1 – Experimental arena used to assess crayfish behaviour. The tank was filled with water up to 3 cm below the level of the terrestrial area. The base of the arena, ramp (incline 30°) and bridge were coated in 1-2 cm of pea gravel.

Fig. 2 – Survival (a, b) and infection levels (c, d) of juvenile signal crayfish infected with *A. astaci* for two weeks. Infection treatments were sham-infection, 1, 10 and 100 zoospores mL⁻¹; a & c) 4 weeks after hatching; b & d) 10 weeks after hatching. Note in b) sham-infection treatment is identical to infection treatment 1 (grey / dashed grey). A subset of juvenile crayfish from each treatment was tested using qPCR, c) sham-infection, 0 zoospores mL⁻¹ [n = 5], 1 [n = 6], 10 [n = 5], 100 [n = 4]; d) 0 [n = 3], 1 [n = 3], 10 [n = 5], 100 [n = 6]. See Appendix Table 1 for absolute values.

Table 1 – Results of GAMLSS (**Generalised Additive Models for Location, Scale and Shape**) statistical analyses and mean proportion of time crayfish spent engaged in different behaviours over 24 h. Significant results ($p < 0.05$) are highlighted in bold. BE = beta, BEZI = beta zero-inflated, BEINF = beta inflated, BI = binomial, SD = standard deviation. * denotes proportion of crayfish, not the mean.

Response variable	Infection treatment group	Mean (%)	Range (%)	GAMLSS Parameter	Family	Variable	DF	LRT	P-value
Proportion of time active	Low	31	21-49	μ	BE	Treatment	1, 28	0.016	0.898
	High	32	18-48			CL		0.024	0.876
Proportion of time out of water	Low	4	0-9	μ	BEZI	Treatment	1, 25	2.095	0.147
	High	1	0-4			CL		0.075	0.784
				ν		Treatment		5.671	0.017
						CL		0.125	0.724
Proportion of time in shelter	Low	21	0-55	μ	BEINF	Treatment	1, 24	0.046	0.830
	High	17	0-75			CL		2.478	0.116
				ν		Treatment		0.043	0.835
						CL		5.514	0.019
Proportion of time stationary	Low	44	11-76	μ	BEINF	Treatment	1, 26	0.383	0.536
	High	50	0-76			CL		5.730	0.017
Proportion of crayfish that tail flipped	Low	75%	na	μ	BI	Treatment	1, 26	4.036	0.045

		High	35%	na			CL	1.607	0.205
Proportion of crayfish that exhibited threat behaviour	Low		67%	na	μ	BI	Treatment	0.177	0.674
	High		65%	na			CL	4.758	0.029

582

583