

# Berry chemoprevention: Do berries decrease the window of opportunity for tumorigenesis

Dr. Stephanie May BSc(Hons), PhD | Connor Parry BSc  | Dr. Lee Parry BSc(Hons), PhD

European Cancer Stem Cell Research Institute,  
School of Biosciences, Cardiff University,  
Cardiff, UK

## Correspondence

Lee Parry, European Cancer Stem Cell  
Research Institute, School of Biosciences,  
Cardiff University, Cardiff CF24 4HQ, UK.  
Email: [ParryL3@cardiff.ac.uk](mailto:ParryL3@cardiff.ac.uk)

## Funding information

Tenovus Cancer Care, Grant/Award Num-  
bers: Phd2013/L25, LP&SM; World Can-  
cer Research Fund, Grant/Award Number:  
(2013/1006; Cancer Research UK Programme,  
Grant/Award Number: C1295/ A15937-LP; Mr  
Lyndon and Mrs Shirley-Ann Wood from the  
Moorhouse Group Ltd, Grant/Award Number:  
LP&SM

## Abstract

In addition to smoking and UV exposure, lifestyle factors, such as diet, nutrition, and physical activity, have been shown to play a significant role for many cancers. It is estimated that up to 50% of some cancer types are preventable; many through lifestyle and dietary changes with the presence or absence of certain dietary components strongly associated with an increased or decreased risk. Here we summarize the work that has been performed with polyphenols, with a focus on those derived from black raspberries. These have been extensively studied for the prevention and treatment of a variety of conditions and diseases. Here we focus on their use for the prevention or treatment of specific cancer types and the impact they have on biological systems. The aim is to highlight the need to improve our understanding of how the environment impacts upon the normal biological processes that affect health and disease, thereby, enabling us to implement smarter prevention and treatment measures.

## KEYWORDS

anthocyanin, black raspberry, cancer, chemoprevention, epigenetics, inflammation, microbiome

## 1 | INTRODUCTION

The global increase in the incidence and mortality of all cancer types is leading to an escalating burden on global economics and healthcare (Bray et al., 2018). The reasons behind this are somewhat complex, but reflect a rapid population growth, an increased ageing population, and alterations to the prevalence and distribution of risk factors attributed to the disease (Bray et al., 2018). Lifestyle factors, such as diet, obesity, and socioeconomic status, have a profound impact on cancer incidence. Combined, these highlight an increasing demand for strategies for cancer prevention, as there is substantive evidence that diet, nutrition, and physical activity play fundamental roles in cancer and disease prevention (World Cancer Research Fund/American Institute for Cancer Research, 2018). Over the past two decades, a substantial


amount of research has focused on the chemopreventative and anti-tumorigenic properties of dietary polyphenols, such as those found in many fruits and vegetables. In this review, we will present the current position on lifestyle factors and cancer, the utilization of polyphenols by the body, summarize the data on polyphenols and specific tumor types, and how they influence the epigenome, microbiome, and immunity.

### 1.1 | Established causes of cancer

There are over 100 different types of cancer, which is complicated further by the several subclassifications within each cancer (World Cancer Research Fund/American Institute for Cancer Research,

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Food Frontiers* published by John Wiley & Sons Australia, Ltd and Nanchang University, Northwest University, Jiangsu University, Zhejiang University, Fujian Agriculture and Forestry University



**FIGURE 1** Endogenous and exogenous risk factors of cancer development. Cancer arises from abnormal and uncontrolled cell division. Cancer progression is the result of a complex interaction between innate host factors (endogenous risk factors), such as inherited mutations and epigenetic signatures, disturbances in microbial diversity due to chronic inflammatory diseases and reduce immune function, with exogenous factors that can increase mutational risk over time such as UV exposure, smoking, physical inactivity, obesity, inadequate dietary intake, and alcohol consumption

2018). Although the exact mechanisms and behaviors behind each cancer is not currently understood, the complex relationship between cancer, host, nutrition, and physical activity are being explored. At a molecular level, cancer arises from an accumulation of DNA damage and as a result of abnormal and uncontrollable cell division. There are several factors identified under the classic “hallmarks” that can compromise normal cell division (Hanahan & Weinberg Robert, 2011; Wu, Zhu, Thompson, & Hannun, 2018), these can be broadly divided into two groups; “endogenous” and “exogenous” factors (World Cancer Research Fund/American Institute for Cancer Research, 2018, Wu et al., 2018; Figure 1).

## 1.2 | “Endogenous” factors

Endogenous factors arise from processes within the individual. Less than 10% of cancers are attributed to inherited genetic mutations often affecting a tumor suppressor gene, for example, patients with familial adenomatous polyposis (FAP), inherit a mutated *Adenomatosis polyposis coli* gene and are predisposed to colorectal cancer (CRC) (Nishisho et al., 1991). Other endogenous factors include imbalances in hormonal and metabolic pathways, chronic inflammation associated with DNA damage through genetic and epigenetic modifications, and oxidative stress (Wu et al., 2018).

### 1.3 | “Exogenous” factors

Exogenous factors are commonly termed as lifestyle factors. These include carcinogens from smoking, excessive alcohol consumption, infectious agents: such as liver cancer from hepatitis B and C infections, and *Helicobacter pylori* a cause of stomach cancer (World Cancer Research Fund/American Institute for Cancer Research, 2018); ionizing radiation from ultraviolet (UV) light and influences from nutrition, body fatness, and physical activity (World Cancer Research Fund/American Institute for Cancer Research, 2018). The relationship between diet or physical activity and cancer is specific for certain cancers, such as the relationship between CRC and diets consisting of red and processed meats (Mosley et al., 2020). However, it is often the case that several exogenous exposures relating to diet and physical activity are attributed to numerous cancer types (World Cancer Research Fund/American Institute for Cancer Research, 2018).

In the UK around 4 in 10 cancer cases a year are linked to exogenous factors. The largest single cause of cancer is smoking, the predominant cause of lung cancer. Body fatness and obesity are estimated to contribute to 6.3% of cancer cases. UV radiation and occupational risks make up 3.8% of UK cancer cases. Dietary factors have significant roles in the etiology of many cancers, particularly CRC, excessive alcohol consumption, insufficient fiber intake and infections are attributable to 2–4% of cases (Brown et al., 2018). To fully understand the chemopreventative mechanisms underpinning nutritional factors in cancer, both large-scale epidemiological and preclinical studies are necessary. One area of intense research has focused on the protective role of bioactive phytochemicals, in particular polyphenols.

### 1.4 | Dietary polyphenols

Polyphenols are naturally occurring compounds enriched in fruits, vegetables, and cereals and are a major source of antioxidants in the human diet. They are secondary metabolites in plants that aid in protecting against UV radiation and aggression by pathogens (Pandey & Ibrahim, 2009). For many years, plant-derived nutritional polyphenols have been investigated for their antitumorigenic (Middleton, Kandaswami, & Theoharides, 2000) properties, protection against cardiovascular disease, diabetes, neurodegenerative disease (Soobrattee, Bahorun, & Aruoma, 2006), anti-inflammatory (Nijveldt et al., 2001), antithrombotic (Manach, Scalbert, Morand, Remesy, & Jimenez, 2004), and antiviral (Lee et al., 2016) properties.

These antioxidants prevent damage to cells by scavenging and neutralizing free radicals and reactive oxygen species (ROS) thereby reducing oxidative stress. Although ROS are naturally produced within the body from normal oxidative metabolism (Diplock et al., 1998), accumulation of these species can be detrimental. High concentrations of free radicals and ROS can result in extensive DNA, membrane lipid, and protein damage which can potentiate the development of cancer (Diplock et al., 1998). Naturally produced endogenous antioxidants remove free radicals and prevent cellular damage in the body. However, the body heavily relies on exogenous sources of antioxidants, mainly

from the diet (Valko et al., 2007). The most abundant antioxidants in our food are the polyphenols.

Based upon their chemical structure, nutritional polyphenols can be divided into several groups: the flavonoids, phenolic acids, stilbenes, and lignans (Bravo, 1998; Manach et al., 2004). The flavonoids can be further divided into six subgroups, flavonols, flavones, isoflavones, flavanones, anthocyanidins, and flavanols (catechins and proanthocyanidins).

### 1.5 | Anthocyanins

The most abundant flavonoid found in fruit and vegetables are the anthocyanins. Anthocyanins are responsible for the variety of colors, most notably red, blue, purple, and black, in the fruits, flowers, leaves, stems, and roots of plants (Wang & Stoner, 2008). Their bright colors attract animals and insects, aiding pollination and seed dispersal of plants. Additionally, their antioxidant properties defend against extreme UV exposure, pathogens, and oxidative stress (Kong, Chia, Goh, Chia, & Brouillard, 2003). In humans, their antioxidant, anticarcinogenic, and anti-inflammatory properties (Seeram, 2008) have been widely investigated as a potential treatment for many cancers.

In the USA, the estimated daily intake of anthocyanins is around 180–215 mg, approximately ninefold more than other dietary flavonoids which are consumed at about 23 mg/day (Wang & Stoner, 2008). These natural anthocyanins are well represented in our diet as they occur in red, blue, and purple produce such as carrots, onions, aubergines, and grapes, but they are most abundant in berry fruits (Table 1; Manach et al., 2004; Mazza, 2007), having as high as 1,480 mg of anthocyanin/100 g fresh weight berry (Wu, Gu, Prior, & McKay, 2004). The concentration of anthocyanins in fruits is dependent on the cultivator, growing conditions (e.g., amount and quality of sunlight, soil composition, salinity, and temperature) and season (Kalt, 2005). There have been over 400 different anthocyanins identified, but the six most common anthocyanidins (the sugar-free counterparts of anthocyanins) are pelargonidin, cyanidin, delphinidin, peonidin, petunidin, and malvidin, with cyanidin being the most abundant in nature. One berry known for its particularly high anthocyanin content is the black raspberry (BRB) (approximately 607 mg/ 100 g fresh weight) (Table 1). Studies have indicated that BRB anthocyanins have chemoprotective roles (Stoner et al., 2007; Kresty, Mallery, & Stoner, 2016) and thus may have a potential use as a chemopreventative strategy against cancer. The main anthocyanins in BRBS are cyanidin-3-glucoside, cyanidin-3-sambubioside, cyanidin-3-xylosylrutinoside, cyanidin-3-rutinoside, and pelargonidin-3-rutinoside (Tian, Giusti, Stoner, & Schwartz, 2006).

### 1.6 | Anthocyanin metabolism

Anthocyanins have poor bioavailability, often with as little as 0.1% of the ingested amount being detectable in urine (Wu et al., 2009), but berry metabolites can be detected in urine, feces, blood, and tissues. It is not fully understood why anthocyanins have a low bioavailability, and

**TABLE 1** Anthocyanin content in edible berry fruits

Berry type	Scientific name	Anthocyanin content (mg/ 100 g FW)	References
Black chokeberry	<i>Aronia melanocarpa</i>	307–1480	Wu et al. (2004); Denev, Kratchanov, Ciz, Lojek, and Kratchanova (2012))
Elderberry	<i>Sambucus nigra</i>	332–1374	Määttä-Riihinen, Kamal-Eldin, Mattila, González-Paramás, and Törrönen (2004); Wu et al. (2004)
Bilberry	<i>Vaccinium myrtillus</i>	300–808	Määttä-Riihinen et al. (2004); Ogawa et al. (2008); Prior et al. (1998)
Black raspberry	<i>Rubus occidentalis</i>	145–607	McGhie, Hall, Ainge, and Mowat (2002); Moyer, Hummer, Finn, Frei, and Wrolstad (2002)
Blueberry	<i>Vaccinium corymbosum</i>	63–430	Corona, Tanga, Vauzoura, Rodriguez-Mateosaand, and Spencer (2011); Moyer et al. (2002); Ogawa et al. (2008); Prior et al. (1998)
Cranberry	<i>Vaccinium macrocarpon</i>	19–360	Corona et al. (2011); Moyer et al. (2002); Ogawa et al. (2008); Prior et al. (1998)
Evergreen blackberry	<i>Rubus laciniatus</i>	91–164	Ogawa et al. (2008); Siriwoharn, Wrolstad, Finn, and Pereira (2004); Wada and Ou (2002)
Marionberry	<i>Rubus ursinus</i>	62–200	Deighton, Brennan, and Davies (2000); Siriwoharn et al. (2004); Wada and Ou (2002)
Red raspberry	<i>Rubus idaeus</i>	28–104	Corona et al. (2011); McGhie et al. (2002); Moyer et al. (2002); Ogawa et al. (2008)
Strawberry	<i>Fragaria × ananassa</i>	13–55	Cordenunsi, Oliveira do Nascimento, Genovese, and Lajolo (2002); Corona et al. (2011); Ogawa et al. (2008)

Anthocyanin content (mg/ 100 g fresh weight (FW)) in selected berries. Range of anthocyanin content found in several berry fruits.

thus this could be seen as a major limitation to the potential efficacy of berry chemopreventative interventions outside of the gastrointestinal tract. However, as ~60% of ingested anthocyanins reach the colon and come into direct contact with the gastrointestinal epithelium, it may explain why berries have higher protective roles against gastrointestinal cancers (Johnson, 2009). Anthocyanin metabolism occurs within three distinct regions of the gastrointestinal tract, the stomach, the small intestine, and the colon. These regions are distinct from one another due to varying pH and microbial diversities. Anthocyanins are most stable in the stomach due to a low pH (pH 1–2), and it is here intact anthocyanins are absorbed into the circulatory system. Most anthocyanins survive intestinal transit where they eventually get metabolized by the colonic microflora into a plethora of phenolic acids and other degradative products before being absorbed into the circulatory system. Anthocyanins can be subclassified into sugar-free anthocyanidin aglycones and anthocyanin glycosides. The intestinal microflora hydrolyze the sugar moieties (glycosides) attached to anthocyanins, the higher pH (neutral) in the colon aids in the metabolism of anthocyanin aglycones to phenolic acids (Johnson, 2009; Wu et al., 2009) (Figure 2).

## 1.7 | Black raspberries

The black raspberry (*Rubus occidentalis*) is a minor but specialty fruit of eastern North America and shares its name with its western North American relative *Rubus leucodermis*. BRBs contain a range of nutrients and bioactive phytochemicals such as vitamin A, C, and E, folic acid, calcium, selenium,  $\beta$ -sitosterol, fiber, a variety of polyphenols such as


ellagic acid, ferulic acid, ellagitannins, most abundantly anthocyanins, and other bioflavonoids (Stoner, 2009). Due to their expansive health benefits and protective properties against many human diseases and cancer, BRBs have been the most studied berry (Johnson, 2009). The chemopreventative properties of BRBs, in particular, their anthocyanin counterparts have been investigated in vitro and in vivo in both human and animal models (Kresty et al., 2016) with the best documented studies being for oral, esophageal and colon cancers (Zikri et al., 2009).

## 1.8 | Chemopreventive studies of black raspberries

### 1.8.1 | Chemopreventive studies of black raspberries in oral cancer

Oral cancer is the most common type of head and neck cancer, and in 90% of cases it presents as a squamous cell carcinoma (SCC) (El-Bayoumy et al., 2017) with strong evidence linking it to exogenous factors such as smoking and excessive alcohol consumption (Hashibe et al., 2009; Torre et al., 2015; El-Bayoumy et al., 2017). In 2006, a study investigated the effects of freeze-dried BRB ethanol extracts at varying concentrations on cell lines derived from human SCC. The results indicated that the BRB extract had antiproliferative effects in a dose-dependent manner and could induce apoptosis, terminal differentiation, and suppress the translation of VEGF (vascular endothelial growth factor; potentially inhibiting angiogenesis which would aid tumor development) (Rodrigo et al., 2006).

In a clinical study with oral intraepithelial neoplasia (IEN), a precursor of oral SCC, topical application of a 10% freeze-dried BRB



**FIGURE 2** Overview of black raspberry metabolism. Schematic representation of the organs involved in the absorption and metabolism of black raspberry polyphenols

mucoadhesive gel was administered (0.5 g) to the tongue four times a day for 6 weeks. The study reported that the lesion grade decreased in 41% of patients, while 23% of patients had an increase in lesion grade and 35% of participants remained at a stable state. Importantly, the study concluded that freeze-dried BRB gel was safe to use as patients enrolled in the clinical trial reported no BRB gel associated toxicities (Shumway et al., 2008). The BRB-induced reduction in size and grade of oral IEN lesions may in part, be attributed to increased levels of proapoptotic and terminal differentiation genes, reduced epithelial levels of the proinflammatory proteins cyclooxygenase-2 (COX-2), and inducible nitric oxide synthase (iNOS). However, the study reported significant interpatient variation (Mallery et al., 2008).

The success of these studies led to the development of a multicenter placebo-controlled trial, investigating the efficacy of the BRB gel over a 3-month period. The previously described dosing regime was utilized in patients with confirmed oral IEN. Compared to placebo-treated patients, BRB gel intervention resulted in significant reductions in both lesion size and grade (Mallery et al., 2014). In a more recent clinical trial, oral cancer patients were treated with dissolvable slow-release BRB troches which equated to a daily dose of 4.3 g freeze-dried BRB powder. Treatment with these BRB troches resulted in the reduction of proinflammatory and prosurvival genes and reduced biomarker expression of genes dysregulated in oral cancers (Knobloch et al., 2016).

Since these clinical trials, substantial work has focused on the development of suitable *in vivo* animal models that better represent carcinogen induced oral dysplasia's in humans, such that the chemopreventive mechanisms of BRBs could be further investigated. In 2012, a novel nonsurgical murine model of oral cancer was developed, whereby animals were exposed to an environmental pollutant and component of tobacco smoke dibenzo[*a,l*]pyrene, and its diol epoxide metabolite. Topical exposure of these carcinogens to the oral cavity of mice resulted in 74% and 100% oral squamous cell carcinoma (OSCC) in tongue and oral tissues, respectively; mediated via the induction of DNA adducts, prerequisites for oral carcinogenesis (Guttenplan et al., 2012; Chen et al., 2013). These mice were fed either a control diet (AIN-93 M) or a diet supplemented with 5% BRB powder for 2 weeks prior to the induction of tumorigenesis in the oral cavity. The study reported that administration of a 5% BRB diet prior to carcinogen exposures resulted in fewer DNA adducts and abrogated tumor incidence from 93% to 66% (Chen et al., 2018). It was later reported that BRB-mediated reduction in oral mutagenesis could be attributed, in part, to epigenetics and hypermethylation of genes involved in epithelial-mesenchymal transition (Chen et al., 2020). These BRB-mediated metabolomic effects in oral cancer are further supported in a study by Knobloch et al. (2019), using a chemically-induced model of oral cancer, (4-nitroquinoline-1-oxide) (El-Bayoumy et al., 2017), whereby 6-week exposure to a 10% BRB diet inhibited enzymes fundamental for the metabolic pathway of

glycolysis, thereby limiting nutrient availability for cancer cells (Knobloch et al., 2019).

### 1.8.2 | Chemopreventive studies of black raspberries in esophageal cancer

Worldwide, esophageal cancer ranks the seventh most common cancer (Bray et al., 2018). Esophageal cancers present as one of the two most common histological subtypes; SSC or adenocarcinoma and is linked to excessive alcohol consumption, tobacco carcinogens, or drinking scalding hot mate (a plant-based infusion drunk in parts of South America), whereas adenocarcinomas are often attributed to higher socioeconomic countries where obesity and gastroesophageal reflux disease (GERD) are on the rise (Blot & Tarone, 2018; World Cancer Research Fund/American Institute for Cancer Research, 2018).

In 2001, *in vivo* studies using the NMBA (*N*-nitrosomethylbenzylamine) rat model of esophageal cancer reported that feeding of 5 and 10% lyophilized BRB diet for 2 weeks prior to NMBA exposure and then throughout a 30-week time period, resulted in significant reduction in tumor multiplicity and inhibited the formation of mutagenic DNA adducts (Kresty et al., 2001). Additional studies with this model have reported that BRB diet reduced rat esophageal cancer by decreasing DNA adducts (specifically  $O^6$ -methylguanine), inhibiting proliferation and dampening down expression of proinflammatory enzyme Cox-2 (Stoner et al., 2006), reversing oxidative stress (Stoner et al., 2006; Shi et al., 2017), and via inhibition of proangiogenesis genes such as *Vegf* (Chen, Rose, Hwang, Nines, & Stoner, 2006). BRBs have also been shown to inhibit NMBA-induced esophageal cancer by inhibiting the Wnt signaling pathway through alleviation of hypermethylation of the Wnt pathway antagonist, *Sfrp4* (secreted frizzled related protein 4), and reduction in mRNA levels of DNA methyltransferases (DNMTs), *Dnmt1* and *Dnmt3b*. These BRB-mediated alterations resulted in significant reductions in nuclear  $\beta$ -catenin localization thus preventing proliferation of rat esophageal squamous cell papilloma (Huang, Gu, Dombkowski, Wang, & Stoner, 2016).

In 2016, BRBs were used in a phase I pilot study in patients with Barrett's esophagus (BE), a precursor lesion for esophageal adenocarcinoma. To determine the long-term tolerability of BRBs in esophageal cancer, a 6-month pilot study was performed. Based on rodent pre-clinical studies, patients confirmed to have BE were administered lyophilized or freeze-dried BRB powder at 32 and 45 g (~1.5 and 2 cups of whole fruit; Kresty et al., 2001; Stoner et al., 2006; Lechner et al., 2008; Zikri et al., 2009). Results showed that lyophilized BRBs were generally well tolerated with a few adverse effects including epigastric pain, diarrhea, and constipation and resulted in increased levels of GST-pi (a marker of detoxification) in over 50% of patients. Suggesting BRBs may help prevent BE progression by detoxification of the mutagenic components involved in gastroesophageal reflux disease (Kresty et al., 2006, 2018).

In addition, the chemopreventive properties of BRBs in esophageal cancer have been linked in part to their effects on inflammation and

immune cell trafficking. Exposure to 6.1% BRB diet, a BRB-derived anthocyanin-enriched fraction (3.8  $\mu$ mol/g) or treatment with protocatechuic acid, a major microbial-generated metabolite of BRB anthocyanins, inhibited NMBA-induced rat esophageal cancer. Mechanistically, this was found to occur through inhibition of inflammatory genes *Cox-2* and *IL-1 $\beta$* , reduced infiltration of inflammatory macrophages into dysplastic lesions, limited neutrophil and cytokine accumulation, and decreased angiogenesis (Peiffer et al., 2014, 2016).

Similar results to both of these studies have been reported using human esophageal microvascular endothelial cells (HEMEC) stimulated with either the proinflammatory cytokine tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) or IL-1 $\beta$ . HEMEC cells were pretreated with BRB ethanol extract for 2 hr before activation with TNF- $\alpha$  or IL-1 $\beta$ . BRB ethanol extract had a strong anti-inflammatory effect by inhibiting nuclear factor- $\kappa$ B (NF- $\kappa$ B) nuclear translocation, reducing COX-2 gene and protein levels in stimulated HEMECs, abolished the proangiogenic effects of VEGF by inhibiting cell migration, proliferation, and survival of VEGF-stimulated HEMECs (Medda et al., 2015).

### 1.8.3 | Chemopreventive studies of black raspberries in CRC

CRC, which encompasses both colon and rectal tumors, is the third most common cancer in men (10% of total cancer cases) and the second most common cancer in women (9.2% of total cancer cases) worldwide (Ferlay et al., 2015). In 2018 alone, it was estimated that there were more than 1.8 million new CRC diagnoses and 881,000 deaths reported (Bray et al., 2018), highlighting the burden CRC puts on the healthcare infrastructure. CRC incidence displays wide geographical variation globally, with similar patterns seen for both men and women. CRC is often considered a marker of socioeconomic development since it mostly occurs in countries undergoing major development and nutritional transition and in those which have a high human development index (Ferlay et al., 2015; Bray et al., 2018; World Cancer Research Fund/American Institute for Cancer Research, 2018).

The majority of CRCs are adenocarcinomas (approximately 95%) which occur sporadically, with only a small proportion of CRC cases attributable to inherited genetic mutations (~10%); mainly FAP and Lynch syndrome. There is evidence that CRC likely develops through an interplay of endogenous (epigenetic, microbial status, inflammation, immune status) and exogenous factors (diet composition, stress, alcohol consumption, smoking, and obesity) which increase mutation rates (Paul et al., 2015). In the UK alone, it is estimated that around half of CRC cases could have been prevented through lifestyle modifications such as alteration to the diet and reduction in body fatness (World Cancer Research Fund/American Institute for Cancer Research, 2018). The combination of damage from carcinogens that directly interact with the intestinal and colonic epithelium, and the mechanical deformation likely underpins the high rate of intestinal stem cell (ISC) division, which is required to maintain an intact epithelial barrier. In turn, this can lead to increased CRC risk due to a buildup of oncogenic DNA mutations in ISCs (Barker, 2014).

In 2001, a study investigated the protective effects of lyophilized BRBs in the azoxymethane (AOM) rat model of CRC (Harris et al., 2001). AOM-induced rodent CRC occurs via the production of DNA adducts following the AOM metabolism, and faithfully models sporadic and hereditary human CRC development sharing similar genetics, histopathological characteristics, and defective mismatch repair to the human disease (Stevens et al., 2007). Twenty-four hours after final treatment with AOM, rats were fed either 0%, 2.5%, 5%, or 10% freeze-dried BRB diet for 9 and 33 weeks. The total tumor multiplicity decreased by 42%, 45%, and 71%, respectively, in a dose-dependent manner, tumor burden also decreased in a similar fashion but was not found to be significantly different. This demonstrates the ability of BRBs to reduce tumorigenesis in a rodent model of colon carcinogenesis (Harris et al., 2001).

In 2010, Bi and colleagues performed *in vivo* experiments in two genetic murine models of human CRC to investigate whether BRBs would inhibit tumorigenesis (Bi, Fang, Wang, Stoner, & Yang, 2010). The two mouse models used were *Apc1638<sup>+/-</sup>*, which develops small intestinal tumors as a result of having only one functional allele of *Apc* with subsequent loss of the wild-type allele leading to aberrant Wnt signaling (Fodde et al., 1994) and the *Muc2<sup>-/-</sup>* mouse which develop spontaneous intestinal and colonic adenomas in response to chronic inflammation (Velcich et al., 2002). A 12-week feeding program of 10% freeze-dried BRB diet significantly decreased tumor incidence and tumor multiplicity in *Apc1638<sup>+/-</sup>* mice by 45% and 60%, respectively. Similarly, in *Muc2<sup>-/-</sup>* mice, feeding of BRBs reduced tumor incidence and multiplicity by 50%. BRBs appeared to inhibit proliferation in both models, whereas terminal differentiation of goblet cells was not altered, potentially due to a relatively short feeding program. Quantification of other cell types, such as enteroendocrine, Paneth, and enterocytes may provide a better insight into the effects of BRBs on intestinal cell differentiation in the future. Mechanistically, it was reported that reductions in tumorigenesis within the two models was via two different biological pathways. BRBs elicited antitumor effects in *Apc1638<sup>+/-</sup>* mice through inhibition of Wnt/ $\beta$ -catenin signaling and its downstream targets, including *c-Myc* and *CyclinD1*, whereas in the *Muc2<sup>-/-</sup>* mouse BRBs downregulated cytokines commonly associated with inflammation, including Cox-2, Tnf- $\alpha$ , IL-1, IL-6, and IL-10 (Bi et al., 2010). These models were the first to suggest potential mechanisms of action of BRBs against *in vivo* CRC development.

The hereditary condition FAP is characterized by an early onset of colonic polyposis which confers an increased risk of developing CRC. A key phase Ib study was conducted to investigate the effects of BRBs on FAP patient rectal polyps (Wang et al., 2014). Patients with at least five,  $\geq 2$  mm, rectal polyps at baseline endoscopy were enrolled into the study and randomly divided into two groups consisting of seven patients each. Group 1 received placebo powder slurry (maltodextrin and dextrose) taken orally three times a day (60 g/day in total) and two BRB rectal suppositories at night (each containing 720 mg BRB powder). Group 2 received a BRB powder slurry three times a day (60 g/day in total) plus two BRB rectal suppositories. BRB intervention was continued for up to 9 months. The study showed that, despite three non-responders, BRB rectal suppositories were sufficient to reduce polyp

burden and regress rectal polyps, but there was no added benefit from oral BRB powder consumption (Wang et al., 2014). Within responding adenomas, BRB treatment decreased cellular proliferation, and positively impacted on epigenetic marks and proteins by decreasing promoter methylation, but had no effect on cell death (Wang et al., 2014). These results suggest that BRBs may be a potential adjuvant treatment to improve the outcome for FAP patients.

### *Black raspberries and epigenetics*

Epigenetics refers to the study of the heritable and dynamic chemical modification of a cell's transcriptional capacity, without any alteration to the genetic sequence (Allis & Jenuwein, 2016). Epigenetic mechanisms include but are not limited to histone modification, modulation of DNA methylation status and miRNA-mediated manipulation of translation; these mechanisms regulate gene expression and can impact upon tumorigenesis. Several major enzymes that catalyze these mechanisms: DNMTs; 10 eleven translocation (TET) enzymes; histone acetyltransferases, and histone deacetylases (HDACs) (Allis & Jenuwein, 2016). Metabolites resulting from processed dietary phytochemicals and fiber can affect these enzymes and epigenetic mechanisms, and ultimately gene expression. Specifically, in the context of cancer, genes involved in tumor suppression are usually hypermethylated resulting in epigenetic gene silencing; a mechanism that supports tumor survival (Costello et al., 2000).

Previous studies by Wang et al. have reported an association between BRB-derived anthocyanins and a decreased hypermethylation status seen in CRC, suggesting that BRBs could play a role in modulating epigenetic events (Wang et al., 2011). Typically, promoter regions of Wnt pathway antagonists such as *SFRP2*, are hypermethylated in human CRC, leading to aberrant Wnt signaling and tumor progression. Human CRC cell lines treated with BRB-derived anthocyanins for 3 days at 0.5, 5, and 25  $\mu\text{g}/\text{mL}$  suppressed DNMT1 (maintenance DNA methyl transferase) and DNMT3b (de novo DNA methyl transferase) activity. Additionally, anthocyanin (AC) exposure resulted in the reactivation of Wnt pathway repressors, *CDKN2A*, *SFRP2*, *SFRP55*, and *WIF1*, via promoter demethylation. This resulted in the induction of apoptosis, cell cycle arrest, and mRNA suppression of key Wnt regulators,  $\beta$ -catenin, and C-MYC (a downstream target of the Wnt pathway), in these CRC cell lines (Wang et al., 2013).

The role of BRBs on epigenetic silencing has been evaluated in pre-cancerous murine models that recapitulate human ulcerative colitis (UC, an intermediary step of colon carcinogenesis that is also influenced by deregulation of the Wnt pathway). IL-10 knockout UC mice fed a 5% BRB diet had less colonic ulceration as a result of reduced nuclear localization of  $\beta$ -catenin than respective control mice (Wang et al., 2013). This study also reported that mRNA expression of Wnt repressors was significantly upregulated in BRB-treated mice, which coincided with hypomethylation of the same gene promoters and decreased protein levels of methylation-regulating proteins DNMT, HDACs, and methyl-binding domain proteins (Mbd2), compared to control fed UC mice (Wang et al., 2013). Studies from the same group using a dextran sodium sulfate (DSS)-induced, an irritant that induces extensive colonic mucosal inflammation, UC model reported that 5% BRB

diet significantly reduced colonic ulceration and decreased protein levels of Dnmt3B, Hdac1, Hdac2, and Mbd2 and reduced promoter methylation of Wnt repressor genes (Wang et al., 2013). In AOM/DSS chemically induced murine cancer models, diets containing 3.5 and 7  $\mu\text{mol/g}$  BRB-derived ACs, which corresponded to the AC concentrations in 5 and 10% freeze-dried BRB powder, respectively, resulted in significant reduction in Dnmt1 protein expression. However, BRB anthocyanins had no effect on TET enzyme mRNA expression (which mediate active DNA demethylation) in the intestinal epithelium at either concentration (Chen et al., 2018), this suggests that BRB-induced demethylation is a passive process in this context. Similarly, to other studies, the cancer preventative effect of BRBs was also linked to the promoter demethylation of *Sfrp2*, and subsequent increase in *Sfrp2* protein concentration negatively regulating Wnt/ $\beta$ -catenin signaling (Chen et al., 2018).

In a phase 1 clinical study of 20 CRC patients who were treated with oral BRB powder (60 g/day) for 1-9 weeks, epigenetic changes were only observed in patients who only received on average 4 weeks of BRB intervention, with reduced promoter methylation of key Wnt pathway inhibitors and decreased expression of DNMT1 (Wang et al., 2011). Epigenetic changes were confirmed in a subsequent phase 1b clinical trial, in which 14 FAP patients received BRB intervention through rectal suppositories with or without the addition of oral supplementation for 9 months. Here the majority of patients (11/14) responded to the BRB treatment with polyp regression and significant decreases in protein levels of DNMT1 with reactivation of the silenced p16 promoter (Wang et al., 2014). Providing direct clinical evidence for berry-mediated chemoprevention.

Aside from BRB anthocyanins, the dietary fiber obtained from sources such as BRBs can also protect against tumorigenesis through several mechanisms. First, increased speed of colonic transit helps minimize epithelial exposure to carcinogens and, second, metabolism of fiber by gut bacteria, such as *Anaerostipes* or *Clostridia*, into short-chain fatty acids (SCFAs) such as butyrate, acetate, and propionate, can cause tumor-influencing epigenetic alterations (Park, Mitrou, Luben, Khaw, & Bingham, 2009). SCFAs are a source of energy but also act as signaling molecules within the body. Butyrate is the main source of energy for colonocytes and can also induce apoptosis in cancer cells (Hague et al., 1993). Intracellular butyrate and propionate can inhibit gut HDAC activity, promoting hyperacetylation of histones (limiting methylation and thus reducing epigenetic silencing of antitumorigenic genes) and can impact on signal transduction (Waldecker, Kautenburger, Daumann, Busch, & Schrenk, 2008). H3 acetylation in the promoter and conserved noncoding (CNS1) regions of the *Foxp3* locus (which controls colonic T-regulatory cell population thought to be cancer preventative) is enhanced by butyrate-mediated HDAC inhibition (Furusawa et al., 2013). Butyrate, along with propionate and acetate, activate free fatty acid receptor 2 (FFAR2), the receptor for SCFAs which regulates whole-body energy homeostasis and is an important regulator of inflammation involved in enhancing T-reg cell accumulation. Stimulation of FFAR2 via SCFAs is essential for innate immunity and chemoattractant-induced responses; SCFA-mediated T-reg accumulation is important in limiting inflammatory responses associated with

tumorigenesis (Maslowski et al., 2009; Furusawa et al., 2013). In disease states or conditions resulting from inadequate diets, such as CRC, the levels of SCFA may be diminished, which can mitigate the protective effects of SCFA-mediated immune regulation (Chen et al., 2013).

In more recent years, there have been emerging roles of microRNAs (miRNAs, small noncoding RNAs) in epigenetic regulation. miRNAs can act as epigenetic regulators posttranscriptionally, altering target mRNAs and interfering with protein translation without modifying gene sequences. This is often via the induction of mRNA degradation or repression, physical interaction of miRNAs with RNA interference machinery (Yao, Chen, & Zhou, 2019). In addition to being epigenetic regulators, miRNAs can themselves be regulated by the epigenetic machinery involved in DNA methylation and RNA and histone modifications. miRNAs have been implicated in several fundamental biological processes such as proliferation and migration. The impact of miRNAs on physiological and pathophysiological processes can be either beneficial or detrimental depending on the specific miRNA and its target, and dysregulation of this complex miRNA-epigenetic feedback loop can contribute to tumorigenesis (Yao et al., 2019). Studies have shown that diet, and the subsequent downstream effects of dietary components on the microbiome and production of metabolites, can influence and interact with miRNAs. miR-92a, a member of the miR-17-92a cluster important in regulating cell cycle, proliferation and apoptosis have been shown to be 7x higher in human CRC tissue compared to adjacent healthy tissue from the same patient (Hu, Liu, Chang, Wang, & Raufman, 2015). miR-92a has been implicated in tumor development through its suppressive interaction with the antiapoptotic molecule, BCL-2-interacting mediator of cell death, in human CRC cell lines, and colonic cancer tissue (Tsuchida et al., 2011). In human CRC cell lines, microbe-derived butyrate has been shown to reduce the levels of miR-92a and other miRNAs belonging to the miR-17-92a cluster. A cancer protective role of butyrate was identified via butyrate's silencing effect on c-Myc protein levels leading to a reduction in c-Myc-induced miR-17-92a promoter activity and ultimately reducing levels of oncogenic miR-92a synthesis (Hu et al., 2015). A recent study has identified a BRB-mediated chemopreventative association with miR-24-1-5p in CRC through inhibition of Wnt/B-catenin signaling and subsequent downstream targets involved in CRC tumorigenesis (Takebe et al., 2013). Utilizing the colitis-induced CRC model (AOM/DSS), expression levels of miR-24-1-5p was found to be low in tissues from AOM/DSS-induced mice but was significantly increased in AOM/DSS mice following 9-week feeding intervention with BRB anthocyanins (7  $\mu\text{mol/g}$  BRB anthocyanins equating to a 10% BRB diet; Zhang et al., 2019). It was shown that upregulation of miR-24-1-5p in human CRC cell lines leads to inhibition of cancer cell proliferation, migration, and survival, by suppressing  $\beta$ -catenin protein levels and modulating downstream target gene expression (Zhang et al., 2019). FAP patients who responded to 9 month BRB suppository intervention had 37 unique miRNAs that were demethylated in adenomas, while in these same patients promoter methylation of Wnt suppressors was not altered, indicating the uniquely demethylated miRNAs regulate the Wnt signaling pathway (Wang et al., 2014). Together these studies highlight a potential mechanism whereby BRBs and their anthocyanin derivatives can

mediate tumor suppression through miRNA epigenetic interaction with  $\beta$ -catenin and downstream Wnt target genes. We are yet to fully establish the chemopreventative relationship between BRBs and epigenetic gene regulation in the colon; however, there is increasing evidence suggesting that BRBs and their microbial-derived metabolites play roles in regulating the epigenome and can influence carcinogenesis.

### *Black raspberries, microbiome, and the metabolome*

The microbiota of the human gut is a symbiotic web consisting of trillions of microorganisms: primarily bacteria, but also viruses, archaea, fungi, and protozoa (Ursell, Metcalf, Parfrey, & Knight, 2012). The human gut microbiome refers to the collective genomes of these microorganisms, in which there are approximately 3.3 million genes encoded (Qin et al., 2012). The importance of the gut microbiota on human physiology has become increasingly apparent over the past decade. With the advent of “omic” technologies and the rapidly developing analytical and sequencing techniques that follow we can now study the impact of the microbiota on health and disease in great detail.

The human microbiota is often referred to as a hidden organ, with its own physiology and pathophysiology (Baquero & Nombela, 2012). It is essential in regulating normal gut homeostasis and health and is a relatively stable entity once it has been established. However, the microbiome can be manipulated and altered through intervention with antibiotics or changes in diet, resulting in dysbiosis and disease. Dysbiosis of the intestinal microbiota, contributes to many disease states including obesity, diabetes, and inflammatory bowel disease, all of which may predispose to CRC (Ramos-Nino, 2013; González et al., 2017).

Excessive consumption of dietary elements such as protein, sugar, and saturated fatty acids can adversely affect the gut microbiota, decreasing microbial diversity and resulting in inflammation and decreased production of butyrate (Tu et al., 2020). Intake of fiber and polyphenols positively influence the microbiota, resulting in increased frequency of beneficial gut microbes that exert biological effects including reduced inflammation, increased antioxidant production, and increased butyrate production (Pan et al., 2017). BRBs, and the abundance of fiber and polyphenols such as anthocyanin and ellagitannin, contained within them, are capable of altering the composition and metabolic potential of the gut microbiota (Gu et al., 2019). A BRB diet has been shown to positively influence the ratio of beneficial gut microbes including those of the genera *Anaerostipes*, *Akkermansia*, *Desulfovibrio*, and members of the phylum *Bacteroidetes* (Pan et al., 2017; Tu et al., 2018).

Metabolomic studies have identified an enrichment in several beneficial metabolic pathways and metabolites, including butyrate, in mice fed a diet rich in BRB (Tu et al., 2020). Rats fed a BRB diet showed an increase in butyrate-producing bacteria such as *Anaerostipes* when compared to a control diet (Pan et al., 2017). It has been shown that FFAR2, the receptor for SCFAs, is required for the beneficial effects of BRBs; indicating bacterial fermentation is an important mechanism underpinning BRB-mediated benefits (Pan et al., 2018). Colonic inflammation and the subsequent cytokine and ROS generation results in

an increase in opportunistic, adhesion-invasive bacterial species such as *Escherichia coli* that further promote inflammation, and a decrease in species from *Akkermansia* and *Desulfovibrio* genera that have anti-inflammatory effects (Tu et al., 2018). Intervention with a BRB diet can reduce inflammation and the increased risk of tumorigenesis associated with it by increasing the abundance of these anti-inflammatory bacterial species and via increased activity of FFAR2, which has been implicated in suppressing colonic inflammation.

Several studies have demonstrated the association of an increased *Firmicutes* to *Bacteroidetes* ratio with diet-induced obesity (Ley et al., 2005). *Firmicutes* are positively selected for in obesity as member species have an increased capacity for processing polysaccharides, a major component of diet-induced obesity (Turnbaugh, Bäckhed, Fulton, & Gordon, 2008). Furthermore, it has been reported that obese individuals have a decreased population of *Akkermansia muciniphila* which, as previously stated, is a bacteria that has anti-inflammatory effects as well as antidiabetic and antiobesity functions, including roles in glucose homeostasis and adipose metabolism (Tu et al., 2018). A BRB-rich diet has been shown to modulate the murine gut microbiota, increasing the abundance of *Bacteroidetes* compared to *Firmicutes* and the abundance of *A. muciniphila* (Tu et al., 2018). This microbial modulation via BRB intervention provides a number of bacterial metabolic alterations, including increased biosynthesis of vitamins and essential amino acids, that provide health benefits to the host such as reduced inflammation and increased cardiac and neural tube protection (Said, 2011; Tu et al., 2018). Regulating inflammation and helping to maintain a healthy weight are some of the key elements of BRB-mediated chemoprevention.

The phytochemicals contained within BRBs are processed and absorbed by the various microbes in the gut, but the subsequent metabolites can be transported through the body and exert systemic beneficial effects. Metabolomic approaches have shown alteration in over 100 metabolites across the colonic mucosa, liver, and the feces of healthy mice fed a BRB-supplemented diet. BRB supplementation decreased certain lipid metabolites in the colon including long-chain fatty acids and omega-6 polyunsaturated fatty acids, which can contribute to insulin resistance, inflammation, and obesity when consumed in excess. Conversely, BRBs led to an increase in cancer protective omega-3 polyunsaturated fatty acids ( $\omega$ -3 PUFA), such as linolenate, in the liver (Montgomery et al., 2013; Pan et al., 2017). Linolenate is converted to docosapentaenoate, docosahexaenoate (DHA), and eicosapentaenoate, each of which has been shown to provide metabolic, anti-inflammatory, and neuroprotective benefits (Dyall, 2015). Clinical studies have also indicated an improvement in body weight and physical, cognitive, and social functions in lung cancer patients given a daily supplement of DHA and EPA, further demonstrating how BRBs exert their chemoprotective effects (Sánchez-Lara et al., 2014).

Several species of bacteria that have generated interest due to their tumorigenic potential in the gastrointestinal tract, namely *Fusobacterium nucleatum*, *Escherichia coli*, and *Bacteroides Fragilis*. In both cell lines and patient-derived xenograft murine models, the presence of *F. nucleatum* increases tumor cell proliferation and overall tumor growth rates (Bullman et al., 2017). *E. coli* that express the genomic island

polyketide synthase can produce DNA-alkylating genotoxins that may result in mutagenic DNA damage leading to tumorigenesis (Wilson et al., 2019). *B. Fragilis*, along with *F. nucleatum* and *E. coli* can create a tumorigenic biofilm that facilitates bacterial epithelial infiltration, recruits immune cells, and promotes inflammation in precancerous colonic lesions (Dejea et al., 2018). BRB intervention has been shown to increase the abundance of butyrate-producing bacteria; butyrate stimulates mucus production and maintains epithelial integrity, possibly protecting against the adverse effects resulting from biofilm generation (Peng, Li, Green, Holzman, & Lin, 2009).

As previously described, the microbiota has the capacity to influence many disease states in the gut. Pathobiotic bacteria, such as *E. coli*, which accumulate in a poor-diet or disease-state gut, can penetrate the gut epithelium and generate toxins that promote colonic inflammation and tumorigenesis (Sheflin, Whitney, & Weir, 2014). Studies using mice have shown that in circumstances such as these, antibiotics can be used to suppress this pathobiotic accumulation, and ultimately, the adverse effects associated with it (Tanaka, Ito, & Isobe, 2016). Conversely, the use of antibiotics can also reduce the frequency of beneficial gut microbes and depletion of the gut microbiota may encourage the growth of antibiotic-resistant bacteria that can further induce inflammation (Dethlefsen & Relman, 2011). The gut microbiota is essential in facilitating the beneficial effects of BRBs, and any perturbation of the gut microbial content has the capacity to interfere with these effects. Administration of a 5% BRB diet to *Apc<sup>Min/+</sup>* mice (which mimics FAP disease) did not affect polyp development in mice that had been treated with antibiotics (1 g/L ampicillin, 1 g/L neomycin, 1 g/L metronidazole, and 0.5 g/L vancomycin in the drinking water) (Pan et al., 2018). This demonstrates that the gut microbiota is indeed required for BRB-mediated chemoprevention. Homozygous deletion of *Apc* within the ISC compartment of mice, that recapitulates the initial stages of CRC initiation, has also been shown to alter the composition of the microbiota as seen with trends of increasing abundance of *Bacteroides* and *Clostridium XVII* (May, McDermott, Marchesi, & Parry, 2020) which can contribute to the tumorigenic biofilm. The gut microbiota is affected very early on in tumorigenesis and has the potential to serve as an early diagnostic tool. Mice fed a diet of 10% BRBs from 2-weeks prior to induction of ISC *Apc* loss have a reduced impact to the gut microbiota upon tumorigenesis, indicating that BRBs can abrogate the initial cancer-mediated alterations to the microbiota and promote a favorable *Bacteroidetes* to *Firmicutes* ratio which supports a leaner bodyweight (May et al., 2020). Further work with larger sample sizes is needed to fully characterize the effect of BRBs on the composition and function of the microbiota in CRC.


### Black raspberries, inflammation, and immunity

Chronic inflammation is a major risk factor for CRC and is considered a hallmark of cancer (Hanahan et al., 2011). Inflammation involves a complex network consisting of numerous immune and inflammatory cells, cytokines, chemokines, and proinflammatory mediators (Lasry, Zinger, & Ben-Neriah, 2016). Inflammation and subsequent accumulation of immune cells in inflammatory areas can lead to the activation of signaling pathways that promote tumor growth and invasion.

Increased activation of inflammatory signaling pathways can often lead to genetic instability, an increased incidence of oncogenic mutations and result in inflammation-mediated carcinogenesis (Ding et al., 2005). BRB intervention can help to reduce inflammatory-mediated tumorigenesis through its inhibition of certain proinflammatory mediators and its regulation of immuno-modulatory cells (Stoner et al., 2007; Kresty et al., 2016).

The proinflammatory mediator COX-2 is elevated in many cancers including CRC and can be rapidly and transiently induced by inflammatory cytokines and growth factors (Eberhart et al., 1994). COX-2 is an enzyme responsible for the production of prostaglandins that enhance tumor growth and can aid in tumor evasion from the immune system by modulating the tumor microenvironment (Zelenay et al., 2015). A further role of COX-2 in tumorigenesis is its capacity to increase tumor invasiveness and metastatic potential (Tsuji, Kawano, & DuBois, 1997). Several studies have demonstrated that BRB diet can suppress COX-2 expression through inhibition of the transcription factor, (NF- $\kappa$ B), which is involved in signaling pathways (e.g., Mitogen-Activated Protein Kinase (MAPK) Pathway signaling), proliferation, cell adhesion, and immune/inflammatory modulation (Huang et al., 2002; Montrose et al., 2011; Chen, Shi, & Afzali, 2019). In murine macrophages stimulated by the endotoxin lipopolysaccharide (LPS), administration of an anthocyanin-rich BRB fraction reduced LPS-induced inflammation (by dampening iNOS expression) and through inhibition of the MAPK pathway (Jo et al., 2015). More recently, a unique phytochemical from berry fruits known as triterpenoid glycosides have been implicated in disease prevention. Specifically, oral administration of a triterpenoid-rich fraction (TFRC; 25, 50, or 100 mg/kg/day) from the Korean black raspberry (*Rubus coreanus*) has been shown to ameliorate colitis in DSS-treated mice, maintaining crypt structures and reducing markers of inflammation including reductions in *Tnf- $\alpha$* , *IL-1 $\beta$* , and *IL-6* mRNA levels. TFRC protected against colitis and in turn carcinogenesis by reducing macrophage accumulation in mice and inhibited nitric oxide and prostaglandin production by suppressing iNOS and Cox-2 levels, downstream inflammatory cytokines, and members of the MAPK/NF- $\kappa$ B pathway in vitro (Shin et al., 2014). Similarly, in DSS-induced colitis mice, a 5% BRB diet reduced macrophage and neutrophil accumulation and suppressed nuclear translocation of NF- $\kappa$ B (Wang et al., 2013).

Natural killer (NK) cells are a type of innate immune lymphocyte that exhibits cytotoxic activity, mediating antitumor, and antiviral responses (Vivier, Tomasello, Baratin, Walzer, & Ugolini, 2008). Depletion of NK cells substantially encourages CRC development, whereas increased NK cell tumor infiltration is associated with a better prognosis, demonstrating the importance of these cells in mitigating tumor progression (Coca et al., 1997). In two colitis-associated CRC mouse models, AOM/DSS and *Apc<sup>Min/+</sup>*/DSS which have a depletion of NK cells in adenomas, a diet supplemented with 5% BRB powder suppressed colon polyp size and frequency while enhancing NK cell tumor infiltration (Pan et al., 2017). Moreover, in colon biopsies from CRC patients treated with oral BRB slurry, BRBs were shown to increase NK cell tumor infiltration (Pan et al., 2017). In addition to NK lymphocytes, another immune cell population that has important immune surveillance roles are T cell lymphocytes. Cytotoxic killer, CD8+, T cells



**FIGURE 3** Summary of the cellular processes involved in BRB-mediated CRC chemoprevention. Black raspberries exert chemopreventative and therapeutic effects against colorectal cancer by acting on several cellular functions, biological processes and signaling pathways that are all interlinked. The protective effects of black raspberry arises from a complex interaction and cross-talk of BRB polyphenols on genes and proteins associated with epigenetics, inflammation, oxidative stress, proliferation, cell cycle regulation, apoptosis, angiogenesis, metastasis, and via modulation of the gut microbiome. So far little is known about the involvement of the intestinal stem cell compartment in BRB-mediated cancer prevention. † = Inhibition; ↓ = Decrease, ↑ = Increase

are involved in immune-mediated cell death and can also act as cytokines, CD4+ T cells, known as helper T cells, secrete cytokines to assist with the immune response, and regulatory T cells mainly act to suppress T cell-mediated immune response. In cancer T cells often become dysfunctional usually aiding cancer progression (Xia, Zhang, Xu, Yin, & Lu, 2019). BRB-ethanol extract and two abundant BRB phytochemical metabolites (cyanidin-3-rutinoside and quercetin-3-rutinoside) can modulate the proliferative capacity of the human CD4+ and CD8+ T lymphocytes. All compounds were able to inhibit activated-T cell proliferation and survival and reduced differentiation of myeloid-derived suppressor cells (MDSC, cells capable of priming the metastatic niche) in vitro, through modulation of the Janus kinase (JAK)/signal transducer and activator of transcription (STAT) pathway which has documented roles in MDSC differentiation. Prior studies have shown that the BRB phytochemicals, Cyanidin-3-glucoside anthocyanin, Quercetin and Rutin can inhibit JAK/STAT and MAPK signaling (Choi et al., 2009; Luo et al., 2012). The beneficial effects of BRBs on inflammation are diverse and numerous, in vitro, the anthocyanin component of BRBs has been more effective at suppressing inflammation

than the commonly used anti-inflammatory drug 5-aminosalicylic acid (Serra, Paixão, Nunes, Dinis, & Almeida, 2013), demonstrating how BRB and their metabolites can be an effective and accessible drug option for treatment of inflammation and inflammation-associated cancer.

## 2 | CONCLUSION

The potential of BRBs as a cancer therapeutic is indisputable, as daily administration of BRBs have shown to reduce colorectal tumor/polyp numbers, oral SCCs, and severity in humans with minimal adverse effects (Shumway et al., 2008; Wang et al., 2011, 2014). Extensive research has revealed that BRBs have pleiotropic effects against all stages of carcinogenesis by promoting apoptosis and differentiation, reducing proliferation, modulating the epigenome, the microbiome, and the metabolome, reducing tumor-mediated inflammation and oxidative stress and inhibiting angiogenesis (Figure 3). Collectively, these studies support the potential of BRBs for therapeutic cancer intervention and highlight the ability of BRBs to limit the window

of opportunity of cancer initiation and progression. However, they highlight the complexity of the involvement of endogenous factors in carcinogenesis and emphasizes the importance of understanding the precise mechanism in which dietary and drug interventions protect against cancer. Moreover, translation into the clinic has not been straightforward, with varied responses occurring among patients (Wang et al., 2011, 2014). This is likely to reflect inter- and intraindividual and tumoral heterogeneity, differences in molecular signaling, and enzyme activity within tumors and variances in individuals' microbiota and metabolism. In addition, the limited bioavailability of anthocyanins, the concentration, frequency, duration, delivery method, and delivery matrix of BRB interventions has differed across studies, which could also be a confounding factor for reporting beneficial outcomes. BRBs have been administered as a slurry of freeze-dried powder suspended in water, as rectal suppositories, as topical agents delivered in a bioadhesive gel, or as dissolvable slow-release BRB troches. It is important to note that the concept of chemoprevention is not to radicalize our lifestyle, but to identify ways in which to adopt a healthier lifestyle and reduce the window of opportunity of cancer development by lessening the risks. Patient compliance is often a rate-limiting step in the success of chemopreventative strategies, and so it is important that such strategies are easy to adhere to. While it remains difficult to accurately estimate the concentration of BRB consumption for individual patients to have beneficial effects, it is necessary that the regime is simple and does not require the consumption of hundreds of berries a day which would not be a sustainable lifestyle change. It is also likely that for each given cancer type and predisposing disease that a specific BRB dose, delivery method and regime is utilized to maximize the efficacy of the treatment.

It is also important to note that the chemopreventative effects of BRBs are not attributed to a single BRB compound but collectively from anthocyanins, ellagic acid, flavonoids, fiber, other antioxidants, and phytochemicals such as quercetin and rutin, and BRB metabolites which may individually impact on specific cellular processes, or work collectively to dampen down signaling pathways and prevent feedback loops and cross-talk among pathways that are fundamental for tumor development. Additionally, edible berries have proven to protect against cancer before the transformation of a premalignant cell, by aiding in the maintenance of a healthy body weight and supporting the growth and abundance of beneficial gut microbes that protect against diseases which predispose individuals to cancer such as obesity, diabetes, gastroesophageal reflux, and inflammatory bowel diseases.

The best documented anticancer effects of BRBs have been identified in oral, esophageal, and colorectal cancers, it is clear that cancer can act systemically affecting the function of other organs and resulting in metastasis. While research has started to focus on investigating the chemopreventative roles of BRBs in breast (Ravoori, Vadhanam, Aqil, & Gupta, 2012) and prostate cancer (Eskra, Schlicht, & Bosland, 2019), future studies should focus on the impact of BRBs and their metabolites on (1) the systemic implications of cancer, (2) the ability of polyphenols to prevent metastasis at secondary sites, (3) the impact of BRBs on liver-intestinal cross-talk due to the roles of the liver in

metabolism, and ultimately (4) the impact of BRB polyphenols on cancer stem cell or stem-cell like populations.

## ACKNOWLEDGMENTS

The authors recognize the assistance of Sarah Dwyer, Adam Higgins, Anna Powell, and Katie Stott for proof reading manuscript. The authors would like to thank the following organizations who have funded their positions during their research into diet and cancer: Tenovus Cancer Care (PhD2013/L25; LP&SM), the [World Cancer Research Fund](#) (2013/1006; LP), Cancer Research UK Programme grant (C1295/A15937-LP), and a philanthropic donation from Mr Lyndon and Mrs Shirley-Ann Wood from the Moorhouse Group Ltd (LP&SM).

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## AUTHOR CONTRIBUTIONS

SM was involved with the literature search, figure design, and drafting and critical revision of the manuscript. CP was involved with the literature search and drafting of the manuscript. LP was involved in drafting and critical revision of the manuscript.

## ROLE OF FUNDING SOURCE

Nothing to disclose

## ORCID

Connor Parry BSc  <https://orcid.org/0000-0002-4467-9196>

## REFERENCES

- Allis, C. D., & Jenuwein, T. (2016). The molecular hallmarks of epigenetic control. *Nature Reviews in Genetics*, 17(8), 487–500.
- Baquero, F., & Nombela, C. (2012). The microbiome as a human organ. *Clinical Microbiology and Infection*, 18, 2–4.
- Barker, N. (2014). Adult intestinal stem cells: Critical drivers of epithelial homeostasis and regeneration. *Nature Reviews Molecular Cell Biology*, 15, 19–33.
- Bi, X., Fang, W., Wang, L.-S., Stoner, G. D., & Yang, W. (2010). Black raspberries inhibit intestinal tumorigenesis in Apc1638<sup>+/−</sup> and Muc2<sup>−/−</sup> mouse models of colorectal cancer. *Cancer Prevention Research*, 3, 1443–1450.
- Blot, W. J., & Tarone, R. E. (2018). Esophageal cancer. In M. J. Thun, M. S. Linet, J. R. Cerhan, C. A. Haiman, & D. Schottenfeld (Eds.), *Cancer epidemiology and prevention* (4 ed., pp. 579–592). New York, NY: Oxford University Press.
- Bravo, L. (1998). Polyphenols: Chemistry, dietary sources, metabolism, and nutritional significance. *Nutrition Reviews*, 56, 317–333.
- Bray, F., Ferlay, J., Soerjomataram, I., Siegel, R. L., Torre, L. A., & Jemal, A. (2018). Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: A Cancer Journal for Clinicians*, 68(6), 394–424.
- Brown, K. F., Rumgay, H., Dunlop, C., Ryan, M., Quartly, F., Cox, A., ... Parkin, D. M. (2018). The fraction of cancer attributable to modifiable risk factors in England, Wales, Scotland, Northern Ireland, and the United Kingdom in 2015. *British Journal of Cancer*, 118(8), 1130–1141.
- Bullman, S., Pedamallu, C. S., Sicinska, E., Clancy, T. E., Zhang, X., Cai, D., ... Meyerson, M. (2017). Analysis of Fusobacterium persistence and antibiotic response in colorectal cancer. *Science (New York, N.Y.)*, 358(6369), 1443–1448.

- Chen, H.-M., Yu, Y.-N., Wang, J.-L., Lin, Y.-W., Kong, X., Yang, C.-Q., & Fang, J. Y. (2013). Decreased dietary fiber intake and structural alteration of gut microbiota in patients with advanced colorectal adenoma. *The American Journal of Clinical Nutrition*, 97(5), 1044–1052.
- Chen, K.-M., Guttenplan, J. B., Sun, Y.-W., Cooper, T., Shalaby, N. A. E., Kosinska, W., ... El-Bayoumy, K. (2018). Effects of black raspberry on dibenzo[a,h]pyrene diol epoxide induced DNA adducts, mutagenesis, and tumorigenesis in the mouse oral cavity. *Cancer Prevention Research*, 11(3), 157.
- Chen, K.-M., Sun, Y.-W., Kawasawa, Y. I., Salzberg, A. C., Zhu, J., Gowda, K., ... El-Bayoumy, K. (2020). Black raspberry inhibits oral tumors in mice treated with the tobacco smoke constituent dibenzo(def,p)chrysene via genetic and epigenetic alterations. *Cancer Prevention Research*, 13(4), 357–366. doi: [10.1158/1940-6207.CANP-19-0496](https://doi.org/10.1158/1940-6207.CANP-19-0496).
- Chen, K.-M., Guttenplan, J. B., Zhang, S.-M., Aliaga, C., Cooper, T. K., Sun, Y.-W., ... El-Bayoumy, K. (2013). Mechanisms of oral carcinogenesis induced by dibenzo[a,h]pyrene: An environmental pollutant and a tobacco smoke constituent. *International Journal of Cancer*, 133(6), 1300–1309.
- Chen, L., Jiang, B., Zhong, C., Guo, J., Zhang, L., Mu, T., ... Bi, X. (2018). Chemoprevention of colorectal cancer by black raspberry anthocyanins involved the modulation of gut microbiota and SFRP2 demethylation. *Carcinogenesis*, 39(3), 471–481.
- Chen, T., Rose, M. E., Hwang, H., Nines, R. G., & Stoner, G. D. (2006). Black raspberries inhibit N-nitrosomethylbenzylamine (NMBA)-induced angiogenesis in rat esophagus parallel to the suppression of COX-2 and iNOS. *Carcinogenesis*, 27(11), 2301–2307.
- Chen, T., Shi, N., & Afzali, A. (2019). Chemopreventive effects of strawberry and black raspberry on colorectal cancer in inflammatory bowel disease. *Nutrients*, 11(6), 1261.
- Choi, J.-S., Kang, S.-W., Li, J., Kim, J. L., Bae, J.-Y., Kim, D. S., & Kang, Y.-H. (2009). Blockade of oxidized LDL-triggered endothelial apoptosis by quercetin and rutin through differential signaling pathways involving JAK2. *Journal of Agricultural and Food Chemistry*, 57(5), 2079–2086.
- Coca, S., Perez-Piqueras, J., Martinez, D., Colmenarejo, A., Saez, M. A., Vallejo, C., ... Moreno, M. (1997). The prognostic significance of intratumoral natural killer cells in patients with colorectal carcinoma. *Cancer*, 79(12), 2320–2328.
- Cordenunsi, B. R., Oliveira do Nascimento, J. R., Genovese, M. I., & Lajolo, F. M. (2002). Influence of cultivar on quality parameters and chemical composition of strawberry fruits grown in Brazil. *Journal of Agricultural and Food Chemistry*, 50(9), 2581–2586.
- Corona, G., Tanga, F., Vauzoura, D., Rodriguez-Mateosaand, A., & Spencer, J. P. E. (2011). Assessment of the anthocyanidin content of common fruits and development of a test diet rich in a range of anthocyanins. *Journal of Berry Research*, 1, 209–216.
- Costello, J. F., Frühwald, M. C., Smiraglia, D. J., Rush, L. J., Robertson, G. P., Gao, X., ... Plass, C. (2000). Aberrant CpG-island methylation has non-random and tumour-type-specific patterns. *Nature Genetics*, 24(2), 132–138.
- Deighton, N., Brennan, R., & Davies, H. V. (2000). Antioxidant properties of domesticated and wild *Rubus* species. *Journal of the Science of Food and Agriculture*, 80, 1307–1313.
- Dejea, C. M., Fathi, P., Craig, J. M., Boleij, A., Taddese, R., Geis, A. L., ... Sears, C. L. (2018). Patients with familial adenomatous polyposis harbor colonic biofilms containing tumorigenic bacteria. *Science*, 359(6375), 592–597.
- Denev, P. N., Kratchanov, C. G., Ciz, M., Lojek, A., & Kratchanova, M. G. (2012). Bioavailability and antioxidant activity of black chokeberry (*Aronia melanocarpa*) polyphenols: In vitro and in vivo evidences and possible mechanisms of action: A review. *Comprehensive Reviews in Food Science and Food Safety*, 11(5), 471–489.
- Dethlefsen, L., & Relman, D. A. (2011). Incomplete recovery and individualized responses of the human distal gut microbiota to repeated antibiotic perturbation. *Proceedings of the National Academy of Sciences of the United States of America*, 108(Suppl 1), 4554–4561.
- Ding, X., Hiraku, Y., Ma, N., Kato, T., Saito, K., Nagahama, M., ... Kawanishi, S. (2005). Inducible nitric oxide synthase-dependent DNA damage in mouse model of inflammatory bowel disease. *Cancer Science*, 96(3), 157–163.
- Diplock, A. T., Charleux, J. L., Crozier-Willi, G., Kok, F. J., Rice-Evans, C., Roberfroid, M., ... Viña-Ribes, J. (1998). Functional food science and defence against reactive oxidative species. *British Journal of Nutrition*, 80, S77–S112.
- Dyall, S. C. (2015). Long-chain omega-3 fatty acids and the brain: A review of the independent and shared effects of, E. P. A., DPA and DHA. *Frontiers in Aging Neuroscience*, 7, 52–52.
- Eberhart, C. E., Coffey, R. J., Radhika, A., Giardiello, F. M., Ferrenbach, S., & DuBois, R. N. (1994). Up-regulation of cyclooxygenase 2 gene expression in human colorectal adenomas and adenocarcinomas. *Gastroenterology*, 107(4), 1183–1188.
- El-Bayoumy, K., Chen, K.-M., Zhang, S.-M., Sun, Y.-W., Amin, S., Stoner, G., ... Guttenplan, J. B. (2017). Carcinogenesis of the oral cavity: Environmental causes and potential prevention by black raspberry. *Chemical Research in Toxicology*, 30(1), 126–144.
- Esakra, J. N., Schlicht, M. J., & Bosland, M. C. (2019). Effects of black raspberries and their ellagic acid and anthocyanin constituents on taxane chemotherapy of castration-resistant prostate cancer cells. *Scientific Reports*, 9(1), 4367.
- Ferlay, J., Soerjomataram, I., Dikshit, R., Eser, S., Mathers, C., Rebelo, M., & Bray, P. (2015). Cancer incidence and mortality worldwide: Sources, methods and major patterns in GLOBOCAN 2012. *International Journal of Cancer*, 136(5), E359–E86.
- Fodde, R., Edelmann, W., Yang, K., van Leeuwen, C., Carlson, C., Renault, B., ... Khan, P. M. (1994). A targeted chain-termination mutation in the mouse *Apc* gene results in multiple intestinal tumors. *Proceedings of the National Academy of Sciences of the United States of America*, 91(19), 8969–8973.
- Furusawa, Y., Obata, Y., Fukuda, S., Endo, T. A., Nakato, G., Takahashi, D., ... Ohno, H. (2013). Commensal microbe-derived butyrate induces the differentiation of colonic regulatory T cells. *Nature*, 504(7480), 446–450.
- González, N., Prieto, I., Del Puerto-Nevado, L., Portal-Nuñez, S., Ardura, J. A., Corton, M., ... Ortiz, A. (2017). 2017 update on the relationship between diabetes and colorectal cancer: Epidemiology, potential molecular mechanisms and therapeutic implications. *Oncotarget*, 8(11), 18456–18485.
- Gu, J., Thomas-Ahner, J. M., Riedl, K. M., Bailey, M. T., Vodovotz, Y., Schwartz, S. J., ... Clinton, S. K. (2019). Dietary black raspberries impact the colonic microbiome and phytochemical metabolites in mice. *Molecular Nutrition & Food Research*, 63(8), 1800636.
- Guttenplan, J. B., Kosinska, W., Zhao, Z.-L., Chen, K.-M., Aliaga, C., DelTondo, J., ... El-Bayoumy, K. (2012). Mutagenesis and carcinogenesis induced by dibenzo[a,h]pyrene in the mouse oral cavity: A potential new model for oral cancer. *International Journal of Cancer*, 130(12), 2783–2790.
- Hague, A., Manning, A. M., Hanlon, K. A., Hart, D., Paraskeva, C., & Huschtscha, L. I. (1993). Sodium butyrate induces apoptosis in human colonic tumour cell lines in a p53-independent pathway: Implications for the possible role of dietary fibre in the prevention of large-bowel cancer. *International Journal of Cancer*, 55(3), 498–505.
- Hanahan, D., & Weinberg Robert, A. (2011). Hallmarks of cancer: The next generation. *Cell*, 144(5), 646–674.
- Harris, G. K., Gupta, A., Nines, R. G., Kresty, L. A., Habib, S. G., Frankel, W. L., ... Stoner, G. D. (2001). Effects of lyophilized black raspberries on azoxymethane-induced colon cancer and 8-hydroxy-2'-deoxyguanosine levels in the Fischer 344 rat. *Nutrition and Cancer*, 40, 125–133.
- Hashibe, M., Brennan, P., Chuang, S.-C., Boccia, S., Castellsague, X., Chen, C., ... Boffetta, P. (2009). Interaction between tobacco and alcohol use and the risk of head and neck cancer: Pooled analysis in the International Head and Neck Cancer Epidemiology Consortium. *Cancer Epidemiol Biomarkers Prev*, 18(2), 541–550.
- Hu, S., Liu, L., Chang, E. B., Wang, J.-Y., & Raufman, J.-P. (2015). Butyrate inhibits pro-proliferative miR-92a by diminishing c-Myc-induced

- miR-17-92a cluster transcription in human colon cancer cells. *Molecular Cancer*, 14, 180–180.
- Huang, C., Huang, Y., Li, J., Hu, W., Aziz, R., Tang, M.-S., & Stoner, G. D. (2002). Inhibition of Benzo(a)pyrene diol-epoxide-induced transactivation of activated protein 1 and nuclear factor  $\kappa$ B by black raspberry extracts. *Cancer Research*, 62(23), 6857–6863.
- Huang, Y.-W., Gu, F., Dombkowski, A., Wang, L.-S., & Stoner, G. D. (2016). Black raspberries demethylate Sfrp4, a WNT pathway antagonist, in rat esophageal squamous cell papilloma. *Molecular Carcinogenesis*, 55(11), 1867–1875.
- Jo, Y.-H., Park, H.-C., Choi, S., Kim, S., Bao, C., Kim, H. W., ... Auh, J.-H. (2015). Metabolomic analysis reveals cyanidins in black raspberry as candidates for suppression of lipopolysaccharide-induced inflammation in murine macrophages. *Journal of Agricultural and Food Chemistry*, 63(22), 5449–5458.
- Johnson, J. (2009). Effect of black raspberry extracts on colon cancer cell proliferation. [Master's thesis]. [Columbus (OH)]. The Ohio State University.
- Kalt, W. (2005). Effects of production and processing factors on major fruit and vegetable antioxidants. *Journal of Food Science*, 70(1), R11–R19.
- Knobloch, T. J., Ryan, N. M., Bruschweiler-Li, L., Wang, C., Bernier, M. C., Somogyi, A., ... Oghumu, S. (2019). Metabolic regulation of glycolysis and AMP activated protein kinase pathways during black raspberry-mediated oral cancer chemoprevention. *Metabolites*, 9(7), 140.
- Knobloch, T. J., Uhrig, L. K., Pearl, D. K., Casto, B. C., Warner, B. M., Clinton, S. K., ... Weghorst, C. M. (2016). Suppression of proinflammatory and pro-survival biomarkers in oral cancer patients consuming a black raspberry phytochemical-rich troche. *Cancer Prevention Research (Philadelphia, PA)*, 9(2), 159–171.
- Kong, J., Chia, L., Goh, N., Chia, T., & Brouillard, R. (2003). Analysis and biological activities of anthocyanins. *Phytochemistry*, 64, 933.
- Kresty, L. A., Frankel, W. L., Hammond, C. D., Baird, M. E., Mele, J. M., Stoner, G. D., ... Fromkes, J. J. (2006). Transitioning from preclinical to clinical chemopreventive assessments of lyophilized black raspberries: Interim results show berries modulate markers of oxidative stress in Barrett's esophagus patients. *Nutrition and Cancer*, 54(1), 148–156.
- Kresty, L. A., Fromkes, J. J., Frankel, W. L., Hammond, C. D., Seeram, N. P., Baird, M., ... Stoner, G. D. (2018). A phase I pilot study evaluating the beneficial effects of black raspberries in patients with Barrett's esophagus. *Oncotarget*, 9, 35356–35372.
- Kresty, L. A., Mallery, S. R., & Stoner, G. D. (2016). Black raspberries in cancer clinical trials: Past, present and future. *Journal of Berry Research*, 6(2), 251–261.
- Kresty, L. A., Morse, M. A., Morgan, C., Carlton, P. S., Lu, J., Gupta, A., ... Stoner, G. D. (2001). Chemoprevention of esophageal tumorigenesis by dietary administration of lyophilized black raspberries. *Cancer Research*, 61(16), 6112.
- Lasry, A., Zinger, A., & Ben-Neriah, Y. (2016). Inflammatory networks underlying colorectal cancer. *Nature Immunology*, 17(3), 230–240.
- Lechner, J. F., Reen, R. K., Dombkowski, A. A., Cukovic, D., Salagrama, S., Wang, L. S., ... Stoner, G. D. (2008). Effects of a black raspberry diet on gene expression in the rat esophagus. *Nutrition Cancer*, 1, 61–69.
- Lee, J.-H., Oh, M., Seok, J. H., Kim, S., Lee, D. B., Bae, G., ... Kim, K. H. (2016). Antiviral effects of black raspberry (*Rubus coreanus*) seed and its gallic acid against Influenza virus infection. *Viruses*, 8, 157.
- Ley, R. E., Bäckhed, F., Turnbaugh, P., Lozupone, C. A., Knight, R. D., & Gordon, J. I. (2005). Obesity alters gut microbial ecology. *Proceedings of the National Academy of Sciences of the United States of America*, 102(31), 11070–11075.
- Luo, X., Fang, S., Xiao, Y., Song, F., Zou, T., Wang, M., ... Ling, W. (2012). Cyanidin-3-glucoside suppresses TNF- $\alpha$ -induced cell proliferation through the repression of Nox activator 1 in mouse vascular smooth muscle cells: Involvement of the STAT3 signaling. *Molecular and Cellular Biochemistry*, 362(1), 211–218.
- Määttä-Riihinen, K. R., Kamal-Eldin, A., Mattila, P. H., González-Paramás, A. M., & Törrönen, A. R. (2004). Distribution and contents of phenolic compounds in eighteen Scandinavian berry species. *Journal of Agricultural and Food Chemistry*, 52(14), 4477–4486.
- Mallery, S. R., Tong, M., Shumway, B. S., Curran, A. E., Larsen, P. E., Ness, G. M., ... Stoner, G. D. (2014). Topical application of a mucoadhesive freeze-dried black raspberry gel induces clinical and histologic regression and reduces loss of heterozygosity events in premalignant oral intraepithelial lesions: Results from a multi-centered, placebo controlled clinical trial. *Clinical Cancer Research*, 20(7), 1910–1924.
- Mallery, S. R., Zwick, J. C., Pei, P., Tong, M., Larsen, P. E., Shumway, B. S., ... Stoner, G. D. (2008). Topical application of a bioadhesive black raspberry gel modulates gene expression and reduces cyclooxygenase 2 protein in human premalignant oral lesions. *Cancer Research*, 68(12), 4945–4957.
- Manach, C., Scalbert, A., Morand, C., Remesy, C., & Jimenez, L. (2004). Polyphenols: Food sources and bioavailability. *The American Journal of Clinical Nutrition*, 79, 727–747.
- Maslowski, K. M., Vieira, A. T., Ng, A., Kranich, J., Sierro, F., Yu, D., ... Mackay, C. R. (2009). Regulation of inflammatory responses by gut microbiota and chemoattractant receptor GPR43. *Nature*, 461(7268), 1282–1286.
- May, S., McDermott, G., Marchesi, J. R., & Parry, L. (2020). Impact of black raspberries on the normal and malignant Apc deficient murine gut microbiome. *Journal of Berry Research*, 10, 61–76.
- Mazza, G. (2007). Anthocyanins and heart health. *Annali dell'Istituto Superiore di Sanità* 43, 369–374.
- McGhie, T. K., Hall, H. K., Ainge, G. D., & Mowat, A. D. (2002). Breeding rubus cultivars for high anthocyanin content and high antioxidant capacity. *Acta Horticulture*, 585, 495–500.
- Medda, R., Lyros, O., Schmidt, J. L., Jovanovic, N., Nie, L., Link, B. J., ... Rafiee, P. (2015). Anti-inflammatory and anti-angiogenic effect of black raspberry extract on human esophageal and intestinal microvascular endothelial cells. *Microvascular Research*, 97, 167–180.
- Middleton, J. E., Kandaswami, C., & Theoharides, T. C. (2000). The effects of plant flavonoids on mammalian cells: Implications for inflammation, heart disease, and cancer. *Pharmacological Reviews*, 52, 673–751.
- Montgomery, M. K., Osborne, B., Brown, S. H. J., Small, L., Mitchell, T. W., Cooney, G. J., & Turner, N. (2013). Contrasting metabolic effects of medium-versus long-chain fatty acids in skeletal muscle. *Journal of Lipid Research*, 54(12), 3322–3333.
- Montrose, D. C., Horelik, N. A., Madigan, J. P., Stoner, G. D., Wang, L.-S., Bruno, R. S., ... Rosenberg, D. W. (2011). Anti-inflammatory effects of freeze-dried black raspberry powder in ulcerative colitis. *Carcinogenesis*, 32(3), 343–350.
- Mosley, D., Su, T., Murff, H. J., Smalley, W. E., Ness, R. M., Zheng, W., & Shrubsole, M. J. (2020). Meat intake, meat cooking methods, and meat-derived mutagen exposure and risk of sessile serrated lesions. *The American Journal of Clinical Nutrition*, 111, 1244–1251.
- Moyer, R. A., Hummer, K. E., Finn, C. E., Frei, B., & Wrolstad, R. E. (2002). Anthocyanins, phenolics, and antioxidant capacity in diverse small fruits: Vaccinium, rubus, and ribes. *Journal of Agricultural and Food Chemistry*, 50(3), 519–525.
- Nijveldt, R. J., van Nood, E., van Hoorn, D. E., Boelens, P. G., van Norren, K., & van Leeuwen, P. A. (2001). Flavonoids: A review of probable mechanisms of action and potential applications. *The American Journal of Clinical Nutrition*, 74, 418–425.
- Nishisho, I., Nakamura, Y., Miyoshi, Y., Miki, Y., Ando, H., Horii, A., ... Hedge, P. (1991). Mutations of chromosome 5q21 genes in FAP and colorectal cancer patients. *Science*, 253(5020), 665–669.
- Ogawa, K., Sakakibara, H., Iwata, R., Ishii, T., Sato, T., Goda, T., ... Kumazawa, S. (2008). Anthocyanin composition and antioxidant activity of the crowberry (*Empetrum nigrum*) and other berries. *Journal of Agricultural and Food Chemistry*, 56(12), 4457–4462.
- Pan, P., Kang, S., Wang, Y., Liu, K., Oshima, K., Huang, Y.-W., ... Wang, L.-S. (2017). Black raspberries enhance natural killer cell infiltration into

- the colon and suppress the progression of colorectal cancer. *Frontiers in Immunology*, 8, 997.
- Pan, P., Lam, V., Salzman, N., Huang, Y.-W., Yu, J., Zhang, J., ... Wang, L. S. (2017). Black raspberries and their anthocyanin and fiber fractions alter the composition and diversity of gut microbiota in F-344 rats. *Nutrition and Cancer*, 69(6), 943–951.
- Pan, P., Oshima, K., Huang, Y.-W., Yearsley, M., Zhang, J., Arnold, M., ... Wang, L.-S. (2018). Gut bacteria are required for the benefits of black raspberries in Apc (Min/+) mice. *Journal of Berry Research*, 8(4), 239–249.
- Pan, P., Skaer, C. W., Wang, H.-T., Kreiser, M. A., Stirdivant, S. M., Oshima, K., ... Wang, L.-S. (2017). Systemic metabolite changes in wild-type C57BL/6 mice fed black raspberries. *Nutrition and Cancer*, 69(2), 299–306.
- Pandey, K. B., & Ibrahim, S. R. (2009). Plant polyphenols as dietary antioxidants in human health and disease. *Oxidative Medicine and Cellular Longevity*, 2, 270–278.
- Park, J. Y., Mitrou, P. N., Luben, R., Khaw, K.-T., & Bingham, S. A. (2009). Is bowel habit linked to colorectal cancer? – Results from the EPIC-Norfolk study. *European Journal of Cancer*, 45(1), 139–145.
- Paul, B., Barnes, S., Demark-Wahnefried, W., Morrow, C., Salvador, C., Skibola, C., ... Tollefsbol, T. O. (2015). Influences of diet and the gut microbiome on epigenetic modulation in cancer and other diseases. *Clinical Epigenetics*, 7, 112.
- Peiffer, D. S., Wang, L.-S., Zimmerman, N. P., Ransom, B. W. S., Carmella, S. G., Kuo, C.-T., ... Stoner, G. D. (2016). Dietary consumption of black raspberries or their anthocyanin constituents alters innate immune cell trafficking in esophageal cancer. *Cancer Immunology Research*, 4(1), 72.
- Peiffer, D. S., Zimmerman, N. P., Wang, L.-S., Ransom, B. W. S., Carmella, S. G., Kuo, C.-T., ... Stoner, G. D. (2014). Chemoprevention of esophageal cancer with black raspberries, their component anthocyanins, and a major anthocyanin metabolite, protocatechuic acid. *Cancer Prevention Research*, 7(6), 574.
- Peng, L., Li, Z.-R., Green, R. S., Holzman, I. R., & Lin, J. (2009). Butyrate enhances the intestinal barrier by facilitating tight junction assembly via activation of AMP-activated protein kinase in Caco-2 cell monolayers. *The Journal of Nutrition*, 139(9), 1619–1625.
- Prior, R. L., Cao, G., Martin, A., Sofic, E., McEwen, J., O'Brien, C., ... Mainland, M. (1998). Antioxidant capacity as influenced by total phenolic and anthocyanin content, maturity, and variety of vaccinium species. *Journal of Agricultural and Food Chemistry*, 46(7), 2686–2693.
- Qin, J., Li, Y., Cai, Z., Li, S., Zhu, J., Zhang, F., ... Wang, J. (2012). A metagenome-wide association study of gut microbiota in type 2 diabetes. *Nature*, 490, 55.
- Ramos-Nino, M. E. (2013). The role of chronic inflammation in obesity-associated cancers. *ISRN Oncology*, 2013, 697521–697521.
- Ravoori, S., Vadhanam, M. V., Aqil, F., & Gupta, R. C. (2012). Inhibition of estrogen-mediated mammary tumorigenesis by blueberry and black raspberry. *Journal of Agricultural and Food Chemistry*, 60(22), 5547–5555.
- Rodrigo, K. A., Rawal, Y., Renner, R. J., Schwartz, S. J., Tian, Q., Larsen, P. E., ... Mallery, S. R. (2006). Suppression of the tumorigenic phenotype in human oral squamous cell carcinoma cells by an ethanol extract derived from freeze-dried black raspberries. *Nutrition and Cancer*, 54(1), 58–68.
- Said, H. M. (2011). Intestinal absorption of water-soluble vitamins in health and disease. *Biochemistry Journal*, 437(3), 357–372.
- Sánchez-Lara, K., Turcott, J. G., Juárez-Hernández, E., Nuñez-Valencia, C., Villanueva, G., Guevara, P., ... Arrieta, O. (2014). Effects of an oral nutritional supplement containing eicosapentaenoic acid on nutritional and clinical outcomes in patients with advanced non-small cell lung cancer: Randomised trial. *Clinical Nutrition*, 33(6), 1017–1023.
- Seeram, N. P. (2008). Berry fruits for cancer prevention: Current status and future prospects. *Journal of Agricultural and Food Chemistry*, 56, 630–635.
- Serra, D., Paixão, J., Nunes, C., Dinis, T. C. P., & Almeida, L. M. (2013). Cyanidin-3-glucoside suppresses cytokine-induced inflammatory response in human intestinal cells: Comparison with 5-aminosalicylic acid. *PLoS One*, 8(9), e73001–e01.
- Sheflin, A. M., Whitney, A. K., & Weir, T. L. (2014). Cancer-promoting effects of microbial dysbiosis. *Current Oncology Reports*, 16(10), 406–406.
- Shi, N., Chen, F., Zhang, X., Clinton, S. K., Tang, X., Sun, Z., ... Chen, T. (2017). Suppression of oxidative stress and NF- $\kappa$ B/MAPK signaling by lyophilized black raspberries for esophageal cancer prevention in rats. *Nutrients*, 9(4), 413.
- Shin, J.-S., Cho, E.-J., Choi, H.-E., Seo, J.-H., An, H.-J., Park, H.-J., ... Leeab, K.-T. (2014). Anti-inflammatory effect of a standardized triterpenoid-rich fraction isolated from *Rubus coreanus* on dextran sodium sulfate-induced acute colitis in mice and LPS-induced macrophages. *Journal of Ethnopharmacology*, 158, 291–300.
- Shumway, B. S., Kresty, L. A., Larsen, P. E., Zwick, J. C., Lu, B., Fields, H. W., ... Mallery, S. R. (2008). Effects of a topically applied bioadhesive berry gel on loss of heterozygosity indices in premalignant oral lesions. *Clinical Cancer Research*, 14(8), 2421–2430.
- Siriwoharn, T., Wrolstad, R. E., Finn, C. E., & Pereira, C. B. (2004). Influence of cultivar, maturity, and sampling on blackberry (*Rubus L. Hybrids*) anthocyanins, polyphenolics, and antioxidant properties. *Journal of Agricultural and Food Chemistry*, 52(26), 8021–8030.
- Soobrattee, M. A., Bahorun, T., & Aruoma, O. I. (2006). Chemopreventive actions of polyphenolic compounds in cancer. *Biofactors*, 27, 19–35.
- Stevens, R. G., Swede, H., Heinen, C. D., Jablonski, M., Grupka, M., Ross, B., ... Levine, J. (2007). Aberrant crypt foci in patients with a positive family history of sporadic colorectal cancer. *Cancer Letters*, 248(2), 262–268.
- Stoner, G. D. (2009). Foodstuffs for preventing cancer: The preclinical and clinical development of berries. *Cancer Prevention Research*, 2(3), 187.
- Stoner, G. D., Chen, T., Kresty, L. A., Aziz, R. M., Reinemann, T., & Nines, R. (2006). Protection against esophageal cancer in rodents with lyophilized berries: Potential mechanisms. *Nutrition and Cancer*, 54(1), 33–46.
- Stoner, G. D., Wang, L.-S., Zikri, N., Chen, T. S., Hecht, S., Huang, C., ... Lechner, J. F. (2007). Cancer prevention with freeze-dried berries and berry components. *Seminars in Cancer Biology*, 17(5), 403–410.
- Takebe, T., Sekine, K., Enomura, M., Koike, H., Kimura, M., Ogaeri, T., ... Taniguchi, H. (2013). Vascularized and functional human liver from an iPSC-derived organ bud transplant. *Nature*, 499(7459), 481–484.
- Tanaka, Y., Ito, S., & Isobe, K.-i. (2016). Vancomycin-sensitive bacteria trigger development of colitis-associated colon cancer by attracting neutrophils. *Scientific Reports*, 6(1), 23920.
- Tian, Q., Giusti, M. M., Stoner, G. D., & Schwartz, S. J. (2006). Characterization of a new anthocyanin in black raspberries (*Rubus occidentalis*) by liquid chromatography electrospray ionization tandem mass spectrometry. *Food Chemistry*, 94, 465–468.
- Torre, L. A., Bray, F., Siegel, R. L., Ferlay, J., Lortet-Tieulent, J., & Jemal, A. (2015). Global cancer statistics, 2012. *CA: A Cancer Journal for Clinicians*, 65(2), 87–108.
- Tsuchida, A., Ohno, S., Wu, W., Borjigin, N., Fujita, K., Aoki, T., ... Takahashi, M. (2011). miR-92 is a key oncogenic component of the miR-17-92 cluster in colon cancer. *Cancer Science*, 102(12), 2264–2271.
- Tsujii, M., Kawano, S., & DuBois, R. N. (1997). Cyclooxygenase-2 expression in human colon cancer cells increases metastatic potential. *Proceedings of the National Academy of Sciences of the United States of America*, 94(7), 3336–3340.
- Tu, P., Bian, X., Chi, L., Gao, B., Ru, H., Knobloch, T. J., ... Lu, K. (2018). Characterization of the functional changes in mouse gut microbiome associated with increased akkermansia muciniphila population modulated by dietary black raspberries. *ACS Omega*, 3(9), 10927–10937.
- Tu, P., Bian, X., Chi, L., Xue, J., Gao, B., Lai, Y., ... Lu, K. (2020). Metabolite profiling of the gut microbiome in mice with dietary administration of black raspberries. *ACS Omega*, 5(3), 1318–1325.
- Turnbaugh, P. J., Bäckhed, F., Fulton, L., & Gordon, J. I. (2008). Diet-induced obesity is linked to marked but reversible alterations in the mouse distal gut microbiome. *Cell Host & Microbe*, 3(4), 213–223.
- Ursell, L. K., Metcalf, J. L., Parfrey, L. W., & Knight, R. (2012). Defining the human microbiome. *Nutrition Reviews*, 70(Suppl 1), S38–S44.

- Valko, M., Leibfritz, D., Moncol, J., Cronin, M. T., Mazur, M., & Telser, J. (2007). Free radicals and antioxidants in normal physiological functions and human disease. *The International Journal of Biochemistry & Cell Biology*, 39, 44–84.
- Velcich, A., Yang, W., Heyer, J., Fragale, A., Nicholas, C., Viani, S., ... Augenthaler, L. (2002). Colorectal cancer in mice genetically deficient in the mucin Muc2. *Science*, 295(5560), 1726.
- Vivier, E., Tomasello, E., Baratin, M., Walzer, T., & Ugolini, S. (2008). Functions of natural killer cells. *Nature Immunology*, 9(5), 503–510.
- Wada, L., & Ou, B. (2002). Antioxidant activity and phenolic content of *Oregon caneberries*. *Journal of Agricultural and Food Chemistry*, 50(12), 3495–3500.
- Waldecker, M., Kautenburger, T., Daumann, H., Busch, C., & Schrenk, D. (2008). Inhibition of histone-deacetylase activity by short-chain fatty acids and some polyphenol metabolites formed in the colon. *The Journal of Nutritional Biochemistry*, 19(9), 587–593.
- Wang, L.-S., Kuo, C.-T., Cho, S.-J., Seguin, C., Siddiqui, J., Stoner, K., ... Huang, Y.-W. (2013). Black raspberry-derived anthocyanins demethylate tumor suppressor genes through the inhibition of DNMT1 and DNMT3B in colon cancer cells. *Nutrition and Cancer*, 65(1), 118–125.
- Wang, L.-S., Kuo, C.-T., Huang, T. H. M., Yearsley, M., Oshima, K., Stoner, G. D., ... Huang, Y.-W. (2013). Black raspberries protectively regulate methylation of wnt pathway genes in precancerous colon tissue. *Cancer Prevention Research (Philadelphia, PA)*, 6(12), 1317–1327.
- Wang, L.-S., Kuo, C.-T., Stoner, K., Yearsley, M., Oshima, K., Yu, J., ... Huang, Y.-W. (2013). Dietary black raspberries modulate DNA methylation in dextran sodium sulfate (DSS)-induced ulcerative colitis. *Carcinogenesis*, 34(12), 2842–2850.
- Wang, L.-S., & Stoner, G. D. (2008). Anthocyanins and their role in cancer prevention. *Cancer Letters*, 269, 281–290.
- Wang, L.-S., Arnold, M., Huang, Y.-W., Sardo, C., Seguin, C., Martin, E., ... Stoner, G. (2011). Modulation of Genetic and epigenetic biomarkers of colorectal cancer in humans by black raspberries: A phase I pilot study. *Clinical Cancer Research*, 17(3), 598.
- Wang, L.-S., Burke, C. A., Hasson, H., Kuo, C.-T., Molmenti, C. L. S., Seguin, C., ... Stoner, G. D. (2014). A phase Ib study of the effects of black raspberries on rectal polyps in patients with familial adenomatous polyposis. *Cancer Prevention Research*, 7(7), 666.
- Wilson, M. R., Jiang, Y., Villalta, P. W., Stornetta, A., Boudreau, P. D., Carrá, A., ... Balskus, E. R. (2019). The human gut bacterial genotoxin colibactin alkylates DNA. *Science*, 363(6428), eaar7785.
- World Cancer Research Fund/American Institute for Cancer Research. (2018). Diet, nutrition, physical activity and cancer: A global perspective. continuous update project expert report. A summary of the Third Expert Report 2018. Retrieved from <https://www.wcrf.org/dietandcancer/resources-and-toolkit>
- Wu, S., Zhu, W., Thompson, P., & Hannun, Y. A. (2018). Evaluating intrinsic and non-intrinsic cancer risk factors. *Nature Communications*, 9(1), 3490.
- Wu, X., Gu, L., Prior, R. L., & McKay, S. (2004). Characterization of anthocyanins and proanthocyanidins in some cultivars of ribes, aronia, and sambucus and their antioxidant capacity. *Journal of Agricultural and Food Chemistry*, 52(26), 7846–7856.
- Wu, X., Pittman, H. E. III, Hager, T., Hager, A., Howard, L., & Prior, R. L. (2009). Phenolic acids in black raspberry and in the gastrointestinal tract of pigs following ingestion of black raspberry. *Molecular Nutrition & Food Research*, 53(S1), S76–S84.
- Xia, A., Zhang, Y., Xu, J., Yin, T., & Lu, X.-J. (2019). T cell dysfunction in cancer immunity and immunotherapy. *Frontiers in Immunology*, 10, 1719–1719.
- Yao, Q., Chen, Y., & Zhou, X. (2019). The roles of microRNAs in epigenetic regulation. *Current Opinion in Chemical Biology*, 51, 11–17.
- Zelenay, S., van der Veen, A. G., Böttcher, J. P., Snelgrove, K. J., Rogers, N., Acton, S. E., ... Reis e Sousa, C. (2015). Cyclooxygenase-dependent tumor growth through evasion of immunity. *Cell*, 162(6), 1257–1270.
- Zhang, H., Guo, J., Mao, L., Li, Q., Guo, M., Mu, T., ... Bi, X. (2019). Up-regulation of miR-24-1-5p is involved in the chemoprevention of colorectal cancer by black raspberry anthocyanins. *British Journal of Nutrition*, 122(5), 518–526.
- Zikri, N. N., Riedl, K. M., Wang, L.-S., Lechner, J. F., Schwartz, S. J., & Stoner, G. D. (2009). Black raspberry components inhibit proliferation, induce apoptosis and modulate gene expression in rat esophageal epithelial cells. *Nutrition and Cancer*, 61(6), 816–826.

**How to cite this article:** May S, Parry C, Parry L. Berry chemoprevention: do berries decrease the window of opportunity for tumorigenesis. *Food Frontiers*. 2020;1:260–275. <https://doi.org/10.1002/fft2.32>