Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/133523/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Soriano, L. F., Poole, H., Ingram, J. R. ORCID: <https://orcid.org/0000-0002-5257-1142> and Abbott, R. A. 2020. Effectiveness of a stop smoking service in reducing cigarette smoking in patients referred from an NHS dermatology centre. *British Journal of Dermatology* 183 (6) , pp. 1119-1121. 10.1111/bjd.19378 file

Publishers page: <http://dx.doi.org/10.1111/bjd.19378>
<<http://dx.doi.org/10.1111/bjd.19378>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.

Effectiveness of a smoking cessation service in reducing cigarette smoking in patients referred from an NHS dermatology centre

British Journal of Dermatology Research Letter

Effectiveness of a smoking cessation service in reducing cigarette smoking in patients referred from an NHS dermatology centre

Word Count: 750 words

Table Count: 0

Figure Count: 1

References: 9

LF Soriano¹, H Poole², JR Ingram³, RA Abbott¹

¹Department of Dermatology, University Hospital of Wales, Cardiff, UK

²Smoking Cessation Clinic, University Hospital of Wales, Cardiff, UK

³Division of Infection & Immunity, Cardiff University, Cardiff, UK

Corresponding author:

Livia Francine Soriano, Department of Dermatology, University Hospital of Wales, Cardiff, UK

Email: liviasoriano@doctors.org.uk

Funding sources: None.

Conflict of interest: JI is the editor of the British Journal of Dermatology.

ORCID

JI 0000-0002-5257-1142

Dear Editor,

Effectiveness of a smoking cessation service in reducing cigarette smoking in patients referred from an NHS dermatology centre

Cigarette smoking is the leading avoidable cause of mortality in the UK. Individuals who start smoking in early adult life lose 10 years of life expectancy.¹ In dermatology, smoking is risk factor for psoriasis (odds ratio (OR) of 1.94)² and hidradenitis suppurativa (OR 1.90).³ Relative to non-smokers, smokers have an increased relative risk (RR) of healing delay (RR 2.07)⁴, surgical site infections (RR 1.79)⁴, wound complications (RR 2.27)⁴, and melanoma (RR 1.7).⁵

The NHS has developed nationwide smoking cessation (SC) programmes, which involve an individual specifically trained in behavioural counselling and the knowledge of pharmacological interventions to reduce withdrawal symptoms to whom healthcare professional can refer their patients.⁶

We evaluated the effectiveness of an SC clinic in reducing cigarette smoking in patients referred from a tertiary dermatology centre. In this single-centre retrospective study, current smokers were identified in Cardiff dermatology clinics and, after obtaining permission, were referred to a structured hospital-based SC clinic between June 2017 and May 2019. Patients were treated with empathy, and the detrimental effects of smoking on physical appearance, skin ageing and their dermatological disease were emphasized.

The SC clinic arranged an appointment, which patients attended, declined or cancelled. Outcomes included stopping smoking, harm reduction (HR), or a single visit (SV) with SC advice. HR refers to an approach used when the patient is adamant he/she is not ready to quit; the patient is encouraged to reduce the number of cigarettes smoked to a minimum, setting a quit date in the near future. After the agreed quit date, patients were contacted at four weeks. Descriptive statistics were performed using Microsoft Excel 2013 (Microsoft Corp, Redmond, VA, USA).

Overall, 83 patients were referred by the dermatology team, of whom 41 (49.4%) were females and 42 (50.6%) males. Age range was 20-77, with a mean of 44.4 years. Dermatological diagnoses of patients are listed in Fig 1a. Sixty-one were referred to the SC clinic by email, three through postal service and 19 through clinic outcome proformas.

Effectiveness of a smoking cessation service in reducing cigarette smoking in patients referred from an NHS dermatology centre

36 patients had prior attempts at smoking cessation, 15 had no previous attempts and unknown in 32 patients. Twenty-four previously consulted another service: 5 consulted secondary care, 14 saw their GP, two contacted their pharmacist and 3 were seen by Stop Smoking Wales (SSW).

Thirty-one (37.3%) patients attended, 3 (3.6%) were referred to other services, 40 (48.2%) did not attend their appointments, and 9 (10.8%) cancelled their appointments (Fig 1b).

Reasons for non-attendance included transport issues (n=7), appointment not received (n=6), unwell (n=5), did not want to stop smoking (n=4), saw GP instead (n=2), preferred self-treatment (n=4), and unknown (n=21). Of the 49 patients who did not attend appointments, 4 (8%) had given up smoking on their own, 16 (33%) remained smokers and 29 (59%) were uncontactable. Of the 16 smokers, 7 decreased their smoking on their own, and 7 requested another appointment at the SC clinic.

Among the 31 patients who attended, 12 (14.5%) were treated smokers, 5 (16.1%) underwent HR, and 14 (45.2%) underwent SV. Patients were contacted at 4 weeks to assess their smoking status. Among treated smokers, 10 (83%) had a sustained abstinence, one failed to stop smoking, and one was lost to follow up. Among the patients who underwent SV and HR, 2 became nonsmokers, 2 reduced their cigarette consumption, 3 remained unchanged and 12 were lost to follow-up. Smoking cessation outcome was unavailable for the 3 patients referred to other services.

Dermatology patients reported being encouraged to attend SC clinics when clinicians used an empathic approach, explaining the benefits of SC to their skin disease.

Not all referrals via clinic outcome forms were carried out due to insufficient awareness of the referral pathway amongst clinic administrators, which explains the low numbers of patients referred via this route.

Our results show that SC following a referral from dermatology clinics was effective with a quit rate of 12% of patients referred. A recent dermatology study from Singapore⁸ demonstrated a quit rate of 15.8% with a pharmacist-led smoking intervention. A Cochrane review found that quit rates could be

Effectiveness of a smoking cessation service in reducing cigarette smoking in patients referred from an NHS dermatology centre

up to 18%–21%, depending on whether behavioural support was administered with pharmacotherapy.⁹

Limitations of our study include a small sample size and short follow-up period of four weeks.

Currently All Wales reports, including Tier 1 target data on smoking cessation, only report on four-week results. As this is the parameter used for all services, this was felt to be the most appropriate to use for our research. Ideally to ascertain long term success, 1 year follow up data would be collected.

In conclusion, our study suggests that dermatology clinics represent a good opportunity for clinicians to engage patients in the cessation or reduction of cigarette smoking. We encourage colleagues to prioritise smoking cessation. Emphasizing detrimental effects of smoking on physical appearance, skin ageing and dermatological diseases were associated with improved SC outcomes.

References:

1. Royal College of Physicians. Hiding in plain sight: treating tobacco dependency in the NHS [Internet]. 2018 [cited 2019 Aug 20];Available from: <https://www.rcplondon.ac.uk/projects/outputs/hiding-plain-sight-treating-tobacco-dependency-nhs>
2. Armstrong AW, Harskamp CT, Dhillon JS, et al. Psoriasis and smoking: A systematic review and meta-analysis. *Br J Dermatol* 2014; **170**: 304–14.
3. Garg A, Papagermanos V, Midura M, et al. Incidence of hidradenitis suppurativa among tobacco smokers: a population-based retrospective analysis in the U.S.A. *Br J Dermatol* 2018; **178**: 709–14.
4. Sørensen LT. Wound healing and infection in surgery: The clinical impact of smoking and smoking cessation: A systematic review and meta-analysis. *Arch Surg* 2012; **147**: 373–83.
5. Carter BD, Abnet CC, Feskanich D, et al. Smoking and Mortality — Beyond Established Causes. *N Engl J Med* 2015; **372**: 631–40.
6. Aubin HJ, Luquiens A, Berlin I. Pharmacotherapy for smoking cessation: Pharmacological

Effectiveness of a smoking cessation service in reducing cigarette smoking in patients referred from an NHS dermatology centre

- principles and clinical practice. *Br J Clin Pharmacol* 2014; **77**: 324–36.
- 7.
 8. Cheng HM, Liu WC, Chua G, et al. Impact of a pharmacy-led smoking cessation clinic in a dermatology centre. *Singapore Med J* 2019;
 9. Stead LF, Koilpillai P, Fanshawe TR, et al. Combined pharmacotherapy and behavioural interventions for smoking cessation. *Cochrane Database Syst Rev* 2016; **3**: CD008286.
 10. Fildes EE, Kapella-Mshigeni S, Campbell-Heider N. Outcomes of a One-Time Telephone Intervention for Smoking Cessation in Adults. *J Addict Nurs* 2015; **26**: 184–90.

Effectiveness of a smoking cessation service in reducing cigarette smoking in patients referred from an NHS dermatology centre

Figure 1. (a) The chart shows the range of dermatological diagnoses seen among patients, and (b) outcomes of patients referred to the smoking cessation clinic.