Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/134277/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Pu, Na, Yang, Qi, Shi, Xiao-Lei, Chen, Wei-Wei, Li, Xiao-Yao, Zhang, Guo-Fu, Li, Gang, Li, Bai-Qiang, Ke, Lu, Tong, Zhi-Hui, Cooper, David N. ORCID: <https://orcid.org/0000-0002-8943-8484>, Chen, Jian-Min, Li, Wei-Qin and Li, Jie-Shou 2020. Gene-environment interaction between APOA5 c.553G>T and pregnancy in hypertriglyceridemia-induced acute pancreatitis. Journal of Clinical Lipidology 14 (4) 10.1016/j.jacl.2020.05.003 file

Publishers page: <http://dx.doi.org/10.1016/j.jacl.2020.05.003>
<<http://dx.doi.org/10.1016/j.jacl.2020.05.003>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.

Gene-environment interaction between *APOA5* c.553G>T and pregnancy in hypertriglyceridemia-induced acute pancreatitis

Running title: *APOA5* c.553G>T and pregnancy in HTG-AP

Na Pu^{1,†}, BM, Qi Yang^{1,†,*}, PhD, Xiao-Lei Shi¹, BM, Wei-Wei Chen^{1,2}, MD, Xiao-Yao Li^{1,3}, PhD, Guo-Fu Zhang¹, BM, Gang Li¹, MD, Bai-Qiang Li¹, MD, Lu Ke¹, PhD, Zhi-Hui Tong¹, PhD, David N. Cooper⁴, PhD, Jian-Min Chen⁵, MD, PhD, Wei-Qin Li^{1,*}, PhD, Jie-Shou Li¹, PhD

¹Surgical Intensive Care Unit (SICU), Department of General Surgery, Jinling Hospital, Medical School of Nanjing University, Nanjing, China.

²Department of Gastroenterology, Clinical Medical College, Yangzhou University, Yangzhou, China.

³Department of Intensive Care Unit, The Affiliated Drum Tower Hospital, Medical School of Nanjing University, Nanjing, China.

⁴Institute of Medical Genetics, School of Medicine, Cardiff University, Cardiff, United Kingdom.

⁵EFS, Univ Brest, Inserm, UMR 1078, GGB, F-29200 Brest, France.

Na Pu, punayeah@163.com

Qi Yang, yangqi_nj@163.com

Xiao-Lei Shi, 15950561608@163.com

Wei-Wei Chen, cww1984@126.com

Xiao-Yao Li, lixiaoyaonju@163.com

Guo-Fu Zhang, njuzgf@163.com

Gang Li, nju8icu@126.com

Bai-Qiang Li, li_baiqiang@aliyun.com

Lu Ke, kkb9832@163.com

Zhi-Hui Tong, njzyantol@hotmail.com

28 David N. Cooper, CooperDN@cardiff.ac.uk

29 Jian-Min Chen, jian-min.chen@univ-brest.fr

30 Wei-Qin Li, njzy_pancrea@163.com

31 Jie-Shou Li, lijieshou2013@sohu.com

32 †Na Pu and Qi Yang contributed equally to this work.

33

34 ***Corresponding authors:**

35 Qi Yang, Department of General Surgery, Jinling Hospital, Nanjing University School of Medicine,

36 Nanjing 210002, China. E-Mail: yangqi_nj@163.com

37 Wei-Qin Li, Department of General Surgery, Jinling Hospital, Nanjing University School of Medicine,

38 Nanjing 210002, China. njzy_pancrea@163.com

39

40 **Highlights (maximum 85 characters including space, for each point)**

- 41 ➤ *APOA5* c.553G>T has been previously associated with altered triglyceride levels.
- 42 ➤ Here we report for the first time an association of *APOA5* c.553G>T with HTG-AP.
- 43 ➤ We provide evidence that *APOA5* c.553G>T interacts with pregnancy in causing HTG-AP.
- 44 ➤ Our findings provide novel insights into the complex etiology of HTG-AP.

45

Abstract (maximum 250 words)

BACKGROUND: The etiology of hypertriglyceridemia (HTG) and, consequently HTG-induced acute pancreatitis (HTG-AP), is complex.

OBJECTIVE: Herein, we explore a possible gene-environment interaction between *APOA5* c.553G>T (p.185Gly>Cys, rs2075291), a common variant associated with altered triglyceride levels, and pregnancy in HTG-AP.

METHODS: We enrolled 318 Han Chinese HTG-AP patients and divided them into three distinct groups: group 1, male patients (n =183); group 2, female patients whose disease was unrelated to pregnancy (n = 105); and group 3, female patients whose disease was related to pregnancy (n =30). *APOA5* rs2075291 genotype status was determined by Sanger sequencing. 362 healthy Han Chinese subjects were used as controls. Data on body mass index, peak triglyceride level, age of disease onset, episode number and clinical severity of HTG-AP were collected from each patient. Multiple comparisons, either between patient groups, between patient groups and controls, or within each patient group, were performed.

RESULTS: A robust association of *APOA5* rs2075291 with HTG-AP in general, and HTG-APIP in particular, was demonstrated. The minor T allele showed a stronger association with group 3 patients than with either group 1 or group 2 patients. This stronger association was due mainly to the much higher frequency of TT genotype in group 3 patients (20%) than that (<6%) in group 1 and group 2 patients. Moreover, the TT genotype was associated with a significantly higher peak triglyceride level in group 3 patients as compared to the GG genotype.

CONCLUSION: Our findings provide evidence for an interaction between *APOA5* rs2075291 and pregnancy in HTG-AP.

KEYWORDS: hypertriglyceridemia-induced acute pancreatitis (HTG-AP); acute pancreatitis in pregnancy (APIP); triglyceride; apolipoprotein A5; *APOA5* c.553G>T variant; gene-environment interaction

Introduction

Acute pancreatitis (AP) occurs with an annual incidence of 4.9–73.4 per 100,000 individuals worldwide.¹ Its incidence is increasing in recent years and its mean mortality rate has reached 2%.² Although biliary diseases, excessive alcohol consumption and hypertriglyceridemia (HTG) are generally thought to constitute the three leading etiologies of AP worldwide,^{3,4} in China more cases have been reported to be caused by HTG than by alcohol abuse.⁵⁻⁷ Irrespective of which population is studied, HTG is often associated with more severe disease than other etiological factors.^{4,8,9}

The etiology of HTG can be broadly divided into two categories, primary and secondary. Primary factors refer to genetic defects that cause or predispose to HTG whereas secondary etiological factors include obesity, alcohol abuse, diabetes mellitus and chronic renal failure.^{10,11} Nonetheless, in most cases, the etiology of HTG is complex.¹² Thus, for example, Dron and colleagues have recently shown that even severe HTG is primarily polygenic.¹³ Moreover, it is increasingly appreciated that there is an interplay between primary and secondary etiological factors in causing severe HTG.^{14,15}

Pregnancy is a physiological state that is normally associated with a 2- to 4-fold increase in serum triglyceride (TG) levels in late gestation.¹⁶ This increase is well tolerated by most women with normal baseline TG levels but may render those with genetic defects in TG metabolism prone to severe HTG and consequently HTG-induced AP (HTG-AP). In this regard, numerous studies have reported the identification of rare pathogenic variants in the lipoprotein lipase (*LPL*; OMIM# 609708),¹⁷⁻²⁶ glycosylphosphatidylinositol-anchored high-density lipoprotein binding protein 1 (*GPIHBP1*; OMIM# 612757) and apolipoprotein C2 (*APOC2*; OMIM# 608083) genes^{24,26} in patients experiencing HTG-AP during pregnancy (HTG-APIP). *LPL*, *GPIHBP1* and *APOC2*, together with *APOA5* (apolipoprotein A5; OMIM# 606368) and *LMF1* (lipase maturation factor 1; OMIM# 611761), constitute the five primary TG-related genes.^{12,27-29} Specifically, *LPL* plays an essential role in TG metabolism; *APOC2* and *APOA5* are essential *LPL* activators; *LMF1* is involved in the folding and expression of *LPL*; *GPIHBP1* mediates the transmembrane transport and binding of *LPL*.

By contrast, data on the possible interactions between pregnancy and common genetic risk factors in any of the five primary TG-related genes in HTG-AP have so far been scarce. We are aware of only

one such study, involving a single-nucleotide polymorphism in the *APOA5* gene, c.553G>T (p.Gly185>Cys; rs2075291), which was found in three (all homozygotes) of five Chinese patients with HTG-AP.²⁴ *APOA5* c.553G>T is common in the Chinese population [minor allele frequency (MAF) = ~4%^{30, 31}] and other Asian populations [e.g., MAF of 6.8% in East Asians according to gnomAD (<https://gnomad.broadinstitute.org/>)], but it is very rare in European populations (MAF of 0.03% according to gnomAD). Importantly, the minor T allele of *APOA5* rs2075291 has been firmly established as being associated with altered TG levels in normal controls and/or HTG patients in different Asian populations;³⁰⁻⁴³ and its functionality and pathogenicity have been supported by different lines of experimental evidence.⁴⁴⁻⁴⁷

Data on the role of *APOA5* c.553G>T in HTG-AP unrelated to pregnancy are also scarce; this is reflected by the very limited number of patients so far analyzed for this variant [one in Arai et al. (2014)⁴⁸ and 11 in Chen et al. (2017)⁴⁹]. Given the high frequency of this variant in the Chinese population and its direct role in altering TG levels, we sought to explore its potential interaction with pregnancy in HTG-AP in particular, and its relationship with the disease in general, using three well-defined Chinese patient cohorts.

Patients and Methods

Ethical compliance

This study was approved by the Ethics Committee of Jinling Hospital. Informed consent was obtained from all participating patients. Blood samples were obtained from the Biobank of Acute Pancreatitis in Jinling Hospital with the approval of the Biobank's Management Committee.

Study subjects

318 HTG-AP patients were included for final analysis (see Fig. 1 for detailed patient selection procedures). The electronic medical records of each patient were systematically evaluated for peak TG level measurement within the first-72 h post-onset of AP, body mass index (BMI), family history of HTG, family history of AP, number of episodes of HTG-AP and clinical severity of the disease.

Diagnosis of AP and classification of disease severity as mild (MAP), moderate severe (MSAP) and severe (SAP) were made in accordance with the Modified Atlanta Classification Standard in 2012.⁵⁰ HTG-AP was diagnosed in AP patients with fasting serum TG level of >1000 mg/dL (>11.3 mmol/L) alone or >500 mg/dL (>5.65 mmol/L) with coincidentally detected milky serum, with secondary factors such as gallstones and alcohol abuse being excluded as etiologies.⁵¹ 362 healthy Han Chinese subjects (60 newly recruited here and 302 published elsewhere³¹) were used as controls.

DNA sequencing

Genomic DNA was extracted from 0.5 ml whole blood by the TIANamp Blood DNA Kit (TIANGEN Biotech, Beijing, China). All coding and proximal intronic regions of the *LPL*, *APOC2*, *APOA5*, *GPIHBP1* and *LMF1* genes were PCR amplified from genomic DNA and subsequently Sanger sequenced as previously described.¹⁴ DNA extraction and PCR amplification were performed in our lab at the Jinling Hospital whilst Sanger sequencing was conducted by the Genewiz Life Science Company (Nanjing, China). All splice-site, missense, nonsense and frameshifting variants that had an allele frequency of <1% in East Asians (according to gnomAD) were considered to be potentially rare pathogenic variants (these data will be published elsewhere); all their corresponding carriers (n = 49) were excluded from this study (Fig. 1).

Statistical analysis

All data were analyzed using the SPSS 24.0 software package. Continuous variables were expressed as mean $\pm$ S.D. and tested using the Student's t test. *Chi*-Square and F-testing were performed to examine categorical variables. Benjamini-Hochberg test was subjected to a correction for multiple comparisons. *P* value < 0.05 was defined as statistically significant. Haldane's correction (i.e., adding 0.5 to all the cells of a contingency table) was applied for comparisons of the TT genotype frequencies in patients and controls because of the absence of TT homozygotes in controls.

Results

Division of patients into three groups

A total of 318 HTG-AP patients, all of Han Chinese origin, were included for final analysis in this study (Fig. 1). To assess the contribution of *APOA5* c.553G>T (rs2075291) to HTG-AP and its possible interaction with pregnancy, we divided the patients into three groups, firstly on the basis of gender and secondly, in the female patients, on the basis of disease occurrence during pregnancy or not (Fig. 1). The main demographic and clinical characteristics of the three groups of patients are summarized in Table 1. It should be noted that if a female patient experienced both HTG-APIP and HTG-AP not related to pregnancy, she was classified as group 3.

Comparisons of demographic and clinical characteristics between patient groups

We compared the demographic and clinical characteristics between the different patient groups. To increase biological relevance and, for the sake of simplicity, we performed comparisons only between group 1 and 2 patients and between group 2 and 3 patients. Only BMI displayed a significant difference between group 1 and 2 patients. By contrast, BMI did not exhibit any difference between group 2 and 3 patients. However, group 3 patients had a higher peak TG level approaching borderline statistical significance ($P = 0.058$), significantly earlier mean ages at disease onset, a significantly higher rate of SAP but a significantly lower rate of disease recurrence as compared to group 2 patients (Table 1). Differences in terms of age of onset of AP and disease recurrence could well have been related to the selection bias inherent in group 3 patients and will therefore not be considered further.

Comparisons of genotype and allele frequencies of rs2075291 between patient groups and between patients and controls

All exons and exon/intron boundaries of the five primary TG-related genes (*LPL*, *APOC2*, *APOA5*, *GPIHBP1* and *LMF1*) were analyzed by Sanger sequencing initially in 367 HTG-AP patients. Of these, 49 patients were found to carry rare putatively pathogenic variants in the five TG-related genes and excluded from analysis. All remaining 318 patients were informative with respect to the genotype status of rs2075291, and all participated in this study (Fig. 1). Genotype and minor T allele frequencies of rs2075291 in each of the three patient groups are summarized in Table 2. We also

sequenced the five primary TG-related genes in 60 Han Chinese healthy controls. The minor T allele frequency of rs2075291 in these controls showed no significant difference with that in 302 Han Chinese controls (0.07% (8/120) vs. 0.04% (24/604), $P = 0.22$) from Tang and colleagues³¹ (for details, see [Supplementary Table 1](#)). We therefore combined our data with those of Tang and colleagues³¹ to generate a single control dataset ([Table 2](#)).

We first compared the genotype and minor T allele frequencies of rs2075291 between patient groups. As in the preceding section, we performed comparisons only between group 1 and 2 patients and between group 2 and 3 patients. No significant difference was found for any parameters in the context of the group 1 and 2 comparisons. By contrast, both the TT genotype and the minor T allele of rs2075291 were significantly overrepresented in group 3 patients as compared to group 2 patients in the unadjusted analysis ($P = 0.015$ and 0.023) and approached or reached borderline statistical significance after Benjamini-Hochberg Correction ($P = 0.060$ and 0.046) ([Table 2](#)).

We then compared the minor T allele frequencies of rs2075291 and minor T-containing genotypes between each patient group and the control group, respectively. The minor T allele frequency of rs2075291 was significantly enriched in all three groups of patients as compared to controls, with the strongest enrichment being observed in group 3 (OR = 10.02, adjusted $P = 3.4E-9$; [Table 2](#)). Both the GT and TT genotypes were significantly enriched in all three groups of patients as compared to the control group, with the TT genotype invariably showing a much stronger effect than the GT genotype and the strongest enrichment being observed in group 3 patients (OR = 192.35, adjusted $P = 2.7E-7$) ([Table 2](#)).

As shown in [Table 2](#), the TT genotype was absent in the 362 controls but present in approximately 6% of both group 1 and group 2 patients and in up to 20% of group 3 patients. For comparison, the frequency of TT homozygotes in East Asians was 0.56% (55/9875) in accordance with the gnomAD database (accessed 03 March 2020).

Evaluation of the possible impact of rs2075291 on peak TG level, age of AP onset, disease severity and recurrence

We tested whether rs2075291 could possibly impact on peak TG level, age of AP onset, disease severity and recurrence in the patients. To this end, we compared these features between the GG, GT and TT genotype carriers in each of the three groups of patients. No significant difference was found for any parameter comparisons in either group 1 or group 2 patients (data not shown). These findings, together with the similar findings between the two groups observed in preceding sections, made it permissible to combine data from groups 1 and 2 (Fig. 2). In group 3 patients, TT genotype carriers exhibited significantly higher peak TG levels than GG genotype carriers (adjusted $P = 0.0042$; Fig. 3a). Although no other comparison showed any significant difference, two observations are worth mentioning. First, the GT genotype showed a trend toward an increase in peak TG level as compared to the GG genotype (Fig. 3a). Second, the TT genotype appeared to be associated with more severe disease as compared to the GG and GT genotypes (Fig. 3c). Impact on disease recurrence was not analyzed in group 3 patients owing to the very small number of patients who had more than one episode of AP in this group ($n = 3$), which was also deemed to be at least partly related to the selection bias inherent in group 3 patients.

Discussion

The etiology of HTG and, consequently HTG-AP, is complex and may involve multiple gene-gene and/or gene-environment interactions. *LPL*, *APOC2*, *APOA5*, *GPIHBP1* and *LMF1* are by far the most extensively studied HTG genes and pregnancy is a well-established environmental factor predisposing to HTG. Interaction between genetic risk factors in the five primary HTG genes and pregnancy in the etiology of HTG-AP has been frequently described in the literature but these studies have almost invariably been anecdotal case reports that have implicated rare pathogenic variants. The sole exception was a report of a possible interaction between the common *APOA5* c.553G>T variant (rs2075291) and pregnancy in HTG-AP;²⁴ this study was however somewhat limited in scope since only five patients with HTG-APIP (all Chinese) were analyzed.

The high frequency of the rs2075291 T allele in the Chinese population ($\sim 4\%$)^{30, 31} and its established association with altered TG levels made rs2075291 a good candidate (with sufficient

statistical power) to evaluate any interaction between this common genetic risk factor and pregnancy in HTG-AP. Therefore, we retrospectively recruited and sequenced a large cohort of Chinese HTG-AP patients for analysis. To perform this analysis properly, we first divided the patients into three groups. Groups 1 (male) and 2 (female; disease not related to pregnancy) served not only as controls *vis à vis* group 3 (HTG-APIP patients) but also as cohorts for evaluating the role of rs2075291 in HTG-AP unrelated to pregnancy. Since it is almost a truism that rare pathogenic variants have strong genetic effects whereas common pathogenic variants generally have only mild or modest genetic effects, to avoid interference from rare pathogenic variants, patients carrying such variants in the five primary HTG genes were excluded from this study.

Group 1 and 2 patients were initially treated as independent cohorts. The two groups showed remarkable similarity in almost all studied parameters, the only exception being a difference of 1.47 in terms of BMI value (Table 1). This suggested that groups 1 and 2 could effectively be considered together as a single group, as exemplified in Fig. 2. By contrast, biologically meaningful differences in terms of peak TG level and disease severity were apparent between group 3 patients and group 2 patients (as well as group 1 patients), with a higher peak TG level and a higher rate of SAP being observed in group 3 patients with HTG-APIP (Table 1).

Intuitively, an interaction between rs2075291 and pregnancy in HTG-AP should be reflected by a higher detection rate of the risk rs2075291 T allele and genotypes in HTG-APIP patients as compared to HTG-AP patients unrelated to pregnancy. Employing 362 healthy Han Chinese subjects as population controls, we demonstrated for the first time a robust association of rs2075291 with HTG-AP in general and HTG-APIP in particular (Table 2). Importantly, a stronger association was observed in group 3 patients than in either group 1 or group 2 patients in terms of the minor T allele and minor T allele-harboring genotype frequencies. This stronger association was mainly due to the much higher frequency of the TT genotype in group 3 patients (20%) than was found in groups 1 and 2 patients (<6%; Table 2). These findings provide the first evidence for an interaction between rs2075291 and pregnancy in HTG-AP.

To delve deeper into the interaction between rs2075291 and pregnancy in HTG-AP, we evaluated the possible impact of minor T allele-harboring genotypes on peak TG level, age of AP onset, disease

severity and recurrence by reference to the GG genotype within each patient group. We found no significant differences in any comparison between group 1 or 2 patients, with the combined data being provided in Fig. 2. However, we found that the TT genotype was associated with a significantly higher peak TG level in group 3 patients as compared to the GG genotype (adjusted $P = 0.0042$; Fig. 3); this constitutes a new line of evidence supporting an interaction between rs2075291 and pregnancy in HTG-AP.

Whilst we provided strong evidence to support an interaction between rs2075291 and pregnancy in HTG-AP, the most remarkable finding emerging from this study was related to the TT genotype. The TT genotype was found to be present at a much higher frequency (20%) in group 3 patients than in either group 1 or 2 patients (<6%) and it was also associated with a significantly higher peak TG level in group 3 patients. This may be understood in terms of the TT genotype comprising two identical risk alleles. Put simply, the homozygous TT genotype might be expected to exert double the genetic effect of a GT genotype, thereby rendering the corresponding carrier more prone to high HTG during pregnancy. Following this line of reasoning, our findings lend support to the mosaic genetic model of HTG, first proposed by Johansen, Kathiresan and Hegele in 2011.⁵² In essence, this model postulates the “stacking” of additional HTG risk alleles on top of a minimum number of genetic risk alleles and highlights a variable combination of genetic determinants and presence of environmental factors in causing HTG and HTG-related diseases.

The strength of this study lies in the analysis of three well-characterized patient groups. However, our study has various limitations. For example, we did not search for copy number variants in the five primary TG-related genes; although this is a very rare type of variant, it can have strong genetic effects on predisposing to HTG.^{13, 53, 54} The number of our group 3 patients is still relatively small and hence the corresponding findings should still be regarded as exploratory at this stage. Interaction between rs2075291 and pregnancy in HTG-AP remains to be confirmed in larger studies.

Conclusion

Here, on the basis of sequencing the five primary TG-related genes in three groups of well-defined Han Chinese participants, we have for the first time demonstrated a robust association of rs2075291

with HTG-AP in general and HTG-APIP in particular. We have also provided strong evidence for an interaction between rs2075291 and pregnancy in HTG-AP. Our findings provide novel insights into the complex etiology of HTG as well as HTG-AP and suggest that analysis of both rare and common pathogenic variants in TG-related genes is likely to be of key importance in risk assessment for HTG-AP in pregnancy.

Authorship contribution statement

Na Pu: Study design, data curation, and paper writing. Qi Yang: Study design, and paper writing. Xiao-Lei Shi and Wei-Wei Chen: Investigation and methodology. Xiao-Yao Li and Guo-Fu Zhang: Methodology and visualization. Gang Li and Bai-Qiang Li: Investigation and data curation. Lu Ke and Zhi-Hui Tong: Formal analysis and software. David N. Cooper: Paper review and critical revision of the manuscript. Jian-Min Chen: Data interpretation and paper writing. Wei-Qin Li and Jie-Shou Li: Study design, conceptualization and funding acquisition. All authors contributed to revision of the manuscript and approved the final manuscript.

Declarations of Competing Interests

The authors have no conflicts of interest to declare.

Role of the funding source

This study was supported by the National Natural Science Foundation of China (Nos. 81570584, 81670588, 81770641 and 81870441), the Key Research and Development Program Foundation of Jiangsu Province of China (Nos. BE2015685 and BE2016749), the Natural Science Foundation of Jiangsu Province (No. BK20190907), and Six Talent Peaks Project of Jiangsu Province (No. WSN-325). The funding sources did not play any role in designing this study, sample collection, analyses, interpretation of the data, or in writing the manuscript. We have not been paid to write this article by any pharmaceutical companies or other agencies. The corresponding authors, Qi Yang and Wei-Qin

Li, had full access to all data in the study and had final responsibility for the decision to submit for publication.

References

1. Tenner S, Baillie J, DeWitt J, Vege SS. American College of Gastroenterology guideline: management of acute pancreatitis. *Am J Gastroenterol*. 2013;108:1400-1415; 1416.
2. Yadav D, Lowenfels AB. The epidemiology of pancreatitis and pancreatic cancer. *Gastroenterology*. 2013;144:1252-1261.
3. Forsmark CE, Vege SS, Wilcox CM. Acute pancreatitis. *N Engl J Med*. 2016;375:1972-1981.
4. Carr RA, Rejowski BJ, Cote GA, Pitt HA, Zyromski NJ. Systematic review of hypertriglyceridemia-induced acute pancreatitis: A more virulent etiology? *Pancreatology*. 2016;16:469-476.
5. Huang YX, Jia L, Jiang SM, Wang SB, Li MX, Yang BH. Incidence and clinical features of hyperlipidemic acute pancreatitis from Guangdong, China: a retrospective multicenter study. *Pancreas*. 2014;43:548-552.
6. Zheng Y, Zhou Z, Li H, et al. A multicenter study on etiology of acute pancreatitis in Beijing during 5 years. *Pancreas*. 2015;44:409-414.
7. Zhu Y, Pan X, Zeng H, et al. A study on the etiology, severity, and mortality of 3260 patients with acute pancreatitis according to the Revised Atlanta Classification in Jiangxi, China over an 8-year period. *Pancreas*. 2017;46:504-509.
8. Simons-Linares CR, Jang S, Sanaka M, et al. The triad of diabetes ketoacidosis, hypertriglyceridemia and acute pancreatitis. How does it affect mortality and morbidity?: A 10-year analysis of the National Inpatient Sample. *Medicine (Baltimore)*. 2019;98:e14378.
9. Li X, Ke L, Dong J, et al. Significantly different clinical features between hypertriglyceridemia and biliary acute pancreatitis: a retrospective study of 730 patients from a tertiary center. *BMC Gastroenterol*. 2018;18:89.
10. Yuan G, Al-Shali KZ, Hegele RA. Hypertriglyceridemia: its etiology, effects and treatment. *Canadian Medical Association Journal*. 2007;176:1113-1120.
11. Rygiel K. Hypertriglyceridemia - Common causes, prevention and treatment strategies. *Curr Cardiol Rev*. 2018;14:67-76.
12. Hegele RA, Ginsberg HN, Chapman MJ, et al. The polygenic nature of hypertriglyceridaemia: implications for definition, diagnosis, and management. *Lancet Diabetes Endocrinol*. 2014;2:655-666.
13. Dron JS, Wang J, Cao H, et al. Severe hypertriglyceridemia is primarily polygenic. *J Clin Lipidol*. 2019;13:80-88.
14. Chen WW, Yang Q, Li XY, et al. Identification of a novel and heterozygous *LMF1* nonsense mutation in an acute pancreatitis patient with severe hypertriglyceridemia, severe obesity and heavy smoking. *Lipids Health Dis*. 2019;18:68.
15. Serveaux Dancer M, Di Filippo M, Marmontel O, et al. New rare genetic variants of *LMF1* gene identified in severe hypertriglyceridemia. *J Clin Lipidol*. 2018;12:1244-1252.
16. Knopp RH, Warth MR, Charles D, et al. Lipoprotein metabolism in pregnancy, fat transport to the fetus, and the effects of diabetes. *Biol Neonate*. 1986;50:297-317.
17. Ma Y, Liu MS, Ginzinger D, Frohlich J, Brunzell JD, Hayden MR. Gene-environment interaction in the conversion of a mild-to-severe phenotype in a patient homozygous for a Ser172-->Cys mutation in the lipoprotein lipase gene. *J Clin Invest*. 1993;91:1953-1958.
18. Ma Y, Ooi TC, Liu MS, et al. High frequency of mutations in the human lipoprotein lipase gene in pregnancy-induced chylomicronemia: possible association with apolipoprotein E2 isoform. *J Lipid Res*. 1994;35:1066-1075.
19. Keilson LM, Vary CP, Sprecher DL, Renfrew R. Hyperlipidemia and pancreatitis during pregnancy

- in two sisters with a mutation in the lipoprotein lipase gene. *Ann Intern Med.* 1996;124:425-428.
20. Henderson H, Leisegang F, Hassan F, Hayden M, Marais D. A novel Glu421Lys substitution in the lipoprotein lipase gene in pregnancy-induced hypertriglyceridemic pancreatitis. *Clinica Chimica acta; international journal of clinical chemistry.* 1998;269:1-12.
 21. Murugasu CG, Armstrong G, Creedon G, Cavanna JS, Galton DJ, Tomkin GH. Acute hypertriglyceridaemic pancreatitis in a pregnant Indian: a new lipoprotein lipase gene mutation. *J R Soc Med.* 1998;91:205-207.
 22. Suga S, Tamasawa N, Kinpara I, et al. Identification of homozygous lipoprotein lipase gene mutation in a woman with recurrent aggravation of hypertriglyceridaemia induced by pregnancy. *J Intern Med.* 1998;243:317-321.
 23. Barthä I, Dinya T, Seres I, et al. Acute hypertriglyceridemic pancreatitis during pregnancy due to homozygous lipoprotein lipase gene mutation. *Clinica Chimica Acta* 2009;400:137-138.
 24. Xie SL, Chen TZ, Huang XL, et al. Genetic variants associated with gestational hypertriglyceridemia and pancreatitis. *PLoS One.* 2015;10:e0129488.
 25. Liu Y, Lun Y, Lv W, Hou X, Wang Y. A Chinese patient with recurrent pancreatitis during pregnancy induced by hypertriglyceridemia associated with compound heterozygosity (Glu242Lys and Leu252Val) in the lipoprotein lipase gene. *J Clin Lipidol.* 2016;10:199-203.e191.
 26. Chyzyk V, Kozmic S, Brown AS, et al. Extreme hypertriglyceridemia: Genetic diversity, pancreatitis, pregnancy, and prevalence. *J Clin Lipidol.* 2019;13:89-99.
 27. Kathiresan S, Willer CJ, Peloso GM, et al. Common variants at 30 loci contribute to polygenic dyslipidemia. *Nat Genet.* 2009;41:56-65.
 28. Johansen CT, Hegele RA. Genetic bases of hypertriglyceridemic phenotypes. *Current opinion in lipidology.* 2011;22:247-253.
 29. Surendran RP, Visser ME, Heemelaar S, et al. Mutations in *LPL*, *APOC2*, *APOA5*, *GPIHBP1* and *LMF1* in patients with severe hypertriglyceridaemia. *J Intern Med.* 2012;272:185-196.
 30. Kao JT, Wen HC, Chien KL, Hsu HC, Lin SW. A novel genetic variant in the apolipoprotein A5 gene is associated with hypertriglyceridemia. *Hum Mol Genet.* 2003;12:2533-2539.
 31. Tang Y, Sun P, Guo D, et al. A genetic variant c.553G > T in the apolipoprotein A5 gene is associated with an increased risk of coronary artery disease and altered triglyceride levels in a Chinese population. *Atherosclerosis.* 2006;185:433-437.
 32. Hsu LA, Ko YL, Chang CJ, et al. Genetic variations of apolipoprotein A5 gene is associated with the risk of coronary artery disease among Chinese in Taiwan. *Atherosclerosis.* 2006;185:143-149.
 33. Zhai G, Wen P, Guo L, Chen L. Association of *APOA5* c.553G>T polymorphism with type 2 diabetes mellitus in a Chinese population. *Clin Chem Lab Med.* 2006;44:1313-1316.
 34. Matsunaga A, Arishima H, Niimura H ea. Strong linkage disequilibrium and association of-1131T>C and c. 553G>T polymorphisms of the apolipoprotein A5 gene with hypertriglyceridemia in a Japanese population. *Circ J.* 2007;71:746-752.
 35. Pullinger CR, Aouizerat BE, Movsesyan I, et al. An apolipoprotein A-V gene SNP is associated with marked hypertriglyceridemia among Asian-American patients. *Journal of lipid research.* 2008;49:1846-1854.
 36. Li YY, Yin RX, Lai CQ, et al. Association of apolipoprotein A5 gene polymorphisms and serum lipid levels. *Nutr Metab Cardiovasc Dis.* 2011;21:947-956.
 37. Yin RX, Li YY, Liu WY, Zhang L, Wu JZ. Interactions of the apolipoprotein A5 gene polymorphisms and alcohol consumption on serum lipid levels. *PLoS One.* 2011;6:e17954.
 38. Hishida A, Morita E, Naito M, et al. Associations of apolipoprotein A5 (*APOA5*), glucokinase (*GCK*) and glucokinase regulatory protein (*GCKR*) polymorphisms and lifestyle factors with the risk of dyslipidemia and dysglycemia in Japanese - a cross-sectional data from the J-MICC Study. *Endocr J.* 2012;59:589-599.
 39. Lee MJ, Chien KL, Chen MF, Stephenson DA, Su TC. Overweight modulates *APOE* and *APOA5* alleles on the risk of severe hypertriglyceridemia. *Clinica Chimica Acta* 2013;416:31-35.

- 418 **40.** Li S, Hu B, Wang Y, Wu D, Jin L, Wang X. Influences of *APOA5* variants on plasma triglyceride
419 levels in Uyghur population. *PLoS One*. 2014;9:e110258.
- 420 **41.** Chiou KR, Chen CY, Charng MJ. Genetic Diagnosis via Whole Exome Sequencing in Taiwanese
421 Patients with Hypertriglyceridemia. *Journal of Atherosclerosis and Thrombosis*. 2015;22:887-
422 900.
- 423 **42.** Khovidhunkit W, Charoen S, Kiateprungvej A, Chartyingcharoen P, Muanpetch S, Plengpanich
424 W. Rare and common variants in *LPL* and *APOA5* in Thai subjects with severe
425 hypertriglyceridemia: A resequencing approach. *J Clin Lipidol*. 2016;10:505-511 e501.
- 426 **43.** Kim M, Kim M, Yoo HJ, Bang YJ, Lee SH, Lee JH. Apolipoprotein A5 gene variants are associated
427 with decreased adiponectin levels and increased arterial stiffness in subjects with low high-
428 density lipoprotein-cholesterol levels. *Clin Genet*. 2018;94:438-444.
- 429 **44.** Dorfmeister B, Zeng WW, Dichlberger A, et al. Effects of six *APOA5* variants, identified in
430 patients with severe hypertriglyceridemia, on in vitro lipoprotein lipase activity and receptor
431 binding. *Arterioscler Thromb Vasc Biol*. 2008;28:1866-1871.
- 432 **45.** Huang YJ, Lin YL, Chiang CI, Yen CT, Lin SW, Kao JT. Functional importance of apolipoprotein A5
433 185G in the activation of lipoprotein lipase. *Clinica Chimica Acta*. 2012;413:246-250.
- 434 **46.** Sharma V, Witkowski A, Witkowska HE, et al. Aberrant hetero-disulfide bond formation by the
435 hypertriglyceridemia-associated p.Gly185Cys *APOA5* variant (rs2075291). *Arterioscler Thromb*
436 *Vasc Biol*. 2014;34:2254-2260.
- 437 **47.** Chang CK, Lin XR, Lin YL, et al. Magnolol-mediated regulation of plasma triglyceride through
438 affecting lipoprotein lipase activity in apolipoprotein A5 knock-in mice. *PLoS One*.
439 2018;13:e0192740.
- 440 **48.** Arai M, Nishimura A, Mori Y, Ebara T, Okubo M. Hypertriglyceridemia and pancreatitis in a
441 patient with apolipoprotein E7 (p.[E244K; E245K])/E4. *Clinica Chimica Acta* 2014;436:188-192.
- 442 **49.** Chen WJ, Sun XF, Zhang RX, et al. Hypertriglyceridemic acute pancreatitis in emergency
443 department: Typical clinical features and genetic variants. *J Dig Dis*. 2017;18:359-368.
- 444 **50.** Banks PA, Bollen TL, Dervenis C, et al. Classification of acute pancreatitis--2012: revision of the
445 Atlanta classification and definitions by international consensus. *Gut*. 2013;62:102-111.
- 446 **51.** Scherer J, Singh VP, Pitchumoni CS, Yadav D. Issues in hypertriglyceridemic pancreatitis: an
447 update. *J Clin Gastroenterol*. 2014;48:195-203.
- 448 **52.** Johansen CT, Kathiresan S, Hegele RA. Genetic determinants of plasma triglycerides. *Journal of*
449 *lipid research*. 2011;52:189-206.
- 450 **53.** Dron JS, Wang J, McIntyre AD, et al. Partial *LPL* deletions: rare copy-number variants
451 contributing towards severe hypertriglyceridemia. *Journal of lipid research*. 2019;60:1953-
452 1958.
- 453 **54.** Iacocca MA, Dron JS, Hegele RA. Progress in finding pathogenic DNA copy number variations
454 in dyslipidemia. *Current opinion in lipidology*. 2019;30:63-70.

Table 1. Main demographic and clinical characteristics of the three groups of Han Chinese HTG-AP patients

	Group 1 (Male)	Group 2 (Female; disease unrelated to pregnancy)	Group 3 (Female; HTG-APIP)
Number	183	105	30
Age (year)	39.8 ± 8.9	39.2 ± 9.6	29.2 ± 4.1 ^{***}
Family history of HTG	4 (2.19%)	2 (1.90%)	0 (0)
Family history of AP	5 (2.73%)	1 (0.95%)	0 (0)
BMI (kg/m ²)	26.97 ± 13.58	25.50 ± 3.90 ^{**}	25.33 ± 3.59
Peak TG (mmol/L)	20.00 ± 17.93	21.11 ± 18.87	29.45 ± 29.35 ^a
Age at AP onset (year) ^b	37.0 ± 8.5	37.6 ± 9.6	29.0 ± 4.0 ^{***}
Disease severity			
MAP	60 (32.79%)	26 (24.76%)	1 (3.33%)
MSAP	61 (33.33%)	39 (37.14%)	10 (33.33%)
SAP	62 (33.88%)	40 (38.10%)	19 (63.34%) [*]
Episodes			
1	103 (56.28%)	67 (63.81%)	27 (90%)
>1	80 (43.72%)	38 (36.19%)	3 (10%) ^{**}

Comparisons between patient groups were performed only between group 1 and group 2 patients and between group 2 and group 3 patients. No significant difference was observed unless specifically indicated. *, $P < 0.05$. **, $P < 0.01$. ***, $P < 0.001$.

^a $P = 0.058$.

^b First episode of disease in case of recurrence in groups 1 and 2 patients; first episode of HTG-APIP in case of group 3 patients.

AP, acute pancreatitis; BMI, body mass index; HTG, hypertriglyceridemia; HTG-AP, HTG-induced AP; HTG-APIP, HTG-AP in pregnancy; MAP, mild AP; MSAP, moderate severe AP; SAP, severe AP.

Table 2. Frequency of the *APOA5* c.553G>T polymorphism and associated genotypes in the three groups of Chinese HTG-AP patients and controls

Genotype	Patient	Control (n = 362)	<i>P</i> value (unadjusted)	<i>P</i> value (after Benjamini-Hochberg correction)	OR
	<i>Group 1 (n = 183)</i>				
GG	134 (73.22%)	330 (91.16%)			
GT	39 (21.31%)	32 (8.84%)	1.2E-5	2.1E-5	3.00
TT	10 (5.47%)	0 (0.00%)	4.4E-5	6.6E-5	43.88
GT+TT	49 (26.78%)	32 (8.84%)	2.7E-8	1.18E-7	3.77
Minor T allele	59 (16.12%)	32 (4.42%)	0.0040	0.0048	4.16
	<i>Group 2 (n = 105)</i>				
GG	73 (69.52%)	330 (91.16%)			
GT	26 (24.76%)	32 (8.84%)	4.0E-6	8.0E-6	3.67
TT	6 (5.72%)	0 (0.00%)	1.7E-4	2.3E-4	47.36
GT+TT	32 (30.48%)	32 (8.84%)	1.4E-8	8.3E-8	4.52
Minor T allele	38 (18.10%)	32 (4.42%)	0.023	0.023	4.78
	<i>Group 3 (n = 30)</i>				
GG	17 (56.67%)	330 (91.16%)			
GT	7 (23.33%)	32 (8.84%)	0.0060	0.0065	4.25
TT	6 (20.00%) ^b	0 (0.00%)	8.9E-8	2.7E-7	192.35
GT+TT	13 (43.33%)	32 (8.84%)	3.0E-6	7.2E-6	7.89
Minor T allele	19 (31.67%) ^c	32 (4.42%)	2.8E-10	3.4E-9	10.02

Genotype and allele frequency comparisons between patient groups were performed either between group 1 and group 2 or between group 2 and group 3. No significant difference was observed unless specifically indicated. Haldane' correction was applied for comparisons of TT genotype frequencies in patients and controls.

^b Unadjusted *P* = 0.015; adjusted *P* = 0.060 (Group 3 vs. Group 2).

^c Unadjusted *P* = 0.023; adjusted *P* = 0.046 (Group 3 vs. Group 2).

OR, odds ratio.

Figure 1 Flow chart of patient selection procedures. HTG-AP, hypertriglyceridemia-induced acute pancreatitis; HTG-APIP, HTG-AP in pregnancy. TG, triglyceride.

Figure 2 Comparison of peak TG level (a), age of onset (b), number of episodes (c) and clinical severity (d) of HTG-AP patients who carried the different *APOA5* c.553G>T genotypes in the combined group 1 and 2 patients. Numbers of the GG, GT and TT genotype patients were 207, 65 and 16, respectively. In panels **c** and **d**, percentages and exact numbers (in brackets) of the subjects with the indicated clinical characteristics are provided in the context of each genotype. AP, acute pancreatitis; MAP, mild AP; MSAP, moderate severe AP; SAP, severe AP; TG, triglyceride.

Figure 3 Comparison of peak TG level (a), age of onset (b) and clinical severity (c) of group 3 (HTG-APIP) patients who carried the different *APOA5* c.553G>T genotypes. Numbers of the GG, GT and TT genotype carriers were 17, 7 and 6, respectively. In panel c, percentages and exact numbers (in brackets) of the subjects with the indicated clinical characteristics are provided in the context of each genotype. **, adjusted $P = 0.0042$ (TT vs. GG). AP, acute pancreatitis; MAP, mild AP; MSAP, moderate severe AP; SAP, severe AP.