Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/136839/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Norris, Joseph M., Allen, Clare, Ball, Rhys, Freeman, Alex, Giganti, Francesco, Kelly, Daniel ORCID: <https://orcid.org/0000-0002-1847-0655>, Kirkham, Alex, Simpson, Benjamin S., Stavriniades, Vasilis, Whitaker, Hayley C. and Emberton, Mark 2021. Prostate cancer undetected by mpMRI: tumor conspicuity is reliant upon optimal scan timing and quality. *Urology* 148 , pp. 316-317. 10.1016/j.urology.2020.11.037 file

Publishers page: <http://dx.doi.org/10.1016/j.urology.2020.11.037>
<<http://dx.doi.org/10.1016/j.urology.2020.11.037>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.

Prostate Cancer Undetected by mpMRI: Tumour Conspicuity is Reliant Upon Optimal Scan Timing and Quality

Joseph M. Norris^{1,2}, Clare Allen³, Rhys Ball⁴, Alex Freeman⁴, Francesco Giganti^{1,3}, Daniel Kelly⁵, Alex Kirkham³, Benjamin S. Simpson¹, Vasilis Stavrinos¹, Hayley C. Whitaker¹, Mark Emberton^{1,2}

¹ UCL Division of Surgery & Interventional Science, University College London, London, UK

² Department of Urology, University College London Hospitals NHS Foundation Trust, London, UK

³ Department of Radiology, University College London Hospitals NHS Foundation Trust, London, UK

⁴ Department of Pathology, University College London Hospitals NHS Foundation Trust, London, UK

⁵ School of Healthcare Sciences, Cardiff University, Wales, UK

Address correspondence to: Joseph M. Norris, BSc (Hons), MSc, BM BS, MRCS (Eng), FESSR. UCL Division of Surgery & Interventional Science, 3rd Floor, Charles Bell House, 43-45 Foley Street, London W1W 7TS, UK. E-mail: joseph.norris@ucl.ac.uk

Article type: Letter-to-the-Editor.

Word count: 558 words.

Ethics: Not required.

Funding: Joseph M. Norris is funded by the Medical Research Council (MRC) (Grant Reference: MR/S00680X/1). Francesco Giganti is funded by the UCL Graduate Research Scholarship and the Brahm PhD scholarship in memory of Chris Adams.

Conflicts of interest: Norris and Stavrinos receive funding from the MRC. Freeman, Kirkham and Emberton have stock interest in Nuada Medical Ltd. Whitaker receives funding from Prostate Cancer UK, the Urology Foundation and the Rosetrees Trust. Simpson receives funding from the Rosetrees Trust. Emberton receives funding from NIHR-i4i, MRC, Sonacare Inc., Trod Medical, Cancer Vaccine Institute and Sophiris Biocorp for trials in prostate cancer. Emberton is a medical consultant to Sonacare Inc., Sophiris Biocorp, Steba Biotech, GSK, Exact Imaging and Profound Medical. Travel allowance was previously provided from Sanofi Aventis, Astellas, GSK, and Sonacare. Emberton is a proctor for HIFU with Sonacare Inc. and is paid for training other surgeons in this procedure.

We commend Alanee and colleagues for their detailed description of prostate cancers not detected by multiparametric magnetic resonance imaging (mpMRI).¹ They analysed radical prostatectomy findings for 33 men lacking mpMRI-conspicuous prostate cancer (defined as Prostate Imaging Reporting and Data System [PI-RADS] scores 1–2) on pre-operative mpMRI. They found 27% (9/33) of men with negative mpMRI had Gleason Grade Group (GGG) 3–5 cancer at prostatectomy, including nine with extraprostatic extension, and one with Gleason grade 5 + 5 disease. Ostensibly, these findings are at odds with contemporary evidence in which mpMRI-invisible disease harbours reassuring molecular and histopathological features (i.e. overall GGG $\leq$ 2).^{2,3} We believe that this discrepancy is attributable to certain methodological concerns, including quality of mpMRI acquisition.⁴

Here, every included patient underwent post-biopsy mpMRI, despite guidelines now requiring mpMRI to precede biopsy.^{1,2} In fact, the 14 men who underwent pre-biopsy mpMRI in this cohort were excluded from further analysis. The authors acknowledge post-biopsy haemorrhage contributed to disease inconspicuity in approximately one third of men (2–3/9) with high-grade disease, however, given this well-established phenomenon, this proportion may be higher.⁵ As post-biopsy haemorrhage was accepted and incorporated into this study, it is possible that other radiological features (e.g. background patchy/diffuse patterns) may have contributed to reduced tumour conspicuity.

In addition to mpMRI quality, other aspects of this study warrant scrutiny. Of note, 45% (15/33) of men had grade reclassification from random 12-core transrectal ultrasound (TRUS)-guided biopsy to radical prostatectomy (18% downgraded, 27% upgraded). This effect may be attributable to an imperfect reference standard (random TRUS-guided biopsy) which demonstrably overlooks significant cancer approximately half the time.² Furthermore, whilst mpMRI were scored according to PI-RADSv2.1 guidelines, it seems unusual that men with ‘negative’ mpMRI had such high prostate specific antigen densities (PSAD; e.g. 1.08, 0.48 and 0.22 ng/mL/mL) which, in other settings, may have raised radiological suspicion. Unfortunately, a number of key details are missing to fully appraise this study including, biopsy core length, tumour size at prostatectomy, age of MRI machines, number of reporting radiologists and their experience in prostate mpMRI reporting (i.e. how many prostate MR scans per year), all of which impact upon tumour detection on mpMRI. Lastly, in their discussion, the authors cite the Prostate MRI Study (PROMIS), proposing that a 10% non-detection rate of significant disease by mpMRI is a “considerable risk,” however, they do not quote the false negative rate of systematic TRUS-guided biopsy (their own reference standard), which had a non-detection rate of over 50%, in the same study.²

Collectively, we should work toward optimal mpMRI-directed pathway delivery, at every juncture, including scan acquisition, reporting, and biopsy. In an attempt to standardise mpMRI quality, the Prostate Imaging Quality (PI-QUAL) score was developed, based on a 1-to-5 Likert scale derived from evaluation of each sequence, against objective quality

criteria in line with the PI-RADSv2 recommendations.⁴ Work is currently underway to evaluate effects of PI-QUAL on tumour conspicuity, however, we hope that this scheme provides a starting point for centres to evaluate quality of mpMRI delivery. Alanee and colleagues should be congratulated for adding to the mpMRI literature, expounding links between histopathology and radiology, however, we believe their findings should be cautiously interpreted in light of the methodological issues highlighted here. We agree that long-term ramifications of mpMRI conspicuity remain pressing avenues for future research and we eagerly await results of ongoing work.

References

1. Alanee S, Deebajah M, Taneja K, Cole D, Pantelic M, Peabody J, et al. Post prostatectomy pathologic findings of patients with clinically significant prostate cancer and no significant PI-RADS lesions on preoperative magnetic resonance imaging. *Urology*. 2020;S0090-4295(20)31127–4.
2. Norris JM, Carmona Echeverria LM, Bott SRJ, Brown LC, Burns-Cox N, Dudderidge T, et al. What type of prostate cancer is systematically overlooked by multiparametric magnetic resonance imaging? an analysis from the PROMIS cohort. *Eur Urol*. 2020;78:163–170.
3. Norris JM, Simpson BS, Parry MA, Allen C, Ball R, Freeman A, et al. Genetic landscape of prostate cancer conspicuity on multiparametric magnetic resonance imaging: a systematic review and bioinformatic analysis. *Eur Urol Open Sci*. 2020;20:37–47.
4. Giganti F, Allen C. Imaging quality and prostate MR: it's time to improve. *Br J Radiol*. 2020;20200934.
5. Caglic I, Barrett T. Optimising prostate mpMRI: prepare for success *Clin Radiol*. 2019;74:831–840.