Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/138475/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Taylor, Mark, Ronald, Angelica, Martin, Joanna ORCID: <https://orcid.org/0000-0002-8911-3479>, Lundström, Sebastian, Hosang, Georgina and Lichtenstein, Paul 2022. Examining the association between autistic traits in childhood and adolescent hypomania: a longitudinal twin study. *Psychological Medicine* 52 (15) , pp. 3606-3615. 10.1017/S0033291721000374 file

Publishers page: <https://doi.org/10.1017/S0033291721000374>
<<https://doi.org/10.1017/S0033291721000374>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.

Supplementary Materials

Paper Title: Examining the association between autistic traits in childhood and adolescent hypomania: a longitudinal twin study

Authors: Mark J. Taylor, Angelica Ronald, Joanna Martin, Sebastian Lundström, Georgina Hosang, & Paul Lichtenstein

Table S1: Neurodevelopmental and Psychiatric Disorders in CATSS Participants

Table S2: Estimates from the Multivariate Cholesky Decomposition of Autistic Traits and Hypomania

Subscale-Specific Analyses

Table S3: Twin Correlations

Table S4: Model Fit Statistics

Table S5: Estimates for Social Interaction Difficulties

Table S6: Estimates for Communication Problems

Table S7: Estimates for Inflexibility

Table S8: Posthoc Analysis of Social Communication Problems

Table S1 Neurodevelopmental and Psychiatric Disorders in CATSS Participants

	CATSS-9/12	CATSS -15		CATSS -18	
	Participant	Participant	Non-Participant	Participant	Non-Participant
Any Disorder	2217 (8.22%)	705 (9.25%)	1512 (7.81%)	544 (9.26%)	1673 (7.93%)
ASD	392 (1.45%)	101 (1.33%)	291 (1.50%)	69 (1.17%)	323 (1.53%)
ADHD	985 (3.65%)	238 (3.12%)	747 (3.86%)	143 (2.43%)	842 (3.99%)
Anxiety	679 (2.52%)	255 (3.35%)	424 (2.19%)	234 (3.98%)	445 (2.11%)
Depression	619 (2.29%)	249 (3.27%)	370 (1.91%)	215 (3.66%)	404 (1.91%)
OCD	133 (0.49%)	44 (0.58%)	89 (0.46%)	29 (0.49%)	104 (0.49%)
Schizophrenia	25 (0.09%)	8 (0.11%)	17 (0.09%)	10 (0.17%)	15 (0.07%)
Bipolar Disorder	78 (0.29%)	32 (0.42%)	46 (0.24%)	41 (0.70%)	37 (0.18%)

All clinical diagnoses were identified from the National Patient Register (NPR), which records all specialist inpatient and outpatient care given to residents of Sweden. Bipolar disorder was additionally defined as having been prescribed lithium according to the Prescribed Drug Register (PDR), which records all prescribed medications dispensed to residents of Sweden since 2005.

Table S2 Estimates from the Multivariate Cholesky Decomposition of Autistic Traits and Hypomania

	A1	A2	A3	C1	C2	C3	E1	E2	E3
<i>Female Estimates</i>									
A-TAC ASD	0.61 (0.56-0.65)	-----	-----	-----	-----	-----	0.39 (0.35-0.44)	-----	-----
CMRS	0.09 (0.06-0.12)	0.44 (0.32-0.55)	-----	-----	0.25 (0.15-0.36)	-----	0.00 (0.00-0.01)	0.22 (0.20-0.26)	-----
MDQ	0.07 (0.04-0.11)	0.09 (0.04-0.17)	0.39 (0.22-0.55)	-----	-----	0.11 (0.00-0.26)	0.00 (0.00-0.01)	0.01 (0.00-0.04)	0.32 (0.28-0.37)
<i>Male Estimates</i>									
A-TAC ASD	0.73 (0.71-0.76)	-----	-----	-----	-----	-----	0.27 (0.24-0.29)	-----	-----
CMRS	0.07 (0.05-0.10)	0.48 (0.37-0.58)	-----	-----	0.22 (0.14-0.33)	-----	0.00 (0.00-0.01)	0.22 (0.19-0.25)	-----
MDQ	0.06 (0.03-0.10)	0.10 (0.05-0.19)	0.44 (0.34-0.52)	-----	-----	0.00 (0.00-0.08)	0.00 (0.00-0.01)	0.01 (0.00-0.04)	0.38 (0.32-0.45)

A-TAC: Autism-Tics, ADHD, and other Comorbidities inventory; CMRS: Child Mania Rating Scale; MDQ: Mood Disorders Questions
A1, C1, E1: additive genetic, shared environmental, and nonshared environmental influences on autistic traits, and their effect on the
two hypomania measures. Note that shared environment was fixed to zero for autistic traits. A2, C2, E2: genetic and environmental
influences on hypomania at age 15, and their effect on hypomania at age 15. These pathways are distinct from the effects that
hypomania at age 15 shares with autistic traits. A3, C3, E3: genetic and environmental influences on hypomania at age 18, which are
entirely unique to hypomania at age 18.

Subscale-Specific Analyses

Table S3 Twin Correlations

	rPH, F	rPH, M	MZF	DZF	MZM	DZM	DZOS
A-TAC Social -> CMRS	0.27 (0.23-0.30)	0.25 (0.21-0.29)	0.20 (0.15-0.25)	0.12 (0.07-0.18)	0.22 (0.18-0.27)	0.15 (0.09-0.20)	0.14 (0.10-0.18)
A-TAC Social -> MDQ	0.25 (0.21-0.29)	0.25 (0.20-0.29)	0.16 (0.10-0.22)	0.10 (0.03-0.16)	0.23 (0.16-0.30)	0.13 (0.05-0.20)	0.11 (0.06-0.17)
A-TAC Language -> CMRS	0.19 (0.15-0.22)	0.19 (0.15-0.23)	0.19 (0.15-0.23)	0.09 (0.04-0.14)	0.17 (0.12-0.21)	0.17 (0.11-0.22)	0.16 (0.12-0.20)
A-TAC Language -> MDQ	0.14 (0.10-0.19)	0.12 (0.08-0.17)	0.16 (0.11-0.22)	0.08 (0.01-0.15)	0.10 (0.04-0.16)	0.11 (0.04-0.18)	0.09 (0.03-0.14)
A-TAC Flexibility -> CMRS	0.21 (0.17-0.25)	0.22 (0.18-0.26)	0.19 (0.14-0.24)	0.10 (0.05-0.15)	0.18 (0.14-0.23)	0.15 (0.09-0.21)	0.12 (0.08-0.16)
A-TAC Flexibility -> MDQ	0.18 (0.13-0.22)	0.22 (0.17-0.27)	0.15 (0.09-0.21)	0.11 (0.04-0.17)	0.22 (0.16-0.28)	0.13 (0.05-0.20)	0.10 (0.05-0.15)

A-TAC: Autism-Tics, ADHD, and other Comorbidities inventory; CMRS: Child Mania Rating Scale; MDQ: Mood Disorders Questions

Table S4 Twin Model Fit Statistics

Model	-2LL	Parameters	df	BIC	Comparison Model	$\Delta\chi^2$	Δdf	p
<i>Social Problems</i>								
Fully Saturated	103216.40	135	38203	-260781.54	-----	-----	-----	-----
ADCE-s	103670.13	39	38299	-261242.50	Fully Saturated	453.73	96	<0.001
ADCE-s Hom	104205.33	22	38316	-260869.28	ADCE-s	535.20	17	<0.001
ACE-s	103670.13	37	38301	-261261.56	ADCE-s	0.00	2	1.00
ADE-s	103731.79	35	38303	-261218.96	ADCE-s	61.66	4	<0.001
AE-s	103731.79	33	38305	-261238.01	ADCE-s	61.66	6	<0.001
AE	103794.38	31	38307	-261194.48	ADCE-s	124.25	8	<0.001
E	107225.31	19	38319	-257877.88	ADCE-s	3555.18	20	<0.001
<i>Communication Difficulties</i>								
Fully Saturated	102511.05	135	38238	-261820.38	-----	-----	-----	-----
ADCE-s	102873.36	39	38334	-262372.75	Fully Saturated	362.31	96	<0.001
ADCE-s Hom	103535.44	22	38351	-261872.65	ADCE-s	662.08	17	<0.001
ACE-s	102888.20	37	38336	-262376.97	ADCE-s	14.84	2	<0.001
ADE-s	102932.36	35	38338	-262351.87	ADCE-s	59.00	4	<0.001
AE-s	102944.71	33	38340	-262358.57	ADCE-s	71.35	6	<0.001
AE	102961.72	31	38342	-262360.62	ADCE-s	88.36	8	<0.001
E	107474.88	19	38354	-257961.8	ADCE-s	4601.52	20	<0.001
<i>Behavioral Inflexibility</i>								
Fully Saturated	102783.48	135	38236	-261528.89	-----	-----	-----	-----
ADCE-s	103388.67	39	38332	-261838.38	Fully Saturated	605.19	96	<0.001
ADCE-s Hom	104506.01	22	38349	-260883.03	ADCE-s	1117.33	17	<0.001
ACE-s	103388.67	37	38334	-261857.44	ADCE-s	0.00	2	1.00
ADE-s	103447.50	35	38336	-261817.67	ADCE-s	58.83	4	<0.001
AE-s	103447.50	33	38338	-261836.73	ADCE-s	58.83	6	<0.001
AE	103553.21	31	38340	-261750.07	ADCE-s	164.54	8	<0.001
E	106880.39	19	38352	-258537.23	ADCE-s	3491.72	20	<0.001

A-TAC: Autism-Tics, ADHD, and other Comorbidities inventory; CMRS: Child Mania Rating Scale; MDQ: Mood Disorders Questions

A: additive genetic influences; D: non-additive genetic influences; C: shared environmental influences; E: nonshared environmental influences; s: sibling interaction parameters; Hom: homogeneity, indicating no sex differences.
*-2LL: fit statistic, which is -2*log-likelihood of the data; df: degrees of freedom; BIC: Bayesian Information Criteria; $\Delta\chi^2$: -2LL difference between two models, distributed chi-square; Δdf : difference in degrees of freedom between models, equal to the difference in number of parameters.*

Table S5 Estimates for Social Difficulties

	A1	A2	A3	C1	C2	C3	E1	E2	E3
<i>Female Estimates</i>									
A-TAC ASD	0.55 (0.49-0.60)	-----	-----	-----	-----	-----	0.45 (0.40-0.51)	-----	-----
CMRS	0.06 (0.04-0.10)	0.48 (0.36-0.57)	-----	-----	0.23 (0.15-0.33)	-----	0.01 (0.00-0.02)	0.22 (0.19-0.25)	-----
MDQ	0.05 (0.02-0.09)	0.10 (0.05-0.18)	0.39 (0.22-0.55)	-----	-----	0.11 (0.00-0.27)	0.01 (0.00-0.03)	0.01 (0.00-0.03)	0.32 (0.27-0.37)
<i>Male Estimates</i>									
A-TAC ASD	0.71 (0.68-0.74)	-----	-----	-----	-----	-----	0.29 (0.26-0.32)	-----	-----
CMRS	0.08 (0.05-0.11)	0.46 (0.35-0.56)	-----	-----	0.25 (0.16-0.35)	-----	0.00 (0.00-0.01)	0.22 (0.19-0.25)	-----
MDQ	0.08 (0.04-0.12)	0.11 (0.05-0.20)	0.42 (0.32-0.51)	-----	-----	0.00 (0.00-0.09)	0.00 (0.00-0.01)	0.01 (0.00-0.04)	0.38 (0.32-0.44)

A-TAC: Autism-Tics, ADHD, and other Comorbidities inventory; CMRS: Child Mania Rating Scale; MDQ: Mood Disorders Questions
A1, C1, E1: additive genetic, shared environmental, and nonshared environmental influences on autistic traits, and their effect on the two hypomania measures. Note that shared environment was fixed to zero for autistic traits. A2, C2, E2: genetic and environmental influences on hypomania at age 15, and their effect on hypomania at age 15. These pathways are distinct from the effects that hypomania at age 15 shares with autistic traits. A3, C3, E3: genetic and environmental influences on hypomania at age 18, which are entirely unique to hypomania at age 18.

Table S6 Estimates for Communication Problems

	A1	A2	A3	C1	C2	C3	E1	E2	E3
<i>Female Estimates</i>									
A-TAC ASD	0.66 (0.62-0.69)	----	----	----	----	----	0.34 (0.31-0.38)	----	----
CMRS	0.05 (0.03-0.07)	0.47 (0.36-0.58)	----	----	0.25 (0.15-0.36)	----	0.00 (0.00-0.00)	0.23 (0.20-0.26)	----
MDQ	0.04 (0.02-0.07)	0.12 (0.06-0.20)	0.39 (0.20-0.55)	----	----	0.11 (0.00-0.27)	0.00 (0.00-0.01)	0.01 (0.00-0.04)	0.33 (0.28-0.38)
<i>Male Estimates</i>									
A-TAC ASD	0.70 (0.66-0.73)	----	----	----	----	----	0.30 (0.27-0.34)	----	----
CMRS	0.04 (0.02-0.06)	0.52 (0.40-0.61)	----	----	0.23 (0.14-0.33)	----	0.00 (0.00-0.00)	0.22 (0.20-0.26)	----
MDQ	0.02 (0.00-0.04)	0.13 (0.07-0.22)	0.46 (0.35-0.54)	----	----	0.00 (0.00-0.08)	0.00 (0.00-0.01)	0.01 (0.00-0.05)	0.38 (0.33-0.45)

A-TAC: Autism-Tics, ADHD, and other Comorbidities inventory; CMRS: Child Mania Rating Scale; MDQ: Mood Disorders Questions
A1, C1, E1: additive genetic, shared environmental, and nonshared environmental influences on autistic traits, and their effect on the
two hypomania measures. Note that shared environment was fixed to zero for autistic traits. A2, C2, E2: genetic and environmental
influences on hypomania at age 15, and their effect on hypomania at age 15. These pathways are distinct from the effects that
hypomania at age 15 shares with autistic traits. A3, C3, E3: genetic and environmental influences on hypomania at age 18, which are
entirely unique to hypomania at age 18.

Table S7 Estimates for Inflexibility

	A1	A2	A3	C1	C2	C3	E1	E2	E3
<i>Female Estimates</i>									
A-TAC ASD	0.58 (0.52-0.62)	-----	-----	-----	-----	-----	0.42 (0.38-0.48)	-----	-----
CMRS	0.06 (0.03-0.09)	0.47 (0.36-0.58)	-----	-----	0.24 (0.15-0.35)	-----	0.00 (0.00-0.00)	0.22 (0.20-0.26)	-----
MDQ	0.05 (0.02-0.08)	0.10 (0.05-0.18)	0.41 (0.22-0.56)	-----	-----	0.10 (0.00-0.26)	0.00 (0.00-0.01)	0.01 (0.00-0.04)	0.32 (0.28-0.38)
<i>Male Estimates</i>									
A-TAC ASD	0.73 (0.70-0.75)	-----	-----	-----	-----	-----	0.27 (0.25-0.30)	-----	-----
CMRS	0.05 (0.03-0.07)	0.49 (0.38-0.59)	-----	-----	0.24 (0.15-0.34)	-----	0.00 (0.00-0.01)	0.22 (0.19-0.25)	-----
MDQ	0.06 (0.03-0.10)	0.11 (0.05-0.20)	0.43 (0.32-0.52)	-----	-----	0.00 (0.00-0.09)	0.00 (0.00-0.00)	0.01 (0.00-0.04)	0.39 (0.33-0.45)

A-TAC: Autism-Tics, ADHD, and other Comorbidities inventory; CMRS: Child Mania Rating Scale; MDQ: Mood Disorders Questions
A1, C1, E1: additive genetic, shared environmental, and nonshared environmental influences on autistic traits, and their effect on the two hypomania measures. Note that shared environment was fixed to zero for autistic traits. A2, C2, E2: genetic and environmental influences on hypomania at age 15, and their effect on hypomania at age 15. These pathways are distinct from the effects that hypomania at age 15 shares with autistic traits. A3, C3, E3: genetic and environmental influences on hypomania at age 18, which are entirely unique to hypomania at age 18.

Table S8 Estimates for Social Communication

	A1	A2	A3	C1	C2	C3	E1	E2	E3
<i>Female Estimates</i>									
A-TAC ASD	0.62 (0.57-0.66)	-----	-----	-----	-----	-----	0.38 (0.34-0.43)	-----	-----
CMRS	0.07 (0.04-0.10)	0.46 (0.34-0.56)	-----	-----	0.25 (0.15-0.35)	-----	0.00 (0.00-0.01)	0.22 (0.20-0.25)	-----
MDQ	0.06 (0.03-0.10)	0.10 (0.05-0.18)	0.39 (0.21-0.54)	-----	-----	0.11 (0.00-0.27)	0.00 (0.00-0.01)	0.01 (0.00-0.04)	0.32 (0.28-0.37)
<i>Male Estimates</i>									
A-TAC ASD	0.74 (0.72-0.77)	-----	-----	-----	-----	-----	0.26 (0.23-0.28)	-----	-----
CMRS	0.07 (0.05-0.10)	0.49 (0.37-0.58)	-----	-----	0.22 (0.14-0.32)	-----	0.00 (0.00-0.00)	0.22 (0.19-0.25)	-----
MDQ	0.05 (0.03-0.09)	0.11 (0.05-0.19)	0.45 (0.34-0.53)	-----	-----	0.00 (0.00-0.08)	0.00 (0.00-0.01)	0.01 (0.00-0.04)	0.38 (0.32-0.45)

A-TAC: Autism-Tics, ADHD, and other Comorbidities inventory; CMRS: Child Mania Rating Scale; MDQ: Mood Disorders Questions
A1, C1, E1: additive genetic, shared environmental, and nonshared environmental influences on autistic traits, and their effect on the two hypomania measures. Note that shared environment was fixed to zero for autistic traits. A2, C2, E2: genetic and environmental influences on hypomania at age 15, and their effect on hypomania at age 15. These pathways are distinct from the effects that hypomania at age 15 shares with autistic traits. A3, C3, E3: genetic and environmental influences on hypomania at age 18, which are entirely unique to hypomania at age 18.

Examining the association between childhood autistic traits and adolescent hypomania: a longitudinal twin study

Mark J. Taylor PhD^{1*}, Angelica Ronald PhD², Joanna Martin PhD^{1,3}, Sebastian Lundström PhD^{4,5}, Georgina M. Hosang PhD⁶, & Paul Lichtenstein¹

¹Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden

²Genes Environment Lifespan Laboratory, Centre for Brain and Cognitive Development, Department of Psychological Science, Birkbeck, University of London, UK

³MRC Centre for Neuropsychiatric Genetics and Genomics, Cardiff University, UK

⁴Gillberg Neuropsychiatry Centre, University of Gothenburg, Sweden

⁵Centre for Ethics, Law and Mental Health, University of Gothenburg, Sweden

⁶Centre for Psychiatry, Wolfson Institute of Preventive Medicine, Barts and the London School of Medicine and Dentistry, Queen Mary, University of London, UK

*Corresponding author: Dr Mark J. Taylor, Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Nobels väg 12A, 171 65 Solna, Sweden (email: mark.taylor@ki.se)

Abstract

Background: There is evidence that autism spectrum disorders (ASDs) co-occur with bipolar disorder (BD) relatively frequently. Individuals with BD often report symptoms of mania and hypomania during adolescence, prior to the age of onset for BD. It is unknown whether these symptoms are associated with ASDs. We examined whether diagnoses of ASDs and autistic traits were associated with hypomania in a large, population-based Swedish twin sample.

Methods: Parental structured interviews assessed autistic traits, and were used to assign screening diagnoses of ASDs, when twins were aged 9 or 12 (N=13,533 pairs). Parents then completed questionnaires assessing hypomania when the twins were aged 15 and 18 (N=3852 pairs at age 15, and 3,013 pairs at age 18). After investigating the phenotypic associations between these measures, we used the classical twin design to test whether genetic and environmental influences on autistic traits influence variation in adolescent hypomania.

Results: Autistic traits and ASD diagnoses in childhood were associated with elevated scores on the measures of adolescent hypomania. Twin analyses indicated that 6-9% of the variance in hypomania was explained by genetic influences that were shared with autistic traits in childhood. When repeating these analyses for specific autistic trait domains, we found a stronger association between social interaction difficulties and hypomania than for other autistic trait domains.

Conclusions: These results indicate a genetic link between autistic traits and hypomania in adolescence. This adds to the growing evidence base of genetic factors associated with ASDs showing links with psychiatric outcomes across childhood and into adulthood.

Introduction

Autism spectrum disorders (ASDs) and bipolar disorder (BD) are highly heritable conditions, with heritability estimates ranging from 46-57% for BD (Song et al., 2018) and as high as 91% for ASDs (Tick et al., 2016). Based on their diagnostic criteria, they have little in common (American Psychiatric Association, 2013), yet there is growing evidence that they co-occur relatively frequently. ASDs and BD each have prevalence rates of approximately 1-2%, yet 3-11% of adults with ASDs have been reported to have BD (Buck et al., 2014; Croen et al., 2015; de Bruin et al., 2007; Hofvander et al., 2009; Verheij et al., 2015), which increases to 25-31% in clinically referred samples (Joshi et al., 2010; Joshi, Wozniak, et al., 2013). Similarly, 30% of individuals with BD have been reported to have ASDs (Joshi et al., 2013). One study of a Japanese clinical sample reported that when ASDs co-occur with mood disorders, in 75% of cases the latter is BD (Munesue et al., 2008), suggesting a particular association between ASDs and BD. A review paper indicated that existing studies of ASDs and BD are based on sample sizes ranging from 35-4,343 individuals, and that up to 21% of individuals with ASDs have been reported to have BD (Skokauskas & Frodl, 2015). This suggests that BD co-occurs relatively commonly with ASDs across the lifespan, thus emphasizing that a better understanding of their association is needed.

Despite this strong phenotypic link, relatively few studies have focused on the association between ASDs and BD, meaning that the reasons behind their association remain poorly understood. One possibility is that the genetic and environmental factors

associated with ASDs also influence BD. There is now robust evidence in ASD research for this hypothesis in relation to other conditions, such as attention-deficit/hyperactivity disorder (ADHD) and anxiety disorders (Hallett et al., 2010; Lundström et al., 2011; Ronald et al., 2008; Taylor et al., 2013). It is, however, less well known whether this applies to ASDs and BD, with only preliminary evidence. Recent common variant genome-wide correlations between these conditions have been estimated at 0.11-0.18, indicating modest overlap between clinical diagnoses of these disorders in adults (Grove et al., 2019; Stahl et al., 2019).

These estimates only account for common genetic variation, thus the full extent of genetic overlap between ASDs and BD is unclear. Twin and family studies have considerable value in understanding genetic overlap between conditions, since they account for all sources of genetic variation. Family studies suggest that siblings of individuals with ASDs are more likely to have BD than siblings of controls, and children of individuals with BD are more likely to have ASDs (Jokiranta-Olkonienmi et al., 2016; Sullivan et al., 2012). In one family study, over 50,000 individuals with BD were identified from the Swedish population; their relatives were diagnosed with ASDs more frequently than relatives of controls and the more closely related individuals were, the more likely they were to have ASDs, suggesting a genetic link between clinical diagnoses of these conditions. In the same study, twin analyses yielded a genetic correlation of 0.24 between these diagnoses (Song et al., 2015).

Nonetheless, genetically informative studies of the association between ASDs and BD remain scarce. Most existing studies report on broad age ranges and focus on clinical

diagnoses (Grove et al., 2019; Song et al., 2015; Stahl et al., 2019). Although BD typically has its onset during adulthood, many individuals with BD report experiencing hypomanic symptoms as teenagers (Perlis et al., 2009). For instance, as many as 10% of adolescents show elevated degrees of hypomania, which may confer increased risk for subsequent BD (Hosang et al., 2017; Hudziak et al., 2005; Waugh et al., 2014). This has led to discussion around whether BD should be considered as a diagnosis in pediatric populations, albeit this has proven to be a contentious issue, with some clinicians remaining hesitant to diagnosis BD in these individuals. Given the vulnerability of individuals with ASDs to mental health difficulties, and the frequent emergence of many persistent mental health problems during adolescence, there is thus considerable value in studying hypomania during adolescence. Whether hypomania is associated with ASDs has not yet been comprehensively examined.

We therefore used a large, longitudinally assessed twin cohort to test whether ASDs were associated with adolescent hypomania. Since there is evidence to suggest that the genetic influences on ASDs also influence autistic traits in the general population, we also examined whether autistic traits in childhood were linked with adolescent hypomania. We then used twin methods to assess whether etiological influences on childhood autistic traits also influence adolescent hypomania. There is evidence to suggest that specific autistic traits, such as communication difficulties and repetitive behavior, are partly distinguishable in terms of etiology (Ronald et al., 2006), and some studies indicate that the strength of the association between autistic traits and other phenotypes differs across these specific domains (Polderman et al., 2014; Ronald et al., 2014; Taylor et al., 2015). We therefore also compared associations with hypomania

across specific autistic trait domains. We expected ASDs (and autistic traits) to be associated with hypomania, both phenotypically and etiologically. We further expected that the magnitude of these associations would differ across specific autistic trait domains.

Method

Participants

When twins are born in Sweden, their parents can voluntarily register them in the Swedish Twin Registry (STR). Based on the STR, all twins born in Sweden since 1992 were invited to participate in CATSS from 2004 when twins were aged 9 (earlier cohorts included 12-year-olds), with an initial response rate of 75% (Anckarsäter et al., 2011). Follow-ups are conducted when the twins are aged 15 or 18. The respective response rates among those who have turned 15 or 18 are 61% and 59%. Supplementary Table S1 shows the numbers of individuals with neurodevelopmental or psychiatric disorders who participated at each wave, and indicates that a greater proportion of participants had at least one of these conditions than those who did not participate. Exclusions were performed for brain injuries (N=194) and chromosomal syndromes (N=27). The sample sizes are shown in Table 1. Zygosity was ascertained using a panel of single nucleotide polymorphisms or an algorithm of twin similarity (Hannellius et al., 2007). Only twins with at least a 95% probability of correct classification have had their zygosity ascertained using the latter method.

CATSS families provide written informed consent prior to participation. CATSS has ethical approval from the Regional Ethical Review Board in Stockholm County (2016/2135-31).

Measures

Autistic Traits: The Autism-Tics, AD/HD, and other Comorbidities inventory (A-TAC) is a 96 item structured interview, completed by parents over the telephone when the twins were aged 9 or 12 (Hansson et al., 2005). ASD is assessed by 17 questions, each of which can be answered 'yes' (scored as 1), 'yes, to some extent' (0.5), or 'no' (0), giving scores ranging from 0-17. Specific autistic traits were measured using three subscales: social interaction difficulties (6 items), communication problems (6 items), and inflexibility (5 items). The total scale was used to assign screening diagnoses of ASD. Scores of 8.5 or more denote a strict screening diagnosis of ASDs, which has sensitivity of 0.71 and specificity of 0.95. A score of 4.5 or more yields a broad screening diagnosis, with specificity of 0.96 and sensitivity of 0.80 (Larson et al., 2010). A longitudinal follow-up study replicated this validity (Larson et al., 2013). The A-TAC was used as a continuous scale and to assign screening diagnoses in the phenotypic analyses, in order to test whether hypomania was associated with different manifestations of ASDs. Due to power concerns, the A-TAC was only used as a continuous scale in the twin analyses.

Clinical Diagnoses of ASDs: CATSS was linked with the Swedish National Patient Register (NPR), which records specialist in-patient and out-patient care given to

residents of Sweden. Clinical diagnoses of ASDs were identified, defined as ICD-10 code F84 (excluding Rett Syndrome, other childhood disintegrative disorders, and overactive disorder associated with intellectual disability and stereotyped movements). These diagnoses have been validated specifically for Stockholm County. These clinical diagnoses were used in the phenotypic analyses, described below.

Hypomania: Two parent report measures assessed hypomania, both of which are among the three most highly recommended instruments for assessing hypomania (Waugh et al., 2014). We relied on parental reports here, as they have been shown to outperform teacher and self-report measures of hypomania in adolescents (Youngstrom et al., 2015). At age 15, the shortened Child Mania Rating Scale (CMRS) was used. The 10 CMRS items are designed to briefly screen for behaviors, thoughts, and emotions characteristic of hypomania and mania, such as irritability, high energy levels, and impulsivity. Questions can be answered 'never' (scored as 0), 'sometimes' (1), 'often' (2), or 'very often' (3), giving a range of scores from 0-30. The shortened CMRS has sensitivity of 0.90 and specificity of 0.96 in detecting individuals with BD, and also distinguishes BD from ADHD with sensitivity of 0.84 and specificity of 0.92. The CMRS was used as a continuous scale in all analyses.

At age 18, parents completed the Mood Disorders Questionnaire (MDQ) (Hirschfeld et al., 2000). The MDQ comprises 13 questions, covering elevated mood, irritability, and high energy levels. A cut-off of seven or more endorsed questions gives sensitivity of 0.73 and specificity of 0.90 in identifying BD (Wagner et al., 2006; Hirschfeld et al., 2000). The MDQ was used as a continuous scale in all analyses.

Data Analyses

Phenotypic Analyses: We first tested whether ASDs showed phenotypic associations with hypomania. Four ASD definitions were used, as described above: autistic traits, broad screening diagnosis, strict screening diagnosis, and clinical diagnosis. These definitions allowed us to test associations between ASDs of differing degrees and hypomania. Regression models tested whether autistic traits and ASDs were associated with hypomania at ages 15 and 18. Linear regressions were implemented as generalized estimating equations (GEEs) with robust standard errors to account for the non-independent sample. Sex and birth year were both adjusted for in the models. Standardized scores were used in these analyses to improve comparability across analyses.

Twin Analyses: We utilized the classical twin design to assess whether genetic and environmental influences on autistic traits at ages 9 or 12 influenced hypomania at ages 15 and 18. The twin design partitions variance in a continuous trait into genetic and environmental components, based on the assumption that monozygotic (MZ) twins share all of their segregating DNA code, compared to approximately 50% in dizygotic (DZ) twins. Higher within-pair resemblance in MZ compared to DZ twins indicates genetic influences on a given trait. Four sources of variance are estimated: additive genetic (termed 'A'), non-additive genetic (D), shared environment (C), and nonshared environment (E), which incorporates measurement error. The multivariate twin model

aims to estimate the degree to which these influences on one trait influence other traits. Initially, twin correlations can be used to determine the degree to which overlap in these components accounts for the covariance between two traits, and involve correlating one twin's score on one trait with their co-twin's score on another trait. Genetic influences on the covariance between these two traits are implied when these correlations are higher in MZ than DZ twins. A more comprehensive overview of the twin design is given elsewhere (Rijsdijk & Sham, 2002).

These twin correlations were initially estimated from a saturated model of the observed data. We then fitted a multivariate Cholesky decomposition to assess the degree to which genetic and environmental influences on autistic traits in childhood influence adolescent hypomania. This estimated the proportion of variance in CMRS and MDQ scores that was explained by genetic and nonshared environmental influences shared with autistic traits, as well as the proportions that were unique to each trait. The MDQ at age 18 was additionally influenced by etiological factors shared with the CMRS at age 15. Only one of C and D was estimated for each measure, as these components confound one another in the classical twin design. Based on previously published univariate results for the measures used here, we allowed A, D, and E to influence autistic traits, and A, C, and E to influence hypomania (Hosang et al., 2019; Taylor et al., 2018). No covariance was estimated for D and C. Furthermore, we have previously detected sibling contrast effects for autistic traits (Taylor et al., 2018), which refer to the influence of one twin's phenotype on their co-twin's phenotype; these effects were thus estimated here for autistic traits. All estimates were initially permitted to differ in magnitude by sex (*quantitative sex limitation*), and were subsequently equated. We then

fitted nested models from whichever of these two models fit best to test the significance of individual groups of parameters. Model fit was assessed using Bayesian Information Criteria (BIC), which has been shown to outperform other fit indices when fitting multivariate twin models to large samples (Markon & Krueger, 2004). Lower BIC values suggest better fitting models, and so the best fitting model was defined by having the lowest BIC value. After performing analyses on the full A-TAC scale, we repeated all twin analyses on the three subscales.

All of the twin analyses were performed using the OpenMx package of R (Neale et al., 2016). Full information maximum-likelihood estimation was used in the analyses, an estimation method that is robust to missing data.

Results

The sample size at each age, descriptive statistics, and the number of individuals with each ASD diagnosis are shown in Table 1.

Phenotypic Analyses

Figure 2 shows the associations between ASDs and hypomania. All ASD definitions were associated with increased hypomania scores at ages 15 and 18. Each standard deviation increase in autistic traits was associated with increases in hypomania of 0.31 and 0.24 standard deviations at ages 15 and 18 respectively. Diagnoses of ASD were

associated with stronger increases in hypomania; for broad ASD, these were 1.19 and 0.84 standard deviations at ages 15 and 18 respectively; for strict ASD, 1.29 and 1.66; and for clinical ASD, 0.90 and 0.71.

Twin Analyses

Phenotypic and twin correlations are shown in Table 2. Phenotypic correlations between autistic traits and hypomania varied from 0.24-0.28, dependent on sex and age. Cross-trait cross-twin correlations were all higher for MZ than DZ twins, indicating genetic influences on the covariance between autistic traits and hypomania. Table 3 shows the twin model fit statistics. The best fitting model included quantitative sex limitation; D could be dropped for autistic traits. The model estimates are shown in Figure 1b (estimates with confidence intervals are shown in Supplementary Table S1). All of the shared variance between autistic traits and hypomania was genetic; all nonshared environmental pathways from autistic traits to hypomania were estimated at 0. Additive genetic influences explained 61% of the variance in autistic traits in females, and 73% in males. These genetic influences on autistic traits explained 9% of the variance in hypomania at age 15 in females, and 7% in males. At age 18, 7% of the variance in hypomania was explained by genetic influences shared with autistic traits in females, and 6% in males. This means that of the 53-55% heritability in hypomania at age 15, 13-17% was accounted for by genetic influences shared between hypomania and autistic traits. At age 18, 10-13% of the 55-60% heritability of hypomania was accounted for by genetic influences shared with autistic traits.

The results of the subscale-specific analyses are shown in Supplementary Tables S2-S6. The strongest phenotypic correlations were between social interaction difficulties and hypomania (0.25-0.27). Phenotypic correlations between inflexibility and hypomania varied from 0.18-0.21, while the weakest phenotypic correlations were between communication problems and hypomania (0.12-0.19). This was reflected in the twin analyses, where 6-8% of the variance in social interaction difficulties was explained by genetic influences shared with autistic traits, compared to 5-6% for inflexibility and 2-5% for communication problems. Confidence intervals overlapped, however. As a posthoc analysis, we reran these analyses using subscales based on DSM-5 criteria for ASDs, which combined the social and communication subscales into one social communication subscale. The results of these analyses are shown in Supplementary Table S8, and indicated that social communication showed stronger associations with hypomania than inflexibility.

Discussion

To our knowledge, our study is the first to show that ASD diagnoses and autistic traits are associated with hypomania during adolescence, prior to the typical age of onset for BD. Genetic factors associated with autistic traits also play a role in adolescent hypomania, with this association particularly driven by social interaction difficulties characteristic of ASDs. This adds to the growing picture of clinical and genetic overlap between ASDs and psychiatric outcomes, extending it to adolescent hypomania.

The limited number of previous studies on ASDs and BD have shown an association between these conditions at the level of clinical diagnoses, especially in adults (Skokauskas & Frodl, 2015), with a modest degree of overlapping genetic factors underlying this association (Grove et al., 2019; Song et al., 2015; Stahl et al., 2019). Our results extend these findings to show that ASDs, and autistic traits, are associated with hypomania during adolescence. Hypomania during adolescence is a potential risk factor for subsequent onset of bipolar disorder (Perlis et al., 2009). More research is clearly needed in order to elucidate whether hypomania in individuals with ASDs follows a trajectory towards clinically impairing symptoms, yet may be useful for clinicians to be aware of hypomania in adolescents with ASDs, even if they do not yet qualify for a diagnosis of BD.

Our results more broadly concur with evidence to indicate that ASDs, and autistic traits, show an association with multiple psychopathological symptoms in adolescence that may, at their extreme, reflect adult onset psychiatric disorders. As well as being associated with hypomania, ASDs are linked with psychotic experiences in adolescence (Cederlöf et al., 2016; Taylor et al., 2015), and subsequently schizophrenia (Joshi et al., 2013). There is now robust evidence to suggest that such psychotic experiences are phenotypically and genetically linked with a variety of adverse outcomes, including mental health problems, substance abuse, and suicide (Cederlöf et al., 2017; McGrath et al., 2016). The evidence on whether hypomania predicts subsequent outcomes remains scant, however. If a similar pattern emerges in future research as is currently the case for psychotic experiences, then it will be important for the full constellation of

adolescent symptoms of adult onset psychiatric disorders to be monitored in adolescents with ASDs, to help with screening, rapid diagnosis and treatment of emerging problems.

Our result also indicated that a model which allowed for sex differences in the genetic and environmental influences underlying autistic traits and hypomania was the best fitting model. This model assumed that the same genetic and environmental factors operate across sexes, but allowed their magnitude to differ. However, the degree to which the variance in hypomania was explained by genetic influences on autistic traits was similar in both sexes. A model allowing for sex differences likely fit better than other models due to the lower heritability of autistic traits in females, meaning that our findings do not indicate that there are sex differences in the association between autistic traits and hypomania. Few existing studies have reported on sex differences in the associations between ASDs, autistic traits, and BD. Croen et al. (2015) reported that 13.8% of women with ASDs also had BD, compared to 9.3% of men. The most recent GWAS of ASDs and BD reported genetic correlations between these conditions collapsed across sex. Thus, more research is needed to fully appreciate whether there are sex differences in the association between ASDs and BD, and autistic traits and hypomania.

In terms of genetic research, our results add to the growing picture that ASDs shows genetic overlap with a multitude of psychiatric and other neurodevelopmental phenotypes. A number of twin studies have focused on the genetic overlap between ASDs, ADHD, and internalizing disorders in childhood (Hallett et al., 2010; Lundström et

al., 2011; Ronald et al., 2008; Taylor et al., 2013). Our results suggest that hypomania in the general population is another psychiatric outcome that shows genetic overlap with autistic traits, indicating that the etiological associations between autistic traits and other phenotypes extends to hypomania. ASDs have thus now been linked to a multitude of psychiatric outcomes using twin methods.

Our results have implications for understanding the origins of adult onset psychiatric disorders. For example, it has been hypothesized that schizophrenia may have neurodevelopmental origins (Owen et al., 2011). This neurodevelopmental hypothesis has received somewhat less attention in relation to BD. In a recent study, we reported genetic links between ADHD traits and adolescent hypomania (Hosang et al., 2019a). Together with the present findings on autistic traits, these studies support the notion that genetic factors associated with neurodevelopmental conditions could influence hypomania. It may thus be beneficial for future research to further test a neurodevelopmental framework for BD using other behavioral genetic methods.

At the same time, we note that the magnitude of the associations observed here suggest that adolescent hypomania is not just an extension of existing childhood autistic traits, which also concurs with our existing work on ADHD and hypomania (Hosang et al., 2019a). The associations we observed here appear to be relatively modest, with a large proportion of the variance in hypomania accounted for by genetic factors that were independent of autistic traits. While the observed associations were likely attenuated to a degree by the gap between measurements of autistic traits and hypomania, the results

do emphasize that there is considerable value in studying hypomania in adolescence in its own right.

Further to this, it is also worth highlighting that the associations we observed differed across different autistic trait domains. Different autistic trait domains are partly independent at an etiological level, and we found social interaction difficulties to be more strongly associated with hypomania. This contrasts somewhat with other studies that have examined the association between specific autistic traits and other conditions, where inflexible behavior and communication problems show stronger genetic overlap with internalizing traits and traits of ADHD than social interaction difficulties (Ghirardi et al., 2019; Hallett et al., 2012; Polderman et al., 2013, 2014; Taylor et al., 2013; Taylor et al., 2015). The social interaction difficulties covered by our measure of autistic traits relate to difficulties interacting with peers, behaving in ways that are not expected by peers, and not sharing interests with others. In future, it will be informative for longitudinal studies with data on hypomania and autistic traits collected concurrently across ages to investigate ways in which these behaviors may contribute to the subsequent development of hypomania.

It is important to caveat our findings by pointing out that the genetic links between autistic traits and hypomania reported here do not necessarily mean that the genes that influence variation in autistic traits directly influence variation in hypomania. Two traits could appear genetically linked with one another for reasons other than that they share these genetic factors. It might be that autistic traits and hypomania are genetically linked through other traits that both appear to share genetic influences with. For example, both

have been shown to be associated with ADHD (Hosang et al., 2019; Ronald et al., 2008a; Taylor, et al., 2015). It will be important for future studies to employ sophisticated multivariate models to examine whether the association between autistic traits and hypomania is independent of such associations. Furthermore, other confounding factors that we did not measure here could also have influenced the association between autistic traits and hypomania, such as current psychotropic medication use and drug use. It will thus be important for future research to further consider whether these factors contribute to the association between autistic traits and hypomania.

There were several strengths and limitations to our study. Our large, longitudinal sample, with follow-ups from middle childhood to emerging adulthood, enabled us to perform the first twin study of autistic traits and hypomania. This lays the foundation for subsequent studies to further understand the development of BD in individuals with ASDs. Our data collection only started at age 9, however. Data on autistic traits, as well as hypomania, at even earlier ages would have allowed for a more comprehensive overview of the association between these phenotypes. One potential challenge regards whether hypomania in adolescence is a valid construct, or simply reflects ADHD. However, we have previously shown that hypomania in adolescence is separable genetically and phenotypically from traits of ADHD, and that their association also extends to inattention, which is phenotypically more distinguishable from hypomania than hyperactivity and impulsivity (Hosang et al., 2019a). We are thus confident that our findings reflect an association between autistic traits and hypomania, rather than being driven by the established strong association between ASDs and ADHD. However, we are unable to rule out the possibility that irritability associated with ASDs drove our

findings. Irritability is commonly reported among individuals with ASDs (Mayes et al., 2011), and was not measured here. Data were collected from twins, which may raise questions about generalizability, albeit there is evidence to suggest that autistic traits and BD are not elevated in twins relative to singletons (Curran et al., 2011; Kläning et al., 2004). Finally, the Cholesky decompositions fit the observed data poorly according to the likelihood-ratio test; it is worth noting, however our models fit well according to an index better suited to multivariate models in larger samples (Markon & Krueger, 2004). Finally, our study considers one hypothesis of the association between ASDs and BD; it will be informative for future studies to consider alternative explanations, such as phenotypic causality.

It is finally worth noting that there are substantial controversies surrounding BD in younger populations, with long-held clinical consensus positing that BD is an adult-onset disorder. This appears particularly so for BD in prepubescent individuals (Duffy et al., 2020). In this study, we focus on an adolescent sample, and thus our results do not address the controversial issue of BD in younger children. Nonetheless, we acknowledge that research on the clinical meaning of adolescent hypomania remains at a highly preliminary stage. While there is emergent evidence to indicate that many individuals with BD report having experienced symptoms earlier in adolescence (Perlis et al., 2008), relatively few studies have examined the nature of hypomania symptoms in large-scale, representative samples. In some studies, the prevalence of elevated hypomania symptoms is substantially higher than that of BD (Hosang et al., 2017), indicating that for many individuals these are likely transient symptoms. Thus, our

findings should be interpreted in the context of the preliminary nature of research on adolescent hypomania.

This study shows that both ASDs and autistic traits in childhood are associated with hypomania as early as adolescence, prior to the typical age of onset for BD. A certain degree of shared genetic liability underlies this association. ASDs and hypomania can thus be expected to co-occur with one another. Future studies need to consider whether this pattern extends beyond adolescence, and whether elevated levels of hypomania put individuals with ASDs at a greater risk for mental health outcomes in adulthood. The results further provide new insights into the shared genetic architecture of different neurodevelopmental and psychiatric phenotypes.

Acknowledgements

We thank the participants in the Child and Adolescent Twin Study in Sweden (CATSS) for their ongoing participation.

Financial Support

CATSS is funded by the Swedish Council for Health, Working Life and Welfare (project number 2012-1678) and the Swedish Research Council (project number 2016-01989). We acknowledge financial support from the Swedish Research Council through the Swedish Initiative for Research on Microdata in the Social and Medical Sciences (SIMSAM) framework (grant number 340-2013-5867).

Conflicts of Interest

Conflicts of interest: none.

Ethical Standards

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

References

- American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders*. American Psychiatric Association.
<https://doi.org/10.1176/appi.books.9780890425596>
- Anckarsäter, H., Lundström, S., Kollberg, L., Kerekes, N., Palm, C., Carlström, E., Långström, N., Magnusson, P. K. E., Halldner, L., Bölte, S., Gillberg, C., Gumpert, C., Råstam, M., & Lichtenstein, P. (2011). The Child and Adolescent Twin Study in Sweden (CATSS). *Twin Research and Human Genetics*, 14(06), 495–508.
<https://doi.org/10.1375/twin.14.6.495>
- Buck, T. R., Viskochil, J., Farley, M., Coon, H., McMahon, W. M., Morgan, J., & Bilder, D. A. (2014). Psychiatric Comorbidity and Medication Use in Adults with Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*, 44(12), 3063–3071. <https://doi.org/10.1007/s10803-014-2170-2>
- Cederlöf, M., Kuja-Halkola, R., Larsson, H., Sjölander, A., Östberg, P., Lundström, S., Kelleher, I., & Lichtenstein, P. (2017). A longitudinal study of adolescent psychotic experiences and later development of substance use disorder and suicidal behavior. *Schizophrenia Research*, 181, 13–16.
<https://doi.org/10.1016/J.SCHRES.2016.08.029>
- Cederlöf, M., Pettersson, E., Sariaslan, A., Larsson, H., Östberg, P., Kelleher, I., Långström, N., Gumpert, C. H., Lundström, S., & Lichtenstein, P. (2016). The association between childhood autistic traits and adolescent psychotic experiences

is explained by general neuropsychiatric problems. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, 171(2), 153–159.

<https://doi.org/10.1002/ajmg.b.32386>

Croen, L. A., Zerbo, O., Qian, Y., Massolo, M. L., Rich, S., Sidney, S., & Kripke, C. (2015). The health status of adults on the autism spectrum. *Autism*, 19(7), 814–823.

<https://doi.org/10.1177/1362361315577517>

Curran, S., Dworzynski, K., Happé, F., Ronald, A., Allison, C., Baron-Cohen, S., Brayne, C., & Bolton, P. F. (2011). No major effect of twinning on autistic traits. *Autism Research*, 4(5), 377–382. <https://doi.org/10.1002/aur.207>

de Bruin, E. I., Ferdinand, R. F., Meester, S., de Nijs, P. F. A., & Verheij, F. (2007). High Rates of Psychiatric Co-Morbidity in PDD-NOS. *Journal of Autism and Developmental Disorders*, 37(5), 877–886. <https://doi.org/10.1007/s10803-006-0215-x>

Dineen Wagner, K., A Hirschfeld, R. M., Emslie, G. J., Findling, R. L., Gracious, B. L., & Reed, M. L. (2006). PRESS, INC. FOCUS ON CHILDHOOD AND ADOLESCENT MENTAL HEALTH Validation of the Mood Disorder Questionnaire for Bipolar Disorders in Adolescents. In *J Clin Psychiatry* (Vol. 67, Issue 5). https://www.psychiatrist.com/JCP/article/_layouts/ppp.psych.controls/BinaryViewer.ashx?Article=/jcp/article/Pages/2006/v67n05/v67n0518.aspx&Type=Article

Duffy, A., Carlson, G., Dubicka, B., & Hillegers, M. H. J. (2020). Pre-pubertal bipolar disorder: origins and current status of the controversy. In *International Journal of Bipolar Disorders* (Vol. 8, Issue 1, p. 18). Springer. <https://doi.org/10.1186/s40345->

- Ghirardi, L., Pettersson, E., Taylor, M. J., Freitag, C. M., Franke, B., Asherson, P., Larsson, H., & Kuja-Halkola, R. (2019). Genetic and environmental contribution to the overlap between ADHD and ASD trait dimensions in young adults: a twin study. *Psychological Medicine*, 49(10), 1713–1721.
<https://doi.org/10.1017/S003329171800243X>
- Grove, J., Ripke, S., Als, T. D., Mattheisen, M., Walters, R. K., Won, H., Pallesen, J., Agerbo, E., Andreassen, O. A., Anney, R., Awashti, S., Belliveau, R., Bettella, F., Buxbaum, J. D., Bybjerg-Grauholm, J., Bækvad-Hansen, M., Cerrato, F., Chambert, K., Christensen, J. H., ... Børglum, A. D. (2019). Identification of common genetic risk variants for autism spectrum disorder. *Nature Genetics*, 51(3), 431–444.
<https://doi.org/10.1038/s41588-019-0344-8>
- Hallett, V., Ronald, A., Rijdsdijk, F., & Happé, F. (2010). Association of Autistic-Like and Internalizing Traits During Childhood: A Longitudinal Twin Study. *American Journal of Psychiatry*, 167(7), 809–817. <https://doi.org/10.1176/appi.ajp.2009.09070990>
- Hallett, V., Ronald, A., Rijdsdijk, F., & Happé, F. (2012). Disentangling the Associations Between Autistic-Like and Internalizing Traits: A Community Based Twin Study. *Journal of Abnormal Child Psychology*, 40(5), 815–827.
<https://doi.org/10.1007/s10802-011-9596-1>
- Hannelius, U., Gherman, L., Mäkelä, V. V., Lindstedt, A., Zucchelli, M., Lagerberg, C., Tybring, G., Kere, J., & Lindgren, C. M. (2007). Large-scale zygosity testing using single nucleotide polymorphisms. *Twin Research and Human Genetics*, 10(4), 604–

625. <https://doi.org/10.1375/twin.10.4.604>

Hansson, S. L., Svanström Röjvall, A., Rastam, M., Gillberg, C., Gillberg, C., & Anckarsäter, H. (2005). Psychiatric telephone interview with parents for screening of childhood autism - tics, attention-deficit hyperactivity disorder and other comorbidities (A-TAC): preliminary reliability and validity. *The British Journal of Psychiatry: The Journal of Mental Science*, 187(3), 262–267. <https://doi.org/10.1192/bjp.187.3.262>

Hirschfeld, R. M. A., Williams, J. B. W., Spitzer, R. L., Calabrese, J. R., Flynn, L., Keck, P. E., Lewis, L., McElroy, S. L., Post, R. M., Rapoport, D. J., Russell, J. M., Sachs, G. S., & Zajecka, J. (2000). Development and Validation of a Screening Instrument for Bipolar Spectrum Disorder: The Mood Disorder Questionnaire. *American Journal of Psychiatry*, 157(11), 1873–1875. <https://doi.org/10.1176/appi.ajp.157.11.1873>

Hofvander, B., Delorme, R., Chaste, P., Nydén, A., Wentz, E., Ståhlberg, O., Herbrecht, E., Stopin, A., Anckarsäter, H., Gillberg, C., Råstam, M., & Leboyer, M. (2009). Psychiatric and psychosocial problems in adults with normal-intelligence autism spectrum disorders. *BMC Psychiatry*, 9(1), 35. <https://doi.org/10.1186/1471-244X-9-35>

Hosang, G. M., Cardno, A. G., Freeman, D., & Ronald, A. (2017). Characterization and structure of hypomania in a British nonclinical adolescent sample. *Journal of Affective Disorders*, 207, 228–235. <https://doi.org/10.1016/J.JAD.2016.08.033>

Hosang, G. M., Lichtenstein, P., Ronald, A., Lundström, S., & Taylor, M. J. (2019a). Association of Genetic and Environmental Risks for Attention-Deficit/Hyperactivity

Disorder With Hypomanic Symptoms in Youths. *JAMA Psychiatry*.

<https://doi.org/10.1001/jamapsychiatry.2019.1949>

Hosang, G. M., Lichtenstein, P., Ronald, A., Lundström, S., & Taylor, M. J. (2019b).

Association of Genetic and Environmental Risks for Attention-Deficit/Hyperactivity Disorder with Hypomanic Symptoms in Youths. *JAMA Psychiatry*, 76(11), 1150–

1158. <https://doi.org/10.1001/jamapsychiatry.2019.1949>

Hudziak, J. J., Althoff, R. R., Derks, E. M., Faraone, S. V., & Boomsma, D. I. (2005).

Prevalence and Genetic Architecture of Child Behavior Checklist–Juvenile Bipolar Disorder. *Biological Psychiatry*, 58(7), 562–568.

<https://doi.org/10.1016/J.BIOPSYCH.2005.03.024>

Jokiranta-Olkonien, E., Cheslack-Postava, K., Sucksdorff, D., Suominen, A.,

Gyllenberg, D., Chudal, R., Leivonen, S., Gissler, M., Brown, A. S., & Sourander, A. (2016). Risk of Psychiatric and Neurodevelopmental Disorders Among Siblings of

Probands With Autism Spectrum Disorders. *JAMA Psychiatry*, 73(6), 622.

<https://doi.org/10.1001/jamapsychiatry.2016.0495>

Joshi, G., Biederman, J., Petty, C., Goldin, R. L., Furtak, S. L., & Wozniak, J. (2013).

Examining the Comorbidity of Bipolar Disorder and Autism Spectrum Disorders. *The Journal of Clinical Psychiatry*, 74(06), 578–586.

<https://doi.org/10.4088/JCP.12m07392>

Joshi, G., Petty, C., Wozniak, J., Henin, A., Fried, R., Galdo, M., Kotarski, M., Walls, S.,

& Biederman, J. (2010). The Heavy Burden of Psychiatric Comorbidity in Youth with Autism Spectrum Disorders: A Large Comparative Study of a Psychiatrically

Referred Population. *Journal of Autism and Developmental Disorders*, 40(11), 1361–1370. <https://doi.org/10.1007/s10803-010-0996-9>

Joshi, G., Wozniak, J., Petty, C., Martelon, M. K., Fried, R., Bolfek, A., Kotte, A., Stevens, J., Furtak, S. L., Bourgeois, M., Caruso, J., Caron, A., & Biederman, J. (2013). Psychiatric Comorbidity and Functioning in a Clinically Referred Population of Adults with Autism Spectrum Disorders: A Comparative Study. *Journal of Autism and Developmental Disorders*, 43(6), 1314–1325. <https://doi.org/10.1007/s10803-012-1679-5>

Klänning, U., Laursen, T. M., Licht, R. W., Kyvik, K. O., Skytthe, A., & Mortensen, P. B. (2004). Is the risk of bipolar disorder in twins equal to the risk in singletons? A nationwide register-based study. *Journal of Affective Disorders*, 81(2), 141–145. [https://doi.org/10.1016/S0165-0327\(03\)00167-8](https://doi.org/10.1016/S0165-0327(03)00167-8)

Larson, T., Anckarsäter, H., Gillberg, C., Ståhlberg, O., Carlström, E., Kadesjö, B., Råstam, M., Lichtenstein, P., & Gillberg, C. (2010). The Autism - Tics, AD/HD and other Comorbidities inventory (A-TAC): further validation of a telephone interview for epidemiological research. *BMC Psychiatry*, 10(1), 1. <https://doi.org/10.1186/1471-244X-10-1>

Larson, T., Lundström, S., Nilsson, T., Selinus, E. N., Råstam, M., Lichtenstein, P., Gumpert, C. H., Anckarsäter, H., & Kerekes, N. (2013). Predictive properties of the A-TAC inventory when screening for childhood-onset neurodevelopmental problems in a population-based sample. *BMC Psychiatry*, 13(1), 233. <https://doi.org/10.1186/1471-244X-13-233>

- Lundström, S., Chang, Z., Kerekes, N., Gumpert, C. H., Råstam, M., Gillberg, C., Lichtenstein, P., & Anckarsäter, H. (2011). Autistic-like traits and their association with mental health problems in two nationwide twin cohorts of children and adults. *Psychological Medicine*, 41(11), 2423–2433.
<https://doi.org/10.1017/S0033291711000377>
- Markon, K. E., & Krueger, R. F. (2004). An Empirical Comparison of Information-Theoretic Selection Criteria for Multivariate Behavior Genetic Models. *Behavior Genetics*, 34(6), 593–610. <https://doi.org/10.1007/s10519-004-5587-0>
- Mayes, S. D., Calhoun, S. L., Murray, M. J., Ahuja, M., & Smith, L. A. (2011). Anxiety, depression, and irritability in children with autism relative to other neuropsychiatric disorders and typical development. *Research in Autism Spectrum Disorders*, 5(1), 474–485. <https://doi.org/10.1016/j.rasd.2010.06.012>
- McGrath, J. J., Saha, S., Al-Hamzawi, A., Andrade, L., Benjet, C., Bromet, E. J., Browne, M. O., Caldas de Almeida, J. M., Chiu, W. T., Demyttenaere, K., Fayyad, J., Florescu, S., de Girolamo, G., Gureje, O., Haro, J. M., ten Have, M., Hu, C., Kovess-Masfety, V., Lim, C. C. W., ... Kessler, R. C. (2016). The Bidirectional Associations Between Psychotic Experiences and DSM-IV Mental Disorders. *American Journal of Psychiatry*, 173(10), 997–1006.
<https://doi.org/10.1176/appi.ajp.2016.15101293>
- Munesue, T., Ono, Y., Mutoh, K., Shimoda, K., Nakatani, H., & Kikuchi, M. (2008). High prevalence of bipolar disorder comorbidity in adolescents and young adults with high-functioning autism spectrum disorder: A preliminary study of 44 outpatients.

Journal of Affective Disorders, 111(2–3), 170–175.

<https://doi.org/10.1016/J.JAD.2008.02.015>

Neale, M. C., Hunter, M. D., Pritikin, J. N., Zahery, M., Brick, T. R., Kirkpatrick, R. M., Estabrook, R., Bates, T. C., Maes, H. H., & Boker, S. M. (2016). OpenMx 2.0: Extended Structural Equation and Statistical Modeling. *Psychometrika*, 81(2), 535–549. <https://doi.org/10.1007/s11336-014-9435-8>

Owen, M. J., O'Donovan, M. C., Thapar, A., & Craddock, N. (2011). Neurodevelopmental hypothesis of schizophrenia. *British Journal of Psychiatry*, 198(3), 173–175. <https://doi.org/10.1192/bjp.bp.110.084384>

Perlis, R. H., Dennehy, E. B., Miklowitz, D. J., DelBello, M. P., Ostacher, M., Calabrese, J. R., Ametrano, R. M., Wisniewski, S. R., Bowden, C. L., Thase, M. E., Nierenberg, A. A., & Sachs, G. (2009). Retrospective age at onset of bipolar disorder and outcome during two-year follow-up: results from the STEP-BD study. *Bipolar Disorders*, 11(4), 391–400. <https://doi.org/10.1111/j.1399-5618.2009.00686.x>

Perlis, R. H., Purcell, S., Fagerness, J., Kirby, A., Petryshen, T. L., Fan, J., & Sklar, P. (2008). Family-based association study of lithium-related and other candidate genes in bipolar disorder. *Archives of General Psychiatry*, 65(1), 53–61. <https://doi.org/10.1001/archgenpsychiatry.2007.15>

Polderman, T. J. C., Hoekstra, R. A., Posthuma, D., & Larsson, H. (2014). The co-occurrence of autistic and ADHD dimensions in adults: an etiological study in 17 770 twins. *Translational Psychiatry*, 4(9), e435–e435. <https://doi.org/10.1038/tp.2014.84>

- Polderman, T. J. C., Hoekstra, R. A., Vinkhuyzen, A. A. E., Sullivan, P. F., van der Sluis, S., & Posthuma, D. (2013). Attentional switching forms a genetic link between attention problems and autistic traits in adults. *Psychological Medicine*, 43(9), 1985–1996. <https://doi.org/10.1017/S0033291712002863>
- Rijsdijk, F. V., & Sham, P. C. (2002). Analytic approaches to twin data using structural equation models. *Briefings in Bioinformatics*, 3(2), 119–133. <https://doi.org/10.1093/bib/3.2.119>
- Ronald, A., Happé, F., Bolton, P., Butcher, L., & Price, T. (2006). Genetic heterogeneity between the three components of the autism spectrum: A twin study. *J Am Acad Child Psy*, 45(6).
- Ronald, Angelica, Larsson, H., Anckarsäter, H., & Lichtenstein, P. (2014). Symptoms of autism and ADHD: A swedish twin study examining their overlap. *Journal of Abnormal Psychology*, 123(2), 440–451. <https://doi.org/10.1037/a0036088>
- Ronald, Angelica, Simonoff, E., Kuntsi, J., Asherson, P., & Plomin, R. (2008a). Evidence for overlapping genetic influences on autistic and ADHD behaviours in a community twin sample. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 49(5), 535–542. <https://doi.org/10.1111/j.1469-7610.2007.01857.x>
- Ronald, Angelica, Simonoff, E., Kuntsi, J., Asherson, P., & Plomin, R. (2008b). Evidence for overlapping genetic influences on autistic and ADHD behaviours in a community twin sample. *Journal of Child Psychology and Psychiatry*, 49(5), 535–542. <https://doi.org/10.1111/j.1469-7610.2007.01857.x>
- Skokauskas, N., & Frodl, T. (2015). Overlap between Autism Spectrum Disorder and

Bipolar Affective Disorder. *Psychopathology*, 48(4), 209–216.

<https://doi.org/10.1159/000435787>

Song, J., Bergen, S. E., Kuja-Halkola, R., Larsson, H., Landén, M., & Lichtenstein, P. (2015). Bipolar disorder and its relation to major psychiatric disorders: a family-based study in the Swedish population. *Bipolar Disorders*, 17(2), 184–193.

<https://doi.org/10.1111/bdi.12242>

Song, J., Kuja-Halkola, R., Sjölander, A., Bergen, S. E., Larsson, H., Landén, M., & Lichtenstein, P. (2018). Specificity in Etiology of Subtypes of Bipolar Disorder: Evidence From a Swedish Population-Based Family Study. *Biological Psychiatry*, 84(11), 810–816. <https://doi.org/10.1016/J.BIOPSYCH.2017.11.014>

Stahl, E. A., Breen, G., Forstner, A. J., McQuillin, A., Ripke, S., Trubetskoy, V., Mattheisen, M., Wang, Y., Coleman, J. R. I., Gaspar, H. A., de Leeuw, C. A., Steinberg, S., Pavlides, J. M. W., Trzaskowski, M., Byrne, E. M., Pers, T. H., Holmans, P. A., Richards, A. L., Abbott, L., ... Sklar, P. (2019). Genome-wide association study identifies 30 loci associated with bipolar disorder. *Nature Genetics*, 51(5), 793–803. <https://doi.org/10.1038/s41588-019-0397-8>

Sullivan, P. F., Magnusson, C., Reichenberg, A., Boman, M., Dalman, C., Davidson, M., Fruchter, E., Hultman, C. M., Lundberg, M., Långström, N., Weiser, M., Svensson, A. C., & Lichtenstein, P. (2012). Family History of Schizophrenia and Bipolar Disorder as Risk Factors for Autism. *Archives of General Psychiatry*, 69(11), 1099–1103. <https://doi.org/10.1001/archgenpsychiatry.2012.730>

Taylor, M. J., Charman, T., Robinson, E. B., Plomin, R., Happé, F., Asherson, P., &

- Ronald, A. (2013). Developmental associations between traits of autism spectrum disorder and attention deficit hyperactivity disorder: a genetically informative, longitudinal twin study. *Psychological Medicine*, 43(08), 1735–1746.
<https://doi.org/10.1017/S003329171200253X>
- Taylor, Mark J., Charman, T., & Ronald, A. (2015). Where are the strongest associations between autistic traits and traits of ADHD? evidence from a community-based twin study. *European Child & Adolescent Psychiatry*, 24(9), 1129–1138.
<https://doi.org/10.1007/s00787-014-0666-0>
- Taylor, Mark J., Gustafsson, P., Larsson, H., Gillberg, C., Lundström, S., & Lichtenstein, P. (2018). Examining the Association Between Autistic Traits and Atypical Sensory Reactivity: A Twin Study. *Journal of the American Academy of Child & Adolescent Psychiatry*, 57(2), 96–102.
<https://doi.org/10.1016/J.JAAC.2017.11.019>
- Taylor, Mark J., Robinson, E. B., Happé, F., Bolton, P., Freeman, D., & Ronald, A. (2015). A longitudinal twin study of the association between childhood autistic traits and psychotic experiences in adolescence. *Molecular Autism*, 6(1), 44.
<https://doi.org/10.1186/s13229-015-0037-9>
- Tick, B., Bolton, P., Happé, F., Rutter, M., & Rijsdijk, F. (2016). Heritability of autism spectrum disorders: a meta-analysis of twin studies. *Journal of Child Psychology and Psychiatry*, 57(5), 585–595. <https://doi.org/10.1111/jcpp.12499>
- Verheij, C., Louwerse, A., van der Ende, J., Eussen, M. L. J. M., Van Gool, A. R., Verheij, F., Verhulst, F. C., & Greaves-Lord, K. (2015). The Stability of Comorbid

Psychiatric Disorders: A 7 Year Follow Up of Children with Pervasive Developmental Disorder-Not Otherwise Specified. *Journal of Autism and Developmental Disorders*, 45(12), 3939–3948. <https://doi.org/10.1007/s10803-015-2592-5>

Waugh, M. J., Meyer, T. D., Youngstrom, E. A., & Scott, J. (2014). A review of self-rating instruments to identify young people at risk of bipolar spectrum disorders. *Journal of Affective Disorders*, 160, 113–121. <https://doi.org/10.1016/J.JAD.2013.12.019>

Youngstrom, E. A., Genzlinger, J. E., Egerton, G. A., & Van Meter, A. R. (2015). Multivariate meta-analysis of the discriminative validity of caregiver, youth, and teacher rating scales for pediatric bipolar disorder: Mother knows best about mania. *Archives of Scientific Psychology*, 3(1), 112–137. <https://doi.org/10.1037/arc0000024>

Table 1 Descriptive statistics

	Overall	MZF	DZF	MZM	DZM	DZOS, F	DZOS, M
<i>Sample Sizes</i>							
N Pairs, CATSS-9/12	13533	2031	2231	1920	2572	4779	4779
N Pairs, CATSS-15	3852	643	653	511	656	1362	1362
N Pairs, CATSS-18	3013	579	498	417	495	1024	1024
<i>Continuous Measure Descriptives</i>							
A-TAC ASD (Mean, SD)	0.79 (1.56)	0.56 (1.22)	0.68 (1.37)	0.89 (1.61)	1.03 (1.95)	0.56 (1.18)	0.99 (1.81)
A-TAC Social (Mean, SD)	0.28 (0.63)	0.21 (0.51)	0.27 (0.60)	0.29 (0.61)	0.38 (0.79)	0.21 (0.52)	0.33 (0.70)
A-TAC Language (Mean, SD)	0.25 (0.61)	0.19 (0.50)	0.19 (0.50)	0.34 (0.67)	0.31 (0.70)	0.19 (0.50)	0.29 (0.66)
A-TAC Flexibility (Mean, SD)	0.25 (0.62)	0.16 (0.46)	0.22 (0.56)	0.27 (0.64)	0.34 (0.74)	0.16 (0.45)	0.37 (0.75)
CMRS (Mean, SD)	1.80 (2.43)	1.98 (2.51)	2.04 (2.66)	1.44 (2.03)	1.68 (2.32)	2.00 (2.52)	1.67 (2.32)
MDQ (Mean, SD)	0.93 (1.85)	0.85 (1.59)	0.93 (1.88)	0.86 (1.78)	0.96 (2.05)	1.01 (1.92)	0.99 (1.98)
<i>Diagnosis Frequencies</i>							
A-TAC Broad ASD (N, %)	948 (3.50%)	49 (1.21%)	105 (2.35%)	111 (2.89%)	241 (4.69%)	88 (1.84%)	251 (5.23%)
A-TAC Strict ASD (N, %)	269 (0.99%)	15 (0.37%)	28 (0.63%)	32 (0.83%)	77 (1.50%)	21 (0.44%)	70 (1.46%)
Diagnosed ASD (N, %)	399 (1.47%)	18 (0.44%)	45 (1.01%)	48 (1.25%)	107 (2.08%)	43 (0.90%)	92 (1.93%)

CATSS: Child and Adolescent Twin Study in Sweden

A-TAC: Autism-Tics, ADHD, and other Comorbidities inventory at age 9 or 12; CMRS: Child Mania Rating Scale at age 15;

MDQ: Mood Disorders Questionnaire at age 18

A-TAC Broad: broad screening diagnosis, based on a score of 4.5 or more

A-TAC Strict: strict screening diagnosis, based on a score of 8.5 or more for ASD

MZF: monozygotic female twins; DZF: dizygotic female twins; MZM: monozygotic male twins; DZM: dizygotic male twins;

DZOS, F: female twins in opposite-sex dizygotic pairs; DZOS, M: male twins in opposite-sex dizygotic pairs.

Table 2 Twin Correlations

Correlation	rPH, F	rPH, M	MZF	DZF	MZM	DZM	DZOS
A-TAC ASD -> CMRS	0.28 (0.24-0.31)	0.28 (0.24-0.31)	0.25 (0.20-0.29)	0.13 (0.08-0.18)	0.24 (0.20-0.28)	0.18 (0.12-0.23)	0.18 (0.14-0.22)
A-TAC ASD -> MDQ	0.25 (0.20-0.29)	0.24 (0.19-0.28)	0.21 (0.16-0.26)	0.12 (0.05-0.18)	0.21 (0.15-0.27)	0.14 (0.07-0.21)	0.12 (0.07-0.17)
CMRS -> MDQ	0.36 (0.30-0.41)	0.36 (0.30-0.41)	0.29 (0.22-0.35)	0.20 (0.11-0.28)	0.28 (0.21-0.35)	0.23 (0.15-0.31)	0.20 (0.14-0.26)

A-TAC: Autism-Tics, ADHD, and other Comorbidities inventory; CMRS: Child Mania Rating Scale; MDQ: Mood Disorders Questionnaire

rPH, F/ rPH, M: phenotypic correlations, in females and males respectively

MZF: female monozygotic twins; DZF: female dizygotic twins; MZM: male monozygotic twins; DZM: male dizygotic twins; DZOS: opposite-sex dizygotic twins

Table 3 Twin Model Fit Statistics for Multivariate Models

Model	-2LL	Parameters	df	BIC	Comparison Model	$\Delta\chi^2$	Δdf	p
Fully Saturated	101533.48	135	38224	-263048.95	-----	-----	-----	-----
ADCE-s	102062.54	39	38320	-263048.95	Fully Saturated	529.07	96	<0.001
ADCE-s Hom	102629.10	22	38337	-262645.60	ADCE-s	566.55	17	<0.001
<i>ACE-s</i>	<i>102062.54</i>	<i>37</i>	<i>38322</i>	<i>-263069.23</i>	<i>ADCE-s</i>	<i>0.00</i>	<i>2</i>	<i>1.00</i>
ADE-s	102120.94	35	38324	-263029.90	ADCE-s	58.39	4	<0.001
AE-s	102120.94	33	38326	-263048.95	ADCE-s	58.39	6	<0.001
AE	102130.44	31	38328	-263058.51	ADCE-s	67.89	8	<0.001
E	106895.84	19	38340	-258407.44	ADCE-s	4833.3	20	<0.001

The best fitting model is highlighted in bold italics.

A-TAC: Autism-Tics, ADHD, and other Comorbidities inventory; CMRS: Child Mania Rating Scale; MDQ: Mood Disorders Questions

A: additive genetic influences; D: non-additive genetic influences; C: shared environmental influences; E: nonshared environmental influences; s: sibling interaction parameters; Hom: homogeneity, indicating no sex differences.

-2LL: fit statistic, which is $-2 \times \log$ -likelihood of the data; df: degrees of freedom; BIC: Bayesian Information Criteria; $\Delta\chi^2$: -2LL difference between two models, distributed chi-square; Δdf : difference in degrees of freedom between models, equal to the difference in number of parameters.

Figure 1 Association between ASDs and hypomania

The above plot shows the standardized regression coefficients from generalized estimates equations, which reflect the mean increase in hypomania at ages 15 and 18 associated with ASD and autistic traits.

Autistic Traits: continuous score on the Autism-Tics, ADHD, and other Comorbidities inventory (A-TAC)

Broad ASD: screening diagnosis of ASD, based on a score of at least 4.5 on the A-TAC

Strict ASD: screening diagnosis of ASD, based on a score of at least 8.5 on the A-TAC

Diagnosed ASD: diagnosis of ASD recorded in the National Patient Register

Error bars depict 95% confidence intervals.

Figure 2 Cholesky decomposition of autistic traits and adolescent hypomania

A: additive genetic influences; C: shared environmental influences; E: nonshared environmental influences. All path estimates are shown as the proportion of variance explained in each phenotype by that path. Numbers in parentheses are 95% confidence intervals. Dashed lines indicate paths that are not statistically significant, based on 95% confidence intervals overlapping with 0.