

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/140636/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Shenderovich, Yulia , Cluver, Lucie, Eisner, Manuel and Murray, Aja Louise 2020. Moderators of treatment effects in a child maltreatment prevention programme in South Africa. *Child Abuse and Neglect* 106 , 104519. 10.1016/j.chiabu.2020.104519

Publishers page: <https://doi.org/10.1016/j.chiabu.2020.104519>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Moderators of Treatment Effects in a Child Maltreatment Prevention Programme  
in South Africa

Yulia Shenderovich,<sup>a, b</sup> Lucie Cluver,<sup>a, c</sup> Manuel Eisner,<sup>b, d</sup> Aja Louise Murray<sup>b, e</sup>

<sup>a</sup> Department of Social Policy and Intervention, Centre for Evidence-Based Intervention,  
University of Oxford, Barnett House, 32 Wellington Square, OX1 2ER, Oxford, United  
Kingdom;

<sup>b</sup> Institute of Criminology, University of Cambridge, Sidgwick Avenue, CB3 9DA,  
Cambridge, United Kingdom;

<sup>c</sup> Department of Psychiatry and Mental Health, University of Cape Town, J-Block, Groote  
Schoor Hospital, Observatory, 7925, Cape Town, South Africa;

<sup>d</sup> Jacobs Center for Productive Youth Development, University of Zurich, Andreasstrasse 15,  
8050, Zürich, Switzerland;

<sup>e</sup> Department of Psychology, University of Edinburgh, Room F16, Psychology Building, 7  
George Square, Edinburgh EH8 9JZ, United Kingdom.

Corresponding author: Yulia Shenderovich ([y.shenderovich@gmail.com](mailto:y.shenderovich@gmail.com);

[yulia.shenderovich@spi.ox.ac.uk](mailto:yulia.shenderovich@spi.ox.ac.uk)); telephone: +44(0)7774955977

## Moderators of Treatment Effects in a Child Maltreatment Prevention Programme in South Africa

**Background.** Previous research has found mixed results on whether the most disadvantaged families benefit as much as less disadvantaged families from parenting interventions designed to reduce child maltreatment, and little is known in low-income settings.

**Objective.** In this study, we test the effects of child, caregiver, household, and community characteristics as treatment moderators of intervention outcomes – child maltreatment and parenting practices. We test characteristics previously examined elsewhere as well as factors relevant to the South African context.

**Participants and Setting.** This analysis includes adolescents (ages 10-18) and their caregivers ( $N=552$  pairs) who participated in a randomised trial of a parenting programme in the Eastern Cape Province of South Africa.

**Methods.** Data from the caregiver and adolescent standardised questionnaires collected at baseline, post-test (1-month post-intervention), and follow-up (5-9 months) were analysed using longitudinal multilevel analyses. We tested seven hypothesised moderators for each of the primary outcomes through interactions of treatment effect with baseline moderators.

**Results.** No moderator effects were statistically significant after correcting for multiple comparisons testing. Hence, in line with several recent studies examining moderation effects in parenting programmes, our study suggests that parenting interventions aiming to reduce child maltreatment and promote parenting skills in low- and middle-income countries may be similarly effective for families facing various levels of economic, social, and health risk factors.

**Conclusions.** It may be useful to explicitly power trials for testing moderator effects, study different types of moderators and use person-centred analyses to further understand variations in treatment effects.

*Keywords:* Parenting, child abuse, maltreatment, adolescents, treatment moderator.

Child maltreatment has profound life-long effects on young people's health and wellbeing. Parenting interventions are a promising approach to improving parenting skills and reducing child maltreatment by caregivers (Barlow, Smailagic, Huband, Roloff, & Bennett, 2014; Mikton & Butchart, 2009; Vlahovicova, Melendez-Torres, Leijten, Knerr, & Gardner, 2017). Parenting interventions can also target a range of other outcomes, such as child conduct problems and parental mental health (Chen & Chan, 2016). Although most evaluations have been conducted in high-income countries (HICs), there are multiple rigorous evaluations of parenting interventions that have been implemented in low- and middle-income countries (LMICs) that suggest positive effects (Gardner, Montgomery, & Knerr, 2015; Knerr, Gardner, & Cluver, 2013). International agencies, such as the World Health Organization, have recommended the use of parenting interventions worldwide as a means towards reduction in violence against children (Butchart & Mikton, 2014).

### **Variation in Treatment Effects in Parenting Programmes**

Most research examining variation in the effects of parenting programmes has focused on programmes addressing child behaviour problems. Only a few studies have examined treatment moderators in relation to maltreatment or harsh parenting (e.g., Puffer et al., 2015). However, there are crucial similarities between programmes focusing primarily on changing child behaviour and preventing child maltreatment. Both tend to draw on social learning theory to teach skills that reduce child and parent aggression, and strengthen caregiver-child relationships (Knerr et al., 2013). Therefore, research on moderators from programmes targeting child behaviour is highly relevant to maltreatment reduction.

While, on average, many parenting programmes are effective for a number of outcomes, they are not equally effective for everyone (Webster-Stratton, Reid, & Hammond, 2004). It is important to examine whether participating families who are most at risk in relation to the intervention outcomes can benefit from the growing number of parenting

interventions in LMICs. For example, the most disadvantaged families may not be able to attend, or engage in, sessions due to pressing demands on their time and attention, such as providing key material necessities for their children. At the same time, primary studies of risk factors suggest that children living in households facing challenges such as exposure to violence, poverty, and illness may be most at risk regarding child maltreatment (Meinck, Cluver, Boyes, & Mhlongo, 2015; Stith et al., 2009).

Even if disadvantaged families can participate equally in an intervention and receive high-quality programming, it may still be more difficult for these families to implement changes in the household due to diminished health, less stable home environment, or conflicts with other caregivers (Eckenrode et al., 2000; Furlong & McGiloway, 2012). Alternatively, higher-risk families can benefit equally or more than lower-risk participants, as the former may perceive a stronger need for change and have greater room for improvement (Pelham, Dishion, Tein, Shaw, & Wilson, 2017). Researchers have termed these compensatory effects (more beneficial for high-risk groups) and leveraging effects (more beneficial for low-risk groups) (Spoth, Shin, Guyll, Redmond, & Azevedo, 2006).

Treatment moderation can be defined as an interaction between treatment and baseline characteristics of the participants or settings (Kraemer & Wilson, 2013). A 2014 review by Shelleby and Shaw (2014) summarised a number of such within-study moderator analyses examining interventions that target child conduct problems from ages 1 through 10, looking at both child behaviour and parenting outcomes. Several earlier reviews had examined heterogeneity of effects in parenting interventions through meta-analytic moderation analyses, comparing intervention results between groups of studies. In such reviews, authors have categorised parenting intervention evaluations based on their samples, such as high or low socio-economic status, comparing intervention effects across subgroups of studies. However, comparing results between studies has several limitations, such as

potential confounding (Bloch, 2014). For example, interventions conducted in research settings may also include more affluent parents, making it difficult to disentangle the two effects to understand what might be driving different treatment effects between samples (Gardner et al., 2017). Therefore, in reviewing previous literature we focus on studies and reviews that examine moderators as interactions of baseline characteristics and treatment.

Moderators identified in studies in HICs may not have the same effects in LMICs, for instance because of differences in common family structures or greater prevalence of some risk factors, such as absolute poverty (Murray et al., 2018). Alongside programme evaluations, there is an emerging body of evidence on moderators of family interventions in LMICs (Puffer, Annan, Sim, Salhi, & Betancourt, 2017; Puffer et al., 2015).

**Caregiver factors.** Previous research has examined a range of risk factors to assess whether families that face more challenges are able to benefit equally from interventions. A common concern is that caregiver psychosocial risks may prevent behaviour change. For example, mothers' exposure to intimate partner violence (IPV) predicted a decreased effect on reduction of child maltreatment in a study of the Nurse-Family Partnership, a home visitation programme in the United States (US) (Eckenrode et al., 2000). Shelleby and Shaw (2014) review found an equal effectiveness across levels of parental characteristics, such as maternal mental health, history of maltreatment as a child, education, and others, suggesting that variation in parental risk factors did not substantially affect intervention effectiveness.

Several studies that evaluated group-based Incredible Years programme found that the families of caregivers experiencing issues such as marital discord and higher levels of depression benefitted more than other participants from the intervention in respect to child behaviour problem reduction (Beauchaine, Webster-Stratton, & Reid, 2005; Gardner, Hutchings, Bywater, & Whitaker, 2010). The pooled data from 14 RCTs of Incredible Years in Europe did not reveal differences in programme effects on disruptive child behaviour based on

family socioeconomic or ethnic minority status, or baseline parenting practices (Gardner et al., 2017). The findings also suggested some compensatory effects, as the intervention effects were greatest for children with more depressed parents.

Caregiver HIV status, while not previously studied to our knowledge in regard to parenting intervention effects, is an important predictor of parenting and maltreatment outcomes within the South African context (Meinck et al., 2017) and may influence the effects of a child maltreatment programme.

**Child factors.** Given the bidirectional relationship between child and caregiver behaviours, externalising child behaviour is an important correlate of harsh parenting and child maltreatment (Meinck, Cluver, Boyes, & Ndhlovu, 2015; Stith et al., 2009) and a commonly studied moderator of treatment effects in parenting programmes. Shelleby and Shaw (2014) review found that higher levels of child problem behaviour at baseline was, in some studies, associated with greater benefits, while in other studies, there was no association. None of the studies included in the review found a negative effect of higher baseline problems on treatment outcomes. Similarly, studies of Incredible Years in a number of European countries found that children with more severe baseline behaviour problems on average benefitted more (Leijten et al., 2017, 2018).

**Household factors.** Families facing greater economic disadvantage may benefit less from parenting interventions, as they often experience multiple chronic stressors. However, previous research suggests mixed results on the role of economic disadvantage on programme effectiveness (Leijten et al., 2018). Although some reviews comparing study-level results suggested diminished results for economically disadvantaged families (e.g. Lundahl, Risser, & Lovejoy, 2006), several studies have found equal effectiveness (McTaggart & Sanders, 2007; Weeland et al., 2017). Poverty is also a key risk factor for child maltreatment identified in research in Africa (Meinck, Cluver, Boyes, & Ndhlovu, 2015).



**Community factors.** As interventions are delivered in multiple contexts, there is a growing interest in examining community-level characteristics that may affect violence prevention interventions, such as levels of income and violence against children in the community (Morris et al., 2017).

Given the above evidence, the current study explores if the characteristics of participating families and communities affected the intervention effects of Sinovuyo Teen on child maltreatment and parenting behaviours within an RCT in South Africa. This study presents an opportunity to examine moderators studied in HICs, such as poverty, in a LMIC context, as well as explore the markers of disadvantage that have not been examined extensively but are relevant to Southern Africa, such as HIV. The selection of moderators was based on existing literature on treatment moderators, risk factors for maltreatment, and the study setting. Our selection of potential intervention effect moderators was also pre-specified in the trial protocol (Cluver, Meinck, Shenderovich, et al., 2016). In addition to the moderators pre-specified in the protocol, we tested the moderation effects of adolescent externalising behaviour, as well as rural versus peri-urban residence. In South Africa, rural areas tend to have lower levels of public services and income in comparison to peri-urban or urban locations (Coovadia et al. 2009), and therefore location is an important community characteristic. Thus, the moderators examined in this paper include factors at the level of the caregiver (caregiver HIV, history of maltreatment, and exposure to IPV), adolescent (problem behaviour), family (poverty) and household (rural/urban).

## **Methods**

### **Participants and Procedures**

**Study setting.** This study was a cluster-randomised two-arm trial conducted in the Eastern Cape, South Africa. A comprehensive description of the trial can be found in the study protocol (Cluver, Meinck, Shenderovich, et al., 2016). In brief, the treatment arm

received the Sinovuyo Teen, an evidence-informed group-based parenting programme that focused on reducing physical and emotional maltreatment against adolescent children and promoting positive parenting. The control arm received a one-day hygiene information intervention. The trial enrolled 552 families in 32 rural and 8 peri-urban study clusters and took place during April 2015 – August 2016 in disadvantaged isiXhosa-speaking communities with high rates of poverty, unemployment, and HIV. The study clusters were randomised within the urban and rural strata (1:1).

**Screening and recruitment of participants.** In each participating family, the study enrolled one adolescent aged 10 to 18 and their primary caregiver (defined as the person mainly responsible for the adolescent and residing in the same household at least four nights a week). The participants were recruited into the study through local community members, schools, social workers, as well as door-to-door recruitment, focusing on families that experience family conflict and high stress. To be enrolled into the study, both adolescent and caregiver had to complete the baseline assessment. All responses were kept confidential, except in cases of participants requesting assistance, or at risk of significant harm, such as adolescents with recent suicide attempts. Families did not receive monetary incentives for participation, but were given packs with snacks, stationery and toiletries to thank them for participating. Key descriptive statistics of the sample are provided in Table 1.

Table 1. *Sample Baseline Characteristics (N=552)*

<i>Sample characteristic</i>	<i>Mean</i>	<i>SD</i>	<i>Observed range</i>	<i>Possible range</i>
Adolescent age	13.7	2.3	10-18	
Adolescent girls, <i>n</i> (%)	228 (41%)			
Caregiver age	49	15.2	18-92	
Female caregivers, <i>n</i> (%)	524 (95%)			
<i>Moderator variables (baseline)</i>				
Adolescent externalising	18.46	11.68	0-56	0-70
Caregiver childhood maltreatment	0.97	1.38	0-7	0-7
Caregiver IPV	0.40	1.14	0-6	0-6
Household necessities	3.83	2.21	0-8	0-8
Caregiver HIV-positive, <i>n</i> (%)	148 (27%)			

**Intervention.** Sinovuyo Teen is group-based manualised programme based on social learning theory. The programme consists of a series of 14 weekly meetings of 1.5-2 hours each. In addition, when a participant was not able to attend a session, for instance due to illness, the programme included facilitators visiting them at home with a brief recap. It was developed drawing on parent training principles emphasized in existing research, such as modelling positive behaviour and collaborative problem solving (Cluver, Lachman, et al., 2016). During the initial programme development and two pilot studies (Cluver, Lachman, et al., 2016; Cluver, Meinck, Yakubovich, et al., 2016), the intervention was designed to suit the South African context. This study, part of the Parenting for Lifelong Health initiative launched in 2012, is a collaboration between academic researchers and collaborators from the WHO, UNICEF and non-governmental organisations (NGOs) to develop and test, in

randomised trials, a suite of non-commercialised parenting programmes for low-resource settings.

The intervention was delivered by Clowns Without Borders South Africa, a South African non-governmental organisation that focuses on the psychosocial wellbeing of children and communities. Facilitators with various backgrounds, such as childcare, social work, and a range of other diverse backgrounds, delivered the group sessions after receiving a week of training as well as on-going supervision throughout the programme delivery.

Facilitator fidelity to the programme at the group sessions was recorded by a team of trained Research Assistants (RAs) at 99% of all delivered sessions (32% were double-rated). The RAs used an observation tool measuring how well the facilitators implemented the core activities in a session, such as guiding and discussing role plays. Average 83% fidelity was recorded. Overall, caregivers attended on average 50% of group sessions and adolescents, 64%. Together with home-based visits, in total the intervention families received 91% of the sessions either via group sessions or home visits (for more details on intervention delivery, see Shenderovich et al., 2018, 2019).

**Data collection.** Primary caregivers and adolescents completed self-report measures at pre-test, one month post-intervention (92% retention from baseline), and 5-9 months post-intervention (97% retention from baseline). Furthermore, due to the high migration in the area, 52 adolescents were no longer living with their original caregivers at follow-up. Only reports regarding original caregivers were included in the current analysis. Details on sample characteristics as well as the CONSORT diagram are available in other reports (Cluver et al., 2018).

**Informed consent.** Ethical approval for the study was obtained from the Research Ethics Committees at [Blinded University Names], and from the South African Departments

of Social Development and Department of Basic Education. Both adolescents and caregivers needed to complete written informed consent for the dyad to participate.

## **Study Measures**

**Outcome measures.** As previous research on moderators in parenting interventions has sometimes found effects in opposing directions, it has been recommended to examine multiple measures of risk and outcomes (Shelleby & Shaw 2014). Therefore, we examine moderation effects for all intervention primary outcomes, described below. As a pragmatic trial (Porzsolt et al., 2015), the study included two clusters of primary outcomes, one related to maltreatment and one to parenting, comprising a total of 14 constructs, described below.

*Adolescent emotional and physical maltreatment, and neglect by caregivers* within the past month were measured using a culturally-adapted version of the ISPCAN Child Abuse Screening Tool, ICAST-C (Meinck et al., 2018); see Supplementary Materials for examples of all measures. At baseline, in this sample, Cronbach's alpha for caregiver-reported physical and emotional maltreatment scale was .78 (14 items), and for adolescent-reported maltreatment – .86 (12 items). For neglect, the alphas were .63 (3 items) in caregiver report, and .78 (6 items) in adolescent report.

Relevant dimensions of *parenting* (poor parental monitoring and supervision, inconsistent discipline, corporal punishment, positive parenting, and involved parenting) in the past month were measured using the Alabama Parenting Questionnaire (parent and child versions), widely used internationally and previously used in South Africa (Frick, 1991; Lachman, Cluver, Boyes, Kuo, & Casale, 2014). In the caregiver report, Cronbach's alphas were .77 for positive parenting (6 items); .77 for involved parenting (10 items); .75 for poor monitoring (10 items); .70 for corporal punishment (3 items); and .58 for inconsistent discipline (6 items). In the adolescent report, Cronbach's alphas were .87 for positive

parenting (6 items), .87 for involved parenting (10 items), .75 for poor monitoring (10 items), .67 for corporal punishment (3 items) and .67 for inconsistent discipline (6 items).

**Possible treatment moderators.** Drawing on the longitudinal trial design, we examined whether the *baseline values of parenting and maltreatment behaviours* moderated the intervention effects on these measures. That is, we evaluated whether the rate of change due to the intervention was related to the initial level of the outcome measures at baseline.

*Caregiver IPV exposure in the past month* (psychological aggression and physical assault by intimate partners) was measured using modified items from the revised Conflict Tactics Scale (Straus, Hamby, Boney-McCoy, & Sugarman, 1996). Cronbach's alpha was .90 (6 items). A summary variable was created to indicate whether the participant experienced different forms of IPV in the past month.

*Caregiver history of childhood maltreatment* was assessed using an adapted version of the ISPCAN Child Abuse Screening Tool-Retrospective (ICAST-R) (Dunne et al., 2009), measuring the occurrence of abusive physical, sexual and emotional victimisation events before the age of 18. Cronbach's alpha was .71 (7 items).

*Caregiver HIV* was identified using a list of signs and symptoms, developed for areas with over 20% HIV-prevalence (Hosegood, Vanneste, & Timaeus, 2004; Lopman et al., 2006). We considered participants HIV-positive if they self-reported as HIV positive or reported three or more AIDS-related symptoms.

*Adolescent behaviour problems* were measured using a composite score of the rule-breaking and aggression sub-scales from the Child Behaviour Checklist (caregiver report) (Rescorla et al., 2012). Cronbach's alpha was .90 (35 items).

*Economic disadvantage (household necessities)* was measured based on caregiver report on access to eight necessities for children, such as three meals a day and school uniform, in the past month. These commodities were perceived as top necessities by over

80% of South African population in a nationally-representative survey (Pillay, Roberts, & Rule, 2006). Cronbach's alpha was .71 (8 items).

*Rural/peri-urban residence* was recorded at cluster level to reflect if the participant lived in a rural village or a peri-urban township.

### **Intervention Treatment Effects**

Trial outcomes on primary and secondary measures are reported in detail elsewhere (Cluver et al., 2018). In brief, the randomised trial found improvements across a range of parenting and family outcomes. Among the primary outcomes, caregivers reported reduced physical and emotional maltreatment, reduced use of corporal punishment and poor monitoring, as well as an increase in positive and involved parenting. Adolescents reported an increase in involved parenting and a decrease in inconsistent discipline. Overall, adolescents reported fewer programme effects than the caregivers. Chance baseline differences between arms were observed on three outcomes and were accounted for by the difference-in-difference approach used for all analyses.

### **Moderator Analysis Strategy**

To be a moderator of treatment, the variable must be a baseline characteristic, uncorrelated with treatment allocation, and have an interactive effect with treatment on the outcome (Kraemer & Wilson, 2013). Given that the trial included repeated measures, we adopt a longitudinal data analysis approach (Moerbeek & Teerenstra, 2016). Therefore, moderator effects were tested through interactions (Hall & Sammons, 2013), using the following model:

$$\begin{aligned} Parenting_{ti} = & \beta_{00} + \beta_1(Treatment_i) + \beta_2(Moderator_i) + \beta_3(PT_{ti}) + \beta_4(FU_{ti}) \\ & + \beta_5(Treatment_i * PT_{ti}) + \beta_6(Treatment_i * FU_{ti}) \\ & + \beta_7(Treatment_i * Moderator_i) + \beta_8(PT_{ti} * Moderator_i) + \beta_9(FU_{ti} \\ & * Moderator_i) + \beta_{10}(Moderator_i * Treatment_i * PT_{ti}) + \beta_{11}(Moderator_i \\ & * Treatment_i * FU_{ti}) + \alpha_1(Stratification_i) + u_{0i} + \varepsilon_{ti}, (1) \end{aligned}$$

where Level 1 = occasion ( $t$ ), Level 2 = individual( $i$ ),  $PT_{ti}$  is the dummy variable that encodes the contrast between baseline and immediate post-test, and  $FU_{ti}$  is the dummy variable that encodes the contrast between baseline and follow-up.  $Parenting_{ti}$  is the estimated value for parenting at time  $t$  for person  $i$ ,  $Treatment_i$  is the treatment allocation received by individuals in study clusters,  $Moderator_i$  is the candidate moderator that is a baseline characteristic of the individual  $i$ ,  $\beta_{00}$  is the grand intercept, and  $\beta_I$  accounts for the mean differences between treatment and control groups at baseline. The three-way cross-level interactions  $\beta_{10}$  and  $\beta_{11}$  are the key parameters of interest used to assess moderation effects. These interactions compare the effect of the predictors in the intervention group to the effect in the control group, assessing whether the moderator affects the difference in the slopes between the two trial arms. The last two terms represent the time- and person-level residuals. All moderator variables were grand-mean centred. We used a negative binomial link function for maltreatment outcomes to account for their skewed distribution and a linear function for parenting outcomes. The outputs of the negative binomial models are presented as incidence rate ratios (IRR).

Random slopes for individual trajectories over time did not improve model fit and thus, were not included in the final model (Barr, 2013). The models were estimated using maximum likelihood estimation, drawing on all available data. Although this was a cluster randomised trial, due to low ICC at cluster level (under 5%) and a design effect under 2.0, we did not include a separate level for clusters in the moderator analyses (Peugh, 2010). We used robust clustered standard errors to account for any cluster level correlations. Given the large number of comparisons, we used the Benjamini-Hochberg procedure to account for multiple hypothesis testing (Benjamini & Hochberg, 1995), treating all tests as one family. For each outcome, we report the unadjusted  $p$ -values and the corrected  $q$ -values.

Preliminary power calculations suggested that the minimum detectable effect size difference for moderation was approximately  $d = 0.23$  for continuous moderators, and  $d =$



0.50 for binary moderators at participant level, with power of .80, two-tailed test with alpha of .05, non-randomly varying slopes, and intra-class correlation coefficient (ICC) of .07 (Dong & Maynard, 2013). Therefore, the study would not have sufficient statistical power to detect smaller difference in effects due to moderators. Furthermore, these power calculations did not account for the use of non-linear link function and the multiple testing procedures, which further reduce the ability to detect effects (Porter, 2017).

## **Results**

On average, in the intervention arm the incidents of physical and emotional maltreatment reported by caregivers reduced from  $M=8.40$  ( $SD=9.81$ ) incidents at baseline to  $M=3.82$  ( $SD=6.17$ ) at 5-9 months follow-up within the past month. Adolescents in the intervention group reported maltreatment scores of  $M=8.99$  ( $SD=12.56$ ) at baseline and  $M=4.84$  ( $SD=8.52$ ) at follow-up. Within that, there was considerable variation in changes over time reported by intervention participants: the changes in maltreatment behaviours towards adolescents reported by caregivers in the intervention condition from baseline to follow-up varied from a decrease of 56 to an increase of 37 in the maltreatment score. Similarly, intervention adolescents reported a wide range of changes following the intervention, from a decrease of 83 points to an increase of 47. Analysis of average trial outcomes is available in other publications (Cluver et al., 2018).

### **Moderation by Baseline Characteristics**

Results of the moderator analyses are provided in Tables 2 and 3. These results showed that, at both post-test and follow-up, there was not a consistent trend of any of the candidate moderators affecting the treatment effect across the primary outcomes. Although there was no overall consistent pattern of individual characteristics moderating treatment effects, there were several statistically significant moderation effects. In particular, there was an effect of adolescent externalising behaviours on treatment effects for adolescent-reported

involved parenting, indicating that families with higher externalising problems at baseline reported a stronger improvement on caregiver involved parenting at follow-up. There was also a steeper improvement for caregiver-reported parental monitoring at follow-up among caregivers who experienced maltreatment as children. Furthermore, among caregivers who have experienced IPV, these analyses suggested a larger improvement on positive parenting (at post-test in caregiver report) and on parental monitoring (at post-test in adolescent report and at follow-up in caregiver report). Finally, HIV-positive caregivers reported a smaller decline in corporal punishment of their adolescents at follow-up.

We also examined the role of one cluster-level predictor: rural and peri-urban location, finding that participants in rural areas reported greater intervention effects on some outcomes at follow-up. In particular, we found that both adolescents and caregivers in rural clusters reported a larger improvement in positive parenting than their counterparts in peri-urban townships at follow-up. Caregivers in villages also reported a higher level of growth in involved parenting and a stronger decrease in physical and emotional maltreatment at follow-up. Adolescents in rural areas reported a larger decrease in caregiver neglect at follow-up.

### **Moderation by Baseline Outcome Levels**

Generally, programme effects did not vary according to the baseline reports of primary outcomes. However, we did observe that caregiver reports of parental monitoring at baseline appear to have an impact on programme effectiveness on monitoring. In particular, the caregivers who, at baseline, reported the least parental monitoring benefitted from the intervention more in comparison to those who reported better monitoring at baseline. On the other hand, levels of neglect reported by adolescents decreased less for those who reported higher neglect at baseline. Finally, due to low prevalence, it was not possible to estimate the effect of baseline severity on caregiver-reported neglect. Consistent with the results from baseline participant characteristics, analysis of baseline levels of primary outcomes suggested

that participants were able to benefit from the intervention at similar levels regardless of these risks.

Overall, out of the thirteen statically significant moderation effects, eleven were in the direction of greater disadvantage predicting stronger effects. However, after the adjustment for multiple testing, none of the relations between treatment effects and baseline moderators remained significant. Thus, the evidence did not suggest differential impacts based on the risk factors studied.

Table 2. *Moderation at Immediate Post-Test*

Outcome \ Moderator	Caregiver intimate partner violence	Caregiver childhood maltreatment	Caregiver HIV-positive	Baseline outcome level	Adolescent externalising	Household necessities	Rural area
Coef. [95% CI]							
Child							
maltreatment	IRR: 1.02	IRR: 1.06	IRR: 1.31	IRR: 1.00	IRR: 0.99	IRR: 0.97	IRR: 0.57
(adolescent report)	[0.86; 1.21]	[0.87; 1.30]	[0.61; 2.82]	[0.96; 1.03]	[0.97; 1.01]	[0.87; 1.07]	[0.28; 1.15]
<i>a</i>							
p-value	0.815	0.557	0.484	0.879	0.357	0.542	0.116
Adjusted p-value	0.954	0.875	0.875	0.961	0.875	0.875	0.700
Child maltreatment	IRR: 0.90	IRR: 1.16	IRR: 0.81	IRR: 1.00	IRR: 0.99	IRR: 1.05	IRR: 0.80
(caregiver report) <i>a</i>	[0.71; 1.14]	[0.89; 1.51]	[0.40; 1.67]	[0.96; 1.03]	[0.96; 1.03]	[0.88; 1.24]	[0.32; 2.02]
p-value	0.364	0.281	0.574	0.920	0.732	0.595	0.639
Adjusted p-value	0.867	0.875	0.875	0.974	0.914	0.875	0.892
Positive parenting	0.23	0.08	-1.43	-0.05	0.00	0.15	3.00

Outcome \ Moderator	Caregiver intimate partner violence	Caregiver childhood maltreatment	Caregiver HIV-positive	Baseline outcome level	Adolescent externalising	Household necessities	Rural area
(adolescent report)	[-0.98; 1.45]	[-0.65; 0.81]	[-4.07; 1.21]	[-0.22; 0.12]	[-0.09; 0.09]	[-0.45; 0.74]	[-1.14; 6.96]
p-value	0.708	0.832	0.290	0.589	0.993	0.630	0.160
Adjusted p-value	0.910	0.953	0.875	0.875	0.993	0.892	0.790
Positive parenting	0.81*	0.10	-1.13	-0.01	0.02	-0.02	0.27
(caregiver report)	[0.15; 1.47]	[-0.64; 0.83]	[-3.27; 1.02]	[-0.17; 0.14]	[-0.08; 0.11]	[-0.54; 0.50]	[-2.76; 3.31]
p-value	0.016	0.794	0.303	0.865	0.719	0.942	0.861
Adjusted p-value	0.541	0.942	0.875	0.953	0.914	0.974	0.953
Involved parenting	0.61	0.01	-1.05	-0.04	0.05	0.49	4.01
(adolescent report)	[-0.91; 2.14]	[-1.10; 1.12]	[-5.37; 3.27]	[-0.19; 0.11]	[-0.05; 0.15]	[-0.22; 1.19]	[2.84; 10.85]
p-value	0.431	0.983	0.633	0.601	0.335	0.179	0.251
Adjusted p-value	0.867	0.989	0.892	0.875	0.875	0.830	0.875
Involved parenting	0.29	-0.90	-0.15	-0.06	0.02	0.24	3.48
(caregiver report)	[-1.04; 1.62]	[-2.34; 0.53]	[-3.91; 3.60]	[-0.22; 0.11]	[-0.12; 0.16]	[-0.46; 0.95]	[0.62; 7.57]

Outcome \ Moderator	Caregiver intimate partner violence	Caregiver childhood maltreatment	Caregiver HIV-positive	Baseline outcome level	Adolescent externalising	Household necessities	Rural area
p-value	0.668	0.218	0.936	0.496	0.778	0.501	0.096
Adjusted p-value	0.889	0.874	0.974	0.875	0.934	0.875	0.700
Poor parental monitoring (adolescent report)	-1.81* [-2.94; -0.70]	0.41 [-0.3; 1.44]	-2.48 [-5.47; 0.51]	-0.01 [-0.16; 0.15]	-0.03 [-0.15; 0.08]	0.49 [-0.45; 1.43]	-0.28 [-6.36; 5.79]
p-value	0.002	0.440	0.104	0.923	0.606	0.311	0.927
Adjusted p-value	0.153	0.875	0.700	0.974	0.875	0.875	0.974
Poor parental monitoring (caregiver report)	-0.80 [-1.97; 0.36]	-0.50 [-1.68; 0.68]	-1.06 [-4.07; 1.94]	-0.13 [-0.27; 0.00]	0.02 [-0.13; 0.16]	0.30 [-0.48; 1.07]	2.43 [3.01; 7.87]
p-value	0.177	0.407	0.488	0.057	0.807	0.454	0.382
Adjusted p-value	0.856	0.875	0.875	0.581	0.950	0.875	0.875

*Note.* <sup>a</sup> – negative binomial models, results presented as incidence rate ratios. Maltreatment includes physical and emotional violence.

\* $p < 0.05$  for  $p$ -values before the correction for multiple comparisons

Table 3. *Moderation at Follow-Up*

Outcome \ Moderator	Caregiver intimate partner violence	Caregiver childhood maltreatment	Caregiver HIV-positive	Baseline outcome level	Adolescent externalising	Household necessities	Rural area
	Coef. [95% CI]						
Child maltreatment (adolescent report) <sup>a</sup>	IRR: 0.92 [0.72; 1.18]	IRR: 0.95 [0.77; 1.16]	IRR: 1.13 [0.56; 2.30]	IRR: 1.03 [0.99; 1.07]	IRR: 0.98 [0.95; 1.00]	IRR: 0.97 [0.87; 1.07]	IRR: 0.47 [0.19; 1.17]
<i>p-value</i>	0.500	0.603	0.728	0.189	0.067	0.866	0.104
Adjusted <i>p-value</i>	0.867	0.875	0.914	0.851	0.641	0.953	0.700
Child maltreatment (caregiver report) <sup>a</sup>	IRR: 0.83 [0.64; 1.06]	IRR: 0.94 [0.77; 1.15]	IRR: 1.53 [0.86; 2.73]	IRR: 1.00 [0.96; 1.03]	IRR: 0.99 [0.98; 1.05]	IRR: 1.07 [0.92; 1.25]	IRR: 0.52* [0.30; 0.90]
<i>p-value</i>	0.132	0.562	0.146	0.840	0.445	0.393	0.019
Adjusted <i>p-value</i>	0.777	0.875	0.790	0.953	0.875	0.875	0.541
Neglect	IRR : 0.84	IRR : 0.71	IRR : 1.55	IRR: 1.13*	IRR: 0.99	IRR: 1.26	IRR: 0.25*

Outcome \ Moderator	Caregiver intimate partner violence	Caregiver childhood maltreatment	Caregiver HIV-positive	Baseline outcome level	Adolescent externalising	Household necessities	Rural area
(adolescent report) <sup>a</sup>	[0.49; 1.45]	[0.47; 1.07]	[0.46; 5.22]	[1.01 ; 1.27]	[0.95; 1.04]	[0.94; 1.68]	[0.07; 0.97]
<i>p-value</i>	0.531	0.101	0.476	0.041	0.817	0.119	0.045
Adjusted <i>p-value</i>	0.867	0.700	0.875	0.541	0.953	0.700	0.541
Neglect (caregiver report) <sup>a</sup>	IRR: 1.49 [0.74 ; 2.98]	IRR: 1.44 [0.67; 3.09]	IRR: 0.31 [0.03; 2.87]	N/A	IRR: 1.04 [0.94; 1.16]	IRR: 0.86 [0.54; 1.38]	IRR : 0.19 [0.01; 2.63]
<i>p-value</i>	0.266	0.349	0.300		0.463	0.536	0.214
Adjusted <i>p-value</i>	0.867	0.875	0.875		0.875	0.875	0.875
Corporal punishment (adolescent report) <sup>a</sup>	IRR: 0.88 0.73 ; 1.07	IRR: 0.94 [0.77 ; 1.15]	IRR: 1.04 [0.56 ; 1.92]	IRR: 1.05 [0.90 ; 1.22]	IRR: 0.98 [0.95; 1.00]	IRR: 1.08 [0.95 ; 1.23]	IRR: 0.71 [0.44 ; 1.14]
<i>p-value</i>	0.194	0.561	0.901	0.574	0.102	0.248	0.155
Adjusted <i>p-value</i>	0.857	0.875	0.971	0.875	0.700	0.875	0.790



Outcome \ Moderator	Caregiver intimate partner violence	Caregiver childhood maltreatment	Caregiver HIV-positive	Baseline outcome level	Adolescent externalising	Household necessities	Rural area
Corporal punishment (caregiver report) <sup>a</sup>	IRR: 1.01 [0.80 ; 1.29]	IRR: 0.91 [0.75; 1.10]	IRR: 2.25* [1.01 ; 4.99]	IRR: 1.00 [0.90 ; 1.12]	IRR: 0.99 [0.96 ; 1.02]	IRR: 1.04 [-0.91 ; 1.19]	IRR: 1.20 [0.62 ; 2.32]
<i>p-value</i>	0.905	0.328	0.046	0.950	0.565	0.545	0.578
Adjusted <i>p-value</i>	0.974	0.875	0.541	0.976	0.875	0.875	0.875
Positive parenting (adolescent report)	-0.57 [-1.75; 0.62]	0.41 [-0.42; 1.25]	-1.03 [-3.32; 1.26]	-0.06 [-0.21; 0.10]	0.02 [-0.08; 0.12]	0.01 [-0.43; 0.45]	3.00* [0.17; 5.83]
<i>p-value</i>	0.348	0.326	0.377	0.480	0.735	0.965	0.038
Adjusted <i>p-value</i>	0.867	0.875	0.875	0.875	0.914	0.984	0.541
Positive parenting (caregiver report)	0.56 [-0.40; 1.51]	0.20 [-0.54; 0.95]	1.09 [-0.83; 3.02]	-0.13 [-0.29; 0.03]	0.06 [-0.04; 0.16]	-0.24 [-0.77; 0.30]	3.42* [0.27; 6.58]
<i>p-value</i>	0.253	0.593	0.266	0.113	0.227	0.389	0.033
Adjusted <i>p-value</i>	0.867	0.875	0.875	0.700	0.875	0.875	0.541

Outcome \ Moderator	Caregiver intimate partner violence	Caregiver childhood maltreatment	Caregiver HIV-positive	Baseline outcome level	Adolescent externalising	Household necessities	Rural area
Involved parenting (adolescent report)	0.64 [-0.81; 2.08]	0.75 [-0.83; 2.32]	-1.55 [-5.24; 2.14]	0.03 [-0.12; 0.19]	0.14* [0.01; 0.27]	0.12 [-0.68; 0.93]	1.18 [-4.37; 6.72]
<i>p-value</i>	0.387	0.352	0.410	0.679	0.032	0.767	0.677
Adjusted <i>p-value</i>	0.867	0.875	0.875	0.903	0.541	0.931	0.903
Involved parenting (caregiver report)	1.18 [-0.61; 2.98]	-1.14 [-2.27; 0.00]	1.65 [-1.35; 4.65]	-0.11 [-0.30; 0.08]	0.08 [-0.09; 0.25]	0.08 [-0.83; 1.00]	3.45* [0.21; 6.70]
<i>p-value</i>	0.196	0.050	0.281	0.274	0.371	0.858	0.037
Adjusted <i>p-value</i>	0.857	0.546	0.875	0.875	0.875	0.953	0.541
Poor parental monitoring (adolescent report)	-0.41 [-1.61; 0.79]	-0.54 [-1.49; 0.41]	0.85 [-2.30; 4.01]	-0.11 [-0.30; 0.09]	-0.07 [-0.19; 0.05]	0.08 [-0.78; 0.94]	-0.99 [-5.97; 3.98]
<i>p-value</i>	0.501	0.266	0.596	0.280	0.250	0.857	0.696
Adjusted <i>p-value</i>	0.867	0.875	0.875	0.875	0.875	0.953	0.914

Outcome \ Moderator	Caregiver intimate partner violence	Caregiver childhood maltreatment	Caregiver HIV-positive	Baseline outcome level	Adolescent externalising	Household necessities	Rural area
Poor parental monitoring (caregiver report)	-1.41* [-2.60; -0.22]	-0.91* [-1.79; -0.02]	1.46 [-2.06; 4.98]	-0.22* [-0.35; -0.09]	0.01 [-0.13; 0.15]	0.20 [-0.50; 0.91]	0.63 [-2.03; 3.30]
<i>p-value</i>	0.021	0.044	0.415	0.001	0.898	0.580	0.641
Adjusted <i>p-value</i>	0.541	0.541	0.875	0.153	0.971	0.875	0.892
Inconsistent discipline (adolescent report)	0.10 [-0.70 ; 0.89]	-0.16 [-0.84 ; 0.53]	0.55 [-2.06; 3.17]	0.06 [-0.06 ; 0.19]	0.00 [-0.10; 0.10]	0.24 [-0.25; 0.73]	-1.32 [-5.18 ; 2.53]
<i>p-value</i>	0.806	0.652	0.678	0.309	0.938	0.334	0.500
Adjusted <i>p-value</i>	0.954	0.899	0.903	0.875	0.974	0.875	0.875
Inconsistent discipline (caregiver report)	-0.19 [-1.08; 0.70]	-0.33 [-1.17 ; 0.51]	0.35 [-1.87 ; 2.56]	0.08 [-0.10 ; 0.29]	0.06 [-0.02; 0.14]	0.32 [-0.03 ; 0.68]	0.02 [1.85; 1.89]

Outcome \ Moderator	Caregiver intimate partner violence	Caregiver childhood maltreatment	Caregiver HIV-positive	Baseline outcome level	Adolescent externalising	Household necessities	Rural area
<i>p-value</i>	0.680	0.442	0.758	0.333	0.157	0.072	0.981
Adjusted <i>p-value</i>	0.889	0.875	0.928	0.875	0.790	0.648	0.989

*Note.* <sup>a</sup> – negative binomial models, results presented as incidence rate ratios

\* $p < 0.05$  for  $p$ -values before the correction for multiple comparisons

## **Sensitivity Analyses**

As suggested by previous research, it is important to examine the cumulative effect of moderators (Weeland et al., 2017). Therefore, we also examine the pattern of results in models with all moderators included at once, finding a similar pattern of results.

## **Discussion**

Our research set out to examine if the most vulnerable families within an at-risk sample benefitted in a similar way as less vulnerable families from a parenting intervention in the Eastern Cape Province of South Africa in a multi-informant pragmatic randomised evaluation including over 500 families. Prior to the correction for multiple testing, there were several statistically significant treatment moderation effects, suggesting that there may be greater benefits for families who report higher levels of disadvantage for some of the outcomes examined, such as caregiver experience of childhood maltreatment, exposure to IPV, and rural residence. There were, however, two exceptions to this, including smaller intervention effects observed for caregiver-reported corporal punishment among HIV-positive caregivers, and for adolescent-reported neglect for families with higher baseline neglect. However, after adjusting for the testing of multiple hypotheses, none of the baseline risk factors had a statistically significant impact on treatment effects. On balance, therefore, our results suggest that there were no clear detectable differences in the impact of the intervention based on the examined moderator variables, such as baseline household poverty and location, adolescent problem behaviour, caregiver exposure to IPV, their HIV status, and childhood maltreatment.

As parenting research has primarily focused on young children, only a few other studies have explored whether family characteristics influence the treatment effects for adolescents and their caregivers. Our findings within this RCT of a child maltreatment prevention programme align with recent studies and reviews of parenting programmes that

target the reduction of child behavioural problems among families with younger children (Gardner et al., 2015; Pelham et al., 2017; Weeland et al., 2017), as well as programmes addressing parenting and harsh punishment (Annan, Sim, Puffer, Salhi, & Betancourt, 2017; Puffer et al., 2015). Notably, these recent studies differ from some of the earlier findings suggesting reduced parenting programme benefits for disadvantaged families and caregivers (e.g., Eckenrode et al., 2000; Lundahl et al., 2006), whereas the more recent studies have either suggested no reduced benefits for disadvantaged families.

There are many potential reasons for the variations in findings, such as different analytical approaches (Coffman, Edelman, & Woolson, 2016). For example, correction for multiple hypotheses testing has been used in some of the recent moderation studies, which can considerably impact results and conclusions, reducing the number of identified moderators (Shelleby & Shaw, 2014). Variation in analytical approaches may help explain some of the differences in findings, together with the variety of interventions, samples, outcomes, and risk measures used. Researchers have also argued for a more nuanced approach, where variation in families' responses to specific techniques or programme components rather than to the programme overall is examined (Leijten et al., 2018).

The reporting of randomised trials increasingly follows standard guidelines, such as CONSORT. Given the importance of understanding variation in treatment effects, it will be useful for guidelines to also be used for moderator analyses (for example, Van Hoorn et al., 2017). While some literature emphasizes the importance of pre-planning analyses, others also highlight the benefits of exploratory moderator analyses (Kraemer, Frank, & Kupfer, 2006). This is especially relevant for outcomes where risk factors are not well established and it may be beneficial to explicitly modify analyses plans, for instance, based on the hypotheses emerging during data collection (Jamal et al., 2015). Furthermore, in a trial with multiple outcomes, it is not immediately clear how many statistically significant moderation results

constitute a pattern. Researchers could address this by adopting pre-planned decision rules regarding moderators in study protocols (Porter, 2017). It may also be useful to define a minimal effect size for a moderator to be clinically meaningful, and the type of effect – for instance, whether the average effect includes benefits to some and harm to other groups, or just differential effectiveness.

There might also be important effects resulting from a combination of risk factors. Person-centred analyses, such as latent class analysis (Pelham et al., 2017), would be better suited to examining combinations of risk factors. Machine-learning inference can help identify groups that benefit most and least from an intervention using a large number of observed characteristics and their combinations that are not captured via simple interactions (Chernozhukov, Demirer, Duflo, & Fernandez-Val, 2018).

Our study design has allowed us to examine the role of pre-intervention family characteristics in the variation in intervention effects. Having examined all pre-specified primary outcomes, this study did not find consistent evidence of treatment effects moderation by baseline characteristics, adding to a growing body of literature suggesting limited explanatory power of baseline sociodemographic moderators.

The limited variation in treatment effects across families with various levels of risk in our study may be related to the design of the intervention and its implementation which may have helped participants at various risk levels to engage and benefit. For example, intervention families were provided with free meals and transportation, as well as home visits for those who missed group sessions. Like Incredible Years and other similar interventions, this programme has a collaborative working style which can help accommodate the needs of different families (Gardner et al., 2017). The intervention was delivered within a pragmatic randomised trial by lay workers following only a week of training, with ongoing weekly training and supervision. Process data suggest that the intervention quality was comparable to

what was intended (e.g. overall fidelity of 83%). Thus, the programme design, fidelity, and high coverage of group sessions combined with home visits may have helped overcome barriers for some families who might struggle to engage if, for example, a lengthy trip had been required to engage with the programme (Whittaker & Cowley, 2012). However, evaluations of family engagement strategies are still limited (Gonzalez, Morawska, & Haslam, 2018), and further research is crucial to establish which support strategies in fact impact family engagement and outcomes (Lachman et al., 2019). Implementation and programme supports should also be examined in routine settings (Alonge et al., 2019), so we can compare delivery in routine implementation to delivery in research studies, as we might expect that similar implementation is required to achieve comparable effects.

The current study targeted families in a high poverty area, focusing specifically on families with existing challenges around family conflict and high stress. Since targeting individual families or participants has a cost in terms of resources for screening as well as a potential to create stigma, it needs to be managed carefully, and delivering the programme on a more generalised basis can be considered. Further research should examine the effects and costs of universal community-wide programme delivery to all families, for instance in similar high-poverty areas as the area of the current study. Moreover, in ongoing service provision, parenting programmes are often rolled out as part of a package of services, which have broader selection criteria not specifically informed by the parenting component. For example, the current intervention is being delivered in some settings within the USAID-PEFPAR DREAMS intervention package, which includes a parenting component, as part of a range of other strategies to support adolescent girls and young women and reduce the rate of new HIV infections in this group (Gourlay et al., 2019).

Among limitations of the current study, it is important to note that the effects of moderators may have been smaller than could be detected in this trial. Future trials interested



in moderation ideally need to be powered for detecting moderation effects of a specific magnitude, in addition to the main intervention effects (Shieh, 2009). It may also be harder to detect moderation effects within a high-risk sample, such as the current study, where there may be range restriction on some of the relevant moderating characteristics.

Another set of limitations is regarding measurement. Some of the constructs used in our evaluation, in particular inconsistent parenting, had relatively low alpha coefficients. Previous studies using Alabama Parenting Questionnaire in South Africa and other contexts (e.g., Boyes, Cluver, Meinck, Casale, & Newnham, 2019; Dadds, Maujean, & Fraser, 2003; Lachman, Cluver, Boyes, Kuo, & Casale, 2014; Topçuoğlu, Eisner, & Ribeaud, 2015) reported alpha coefficients in the range of 0.6-0.9. This is perhaps due to a small number of items and a broad range of parenting behaviours covered by the subscales, resulting in a content validity-reliability trade-off. As using observational measures with adolescents is challenging, particularly in a low-resource setting, we relied on the parent and child reports of parenting practices, which may be subject to social desirability and recall biases. As long as there is variation in treatment effects among participants, it remains a question whether there are any other risk factors, or combinations of risk factors, that could explain this variation. Understanding the mechanisms behind heterogeneity of treatment effects can be used for targeting and refining interventions as parenting interventions are rolled out to new settings.

Supplementary Table 1. *Overview of measures used in the questionnaires*

Construct	Measure	Sample items (adolescent report)	Sample items (caregiver report)	Response codes
<b>Child maltreatment</b>	ISPCAN Child Abuse	- “In the past month, how often did	- “In the past month, how often	0- zero
	Screening Tool for use in	an adult in your house say that	did you tell your teen that you	1- one
	trials (ICAST-Trial)	they wished you were dead or had	wished he had never been	2- two
	- physical and	never been born?” (emotional)	born?” (emotional)	3- three
	emotional	- “In the past month, how often did	- “In the past month, how often	4- four
	maltreatment	an adult in your house push, grab,	did you push, grab or kick	5- five
		or kick you?” (physical)	your teen?” (physical)	6- six
	- neglect	- “In the past month, were you not	- “In the past month, how many	7- seven
		taken care of when you were sick-	times did you not get medical	8- eight or more
		for example not taken to see a	care for an injury or illness	
		doctor when you were hurt or not	that your teen needed at the	
		given the medicines you needed?”	time even though there was	
			money to pay for it?”	
<b>Parenting</b>	Alabama Parenting	- “In the past month, how often does	- “In the past month, how often	0 – Never
	Questionnaire	your caregiver tell you that he/she	do you tell your teen that you	

<b>Construct</b>	<b>Measure</b>	<b>Sample items (adolescent report)</b>	<b>Sample items (caregiver report)</b>	<b>Response codes</b>
	- positive parenting	likes it when you help around the house?"	like it when he/she helps around the house?"	1 – Almost never 2 – Sometimes
	- involved parenting	- "... you play games or do other fun things with your caregiver?"	- "...you play games or do other fun things with your teen?"	3 – Often 4 – Always
	- poor monitoring	- "... you stay out in the evening past the time you are supposed to be home"	- "... your teen stays out in the evening past the time your teen is supposed to be home"	
	- corporal punishment	- "... your teen is not punished when your teen has done something wrong"	- "... your teen talks you out of being punished after your teen has done something wrong"	
	- inconsistent discipline	- "... your caregiver slaps you when you have done something wrong."	- "... you slap your teen when your teen has done something wrong."	
<b>Household necessities</b>	South African Social Attitudes survey	Not used in the current analysis	"Please tick if you could afford 3 meals a day in the past month"	0 – No 1 – Yes

Construct	Measure	Sample items (adolescent report)	Sample items (caregiver report)	Response codes
<b>Adolescent behaviour problems</b>	Child Behaviour	Not used in the current analysis	“Your teen drinks alcohol or uses drugs without your approval”	0- Not True
	Checklist			1- Sometimes
	- rule-breaking and			True
	aggression			2 -Very True
<b>Caregiver IPV exposure</b>	Conflict Tactics Scale	N/A	“In the past month, my partner twisted	0- Never
			my arm, my hair or threw something	1- Once or twice
			at me that could hurt.”	2- 3-5 times
				3- More than 5 times
<b>Caregiver history of maltreatment</b>	ISPCAN Child Abuse	N/A	“When you were growing up (before you were 18 years old) did anyone ever hit, punch, kick, beat or shake you very hard so that it hurt you?”	0 – No
	Screening Tool- Retrospective			1 – Yes
<b>Caregiver HIV</b>	Self-report as HIV positive or reported three			

Construct	Measure	Sample items (adolescent report)	Sample items (caregiver report)	Response codes
	or more AIDS-related symptoms			
<b>Rural/peri-urban residence</b>	Determined by the research team based on participant's residence			

Supplementary Table 2. *Intra-class Correlation Coefficients for Study Outcomes*

Primary study outcomes	ICC at person level	ICC at cluster level
Physical and emotional maltreatment (adolescent report)	29%	0.35%
Physical and emotional maltreatment (caregiver report)	15%	2%
Neglect (adolescent report)	22%	0%
Neglect (caregiver report)	3%	3%
Corporal punishment (adolescent report)	7%	0%
Corporal punishment (caregiver report)	10%	0%
Positive parenting (adolescent report)	25%	1%
Positive parenting (caregiver report)	22%	5%
Involved parenting (adolescent report)	27%	3%
Involved parenting (caregiver report)	25%	4%
Poor monitoring (adolescent report)	22%	3%
Poor monitoring (caregiver report)	22%	5%
Inconsistent discipline (adolescent report)	1%	1%
Inconsistent discipline (caregiver report)	8%	1%

## References

- Alonge, O., Rodriguez, D. C., Brandes, N., Geng, E., Reveiz, L., & Peters, D. H. (2019). How is implementation research applied to advance health in low-income and middle-income countries. *BMJ Global Health*, 4(2), 1–10. <https://doi.org/10.1136/bmjgh-2018-001257>
- Annan, J., Sim, A., Puffer, E. S., Salhi, C., & Betancourt, T. S. (2017). Improving mental health outcomes of Burmese migrant and displaced children in Thailand: A community-based randomized controlled trial of a parenting and family skills intervention. *Prevention Science*, 18(7), 793–803. <https://doi.org/10.1007/s11121-016-0728-2>
- Barlow, J., Smailagic, N., Huband, N., Roloff, V., & Bennett, C. (2014). Group-based parent training programmes for improving parental psychosocial health. *The Cochrane Library*.
- Barr, D. J. (2013). Random effects structure for testing interactions in linear mixed-effects models. *Frontiers in Psychology*, 4, 328. <https://doi.org/10.3389/fpsyg.2013.00328>
- Beauchaine, T. P., Webster-Stratton, C., & Reid, M. J. (2005). Mediators, moderators, and predictors of 1-year outcomes among children treated for early-onset conduct problems: a latent growth curve analysis. *Journal of Consulting and Clinical Psychology*, 73(3), 371–388. <https://doi.org/10.1037/0022-006X.73.3.371>
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society*, 57, 289–300. <https://doi.org/10.2307/2346101>
- Bloch, M. H. (2014). Meta-analysis and moderator analysis: Can the field develop further? *Journal of the American Academy of Child and Adolescent Psychiatry*, 53(2), 135–137. <https://doi.org/10.1016/j.jaac.2013.12.001>
- Boyes, M. E., Cluver, L. D., Meinck, F., Casale, M., & Newnham, E. (2019). Mental health in South African adolescents living with HIV: correlates of internalising and

- externalising symptoms. *AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV*, 31(1), 95–104. <https://doi.org/10.1080/09540121.2018.1524121>
- Butchart, A., & Mikton, C. R. (2014). *Global status report on violence prevention. World Health Organization*. <https://doi.org/10.1007/s13398-014-0173-7.2>
- Chen, M., & Chan, K. L. (2016). Effects of parenting programs on child maltreatment prevention a meta-analysis. *Trauma, Violence, & Abuse*, 17(1), 88–104. <https://doi.org/10.1177/1524838014566718>
- Chernozhukov, V., Demirer, M., Duflo, E., & Fernandez-Val, I. (2018). *Generic machine learning inference on heterogenous treatment effects in randomized experiments* (No. 1712.04802). <https://doi.org/10.3386/w24678>
- Cluver, L. D., Lachman, J. M., Ward, C. L., Gardner, F., Peterson, T., Hutchings, J. M., ... Redfern, A. A. (2016). Development of a parenting support program to prevent abuse of adolescents in South Africa: Findings from a pilot pre-post study. *Research on Social Work Practice*, 27(7), 758–766. <https://doi.org/10.1177/1049731516628647>
- Cluver, L. D., Meinck, F., Shenderovich, Y., Ward, C. L., Romero, R. H., Redfern, A., ... Lechowicz, M. (2016). A parenting programme to prevent abuse of adolescents in South Africa: Study protocol for a randomised controlled trial. *Trials*, 17(328), 1–10. <https://doi.org/10.1186/s13063-016-1452-8>
- Cluver, L. D., Meinck, F., Steinert, J., Shenderovich, Y., Doubt, J., Herrero-Romero, R., ... Gardner, F. (2018). Parenting for Lifelong Health: A pragmatic cluster randomised controlled trial of a non-commercialised parenting programme for adolescents and their families in South Africa. *BMJ Global Health*, 3(1), 1–16. <https://doi.org/doi:10.1136/bmjgh-2017-000539>
- Cluver, L. D., Meinck, F., Yakubovich, A., Doubt, J., Redfern, A., Ward, C., ... Gardner, F. (2016). Reducing child abuse amongst adolescents in low- and middle-income countries:



- A pre-post trial in South Africa. *BMC Public Health*, 16(567), 1–11.  
<https://doi.org/10.1186/s12889-016-3262-z>
- Coffman, C. J., Edelman, D., & Woolson, R. F. (2016). To condition or not condition? Analysing ‘change’ in longitudinal randomised controlled trials. *BMJ Open*, 6(12), e013096. <https://doi.org/10.1136/bmjopen-2016-013096>
- Dadds, M. R., Maujean, A., & Fraser, J. A. (2003). Parenting and Conduct Problems in Children: Australian Data and Psychometric Properties of the Alabama Parenting Questionnaire. *Australian Psychologist*, 38(3), 238–241.  
<https://doi.org/10.1080/00050060310001707267>
- Davis, J. M. V., & Heller, S. (2017). *Rethinking the benefits of youth employment programs: The heterogeneous effects of summer jobs* (No. 23443). *NBER Working Paper Series*.  
<https://doi.org/10.3386/w23443>
- Dong, N., & Maynard, R. (2013). PowerUp!: A tool for calculating minimum detectable effect sizes and minimum required sample sizes for experimental and quasi-experimental design studies. *Journal of Research on Educational Effectiveness*, 6(1), 24–67.
- Dunne, M. P., Zolotor, A. J., Runyan, D. K., Andrevia-Miller, I., Choo, W. Y., Dunne, S. K., ... Youssef, R. (2009). ISPCAN Child Abuse Screening Tools Retrospective version (ICAST-R): Delphi study and field testing in seven countries. *Child Abuse & Neglect*, 33(11), 815–825. <https://doi.org/10.1016/j.chiabu.2009.09.005>
- Eckenrode, J., Ganzel, B., Henderson, C. R., Smith, E., Olds, D. L., Powers, J., ... Sidora, K. (2000). Preventing child abuse and neglect with a program of nurse home visitation. *JAMA*, 284(11), 1385–1391. <https://doi.org/10.1001/jama.284.11.1385>
- Frick, P. J. (1991). Alabama Parenting Questionnaire. *Unpublished Rating Scale, University of Alabama*.

- Furlong, M., & McGilloway, S. (2012). The Incredible Years Parenting program in Ireland: A qualitative analysis of the experience of disadvantaged parents. *Clinical Child Psychology and Psychiatry*, 17(4), 616–630. <https://doi.org/10.1177/1359104511426406>
- Gardner, F., Hutchings, J., Bywater, T., & Whitaker, C. (2010). Who benefits and how does it work? Moderators and mediators of outcome in an effectiveness trial of a parenting intervention. *Journal of Clinical Child & Adolescent Psychology*, 39(4), 568–580. <https://doi.org/10.1080/15374416.2010.486315>
- Gardner, F., Leijten, P., Mann, J., Landau, S., Harris, V., Beecham, J., ... Scott, S. (2017). Could scale-up of parenting programmes improve child disruptive behaviour and reduce social inequalities? Using individual participant data meta-analysis to establish for whom programmes are effective and cost-effective. *Public Health Research*, 5(10). <https://doi.org/10.3310/phr05XXX>
- Gardner, F., Montgomery, P., & Knerr, W. (2015). Transporting evidence-based parenting programs for child problem behavior (age 3–10) between countries: Systematic review and meta-analysis. *Journal of Clinical Child & Adolescent Psychology*, 45, 749–762. <https://doi.org/10.1080/15374416.2015.1015134>
- Gonzalez, C., Morawska, A., & Haslam, D. M. (2018). Enhancing Initial Parental Engagement in Interventions for Parents of Young Children: A Systematic Review of Experimental Studies. *Clinical Child and Family Psychology Review*, 21(3), 415–432. <https://doi.org/10.1007/s10567-018-0259-4>
- Gourlay, A., Birdthistle, I., Mthiyane, N. T., Orindi, B. O., Muuo, S., Kwaro, D., ... Floyd, S. (2019). Awareness and uptake of layered HIV prevention programming for young women: analysis of population-based surveys in three DREAMS settings in Kenya and South Africa. *BMC Public Health*, 19(1), 1417. <https://doi.org/10.1186/s12889-019-7766-1>

- Hall, J., & Sammons, P. (2013). Mediation, moderation & interaction. In *Handbook of Quantitative Methods for Educational Research* (pp. 267–286). Sense Publishers.
- Hosegood, V., Vanneste, A., & Timaeus, I. (2004). Levels and causes of adult mortality in rural South Africa: The impact of AIDS. *AIDS*, 5, 663–671.
- Jamal, F., Fletcher, A., Shackleton, N., Elbourne, D., Viner, R., & Bonell, C. (2015). The three stages of building and testing mid-level theories in a realist RCT: a theoretical and methodological case-example. *Trials*, 16(1), 466. <https://doi.org/10.1186/s13063-015-0980-y>
- Knerr, W., Gardner, F., & Cluver, L. D. (2013). Improving positive parenting skills and reducing harsh and abusive parenting in low- and middle-income countries: A systematic review. *Prevention Science*, 14(4), 352–363. <https://doi.org/10.1007/s11121-012-0314-1>
- Kraemer, H. C., Frank, E., & Kupfer, D. J. (2006). Moderators of treatment outcomes. *JAMA*, 296(10), 1286–1289. <https://doi.org/10.1001/jama.296.10.1286>
- Kraemer, H. C., & Wilson, G. T. (2013). Mediators and moderators of treatment effects in randomized clinical trials. *Archives of General Psychiatry*, 59, 877–883.
- Lachman, J. M., Cluver, L. D., Boyes, M. E., Kuo, C., & Casale, M. (2014). Positive parenting for positive parents: HIV/AIDS, poverty, caregiver depression, child behavior, and parenting in South Africa. *AIDS Care*, 26(3), 304–313. <https://doi.org/10.1080/09540121.2013.825368>
- Lachman, J. M., Heinrichs, N., Jansen, E., Brühl, A., Taut, D., Fang, X., ... Foran, H. M. (2019). Preventing child mental health problems through parenting interventions in Southeastern Europe (RISE): Protocol for a multi-country cluster randomized factorial study. *Contemporary Clinical Trials*, 86, 105855. <https://doi.org/10.1016/j.cct.2019.105855>

- Leijten, P., Gardner, F., Landau, S., Harris, V., Mann, J., Hutchings, J., ... Scott, S. (2017). Research Review: Harnessing the power of individual participant data in a meta-analysis of the benefits and harms of the Incredible Years parenting program. *Journal of Child Psychology and Psychiatry*. <https://doi.org/10.1111/jcpp.12781>
- Leijten, P., Raaijmakers, M., Wijngaards, L., Matthys, W., Menting, A., Hemink-van Putten, M., & Orobio de Castro, B. (2018). Understanding Who Benefits from Parenting Interventions for Children's Conduct Problems: an Integrative Data Analysis. *Prevention Science*, 1–10. <https://doi.org/10.1007/s11121-018-0864-y>
- Lopman, B., Barnabas, R., Boerma, T., Chawira, J., Gaitskell, K., Harrop, T., ... Chawira, G. (2006). Creating and validating an algorithm to measure AIDS mortality in the adult population using verbal autopsy. *Public Library of Science Medicine*, 3, e312.
- Lundahl, B., Risser, H. J., & Lovejoy, M. C. (2006). A meta-analysis of parent training: moderators and follow-up effects. *Clinical Psychology Review*, 26(1), 86–104. <https://doi.org/10.1016/j.cpr.2005.07.004>
- McKenzie, D. (2012). Beyond baseline and follow-up: The case for more T in experiments. *Journal of Development Economics*, 99(2), 210–221. <https://doi.org/10.1016/j.jdeveco.2012.01.002>
- McTaggart, P., & Sanders, M. R. (2007). Mediators and moderators of change in dysfunctional parenting in a school-based universal application of the Triple-P Positive Parenting Programme. *Journal of Children's Services*, 2(1), 4–17. <https://doi.org/10.1108/17466660200700002>
- Meinck, F., Boyes, M., Cluver, L. D., Ward, C., Schmidt, P., DeStone, S., & Dunne, M. (2018). Adaptation and psychometric properties of the ISPCAN Child Abuse Screening Tool for use in trials (ICAST-Trial) among South African adolescents and their primary caregivers. *Child Abuse and Neglect*, 82, 45–58.

- Meinck, F., Cluver, L. D., Boyes, M. E., & Mhlongo, E. L. (2015). Risk and protective factors for physical and sexual abuse of children and adolescents in Africa: A review and implications for practice. *Trauma Violence & Abuse, 16*(1), 81–107.  
<https://doi.org/10.1177/1524838014523336>
- Meinck, F., Cluver, L. D., Boyes, M. E., & Ndhlovu, L. D. (2015). Risk and protective factors for physical and emotional abuse victimisation amongst vulnerable children in South Africa. *Child Abuse Review, 24*(3), 182–197. <https://doi.org/10.1002/car.2283>
- Meinck, F., Cluver, L. D., Orkin, F. M., Kuo, C., Sharma, A. D., Hensels, I. S., & Sherr, L. (2017). Pathways from family disadvantage via abusive parenting and caregiver mental health to adolescent health risks in South Africa. *Journal of Adolescent Health, 60*(1), 57–64. <https://doi.org/10.1016/j.jadohealth.2016.08.016>
- Mikton, C. R., & Butchart, A. (2009). Child maltreatment prevention: A systematic review of reviews. *Bulletin of the World Health Organization, 87*(5), 353–361.  
<https://doi.org/10.2471/BLT.08.057075>
- Moerbeek, M., & Teerenstra, S. (2016). *Power analysis of trials with multilevel data*. Boca Raton, FL: CRC Press.
- Morris, M. C., Kouros, C. D., Janecek, K., Freeman, R., Mielock, A., & Garber, J. (2017). Community-level moderators of a school-based childhood sexual assault prevention program. *Child Abuse & Neglect, 63*(3), 295–306.  
<https://doi.org/10.1016/j.chiabu.2016.10.005>
- Murray, J., Shenderovich, Y., Gardner, F., Mikton, C., Derzon, J. H., Liu, J., & Eisner, M. (2018). Risk Factors for Antisocial Behavior in Low- and Middle-Income Countries: A Systematic Review of Longitudinal Studies. *Crime and Justice, 47*(1), 255–364.  
<https://doi.org/10.1086/696590>
- Pelham, W. E., Dishion, T. J., Tein, J. Y., Shaw, D. S., & Wilson, M. N. (2017). What

- doesn't work for whom? Exploring heterogeneity in responsiveness to the Family Check-Up in early childhood using a mixture model approach. *Prevention Science*, 1–12. <https://doi.org/10.1007/s11121-017-0805-1>
- Peugh, J. L. (2010). A practical guide to multilevel modeling. *Journal of School Psychology*, 48(1), 85–112. <https://doi.org/10.1016/j.jsp.2009.09.002>
- Pillay, U., Roberts, B., & Rule, S. (2006). *South African social attitudes: changing times, diverse voices*. Cape Town: HSRC Press.
- Porter, K. E. (2017). Statistical power in evaluations that investigate effects on multiple outcomes: A guide for researchers. *Journal of Research on Educational Effectiveness*, 1–29. <https://doi.org/10.1080/19345747.2017.1342887>
- Porzsolt, F., Galito, N., Toledo-Arruda, A., Thomaz, T., Moraes, C., Guerra, T., ... Weiß, C. (2015). Efficacy and effectiveness trials have different goals, use different tools, and generate different messages. *Pragmatic and Observational Research*, 47. <https://doi.org/10.2147/POR.S89946>
- Puffer, E. S., Annan, J., Sim, A. L., Salhi, C., & Betancourt, T. S. (2017). The impact of a family skills training intervention among Burmese migrant families in Thailand: A randomized controlled trial. *Plos One*, 12(3), e0172611. <https://doi.org/10.1371/journal.pone.0172611>
- Puffer, E. S., Green, E. P., Chase, R. M., Sim, A. L., Zayzay, J., Friis, E., ... Boone, L. (2015). Parents make the difference: A randomized-controlled trial of a parenting intervention in Liberia. *Global Mental Health*, 2, e15. <https://doi.org/10.1017/gmh.2015.12>
- Rescorla, L. A., Achenbach, T. M., Ivanova, M. Y., Bilenberg, N., Bjarnadottir, G., Denner, S., ... Verhulst, F. C. (2012). Behavioral/emotional problems of preschoolers: Caregiver/teacher reports from 15 societies. *Journal of Emotional and Behavioral*

- Disorders*, 20(2), 68–81. <https://doi.org/10.1177/1063426611434158>
- Shelleby, E. C., & Shaw, D. S. (2014). Outcomes of parenting interventions for child conduct problems: A review of differential effectiveness. *Child Psychiatry and Human Development*, 45(5), 628–645. <https://doi.org/10.1007/s10578-013-0431-5>
- Shenderovich, Y., Eisner, M., Cluver, L., Doubt, J., Berezin, M., Majokweni, S., & Murray, A. L. (2018). What affects attendance and engagement in a parenting program in South Africa? *Prevention Science*, 19(7), 977–986. <https://doi.org/10.1007/s11121-018-0941-2>
- Shenderovich, Y., Eisner, M., Cluver, L., Doubt, J., Berezin, M., Majokweni, S., & Murray, A. L. (2019). Delivering a Parenting Program in South Africa: The Impact of Implementation on Outcomes. *Journal of Child and Family Studies*. <https://doi.org/10.1007/s10826-018-01319-y>
- Shieh, G. (2009). Detecting interaction effects in moderated multiple regression with continuous variables: Power and sample size considerations. *Organizational Research Methods*, 12(3), 510–528. <https://doi.org/10.1177/1094428108320370>
- Snell-Johns, J., Mendez, J. L., & Smith, B. H. (2004). Evidence-based solutions for overcoming access barriers, decreasing attrition, and promoting change with underserved families. *Journal of Family Psychology*, 18(1), 19–35. <https://doi.org/10.1037/0893-3200.18.1.19>
- Spoth, R., Shin, C., Gyll, M., Redmond, C., & Azevedo, K. (2006). Universality of effects: An examination of the comparability of long-term family intervention effects on substance use across risk-related subgroups. *Prevention Science*, 7(2), 209–224. <https://doi.org/10.1007/s11121-006-0036-3>
- Stith, S. M., Liu, T., Davies, L. C., Boykin, E. L., Alder, M. C., Harris, J. M., ... Dees, J. E. M. E. G. (2009). Risk factors in child maltreatment: A meta-analytic review of the literature. *Aggression and Violent Behavior*, 14(1), 13–29.

<https://doi.org/10.1016/j.avb.2006.03.006>

- Straus, M. A., Hamby, S. L., Boney-McCoy, S., & Sugarman, D. B. (1996). The Revised Conflict Tactics Scales (CTS2): Development and preliminary psychometric Data. *Journal of Family Issues*, 17(3), 283–316. <https://doi.org/10.1177/019251396017003001>
- Topçuoğlu, T., Eisner, M., & Ribeaud, D. (2015). Sex differences in the effects of parents' use of corporal punishment on children's aggressive behavior. *Annales de La Faculté de Droit d'Istanbul*, 46, 185–218. Retrieved from <https://dergipark.org.tr/en/pub/iaufdi/issue/727/7851>
- Van Hoorn, R., Tummers, M., Booth, A., Gerhardus, A., Rehfuess, E., Hind, D., ... Kievit, W. (2017). The development of CHAMP: A checklist for the appraisal of moderators and predictors. *BMC Medical Research Methodology*, 17(1), 1–9. <https://doi.org/10.1186/s12874-017-0451-0>
- Vlahovicova, K., Melendez-Torres, G. J., Leijten, P., Knerr, W., & Gardner, F. (2017). Parenting programs for the prevention of child physical abuse recurrence: A Systematic review and meta-analysis. *Clinical Child and Family Psychology Review*, 20(3), 1–15. <https://doi.org/10.1007/s10567-017-0232-7>
- Webster-Stratton, C., Reid, M. J., & Hammond, M. (2004). Treating children with early-onset conduct problems: Intervention outcomes for parent, child, and teacher training. *Journal of Clinical Child and Adolescent Psychology*, 33(1), 105–124. <https://doi.org/10.1207/S15374424JCCP3301>
- Weeland, J., Chhangur, R. R., van der Giessen, D., Matthys, W., de Castro, B. O., & Overbeek, G. (2017). Intervention effectiveness of The Incredible Years: New insights into sociodemographic and intervention-based moderators. *Behavior Therapy*, 48(1), 1–18. <https://doi.org/10.1016/j.beth.2016.08.002>
- Whittaker, K. A., & Cowley, S. (2012). An effective programme is not enough: A review of



factors associated with poor attendance and engagement with parenting support programmes. *Children and Society*, 26(2), 138–149. <https://doi.org/10.1111/j.1099-0860.2010.00333.x>

## Acknowledgements

Many research assistants contributed to the execution of this study. The authors also wish to thank Rocio Herrero Romero, Janina Steinert, Jenny Doubt, Franziska Meinck, and the rest of the research and implementation teams for their dedicated work. Thank you to Andres Molano and Kate Schwartz for their advice and feedback on earlier versions of the manuscript. Special thanks to all the participants and their families. Yulia Shenderovich is supported by the Cambridge Commonwealth, European & International Trust, Additional Insights 2016-2017 Fellowship, Smuts Memorial Fund, managed by the University of Cambridge in memory of Jan Christiaan Smuts, and St John's College, Cambridge. The Sinovuyo Teen Study was funded by the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013, grant agreement 313421), UNICEF Innocenti Office of Research, the Leverhulme Trust (grant number PLP-2014-095), the University of Oxford's Economic and Social Research Council Impact Acceleration Account (K1311-KEA-004 & 1602-KEA-189), and the John Fell Fund (103/757).

## Conflict of Interest

The first author (Shenderovich) worked as a project manager on the randomised trial described in the paper and drew on this research as part of her PhD under the supervision of the second and third authors. The second author (Cluver) was involved in developing the Sinovuyo Caring Families Programme for Parents and Teens, which is licensed under a Creative Commons 4.0 Non-commercial No Derivatives license. Other authors declare no financial relationships with any organisations that might have an interest in the submitted work and no other relationships or activities that could appear to have influenced the submitted work.