

# Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/140743/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Smith, Pamela, Daniel, Rhian, Murray, Rachael L, Moore, Graham, Nelson, Annmarie and Brain, Kate 2021. Psychosocial determinants of quit motivation in older smokers from deprived backgrounds: a cross-sectional survey. *BMJ Open* 11 (5) , e044815. 10.1136/bmjopen-2020-044815 file

Publishers page: <https://doi.org/10.1136/bmjopen-2020-044815>  
<<https://doi.org/10.1136/bmjopen-2020-044815>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1           **Psychosocial Determinants of Quit Motivation in Older Smokers from Deprived**  
2                                   **Backgrounds: A Cross-Sectional Survey**

3           Authors: Pamela Smith\*<sup>a</sup>, Rhian Daniel<sup>a</sup>, Rachael L Murray<sup>b</sup>, Graham Moore<sup>c</sup>, Annmarie  
4                                   Nelson<sup>d</sup>, Kate Brain<sup>a</sup>

5           \*corresponding author

6           [Smithp18@cardiff.ac.uk](mailto:Smithp18@cardiff.ac.uk)

7           8<sup>th</sup> Floor, Neuadd Meirionnydd, Heath Park, CF14 4YS  
8           02920 687695

9  
10           <sup>a</sup> Division of Population Medicine, Cardiff University, Heath Park, Cardiff, UK, CF14 4YS

11           <sup>b</sup> Division of Epidemiology and Public Health Clinical Sciences, University of Nottingham,  
12           Nottingham City Hospital, NG5 1PB

13           <sup>c</sup> Cardiff University, School of Social Sciences, 1-3 Museum Place, Cardiff, UK CF10 3BD

14           <sup>d</sup> Cardiff University, Marie Curie Research Centre, Cardiff, UK, CF14 4YS

15

1 Objectives

2 To identify psychosocial determinants of quit motivation in older deprived smokers. The  
3 evidence may be used to optimise smoking cessation interventions for the target population.

4 Design

5 Cross-sectional survey using online recruitment methods including Facebook-targeted  
6 advertising.

7 Setting

8 United Kingdom, 2019.

9 Participants

10 Current smokers aged 50 years or older and from a socioeconomically deprived background.

11 Main outcome measures

12 Measures included motivation to stop smoking, smoking history, perceived social support,  
13 self-efficacy for quitting, self-exempting beliefs and lung cancer risk perception. Multivariable  
14 regression was used to analyse factors associated with quit motivation.

15 Results

16 Of a total 578 individuals who consented to take part, 278 (48.1%) did not meet the inclusion  
17 criteria. Of the 300 eligible participants most were recruited using Facebook (94.0%), were  
18 aged 50-64 years (83.7%) and female (85.7%). Most participants were renting from a  
19 housing association (72.0%) and had low education (61.0%). Higher motivation to quit was  
20 statistically significantly associated with a higher intensity of previous quit attempts ( $p=0.03$ ),  
21 higher quit confidence ( $p=0.01$ ), higher smoking self-efficacy ( $p=0.01$ ), a lower risk-  
22 minimising beliefs score ( $p=0.01$ ) and using traditional nicotine replacement therapy when  
23 trying to stop smoking or cut down ( $p<0.001$ ).

1 Conclusion

2 Older smokers from deprived backgrounds face complex barriers to quitting smoking.  
3 Interventions are needed to increase self-efficacy for quitting, modify risk-minimising beliefs  
4 and target elements of previous quit attempts (i.e. the use of nicotine replacement therapy)  
5 that are associated with motivation to stop smoking.

6 Strengths and limitations of this study

- 7       • This research adds knowledge to the context of targeted smoking cessation  
8       interventions for a population who are at high-risk of developing a variety of smoking  
9       related diseases, including lung cancer
- 10       • An online survey, disseminated through Facebook advertising was useful for  
11       recruiting the target population
- 12       • There was an underrepresentation of older individuals (i.e. 65 years and older) and  
13       males
- 14       • Due to the cross-sectional study design, caution is needed when interpreting  
15       associations with quit motivation

16

1 **INTRODUCTION**

2 Tobacco smoking is estimated to cause the premature deaths of approximately six million  
3 individuals worldwide and 96,000 within the UK per year (1), resulting in an average of ten  
4 life years lost (2). Smokers who do not stop smoking also begin to suffer diseases of old age  
5 around ten years earlier in comparison to non-smokers (3). Most deaths related to smoking  
6 arise from cancers, respiratory disease and cardiovascular disease (1). For example, lung  
7 cancer mortality is around fifteen times higher in current smokers compared to never-  
8 smokers and increases with smoking amount and duration (4).

9 Smoking is a major health inequality concern due to higher smoking rates among people  
10 from low socioeconomic groups in developed countries. The high prevalence of cigarette  
11 smoking among low socioeconomic groups demonstrates the striking relationship between  
12 social context and health behaviour (5, 6).

13 Quit attempts by older smokers from deprived backgrounds are less likely to be successful  
14 due to higher levels of nicotine dependence and/or a lack of self-efficacy (7, 8). Research  
15 has indicated that smokers from deprived backgrounds find quitting more difficult due to  
16 having less social support when making a quit attempt (9, 10). Self-exempting beliefs, also  
17 known as risk-minimising beliefs, have also been shown to influence continued smoking  
18 among individuals from low socioeconomic groups. These beliefs may be adopted as a  
19 mechanism to help rationalise or justify smoking despite the well-known harms (11).

20 Perceived risk of smoking-related diseases, such as lung cancer, has been linked to  
21 smoking, with absolute and relative lung cancer risk perception being positively related to  
22 cigarette-smoking behaviours (12). Furthermore, previous research involving deprived  
23 smokers aged 40 years or older has found perceived lung cancer risk to be higher among  
24 this population compared to non-smokers (13).

25 A systematic review of behavioural smoking cessation interventions indicated that offering  
26 incentives, use of peer facilitators and more intensive behavioural counselling are promising  
27 for encouraging smoking cessation in older smokers from deprived backgrounds (14).

1 However, further evidence is needed regarding psychosocial targets for smoking cessation  
2 interventions aimed at this population (14).

3 In this study, we aimed to identify modifiable psychosocial determinants of motivation to stop  
4 smoking among older smokers from socioeconomically deprived backgrounds. The selected  
5 survey measures were driven by a variety of theories including PRIME Theory of Addiction  
6 and the Extended Parallel Process Model (15, 16). It was anticipated that higher motivation  
7 to quit smoking would be associated with lower nicotine dependence, higher self-efficacy,  
8 higher perceived social support, lower self-exempting beliefs and higher perceived lung  
9 cancer risk. The evidence may be used to improve smoking cessation interventions for the  
10 target population.

## 11 **METHODS**

### 12 **Participants and recruitment methods**

13 Sample size was determined on the basis of (i) tobacco smoking rates in the UK population  
14 of older smokers (17) and (ii) expected response rate among smokers (18, 19). Although we  
15 planned from the outset to measure the primary outcome of motivation to stop smoking on a  
16 7-point scale, it was impossible to know in advance whether its conditional distribution given  
17 predictors would allow analysis using linear regression. The decision was thus taken to  
18 calculate the sample size based on a dichotomisation of the outcome, which is conservative  
19 given that dichotomisation decreases power. A sample of 300 respondents was chosen to  
20 provide 80% power to detect an odds ratio of 2.0 in the dichotomised primary outcome at a  
21 5% significance level (20).

22 Participants were recruited to an online cross-sectional survey pragmatically and were  
23 current smokers, aged 50 years or more, from deprived backgrounds. Participants were  
24 given the option to be entered into a £50 shopping voucher prize draw and consented to be  
25 contacted if they won a voucher. A link to the online questionnaire was disseminated using  
26 Facebook, a population survey platform (HealthWise Wales:

1 [www.healthwisewales.gov.wales](http://www.healthwisewales.gov.wales)) and a Wales based cancer charity (Tenovus Cancer  
2 Care). Snowball sampling was also used to recruit the target sample.

3 To determine eligibility, participants completed five questions based on socioeconomic  
4 deprivation, smoking status and age. Participants were eligible if they were  $\geq 50$  years of age,  
5 a current smoker and met the threshold for at least two out of three individual-level  
6 deprivation indicators. Thresholds for deprivation indicators were: 1) *Education* - educated to  
7 O-level or GCSE equivalent (mostly grade D-G) or below; 2) *Household income* - casual  
8 labourer, pensioner, student, unemployed (e.g. pensioner without private pension and  
9 anyone living on basic benefits), semi-skilled and unskilled manual worker, skilled manual  
10 worker and those with no previous or current employment within the household; 3) *Home*  
11 *ownership* - renting from local authority/housing association or living with family/friends.

## 12 **Ethical approval**

13 The study received ethical approval from Cardiff University School of Medicine (SMREC  
14 Reference Number: 19/06).

## 15 **Patient and public involvement**

16 Public involvement was integrated according to national standards (21), and used during the  
17 development and design of the survey questions and structure. An appointed research  
18 partner presented feedback on the survey at various stages including the final version, prior  
19 to recruitment commencing. The research partner also assisted in the recruitment of  
20 participants for cognitive interviews used for survey development.

## 21 **Measures**

22 Cognitive interviews were first carried out with the target population to ascertain the  
23 suitability of the questionnaire. Interviews were conducted with five participants (three female  
24 and two male, aged 50-88 years) from a socioeconomically deprived area of South Wales.  
25 Content validity of unvalidated items was assessed by a panel of six academics with

1 experience in the field of health behaviours. These steps were conducted in order to assess  
2 acceptability and comprehension of items, and resulted in adaptations to item wording and  
3 inclusion of selected items for the final version of the survey (see Supplementary Material).

#### 4 *Demographic characteristics*

5 Data were collected on gender, age, relationship status, highest level of education, home  
6 ownership, household employment and co-morbid conditions (22). A total score was created  
7 for number of comorbid conditions, with a higher score indicating the presence of more  
8 comorbid conditions (score range 0-14).

#### 9 *Smoking characteristics*

10 Nicotine dependence was assessed using the Fagerstrom Test for Nicotine Dependence  
11 (23) in order to create a total nicotine dependence score. Number of previous quit attempts  
12 (24) and intensity of previous quit attempts (number of attempts combined with duration of  
13 the longest previous quit attempt) were also measured. Participants were asked whether  
14 they were currently trying to cut down and were asked to select multiple options from a list of  
15 smoking cessation aids (nicotine gum, lozenges, patches, inhalator, mouth spray, electronic  
16 cigarette, behavioural counselling or not using anything to help stop smoking).

#### 17 *Motivation to stop smoking (primary outcome)*

18 The Motivation to Stop Scale (MTSS) (25) is a validated measure that assesses the key  
19 elements of quit motivation: belief, desire and intention (15). The MTSS is a single-item  
20 instrument and assesses a smoker's motivation to stop smoking. The scale score ranged  
21 from 1-7: (1) 'I don't want to stop smoking', (2) 'I think I should stop smoking but don't really  
22 want to', (3) 'I want to stop smoking but haven't thought about when', (4) 'I REALLY want to  
23 stop smoking but I don't know when I will', (5) 'I want to stop smoking and hope to soon', (6)  
24 'I REALLY want to stop smoking and intend to in the next 3 months' and (7) 'I REALLY want  
25 to stop smoking and intend to in the next month'.



1 *Social support and smoking*

2 The social support and smoking questions (26) measured how supported the participant felt  
3 during a previous quit attempt. Respondents were asked about various forms of social  
4 support experienced during a previous quit attempt, for example “*Encourage you to keep at*  
5 *quitting*”, “*Celebrate your quitting with you*”, “*Say you were going to start smoking again*”.  
6 Response options were rated on a scale of 1 to 5 (1=Never, 5=Very often). Principal  
7 components analysis (PCA) of the adapted social support items yielded two factor-derived  
8 subscales that were labelled positive smoking social support (8 items,  $\alpha=0.85$ ) and negative  
9 smoking social support (3 items,  $\alpha=0.64$ ). Total possible score range for positive social  
10 support was 8-40 with higher scores indicating higher positive social support during a  
11 previous quit attempt. Total possible score range for negative social support was 3-15 with  
12 higher scores indicating higher negative social support during a previous quit attempt.  
13 Participants were also asked how supported they felt by their partner, friends, colleague and  
14 GP/ healthcare professional during a previous quit attempt, with response options ranging  
15 from 1-5 (1=not at all, 5=extremely).

16 *Self-exempting beliefs*

17 The self-exempting beliefs measure comprises 16 items, with two subscales for self-  
18 exempting (“bulletproof”) beliefs and risk-minimising (“jungle”, “skeptic”, “worth it”) beliefs  
19 (27). Examples of risk-minimising beliefs items included: “*Smoking cannot be all that bad for*  
20 *you because many people who smoke live long lives*”, “*I would rather live a shorter life and*  
21 *enjoy it than a longer one where I will be deprived of the pleasure of smoking*” and “*You*  
22 *have got to die of something, so why not enjoy yourself and smoke*”. Agreement with each  
23 item was rated on a scale of 1 to 5 (1=totally disagree, 5=totally agree). Two factor-derived  
24 subscales labelled risk-minimising beliefs (5 items,  $\alpha=0.83$ ) and scepticism (2 items,  $\alpha=0.54$ )  
25 were derived from PCA. Total possible score ranges were 5-25 for the risk-minimising beliefs  
26 scale and 2-10 for the scepticism scale, with higher scores indicating higher levels of  
27 agreement.

1 *Smoking self-efficacy*

2 The eight-item smoking self-efficacy questionnaire (SEQ-12) (28) was originally designed to  
3 measure current smokers' confidence in their ability to abstain from smoking in high-risk  
4 situations. The instrument presents participants with situations in which people might be  
5 tempted to smoke such as when they feel nervous, depressed, angry, have the urge to  
6 smoke, are with other smokers and when they are having a tea or coffee. Participants  
7 responded on a 5-point Likert scale from 1 ("not at all sure") to 5 ("absolutely sure"). Total  
8 score range for this measure was 6-30 with higher scores indicating greater smoking self-  
9 efficacy ( $\alpha=0.88$ ). After PCA, two items were removed and a smoking self-efficacy scale was  
10 created.

11 *Quit confidence*

12 Quit confidence was measured using a single item (29): "*How confident are you that you*  
13 *could quit smoking for good if you wanted to?*". The response options were on a scale of 1 to  
14 5 where 1="not at all" and 5="extremely".

15 *Lung cancer experience*

16 Lung cancer experience was measure using seven items to assess personal experience of  
17 lung cancer as well as experience of social contacts (family, friends, member of community)  
18 (30). Response options for this measure were 'yes', 'no', 'don't know' and 'prefer not to say'.

19 *Lung cancer risk perception*

20 Three items were used to assess lung cancer risk perception (31). Absolute risk perception  
21 was measured using the item "*How likely do you think it is that you will develop lung cancer*  
22 *in your lifetime?*" with response options from 1 ("very unlikely") to 5 ("very likely").

23 Comparative risk perception was measured by asking "*Compared to others your age and*  
24 *sex, what do you think is your chance of getting lung cancer in your lifetime?*" with response  
25 options from 1-5 ('much lower' to 'much higher'). Affective risk perception was measured

1 using the question “*How worried are you about getting lung cancer in your lifetime?*”, with  
2 response options on a scale of 1 (“not at all”) to 4 (“extremely”).

### 3 **Statistical analyses**

4 Data were analysed using SPSS version 25. In order to deal with missing data in preparation  
5 for univariable and multivariable analysis, multiple imputation using chained equations was  
6 conducted on the original dataset (32). All the incomplete variables were either dichotomous  
7 or (treated as) continuous. For each incomplete variable being imputed, all other variables  
8 were included in its imputation model. Where the incomplete variables were constituent parts  
9 of a composite score (e.g. positive and negative social support scale), the imputation was  
10 performed on the individual questionnaire responses, and these imputed variables were then  
11 summed to create the imputed composite score. Those who responded with ‘*Don’t know*’ to  
12 MTSS were removed from analysis.

13 PCA was used to identify the underlying factor structure of unvalidated measures (social  
14 support and smoking, self-exempting beliefs, smoking self-efficacy) that had been adapted  
15 on the basis of cognitive interviewing and content validity assessment.

16 Univariable associations between MTSS and demographic (age, gender, education, income,  
17 living situation, presence of comorbid conditions, lung cancer experience), smoking-related  
18 factors (nicotine dependence, previous quit attempts, smoking abstinence, smoking  
19 cessation aids) and psychosocial variables (self-efficacy, quit confidence, perceived social  
20 support for quitting, self-exempting beliefs, lung cancer risk perception) were assessed.

21 Independent T-tests, Spearman’s Rank Correlation and Pearson’s Correlation were used to  
22 test for univariable associations. A multivariable linear regression analysis was conducted, in  
23 which all variables were entered in a single step. Confidence intervals of 95% were  
24 calculated for the associations between MTSS and variables that were statistically significant  
25 at  $p \leq 0.05$  in univariable analyses.

1 MTSS was treated as a continuous variable, with a low score demonstrating an absence of  
2 any belief, desire or intention to stop smoking and a high score demonstrating strong desire  
3 and short-term intention. Examination of the distribution of the outcome variable, and its  
4 mean and variance given predictors, suggested that a multivariable linear regression model  
5 was a suitable choice.

## 6 **RESULTS**

### 7 **Sample characteristics**

8 A total of 578 individuals consented to take part in the survey. Of those who consented, 278  
9 (48.1%) did not meet the inclusion criteria due to age (n=2), smoking status (n=24) and level  
10 of deprivation (n=252). Of 300 eligible participants, most were recruited using Facebook  
11 advertising (94%). The majority of participants were aged between 50 and 64 years of age  
12 (83.7%) and female (85.7%).

13 Almost half of participants (43.7%) reported that they did not want to stop smoking (see  
14 Table 1). Most participants had made a serious previous quit attempt (73.3%) and were  
15 currently trying to cut down on how much they smoked (66%). The most commonly used  
16 smoking cessation aid when trying to cut down/stop smoking was e-cigarettes (28.3%), with  
17 over half the sample (51.1%) not using anything to help them cut down/stop smoking.

### 18 **Univariable analysis of factors associated with motivation to stop smoking**

19 At univariable analysis, MTSS was significantly associated with 12 variables (see Table 2  
20 and 3). As depicted in Table 2, higher motivation to stop smoking was statistically  
21 significantly associated with lower nicotine dependence ( $p=0.05$ ), higher intensity of previous  
22 quit attempts ( $p<0.001$ ), currently trying to cut down on smoking ( $p<0.001$ ), using a smoking  
23 cessation aid when trying to help cut down/quit smoking ( $p<0.001$ ) and trying to cut  
24 down/quit without any smoking cessation aid ( $p=0.04$ ) (Table 2). Associations between  
25 MTSS and gender, age, relationship, education, employment, housing and comorbidity were  
26 not statistically significant.

1 As shown in Table 3, higher motivation to stop smoking was statistically significantly  
2 associated with higher positive social support when making a quit attempt ( $p=0.04$ ), higher  
3 smoking self-efficacy ( $p<0.001$ ), higher quit confidence ( $p<0.001$ ) and lower risk-minimising  
4 beliefs ( $p<0.001$ ). Higher motivation to stop smoking was statistically significantly associated  
5 with higher perceived comparative ( $p=0.01$ ) and affective ( $p<0.001$ ) lung cancer risk, and  
6 having experienced lung cancer in a close friend ( $p=0.01$ ).

### 7 **Multivariable regression of factors associated with motivation to stop smoking**

8 Conditional on all other predictors in the model, a unit increase in intensity of previous quit  
9 attempts was associated with an estimated 0.17 increase in MTSS ( $p=0.03$ , 95% CI=0.02,  
10 0.32). A unit increase for quit confidence was associated with an estimated 0.20 increase in  
11 MTSS ( $p=0.01$ , 95% CI=0.06, 0.35), self-efficacy was associated with an estimated 0.22  
12 increase in MTSS ( $p=0.01$ , 95% CI=0.06, 0.37) and a unit increase in risk-minimising beliefs  
13 was associated with an estimated 0.18 decrease in MTSS ( $p=0.01$ , 95% CI= 0.04, 0.32).  
14 Those who reported using traditional nicotine replacement therapy (NRT) when trying to stop  
15 smoking or cut down scored on average a 0.26 higher MTSS score ( $p<0.001$ , 95% CI=0.12,  
16 0.39) compared to those who did not (Table 4).

## 1 **DISCUSSION**

2 To our knowledge this is the first study to examine psychosocial determinants of quit  
3 motivation in a sample of older smokers, aged 50+ from deprived backgrounds. Higher  
4 motivation to quit was associated with higher intensity of previous quit attempts, higher quit  
5 confidence, higher smoking self-efficacy, lower risk-minimising beliefs and using traditional  
6 nicotine replacement therapy when trying to stop smoking or cut down.

7 Previous studies suggest that smokers from deprived backgrounds are equally motivated to  
8 try and quit smoking but less likely to be successful during a quit attempt compared to more  
9 affluent smokers (33-36). This paradox results in lower quit rates and potentially exacerbates  
10 inequalities in the prevalence of smoking. An important factor that contributes to higher  
11 smoking rates among deprived smokers lies in the lack of research output that target this  
12 population (37) and the general scarcity of literature for effective smoking cessation  
13 interventions for more disadvantaged smokers (38).

14 The current findings suggest that the salient determinant of quit motivation in the target  
15 population was the use of traditional NRT (e.g. patches, gum, lozenges) during a previous  
16 attempt to cut down or quit smoking. Use of NRT has been shown to increase cessation  
17 success (39) and is more effective at promoting cessation in older age groups compared to  
18 younger groups (40, 41). Our findings confirm that NRT is an important tool in encouraging  
19 motivation to stop smoking in older smokers from deprived backgrounds. However, evidence  
20 for which type of NRT works best for this population is limited and the provision of NRT may  
21 need to be tailored in order to fit the needs and preferences of this population.

22 Previous studies have found deprived smokers to be less likely to use NRTs and that one of  
23 the barriers to quitting includes inability to access treatment (42). Making NRTs easily  
24 available to this population could increase motivation to stop smoking and encourage older  
25 smokers from deprived backgrounds to make serious quit attempts that are more likely to be  
26 successful. Although an association between quit motivation and the presence of comorbid

1 conditions was not observed in the current study, it has been proposed that smoking  
2 cessation interventions (specifically NRTs) for older adults should be personalised due to the  
3 presence of comorbidities and medication that can reduce the use and effectiveness of NRT  
4 (43).

5 Furthermore, the current research suggests that a higher intensity of previous quit attempts  
6 is associated with higher motivation to stop smoking. Future interventions should focus on  
7 encouraging the target population to persist with their quit attempts as well as promote the  
8 use of behavioural support and NRT in order to improve motivation to quit.

9 The current study further illustrates the importance of quitting self-efficacy in relation to  
10 improving motivation to stop smoking among older smokers from deprived backgrounds.  
11 Previous research from Siahpush et al (44) showed that low self-efficacy can explain the  
12 lower smoking cessation rate observed among low socioeconomic smokers. Individuals from  
13 deprived backgrounds are more likely to be exposed to stressful and disadvantaged lives  
14 and have less access to material and social resources, which in turn can reduce self-efficacy  
15 and motivation to stop smoking (45, 46).

16 Risk-minimising beliefs may be adopted in order to justify continued smoking despite the  
17 risks to health (27, 47). Similarly, previous studies indicate that individuals from socially  
18 deprived backgrounds are more likely to hold self-exempting beliefs (27, 47-49). Older  
19 smokers who have previously been shown to hold these beliefs consider themselves as  
20 '*survivors*', resulting in a lower motivation to stop smoking (1).

21 Methodological limitations of the current study are acknowledged. Although we were able to  
22 demonstrate representation of smokers from a low socioeconomic background in the sample  
23 there are issues of generalisability. Men and those aged 65 years or older were under-  
24 represented which could be due to older adults being less likely to have access to social  
25 media or the internet (50) and men being traditionally more difficult to recruit to research  
26 (51). Due to the cross-sectional study design, caution is needed when interpreting

1 associations due to shared-method variance and inability to infer causal relationships (52).  
2 Further limitations include the variable internal validity of factor-derived subscales and the  
3 use of self-reported measures. Suitable validated measures are needed to better capture  
4 evidence of psychosocial influences on smoking behaviour in older smokers from deprived  
5 backgrounds.

6 This research is valuable in the context of targeted smoking cessation interventions for a  
7 population who are at high-risk of developing a variety of smoking-related diseases,  
8 including lung cancer. The target population face a breadth of complex barriers to quitting  
9 smoking and interventions are needed to increase self-efficacy for quitting, dispel risk-  
10 minimising beliefs and target elements of previous quit attempts (i.e. NRTs used) that are  
11 associated with motivation to stop smoking. Findings from this research could be used in the  
12 adaptation of behavioural smoking cessation interventions by addressing identified  
13 psychosocial determinants. This has the potential to improve quit motivation and could  
14 encourage a smoking cessation attempt in older smokers from deprived backgrounds.

15 Contributors: PS, KB, AN, RM and GM were responsible for the concept, overall design and  
16 conduct of the study. RD gave additional support and advice on multiple imputation for  
17 missing data and statistical analysis. PS was responsible for collection of data and  
18 manuscript preparation. All authors were involved in the interpretation of the results and  
19 approved the final version of the manuscript. No competing interests were declared by  
20 authors.

21 Funding: This study was funded by School of Medicine, Cardiff University. AN's post was  
22 supported by Marie Curie core grant funding (grant reference: MCCCFCO-11-C). The work  
23 was also conducted under the auspices of the Wales Cancer Research Centre (Health and  
24 Care Research Wales grant reference: 517190).

## 25 **DATA SHARING AND COMPETING INTERESTS**



1 All data relevant to the study are included in the article or uploaded as supplementary  
2 information. The authors declare that they have no conflict of interest.

3

1 **Table 1.** Sample characteristics (N=300)

Variable	Descriptive statistic	
	n	%
<b>Age, years</b>		
50-64 years	251	83.7
65 years or older	49	16.3
Missing	0	0
<b>Gender</b>		
Male	43	14.3
Female	257	85.7
Missing	0	0
<b>Highest level of education</b>		
Finished school at or before age fifteen	76	25.3
No qualifications/ left school at 16	107	35.7
Completed CSEs, O-levels or equivalent (Mostly grade A-C)	56	18.7
Completed CSEs, O-levels or equivalent (Mostly grade D-G)	22	7.3
Completed A levels or equivalent	7	2.3
Completed further education but not degree	30	10
Completed a Bachelor's degree/Masters/PhD	2	0.7
Missing	0	0
<b>Home/Living arrangement</b>		
Own outright	19	6.3
Own mortgage	19	6.3
Rent from local authority/housing association	216	72
Rent privately	35	11.7
Living with family or friends	11	3.7
Missing	0	0
<b>Employment</b>		
Casual labourer, pensioner, student, unemployed (e.g. pensioner without private pension and anyone living on basic benefits)	99	33
Semi-skilled and unskilled manual worker (e.g. assembly line worker, refuse collector, messenger)	88	29.3
Skilled manual worker (e.g. electrician, carpenter)	45	15
Supervisory/ clerical/ junior managerial/ professional/ administrative (e.g. supervisor, bank clerk, salesperson)	19	6.3
Intermediate managerial/professional/ administrative (e.g. middle management, bank manager, teacher)	4	1.3
Higher managerial/professional/administrator (e.g. Chief executive, senior civil servant, surgeon)	0	0
No previous or current employment within the household	45	15
<b>Relationship status</b>		
Married or in a civil partnership	90	30
Living with my partner	38	12.7
Single (never married and not living with a partner)	43	14.3
Divorced or separated and not living with another partner	92	30.7
Widowed and not living with another partner	32	10.7
Prefer not to say	4	1.3
Missing	1	0.3
<b>Pre-existing health conditions</b>		
Heart disease	28	9.3
High blood pressure	65	21.7

Lung disease	68	22.7
Diabetes	43	14.3
An ulcer or stomach disease	29	9.7
Kidney disease	7	2.3
Liver disease	5	1.7
Anaemia or other blood disease	17	5.7
Cancer	5	1.7
Depression	154	51.3
Osteoarthritis or degenerative	90	30
Back pain	167	55.7
Rheumatoid arthritis	42	14
<b>Motivation to Stop Smoking</b>		
I don't want to stop smoking	41	13.7
I think I should stop smoking but don't really want to	90	30.0
I want to stop smoking but haven't thought about when	40	13.3
I REALLY want to stop smoking but don't know when I will	64	21.3
I want to stop smoking and hope to soon	19	6.3
I REALLY want to stop smoking and intend to in the next 3 months	14	4.7
I REALLY want to stop smoking and intend to in the next month	16	5.3
Don't know*	16	5.3
<b>Previous serious quit attempt</b>		
Yes	220	73.3
No	79	26.3
Missing	1	0.3
<b>Currently trying to cut down smoking</b>		
Yes	198	66.0
No	100	33.3
Missing	2	0.7
<b>Smoking cessation aids (n= 198)</b>		
Nicotine gum	4	1.8
Nicotine lozenge	5	2.3
Nicotine patch	14	6.4
Nicotine inhaler/inhalator	6	2.7
Another nicotine product	3	1.4
Electronic cigarette	62	28.3
Nicotine mouth spray	3	1.4
Behavioural counselling (e.g. group sessions, telephone support, individual support)	2	0.9
I am not using anything to help me stop smoking	112	51.1

1 \*Participants who responded 'don't know' were removed from further analysis

**Table 2.** Univariable associations between Motivation To Stop Smoking and demographic/smoking characteristics (n=284)

<b>Variable</b>	<b>N</b>	<b>MTSS mean (SD)</b>	<b>Test statistic/ Correlation Coefficient</b>	<b>P value</b>
<b>Gender<sup>1</sup></b>				
Female	243	3.14 (1.62)	0.22	0.82
Male	41	3.07 (1.84)		
<b>Age<sup>1</sup></b>				
50-64 years	239	3.10 (1.65)	-0.72	0.47
65 +	45	3.29 (1.69)		
<b>Relationship<sup>1</sup></b>				
Widowed/divorced/single	161	3.14 (1.65)	0.14	0.89
Married/living with a partner	120	3.11 (1.66)		
Missing data	5			
<b>Education<sup>1</sup></b>				
Lower education	193	3.11 (1.62)	-0.27	0.79
Higher education	91	3.16 (1.72)		
<b>Employment<sup>1</sup></b>				
Lower employment level	262	3.13 (1.64)	0.24	0.81
Higher employment level	22	3.05 (1.81)		
<b>Housing<sup>1</sup></b>				
Own house/ mortgage/ rent privately	70	3.36 (1.70)	1.35	0.18
Rent from housing association/ living with friends or family	214	3.05 (1.63)		
<b>Comorbidity score<sup>2</sup></b>	284		0.00	0.97
<b>Nicotine dependence score<sup>2</sup></b>	284		-0.11	<b>0.05</b>
<b>Intensity of previous quit attempt score<sup>2</sup></b>	209		0.26	<b>&lt;0.001</b>
<b>Currently trying to cut down on smoking<sup>1</sup></b>				
Yes	185	3.44 (1.57)	4.53	<b>&lt;0.001</b>
No	97	2.54 (1.64)		

Variable	N	MTSS mean (SD)	Test statistic	P value
<b>Using traditional NRTs to help cut down/quit<sup>1</sup></b>				
Yes	23	4.87 (1.55)	-4.95	<b>&lt;0.001</b>
No	162	3.24 (1.47)		
<b>Using electronic cigarette to help cut down/quit<sup>1</sup></b>				
Yes	58	3.21 (1.46)	1.37	0.17
No	226	3.55 (1.61)		
<b>Trying to cut down/quit without any smoking cessation aid<sup>1</sup></b>				
Yes	104	3.23 (1.44)	2.07	<b>0.04</b>
No	81	3.72 (1.70)		

<sup>1</sup> Independent t-test

<sup>2</sup> Pearson's correlation coefficient

**Table 3.** Univariable associations between Motivation To Stop Smoking and psychosocial variables

Variable	N	MTSS mean (SD)	Test Statistic/ Correlation Coefficient	P value
<b>Psychological variables</b>				-
Risk-minimising beliefs <sup>1</sup>	284	-	-0.27	<0.001
Scepticism <sup>1</sup>	284	-	-0.07	0.26
Quit confidence <sup>2</sup>	284	-	0.40	<0.001
Smoking self-efficacy <sup>1</sup>	284	-	0.21	<0.001
<b>Smoking and social support during a previous quit attempt</b>				
Someone to turn during a quit attempt <sup>2</sup>		-	-0.04	0.53
Positive smoking social support <sup>1</sup>	211	-	0.14	<b>0.04</b>
Negative smoking social support <sup>1</sup>	210	-	-0.01	0.91
Support from partner during previous quit attempt <sup>2</sup>	180	-	-0.01	0.89
Support from family during previous quit attempt <sup>2</sup>	200	-	0.03	0.63
Support from friends during previous quit attempt <sup>2</sup>	198	-	0.01	0.92
Support from colleagues during previous quit attempt <sup>2</sup>	187	-	-0.00	0.96
Support from GP or HCP during previous quit attempt <sup>2</sup>	188	-	-0.06	0.44
Someone to turn during if finding quitting difficult <sup>2</sup>	284	-	-0.04	0.53
<b>Lung cancer experience <sup>3</sup></b>				
<i>Self</i>				-
Yes	19	3.19 (1.85)	0.22	0.83
No	246	3.08 (1.63)		
<i>Partner</i>				
Yes	19	3.58 (2.10)	0.77	0.45
No	223	3.13 (1.64)		
<i>Close family member</i>				
Yes	98	3.30 (1.58)	1.13	0.26
No	157	3.05 (1.65)		
<i>Other family member</i>				
Yes	63	3.29 (1.56)	1.38	0.17
No	174	2.96 (1.58)		
<i>Close friend</i>				
Yes	63	3.62 (1.68)	2.78	<b>0.01</b>
No	173	2.88 (1.62)		

<b>Variable</b>	<b>N</b>	<b>MTSS mean (SD)</b>	<b>Test Statistic</b>	<b>P value</b>
<i>Other friend</i>				
Yes	53	3.34 (1.60)	1.42	0.16
No	174	2.96 (1.55)		
<i>A member of your community</i>				
Yes	87	3.29 (1.66)	1.84	0.07
No	124	2.87 (1.52)		
<b>Lung cancer risk perception</b> <sup>2</sup>				
Absolute risk	284	-	0.11	0.08
Comparative risk	284	-	0.15	<b>0.01</b>
Affective risk	284	-	0.36	<b>&lt;0.001</b>

<sup>1</sup> *Pearson's correlation coefficient*

<sup>2</sup> *Spearman's Rank Correlation Coefficient*

<sup>3</sup> *Independent t-test*

**Table 4.** Multivariable associations with Motivation To Stop Smoking

<b>Variable</b>	<b>Estimated adjusted mean difference</b>	<b>Standard Error</b>	<b>95% Confidence Interval</b>	<b>P value</b>
Intensity of previous quit attempts	0.17	0.08	0.02, 0.32	<b>0.03</b>
Currently trying to cut down on smoking	-0.02	0.31	-0.20, 0.16	0.81
Positive social support during a previous quit attempt	0.00	0.02	-0.14, 0.14	1.00
Smoking self-efficacy	0.22	0.03	0.06, 0.37	<b>0.01</b>
Quit confidence	0.20	0.10	0.06, 0.35	<b>0.01</b>
Risk-minimising beliefs	-0.18	0.03	-0.32, -0.04	<b>0.01</b>
Close friend with a diagnosis of lung cancer**	0.13	0.26	-0.01, 0.26	0.07
Comparative lung cancer risk	-0.04	0.15	-0.18, 0.11	0.64
Affective lung cancer risk	0.12	0.10	-0.02, 0.27	0.10
Using traditional NRTs to help cut down/quit*	0.26	0.43	0.12, 0.39	<b>&lt;0.001</b>
Not using any smoking cessation aids to help cut down*	0.02	0.27	-0.14, 0.18	0.78
Nicotine dependence	0.06	0.05	-0.08, 0.20	0.30

\*Dichotomous variable (coded yes/no)



## References

1. Action on Smoking and Health 2016 [Available from: <http://ash.org.uk/information-and-resources/fact-sheets/smoking-statistics-who-smokes-and-how-much/>].
2. Office of the Surgeon G, Office on S, Health. Reports of the Surgeon General. The Health Consequences of Smoking: A Report of the Surgeon General. Atlanta (GA): Centers for Disease Control and Prevention (US); 2004.
3. Jha P, Peto R. Global effects of smoking, of quitting, and of taxing tobacco. *N Engl J Med*. 2014;370(1):60-8.
4. Doll R, Peto R, Boreham J, Sutherland I. Mortality from cancer in relation to smoking: 50 years observations on British doctors. *Br J Cancer*. 2005;92(3):426-9.
5. Cavelaars AE, Kunst AE, Geurts JJ, Crialesi R, Grotvedt L, Helmert U, et al. Educational differences in smoking: international comparison. *BMJ*. 2000;320(7242):1102-7.
6. Huisman M, Kunst AE, Mackenbach JP. Inequalities in the prevalence of smoking in the European Union: comparing education and income. *Prev Med*. 2005;40(6):756-64.
7. Benowitz NL. Nicotine addiction. *N Engl J Med*. 2010;362(24):2295-303.
8. Fiore MC, Jaen CR, Baker TB, Bailey WC, Benowitz NL, Curry SJ, et al. Treating tobacco use and dependence: 2008 update US Public Health Service Clinical Practice Guideline executive summary. *Respiratory Care*. 2008;53(9):1217-22.
9. Chandola T, Head J, Bartley M. Socio-demographic predictors of quitting smoking: how important are household factors? *Addiction*. 2004;99(6):770-7.
10. Lacey LP, Manfredi C, Balch G, Warnecke RB, Allen K, Edwards C. Social support in smoking cessation among black women in Chicago public housing. *Public Health Reports*. 1993;108(3):387-94.
11. Guillaumier A, Bonevski B, Paul C, D'Este C, Twyman L, Palazzi K, et al. Self-Exempting Beliefs and Intention to Quit Smoking within a Socially Disadvantaged Australian Sample of Smokers. 2016.
12. Chen L-S, Kaphingst KA, Tseng T-S, Zhao S. How are lung cancer risk perceptions and cigarette smoking related?-testing an accuracy hypothesis. *Transl Cancer Res*. 2016;5(Suppl 5):S964-S71.
13. Quaife SL, Marlow LAV, McEwen A, Janes SM, Wardle J. Attitudes towards lung cancer screening in socioeconomically deprived and heavy smoking communities: informing screening communication. *Health Expect*. 2017;20(4):563-73.
14. Smith P, Poole R, Mann M, Nelson A, Moore G, Brain K. Systematic review of behavioural smoking cessation interventions for older smokers from deprived backgrounds. *BMJ Open*. 2019;9(11):e032727.
15. West R, Brown J. Theory of addiction. Chichester, West Sussex, UK: Wiley Blackwell/Addiction Press,; 2014.
16. Witte K. Putting the Fear Back into Fear Appeals - the Extended Parallel Process Model. *Communication Monographs*. 1992;59(4):329-49.
17. Office for National Statistics- Adults smoking habits in the UK 2017 [Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/healthandlifeexpectancies/bulletins/adultsmokinghabitsingreatbritain/2017>].
18. Brodar KE, Hall MG, Butler EN, Parada H, Stein-Seroussi A, Hanley S, et al. Recruiting Diverse Smokers: Enrollment Yields and Cost. *Int J Environ Res Public Health*. 2016;13(12).
19. Thompson TP, Aveyard P, Ayres R, Taylor R, Warren F, Campbell J, et al. Strategies to recruit hard-to-reach smokers into Exercise Assisted Reduction then Stop (EARS): a pragmatic pilot randomised trial. *The Lancet*. 2012;380:S73.
20. Dupont WD, Plummer WD. Power and Sample Size Calculations for Studies Involving Linear Regression. *Controlled Clinical Trials*. 1998;19(6):589-601.
21. NIHR. National Standards for Public Involvement (website). 2018; Available from: <https://sites.google.com/nih.ac.uk/pi-standards/home>. Accessed 10 June 2018.

22. Sangha O, Stucki G, Liang MH, Fossel AH, Katz JN. The Self-Administered Comorbidity Questionnaire: a new method to assess comorbidity for clinical and health services research. *Arthritis Rheum.* 2003;49(2):156-63.
23. Heatherton TF, Kozlowski LT, Frecker RC, Fagerstrom KO. The Fagerstrom Test for Nicotine Dependence: a revision of the Fagerstrom Tolerance Questionnaire. *Br J Addict.* 1991;86(9):1119-27.
24. The Smoking Toolkit Study [cited 2018 September]. Available from: <http://www.smokinginengland.info/sts-documents/>.
25. Kotz D, Brown J, West R. Predictive validity of the Motivation To Stop Scale (MTSS): a single-item measure of motivation to stop smoking. *Drug Alcohol Depend.* 2013;128(1-2):15-9.
26. NCSCT. National Centre for Smoking Cessation and Training.; [cited 2018 September]. Available from: <https://www.ncsct.co.uk/usr/pub/Social%20support.pdf>.
27. Oakes W, Chapman S, Borland R, Balmford J, Trotter L. "Bulletproof skeptics in life's jungle": which self-exempting beliefs about smoking most predict lack of progression towards quitting? *Prev Med.* 2004;39(4):776-82.
28. Etter JF, Bergman MM, Humair JP, Perneger TV. Development and validation of a scale measuring self-efficacy of current and former smokers. *Addiction.* 2000;95(6):901-13.
29. Kaufman AR, Coa KI, Nguyen AB. Cigarette smoking risk-reducing beliefs: Findings from the United States Health Information National Trends Survey. *Preventive Medicine: An International Journal Devoted to Practice and Theory.* 2017;102:39-43.
30. UK CR. Lung Cancer Awareness Measure (Lung CAM) Toolkit. [Available from: [https://www.cancerresearchuk.org/sites/default/files/health\\_professional\\_lung\\_cam\\_toolkit\\_version\\_2.1\\_09.02.11.pdf](https://www.cancerresearchuk.org/sites/default/files/health_professional_lung_cam_toolkit_version_2.1_09.02.11.pdf)].
31. Kaufman AR, Koblitz AR, Persoskie A, Ferrer RA, Klein WM, Dwyer LA, et al. Factor Structure and Stability of Smoking-Related Health Beliefs in the National Lung Screening Trial. *Nicotine Tob Res.* 2016;18(3):321-9.
32. Van Buuren S, Boshuizen HC, Knook DL. Multiple imputation of missing blood pressure covariates in survival analysis. *Statistics in medicine.* 1999;18(6):681-94.
33. Caleyachetty A, Lewis S, McNeill A, Leonardi-Bee J. Struggling to make ends meet: exploring pathways to understand why smokers in financial difficulties are less likely to quit successfully. *Eur J Public Health.* 2012;22 Suppl 1:41-8.
34. Hiscock R, Judge K, Bauld L. Social inequalities in quitting smoking: what factors mediate the relationship between socioeconomic position and smoking cessation? *J Public Health (Oxf).* 2011;33(1):39-47.
35. Kotz D, Fidler J, West R. Factors associated with the use of aids to cessation in English smokers. *Addiction.* 2009;104(8):1403-10.
36. Levy DT, Romano E, Mumford E. The relationship of smoking cessation to sociodemographic characteristics, smoking intensity, and tobacco control policies. 2005:387-96, 2005 Jun.
37. Courtney RJ, Naicker S, Shakeshaft A, Clare P, Martire KA, Mattick RP. Smoking Cessation among Low-Socioeconomic Status and Disadvantaged Population Groups: A Systematic Review of Research Output. *Int J Environ Res Public Health.* 2015;12(6):6403-22.
38. Bryant J, Bonevski B, Paul C, McElduff P, Attia J. A systematic review and meta-analysis of the effectiveness of behavioural smoking cessation interventions in selected disadvantaged groups. 2011:1568-85, 2011 Sep.
39. Stead LF, Perera R, Bullen C, Mant D, Hartmann-Boyce J, Cahill K, et al. Nicotine replacement therapy for smoking cessation. *Cochrane Database Syst Rev.* 2012;11:CD000146.
40. Cawkwell PB, Blaum C, Sherman SE. Pharmacological Smoking Cessation Therapies in Older Adults: A Review of the Evidence. *Drugs Aging.* 2015;32(6):443-51.
41. Scholz J, Santos PC, Buzo CG, Lopes NH, Abe TO, Gaya PV, et al. Effects of aging on the effectiveness of smoking cessation medication. *Oncotarget.* 2016;7(21):30032-6.

42. Roddy E, Romilly N, Challenger A, Lewis S, Britton J. Use of nicotine replacement therapy in socioeconomically deprived young smokers: a community-based pilot randomised controlled trial. *Tob Control*. 2006;15(5):373-6.
43. Appel DW, Aldrich TK. Smoking cessation in the elderly. *Clin Geriatr Med*. 2003;19(1):77-100.
44. Siahpush M, McNeill A, Borland R, Fong GT. Socioeconomic variations in nicotine dependence, self-efficacy, and intention to quit across four countries: findings from the International Tobacco Control (ITC) Four Country Survey. *Tob Control*. 2006;15 Suppl 3:iii71-5.
45. Businelle MS, Kendzor DE, Reitzel LR, Costello TJ, Cofta-Woerpel L, Li Y, et al. Mechanisms linking socioeconomic status to smoking cessation: a structural equation modeling approach. *Health Psychol*. 2010;29(3):262-73.
46. Turner LR, Mermelstein R, Hitsman B, Warnecke RB. Social support as a moderator of the relationship between recent history of depression and smoking cessation among lower-educated women. *Nicotine Tob Res*. 2008;10(1):201-12.
47. Borland R, Yong HH, Balmford J, Fong GT, Zanna MP, Hastings G. Do risk-minimizing beliefs about smoking inhibit quitting? Findings from the International Tobacco Control (ITC) Four-Country Survey. *Preventive Medicine*. 2009;49(2-3):219-23.
48. Chapman S, Wong WL, Smith W. Self-exempting beliefs about smoking and health: differences between smokers and ex-smokers. *Am J Public Health*. 1993;83(2):215-9.
49. Peretti-Watel P, Halfen S, Gremy I. Risk denial about smoking hazards and readiness to quit among French smokers: an exploratory study. *Addict Behav*. 2007;32(2):377-83.
50. Office for National Statistics. Exploring the UK's digital divide. The scale of digital exclusion in the UK; those who aren't currently using the internet, how digital skills vary for different groups of the population and some of the barriers to digital inclusion.; 2019.
51. Ryan J, Lopian L, Le B, Edney S, Van Kessel G, Plotnikoff R, et al. It's not raining men: a mixed-methods study investigating methods of improving male recruitment to health behaviour research. *BMC Public Health*. 2019;19(1):814.
52. Setia MS. Methodology Series Module 3: Cross-sectional Studies. *Indian J Dermatol*. 2016;61(3):261-4.