

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/141841/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Jakovljevic, A., Sljivancanin Jakovljevic, T., Duncan, H. F., Nagendrababu, V., Jacimovic, J., Aminoshariae, A., Milasin, J. and Dummer, P. M. H. 2021. The association between apical periodontitis and adverse pregnancy outcomes: a systematic review. *International Endodontic Journal* 54 (9) , pp. 1527-1537. 10.1111/iej.13538

Publishers page: <https://doi.org/10.1111/iej.13538>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



# **The association between apical periodontitis and adverse pregnancy outcomes: a systematic review**

A. Jakovljevic<sup>1</sup>, T. Sljivancanin Jakovljevic<sup>2</sup>, H. F. Duncan<sup>3</sup>, V. Nagendrababu<sup>4</sup>, J. Jacimovic<sup>5</sup>, A. Aminoshariae<sup>6</sup>, J. Milasin<sup>7</sup>, & P. M. H. Dummer<sup>8</sup>

<sup>1</sup> Department of Pathophysiology, School of Dental Medicine, University of Belgrade, Belgrade, Serbia; <sup>2</sup> Department of Neonatology, Obstetrics and Gynecology Clinic "Narodni Front", Belgrade, Serbia; <sup>3</sup> Division of Restorative Dentistry & Periodontology, Dublin Dental University Hospital, Trinity College Dublin, Dublin, Ireland; <sup>4</sup> Department of Preventive and Restorative Dentistry, College of Dental Medicine, University of Sharjah, Sharjah, UAE; <sup>5</sup> Central Library, School of Dental Medicine, University of Belgrade, Belgrade, Serbia; <sup>6</sup> Department of Endodontics, School of Dental Medicine, Case Western Reserve University, Cleveland, Ohio, USA; <sup>7</sup> Department of Human Genetics, School of Dental Medicine, University of Belgrade, Belgrade, Serbia; and <sup>8</sup> School of Dentistry, College of Biomedical and Life Sciences, Cardiff University, Cardiff, UK

**Running title:** Apical periodontitis and pregnancy outcomes

**Keywords:** adverse pregnancy outcomes, apical periodontitis, low birth weight, preeclampsia, preterm birth, systematic review.

## **Correspondence:**

Aleksandar Jakovljevic, DDS, PhD

Department of Pathophysiology, School of Dental Medicine, University of Belgrade, dr Subotica 8, 11 000 Belgrade, Serbia

e-mail: [dr.sasuli@hotmail.com](mailto:dr.sasuli@hotmail.com), [a.jakovljevic@stomf.bg.ac.rs](mailto:a.jakovljevic@stomf.bg.ac.rs)

## **Conflict of Interest statement**

The authors have stated explicitly that there are no conflicts of interest in connection with this article.

## Introduction

Apical periodontitis is a chronic inflammatory process within the periapical tissues of teeth with an infected root canal system (Nair 2006). Although, it can be caused by several aetiological factors (e.g. physical, chemical, iatrogenic), it is generally accepted that microorganisms within the root canal system are the primary cause of pulp necrosis and the subsequent inflammatory reaction in the periapical region (Nair 2006). The principal radiographic feature of apical periodontitis is the destruction of periradicular tissues, evident as a radiolucency around the roots of the affected tooth. The destruction of the apical tissues is the consequence of a complex interplay between microorganisms and the activated innate and adaptive immune system of the host, as well as microbial by-products, e.g. virulence factors, cell-specific mediators (Márton & Kiss 2014).

Recent epidemiological data indicates a high global burden of apical periodontitis in the general adult population (Jakovljevic *et al.* 2020a), which correlates well with an increased global prevalence of untreated caries in the permanent dentition (Peres *et al.* 2019). A recent systematic review and meta-analysis on the prevalence of apical periodontitis (Jakovljevic *et al.* 2020a) revealed an increase in both endodontically treated and untreated teeth compared with a previous systematic review (Pak *et al.* 2012). These findings are in accordance with the continuous increase in the global age-standardized incidence of dental caries evident over the last 30 years (Kassebaum *et al.* 2015).

There is an increasing evidence linking general health and apical periodontitis, which emphasises the potential importance of oral health on general health (Murray & Saunders 2000, Segura-Egea *et al.* 2015). Although perceived as the local destruction of periodontal tissues, apical periodontitis has additional systemic inflammatory ramifications (Georgiou *et al.* 2019). A recent systematic review and meta-analysis revealed that apical periodontitis was associated with increased systemic inflammation, including increased immunoglobulin [Ig] A, IgM, IgG, C-reactive protein (CRP), interleukin (IL) 6, asymmetric dimethylarginine, C3 levels, amongst others (Georgiou *et al.* 2019). The likely cause of this generalised problem stems from an oral infection, which spreads through the blood system and activates the systemic immune response, leading to the development of generalized low-grade inflammation.

Previous studies have investigated the potential association between the presence or progression of apical periodontitis and several systemic diseases (Nagendrababu *et al.* 2020, Jakovljevic *et al.* 2020b). In an umbrella review, Nagendrababu *et al.* (2020) reported that diabetes mellitus was associated with a reduced outcome for root canal treatment and should be considered as a negative preoperative prognostic factor. In a separate umbrella review, Jakovljevic *et al.* (2020b) reported the existence of a weak association between apical periodontitis and cardiovascular diseases, with the weak association probably due to the small number of primary studies and the significant methodological inconsistencies between them.

Adverse pregnancy outcome (APO) is a broad term that encompasses several clinical outcomes (Athukorala *et al.* 2010, Lean *et al.* 2017, Pinheiro *et al.* 2019, Søndergaard *et al.* 2020), including (i) stillbirth (defined as intrauterine death of a child after 20 weeks of gestation or weighing  $\geq 350$  g if gestational age is unknown), (ii) small for gestational age (SGA) (defined as a birthweight below 10<sup>th</sup> percentile adjusted for gestational age) with or without intrauterine growth restriction (IUGR), (iii) neonatal death, (iv) low birth weight (LBW) (<2500 g) or very low birth weight (VLBW) (<1500 g), (v) admissions to neonatal intensive care unit (NICU), (vi) neonatal acidosis (umbilical artery pH <7.0–7.2), (vii) pre-eclampsia (PE) (viii) placental abruption, (ix) preterm birth (PTB) (<37 weeks gestation, or very preterm <32 weeks) and/or (x) gestational diabetes mellitus (GDM). APOs are a significant public health problem with considerable personal, social and financial implications worldwide (Kramer 2003, Poon *et al.* 2018).

Mechanistically during pregnancy, there is a shift from T helper (Th)1 and Th17 towards a Th2 and T regulatory cell immune response, which occurs both in the peripheral blood and at the foeto-maternal interface. It has been reported that any disturbance of this immune response increases significantly the risk of APOs occurrence (Sykes *et al.* 2012). In addition, APOs are significantly associated with elevated maternal local and systemic inflammatory mediators, which might be accompanied by intrauterine infections (Sykes *et al.* 2012). As a result, previous oral investigations have hypothesized that periodontal disease is a significant risk factor for the development of APOs. This link can be explained by two major mechanisms: (i) direct - the translocation of periodontal pathogens to the foeto-placental unit via haematogenous dissemination, and/or (ii) indirect, i.e. via the effect of inflammatory mediators on the foetal-placental unit (Figuro *et al.* 2020). Moreover, a recent umbrella review revealed positive

associations between periodontal disease and preterm birth (relative risk (RR) = 1.6; 95% confidence interval (CI) (1.3-2.0)), low birth weight (RR=1.7, 95% CI (1.3-2.1)), and preeclampsia (RR=2.2, 95% CI (1.4-3.4)) (Daalderop *et al.* 2018).

Apical periodontitis and periodontal disease share a similar inflammatory response (Gomes *et al.* 2015, Das *et al.* 2020), and many systemic diseases linked to periodontal disease have shown comparable associations with apical periodontitis (e.g. diabetes mellitus [Nagendrababu *et al.* 2020]). As apical periodontitis is not just a local event, it was reasonable to investigate whether generalized low-grade inflammation and/or bloodstream infection could potentially contribute to APOs. Notably, the findings of the existing primary studies appear inconsistent (Harjunmaa *et al.* 2015, Leal *et al.* 2015, Khalighinejad *et al.* 2017); however, the association of APOs and apical periodontitis has not been examined rigorously or reviewed. Therefore, this systematic review aimed to critically evaluate the available evidence on the association of maternal apical periodontitis with several APOs.

## **Materials and Methods**

This systematic review was reported according to the principles recommended by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Protocols (PRISMA) statement (Moher *et al.* 2009). The protocol of the review was registered *a priori* with the PROSPERO international prospective register of systematic reviews (CRD42020191987).

### *Review question*

In pregnant women (P), does the presence (I) or absence (C) of apical periodontitis modify the prevalence of adverse pregnancy outcomes (O) assessed from observational longitudinal clinical trials, cohort, case-control, and cross-sectional studies (S)?

### *Eligibility criteria*

#### *Inclusion criteria*

- Pregnant women.
- Previously healthy women (American Society of Anesthesiology I or II) diagnosed with APOs compared with uncomplicated pregnancies.
- Prevalence, confirmed radiographically, of apical periodontitis associated with or without root filled teeth.
- Observational studies including longitudinal clinical trials, cohort, case-control, or cross-sectional studies of prospective and retrospective design.

#### *Exclusion criteria*

- Articles presenting duplicate or overlapping results, abstract-only papers, case reports, case series, animal studies and reviews.

#### *Literature search process*

To identify all types of studies that examined an association between apical periodontitis and adverse pregnancy outcomes, several international and regional databases were searched systematically. Clarivate Analytics' Web of Science (including Web of Science Core Collection - WoS, Korean Journal Database - KJD, Russian Science Citation Index - RSCI, SciELO Citation Index - SciELO) [1980-2020], Scopus [1960-2020], PubMed [1964-2020], and Cochrane Central Register of Controlled Trials (CENTRAL) [1996-2020] were explored up to 25 February 2021, without language restrictions. The basic search strategy, developed on the formulated research question, was used for preliminary searches to avoid duplication and identify possible previously published systematic reviews, to validate the proposed idea and identify relevant articles, to determine controlled vocabulary (e.g. Medical Subject Headings – MeSH), free-text, or synonymous key terms for apical periodontitis and adverse pregnancy outcomes, and to evaluate and develop the most optimum information retrieval approach. The central search strategy was modified according to the specific characteristics of each selected database to increase sensitivity, using various combinations of previously identified keywords,

Boolean, truncation, and proximity operators. The detailed electronic search strategy for each database is presented in Supplemental Table 1. Furthermore, to identify unpublished manuscripts, research reports, conference papers, doctoral dissertations, and other grey literature, additional searches of available digital repositories (e.g. OpenGrey, Networked Digital Library of Theses and Dissertations, Open Access Theses and Dissertations, DART-Europe E-theses Portal – DEEP, Opening access to UK theses – EThOS) and Google Scholar (first 100 returns) were performed. Finally, to assure the reliability of the data collected, the electronic search was further supplemented with additional citation searching through the reference lists of identified studies and relevant reviews. To ensure the inclusion of eligible studies that have not yet been indexed by the databases, the most current issues and articles accepted for publication in journals which published key articles (*International Endodontic Journal*, *Journal of Endodontics*, *Australian Endodontic Journal*, *Journal of Dental Research*, *Clinical Oral Investigations*, and *Archives of Oral Biology*) were also considered. Additional search during the final drafting of the paper indicated no new relevant studies had been published after completion of the literature search.

For duplicate removal and screening of search results, all records obtained were exported automatically from the used databases and imported into the Rayyan (Qatar Computing Research Institute (Data Analytics), Doha, Qatar) (Ouzzani *et al.* 2016), a free web app (<https://rayyan.qcri.org/welcome>) aimed to facilitate collaboration among reviewers, to expedite the initial process of title/abstract screening and record the study eligibility decisions and exclusion reasons. The selection of studies included in the systematic review was performed using a two-stage screening process. To identify potential primary studies during the first phase, two independent reviewers (T.S.J. and J.J) performed the initial screening of titles and abstracts of previously identified articles, documenting reasons for exclusion. In the second screening, reviewers considered the full texts of studies that were classified as eligible in the first phase. Disagreement between the two independent reviewers was resolved by discussion with the third reviewer (V.N.). All studies that did not meet the inclusion criteria were excluded from the analysis.

### *Data extraction*

The following details were extracted independently by two reviewers (T.S.J. and A.J.) from each study included in the final review: name of the first author, year published, type of study design, total number of participants with age distribution, population characteristics, exposure evaluation method, investigated outcomes, outcome evaluation method, main results, adjustment, limitations and quality of included studies. Disagreements were resolved by discussing with a third reviewer (V.N.). All extracted data were stored in tables created using Microsoft Office software (Microsoft Inc., Redmond, WA, USA).

### *Quality of studies*

The Newcastle-Ottawa Scale (NOS) (Wells *et al.* 2019) and theorem of the NOS adapted for cross-sectional studies (Patra *et al.* 2015), were used to assess the risk of bias for case-control and cross-sectional studies, respectively. Critical appraisal of the included studies was performed by two independent reviewers (A.A. and A.J.) using the NOS star rating system, where each study was evaluated for sample selection, comparability of the groups, and the outcome assessment. Studies with 7-9 and 4-6 stars assigned were considered to be of "Good" and "Fair" quality, respectively, while studies with less than 3 stars were regarded as "Poor" quality studies (McPheeters *et al.* 2012). Disagreements during the assessment were discussed and resolved by a third reviewer (V.N.).

### *Data synthesis (qualitative synthesis)*

The conceptual framework for data synthesis was developed using a modified narrative synthesis approach (Popay *et al.* 2005). Due to the differing primary and secondary outcomes of the included studies, the extracted data were narratively synthesized through textual descriptions and the development of a preliminary synthesis of the findings from the included studies, examining relationships in the data, and the evaluation of synthesis robustness.

## **Results**

### *Literature search and characteristics of the included studies*

The study selection process is summarised in Figure 1. The literature search of the chosen databases and other relevant sources retrieved a total of 523 records for potential inclusion in the



systematic review. After 49 duplicates were removed, 470 studies were excluded, while four were eligible for full-text assessment. Finally, one study was excluded because the outcomes studied were placental and systemic markers rather than adverse pregnancy complications (Harjunmaa *et al.* 2018), while three studies were included in the current review (Harjunmaa *et al.* 2015, Leal *et al.* 2015, Khalighinejad *et al.* 2017). The characteristics of the included clinical studies are described in Table 1. The two case-control and one cross-sectional study were carried out in Africa, Brazil (Harjunmaa *et al.* 2015, Leal *et al.* 2015) and USA (Khalighinejad *et al.* 2017), respectively. The included studies were published between 2015 and 2017. A total of 1187 individuals participated in the three clinical studies, with an approximate age range of 15 to 40 years. The included studies reported different outcomes, hence it was not possible to perform a meta-analysis.

### *Quality of studies*

One study did not report an adequate sample size calculation (Khalighinejad *et al.* 2017), and the other one did not specify the criteria for the evaluation of apical periodontitis (Harjunmaa *et al.* 2015). Moreover, in these studies investigators were not blinded during the evaluations undertaken (Supplemental Tables 2, 3). Therefore, the overall quality of the evidence for these investigations was considered as “Fair” (Supplemental Tables 2, 3).

On the other hand, the study performed by Leal *et al.* (2015) included an adequate sample size calculation, and reported the radiographic criteria for the evaluation of apical periodontitis. Also, the authors excluded potential confounding groups (smoking, infection during pregnancy, diabetes mellitus, stillbirths with less than 28 weeks or serious physical defects and periodontal disease) and performed adequate statistical analysis. Therefore, the overall quality of the evidence for this study was considered as “Good” (Supplemental Table 3).

### *Principal findings*

#### *Pregnancy duration, birthweight, length and head circumference of their infants, preterm birth*

Pregnant women with apical periodontitis had a significantly shorter mean pregnancy duration and delivered infants with lower birth rate and shorter neonatal length and head circumference than women without periapical infection ( $P = 0.014$ ,  $P = 0.019$ ,  $P = 0.002$ ,  $P =$

0.033, respectively). The incidence of preterm birth was higher (10.0%) in pregnant women with periapical infection compared to women without periapical infection (7.3%) (Harjunmaa *et al.* 2015).

The univariate analysis of the association of the variables indicates that women with apical periodontitis were five times more likely to deliver a child with low-birth weight and have a preterm birth than women without periapical lesions (crude OR, 4.80; 95% CI, 1.55–14.81). After the multivariate analysis the presence of periapical lesions in postpartum women remained associated as a risk factor for low-birth weight and preterm birth (adjusted OR, 3.52; 95% CI, 1.01–12.32). (Leal *et al.* 2015)

### *Preeclampsia*

Pregnant women with apical periodontitis were significantly more susceptible to develop preeclampsia than women without apical periodontitis (Odds ratio (OR) = 2.49; 95% CI (1.1–5.62),  $P = 0.002$ ) (Khalighinejad *et al.* 2017).

## **Discussion**

The latest epidemiological data suggest that APOs are an increasingly important global health problem (Chawanpaiboon *et al.* 2019, Blencowe *et al.* 2019). There were almost 15 million worldwide preterm live births in 2014 (Chawanpaiboon *et al.* 2019) and 20.5 million new-born children in 2015 had a birthweight of less than the threshold 2500 g (Blencowe *et al.* 2019). A previous systematic review reported that the model-based incidence of preeclampsia was 4.6% for all deliveries, with wide variation across regions of the world (Abalos *et al.* 2013). In addition, APOs remain a significant cause of maternal and foetal morbidity and/or mortality, especially in low- and middle-income countries (Chawanpaiboon *et al.* 2019, Blencowe *et al.* 2019). Separately, a recent systematic review with meta-analysis also revealed an increased prevalence of apical periodontitis in the global adult population (Jakovljevic *et al.* 2020a). Despite evidence linking periodontal disease and APOs (Pitiphat *et al.* 2008), the importance of endodontic disease during pregnancy is insufficiently recognized by dental and obstetric health professionals. Therefore, given the global disease burden of apical periodontitis and APOs, it is

important to clarify their potential association in order to develop adequate preventive and therapeutic strategies at earlier stages of pregnancy in the future.

The aetiology of APOs is multifactorial, with several risk factors potentially contributing to their development (e.g. heredity, environmental, nutritional, lifestyle, socio-economic, and foetal-related factors) (Lawn *et al.* 2016, Escañuela Sánchez *et al.* 2019). In addition, the development of APOs is significantly associated with elevated parameters of local and systemic inflammation with or without various intra-uterine infections (Sykes *et al.* 2012, Figuero *et al.* 2020). Importantly, altered levels of female sex hormones during pregnancy increases vascular permeability, which leads to the spreading of inflammation with possible haematogenous dissemination of infection (Sykes *et al.* 2012, Figuero *et al.* 2020).

Previous investigations hypothesized that severe periodontitis is a potential novel risk factor for APOs (Madianos *et al.* 2013, Bobetsis *et al.* 2020). This association is explained by the fact that both conditions are correlated with microbial infections and increased levels of local and systemic inflammatory mediators (Madianos *et al.* 2013, Bobetsis *et al.* 2020,). In this context, consistent evidence from previous systematic reviews with a low risk of bias indicates that pregnant women with periodontal disease are at increased risk of developing preeclampsia (Sgolastra *et al.* 2013) and delivering preterm and/or low birth weight newborns (Corbella *et al.* 2016). These findings were confirmed in an umbrella review (Daalderop *et al.* 2018).

In general, apical periodontitis and marginal periodontitis share a similar inflammatory response (Gomes *et al.* 2015, Das *et al.* 2020). Both conditions are characterized by systemic low-grade inflammation and potential microbial dissemination to remote organs (Hasturk & Kantarci 2015, Georgiou *et al.* 2019). Moreover, many systemic diseases linked to periodontal disease have similar associations with apical periodontitis (e.g. diabetes mellitus). From the above results, it is interesting to note that, so far, only a limited number of studies have investigated the potential association between the presence of apical periodontitis and the development of APOs in pregnant women (Harjunmaa *et al.* 2015, Leal *et al.* 2015, Khalighinejad *et al.* 2017). Additionally, their association has not previously been analysed systematically.

This systematic review for the first time critically evaluated the potential association between apical periodontitis and APOs. Based on strict eligibility criteria, only three studies were included in the review (Harjunmaa *et al.* 2015, Leal *et al.* 2015,, Khalighinejad *et al.* 2017).

The qualitative synthesis of data indicates that pregnant women with apical periodontitis were significantly more susceptible to develop preeclampsia (Khalighinejad *et al.* 2017), had a remarkably shorter mean pregnancy duration and delivered infants with a lower birth rate, shorter neonatal length, and head circumference (Harjunmaa *et al.* 2015, Leal *et al.* 2015,) than women without periapical disease. These results are in accordance with previous systematic reviews that investigated the association between periodontal disease and APOs (Sgolastra *et al.* 2013, Corbella *et al.* 2016). However, the NOS was categorised as “Fair” quality for two out of three included studies (Harjunmaa *et al.* 2015, Khalighinejad *et al.* 2017),, and caution should be exercised in interpreting the results of this systematic review because many covariates were not considered in the original investigations (i.e. oral hygiene, periodontal status, smoking, body mass index, alcohol intake, quality of restorations, number of carious lesions, cracked teeth, trauma, etc.). Additionally, two out of the three included primary studies (Harjunmaa *et al.* 2015, Khalighinejad *et al.* 2017) had many methodological inconsistencies and flaws (e.g. unjustified sample size, an unspecified method for evaluation of apical periodontitis, using the same sample in two investigations, etc.) that render the conclusions as preliminary with a low level of evidence in these cases

On the other hand, using the NOS evaluation scale, the study performed by Leal *et al.* (2015) was categorised as “Good” quality supporting the fact, based on the high level of evidence, that apical periodontitis in a multivariate analysis is associated with low-birth weight and preterm birth. Leal *et al.* (2015) effectively controlled the potential confounders, calculated sample size adequately, used an appropriate radiographic method for assessing the periapical status, and performed suitable statistical analysis.

Moreover, previous biological experimentations in animal models also provide evidence to support a potential link between maternal apical periodontitis and APOs. Experiments on Sprague–Dawley rats (Bain *et al.* 2009, 2013) concluded that the development of uncontrolled gestational diabetes mellitus, increased blood glucose and serum levels of insulin, Th-1 pro-inflammatory cytokines, myelin basic protein, and norepinephrine concentrations in pregnant rats with periapical abscess compared to a control group of animals. Based on their experiments, Bain *et al.* (2009, 2013) suggested that maternal periapical inflammation could serve as a modifiable risk factor of APOs. In another animal model, the pulp chambers of first molars of C57BL/6J mice were directly infected with the w83 strain of *Porphyromonas gingivalis* (Ao *et*

*al.* 2015). The authors demonstrated significant preterm birth and low birth weight in infected mice compared to the control group. They also, immunohistochemically confirmed the translocation of *P. gingivalis* to placental tissues, histologically assessed defects in placental tissue, and found increased circulating and local pro-inflammatory markers (tumour necrosis factor – alpha [TNF- $\alpha$ ], IL-1 $\beta$ , IL-6, and IL-17) (Ao *et al.* 2015). Similarly, based on this animal model, the authors concluded that dental infections represented a predisposing factor for preterm birth and low birth weight. It is important to note here that rodent models provide only surrogate findings that have to be taken in context and may not reflect the likely outcome in pregnant human females.

An association between apical periodontitis and impairment of general health remains a matter of debate. As the highest level of secondary evidence, only two umbrella reviews dealt with the potential bi-directional association between apical periodontitis and systemic diseases. Nagendrababu *et al.* (2020) revealed that diabetes mellitus is associated with a reduced outcome for root canal treatment, while Jakovljevic *et al.* (2020b) demonstrated the existence of a weak association between apical periodontitis and cardiovascular diseases. The principle problems facing systematic reviews in this field are related to methodological shortcomings of the available primary studies, including inappropriate study designs, unjustified sample sizes, unmatched study groups, inadequate markers and/or surrogate models, unadjusted confounding factors, incomparable results, amongst others. Other important problems are related to ethical considerations of the most appropriate study designs in humans and eventual translation of findings revealed in experimental animal models to humans. Promising results were reported in the most recent prospective longitudinal interventional study (Poornima *et al.* 2020) who investigated the impact of root canal treatment on serum high-sensitivity C-reactive protein (hsCRP) levels in adults with apical periodontitis. The authors concluded that root canal treatment reduced serum hsCRP levels in individuals with apical periodontitis, suggesting that root canal treatment could influence inflammation in the human body.

The current systematic review has several limitations, comprising

- (i) a small number of included studies,
- (ii) two included studies were case-control and one was a cross-sectional study,
- (iii) the unjustified sample size in one included study,
- (iv) unspecified method for evaluation of apical periodontitis in two included studies, and

- (v) lack of control for periodontal diseases, oral hygiene, caries, and smoking as confounding factors for the development of apical periodontitis were not adjusted for two out of three included studies.

However, the following parameters were considered as the strengths of the current review:

- (i) an *a priori* developed and registered protocol in the PROSPERO database,
- (ii) a comprehensive literature search performed with no language restriction in four electronic databases, including the grey literature,
- (iii) literature search and data extraction process conducted by two independent reviewers, and
- (iv) critical appraisal of included studies using the NOS, also conducted by two reviewers independently.

## **Conclusion**

This investigation for the first time systematically reviewed the potential association between apical periodontitis and APOs, and clearly highlighted a significant gap in the endodontic literature related to this issue. Based on a limited volume and “Fair” and “Good” quality of evidence, a positive association between maternal apical periodontitis and APOs was observed. In future, more “Good” quality clinical studies are required to confirm the results of the current systematic review.

## References

- Abalos E, Cuesta C, Grosso AL, Chou D, Say L (2013) Global and regional estimates of preeclampsia and eclampsia: a systematic review. *European Journal of Obstetrics & Gynecology and Reproductive Biology* **170**, 1-7.
- Ao M, Miyauchi M, Furusho H *et al.* (2015) Dental Infection of Porphyromonas gingivalis Induces Preterm Birth in Mice. *PLoS One* **10**, e0137249.
- [Athukorala C, Rumbold AR, Willson KJ, Crowther CA \(2010\) The risk of adverse pregnancy outcomes in women who are overweight or obese. \*BMC Pregnancy and Childbirth\* \*\*10\*\*, 56.](#)
- Bain JL, Lester SR, Henry WD, Naftel JP, Johnson RB (2009) Effects of induced periapical abscesses on rat pregnancy outcomes. *Archives of Oral Biology* **54**, 162-71.
- Bain JL, Lester SR, Henry WD, Pongetti JL, Blackman ME, Johnson RB (2013) Association between maternal periapical lesions and brain inflammation in rat pups. *Archives of Oral Biology* **58**, 266-71.
- Blencowe H, Krasevec J, de Onis M *et al.* (2019) National, regional, and worldwide estimates of low birthweight in 2015, with trends from 2000: a systematic analysis. *The Lancet Global Health* **7**, e849-e860.
- Bobetsis YA, Graziani F, Gürsoy M, Madianos PN (2020) Periodontal disease and adverse pregnancy outcomes. *Periodontology 2000* **83**, 154-74.
- Chawanpaiboon S, Vogel JP, Moller AB *et al.* (2019) Global, regional, and national estimates of levels of preterm birth in 2014: a systematic review and modelling analysis. *The Lancet Global Health* **7**, e37-e46.
- Corbella S, Taschieri S, Del Fabbro M, Francetti L, Weinstein R, Ferrazzi E (2016) Adverse pregnancy outcomes and periodontitis: A systematic review and meta-analysis exploring potential association. *Quintessence International* **47**, 193-204.
- Daalderop LA, Wieland BV, Tomsin K *et al.* (2018) Periodontal Disease and Pregnancy Outcomes: Overview of Systematic Reviews. *JDR Clinical & Translational Research* **3**, 10-27.
- Das AC, Sahoo SK, Parihar AS, Bhardwaj SS, Babaji P, Varghese JG (2020) Evaluation of role of periodontal pathogens in endodontic periodontal diseases. *Journal of Family Medicine and Primary Care* **9**, 239-42.

- Escañuela Sánchez T, Meaney S, O'Donoghue K (2019) Modifiable risk factors for stillbirth: a literature review. *Midwifery* **79**, 102539.
- Figuero E, Han YW, Furuichi Y (2020) Periodontal diseases and adverse pregnancy outcomes: Mechanisms. *Periodontology 2020* **83**, 175-88.
- Georgiou AC, Crielaard W, Armenis I, de Vries R, van der Waal SV (2019) Apical periodontitis is associated with elevated concentrations of inflammatory mediators in peripheral blood: a systematic review and meta-analysis. *Journal of Endodontics* **45**, 1279–95.
- Gomes BP, Berber VB, Kokaras AS, Chen T, Paster BJ (2015) Microbiomes of Endodontic-Periodontal Lesions before and after Chemomechanical Preparation. *Journal of Endodontics* **41**, 1975-84.
- Harjunmaa U, Doyle R, Järnstedt J *et al.* (2018) Periapical infection may affect birth outcomes via systemic inflammation. *Oral Diseases* **24**, 847-55.
- Harjunmaa U, Järnstedt J, Alho L *et al.* (2015) Association between maternal dental periapical infections and pregnancy outcomes: results from a cross-sectional study in Malawi. *Tropical Medicine & International Health* **20**, 1549-58.
- Hasturk H, Kantarci A (2015) Activation and resolution of periodontal inflammation and its systemic impact. *Periodontology 2000* **69**, 255-73.
- Jakovljevic A, Nikolic N, Jacimovic J *et al.* (2020a) Prevalence of Apical Periodontitis and Conventional Nonsurgical Root Canal Treatment in General Adult Population: An Updated Systematic Review and Meta-analysis of Cross-sectional Studies Published between 2012 and 2020. *Journal of Endodontics* **46**, 1371-86.e8.
- Jakovljevic A, Duncan HF, Nagendrababu V, Jacimovic J, Milasin J, Dummer PMH (2020b) Association between cardiovascular diseases and apical periodontitis: an umbrella review. *International Endodontic Journal* **53**, 1374-86.
- Kamer MS (2003) The epidemiology of adverse pregnancy outcomes: an overview. *Journal of Nutrition* **133**, 1592S-6S.
- Kassebaum NJ, Bernabé E, Dahiya M, Bhandari B, Murray CJ, Marcenes W (2015) Global burden of untreated caries: a systematic review and metaregression. *Journal of Dental Research* **94**, 650-8.
- Khalighinejad N, Aminoshariae A, Kulild JC, Mickel A (2017) Apical Periodontitis, a Predictor Variable for Preeclampsia: A Case-control Study. *Journal of Endodontics* **43**, 1611-4.



- Lawn JE, Blencowe H, Waiswa P *et al.* (2016) Stillbirths: rates, risk factors, and acceleration towards 2030. *Lancet* **387**, 587-603.
- Leal ASM, de Oliveira AEF, Brito LMO *et al.* (2015) Association between chronic apical periodontitis and low-birth-weight preterm births. *Journal of Endodontics* **41**, 353–7.
- Lean SC, Derricott H, Jones RL, Heazell AEP (2017) Advanced maternal age and adverse pregnancy outcomes: A systematic review and meta-analysis. *PLoS One* **12**, e0186287.
- Madianos PN, Bobetsis YA, Offenbacher S (2013) Adverse pregnancy outcomes (APOs) and periodontal disease: pathogenic mechanisms. *Journal of Clinical Periodontology* **40**, S170-80.
- Márton IJ, Kiss C (2014) Overlapping protective and destructive regulatory pathways in apical periodontitis. *Journal of Endodontics* **40**, 155–63.
- McPheeters ML, Kripalini S, Peterson NB *et al.* (2012) Quality improvement interventions to address health disparities. Evidence report/technology assessment. Rockville (MD): Agency for Healthcare Research and Quality (US). URL <http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0049222/pdf/TOC.pdf>. [Accessed on 7 November 2020]
- Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Annals of Internal Medicine* **151**, 264–9.
- Murray CA, Saunders WP (2000) Root canal treatment and general health: a review of the literature. *International Endodontic Journal* **33**, 1-18.
- Nagendrababu V, Segura-Egea JJ, Fouad AF, Pulikkotil SJ, Dummer PMH (2020) Association between diabetes and the outcome of root canal treatment in adults: an umbrella review. *International Endodontic Journal* **53**, 455–66.
- Nair PN (2006) On the causes of persistent apical periodontitis: a review. *International Endodontic Journal* **39**, 249–81.
- Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A (2016) Rayyan: A web and mobile app for systematic reviews. *Systematic Reviews* **5**, 210.
- Ørstavik D, Kerekes K, Eriksen HM (1986) The periapical index: a scoring system for radiographic assessment of apical periodontitis. *Endodontics & Dental Traumatology* **2**, 20-34.
- Pak JG, Fayazi S, White SN (2012) Prevalence of periapical radiolucency and root canal treatment: a systematic review of cross-sectional studies. *Journal of Endodontics* **38**, 1170–6.

Patra J, Bhatia M, Suraweera W *et al.* (2015) Exposure to second-hand smoke and the risk of tuberculosis in children and adults: a systematic review and meta-analysis of 18 observational studies. *PLoS Medicine* **12**, e1001835.

Peres MA, Macpherson LMD, Weyant RJ *et al.* (2019) Oral diseases: a global public health challenge. *Lancet* **394**, 249–60.

Pinheiro RL, Areia AL, Mota Pinto A, Donato H (2019) Advanced Maternal Age: Adverse Outcomes of Pregnancy, A Meta-Analysis. *Acta Médica Portuguesa* **32**, 219-26.

Pitiphat W, Joshipura KJ, Gillman MW, Williams PL, Douglass CW, Rich-Edwards JW (2008) Maternal periodontitis and adverse pregnancy outcomes. *Community Dentistry and Oral Epidemiology* **36**, 3-11.

Poon LC, McIntyre HD, Hyett JA, da Fonseca EB, Hod M; FIGO Pregnancy and NCD Committee (2018) The first-trimester of pregnancy - A window of opportunity for prediction and prevention of pregnancy complications and future life. *Diabetes Research and Clinical Practice* **145**, 20-30.

Poornima L, Ravishankar P, Abbott PV, Subbiya A, PradeepKumar AR (2020) Impact of root canal treatment on high-sensitivity C-reactive protein levels in systemically healthy adults with apical periodontitis - a preliminary prospective, longitudinal interventional study. *International Endodontic Journal* [Epub ahead of print] doi: 10.1111/iej.13444.

Popay J, Roberts H, Sowden A *et al.* (2005) Developing guidance on the conduct of narrative synthesis in systematic reviews. *Journal of Epidemiology and Community Health* **59**, A7.

Segura-Egea JJ, Martín-González J, Castellanos-Cosano L (2015) Endodontic medicine: connections between apical periodontitis and systemic diseases. *International Endodontic Journal* **48**, 933-51.

Sgolastra F, Petrucci A, Severino M, Gatto R, Monaco A (2013) Relationship between periodontitis and pre-eclampsia: a meta-analysis. *PLoS One* **8**, e71387.

Sykes L, MacIntyre DA, Yap XJ, Teoh TG, Bennett PR (2012) The Th1:th2 dichotomy of pregnancy and preterm labour. *Mediators of Inflammation* **2012**, 967629.


Søndergaard MM, Hlatky MA, Stefanick ML *et al.* (2020) Association of Adverse Pregnancy Outcomes With Risk of Atherosclerotic Cardiovascular Disease in Postmenopausal Women. *JAMA Cardiology* **5**, 1390–8.

Wells GA, Shea B, O'Connell D *et al.* (2019) The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Available at: [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp). Accessed November 7, 2020.

## Legends

Figure 1. PRISMA flow diagram of the study search and identification of relevant studies.

Table 1. The characteristics, main results, limitations and quality of included studies in the systematic review



**Table 1. The characteristics, main results, limitations and quality of included studies in the systematic review**

Authors, year	Study design	Number of participants  / mean age  ± SD	Population characteristics	Exposure evaluation  method/AP  definition	Investigated outcomes	Outcome evaluation  method	Main results	Adjustment	Limitations	Quality of studies
Harjunmaa <i>et al.</i> 2015	Cross- sectional	1024/  25±6.2	A rural population from Mangochi district in Malawi, Africa	Digital panoramic radiographs/  Not specified	Continuous birth outcomes (duration of pregnancy, birthweight, neonatal weight-for-age, length-for-age, head circumference-for- age);  Dichotomous birth outcomes (Incidence of preterm birth, low birthweight, prevalence of neonatal underweight,	Anthropometric measurements were gained with an electronic scale and plastic tape; standardized indices were assessed using the WHO Child Growth Standards.	Women with AP had mean (95% CI) pregnancy duration 0.4 weeks (0.1– 0.8) shorter and infants with 79 g (13–145) lower BW, 0.5 cm (0.2–0.9) shorter neonatal length, 0.27 units (0.11– 0.44) lower LA and 0.18 units (0.01–0.35) smaller head circumference z-score, increased prevalence of	Maternal age, maternal height, BMI, HIV status, malaria status and anaemia at enrolment, number of previous pregnancies, study site, socio- economic score, periodontitis, number of teeth, time between delivery and examination, and	Method evaluating AP is not specified.	Fair*

					neonatal stunting,  small head  circumference)		neonatal stunting  (adjusted RR = 1.68;  P=0.007) and small  head circumference  (adjusted RR = 2.52;  P=0.012) compared to  women without the  AP.	intervention.  Patients were not  smokers.		
Leal <i>et al.</i>  2015	Case -  control	Cases - 33  mothers who  had preterm  infants  (more than  27 and  less than 37  gestation  weeks) and  weighing  less than  2500 g./  between 15	An urban  population  from São Luís,  Maranhão,  Brazil.	A full-mouth  set of  periapical  radiographs/T  he periapical  index  introduced by  Ørstavik et al.  (1986).	The presence of  radiographically  assessed periapical  lesions in women  with low-birth-weight  preterm births  compared to women  without pregnancy  complications.	The periapical  and endodontic  status in both  groups was  evaluated using a  full-mouth set of  periapical  radiographs in  postpartum period  and analysed by 2  previously  calibrated  endodontists who	The univariate analysis  of association of the  variables indicates that  women with AP had  about 5 times more  odds of presenting  LBWPB than women  without periapical  lesion  (crude OR, 4.80; 95%  CI, 1.55–14.81). After  the multivariate  analysis the presence	Women with  diabetes mellitus,  multiple  pregnancies, and  with the presence of  periodontitis were  excluded from the  sample. Stillbirths  with less than 28  weeks or serious  physical defects  that could affect the  weight or survival	-	Good**

		and 40 years of age Controls – 30 mothers of newborns at term (more than 37 and less than 42 gestation weeks) and weighing more than 2500 g. / between 15 and 40 years of age				were blinded and had more than 5 years of clinical experience.	of periapical lesion in postpartum women remained associated as a risk factor for LBWPB (adjusted OR, 3.52; 95% CI, 1.01–12.32),	of the newborn were also excluded.		
Khalighinejad et al. 2017	Case-	50 in	An urban	digital	Endodontic variables	The periapical	AP was significantly	Maternal AP	Sample size not	Fair**

control	experimenta	population	panoramic	(presence/absence of	and endodontic	more common in PE	maternal	justified.
	l group	form	radiographs/	AP, average number	status in both	compared to healthy	periodontitis, and	Presence of
	(preeclampsia)/	Cleveland, OH,	The	of teeth with AP,	groups was	pregnancies (OR =	endodontic	smokers and
	26 ± 3.2	USA	periapical	presence/absence of	evaluated using	2.49; 95% CI, 1.1–	treatment.	women with
			index	endodontically	digital panoramic	5.62). PE group has a		diagnosed
	50 in control		introduced by	treated tooth)	radiographs that	significantly higher		periodontitis.
	group		Ørstavik <i>et</i>		were taken before	number of teeth with		
	(women		<i>al.</i> (1986)		pregnancies	AP compared to		
	with an					control (P = 0.001).		
	uncomplicat					Binary logistic		
	ed course of					regression model		
	pregnancy) /					showed the highly		
	24 ± 2.8					significant association		
						of AP with PE was		
						(adjusted OR = 2.23;		
						95% CI, 1.92–6.88).		

\* Evaluated with theorem of the Newcastle-Ottawa Scale (NOS) adapted for cross-sectional studies (Patra et al. 2015); \*\* Evaluated with the NOS for case-control studies (Wells et al. 2019); AP, apical periodontitis; PAI, periapical index; USA, Unites States of America; PCR, Polymerase Chain Reaction; WHO, World Health Organization; OR, odds ratio, CI, confidence interval; PE, preeclampsia; BMI, body mass index; HIV; human immunodeficiency virus; BW, birth weight; gw, gestational week; LA, length-for-age; SD, standard deviation, LBWPB, low birth weight and preterm birth.



Supplemental Table 1. Electronic Databases and Search Strategy

Database (n)		Search strategy #1 AND #2
WoS (n=33)	#1	TS = (((apical OR periapical OR periradicular OR radicular OR pulp*) NEAR/1 (periodontitis OR disease\$ OR abscess* OR granuloma\$ OR infection\$
KJD (n=0)		OR lesion\$ OR patho* OR inflammat* OR condition\$ OR process*)) OR pulpitis OR root canal OR endodont*) (n=33,933)
RSCI (n=0)	#2	TS = (((((pregnan* OR birth) NEAR/1 (outcome\$ OR complication\$)) OR ((premature OR preterm) NEAR/1 birth) OR pre-Eclampsia OR
SCIELO (n=0)		preeclampsia OR (hypertension\$ NEAR/1 (gestational OR pregnancy)) OR ((fetal OR intrauterine) NEAR/1 growth NEAR/1 (retardation OR
		restriction)) OR low birth weight OR birthweight OR chorioamnionitis OR stillbirth OR neonatal sepsis OR (duration NEAR/1 pregnancy) OR
		((neonatal OR birth) NEAR/1 (size OR length OR weight OR underweight OR stunting)))) (n=289,653)
Scopus	#1	TITLE-ABS-KEY ( ( ( apical OR periapical OR periradicular OR radicular OR pulp* ) W/1 ( periodontitis OR disease OR abscess OR granuloma
(n=66)		OR infection OR lesion OR patho* OR inflammat* OR condition OR process ) ) OR pulpitis OR {root canal} OR endodont* ) (n=71,716)
	#2	TITLE-ABS-KEY ( ( ( pregnan* OR birth ) W/1 ( outcome OR complication ) ) OR ( ( premature OR preterm ) W/1 birth ) OR pre-eclampsia OR
		( hypertension W/1 ( gestational OR pregnancy ) ) OR preeclampsia OR ( ( fetal OR intrauterine ) W/1 growth W/1 ( retardation OR restriction ) )
		OR "low birth weight" OR birthweight OR chorioamnionitis OR stillbirth OR "neonatal sepsis" OR ( duration W/1 pregnancy ) OR ( ( neonatal
		OR birth ) W/1 ( size OR length OR weight OR underweight OR stunting ) ) ) (n=425,200)
PubMed	#1	"periapical periodontitis"[MeSH Terms] OR ("apical"[Title/Abstract] OR "periapical"[Title/Abstract] OR "periradicular"[Title/Abstract] OR
(n=373)		"radicular"[Title/Abstract] OR "pulp*"[Title/Abstract]) AND ("periodontitis"[Title/Abstract] OR "disease*"[Title/Abstract] OR "abscess*"[Title/Abstract]
		OR "granuloma*"[Title/Abstract] OR "infection*"[Title/Abstract] OR "lesion*"[Title/Abstract] OR "patho*"[Title/Abstract] OR
		"inflammat*"[Title/Abstract] OR "condition*"[Title/Abstract] OR "process*"[Title/Abstract]) OR "pulpitis"[Title/Abstract] OR "root canal"[Title/Abstract]
		OR “endodont*”[Title/Abstract] (n=83,570)

	#2	"Pregnancy Outcome"[Mesh] OR "Pregnancy Complications"[Mesh] OR "Pre-Eclampsia"[Mesh] OR "Premature Birth"[Mesh] OR "Fetal Growth Retardation"[Mesh] OR "Infant, Low Birth Weight"[Mesh] OR "Chorioamnionitis"[Mesh] OR "Stillbirth"[Mesh] OR "Neonatal Sepsis"[Mesh] OR ("pregnan*" [Title/Abstract] OR "birth"[All Fields]) AND ("outcome*" [Title/Abstract] OR "complication*" [All Fields])) OR (("Premature" [Title/Abstract] OR "preterm" [All Fields]) AND "Birth" [Title/Abstract]) OR "Pre-Eclampsia" [Title/Abstract] OR "Preeclampsia" [Title/Abstract] OR (hypertension [Title/Abstract] AND (gestational [Title/Abstract] OR pregnancy [Title/Abstract])) OR (("Fetal" [Title/Abstract] OR "intrauterin" [All Fields] OR "intrauterine" [All Fields]) AND "Growth" [Title/Abstract] AND ("Retardation" [Title/Abstract] OR "restriction" [All Fields])) OR "low birth weight" [Title/Abstract] OR "Chorioamnionitis" [Title/Abstract] OR "Stillbirth" [Title/Abstract] OR "Neonatal Sepsis" [Title/Abstract] OR (“duration” [Title/Abstract] AND “pregnancy” [Title/Abstract]) OR ((“neonatal” [Title/Abstract] OR “birth” [Title/Abstract]) AND (“size” [Title/Abstract] OR “length” [Title/Abstract] OR “weight” [Title/Abstract] OR “underweight” [Title/Abstract] OR “stunting” [Title/Abstract])) (n=659,687)
CENTRAL (n=51)	#1	[mh "periapical periodontitis"] OR ((apical OR periapical OR periradicular OR radicular OR pulp*) NEAR/1 (periodontitis OR disease? OR abscess? OR granuloma? OR infection? OR lesion? OR patho* OR inflammat* OR condition? OR process*)):ti,ab,kw (n=1,192)
	#2	([mh "Pregnancy Outcome"] OR [mh "Pregnancy Complications"] OR [mh "Pre-Eclampsia"] OR [mh "Premature Birth"] OR [mh "Fetal Growth Retardation"] OR [mh "Infant, Low Birth Weight"] OR [mh "Chorioamnionitis"] OR [mh "Stillbirth"] OR [mh "Neonatal Sepsis"]) OR (((Pregnan* OR birth) AND (Outcome? OR Complication?)) OR ((Premature OR Preterm) AND Birth) OR Pre-Eclampsia OR Preeclampsia OR (hypertension? NEAR/1 (gestational OR pregnancy)) OR ((Fetal OR intrauterine) AND Growth AND (Retardation OR Restriction)) OR "low birth weight" OR Chorioamnionitis OR Stillbirth OR "Neonatal Sepsis" OR (duration NEAR/1 pregnancy) OR ((neonatal OR birth) NEAR/1 (size OR length OR weight OR underweight OR stunting))):ti,ab,kw (n=61,848)

n - number of hits, WoS - Web of Science Core Collection, KJD - Korean Journal Database, RSCI - Russian Science Citation Index, SCIELO - SciELO Citation Index, CENTRAL - Cochrane Central Register of Controlled Trials, TS - Topic (article title, abstract and keywords).

**Supplemental Table 2. Critical appraisal of included cross-sectional study via adapted Newcastle-Ottawa Scale tool**

Study (year)	Selection	Selection	Selection	Comparability	Comparability	Outcome	Outcome	Statistical	Stars	Results *
	Representative	Non-respondents	Exposure	Appropriate control	Other Factors	Independent	Record Linkage	Test		
Harjunmaa <i>et al.</i> 2015	*	*	AP not specified	*	Smokers and	Not blinded	*	*	5	Fair
					Periodontitis					

\* The appraisal was based on the 2 stars assessed within the selection domain, the 1 or 2 stars awarded in comparability domain and 2 or 3 stars in outcome/exposure domain (McPheeters *et al.* 2012 ); AP, apical periodontitis;

Supplemental Table 3. Critical appraisal of included case control studies via Newcastle-Ottawa Scale tool

Study (year)	Selection	Selection	Selection	Selection	Comparability	Comparability	Exposure Secure	Exposure Blinded	Exposure Same	Exposure Same	Results *
	Definition	Representativeness	Control	Definition of	Appropriate	Other Factors	Record		Methodology	Rate	
	Adequate		(community control)	Control	control						
Leal <i>et al.</i> 2015	*	*	*	*	*	Smoking, alcohol, infection during pregnancy, diabetes mellitus, and periodontal disease were not confounders	*	*	*	*	Good
Khalighinejad <i>et al.</i> 2017	*	Sample size not justified	*	*	*	Smokers and periodontitis	Not secure record	Not blinded or masked	*	*	Fair

\* The appraisal was based on the 2 stars assessed within the selection domain, the 1 or 2 stars awarded in comparability domain and 2 or 3 stars in outcome/exposure domain (McPheeters *et al.* 2012);