

# New insights into large-scale trends of apparent organic matter reactivity in marine sediments and patterns of benthic carbon transformation

5 Felipe S. Freitas<sup>1,3,a</sup>, Philip A. Pika<sup>2,4,b††</sup>, Sabine Kasten<sup>5,6,7</sup>, Bo B. Jørgensen<sup>8</sup>, Jens Rassmann<sup>9</sup>,  
Christophe Rabouille<sup>9</sup>, Shaun Thomas<sup>10,c †††</sup>, Henrik Sass<sup>10</sup>, Richard D. Pancost<sup>1,3</sup>, Sandra Arndt<sup>4</sup>

<sup>1</sup>Organic Geochemistry Unit, School of Earth Sciences & School of Chemistry, University of Bristol, Bristol, BS8 1TS, United Kingdom

<sup>2</sup>BRIDGE, School of Geographical Sciences, University of Bristol, Bristol, BS8 1RL, United Kingdom

<sup>3</sup>Cabot Institute for the Environment, University of Bristol, Bristol, BS8 1UH, United Kingdom

10 <sup>4</sup>Biogeochemistry and Earth System Modeling, Geosciences, Environment and Society Department, Université Libre de Bruxelles, Brussels, CP160/03 1050, Belgium

<sup>5</sup>Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, 27570, Germany

<sup>6</sup>Faculty of Geosciences, University of Bremen, 28359 Bremen, Germany

<sup>7</sup>MARUM – Center for Marine Environmental Sciences, University of Bremen, 28359 Bremen, Germany

15 <sup>8</sup>Section for Microbiology, Department of Biology, Aarhus University, 8000 Aarhus C, Denmark

<sup>9</sup>Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ-Université Paris Saclay, 91198 Gif-sur-Yvette, France

<sup>10</sup>School of Earth and Ocean Sciences, Cardiff University, Cardiff, CF10 3AT, United Kingdom

20 <sup>a</sup>Current address: School of Earth Sciences, University of Bristol, Bristol, BS8 1RJ, United Kingdom

<sup>b</sup>Current address: Department of Earth Sciences, VU University of Amsterdam, 1081 HV Amsterdam, The Netherlands

<sup>c</sup>Current address: RSR Ltd, Parc Ty Glas, Llanishen, Cardiff, CF14 5DU, United Kingdom

*Correspondence to:* Felipe S. Freitas ([felipe.salesdefreitas@bristol.ac.uk](mailto:felipe.salesdefreitas@bristol.ac.uk))

## Abstract

25 Constraining the mechanisms controlling organic matter (OM) reactivity and, thus, degradation, preservation and burial in marine sediments across spatial and temporal scales is key to understanding carbon cycling in the past, present, and future. However, we still lack a detailed quantitative understanding of what controls OM reactivity in marine sediments and, consequently, a general framework that would allow model parametrization in data poor areas. To fill this gap, we quantify apparent OM reactivity (i.e., OM degradation rate constants) by extracting reactive continuum model (RCM) parameters ( $a$   
30 and  $v$ , which define the shape and scale of OM reactivity profiles, respectively) from observed benthic organic carbon and sulfate dynamics across 14 contrasting depositional settings distributed over five distinct benthic provinces. We further complement the newly derived parameter set with a compilation of 37 previously published RCM  $a$  and  $v$  estimates to explore large-scale trends in OM reactivity. Our analysis shows that the large-scale variability in apparent OM reactivity is largely driven by differences in parameter  $a$  ( $10^{-3}$ – $10^7$ ) with a high frequency of values in the range  $10^0$ – $10^4$  years. In contrast, and in  
35 broad agreement with previous findings, inversely determined  $v$ -values fall within a narrow range (0.1–0.2). Results also show

that the variability in parameter  $a$  and, thus, in apparent OM reactivity is a function of the whole depositional environment, rather than traditionally proposed, single environmental controls (e.g., water depth, sedimentation rate, OM fluxes). Thus, we caution against the simplifying use of a single environmental control for predicting apparent OM reactivity beyond a specific local environmental context (i.e., well defined geographic scale). Additionally, model results indicate that, while OM fluxes exert a dominant control on depth-integrated OM degradation rates across most depositional environments, apparent OM reactivity becomes a dominant control in depositional environments that receive exceptionally reactive OM. Furthermore, model results show that apparent OM reactivity exerts a key control on the relative significance of OM degradation pathways, the redox zonation of the sediment, and rates of anaerobic oxidation of methane. In summary, our large-scale assessment i) further supports the notion of apparent OM reactivity as a dynamic ecosystem property; ii) consolidates the distributions of RCM parameters; and iii) provides quantitative constraints on how OM reactivity governs benthic biogeochemical cycling and exchange. Therefore, it provides important global constraints on the most plausible range of RCM parameters  $a$  and  $v$  and largely alleviates the difficulty of determining OM reactivity in RCM by constraining it to only one variable, i.e., the parameter  $a$ . It thus represents an important advance for model parameterization in data-poor areas.

**Key words:** Reaction-Transport Model; seafloor; diagenesis; degradation rates, anaerobic oxidation of methane; sediment-water interface fluxes.

## 1 Introduction

Organic matter (OM) buried in marine sediments represents the largest reactive reservoir of reduced carbon on Earth (e.g., Hedges, 1992). The majority of OM that reaches modern global ocean seafloors originates from contemporary primary productivity (PP) in terrestrial (global net primary productivity, NPP, =  $\sim 59 \text{ Pg C y}^{-1}$ ) or marine environments (global NPP =  $\sim 49 \text{ Pg C y}^{-1}$ ) (Hedges et al., 1997; Hedges and Oades, 1997; Kharbush et al., 2020). Additionally, it comprises OM derived from chemoautotrophy (Lengger et al., 2019; Vasquez-Cardenas et al., 2020) and ancient PP, the latter delivered through rock weathering, soil erosion, and thawing permafrost (Hedges et al., 1997; Kusch et al., 2021; Grotheer et al., 2020). Since OM production slightly exceeds degradation, a small fraction of the photosynthetically produced organic carbon escapes heterotrophic degradation and is buried in sediments (Berner, 2003; Burdige, 2006). This relatively small imbalance connects the short- and long-term carbon cycles, controls the long-term evolution of atmospheric  $\text{CO}_2$  and has enabled the accumulation of oxygen in the atmosphere (Berner, 2003; Hedges, 1992). Nevertheless, several data compilations reveal that rates of OM degradation and preservation vary significantly in space (Arndt et al., 2013; Burdige, 2006; Emerson and Hedges, 1988; Middelburg, 2019) and time (Berner, 2003). Significant progress has been made over the past decades in identifying the potential environmental controls of this spatio-temporal variability (Arndt et al., 2013; Bogus et al., 2012; LaRowe et al., 2020b; Zonneveld et al., 2010). It is becoming increasingly clear that the susceptibility of OM to heterotrophic degradation is not solely an inherent characteristic of the OM itself, but also depends on its environmental context (Kayler et al., 2019;

LaRowe et al., 2020b; Mayer, 1995; Schmidt et al., 2011; Zonneveld et al., 2010). In fact, OM reactivity is determined by the dynamic interplay of a plethora of different factors (Arndt et al., 2013; Burdige, 2006; Middelburg, 2019; LaRowe et al., 2020b) (Table 1 and references therein). However, the relative significance of each of these factors remains poorly quantified. Consequently, we currently lack a general framework that allows quantifying the reactivity of OM in a given environmental context. This knowledge gap seriously compromises our ability to constrain OM reactivity in both space and time (Arndt et al., 2013; Lessin et al., 2018). The current lack of such a general framework ultimately limits our ability to quantify pathways and rates of biogeochemical processes over different temporal and spatial scales and, thus, predict the impact and feedbacks of past, present, and future climate change on global biogeochemical cycles and climate (Hülse et al., 2018).

Traditionally, apparent (i.e., estimated representation of) OM reactivity has been determined through laboratory-based incubation experiments in shallow (few centimetres deep) and mixed sediments (e.g., Dai et al., 2009; Grossi et al., 2003; Westrich and Berner, 1984) or integrated model-data approaches in deeply buried sediments (hundreds of centimetres to meters deep and hundreds to thousands and million years old) (e.g. Boudreau and Ruddick, 1991; Middelburg, 1989; Wehrmann et al., 2013), using Reaction-Transport Models (RTM). The latter are powerful tools since they allow the extraction of quantitative measures of OM reactivity from comprehensive multi-species porewater datasets integrated over greater sediment depths/ages (e.g., Middelburg, 2019). However, due to our limited quantitative understanding of the environmental controls on OM reactivity, the application of RTM approaches to data-poor settings is still complicated by the difficulty in constraining model parameters.

The need for predictive frameworks that constrain OM degradation model parameters has been recognized since the early days of RTM. Based on the rationale that the reactivity of OM settling onto the sediment is controlled by the degree of degradation, and thus its residence time in the oxygenated water column, early efforts have explored correlations between OM reactivity and water depth, sedimentation rate or OM fluxes (Boudreau, 1997; Boudreau and Ruddick, 1991; Müller and Suess, 1979; Tromp et al., 1995). However, the derived relationships were based on limited datasets and a more recent compilation of global data did not reveal statistically significant correlations between OM reactivity and single depositional environmental descriptors (Arndt et al., 2013). However, the latter combined OM degradation model parameters derived from structurally different models and used observational data of different complexity, thus somewhat compromising the comparability of their findings. Nevertheless, it emphasised the role of a complex interplay of environmental controls on reactivity (Kayler et al., 2019; LaRowe et al., 2020b; Mayer, 1995; Schmidt et al., 2011; Zonneveld et al., 2010).

The efforts by Arndt et al. (2013) to develop a general predictive framework highlighted two critical issues: OM reactivity is variable within and across settings with no significant global trends; and there is a crucial need to resolve the lack of comparative studies. Therefore, we here use a common, integrated model-data approach to determine inversely the model parameters that control apparent OM reactivity from contemporary sediment depth profiles (Sect. 3) at 14 different sites from

five different depositional environments (Sect. 2). OM degradation is formulated according to the Reactive Continuum Model (RCM) (Boudreau and Ruddick, 1991), which represents the most suitable degradation model type to explore links between reactivity parameters and environmental controls (see Sect. 3.2 for details). Derived RCM parameters are discussed in the environmental context of depositional settings. We further complement the newly derived parameter set with a compilation of previously published RCM parameter values to explore the distribution of model parameters and provide first order constraints on model parametrization in data poor area. Finally, we evaluate the control of OM reactivity on benthic cycling and fluxes by linking RCM parameters to estimated OM degradation rates and relative contributions of metabolic pathways, anaerobic oxidation of methane (AOM) coupled to sulfate reduction, as well as fluxes of dissolved species across the sediment-water interface (SWI) at each of the 14 studied sites.

## 2 Study sites

Our data-model study covers a wide range of depositional environments, from coastal to shelf and slope settings characterized by widely different environmental conditions (Fig. 1, Table 2). It comprises the following benthic provinces (Seiter et al., 2004): Northern European margin (EUR1); NW Mediterranean margin off Rhone delta (EUR2); Western Arabian Sea (WARAB); Bering Sea (NWPAC); and SW Atlantic off La Plata river (RIOPLATA). Most of those regions are characterised by predominantly temperate climate regimes, except for the Arabian Sea region (Arid) and the Bering Sea area (Snow/Polar) (Chen and Chen, 2013). OM characteristics also cover a broad spectrum, ranging from organic rich ( $TOC > 2$  wt%) to lean ( $TOC < 1$  wt%) sediments (Premuzic et al., 1982; Seiter et al., 2004), and from predominantly terrestrial (coast), mixed terrestrial and marine (coast/shelf) to predominantly marine (shelf/slope) OM (Hedges et al., 1997; Meyers, 1997). The pelagic ecosystem structure reflects local and regional characteristics. Overall, as a common characteristic they display the occurrence of spring/summer phytoplankton blooms (e.g., Cowie, 2005; Rixen et al., 2019; Fleming-Lehtinen and Laamanen, 2012; Jensen et al., 1990; Coyle et al., 2008). This is also reflected by the composition of the phytoplankton community, and thus OM production at the photic zone and vertical transport efficiency to sediments (e.g., Bach et al., 2019). This wide spectrum of depositional environments (Table 2 and S1) offers a unique opportunity to quantify OM reactivity and explore its multidisciplinary links with environmental drivers.

## 3 Methods

Here, we adopt an integrated model-data approach to determine inversely OM reactivity parameters from contemporary total organic carbon ( $TOC$ ) and sulfate ( $SO_4^{2-}$ ) profiles measured at 14 sites across the five different depositional environments described in Sect. 2. The following sections provide an overview of the observational data (Sect. 3.1), a brief description of the applied RTM (Sect. 3.2.1; for further details see Supplements, Sect. S2), a detailed description of the inverse model

approach used to extract RCM parameters from the observed depth profiles (Sect. 3.2.4), as well as a summary of the analysed model output (section 3.2.5).

### 3.1 Observational Data

135 We benefit from previously published datasets to develop our large-scale OM reactivity assessments. *TOC* and porewater  
*SO<sub>4</sub><sup>2-</sup>* depth-profiles, as well as site-specific physical characteristics and supporting data are available for all sites and were  
compiled from the PANGAEA database ([www.pangaea.de](http://www.pangaea.de)) (except from Rhone delta and Severn estuary datasets), as well as  
corresponding published literature (see below). Specific details about sampling strategies and analytical methods can be found  
in the respective publications. The dataset for the Severn estuary (Table 2; row A) are from Thomas (2014). Rhone delta data  
140 (Table 2; rows B and C) are from the DICASE project (Rassmann et al., 2016). Data for the Northern European margin sites  
Aarhus Bay, Arkona Basin, and the Skagerrak transect (Table 2; rows D, E, G, H, and I) are part of the European funded  
project METROL (Aquilina et al., 2010; Dale et al., 2008a, b; Knab et al., 2008; Mogollón et al., 2012). Data from the  
Helgoland Mud Area, North Sea (Table 2; row F) (Henkel and Kulkarni, 2015), the Arabian Sea transect (Table 2; rows J, K,  
and L) (Bohrmann and cruise participants, 2010a, b, c), Bering Sea (Table 2; row M) (Gersonde, 2009), and Argentine Basin  
145 (Table 2; row N) (Henkel et al., 2011) are presented on PANGAEA database (see above). We use those site-specific datasets  
to inform our data-model analysis and to constrain OM reactivity and related benthic processes.

### 3.2 Model Description

#### 3.2.1 General RTM Framework

150 The Biogeochemical Reaction Network Simulator (BRNS) (Aguilera et al., 2005; Regnier et al., 2002) is an adaptive  
simulation environment that has been successfully employed to reproduce and quantify diagenetic processes in marine  
sediments across a wide range of depositional environments and timescales (Dale et al., 2008a; Thullner et al., 2009;  
Wehrmann et al., 2013). The BRNS is suitable for large, mixed kinetic-equilibrium reaction networks (Dale et al., 2009;  
Jourabchi, 2005; Thullner et al., 2009). The concentration depth profiles of solid and dissolved species in marine sediments  
155 are calculated according to the vertically-resolved mass conservation equation of solid and dissolved species in porous media  
(Berner, 1980; Boudreau, 1997):

$$\frac{\partial \sigma C_i}{\partial t} = \frac{\partial}{\partial z} \left( D_{bio} \sigma \frac{\partial C_i}{\partial z} + D_i \sigma \frac{\partial C_i}{\partial z} \right) - \frac{\partial \sigma \omega C_i}{\partial z} + \alpha_i \sigma (C_i(0) - C_i) + \sum_n s_i^n r_n \quad (1)$$

160 The first three terms on the right-hand side represent the transport process (bioturbation and molecular diffusion, advection,  
and bioirrigation), whereas the last term denotes the sum of all reactions (production and consumption) affecting species  $i$ .  
Table 2 summarizes all symbols employed here.

Briefly, the implemented reaction network encompasses the most pertinent primary and secondary redox reactions found in  
165 the upper layers of marine sediments (e.g., Aguilera et al., 2005; Thullner et al., 2009; Van Cappellen and Wang, 1996;). It  
explicitly accounts for the heterotrophic degradation of OM coupled to the consumption of oxygen (aerobic OM degradation),  
nitrate (denitrification), sulfate (organoclastic sulfate reduction), as well as methanogenesis. Additionally, the reaction network  
accounts for secondary redox reactions, i.e., re-oxidation of reduced species produced during primary redox reactions. It  
explicitly resolves nitrification, sulfide re-oxidation by  $O_2$ , anaerobic oxidation of methane (AOM) coupled to sulfate reduction  
170 and  $CH_4$  reoxidation by  $O_2$ .

The model transport and reactions equations were solved sequentially according to Regnier et al. (1998) The size of model  
domain was fixed at 1,000 cm for all sites, except for the Bering Sea, in which the model domain was extended to 1,500 cm,  
due to the low sedimentation rate assumed for this site (Table 2). This choice is based on initial tests and ensures that the model  
175 domain covers the diagenetically most active zone, thus reducing the influence of biogeochemical dynamics in underlying  
sediments on biogeochemical dynamics within the model domain. The model equations were solved on an uneven grid and  
with a time-adapting timestep ( $\Delta t < 0.01$ ) run until steady state was reached. The supplementary information includes a detailed  
description of the general RTM framework used here, its parametrization and numerical solution (Sect. S2).

### 180 3.2.2 OM degradation model

OM is composed of a complex and dynamic mixture of compounds that are distributed over a wide, continuous spectrum of  
reactivities. Thus, OM degradation is described by the reactive continuum model (RCM) (Boudreau and Ruddick, 1991), which  
assumes a continuous distribution of OM compounds over the entire reactivity spectrum. The RCM assumes a gamma  
distribution to describe the probability density function of OM distribution,  $om(k, t)$  (Aris, 1968; Boudreau and Ruddick,  
185 1991; Ho and Aris, 1987). The overall rate of OM degradation  $R_{TOC}$  is given by the integral:

$$r_n = R_{TOC} = - \int_0^{\infty} k \cdot om(k, t) dk \quad (2)$$

Here,  $om(k, t)$  determines the concentration of OM having a degradability between  $k$  and  $k + dk$  at time  $t$ , with  $k$  being  
190 equivalent to the first-order degradation rate constant. The initial distribution ( $t = 0$ ) of OM is given by:

$$om(k, 0) = \frac{TOC(0) \cdot a^v \cdot k^{v-1} \cdot e^{-a \cdot k}}{\Gamma(v)} \quad (3)$$

where  $TOC(0)$  denotes the initial OM content,  $\Gamma(v)$  is the gamma distribution,  $a$  is the average lifetime of the more reactive components of bulk OM, and  $v$  represents the dimensionless scaling parameter of the distribution near  $k = 0$ . The free, positive parameters  $a$  and  $v$  delineate the shape of the initial distribution of OM compounds along the range of  $k$ , and thus the overall reactivity of bulk OM. As such, the RCM approach requires the definition of two parameters that will define the shape of the OM distribution over reactivity  $k$  (Sect. 3.2.5, Eq. 5).

Due to the rapid depletion of the most reactive compounds, the reactivity of the bulk material decreases during degradation, reflecting the widely observed reactivity decrease with burial time/depth/age (Boudreau and Ruddick, 1991; Middelburg, 1989). This indicates that degradation of OM proceeds at different rates in parallel. Interactions between different compounds or transformations of compounds can change the reactivity of a given compound. While the RCM does not explicitly account for such interactions/transformations, the overall OM profiles take these interactions implicitly into account. Thus, OM compounds are continuously and dynamically distributed over a range of reactivities that captures the decrease in apparent reactivity with burial age/depth as the most reactive compounds are successively degraded. The interplay of  $a$  and  $v$  drives the bulk OM reactivity depth profiles, and consequently the yielded profiles from the reaction network implemented in the BRNS. The evolution of OM concentration as a function of depth  $z$  is given by:

$$TOC(z) = TOC(0) \cdot \left( \frac{a}{a + age(z)} \right)^v \quad (4)$$

where  $a$  and  $v$  are the free RCM parameters (Arndt et al., 2013; Boudreau and Ruddick, 1991) and  $age(z)$  denotes the age of the sediment layer at depth  $z$ . For non-bioturbated sediments ( $z > z_{bio}$ ) the burial  $age(z)$  can be calculated as a function of the burial velocity (Sect. S2.3, Eq. S8). However, within the bioturbated upper sediment layers, the age distribution of reactive species is controlled by both sedimentation, bioturbation, and the reactivity  $k$  of reactive species (Meile and Van Cappellen, 2005). In such cases, we apply a multi-G approximation for the RCM in the bioturbated sediments (Sect. S2.3, Eq. S9–S18).

### 3.2.3 Boundary conditions

Boundary conditions place the BRNS in the environmental context of each of the study sites (Table 4). The boundary conditions are constrained based on either site-specific measurements or alternatively on published data if direct observations were not available. Here, we assume a fixed boundary concentration of OM at the SWI. Concentrations of  $O_2$  are based on measurements in Skagerrak (Canfield et al., 1993) and Rhone delta (Rassmann et al., 2016) sediments. Similarly,

concentrations of  $NO_3^-$  are based on measurements in Skagerrak sediments (Canfield et al., 1993). Because of the lack of site-specific information, these  $O_2$  and  $NO_3^-$  are adopted at all oxic sites (where  $SO_4^{2-} > 28$  mM) since they are representative for coastal and shallow slope sites and their exact value exerts no influence on deep  $SO_4^{2-}$  and  $CH_4$  concentrations. For anoxic sites (where  $SO_4^{2-} < 28$  mM),  $O_2$  and  $NO_3^-$  bottom water concentrations are set to zero.  $SO_4^{2-}$  concentrations were constrained based on measurements for each simulated site. Concentrations of  $CH_4$  and  $HS^-$  were set to zero at SWI, as those species are rapidly oxidised in the overlying bottom water. A no flux ( $\frac{\partial C}{\partial z} = 0$ ) condition was implemented for all species at the lower boundary, assuming that biogeochemical dynamics in underlying sediments exert no influence on diagenetic processes in the model domain (e.g., Thullner et al., 2009).

### 3.2.4 Inverse modeling

We use an inverse model approach to extract the optimal OM reactivity parameter set  $a$  and  $v$  (i.e., the apparent OM reactivity) by assuming that the rank of a parameter set depends on the similarity between simulated and measured data. In general, all inverse modeling approaches are ultimately affected by a few limitations. Among others, the quality of the inverse modeling results is directly affected by the quality of the observational data. In particular, the lack of observations/data at the SWI due to core top loss during sediment sampling by gravity and piston corers can impact results. Additionally, processes that are not explicitly described in the applied model can impact inverse model results because reaction rate parameters implicitly account for those processes. Furthermore, downcore changes in concentration-depth profiles may also be influenced by transient features in bottom water conditions and/or fluxes, and the validity of the steady state assumption thus needs to be carefully evaluated. Finally, multiple parameter sets might fit the observations equally well. Because both  $a$  and  $v$  exert an influence on the apparent OM reactivity and its evolution with depth (see below, Eq. 5), they are not completely independent parameters. For instance, a decrease in  $v$  (decreasing reactivity) can be compensated by a simultaneous decrease in parameter  $a$  (increasing reactivity). Consequently, different  $a$  and  $v$  parameter couples, might result in equally statistically satisfying fits between the simulated and observed profile.

These limitations can be alleviated by using comprehensive, multi-component observational data sets, because the inclusion of additional information adds further constraints. However, such data sets are often not available. For instance, the first efforts to quantify  $a$  and  $v$  parameter values, and thus reactivity, on the basis of observations solely focused on downcore *TOC* profiles (Boudreau and Ruddick, 1991). Such approach was instrumental for the development of the RCM and for producing the first large-scale dataset of  $a$  and  $v$ . However, the increasing availability and accessibility of multi-species data sets allow for the use of additional information that is available across different sites. Because OM heterotrophic degradation is the driver behind biogeochemical dynamics in marine sediments, downcore profiles of TEAs (e.g.,  $SO_4^{2-}$ ) (e.g., Bowles et al., 2014; Jørgensen et al., 2019b) or reduced products of OM degradation (such as  $CH_4$  and  $NH_4^+$ ) could be used as further constraints.

255 However, data availability is often still limited; therefore, it is important to identify a minimum set of observational data that  
are widely available and comparably easy to measure. We tested the suitability of different artificial porewater data sets for  
the inverse determination of apparent OM reactivity. Using an artificially generated dataset (see Supplements, Sect. S3), we  
ran the BRNS in the same fashion as we did for the site-specific data-model approach (see below). Our sensitivity analysis  
confirms that when  $TOC$  is considered as a single constraint, multiple pairs of  $a$  and  $v$  produce equivalent  $TOC$  depth-profiles  
260 (Supplements, Fig. S1). Model results show that including  $SO_4^{2-}$  depth profiles as an additional constraint (Supplements, Fig.  
S2), facilitates the identification of a best fit  $a$  and  $v$  parameter set (Supplements, Fig. S3).  $CH_4$  profiles were used as an  
additional qualitative constraint when the data were available, although measured *ex situ*  $CH_4$  profiles are often not reliable  
due to sampling and measurement uncertainties in deeper sediment layers where concentrations exceed millimolar levels (Dale  
et al., 2008a; Hensen et al., 2003; Hilligsøe et al., 2018). The dynamics of  $SO_4^{2-}$  and  $CH_4$  are solely controlled by OM  
265 degradation and AOM (e.g., Bowles et al., 2014; Egger et al., 2018; Regnier et al., 2011), although their distributions can be  
also affected by bioirrigation in particularly shallow sulfate-methane transition zones (SMTZs) (e.g., Dale et al., 2019). In  
anoxic settings and deeply buried sediments,  $SO_4^{2-}$  is the dominant TEA and  $CH_4$  is the most common reduced species  
(Jørgensen et al., 2019a, b). Thus, a combination of  $TOC$ ,  $SO_4^{2-}$ , and  $CH_4$  (if available to verify the depths of the SMTZ) depth  
profiles incorporates the information contained in the observed benthic sulfur and carbon dynamics and is sufficient to extract  
270 robust estimates of apparent OM reactivity and its evolution from the sediment-water interface down to the SMTZ. In addition,  
because changes in  $a-v$  exert different effects on  $TOC$  and  $SO_4^{2-}$  depth profiles (see Fig. S1 & S2), including these two  
species reduces the impact of the  $a-v$  correlation on the uniqueness of fit. Thus, here we perform a site-specific data-model  
fit based on  $TOC$  and  $SO_4^{2-}$  (and  $CH_4$ ) depth profiles.

275 The optimal parameter set was then determined for each site by assuming that the rank of a parameter set depends on the  
similarity between simulated and measured data. Best fit parameter couples were found by minimizing the misfit between  
model results and simulations. We quantified the similarity between the simulated and observed  $TOC$  and  $SO_4^{2-}$  depth profiles  
in terms of correlation coefficient ( $r$ ), centred root mean square difference (RMSD), and standard deviations (SD). All these  
measures can be graphically represented in a single Taylor diagram (Taylor, 2001) that statistically quantifies the misfit  
280 between the observed and simulated data and visualizes how closely the simulated depth profile resembles the observed one.

The best fit RCM parameters were inversely determined by first running the model for each site with a set of  $a-v$  couples  
over the entire range of previously published  $a$  ( $a = 10^{-3}$ – $10^7$  yr) and  $v$  ( $v = 10^{-2}$ – $10^0$ ) values (Arndt et al., 2013 and references  
therein). The approach adopted for the sampling of the parameter space is a critical step in inverse modeling since the chosen  
285 statistical measures may reveal, in addition to a global optimum, multiple local optima. Because of the comparably low  
computational cost of the forward modeling, the low number of unknown parameters (here,  $a$  and  $v$ ), as well as the weak  
nonlinearity of the problem, we used a simple two-step nested regular sampling of the two-dimensional parameter space. First,

the entire plausible  $a-v$  space (see above) was sampled on a coarse regular grid ( $\Delta \log(a) = 1$ ,  $\Delta v = 0.02$ ) and, for each of these runs,  $r$ , RMSD, and SD were calculated for both  $TOC$  and  $SO_4^{2-}$  profile. Based on these measures, a new, more finely resolved  $a-v$  grid (10x10) was defined around the  $a-v$  couples that showed the smallest combined misfit between simulated and observed  $TOC$  and  $SO_4^{2-}$  profiles. On this grid,  $a-v$  couples were again sampled regularly and the optimized  $a-v$  couple was determined by the lowest combined misfit (Table 5).

### 3.2.5 Quantification of organic matter reactivity and associated benthic processes

The apparent reactivity of bulk OM  $k$  for each site is calculated based on the RCM reactivity parameters  $a$  and  $v$  (Boudreau and Ruddick, 1991) that yield the best data-model fit and is given by:

$$k(z) = \frac{v}{(a + a_{ge}(z))} \quad (5)$$

The apparent reactivity of bulk OM at the sediment water interface ( $z = 0$ ) is thus given by:

$$k(0) = \frac{v}{a} \quad (6)$$

The BRNS simulates downcore concentration profiles for each species  $i$  that is explicitly resolved in the reaction network, as well as reaction rates  $r_n$  (Table S4). From those reaction rates, the total rate of organic carbon oxidation, i.e. the depth integral of OM degradation for each primary redox reaction  $n$  can be calculated by integrating the rate over the entire model domain  $L$  (e.g., Thullner et al., 2009):

$$R_{TOC,total} = \sum_{n=1-4} \int_0^L r_n dx = - \int_0^\infty k \cdot om(k, t) dk \quad (7)$$

310

Where  $n = 1, 2, 3$ , and 4 denotes aerobic OM degradation, denitrification, organoclastic sulfate reduction, and methanogenesis, respectively (Table S4). The OM fluxes ( $J_{in}$ ) at the SWI are calculated as the sum of OM out of the sediment ( $J_{OM,out}$ ) and the burial fluxes at the base of the model domain ( $J_{OM,bur}$ ) (e.g. Burdige, 2006). Here, we assume that  $J_{OM,out}$  is equivalent to the depth-integrated rates of OM degradation (Eq. 7) and  $J_{OM,bur}$  is solely governed by advective processes at depth  $z = 1,000$  cm.

315 As such:

$$J_{OM,out} = \sum_{n=1-4} \int_0^L r_n dx \quad (8)$$

$$J_{OM,bur} = (1 - \varphi_z) \cdot \omega_z \cdot TOC_z \quad (9)$$

320

$$J_{OM,in} = J_{OM,out} + J_{OM,bur} \quad (10)$$

Additionally, the fluxes of species  $i$  at the SWI, i.e., the benthic-pelagic fluxes, are calculated from the model-derived depth profiles of species  $i$  according to (e.g. Burdige, 2006):

325

$$J_i = J_{i,Advection} + J_{i,Diffusion} + J_{i,Bioturbation} + J_{i,Bioirrigation} \quad (11)$$

Positive  $J_i$  values indicate upward fluxes from the sediment, whereas negative values represent downward fluxes into the sediment.

## 330 **4 Results and discussion**

The integrated data-model analysis yielded a comprehensive picture of OM reactivity parameters ( $a$ ,  $v$ , and  $k(z)$ ), as well as OM degradation dynamics and benthic-pelagic coupling for our large-scale compilation of depositional environments (Table 6). Because model parameters implicitly account for all the processes that are not explicitly described in the model, this variability provides important insights into the environmental controls on OM reactivity.

335

### **4.1 Constraining apparent OM reactivity**

#### **4.1.1 Inverse Modelling**

We inversely determined the set of RCM parameters  $a$  and  $v$  that produces the best model fit to the observed  $TOC$  (Fig. 2) and  $SO_4^{2-}$  (Fig. 3) depth-profiles. Table 5 shows the quantitative assessments of these model fits and associated uncertainties.

340

Overall,  $TOC$  profiles display a satisfying fit between model and observations. However, the quality of fit is often compromised by a lack of observations/data at the SWI due to core loss during sediment sampling by gravity and piston corers, poor downcore sampling resolution, as well as possible non-steady state conditions. Generally, the latter may be problematic when trying to reproduce sedimentary conditions of dynamic shallow systems and long-term burial in shelf and deep-sea sites.

345

At estuarine sites, steady-state conditions may be impacted by the highly dynamic nature of such settings. The Severn estuary exhibits strong tidal dynamics that result in continuous cycles of sediment resuspension (Manning et al., 2010). The Rhone delta experiences strong pulses of fresh water and sediments associated to flood events, as well as variable sedimentation rates (Antonelli et al., 2008; Cathalot et al., 2010; Zebracki et al., 2015). In shelf and slope settings, steady-state assumptions can be impacted by fluctuations on sedimentation rates and lateral transport of sediments across shelves (e.g., Henkel et al., 2011;

Riedinger et al., 2014, 2017; Hensen et al., 2003; Cowie, 2005). Further, due to the possible loss of the upper few millimetres  
350 to centimetres during sampling with gravity and piston cores, it can be challenging to quantify OM contents at the SWI. Some  
of our sites (e.g., northern European sites; Aquilina et al., 2010; Knab et al., 2008) are likely to be impacted by such factors.  
Overall, such limitations in inverse model results are highlighted by the somewhat low correlation coefficients for *TOC* profiles  
in most cases ( $r_{TOC} < 0.75$ ; Table 3). We find that by adding additional constraints to the model-data fit relieves such limitation.  
The  $SO_4^{2-}$  depth-profiles (Fig. 3) display a good agreement between present-day observations and model simulations, with the  
355 majority of sites exhibiting high correlation coefficients ( $r_{SO_4} > 0.95$ ; Table 3). Nevertheless, Fig. 3 shows cases where the  
model-derived sulfate depth-profiles display some mismatches relative to observations (Arabian Sea below OMZ, Fig. 3l;  
Bering Sea, Fig. 3m). At the Bering Sea, the data-model discrepancy is likely associated to kinks in both *TOC* (Fig. 2m) and  
 $SO_4^{2-}$  (Fig. 3m) profiles. They likely represent changes in the depositional regime over the studied timescale (e.g., Hensen et  
al., 2003) that are not captured by the model. The mismatch in the Arabian Sea at the site below the OMZ (Fig. 3l) could  
360 originate from  $SO_4^{2-}$  recycling through  $H_2S$  oxidation by  $MnO_2$  and  $Fe(OH)_3$  (e.g., Rassmann et al., 2020; Schulz et al.,  
1994), however we are unable to further explore this hypothesis due to a lack of metal oxide data to constrain those  
biogeochemical reactions.

#### 4.1.2 Apparent OM reactivity parameters $\nu$ and $a$

365 The inversely determined OM reactivity parameters  $a$  and  $\nu$  (Sect. 3.2.4) derived from the best fit model solutions (Table 6)  
reveal important differences in apparent OM reactivity and its distribution on a global scale. Alongside  $a$  and  $\nu$  parameters  
compiled from the literature (Arndt et al., 2009; Boudreau and Ruddick, 1991; Contreras et al., 2013; Henkel et al., 2012;  
Mewes et al., 2016; Mogollón et al., 2012, 2016; Wehrmann et al., 2013; Ye et al., 2016), they provide useful constraints on  
the plausible range of  $a$  and  $\nu$ , as well as their control on the global variability in apparent OM reactivity.

370

Parameter  $\nu$  exerts a scaling influence on OM reactivity distribution. High  $\nu$ -values result in a high initial OM reactivity,  
whereas low  $\nu$ -values produce low initial reactivity (Arndt et al., 2013). For instance, an extremely low  $\nu$ -value of  $\leq 0.01$  for  
OM deposited during the late Cretaceous Oceanic Anoxic Event 2 at Demerara Rise (Arndt et al., 2009) has been linked to  
intense and rapid sulfurization of OM in the strongly euxinic subtropical Atlantic Ocean (Hülse et al., 2019). While previously  
375 published values for contemporary ocean sediments fall within the range 0.1 and 1.08 (Arndt et al., 2013; Boudreau and  
Ruddick, 1991), inversely determined  $\nu$  values predominantly fall within the range of 0.1–0.2 (apparent 6<sup>th</sup> – 11<sup>th</sup> order of  
reaction) – although lower and higher  $\nu$  values were also determined (Fig. 4a). Newly determined  $\nu$  parameters thus confirm  
the previously observed dominance of  $\nu$  values between 0.1 and 0.2 (Boudreau and Ruddick, 1991) and suggest that  $\nu$  is  
relatively constant across a wide range of different depositional environments encountered in the contemporary ocean. These

380 findings also agree with the parametrization of another, empirically derived continuum model description of OM degradation (Middelburg, 1989), which applies a globally constant parameter of 0.16:

$$k(z) = 0.16 \cdot (a + \text{age}(z))^{-0.95} \quad (12)$$

385 Although the parameters of the so-called power model cannot be directly compared to those of the RCM due to different exponents, the RCM is nevertheless equivalent to a general power law with an exponent of  $-1$ :

$$k(z) = v \cdot (a + \text{age}(z))^{-1} \quad (13)$$

390 Thus, the  $v$  parameter range of the RCM is in good agreement with the 0.16 value imposed in the power law model.

Nevertheless, our results also reveal exceptions. Particularly high  $v$ -values ( $v > 0.25$ ) were determined for the Arabian Sea and Bering Sea. Similarly, Boudreau and Ruddick (1991) also report exceptionally high values ( $v \approx 1$ ) for the Peru Margin and North Philippine Sea but invoked the possibility of non-steady state depositional conditions as a possible explanation. In  
395 our study,  $v > 0.25$  also coincides in part with the poorest data-model fits, such that the robustness of such high  $v$  values is uncertain (Table 5). Especially, the high  $v$ -value in the Bering Sea (Table 6) might result from uncertainties (see above, Sect. 4.1.1) associated with boundary conditions and transient features (e.g. Henkel et al., 2011; Hensen et al., 2003). However, the model-derived  $v > 0.25$  for the Arabian Sea agrees with previous diagenetic model results for that region. Luff et al. (2000) developed a 3G-model to investigate OM dynamics and found that bulk OM contains exceptionally reactive compounds with  
400 first order  $k$  of 0.2–30 yr<sup>-1</sup>. In the Arabian Sea, the OM flux to the sediment is supported by high primary production rates during the Monsoon season (e.g., Cowie, 2005), complemented by deep carbon fixation by microorganisms (Lengger et al., 2019). Observations indicate an efficient vertical transport of diatom and haptophyte detritus to the sediment (Barlow et al., 1999; Cowie, 2005; Latasa and Bidigare, 1998; Shalapyonok et al., 2001). High transport rates and pronounced oxygen minimum zones (OMZs) might limit heterotrophic degradation of the sinking OM in the water column, thus delivering  
405 comparably unaltered OM to the sediment. Biomarker analysis of benthic OM indicates the deposition of highly reactive OM (Koho et al., 2013) and a reactivity decrease with extended exposure to oxic conditions in the water column (Vandewiele et al., 2009). Therefore, the somewhat atypical high  $v$ -values determined (together with comparably low  $a$ -values; see below) for the Arabian Sea transect is supported by interdisciplinary observations. Thus,  $v$  has a relatively narrow range of 0.1 to 0.2 with higher  $> 0.2$  values generally restricted to depositional environments that are characterized by high primary production  
410 rates, anoxic water conditions, and enhanced vertical OM fluxes within the OMZ.

Parameter  $a$  determines both the overall apparent reactivity, as well as the shape of the reactivity decrease with depth. High  $a$ -values result in an overall low OM reactivity, but a slow decrease in reactivity with depth. In contrast, low  $a$ -values result in a higher reactivity, but also a more rapid decrease in reactivity with sediment depth (Arndt et al., 2013). Parameter  $a$  is often related to the degree of “pre-ageing” or degradation (e.g., Middelburg, 1989; Middelburg et al., 1993). Very fresh, marine OM has been shown to be characterized by substantially low  $a$  values ( $a = 3.1 \cdot 10^{-4}$  yr) (Boudreau and Ruddick, 1991), while the heavily sulfurized OM deposited during Cretaceous Oceanic Anoxic Event 2 in the strongly euxinic subtropical Atlantic Ocean is characterized by very high  $a$  values ( $a = 10^4$  yr) (Arndt et al., 2009). Our inversely derived  $a$ -values span several orders of magnitude (Table 6; Fig. 4b). They thus corroborate the high variability observed in previously published  $a$ -values compilations (e.g., Arndt et al., 2013; Boudreau and Ruddick, 1991) and suggest that the variability of parameter  $a$  exerts the major control on the spatial variability in OM reactivity. Additionally, inversely determined  $a$  values also broadly agree with previously observed global patterns. Low values ( $a < 10^2$  yr; i.e. high apparent OM reactivity) are generally determined for depositional environments that are characterized by the rapid deposition of predominantly fresh, marine OM (e.g. Arabian Sea and shallow sites at the Northern European margin), while depositional settings that receive pre-degraded OM and/or complex mixtures of OM (e.g. Skagerrak) exhibit high  $a$  ( $a > 10^3$  yr). However, despite the broad range of  $a$ -values, the ensemble of inversely determined  $a$  values shows that most values fall into the range  $10^0$ – $10^4$  years, thus narrowing the range of  $a$ -values.

#### 4.1.3 OM reactivity distributions

Inversely determined parameters  $a$  and  $v$  also allow the assessment of the apparent OM reactivity evolution during burial. Here, we calculate the probability distribution of OM over  $k$  at the time of deposition on the SWI, as well as after burial at 10, 100, and 1,000 cmbsf (centimetres below seafloor). Since  $v$ -values are broadly similar across depositional settings, the differences in  $k$  distributions are mainly driven by the range of  $a$ -values (Fig. 5). However, it is important to note that sedimentation rates  $\omega$ , as well as OM contents also influence the evolution of these distributions as they co-control degradation rates (Eq. 5, Eq. S8). Therefore, the changes in OM–reactivity distributions at the considered burial depths are not entirely driven by differences in reactivity, but more broadly reflect different exposure to heterotrophic degradation across sites.

The Arabian Sea region (Fig. 5 j – l) is characterized by the deposition of highly reactive OM, the concentration of which rapidly decreases with sediment depth (high  $v$  and low  $a$ ; Table 6). The OM reactivity distributions show a clear depletion of the most reactive OM compounds in the uppermost 10 cm. Below 100 cmbsf the remaining bulk OM is dominated by less reactive compounds ( $k(z) < 10^{-5}$  yr<sup>-1</sup>). The rapid decrease in apparent OM reactivity reflects the often rapid deposition of predominantly marine OM (Cowie, 2005; Seiter et al., 2005), whose degradation during settling is limited by short residence times in combination with locally anoxic conditions (Luff et al., 2000). The shallow, sub-oxic/anoxic sediments of the Northern European margin (Aarhus Bay – Fig. 5d; Arkona Basin – Fig. 5e) reveal a similar OM– $k$  distribution and depth profile.

445 However, the loss of the most reactive fractions within the uppermost 10 cm is less pronounced. Such pattern may be attributed  
to the burial of terrestrial, yet reactive OM (Aquilina et al., 2010). In contrast, Skagerrak sediments (S10 – Fig. 5g; S13 – Fig.  
5i) are characterised by the most homogeneous, albeit comparably unreactive OM mixture (high  $a$ -values; Table 4). The OM–  
 $k$  distributions reveal small changes from the SWI down to 1,000 cmbsf, reflecting a low overall reactivity, and thus low  
degradation rates. This somewhat surprising finding can be explained with the atypical character of the Skagerrak – a shallow  
environment that connects the brackish Baltic Sea with the saline North Sea and is characterized by an estuarine-like  
450 circulation. Due to this circulation pattern, the Skagerrak acts as the final sink for marine OM produced in the North Sea (Lohse  
et al., 1995; Van Weering et al., 1987). Consequently, more than 90% of the OM deposited in the Skagerrak is derived from  
re-suspended and pre-degraded marine OM or terrestrial OM (ca. 20% of the OM buried) (de Haas and van Weering, 1997) of  
generally low quality (Dauwe et al., 1999a, b).

455 Between those two endmember cases (high reactivity Arabian Sea and low reactivity Skagerrak), sediments from other sites  
generally reveal small changes in the continuous OM– $k$  distributions within the uppermost 10 cm. At all sites,  $k$ -values have  
converged to  $< 10^{-4} \text{ yr}^{-1}$ , below 100 cmbsf, where  $age(z) \gg a$  and changes in OM degradation rate constants with depth  
are thus largely controlled by  $age(z)$  (Eq. 5; Eq. S8). As such, only the least reactive fractions are preserved, leading to an  
overall low apparent OM reactivity in deeper sediments that is largely independent of the initial apparent OM reactivity  $k(0)$ .  
460 Sediments from the Bering Sea (Fig. 5m) are a notable exception. OM reactivity decreases gradually throughout the sediment  
column, reflecting both unusually high  $a$  and  $v$  values (Table 6). The Bering Sea is one of the most productive ocean regions  
and OM deposition is mostly fuelled by spring and summer blooms (Coyle et al., 2008; Stabeno and Hunt, 2002; Stockwell et  
al., 2001). However, due to great water depths and low sedimentation rates, OM can experience extensive ageing and pre-  
processing prior to burial and result in the OM distribution pattern observed here (Fig. 5m).

465

#### 4.1.4 Emerging environmental patterns in OM reactivity

Parameter  $a$  distributions exert the main control on the global variability in apparent OM reactivity. It also displays plausible  
values spanning several orders of magnitude (Fig. 4b, Table 6). Thus, our ability to quantify and predict benthic OM dynamics  
strongly depends on a general framework that allows constraining parameter  $a$  when observational data is scarce or unavailable  
470 (e.g., global, future and past model scenarios). Therefore, we further explore these links combining our newly generated dataset  
with those compiled from previously published RCM parameters (Fig. 6).

Despite the large scatter, broad trends emerge that generally agree with previously proposed relationships  $k$ – $\omega$  (Boudreau,  
1997; Tromp et al., 1995),  $k$ – $J_{OM,in}$  (Boudreau, 1997; Müller and Suess, 1979), and  $a$ – $\omega$  (Boudreau and Ruddick, 1991)  
475 (Fig. 6). Depositional environments characterised by low sedimentation rates, low OM fluxes, or great water depths generally

display high  $a$ -values (i.e., low apparent reactivity), whereas settings characterized by high sedimentation rates, high OM fluxes, or shallow water depth generally display lower  $a$ -values (high apparent reactivity). While these trends are extremely weak on a global scale, they are relatively more pronounced within a given geographical setting (i.e., on a local/regional scale). However, in agreement with the previous global assessment (Arndt et al., 2013), we do not find any statistically significant global or regional correlation between the reactivity parameters ( $a$  and  $k(0)$ ) and master characteristics of depositional environments. Earlier proposed empirical relationships were based on a limited number of observations available at that time and our results, as well as previously compiled parameter values, thus confirm cautious statements regarding the global validity of previously published relationships (Boudreau, 1997; Müller and Suess, 1979; Tromp et al., 1995). These results further emphasize that the spatial variability in  $a$  and thus apparent OM reactivity is not determined by one single characteristic of the depositional environment. Instead, it results from a complex and dynamic interplay of multiple environmental controls (e.g., Kayler et al., 2019; LaRowe et al., 2020b; Mayer, 1995; Zonneveld et al., 2010; Schmidt et al., 2011) and the ensemble of results indicates that sedimentation rates and OM deposition fluxes are poor proxies for the complex interplay of environmental controls on OM reactivity (Fig. 6 c,f).

However, Fig. 6b indicates that water depth seems to be a useful and easily accessible first-order proxy for this complex interplay of environmental controls on OM reactivity. Thus, it may serve as an acceptable predictor for first-order global reactivity patterns in the absence of more suitable information. At the global scale, inversely determined apparent OM reactivity for the deep, well-oxygenated sites (e.g., Bering Sea and Argentine Basin) fall in the lower end of the reactivity spectrum, whereas shallow coastal environments, such as the Northern European region, reveal generally higher apparent OM reactivities. In addition, within the Northern European region and along the Rhone delta transect, parameter  $a$  generally increases (i.e., decrease in apparent reactivity) with increasing water depth (Fig. 6 b,e). These trends can be explained with the broad large-scale patterns of environmental conditions along the global hypsometry. In deeper depositional environments, prolonged pelagic degradation (Cram et al., 2018; Maerz et al., 2019; Müller and Suess, 1979; Weber et al., 2016), exposure to oxygen (Hartnett et al., 1998), and the continuous ageing during settling (Griffith et al., 2010) result in a low apparent OM reactivity in underlying sediments. In contrast, several processes occurring in shallow environments, such as rapid settling, intense lateral transport, limited degradation in the water column, dynamic reworking through intense macrofaunal activity, and fluctuating redox conditions (e.g., Aller, 1994; Canfield, 1994; Hedges et al., 1997; Burdige, 2007) contribute to higher apparent OM reactivities. Local reactivity–water depth trends in shallow environments are driven by some of the same controls.

The weak water depth–reactivity trends and numerous exceptions at both global and regional level caution against the uncritical use of simplistic  $k$ – $h$  relationships. Further important controls often complicate and sometimes inhibit the use of water depth as a predictor for apparent OM reactivity. For instance, the Severn estuary is an exception to this observed local, shallow environmental trend. This region is characterized by one of the largest tidal ranges on Earth resulting in pronounced

510 environmental gradients along the dynamic land-ocean transition. OM of terrestrial, anthropogenic and marine sources mixes along the gradient and interacts with high loads of fine, suspended particular material. Additionally, intense tidally driven deposition-resuspension cycles (Dyer, 1984; Jonas and Millward, 2010; Manning et al., 2010; Uncles, 2010) result in a continuous re-exposure of benthic OM, which might cause the lower apparent reactivity of the buried OM. The Arabian Sea is another example where the  $k-h$  relationship deviates from the general global trends. Here, reactivity is 2–3 orders of magnitude higher than those derived for other deep-sea sites. Instead, reactivity is rather comparable to values determined for shallow sites in the Northern European margin and the Rhone delta. In addition, across the Arabian Sea OMZ transect, only a modest regional depth trend is observed, with a decrease in reactivity (i.e. increase in  $a$ ) from the OMZ to increasing water depths and bottom water oxygen concentrations (Cowie, 2005; Luff et al., 2000). While distinct from the overall  $k-h$  discussed above, the Arabian Sea OMZ region exhibits relatively similar patterns to other high productive, anoxic water column systems, e.g., Peru margin (Boudreau and Ruddick, 1991), Blake ridge (Marquardt et al., 2010; Wallmann et al., 2006), as well as California, Chile, Costa Rica, and Namibia margins (Marquardt et al., 2010). This suggests a general separation from anoxic and oxic water columns shaping benthic OM reactivity.

## 4.2 Influence of OM reactivity on benthic-pelagic coupling

525 Both quantity and quality of the deposited OM exerts an important control on benthic-pelagic coupling processes with potential implications for global biogeochemical cycles and climate. While the influence of OM quantity on benthic-pelagic exchange along the global hypsometry has been previously investigated (Krumins et al., 2013; Soetaert et al., 1996, 1998; Thullner et al., 2009), the role of reactivity on benthic-pelagic fluxes remains less constrained.

### 530 4.2.1 Control of OM reactivity on OM degradation rates and respiration pathways

In agreement with previous findings (Henrichs, 1992), depth-integrated OM degradation rates ( $R_{TOC,total}$ ; Eq. 7) show a strong correlation with OM fluxes ( $J_{OM,in}$ ) (Fig. 7a,  $r^2 = 0.97$ ;  $p < 0.01$ ;  $n = 14$ ). In contrast, they do not reveal a strong link with apparent OM reactivity (i.e., parameter  $a$ ), thus indicating that the magnitude of the OM deposition flux rather than its quality often exerts a first order control on depth-integrated rates of OM degradation. In our dataset, apparent OM reactivity only becomes a dominant control on  $R_{TOC,total}$  in depositional environments that receive exceptionally reactive OM (i.e.,  $a < 10^1$  yr), such as the Arabian Sea region (Table 6).

OM reactivity does exert an important impact on the relative contributions of metabolic pathways (Fig. 7). Overall, rates of OM degradation coupled to consumption of oxygen (aerobic OM degradation pathway) and nitrate (denitrification) make a small relative contribution (< 10%) to total degradation rates in the investigated oxic depositional environments (Rhone,

Skagerrak, Bering Sea, and Argentine Basin; Fig. 7b). The relative contributions quantified here are generally lower than previous estimates for coastal and shelf sediments (> 20%) (e.g., Long Island Sound, Mackin and Swider, 1989; Skagerrak, Canfield et al., 1993; Eastern Canadian continental margin, Boudreau et al., 1998; Barents Sea, Freitas et al., 2020; and Northern Gulf of Mexico, Owings et al., 2021). Additionally, estimates are consistently lower than values in the deep-sea (> 70%) (e.g., Cape Hatteras continental rise, Heggie et al., 1987), as well as over the global-scale hypsometry (Thullner et al., 2009). However, the overall dynamic nature of the considered environments, which favours the dominance of organoclastic sulfate reduction (Bowles et al., 2014; Jørgensen and Kasten, 2006; Thullner et al., 2009), as well as the length of sediment column (1,000 cmbsf) considered here renders a direct comparisons difficult. Here, the contribution of the aerobic OM degradation pathway further decreases with a decrease in OM apparent reactivity (i.e., increase in  $a$ ) as a consequence of the resulting higher burial of OM to sub-oxic or anoxic sediments (e.g., Meister et al., 2013). Additionally, the enhanced production of reduced species at depth further decreases the relative contribution of the aerobic OM degradation pathway to overall OM degradation rates by channelling more oxygen consumption through the re-oxidation of upward diffusing reduced species (e.g., ammonium and hydrogen sulfide) (Glud, 2008; Hensen et al., 2006).

It has been suggested that anoxic metabolic pathways represent significant pathways of OM degradation (e.g., Bowles et al., 2014; Jørgensen and Kasten, 2006; Thullner et al., 2009). Our results show that sulfate reduction is the main oxidative pathway in regions characterised by high apparent OM reactivity ( $a < 10^2$  yr) and intermediate OM fluxes ( $J_{OM,in} \sim 10^{-4}$ – $10^{-3}$  mol C  $\text{cm}^{-2}$   $\text{yr}^{-1}$ ), e.g., Arabian Sea region (Cowie, 2005; Luff et al., 2000), Arkona Basin and Aarhus Bay (Dale et al., 2008b; Jørgensen et al., 2019a; Mogollón et al., 2012). Additionally, high relative contributions of sulfate reduction are also simulated for environments that are characterized by a reduced supply of less reactive ( $a > 10^3$  yr) OM deposition fluxes, such as the Argentine Basin and Bering Sea sites (e.g., Hensen et al., 2003). While high apparent OM reactivity enhances sulfate consumption in the upper sediment, where sulfate is efficiently replenished by exchange with bottom waters, a low OM reactivity allows for a deeper sulfate penetration (Fig. 3 m-n) that sustains sulfate reduction over a greater sediment volume (e.g., Bowles et al., 2014; Meister et al., 2013; Riedinger et al., 2005, 2014, 2017). In contrast, methanogenesis is an important degradation pathway in environments that are characterized by a high supply ( $J_{OM,in} > 10^{-3}$ – $10^{-2}$  mol C  $\text{cm}^{-2}$   $\text{yr}^{-1}$ ) of both highly reactive ( $a = 10^0$ – $10^1$  yr) OM (Rhone proximal zone and Helgoland Mud Area, North Sea), as well as the least reactive ( $a > 10^3$  yr) OM (Skagerrak). The fast burial of reactive OM drives the development of a shallow methanogenic zone in the sediment (shallow sulfate penetration; Fig. 3 b,f) (e.g. Aromokeye et al., 2020; Oni et al., 2015a, b; Rassmann et al., 2020), whereas very low apparent reactivity OM escapes sulfate reduction and is degraded deeper in the sediment via methanogenesis (e.g., Dale et al., 2008a; Knab et al., 2008; Meister et al., 2013). In both cases, the produced methane diffuses up and supports an enhanced consumption of sulfate through AOM (Sect. 4.2.2). Consequently, the relative contribution of organoclastic sulfate reduction decreases due to the presence of methane (e.g., Bowles et al., 2014; Niewöhner et al., 1998; Regnier et al., 2011; Jørgensen and Kasten, 2006; Jørgensen et al., 2019a; Riedinger et al., 2005, 2014, 2017).

575 We do not consider the degradation pathways mediated by metal oxides due to a lack of SWI data that would allow constraining  
boundary conditions of those processes. This might result in a slight overrepresentation of sulfate reduction (e.g., at the  
Skagerrak; Canfield et al., 1993; Rysgaard et al., 2001). It has been demonstrated that metal oxide pathways represent a minor  
contribution to total OM heterotrophic degradation on a global scale (Thullner et al., 2009), although in continental margin  
sediments iron hydroxide reduction may represent an important OM degradation pathway (Beckler et al., 2016).

580

#### 4.2.2 Anaerobic oxidation of methane dynamics

AOM occurs at the sulfate-methane transition zone (SMTZ), which can vary in sediment depth from centimetres to hundreds  
of meters as a function of different environmental controls, such as OM quantity and quality, sedimentation rate, active fluid  
flow, or microbial growth dynamics (Puglino et al., 2020; Regnier et al., 2011). Reflecting the diversity of the depositional  
585 environments studied here, we observe a broad range of SMTZ depths (Fig. 8 a-c). Generally and in agreement with previous  
findings (Egger et al., 2018; Regnier et al., 2011), we observe a deepening of the SMTZ with decreasing sedimentation rates  
in addition to an important control of OM reactivity on the depth of the SMTZ (Fig. 8b) and the depth-integrated AOM rates  
( $\Sigma AOM$ , Fig. 8d).

590 The lowest  $\Sigma AOM$  and deepest SMTZ depths are determined for sites that are characterized by low apparent OM reactivity  
(i.e., deep-sea, well oxygenated bottom waters regions; Table 6). In those regions, low OM reactivity results in low rates of  
heterotrophic degradation (Fig. 7a) and, thus, deep  $SO_4^{2-}$  penetration (Fig. 3 m–n) or no  $SO_4^{2-}$  depletion at all. In contrast, high  
OM reactivity in combination with high depositional fluxes (Fig. 8 b–e) result in the complete depletion of TEAs in the upper  
sediment ( $z < 50$  cm), allowing the establishment of methanogenesis at relatively shallow sediment depth. For instance,  
595 sediments from the Helgoland Mud area, North Sea, are characterized by a shallow SMTZ and intense  $\Sigma AOM$  (Oni et al.,  
2015a, b). Yet, high OM reactivity does not always result in shallow SMTZ and intense AOM rates. Associated with moderate  
depositional fluxes, high apparent OM reactivity reduces the amount of reactive OM that reaches the methanogenic zone,  
inducing a deepening of the SMTZ and a decrease in  $\Sigma AOM$ . For instance, sediments from the Aarhus Bay (Flury et al., 2016)  
and Arkona Basin (Mogollón et al., 2012) display high OM reactivity ( $a < 10^1$  yr), which supports high rates of degradation  
600 close to the SWI (Fig. 7a) where  $SO_4^{2-}$  is constantly replenished from bottom waters and limits OM fluxes to the deeper,  
methanogenic sediment (e.g., Flury et al., 2016; Meister et al., 2013; Regnier et al., 2011). For moderate depositional fluxes,  
associated with lower OM reactivity ( $a > 10^2$  yr), e.g., the Skagerrak sites, pre-aged OM buried in these sediments (Aquilina  
et al., 2010) is predominantly degraded via methanogenesis (Fig. 7b). Consequently, a shoaling of the SMTZ is observed (Dale  
et al., 2008a; Knab et al., 2008), and higher  $\Sigma AOM$  are determined (Table 6).

605

### 4.2.3 OM reactivity controls on sediment-water interface fluxes patterns

Model results suggest that apparent OM reactivity also influences nutrient recycling at the seafloor. Thus, we estimate benthic-pelagic fluxes of dissolved species ( $O_2$ ,  $NO_3^-$ ,  $SO_4^{2-}$ ,  $NH_4^+$ , and  $PO_4^{3-}$ ). We emphasise that these flux calculations are based on simulated depth profiles at steady-state and are thus only weakly affected by sampling limitations (e.g., possible loss of uppermost sediment layers associated to the choice of sampling devices; Hensen et al., 2006). Our estimates allow us to investigate the links between nutrient recycling and OM reactivity and to assess qualitative differences across sites.

Apparent OM reactivity broadly controls the spatial patterns of benthic-pelagic fluxes, as well as the relative importance of different transport processes (Glud, 2008; Bourgeois et al., 2017). Overall, molecular diffusion is the main transport pathway (Fig. 9 f–j). However, the relative importance of bioirrigation increases at environments characterized by low deposition of OM and less reactive OM. This occurs because of the relative decrease in diffusive fluxes as a consequence of weak concentration gradients across the SWI. Bioturbation fluxes are generally low and reflect our assumption of depth-dependent  $D_{bio}$  (Middelburg et al., 1997). Advective fluxes become an important transport mechanism driving downward sulfate fluxes, especially in rapidly accumulating sediments and at sites characterized by low OM reactivity and moderate depositional rate.

Overall, high apparent OM reactivity drives high sedimentary uptake fluxes of dissolved TEA (Table 6; Fig. 9 a–c). Sulfate largely dominates the benthic TEA uptake, particularly at sites where oxygen and nitrate are unavailable TEA (Fig. 9c). At the Rhone proximal zone, unusually high sedimentation rates (Pastor et al., 2011) deliver fresh and highly reactive OM (Cathalot et al., 2010; Pruski et al., 2015), supporting high OM fluxes and degradation rates (Fig. 7a; Table 6). Consequently, intense oxygen consumption occurs at the SWI (Rassmann et al., 2016) driven by both aerobic degradation of OM and re-oxidation of reduced species. Similarly, in the Arabian Sea sediments below the OMZ the high input of fresh phytoplankton debris (Cowie, 2005; Koho et al., 2013; Rixen et al., 2019; Vandewiele et al., 2009) and potentially chemoautotroph biomass (Lengger et al., 2019) associated with intense degradation rates in the uppermost sediment layers (Luff et al., 2000), result in a high relative contributions of aerobic degradation of OM and denitrification to total OM oxidation. Consequently, both benthic oxygen and nitrate uptake fluxes are comparably high. In contrast, at regions characterised by deposition of less reactive OM ( $k < 10^{-4} \text{ yr}^{-1}$ ; Skagerrak, Argentine Basin, and Bering Sea), benthic-pelagic fluxes of TEAs are generally low.

Benthic fluxes exert a crucial role in nutrient recycling. In the Arabian Sea region, large phytoplankton blooms are associated with monsoon conditions, which result in upwelling of nutrient-rich bottom water (SW monsoon) and deepening of the mixed-layer (NE monsoon) (Cowie, 2005; Luff et al., 2000; Rixen et al., 2019). Similarly, across the Northern European margin, spring and summer diatom blooms are common (Fleming-Lehtinen and Laamanen, 2012; Jensen et al., 1990; Karlson et al., 1996; Lomstein et al., 1990; Wiltshire and Manly, 2004) and maintained by benthic nutrient fluxes. Additionally, bottom water upwelling and nutrient recycling is an important mechanism sustaining spring and summer PP at the Bering Sea (Coyle et al.,

2008). Benthic ammonium and phosphate recycling fluxes mirror the reactivity and OM degradation patterns (Fig. 9 d–e). The largest fluxes are determined for the Arabian Sea region, as well as the shallow Aarhus Bay and Arkona Basin (Table 6). In contrast, the lowest nutrient recycling fluxes are observed in regions characterized by the deposition of less reactive OM (phosphate fluxes are negligible at Skagerrak sites S10 and S13), where heterotrophic degradation rates are slow.

## 5. Conclusions

We developed and applied an inverse modelling approach to quantify apparent OM reactivity (i.e., parameters  $a$  and  $v$ ) based on a common minimum set of benthic observational data, as well as a common RTM approach for 14 different sites across five different depositional environments. We provide important first-order constraints on the Reactive Continuum Model (RCM) parameterization that can inform on the choice of parameter values in data poor areas, including global-scale and timescale scenarios.

Our findings corroborate previous results (Arndt et al., 2013; Boudreau and Ruddick, 1991; Forney and Rothman, 2012; Middelburg, 1989) that the RCM parameter  $v$  is globally relatively constant ( $v = 0.1–0.2$ ). Exceptionally high  $v > 0.2$  are often associated with deposition and burial of highly reactive phytoplankton debris in high productivity regions associated with well-established OMZs. In contrast, in agreement with previous findings, RCM parameter  $a$  can span several orders of magnitude at a global scale, suggesting that the parameter  $a$  is the main parameter describing the variability of apparent OM reactivity. Consequently, future modelling efforts to quantify OM reactivity on a global scale could be reduced to one main reactivity variable. Based on our results and previous findings, we consolidate the range of predominant  $a$  distribution to  $10^0–10^4$  years. This is a valuable constraint when dealing with data poor regions and timescales (e.g., Hülse et al., 2018), since it excludes extreme values at both ends of the  $a$ -range.

Additionally, results indicate that the large variability in apparent OM reactivity is linked to a combination of multiple environmental drivers that control the quality of OM delivered to sediments, and the timescale of settling and burial. Therefore, our findings contribute to the notion that apparent OM reactivity is controlled by a complex and dynamic interplay of environmental drivers, which are measurable and allow the quantification of  $a$ ,  $v$ , and  $k$ . Our model results caution against the use of single environmental master variable such as water depth, sedimentation rate or OM deposition fluxes beyond well-defined geographical scales. Yet, results show that, if no other information is available, water depth can serve as a useful proxy for the complex and dynamic interplay of environmental drivers and can be used to predict first-order, large scale OM reactivity patterns. However, more holistic, interdisciplinary exploration of OM reactivity in its entire environmental context is needed to mechanistically understand benthic carbon and nutrient dynamics.

670 Finally, results also show that OM reactivity exerts a dominant control on the redox zonation of the sediment, the depth of the SMTZ, depth-integrated AOM rates and benthic-pelagic exchange fluxes indicating that these processes could serve as predictor variables for OM reactivity. In contrast, depth-integrated OM degradation rates are largely controlled by the magnitude rather than the quality of OM deposition to the sediment.

### **Acknowledgements**

675 FSF was supported by a PhD scholarship from the Science without Borders Programme (Ciência sem Fronteiras), sponsored by the CAPES Foundation within Ministry of Education, Brazil, grant number 9999.009541/2013-06. FSF also thanks funding from the UKRI Natural Environment Research Council (NERC), grant number NE/P006493/1 (Changing Arctic Ocean Seafloor – ChAOS). RDP acknowledges the advanced ERC grant ‘The greenhouse earth system’ (T-GRES, project reference 340923) and the Royal Society Wolfson Research Merit Award. SA and PAP acknowledge funding from the European Union’s  
680 Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement number 643052 745 (C-CASCADES). ST PhD was funded by a President’s Scholarship, Cardiff University. CR and JR were supported by the INSU/EC2CO program MissRhoDia and by the French state program “Investissement d’avenir” run by the National Research Agency (AMORAD project ANR-11-RSNR-0002). The authors also thank Bernard Boudreau and an anonymous reviewer, as well Associate Editor Marilaure Grégoire for their constructive comments on early versions of the manuscript. Finally, we  
685 thank Co-Editor in Chief Steve Bouillon for granting us a discount on the APC.

### **Authors contributions**

FSF, SA, and RDP designed this study. PAP developed and implemented changes to model description (multi-G approximation of the RCM). SK, BBJ, JR, CR, ST, and HS provided porewater and sediment data to inform model experiments and contributed to discussing and interpreting the data and model results. FSF compiled data, ran all model simulations, and  
690 performed data-model integration. FSF wrote the manuscript with significant contributions from SA. All co-authors edited and approved the manuscript.

### **Code/data availability**

The model code is available at the GitHub repository ([https://github.com/felipesalesdefreitas/BG\\_2020\\_435](https://github.com/felipesalesdefreitas/BG_2020_435)). The data is available on PANGAEA database ([www.pangaea.de](http://www.pangaea.de)) where indicated, or at cited literature along the manuscript.

### **695 Competing interests**

The authors declare no conflicts of interest.

**Table 1. Environmental factors influencing organic matter reactivity in marine sediments.**

<b>Description</b>	<b>Reference</b>
Sources and composition	Burdige (2006); Canfield (1994); Hedges et al. (1997); Meyers (1997); Middelburg (2019)
Oxygen exposure time (settling and burial)	Hartnett et al. (1998); Hoefs et al. (2002); Sinninghe Damsté et al. (2002); Huguet et al. (2008); Sun and Wakeham (1994); Wakeham et al. (1997)
Terminal electron acceptors availability	Aller (1994); Aller and Aller (1998); Canfield et al. (1993); Canfield (1994)
Microbial activity	LaRowe et al. (2020a, b); Stevenson et al. (2020); Bradley et al. (2020)
Sediment biological mixing (bioturbation and bioirrigation)	Aller (1980, 1994); Aller and Aller (1998); Boudreau (1994); Grossi et al. (2003); Middelburg (2018); Aller and Cochran (2019)
Ageing and transport history	Bianchi et al. (2018); Cathalot et al. (2010); Griffith et al. (2010); Mollenhauer et al. (2003, 2007); Ohkouchi et al. (2002); Ausín et al. (2021)
OM—sediment association	Bianchi et al. (2018); Hemingway et al. (2019); Keil and Cowie (1999); Mayer (1994b, a)
OM—iron minerals adsorption	Salvadó et al. (2015); Shields et al. (2016); Wang et al. (2019); Faust et al. (2021)
Priming	van Nugteren et al. (2009); Guenet et al. (2010); Bianchi (2011); Bianchi et al. (2015); Ragavan and Kumar (2021)
Hydrogen sulfide exposure time (sulfurization)	Sinninghe Damsté et al. (1988, 1998); Hülse et al. (2019)

700

705

710

**Table 2. Oceanographic context of depositional environments studied in the large-scale organic matter reactivity quantification.**

	Site	Lat (°)	Climate zone	Setting	Sed. rate (cm yr <sup>-1</sup> )	SWI redox condition	TOC (wt%)	Δ <sup>14</sup> C (‰)	OM Type
	Core ID	Lon (°)	Benthic Province	Depth (m)	Bottom temp. (° C)		OC:SA (mg m <sup>-2</sup> )	F <sub>mod</sub>	Input
A	Severn estuary	51.50 N	Temperate	Coast	4.3·10 <sup>-1</sup>	Sub-oxic	3.0	N/A	Largely Terrestrial
	ST-1	3.07 W	EUR1	8	18.0		N/A	N/A	Fluvial
B	Rhone proximal zone	43.31 N	Temperate / Mediterranean	Coast	1.0·10 <sup>1</sup>	Oxic	2.0	143 ± 14	Largely terrestrial and aquatic
	A	4.85 E	EUR2	19	15.0		N/A	N/A	Fluvial
C	Rhone shelf	43.27 N	Temperate / Mediterranean	Shelf	5.0·10 <sup>-1</sup>	Oxic	1.3	-218 ± 9	Mixed terrestrial and marine
	C	4.77 E	EUR2	74	14.0		0.4 – 0.7	N/A	Fluvial and pelagic
D	Aarhus Bay	56.11 N	Temperate	Coast	3.2·10 <sup>-1</sup>	Sub-oxic / anoxic (summer)	3.5	N/A	Largely terrestrial with marine contribution
	173GC 174GC	10.34 E	EUR1	15	9.0		N/A	N/A	Fluvial and pelagic
E	Arkona Basin	54.80 N	Temperate	Shelf	7.4·10 <sup>-3</sup>	Sub-oxic	5.0	N/A	Mostly terrestrial
	348GC 351GC	13.78 E	EUR1	43	5.0		N/A	N/A	Fluvial
F	Helgoland Mud Area, North Sea	54.08 N	Temperate	Shelf	1.3·10 <sup>0</sup>	Anoxic	1.1	-174 – 195	Largely terrestrial with marine contribution
	HE443-010	7.97 E	EUR1	29	10.0		N/A	N/A	Pelagic and fluvial
G	Skagerrak S10	57.92 N	Temperate	Shelf	5.0·10 <sup>-1</sup>	Oxic	1.3	N/A	Largely terrestrial
	807GC 808GC	9.75 E	EUR1	86	8.7		0.6 – 0.7	N/A	Fluvial and pelagic

H	Skagerrak S11	57.95 N	Temperate	Shelf	$5.0 \cdot 10^{-1}$	Oxic	0.5	N/A	Largely terrestrial
	816GC 836GC	9.70 E	EUR1	150	9.8		0.6 – 0.7	N/A	Fluvial and pelagic
I	Skagerrak S13	58.05 N	Temperate	Slope	$5.0 \cdot 10^{-1}$	Oxic	1.8	N/A	Largely terrestrial
	786GC 789GC	9.60 E	EUR1	386	4.5		0.6 – 0.7	N/A	Fluvial and pelagic
J	Arabian Sea (Core of OMZ)	24.88 N	Arid	Slope	$5.0 \cdot 10^{-2}$	Anoxic	0.92	N/A	Largely marine
	GeoB 12312	63.02 E	WARAB	654	11.6		1.1 – 2.3	0.800 – 0.920	Pelagic
K	Arabian Sea (OMZ transition)	24.81 N	Arid	Slope	$5.0 \cdot 10^{-2}$	Anoxic	0.89	N/A	Largely marine
	GeoB 12309	62.99 E	WARAB	957	9.6		1.1 – 2.3	0.800 – 0.920	Pelagic
L	Arabian Sea (Below OMZ)	24.71 N	Arid	Slope	$5.0 \cdot 10^{-2}$	Oxic	0.87	N/A	Largely marine
	GeoB 12308	62.99 E	WARAB	1,586	5.1		0.4 – 1.1	0.800 – 0.920	Pelagic
M	Bering Sea	54.57 N	Snow / Polar	Slope	$1.6 \cdot 10^{-3}$	Oxic	1.3	-143	Largely marine
	SO202-22	168.81 W	NWPAC	1,476	2.0		N/A	0.857	Pelagic
N	Argentine Basin	39.31 S	Temperate	Slope	$8.0 \cdot 10^{-3}$	Oxic	1.0	N/A	Mixed marine and terrestrial
	GeoB 13863	53.95 W	RIOPLAT	3,687	3.5		N/A	0.777	Pelagic and lateral transport

**Table 3. Summary of model elements incorporated in the BRNS.**

<b>Symbol</b>	<b>Description</b>
	<i>Chemical species, i</i>
TOC	Total organic carbon
CH <sub>2</sub> O	Organic matter (simplified stoichiometry)
O <sub>2</sub>	Oxygen
NO <sub>3</sub> <sup>-</sup>	Nitrate
SO <sub>4</sub> <sup>2-</sup>	Sulfate
CH <sub>4</sub>	Methane
NH <sub>4</sub> <sup>+</sup>	Ammonium
PO <sub>4</sub> <sup>3-</sup>	Phosphate
HS <sup>-</sup>	Sulfides
	<i>Model parameters</i>
C <sub>i</sub>	Concentration of species <i>i</i>
<i>t</i>	Time
<i>z</i>	Sediment depth
<i>L</i>	Length of model domain
<i>T</i>	Temperature
<i>S</i>	Salinity
<i>h</i>	Water depth
	Porosity term
$\sigma$	Solid species, $\sigma = 1 - \varphi$ Dissolved species, $\sigma = \varphi$
$\varphi$	Sediment porosity
D <sub>i</sub>	Effective molecular diffusion coefficient of dissolved species <i>i</i>
D <sub>bio</sub>	Bioturbation diffusion coefficient
z <sub>bio</sub>	Depth of bioturbated zone – sediment mixed layer
$\omega$	Burial velocity – sedimentation rates
$\alpha_i$	Bioirrigation coefficient
s <sub>i</sub> <sup>n</sup>	Stoichiometric coefficient of specie <i>i</i>
<i>n</i>	Kinetically controlled reaction
$\tau_n$	Reaction rate
k <sub>i</sub>	First-order reaction rate constant
$\Gamma(v)$	Gamma distribution
<i>a</i>	Reactive continuum model shaping parameter
<i>v</i>	Reactive continuum model scaling parameter
om( <i>k, t</i> )	Probability density function of OM distribution
	<i>Quantification of model-derived parameters</i>
k( <i>z</i> )	Apparent organic matter reactivity at a given depth
R <sub>TOC</sub>	Organic matter degradation rate
R <sub>TOC,total</sub>	Depth integrated organic matter degradation rates
J <sub>OM,in</sub>	Fluxes of organic matter at the sediment-water interface
J <sub>OM,out</sub>	Fluxes of organic matter consumed by heterotrophic degradation
J <sub>OM,bur</sub>	Fluxes of organic matter buried (below model domain, <i>L</i> )
J <sub>i</sub>	Fluxes of species <i>i</i> at the sediment-water interface

**Table 4. Site-specific transport parameters and upper boundary conditions implemented in the Reaction-Transport Model.**

Site	<i>Site-specific transport parameters</i>							<i>Site-specific upper boundary conditions</i>					
	$\varphi$	$D_{bio}$ cm <sup>2</sup> yr <sup>-1</sup>	$z_{bio}$ cm	$\omega$ cm yr <sup>-1</sup>	$T$ °C	$S$	$h$ m	$TOC$ wt%	$O_2$ μM	$NO_3^-$ μM	$SO_4^{2-}$ mM	$CH_4$ μM	$HS^-$ mM
A	0.70	29.8	9	$4.3 \cdot 10^{-1}$	18.0	35	8	3.0	0	30	23	0	0
B	0.80	0.50	10	$1.0 \cdot 10^1$	16.8	35	18	2.0	245	25	28	0	0
C	0.83	10.0	10	$5.0 \cdot 10^{-1}$	14.4	35	75	1.3	225	25	29	0	0
D	0.86	0	0	$3.2 \cdot 10^{-1}$	12.0	35	15	3.5	0	0	20	0	0
E	0.86	0	0	$7.4 \cdot 10^{-3}$	5.0	35	43	5.0	0	0	6	0	0
F	0.70	0	0	$1.3 \cdot 10^0$	10.0	35	30	1.1	0	0	26	0	0
G	0.44	27.8	10	$5.0 \cdot 10^{-1}$	8.7	35	86	1.3	250	25	29	0	0
H	0.47	26.3	10	$5.0 \cdot 10^{-1}$	9.8	35	147	0.5	250	25	29	0	0
I	0.50	0	0	$5.0 \cdot 10^{-1}$	4,5	35	386	1.8	0	0	22	0	0
J	0.60	0	0	$5.0 \cdot 10^{-2}$	11.6	35	632	5.0	0	0	25	0	0
K	0.60	0	0	$5.0 \cdot 10^{-2}$	9.6	35	930	4.4	0	0	28	0	0
L	0.70	7.3	10	$5.0 \cdot 10^{-2}$	5.1	35	1,552	4.3	250	25	29	0	0
M	0.60	7.8	10	$1.6 \cdot 10^{-3}$	2.0	35	1,469	1.3	250	25	28	0	0
N	0.70	1.0	10	$8.0 \cdot 10^{-3}$	3.5	35	3,687	1.0	250	25	28	0	0

720

725

730


**Table 5. Summary of statistical tests for the best fit  $\alpha$ — $\nu$  pair at each site derived from Taylor Diagrams.**

Site	Reactivity parameters			TOC				$SO_4^{2-}$			
	$\alpha$	$\nu$		Mean	SD	RMSD	$r$	Mean	SD	RSMD	$r$
A	$5.0 \cdot 10^1$	0.100	data	2.97	0.03			18.49	3.57		
			model	4.90	1.62	1.64	0.513	17.51	3.44	0.82	0.947
B	$3.0 \cdot 10^0$	0.125	data	1.96	0.02			14.10	8.07		
			model	1.92	0.16	0.14	0.640	12.59	9.18	1.79	0.974
C	$1.0 \cdot 10^2$	0.125	data	1.28	0.01			27.51	0.80		
			model	1.29	0.01	0.005	0.9998	27.76	0.89	0.37	0.823
D	$1.0 \cdot 10^0$	0.060	data	2.41	0.11			6.00	6.18		
			model	1.63	0.86	0.77	0.738	5.51	5.83	0.57	0.994
E	$1.0 \cdot 10^0$	0.155	data	2.17	0.60			1.54	1.83		
			model	1.79	0.46	0.40	0.561	1.35	1.84	0.23	0.984
F	$3.0 \cdot 10^1$	0.125	data	0.89	0.22			1.71	4.93		
			model	0.89	0.08	0.25	0.014	1.75	5.18	0.76	0.980
G	$1.8 \cdot 10^3$	0.100	data	1.12	0.45			9.54	9.15		
			model	1.34	0.00	0.45	0.597	9.70	10.08	1.52	0.984
H	$1.0 \cdot 10^1$	0.130	data	0.47	0.18			4.40	7.10		
			model	0.38	0.04	0.20	0.240	5.88	8.84	2.73	0.931
I	$3.5 \cdot 10^3$	0.110	data	1.69	0.34			3.51	5.84		
			model	1.84	0.00	0.34	0.002	3.35	6.23	0.64	0.993
J	$1.0 \cdot 10^0$	0.250	data	0.67	0.13			3.26	5.67		
			model	0.78	0.22	0.19	0.289	2.81	5.21	0.50	0.999
K	$1.0 \cdot 10^1$	0.390	data	0.56	0.12			5.30	7.79		
			model	0.48	0.24	0.22	0.154	4.73	6.97	0.90	0.998
L	$1.0 \cdot 10^1$	0.330	data	0.49	0.15			17.67	7.45		
			model	0.78	0.42	0.33	0.544	9.62	6.14	2.74	0.878
M	$5.0 \cdot 10^3$	0.270	data	0.82	0.34			10.38	9.93		
			model	0.94	0.17	0.24	0.566	7.52	7.74	2.79	0.961
N	$2.0 \cdot 10^3$	0.125	data	0.84	0.07			7.56	8.89		
			model	0.78	0.12	0.10	0.298	7.32	8.72	0.62	0.995


735 **Table 6. Benthic organic matter dynamics derived from the coupled model-data assessments across the depositional settings.**

Site	<i>OM reactivity</i>			<i>OM degradation dynamics</i>				<i>Benthic-pelagic fluxes</i>				
	$\alpha$	$\nu$	$k(0)$	$J_{OM,in}$	$R_{TOC,total}$	$\Sigma AOM$	$z_{AOM}$	$J_{O_2}$	$J_{NO_3}$	$J_{SO_4}$	$J_{NH_4}$	$J_{PO_4}$
	yr		yr <sup>-1</sup>	mol cm <sup>-2</sup> yr <sup>-1</sup>			cmbsf	μmol cm <sup>-2</sup> yr <sup>-1</sup>				
A	5.0 ·10 <sup>1</sup>	0.100	2.0 ·10 <sup>-3</sup>	1.55 ·10 <sup>-3</sup>	9.89 ·10 <sup>-4</sup>	9.2 ·10 <sup>-5</sup>	35	n.d	-145	-178	39.9	0.93
B	3.0 ·10 <sup>0</sup>	0.125	4.2 ·10 <sup>-2</sup>	2.04 ·10 <sup>-2</sup>	1.50 ·10 <sup>-2</sup>	1.6 ·10 <sup>-4</sup>	24	-1,649	-132	-612	110	0.79
C	1.0 ·10 <sup>2</sup>	0.125	1.3 ·10 <sup>-3</sup>	5.76 ·10 <sup>-4</sup>	4.19 ·10 <sup>-4</sup>	4.8 ·10 <sup>-6</sup>	537	-81.2	-1.3	-30.3	9.8	0.03
D	1.0 ·10 <sup>0</sup>	0.060	6.0 ·10 <sup>-2</sup>	1.05 ·10 <sup>-3</sup>	8.55 ·10 <sup>-4</sup>	1.2 ·10 <sup>-5</sup>	185	n.d	n.d	-462	278	5.6
E	1.0 ·10 <sup>0</sup>	0.155	1.6 10 <sup>-1</sup>	7.42 ·10 <sup>-4</sup>	7.17 ·10 <sup>-4</sup>	8.8 ·10 <sup>-6</sup>	68	n.d	n.d	-715	467	9.1
F	3.0 ·10 <sup>1</sup>	0.125	4.2 ·10 <sup>-3</sup>	1.59 ·10 <sup>-3</sup>	1.00 ·10 <sup>-3</sup>	5.0 ·10 <sup>-5</sup>	58	n.d	n.d	-98.7	24.1	0.22
G	1.8 ·10 <sup>3</sup>	0.100	5.6 ·10 <sup>-5</sup>	8.27 ·10 <sup>-4</sup>	1.00 ·10 <sup>-4</sup>	1.7 ·10 <sup>-5</sup>	145	-36.7	-0.77	-26.2	4.1	0
H	1.0 ·10 <sup>1</sup>	0.130	1.3 ·10 <sup>-2</sup>	4.10 ·10 <sup>-4</sup>	2.74 ·10 <sup>-4</sup>	3.2 ·10 <sup>-5</sup>	68	-472	-2.7	-108	65.5	1.4
I	3.5 ·10 <sup>3</sup>	0.110	3.1 ·10 <sup>-5</sup>	9.9 ·10 <sup>-4</sup>	9.04 ·10 <sup>-5</sup>	1.8 ·10 <sup>-5</sup>	98	n.d	n.d	-16.8	1.9	0
J	1.0 10 <sup>0</sup>	0.250	2.5 ·10 <sup>-1</sup>	7.51 ·10 <sup>-4</sup>	7.30 ·10 <sup>-4</sup>	7.8 ·10 <sup>-6</sup>	195	n.d	n.d	-2,816	1,824	36.9
K	1.0 ·10 <sup>1</sup>	0.390	3.9 ·10 <sup>-2</sup>	4.88 ·10 <sup>-4</sup>	4.78 ·10 <sup>-4</sup>	5.1 ·10 <sup>-6</sup>	255	n.d	n.d	-968	611	12.9
L	1.0 ·10 <sup>1</sup>	0.330	3.3 ·10 <sup>-2</sup>	1.31 ·10 <sup>-3</sup>	1.30 ·10 <sup>-3</sup>	2.4 ·10 <sup>-6</sup>	410	-1,021	-96.8	-642	439	10.8
M	5.0 ·10 <sup>3</sup>	0.270	5.4 ·10 <sup>-5</sup>	3.72 ·10 <sup>-5</sup>	2.82 ·10 <sup>-5</sup>	1.4 ·10 <sup>-6</sup>	705	-17.3	-0.42	-5.1	2.7	0.03
N	2.0 ·10 <sup>3</sup>	0.125	6.3 ·10 <sup>-5</sup>	6.0 ·10 <sup>-5</sup>	3.06 ·10 <sup>-5</sup>	2.3 ·10 <sup>-6</sup>	527	-17.7	-0.36	-5.0	2.5	0.01

n.d.: not determined (oxygen and nitrate benthic fluxes not determined for locations where SWI concentrations of those TEA were assumed to be zero).




745 **Figure 1. Depositional environment geographic locations. (a) Global scale; (b) European margin; (c) Arabian Sea region. Map produced with Ocean Data View (Schlitzer, 2019).**




**Figure 2. Total organic carbon data-model best fit: (a) Severn estuary; (b) Rhone pro-delta; (c) Rhone shelf; (d) Aarhus Bay; (e) Arkona Basin; (f) Helgoland Mud Area, North Sea; (g) Skagerrak – S10; (h) Skagerrak – S11; (i) Skagerrak – S13; (j) Arabian Sea – OMZ core; (k) Arabian Sea – OMZ transition; (l) Arabian Sea – below OMZ; (m) Bering Sea; (n) Argentine Basin.**

750




755 **Figure 3. Sulfate data-model best-fit: (a) Severn estuary; (b) Rhone pro-delta; (c) Rhone shelf; (d) Aarhus Bay; (e) Arkona Basin; (f) Helgoland Mud Area, North Sea; (g) Skagerrak – S10; (h) Skagerrak – S11; (i) Skagerrak – S13; (j) Arabian Sea – OMZ core; (k) Arabian Sea – OMZ transition; (l) Arabian Sea – below OMZ; (m) Bering Sea; (n) Argentine Basin.**

760




**Figure 4. Frequency histogram of organic matter reactivity parameters distribution in a global scale: (a) scaling parameter —  $\nu$ ; (b) shaping parameter —  $a$ .**




**Figure 5. Apparent organic matter reactivity  $k$  initial probability distributions and downcore evolution as function of burial depths: (a) Severn estuary; (b) Rhone pro-delta; (c) Rhone shelf; (d) Aarhus Bay; (e) Arkona Basin; (f) Helgoland Mud Area, North Sea; (g) Skagerrak – S10; (h) Skagerrak – S11; (i) Skagerrak – S13; (j) Arabian Sea – OMZ core; (k) Arabian Sea – OMZ transition; (l) Arabian Sea – below OMZ; (m) Bering Sea; (n) Argentine Basin.**


770



**Figure 6.** Depositional environment controls on organic matter reactivity parameters  $a$  (a – d) and  $k$  (e – g): (a)  $v$  vs.  $a$ ; (b, e) water depth  $h$ ; (c, f) sedimentation rate  $\omega$ ; (d, g) organic matter fluxes  $J_{OM,in}$ .




**Figure 7. Benthic organic matter degradation dynamics. (a) organic matter fluxes vs. depth-integrated organic matter degradation rates; (b) Relative contribution of each metabolic pathway to heterotrophic organic matter degradation.**



780

**Figure 8. Anaerobic oxidation of methane (AOM) patterns as a function of apparent organic matter reactivity and depositional rates. (a) AOM rates vs. SMTZ depth; SMTZ depths vs. (b) reactivity parameter  $a$  and (c) sedimentation rates  $\omega$ ; AOM rates vs. (d) reactivity parameter  $a$  and (e) sedimentation rates  $\omega$ .**

785



**Figure 9. Benthic fluxes of dissolved species (a–e) and transport mechanism relative contributions (f–j): (a,f) oxygen; (b,g) nitrate; (c,h) sulfate; (d,i) ammonium; (e,j) phosphate.**

## References

- 790 Aguilera, D. R., Jourabchi, P., Spiteri, C., and Regnier, P.: A knowledge-based reactive transport approach for the simulation of biogeochemical dynamics in Earth systems, *Geochim. Geophys. Geosystems*, 6, 1–18, <https://doi.org/10.1029/2004GC000899>, 2005.
- Aller, R. C.: Diagenetic Processes Near the Sediment-Water Interface of Long Island Sound. I., in: *Advances in Geophysics*, vol. 22, Elsevier, 237–350, [https://doi.org/10.1016/S0065-2687\(08\)60067-9](https://doi.org/10.1016/S0065-2687(08)60067-9), 1980.
- 795 Aller, R. C.: Bioturbation and remineralization of sedimentary organic matter: effects of redox oscillation, *Chem. Geol.*, 114, 331–345, [https://doi.org/10.1016/0009-2541\(94\)90062-0](https://doi.org/10.1016/0009-2541(94)90062-0), 1994.
- Aller, R. C. and Aller, J. Y.: The effect of biogenic irrigation intensity and solute exchange on diagenetic reaction rates in marine sediments, *J. Mar. Res.*, 56, 905–936, <https://doi.org/10.1357/002224098321667413>, 1998.
- Aller, R. C. and Cochran, J. K.: The Critical Role of Bioturbation for Particle Dynamics, Priming Potential, and Organic C  
800 Remineralization in Marine Sediments: Local and Basin Scales, *Front. Earth Sci.*, 7, 157, <https://doi.org/10.3389/feart.2019.00157>, 2019.
- Antonelli, C., Eyrolle, F., Rolland, B., Provansal, M., and Sabatier, F.: Suspended sediment and <sup>137</sup>Cs fluxes during the exceptional December 2003 flood in the Rhone River, southeast France, *Geomorphology*, 95, 350–360, <https://doi.org/10.1016/j.geomorph.2007.06.007>, 2008.
- 805 Aquilina, A., Knab, N. J., Knittel, K., Kaur, G., Geissler, A., Kelly, S. P., Fossing, H., Boot, C. S., Parkes, R. J., Mills, R. A., Boetius, A., Lloyd, J. R., and Pancost, R. D.: Biomarker indicators for anaerobic oxidizers of methane in brackish-marine sediments with diffusive methane fluxes, *Org. Geochem.*, 41, 414–426, <https://doi.org/10.1016/j.orggeochem.2009.09.009>, 2010.
- Aris, R.: Prolegomena to the rational analysis of systems of chemical reactions II. Some addenda, *Arch. Ration. Mech. Anal.*, 27, 356–364, <https://doi.org/10.1007/BF00251438>, 1968.
- 810 Arndt, S., Hetzel, A., and Brumsack, H.-J.: Evolution of organic matter degradation in Cretaceous black shales inferred from authigenic barite: A reaction-transport model, *Geochim. Cosmochim. Acta*, 73, 2000–2022, <https://doi.org/10.1016/j.gca.2009.01.018>, 2009.
- Arndt, S., Jørgensen, B. B., LaRowe, D. E., Middelburg, J. J., Pancost, R. D., and Regnier, P.: Quantifying the degradation  
815 of organic matter in marine sediments: A review and synthesis, *Earth-Sci. Rev.*, 123, 53–86, <https://doi.org/10.1016/j.earscirev.2013.06.001>, 2013.
- Aromokeye, D. A., Oni, O. E., Tebben, J., Yin, X., Richter-Heitmann, T., Wendt, J., Nimzyk, R., Littmann, S., Tienken, D., Kulkarni, A. C., Henkel, S., Hinrichs, K.-U., Elvert, M., Harder, T., Kasten, S., and Friedrich, M. W.: Crystalline iron oxides stimulate methanogenic benzoate degradation in marine sediment-derived enrichment cultures, *ISME J.*, <https://doi.org/10.1038/s41396-020-00824-7>, 2020.
- 820 Ausín, B., Bruni, E., Haghipour, N., Welte, C., Bernasconi, S. M., and Eglinton, T. I.: Controls on the abundance, provenance and age of organic carbon buried in continental margin sediments, *Earth Planet. Sci. Lett.*, 558, 116759, <https://doi.org/10.1016/j.epsl.2021.116759>, 2021.
- Bach, L. T., Stange, P., Taucher, J., Achterberg, E. P., Algueró-Muñoz, M., Horn, H., Esposito, M., and Riebesell, U.: The  
825 Influence of Plankton Community Structure on Sinking Velocity and Remineralization Rate of Marine Aggregates, *Glob. Biogeochem. Cycles*, 33, 971–994, <https://doi.org/10.1029/2019GB006256>, 2019.

- Barlow, R. G., Mantoura, R. F. C., and Cummings, D. G.: Monsoonal influence on the distribution of phytoplankton pigments in the Arabian Sea, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 46, 677–699, [https://doi.org/10.1016/S0967-0645\(98\)00123-4](https://doi.org/10.1016/S0967-0645(98)00123-4), 1999.
- 830 Beckler, J. S., Kiriazis, N., Rabouille, C., Stewart, F. J., and Taillefert, M.: Importance of microbial iron reduction in deep sediments of river-dominated continental-margins, *Mar. Chem.*, 178, 22–34, <https://doi.org/10.1016/j.marchem.2015.12.003>, 2016.
- Berner, R. A.: *Early diagenesis: a theoretical approach*, Princeton University Press, Princeton, N.J., 241 pp., 1980.
- Berner, R. A.: The long-term carbon cycle, fossil fuels and atmospheric composition, *Nature*, 426, 323–326, <https://doi.org/10.1038/nature02131>, 2003.
- 835 Bianchi, T. S.: The role of terrestrially derived organic carbon in the coastal ocean: A changing paradigm and the priming effect, *Proc. Natl. Acad. Sci.*, 108, 19473, <https://doi.org/10.1073/pnas.1017982108>, 2011.
- Bianchi, T. S., Thornton, D. C. O., Yvon-Lewis, S. A., King, G. M., Eglinton, T. I., Shields, M. R., Ward, N. D., and Curtis, J.: Positive priming of terrestrially derived dissolved organic matter in a freshwater microcosm system, *Geophys. Res. Lett.*, 42, 5460–5467, <https://doi.org/10.1002/2015GL064765>, 2015.
- 840 Bianchi, T. S., Cui, X., Blair, N. E., Burdige, D. J., Eglinton, T. I., and Galy, V.: Centers of organic carbon burial and oxidation at the land-ocean interface, *Org. Geochem.*, 115, 138–155, <https://doi.org/10.1016/j.orggeochem.2017.09.008>, 2018.
- Bogus, K. A., Zonneveld, K. A. F., Fischer, D., Kasten, S., Bohrmann, G., and Versteegh, G. J. M.: The effect of meter-scale lateral oxygen gradients at the sediment-water interface on selected organic matter based alteration, productivity and temperature proxies, *Biogeosciences*, 9, 1553–1570, <https://doi.org/10.5194/bg-9-1553-2012>, 2012.
- 845 Bohrmann, G. and cruise participants: Documentation of sediment core GeoB12308-5, PANGAEA, <https://doi.org/10.1594/PANGAEA.737529>, 2010a.
- Bohrmann, G. and cruise participants: Documentation of sediment core GeoB12309-3, PANGAEA, <https://doi.org/10.1594/PANGAEA.737530>, 2010b.
- 850 Bohrmann, G. and cruise participants: Documentation of sediment core GeoB12312-3, PANGAEA, <https://doi.org/10.1594/PANGAEA.737532>, 2010c.
- Boudreau, B. P.: Is burial velocity a master parameter for bioturbation?, *Geochim. Cosmochim. Acta*, 58, 1243–1249, [https://doi.org/10.1016/0016-7037\(94\)90378-6](https://doi.org/10.1016/0016-7037(94)90378-6), 1994.
- 855 Boudreau, B. P.: *Diagenetic Models and Their Implementation: Modelling Transport and Reactions in Aquatic Sediments*, Springer Berlin Heidelberg, Berlin, Heidelberg, 1997.
- Boudreau, B. P. and Ruddick, B. R.: On a reactive continuum representation of organic matter diagenesis, *Am. J. Sci.*, 291, 507–538, <https://doi.org/10.2475/ajs.291.5.507>, 1991.
- Boudreau, B. P., Mucci, A., Sundby, B., Luther, G. W., and Silverberg, N.: Comparative diagenesis at three sites on the Canadian continental margin, *J. Mar. Res.*, 56, 1259–1284, <https://doi.org/10.1357/002224098765093634>, 1998.

- 860 Bourgeois, S., Archambault, P., and Witte, U.: Organic matter remineralization in marine sediments: A Pan-Arctic synthesis: Pan-Arctic Benthic Remineralization, *Glob. Biogeochem. Cycles*, 31, 190–213, <https://doi.org/10.1002/2016GB005378>, 2017.
- Bowles, M. W., Mogollon, J. M., Kasten, S., Zabel, M., and Hinrichs, K.-U.: Global rates of marine sulfate reduction and implications for sub-sea-floor metabolic activities, *Science*, 344, 889–891, <https://doi.org/10.1126/science.1249213>, 2014.
- 865 Bradley, J. A., Arndt, S., Amend, J. P., Burwicz, E., Dale, A. W., Egger, M., and LaRowe, D. E.: Widespread energy limitation to life in global subseafloor sediments, *Sci. Adv.*, 6, eaba0697, <https://doi.org/10.1126/sciadv.aba0697>, 2020.
- Burdige, D. J.: *Geochemistry of marine sediments*, Princeton University Press, Princeton, NJ, 609 pp., 2006.
- Burdige, D. J.: Preservation of Organic Matter in Marine Sediments: Controls, Mechanisms, and an Imbalance in Sediment Organic Carbon Budgets?, *Chem. Rev.*, 107, 467–485, <https://doi.org/10.1021/cr050347q>, 2007.
- 870 Canfield, D. E.: Factors influencing organic carbon preservation in marine sediments, *Chem. Geol.*, 114, 315–329, [https://doi.org/10.1016/0009-2541\(94\)90061-2](https://doi.org/10.1016/0009-2541(94)90061-2), 1994.
- Canfield, D. E., Jørgensen, B. B., Fossing, H., Glud, R., Gundersen, J., Ramsing, N. B., Thamdrup, B., Hansen, J. W., Nielsen, L. P., and Hall, P. O. J.: Pathways of organic carbon oxidation in three continental margin sediments, *Mar. Geol.*, 113, 27–40, [https://doi.org/10.1016/0025-3227\(93\)90147-N](https://doi.org/10.1016/0025-3227(93)90147-N), 1993.
- 875 Cathalot, C., Rabouille, C., Pastor, L., Deflandre, B., Viollier, E., Buscail, R., Grémare, A., Treignier, C., and Pruski, A.: Temporal variability of carbon recycling in coastal sediments influenced by rivers: assessing the impact of flood inputs in the Rhône River prodelta, *Biogeosciences*, 7, 1187–1205, <https://doi.org/10.5194/bg-7-1187-2010>, 2010.
- Chen, D. and Chen, H. W.: Using the Köppen classification to quantify climate variation and change: An example for 1901–2010, *Environ. Dev.*, 6, 69–79, <https://doi.org/10.1016/j.envdev.2013.03.007>, 2013.
- 880 Contreras, S., Meister, P., Liu, B., Prieto-Mollar, X., Hinrichs, K.-U., Khalili, A., Ferdelman, T. G., Kuypers, M. M. M., and Jørgensen, B. B.: Cyclic 100-ka (glacial-interglacial) migration of subseafloor redox zonation on the Peruvian shelf, *Proc. Natl. Acad. Sci.*, 110, 18098–18103, <https://doi.org/10.1073/pnas.1305981110>, 2013.
- Cowie, G.: The biogeochemistry of Arabian Sea surficial sediments: A review of recent studies, *Prog. Oceanogr.*, 65, 260–289, <https://doi.org/10.1016/j.pocean.2005.03.003>, 2005.
- 885 Coyle, K. O., Pinchuk, A. I., Eisner, L. B., and Napp, J. M.: Zooplankton species composition, abundance and biomass on the eastern Bering Sea shelf during summer: The potential role of water-column stability and nutrients in structuring the zooplankton community, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 55, 1775–1791, <https://doi.org/10.1016/j.dsr2.2008.04.029>, 2008.
- Cram, J. A., Weber, T., Leung, S. W., McDonnell, A. M. P., Liang, J.-H., and Deutsch, C.: The Role of Particle Size, Ballast, Temperature, and Oxygen in the Sinking Flux to the Deep Sea, *Glob. Biogeochem. Cycles*, 32, 858–876, <https://doi.org/10.1029/2017GB005710>, 2018.
- 890 Dai, J., Sun, M.-Y., Culp, R. A., and Noakes, J. E.: A laboratory study on biochemical degradation and microbial utilization of organic matter comprising a marine diatom, land grass, and salt marsh plant in estuarine ecosystems, *Aquat. Ecol.*, 43, 825–841, <https://doi.org/10.1007/s10452-008-9211-x>, 2009.

- 895 Dale, A. W., Regnier, P., Knab, N. J., Jørgensen, B. B., and Van Cappellen, P.: Anaerobic oxidation of methane (AOM) in marine sediments from the Skagerrak (Denmark): II. Reaction-transport modeling, *Geochim. Cosmochim. Acta*, 72, 2880–2894, <https://doi.org/10.1016/j.gca.2007.11.039>, 2008a.
- Dale, A. W., Aguilera, D. R., Regnier, P., Fossing, H., Knab, N. J., and Jørgensen, B. B.: Seasonal dynamics of the depth and rate of anaerobic oxidation of methane in Aarhus Bay (Denmark) sediments, *J. Mar. Res.*, 66, 127–155,  
900 <https://doi.org/10.1357/002224008784815775>, 2008b.
- Dale, A. W., Brüchert, V., Alperin, M., and Regnier, P.: An integrated sulfur isotope model for Namibian shelf sediments, *Geochim. Cosmochim. Acta*, 73, 1924–1944, <https://doi.org/10.1016/j.gca.2008.12.015>, 2009.
- Dale, A. W., Flury, S., Fossing, H., Regnier, P., Røy, H., Scholze, C., and Jørgensen, B. B.: Kinetics of organic carbon mineralization and methane formation in marine sediments (Aarhus Bay, Denmark), *Geochim. Cosmochim. Acta*, 252, 159–  
905 178, <https://doi.org/10.1016/j.gca.2019.02.033>, 2019.
- Dauwe, B., Middelburg, J. J., Van Rijswijk, P., Sinke, J., Herman, P. M. J., and Heip, C. H. R.: Enzymatically hydrolyzable amino acids in North Sea sediments and their possible implication for sediment nutritional values, *J. Mar. Res.*, 57, 109–134, <https://doi.org/10.1357/002224099765038580>, 1999a.
- Dauwe, B., Middelburg, J. J., Herman, P. M. J., and Heip, C. H. R.: Linking diagenetic alteration of amino acids and bulk  
910 organic matter reactivity, *Limnol. Oceanogr.*, 44, 1809–1814, <https://doi.org/10.4319/lo.1999.44.7.1809>, 1999b.
- Dyer, K. R.: Sedimentation processes in the Bristol Channel/Severn Estuary, *Mar. Pollut. Bull.*, 15, 53–57, [https://doi.org/10.1016/0025-326X\(84\)90462-4](https://doi.org/10.1016/0025-326X(84)90462-4), 1984.
- Egger, M., Riedinger, N., Mogollón, J. M., and Jørgensen, B. B.: Global diffusive fluxes of methane in marine sediments, *Nat. Geosci.*, 11, 421–425, <https://doi.org/10.1038/s41561-018-0122-8>, 2018.
- 915 Emerson, S. and Hedges, J. I.: Processes controlling the organic carbon content of open ocean sediments, *Paleoceanography*, 3, 621–634, <https://doi.org/10.1029/PA003i005p00621>, 1988.
- Faust, J. C., Tessin, A., Fisher, B. J., Zindorf, M., Papadaki, S., Hendry, K. R., Doyle, K. A., and März, C.: Millennial scale persistence of organic carbon bound to iron in Arctic marine sediments, *Nat. Commun.*, 12, 275, <https://doi.org/10.1038/s41467-020-20550-0>, 2021.
- 920 Fleming-Lehtinen, V. and Laamanen, M.: Long-term changes in Secchi depth and the role of phytoplankton in explaining light attenuation in the Baltic Sea, *Estuar. Coast. Shelf Sci.*, 102–103, 1–10, <https://doi.org/10.1016/j.ecss.2012.02.015>, 2012.
- Flury, S., Røy, H., Dale, A. W., Fossing, H., Tóth, Z., Spiess, V., Jensen, J. B., and Jørgensen, B. B.: Controls on subsurface methane fluxes and shallow gas formation in Baltic Sea sediment (Aarhus Bay, Denmark), *Geochim. Cosmochim. Acta*, 188,  
925 297–309, <https://doi.org/10.1016/j.gca.2016.05.037>, 2016.
- Forney, D. C. and Rothman, D. H.: Inverse method for estimating respiration rates from decay time series, *Biogeosciences*, 9, 3601–3612, <https://doi.org/10.5194/bg-9-3601-2012>, 2012.
- Freitas, F. S., Hendry, K. R., Henley, S. F., Faust, J. C., Tessin, A. C., Stevenson, M. A., Abbott, G. D., März, C., and Arndt, S.: Benthic-pelagic coupling in the Barents Sea: an integrated data-model framework, *Phil. Trans. R. Soc. A.*, 378, 1–15,  
930 <http://dx.doi.org/10.1098/rsta.2019.0359>, 2020.

- Gersonde, R.: The expedition of the research vessel “Sonne” to the subpolar North Pacific and the Bering Sea in 2009 (SO202-INOPEX) (Reports on polar and marine research No. 643), Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, 2009.
- 935 Glud, R. N.: Oxygen dynamics of marine sediments, *Mar. Biol. Res.*, 4, 243–289, <https://doi.org/10.1080/17451000801888726>, 2008.
- Griffith, D. R., Martin, W. R., and Eglinton, T. I.: The radiocarbon age of organic carbon in marine surface sediments, *Geochim. Cosmochim. Acta*, 74, 6788–6800, <https://doi.org/10.1016/j.gca.2010.09.001>, 2010.
- Grossi, V., Caradec, S., and Gilbert, F.: Burial and reactivity of sedimentary microalgal lipids in bioturbated Mediterranean coastal sediments, *Mar. Chem.*, 81, 57–69, [https://doi.org/10.1016/S0304-4203\(02\)00139-1](https://doi.org/10.1016/S0304-4203(02)00139-1), 2003.
- 940 Grotheer, H., Meyer, V., Riedel, T., Pfalz, G., Mathieu, L., Hefter, J., Gentz, T., Lantuit, H., Mollenhauer, G., and Fritz, M.: Burial and Origin of Permafrost-Derived Carbon in the Nearshore Zone of the Southern Canadian Beaufort Sea, *Geophys. Res. Lett.*, 47, e2019GL085897, <https://doi.org/10.1029/2019GL085897>, 2020.
- Guenet, B., Danger, M., Abbadie, L., and Lacroix, G.: Priming effect: bridging the gap between terrestrial and aquatic ecology, *Ecology*, 91, 2850–2861, <https://doi.org/10.1890/09-1968.1>, 2010.
- 945 de Haas, H. and van Weering, T. C. E.: Recent sediment accumulation, organic carbon burial and transport in the northeastern North Sea, *Mar. Geol.*, 136, 173–187, [https://doi.org/10.1016/S0025-3227\(96\)00072-2](https://doi.org/10.1016/S0025-3227(96)00072-2), 1997.
- Hartnett, H. E., Keil, R. G., Hedges, J. I., and Devol, A. H.: Influence of oxygen exposure time on organic carbon preservation in continental margin sediments, *Nature*, 391, 572–575, <https://doi.org/10.1038/35351>, 1998.
- Hedges, J. I.: Global biogeochemical cycles: progress and problems, *Mar. Chem.*, 39, 67–93, [https://doi.org/10.1016/0304-4203\(92\)90096-S](https://doi.org/10.1016/0304-4203(92)90096-S), 1992.
- 950 Hedges, J. I. and Oades, J. M.: Comparative organic geochemistries of soils and marine sediments, *Org. Geochem.*, 27, 319–361, [https://doi.org/10.1016/S0146-6380\(97\)00056-9](https://doi.org/10.1016/S0146-6380(97)00056-9), 1997.
- Hedges, J. I., Keil, R. G., and Benner, R.: What happens to terrestrial organic matter in the ocean?, *Org. Geochem.*, 27, 195–212, [https://doi.org/10.1016/S0146-6380\(97\)00066-1](https://doi.org/10.1016/S0146-6380(97)00066-1), 1997.
- 955 Heggie, D., Maris, C., Hudson, A., Dymond, J., Beach, R., and Cullen, J.: Organic carbon oxidation and preservation in NW Atlantic continental margin sediments, *Geol. Soc. Lond. Spec. Publ.*, 31, 215, <https://doi.org/10.1144/GSL.SP.1987.031.01.15>, 1987.
- Hemingway, J. D., Rothman, D. H., Grant, K. E., Rosengard, S. Z., Eglinton, T. I., Derry, L. A., and Galy, V. V.: Mineral protection regulates long-term global preservation of natural organic carbon, *Nature*, 570, 228–231, <https://doi.org/10.1038/s41586-019-1280-6>, 2019.
- 960 Henkel, S. and Kulkarni, A.: Documentation of sediment core HE443/10-3, <https://doi.org/10.1594/PANGAEA.855884>, 2015.
- Henkel, S., Strasser, M., Schwenk, T., Hanebuth, T. J. J., Hüsener, J., Arnold, G. L., Winkelmann, D., Formolo, M., Tomasini, J., Krastel, S., and Kasten, S.: An interdisciplinary investigation of a recent submarine mass transport deposit at the continental margin off Uruguay, *Geochem. Geophys. Geosystems*, 12, 1–19, <https://doi.org/10.1029/2011GC003669>, 2011.

- Henkel, S., Mogollón, J. M., Nöthen, K., Franke, C., Bogus, K., Robin, E., Bahr, A., Blumenberg, M., Pape, T., Seifert, R., März, C., de Lange, G. J., and Kasten, S.: Diagenetic barium cycling in Black Sea sediments – A case study for anoxic marine environments, *Geochim. Cosmochim. Acta*, 88, 88–105, <https://doi.org/10.1016/j.gca.2012.04.021>, 2012.
- 970 Henrichs, S. M.: Early diagenesis of organic matter in marine sediments: progress and perplexity, *Mar. Chem.*, 39, 119–149, [https://doi.org/10.1016/0304-4203\(92\)90098-U](https://doi.org/10.1016/0304-4203(92)90098-U), 1992.
- Hensen, C., Zabel, M., Pfeifer, K., Schwenk, T., Kasten, S., Riedinger, N., Schulz, H. D., and Boetius, A.: Control of sulfate pore-water profiles by sedimentary events and the significance of anaerobic oxidation of methane for the burial of sulfur in marine sediments, *Geochim. Cosmochim. Acta*, 67, 2631–2647, [https://doi.org/10.1016/S0016-7037\(03\)00199-6](https://doi.org/10.1016/S0016-7037(03)00199-6), 2003.
- 975 Hensen, C., Zabel, M., and Schulz, H. N.: Benthic Cycling of Oxygen, Nitrogen and Phosphorus, in: *Marine Geochemistry*, edited by: Schulz, H. D. and Zabel, M., Springer-Verlag, Berlin/Heidelberg, 207–240, [https://doi.org/10.1007/3-540-32144-6\\_6](https://doi.org/10.1007/3-540-32144-6_6), 2006.
- Hilligsøe, K. M., Jensen, J. B., Ferdelman, T. G., Fossing, H., Lapham, L., Røy, H., and Jørgensen, B. B.: Methane fluxes in marine sediments quantified through core analyses and seismo-acoustic mapping (Bornholm Basin, Baltic Sea), *Geochim. Cosmochim. Acta*, 239, 255–274, <https://doi.org/10.1016/j.gca.2018.07.040>, 2018.
- 980 Ho, T. C. and Aris, R.: On apparent second-order kinetics, *AIChE J.*, 33, 1050–1051, <https://doi.org/10.1002/aic.690330621>, 1987.
- Hoefs, M. J. L., Rijpstra, W. I. C., and Sinninghe Damsté, J. S.: The influence of oxic degradation on the sedimentary biomarker record I: evidence from Madeira Abyssal Plain turbidites, *Geochim. Cosmochim. Acta*, 66, 2719–2735, [https://doi.org/10.1016/S0016-7037\(02\)00864-5](https://doi.org/10.1016/S0016-7037(02)00864-5), 2002.
- 985 Huguet, C., de Lange, G. J., Gustafsson, Ö., Middelburg, J. J., Sinninghe Damsté, J. S., and Schouten, S.: Selective preservation of soil organic matter in oxidized marine sediments (Madeira Abyssal Plain), *Geochim. Cosmochim. Acta*, 72, 6061–6068, <https://doi.org/10.1016/j.gca.2008.09.021>, 2008.
- Hülse, D., Arndt, S., Daines, S., Regnier, P., and Ridgwell, A.: OMEN-SED 0.9: A novel, numerically efficient organic mattersediment diagenesis module for coupling to Earth system models, *Climate and Earth System Modeling*, <https://doi.org/10.5194/gmd-2017-296>, 2018.
- 990 Hülse, D., Arndt, S., and Ridgwell, A.: Mitigation of Extreme Ocean Anoxic Event Conditions by Organic Matter Sulfurization, *Paleoceanogr. Paleoclimatology*, 34, 476–489, <https://doi.org/10.1029/2018PA003470>, 2019.
- Jensen, M., Lomstein, E., and Sørensen, J.: Benthic NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> flux following sedimentation of a spring phytoplankton bloom in Aarhus Bight, Denmark, *Mar. Ecol. Prog. Ser.*, 61, 87–96, <https://doi.org/10.3354/meps061087>, 1990.
- 995 Jonas, P. J. C. and Millward, G. E.: Metals and nutrients in the Severn Estuary and Bristol Channel: Contemporary inputs and distributions, *Mar. Pollut. Bull.*, 61, 52–67, <https://doi.org/10.1016/j.marpolbul.2009.12.013>, 2010.
- Jørgensen, B. B. and Kasten, S.: Sulfur Cycling and Methane Oxidation, in: *Marine Geochemistry*, edited by: Schulz, H. D. and Zabel, M., Springer-Verlag, Berlin/Heidelberg, 271–309, [https://doi.org/10.1007/3-540-32144-6\\_8](https://doi.org/10.1007/3-540-32144-6_8), 2006.
- 1000 Jørgensen, B. B., Beulig, F., Egger, M., Petro, C., Scholze, C., and Røy, H.: Organoclastic sulfate reduction in the sulfate-methane transition of marine sediments, *Geochim. Cosmochim. Acta*, 254, 231–245, <https://doi.org/10.1016/j.gca.2019.03.016>, 2019a.

- Jørgensen, B. B., Findlay, A. J., and Pellerin, A.: The Biogeochemical Sulfur Cycle of Marine Sediments, *Front. Microbiol.*, 10, 849, <https://doi.org/10.3389/fmicb.2019.00849>, 2019b.
- 1005 Jourabchi, P.: Quantitative interpretation of pH distributions in aquatic sediments: A reaction-transport modeling approach, *Am. J. Sci.*, 305, 919–956, <https://doi.org/10.2475/ajs.305.9.919>, 2005.
- Karlson, B., Edler, L., Granéli, W., Sahlsten, E., and Kuylenstierna, M.: Subsurface chlorophyll maxima in the Skagerrak-processes and plankton community structure, *J. Sea Res.*, 35, 139–158, [https://doi.org/10.1016/S1385-1101\(96\)90742-X](https://doi.org/10.1016/S1385-1101(96)90742-X), 1996.
- 1010 Kayler, Z. E., Premke, K., Gessler, A., Gessner, M. O., Griebler, C., Hilt, S., Klemmedtsson, L., Kuzyakov, Y., Reichstein, M., Siemens, J., Totsche, K.-U., Tranvik, L., Wagner, A., Weitere, M., and Grossart, H.-P.: Integrating Aquatic and Terrestrial Perspectives to Improve Insights Into Organic Matter Cycling at the Landscape Scale, *Front. Earth Sci.*, 7, 127, <https://doi.org/10.3389/feart.2019.00127>, 2019.
- 1015 Keil, R. G. and Cowie, G. L.: Organic matter preservation through the oxygen-deficient zone of the NE Arabian Sea as discerned by organic carbon:mineral surface area ratios, *Mar. Geol.*, 161, 13–22, [https://doi.org/10.1016/S0025-3227\(99\)00052-3](https://doi.org/10.1016/S0025-3227(99)00052-3), 1999.
- Kharbush, J. J., Close, H. G., Van Mooy, B. A. S., Arnosti, C., Smittenberg, R. H., Le Moigne, F. A. C., Mollenhauer, G., Scholz-Böttcher, B., Obrecht, I., Koch, B. P., Becker, K. W., Iversen, M. H., and Mohr, W.: Particulate Organic Carbon Deconstructed: Molecular and Chemical Composition of Particulate Organic Carbon in the Ocean, *Front. Mar. Sci.*, 7, 518, <https://doi.org/10.3389/fmars.2020.00518>, 2020.
- 1020 Knab, N. J., Cragg, B. A., Borowski, C., Parkes, R. J., Pancost, R., and Jørgensen, B. B.: Anaerobic oxidation of methane (AOM) in marine sediments from the Skagerrak (Denmark): I. Geochemical and microbiological analyses, *Geochim. Cosmochim. Acta*, 72, 2868–2879, <https://doi.org/10.1016/j.gca.2008.03.016>, 2008.
- Koho, K. A., Nierop, K. G. J., Moodley, L., Middelburg, J. J., Pozzato, L., Soetaert, K., van der Plicht, J., and Reichart, G.-J.: Microbial bioavailability regulates organic matter preservation in marine sediments, *Biogeosciences*, 10, 1131–1141, <https://doi.org/10.5194/bg-10-1131-2013>, 2013.
- 1025 Krumins, V., Gehlen, M., Arndt, S., Van Cappellen, P., and Regnier, P.: Dissolved inorganic carbon and alkalinity fluxes from coastal marine sediments: model estimates for different shelf environments and sensitivity to global change, *Biogeosciences*, 10, 371–398, <https://doi.org/10.5194/bg-10-371-2013>, 2013.
- 1030 Kusch, S., Rethemeyer, J., Ransby, D., and Mollenhauer, G.: Permafrost Organic Carbon Turnover and Export Into a High-Arctic Fjord: A Case Study From Svalbard Using Compound-specific <sup>14</sup>C Analysis, *J. Geophys. Res. Biogeosciences*, 126, e2020JG006008, <https://doi.org/10.1029/2020JG006008>, 2021.
- LaRowe, D. E., Arndt, S., Bradley, J. A., Burwicz, E., Dale, A. W., and Amend, J. P.: Organic carbon and microbial activity in marine sediments on a global scale throughout the Quaternary, *Geochim. Cosmochim. Acta*, 286, 227–247, <https://doi.org/10.1016/j.gca.2020.07.017>, 2020a.
- 1035 LaRowe, D. E., Arndt, S., Bradley, J. A., Estes, E. R., Hoarfrost, A., Lang, S. Q., Lloyd, K. G., Mahmoudi, N., Orsi, W. D., Shah Walter, S. R., Steen, A. D., and Zhao, R.: The fate of organic carbon in marine sediments - New insights from recent data and analysis, *Earth-Sci. Rev.*, 204, 103146, <https://doi.org/10.1016/j.earscirev.2020.103146>, 2020b.

- 1040 Latasa, M. and Bidigare, R. R.: A comparison of phytoplankton populations of the Arabian Sea during the Spring Intermonsoon and Southwest Monsoon of 1995 as described by HPLC-analyzed pigments, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 45, 2133–2170, [https://doi.org/10.1016/S0967-0645\(98\)00066-6](https://doi.org/10.1016/S0967-0645(98)00066-6), 1998.
- 1045 Lengger, S. K., Rush, D., Mayser, J. P., Blewett, J., Schwartz-Narbonne, R., Talbot, H. M., Middelburg, J. J., Jetten, M. S. M., Schouten, S., Sinninghe Damsté, J. S., and Pancost, R. D.: Dark carbon fixation in the Arabian Sea oxygen minimum zone contributes to sedimentary organic carbon (SOM), *Glob. Biogeochem. Cycles*, 2019GB006282, <https://doi.org/10.1029/2019GB006282>, 2019.
- Lessin, G., Artioli, Y., Almroth-Rosell, E., Blackford, J. C., Dale, A. W., Glud, R. N., Middelburg, J. J., Pastres, R., Queirós, A. M., Rabouille, C., Regnier, P., Soetaert, K., Solidoro, C., Stephens, N., and Yakushev, E.: Modelling Marine Sediment Biogeochemistry: Current Knowledge Gaps, Challenges, and Some Methodological Advice for Advancement, *Front. Mar. Sci.*, 5, 19, <https://doi.org/10.3389/fmars.2018.00019>, 2018.
- 1050 Lohse, L., Malschaert, J. F. P., Slomp, C. P., Helder, W., and van Raaphorst, W.: Sediment-water fluxes of inorganic nitrogen compounds along the transport route of organic matter in the North Sea, *Ophelia*, 41, 173–197, <https://doi.org/10.1080/00785236.1995.10422043>, 1995.
- Lomstein, E., Jensen, M. H., and Sørensen, J.: Intracellular NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> pools associated with deposited phytoplankton in a marine sediment (Aarhus Bight, Denmark), *Mar. Ecol. Prog. Ser.*, 61, 97–105, 1990.
- 1055 Luff, R., Wallmann, K., Grandel, S., and Schlüter, M.: Numerical modeling of benthic processes in the deep Arabian Sea, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 47, 3039–3072, [https://doi.org/10.1016/S0967-0645\(00\)00058-8](https://doi.org/10.1016/S0967-0645(00)00058-8), 2000.
- Mackin, J. E. and Swider, K. T.: Organic matter decomposition pathways and oxygen consumption in coastal marine sediments, *J. Mar. Res.*, 47, 681–716, <https://doi.org/10.1357/002224089785076154>, 1989.
- 1060 Maerz, J., Six, K. D., Stemmler, I., Ahmerkamp, S., and Ilyina, T.: Microstructure and composition of marine aggregates as co-determinants for vertical particulate organic carbon transfer in the global ocean, *Earth System Science/Response to Global Change: Climate Change*, <https://doi.org/10.5194/bg-2019-378>, 2019.
- Manning, A. J., Langston, W. J., and Jonas, P. J. C.: A review of sediment dynamics in the Severn Estuary: Influence of flocculation, *Mar. Pollut. Bull.*, 61, 37–51, <https://doi.org/10.1016/j.marpolbul.2009.12.012>, 2010.
- 1065 Marquardt, M., Hensen, C., Piñero, E., Wallmann, K., and Haeckel, M.: A transfer function for the prediction of gas hydrate inventories in marine sediments, *Biogeosciences*, 7, 2925–2941, <https://doi.org/10.5194/bg-7-2925-2010>, 2010.
- Mayer, L. M.: Relationships between mineral surfaces and organic carbon concentrations in soils and sediments, *Chem. Geol.*, 114, 347–363, [https://doi.org/10.1016/0009-2541\(94\)90063-9](https://doi.org/10.1016/0009-2541(94)90063-9), 1994a.
- Mayer, L. M.: Surface area control of organic carbon accumulation in continental shelf sediments, *Geochim. Cosmochim. Acta*, 58, 1271–1284, [https://doi.org/10.1016/0016-7037\(94\)90381-6](https://doi.org/10.1016/0016-7037(94)90381-6), 1994b.
- 1070 Mayer, L. M.: Sedimentary organic matter preservation: an assessment and speculative synthesis—a comment, *Mar. Chem.*, 49, 123–126, [https://doi.org/10.1016/0304-4203\(95\)00011-F](https://doi.org/10.1016/0304-4203(95)00011-F), 1995.
- Meile, C. and Van Cappellen, P.: Particle age distributions and O<sub>2</sub> exposure times: Timescales in bioturbated sediments, *Glob. Biogeochem. Cycles*, 19, 1–12, <https://doi.org/10.1029/2004GB002371>, 2005.

- 1075 Meister, P., Liu, B., Ferdelman, T. G., Jørgensen, B. B., and Khalili, A.: Control of sulphate and methane distributions in marine sediments by organic matter reactivity, *Geochim. Cosmochim. Acta*, 104, 183–193, <https://doi.org/10.1016/j.gca.2012.11.011>, 2013.
- Mewes, K., Mogollón, J. M., Picard, A., Rühlemann, C., Eisenhauer, A., Kuhn, T., Ziebis, W., and Kasten, S.: Diffusive transfer of oxygen from seamount basaltic crust into overlying sediments: An example from the Clarion–Clipperton Fracture Zone, *Earth Planet. Sci. Lett.*, 433, 215–225, <https://doi.org/10.1016/j.epsl.2015.10.028>, 2016.
- 1080 Meyers, P. A.: Organic geochemical proxies of paleoceanographic, paleolimnologic, and paleoclimatic processes, *Org. Geochem.*, 27, 213–250, [https://doi.org/10.1016/S0146-6380\(97\)00049-1](https://doi.org/10.1016/S0146-6380(97)00049-1), 1997.
- Middelburg, J. J.: A simple rate model for organic matter decomposition in marine sediments, *Geochim. Cosmochim. Acta*, 53, 1577–1581, [https://doi.org/10.1016/0016-7037\(89\)90239-1](https://doi.org/10.1016/0016-7037(89)90239-1), 1989.
- 1085 Middelburg, J. J.: Reviews and syntheses: to the bottom of carbon processing at the seafloor, *Biogeosciences*, 15, 413–427, <https://doi.org/10.5194/bg-15-413-2018>, 2018.
- Middelburg, J. J.: *Marine Carbon Biogeochemistry: A Primer for Earth System Scientists*, Springer International Publishing, Cham, <https://doi.org/10.1007/978-3-030-10822-9>, 2019.
- Middelburg, J. J., Vlug, T., Jaco, F., and van der Nat, W. A.: Organic matter mineralization in marine systems, *Glob. Planet. Change*, 8, 47–58, [https://doi.org/10.1016/0921-8181\(93\)90062-S](https://doi.org/10.1016/0921-8181(93)90062-S), 1993.
- 1090 Middelburg, J. J., Soetaert, K., and Herman, P. M. J.: Empirical relationships for use in global diagenetic models, *Deep Sea Res. Part Oceanogr. Res. Pap.*, 44, 327–344, [https://doi.org/10.1016/S0967-0637\(96\)00101-X](https://doi.org/10.1016/S0967-0637(96)00101-X), 1997.
- Mogollón, J. M., Dale, A. W., Fossing, H., and Regnier, P.: Timescales for the development of methanogenesis and free gas layers in recently-deposited sediments of Arkona Basin (Baltic Sea), *Biogeosciences*, 9, 1915–1933, <https://doi.org/10.5194/bg-9-1915-2012>, 2012.
- 1095 Mogollón, J. M., Mewes, K., and Kasten, S.: Quantifying manganese and nitrogen cycle coupling in manganese-rich, organic carbon-starved marine sediments: Examples from the Clarion-Clipperton fracture zone, *Geophys. Res. Lett.*, 43, 7114–7123, <https://doi.org/10.1002/2016GL069117>, 2016.
- 1100 Mollenhauer, G., Eglinton, T. I., Ohkouchi, N., Schneider, R. R., Müller, P. J., Grootes, P. M., and Rullkötter, J.: Asynchronous alkenone and foraminifera records from the Benguela Upwelling System, *Geochim. Cosmochim. Acta*, 67, 2157–2171, [https://doi.org/10.1016/S0016-7037\(03\)00168-6](https://doi.org/10.1016/S0016-7037(03)00168-6), 2003.
- Mollenhauer, G., Inthorn, M., Vogt, T., Zabel, M., Sinninghe Damsté, J. S., and Eglinton, T. I.: Aging of marine organic matter during cross-shelf lateral transport in the Benguela upwelling system revealed by compound-specific radiocarbon dating, *Geochem. Geophys. Geosystems*, 8, 1–16, <https://doi.org/10.1029/2007GC001603>, 2007.
- 1105 Müller, P. J. and Suess, E.: Productivity, sedimentation rate, and sedimentary organic matter in the oceans—I. Organic carbon preservation, *Deep Sea Res. Part Oceanogr. Res. Pap.*, 26, 1347–1362, [https://doi.org/10.1016/0198-0149\(79\)90003-7](https://doi.org/10.1016/0198-0149(79)90003-7), 1979.
- Niewöhner, C., Hensen, C., Kasten, S., Zabel, M., and Schulz, H. D.: Deep Sulfate Reduction Completely Mediated by Anaerobic Methane Oxidation in Sediments of the Upwelling Area off Namibia, *Geochim. Cosmochim. Acta*, 62, 455–464, [https://doi.org/10.1016/S0016-7037\(98\)00055-6](https://doi.org/10.1016/S0016-7037(98)00055-6), 1998.

- 1110 van Nugteren, P., Moodley, L., Brummer, G.-J., Heip, C. H. R., Herman, P. M. J., and Middelburg, J. J.: Seafloor ecosystem functioning: the importance of organic matter priming, *Mar. Biol.*, 156, 2277–2287, <https://doi.org/10.1007/s00227-009-1255-5>, 2009.
- Ohkouchi, N., Eglinton, T. I., Keigwin, L. D., and Hayes, J. M.: Spatial and Temporal Offsets Between Proxy Records in a Sediment Drift, *Science*, 298, 1224, <https://doi.org/10.1126/science.1075287>, 2002.
- 1115 Oni, O., Miyatake, T., Kasten, S., Richter-Heitmann, T., Fischer, D., Wagenknecht, L., Kulkarni, A., Blumers, M., Shylin, S. I., Ksenofontov, V., Costa, B. F. O., Klingelhöfer, G., and Friedrich, M. W.: Distinct microbial populations are tightly linked to the profile of dissolved iron in the methanic sediments of the Helgoland mud area, North Sea, *Front. Microbiol.*, 06, <https://doi.org/10.3389/fmicb.2015.00365>, 2015a.
- 1120 Oni, O. E., Schmidt, F., Miyatake, T., Kasten, S., Witt, M., Hinrichs, K.-U., and Friedrich, M. W.: Microbial Communities and Organic Matter Composition in Surface and Subsurface Sediments of the Helgoland Mud Area, North Sea, *Front. Microbiol.*, 6, <https://doi.org/10.3389/fmicb.2015.01290>, 2015b.
- Owings, S. M., Bréthous, L., Eitel, E. M., Fields, B. P., Boever, A., Beckler, J. S., Bombled, B., Lansard, B., Metzger, E., Rabouille, C., and Taillefert, M.: Differential manganese and iron recycling and transport in continental margin sediments of the Northern Gulf of Mexico, *Mar. Chem.*, 229, 103908, <https://doi.org/10.1016/j.marchem.2020.103908>, 2021.
- 1125 Pastor, L., Deflandre, B., Viollier, E., Cathalot, C., Metzger, E., Rabouille, C., Escoubeyrou, K., Lloret, E., Pruski, A. M., Vétion, G., Desmalades, M., Buscail, R., and Grémare, A.: Influence of the organic matter composition on benthic oxygen demand in the Rhône River prodelta (NW Mediterranean Sea), *Cont. Shelf Res.*, 31, 1008–1019, <https://doi.org/10.1016/j.csr.2011.03.007>, 2011.
- 1130 Premuzic, E. T., Benkovitz, C. M., Gaffney, J. S., and Walsh, J. J.: The nature and distribution of organic matter in the surface sediments of world oceans and seas, *Org. Geochem.*, 4, 63–77, [https://doi.org/10.1016/0146-6380\(82\)90009-2](https://doi.org/10.1016/0146-6380(82)90009-2), 1982.
- Pruski, A. M., Buscail, R., Bourgeois, S., Vétion, G., Coston-Guarini, J., and Rabouille, C.: Biogeochemistry of fatty acids in a river-dominated Mediterranean ecosystem (Rhône River prodelta, Gulf of Lions, France): Origins and diagenesis, *Org. Geochem.*, 83–84, 227–240, <https://doi.org/10.1016/j.orggeochem.2015.04.002>, 2015.
- 1135 Puglini, M., Brovkin, V., Regnier, P., and Arndt, S.: Assessing the potential for non-turbulent methane escape from the East Siberian Arctic Shelf, *Biogeosciences*, 17, 3247–3275, <https://doi.org/10.5194/bg-17-3247-2020>, 2020.
- Ragavan, P. and Kumar, S.: Potential role of priming effect in the open ocean oxygen minimum zones: an outlook, *Hydrobiologia*, <https://doi.org/10.1007/s10750-021-04574-1>, 2021.
- 1140 Rassmann, J., Lansard, B., Pozzato, L., and Rabouille, C.: Carbonate chemistry in sediment porewaters of the Rhône River deltadriven by early diagenesis (northwestern Mediterranean), *Biogeosciences*, 13, 5379–5394, <https://doi.org/10.5194/bg-13-5379-2016>, 2016.
- Rassmann, J., Eitel, E. M., Lansard, B., Cathalot, C., Brandily, C., Taillefert, M., and Rabouille, C.: Benthic alkalinity and dissolved inorganic carbon fluxes in the Rhône River prodelta generated by decoupled aerobic and anaerobic processes, *Biogeosciences*, 17, 13–33, <https://doi.org/10.5194/bg-17-13-2020>, 2020.
- 1145 Regnier, P., Mouchet, A., Wollast, R., and Runday, F.: A discussion of methods for estimating residual fluxes in strong tidal estuaries, *Cont. Shelf Res.*, 18, 1543–1571, [https://doi.org/10.1016/S0278-4343\(98\)00071-5](https://doi.org/10.1016/S0278-4343(98)00071-5), 1998.

- Regnier, P., O’Kane, J. P., Steefel, C. I., and Vanderborght, J. P.: Modeling complex multi-component reactive-transport systems: towards a simulation environment based on the concept of a Knowledge Base, *Appl. Math. Model.*, 26, 913–927, [https://doi.org/10.1016/S0307-904X\(02\)00047-1](https://doi.org/10.1016/S0307-904X(02)00047-1), 2002.
- 1150 Regnier, P., Dale, A. W., Arndt, S., LaRowe, D. E., Mogollón, J., and Van Cappellen, P.: Quantitative analysis of anaerobic oxidation of methane (AOM) in marine sediments: A modeling perspective, *Earth-Sci. Rev.*, 106, 105–130, <https://doi.org/10.1016/j.earscirev.2011.01.002>, 2011.
- Riedinger, N., Pfeifer, K., Kasten, S., Garming, J. F. L., Vogt, C., and Hensen, C.: Diagenetic Alteration of Magnetic Signals by Anaerobic Oxidation of Methane Related to a Change in Sedimentation Rate, *Geochim. Cosmochim. Acta*, 69, 4117–4126, <https://doi.org/10.1016/j.gca.2005.02.004>, 2005.
- 1155 Riedinger, N., Formolo, M. J., Lyons, T. W., Henkel, S., Beck, A., and Kasten, S.: An inorganic geochemical argument for coupled anaerobic oxidation of methane and iron reduction in marine sediments, *Geobiology*, 12, 172–181, <https://doi.org/10.1111/gbi.12077>, 2014.
- 1160 Riedinger, N., Brunner, B., Krastel, S., Arnold, G. L., Wehrmann, L. M., Formolo, M. J., Beck, A., Bates, S. M., Henkel, S., Kasten, S., and Lyons, T. W.: Sulfur Cycling in an Iron Oxide-Dominated, Dynamic Marine Depositional System: The Argentine Continental Margin, *Front. Earth Sci.*, 5, 33, <https://doi.org/10.3389/feart.2017.00033>, 2017.
- Rixen, T., Gaye, B., and Emeis, K.-C.: The monsoon, carbon fluxes, and the organic carbon pump in the northern Indian Ocean, *Prog. Oceanogr.*, 175, 24–39, <https://doi.org/10.1016/j.pcean.2019.03.001>, 2019.
- Rysgaard, S., Fossing, H., and Jensen, M. M.: Organic matter degradation through oxygen respiration, denitrification, and manganese, iron, and sulfate reduction in marine sediments (the Kattegat and the Skagerrak), *Ophelia*, 55, 77–91, <https://doi.org/10.1080/00785236.2001.10409475>, 2001.
- 1165 Salvadó, J. A., Tesi, T., Andersson, A., Ingri, J., Dudarev, O. V., Semiletov, I. P., and Gustafsson, Ö.: Organic carbon remobilized from thawing permafrost is resequenced by reactive iron on the Eurasian Arctic Shelf, *Geophys. Res. Lett.*, 42, 8122–8130, <https://doi.org/10.1002/2015GL066058>, 2015.
- Schlitzer, R.: Ocean data view, 2019.
- 1170 Schmidt, M. W. I., Torn, M. S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I. A., Kleber, M., Kögel-Knabner, I., Lehmann, J., Manning, D. A. C., Nannipieri, P., Rasse, D. P., Weiner, S., and Trumbore, S. E.: Persistence of soil organic matter as an ecosystem property, *Nature*, 478, 49–56, <https://doi.org/10.1038/nature10386>, 2011.
- 1175 Schulz, H. D., Dahmke, A., Schinzel, U., Wallmann, K., and Zabel, M.: Early diagenetic processes, fluxes, and reaction rates in sediments of the South Atlantic, *Geochim. Cosmochim. Acta*, 58, 2041–2060, [https://doi.org/10.1016/0016-7037\(94\)90284-4](https://doi.org/10.1016/0016-7037(94)90284-4), 1994.
- Seiter, K., Hensen, C., Schröter, J., and Zabel, M.: Organic carbon content in surface sediments—defining regional provinces, *Deep Sea Res. Part Oceanogr. Res. Pap.*, 51, 2001–2026, <https://doi.org/10.1016/j.dsr.2004.06.014>, 2004.
- Seiter, K., Hensen, C., and Zabel, M.: Benthic carbon mineralization on a global scale, *Glob. Biogeochem. Cycles*, 19, 1–26, <https://doi.org/10.1029/2004GB002225>, 2005.
- 1180 Shalapyonok, A., Olson, R. J., and Shalapyonok, L. S.: Arabian Sea phytoplankton during Southwest and Northeast Monsoons 1995: composition, size structure and biomass from individual cell properties measured by flow cytometry, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 48, 1231–1261, [https://doi.org/10.1016/S0967-0645\(00\)00137-5](https://doi.org/10.1016/S0967-0645(00)00137-5), 2001.

- 1185 Shields, M. R., Bianchi, T. S., Gélinas, Y., Allison, M. A., and Twilley, R. R.: Enhanced terrestrial carbon preservation promoted by reactive iron in deltaic sediments, *Geophys. Res. Lett.*, 43, 1149–1157, <https://doi.org/10.1002/2015GL067388>, 2016.
- Sinninghe Damsté, J. S., Irene, W., Rijpstra, C., de Leeuw, J. W., and Schenck, P. A.: Origin of organic sulphur compounds and sulphur-containing high molecular weight substances in sediments and immature crude oils, *Org. Geochem.*, 13, 593–606, [https://doi.org/10.1016/0146-6380\(88\)90079-4](https://doi.org/10.1016/0146-6380(88)90079-4), 1988.
- 1190 Sinninghe Damsté, J. S., Kok, M. D., Köster, J., and Schouten, S.: Sulfurized carbohydrates: an important sedimentary sink for organic carbon?, *Earth Planet. Sci. Lett.*, 164, 7–13, [https://doi.org/10.1016/S0012-821X\(98\)00234-9](https://doi.org/10.1016/S0012-821X(98)00234-9), 1998.
- Sinninghe Damsté, J. S., Rijpstra, W. I. C., and Reichart, G.: The influence of oxic degradation on the sedimentary biomarker record II. Evidence from Arabian Sea sediments, *Geochim. Cosmochim. Acta*, 66, 2737–2754, [https://doi.org/10.1016/S0016-7037\(02\)00865-7](https://doi.org/10.1016/S0016-7037(02)00865-7), 2002.
- 1195 Soetaert, K., Herman, P. M. J., and Middelburg, J. J.: A model of early diagenetic processes from the shelf to abyssal depths, *Geochim. Cosmochim. Acta*, 60, 1019–1040, [https://doi.org/10.1016/0016-7037\(96\)00013-0](https://doi.org/10.1016/0016-7037(96)00013-0), 1996.
- Soetaert, K., Herman, P. M. J., Middelburg, J. J., and Heip, C.: Assessing organic matter mineralization, degradability and mixing rate in an ocean margin sediment (Northeast Atlantic) by diagenetic modeling, *J. Mar. Res.*, 56, 519–534, <https://doi.org/10.1357/002224098321822401>, 1998.
- 1200 Stabeno, P. J. and Hunt, G. L.: Overview of the Inner Front and Southeast Bering Sea Carrying Capacity Programs, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 49, 6157–6168, [https://doi.org/10.1016/S0967-0645\(02\)00339-9](https://doi.org/10.1016/S0967-0645(02)00339-9), 2002.
- Stevenson, M. A., Faust, J. C., Andrade, L. L., Freitas, F. S., Gray, N. D., Tait, K., Hendry, K. R., Hilton, R. G., Henley, S. F., Tessin, A., Leary, P., Papadaki, S., Ford, A., März, C., and Abbott, G. D.: Transformation of organic matter in a Barents Sea sediment profile: coupled geochemical and microbiological processes, *Philos. Trans. R. Soc. Math. Phys. Eng. Sci.*, 378, 20200223, <https://doi.org/10.1098/rsta.2020.0223>, 2020.
- 1205 Stockwell, D. A., Whitley, T. E., Zeeman, S. I., Coyle, K. O., Napp, J. M., Brodeur, R. D., Pinchuk, A. I., and Hunt, G. L.: Anomalous conditions in the south-eastern Bering Sea, 1997: nutrients, phytoplankton and zooplankton, *Fish. Oceanogr.*, 10, 99–116, <https://doi.org/10.1046/j.1365-2419.2001.00158.x>, 2001.
- Sun, M.-Y. and Wakeham, S. G.: Molecular evidence for degradation and preservation of organic matter in the anoxic Black Sea Basin, *Geochim. Cosmochim. Acta*, 58, 3395–3406, [https://doi.org/10.1016/0016-7037\(94\)90094-9](https://doi.org/10.1016/0016-7037(94)90094-9), 1994.
- 1210 Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, *J. Geophys. Res. Atmospheres*, 106, 7183–7192, <https://doi.org/10.1029/2000JD900719>, 2001.
- Thomas, S.: The response of anaerobic prokaryotic processes and communities in the Severn Estuary sediments to environmental changes, Cardiff University, Cardiff, 2014.
- 1215 Thullner, M., Dale, A. W., and Regnier, P.: Global-scale quantification of mineralization pathways in marine sediments: A reaction-transport modeling approach, *Geochem. Geophys. Geosystems*, 10, 1–24, <https://doi.org/10.1029/2009GC002484>, 2009.
- Tromp, T. K., Van Cappellen, P., and Key, R. M.: A global model for the early diagenesis of organic carbon and organic phosphorus in marine sediments, *Geochim. Cosmochim. Acta*, 59, 1259–1284, [https://doi.org/10.1016/0016-7037\(95\)00042-X](https://doi.org/10.1016/0016-7037(95)00042-X), 1995.

- 1220 Uncles, R. J.: Physical properties and processes in the Bristol Channel and Severn Estuary, *Mar. Pollut. Bull.*, 61, 5–20, <https://doi.org/10.1016/j.marpolbul.2009.12.010>, 2010.
- Van Cappellen, P. and Wang, Y.: Cycling of iron and manganese in surface sediments; a general theory for the coupled transport and reaction of carbon, oxygen, nitrogen, sulfur, iron, and manganese, *Am. J. Sci.*, 296, 197–243, <https://doi.org/10.2475/ajs.296.3.197>, 1996.
- 1225 Van Weering, T. C. E., Berger, G. W., and Kalf, J.: Recent sediment accumulation in the Skagerrak, Northeastern North Sea, *Neth. J. Sea Res.*, 21, 177–189, [https://doi.org/10.1016/0077-7579\(87\)90011-1](https://doi.org/10.1016/0077-7579(87)90011-1), 1987.
- Vandewiele, S., Cowie, G., Soetaert, K., and Middelburg, J. J.: Amino acid biogeochemistry and organic matter degradation state across the Pakistan margin oxygen minimum zone, *Deep Sea Res. Part II Top. Stud. Oceanogr.*, 56, 376–392, <https://doi.org/10.1016/j.dsr2.2008.05.035>, 2009.
- 1230 Vasquez-Cardenas, D., Meysman, F. J. R., and Boschker, H. T. S.: A Cross-System Comparison of Dark Carbon Fixation in Coastal Sediments, *Glob. Biogeochem. Cycles*, 34, e2019GB006298, <https://doi.org/10.1029/2019GB006298>, 2020.
- Wakeham, S. G., Lee, C., Hedges, J. I., Hernes, P. J., and Peterson, M. J.: Molecular indicators of diagenetic status in marine organic matter, *Geochim. Cosmochim. Acta*, 61, 5363–5369, [https://doi.org/10.1016/S0016-7037\(97\)00312-8](https://doi.org/10.1016/S0016-7037(97)00312-8), 1997.
- 1235 Wallmann, K., Aloisi, G., Haeckel, M., Obzhairov, A., Pavlova, G., and Tishchenko, P.: Kinetics of organic matter degradation, microbial methane generation, and gas hydrate formation in anoxic marine sediments, *Geochim. Cosmochim. Acta*, 70, 3905–3927, <https://doi.org/10.1016/j.gca.2006.06.003>, 2006.
- Wang, D., Zhu, M.-X., Yang, G.-P., and Ma, W.-W.: Reactive Iron and Iron-Bound Organic Carbon in Surface Sediments of the River-Dominated Bohai Sea (China) Versus the Southern Yellow Sea, *J. Geophys. Res. Biogeosciences*, 124, 79–98, <https://doi.org/10.1029/2018JG004722>, 2019.
- 1240 Wang, Y. and Van Cappellen, P.: A multicomponent reactive transport model of early diagenesis: Application to redox cycling in coastal marine sediments, *Geochim. Cosmochim. Acta*, 60, 2993–3014, [https://doi.org/10.1016/0016-7037\(96\)00140-8](https://doi.org/10.1016/0016-7037(96)00140-8), 1996.
- Weber, T., Cram, J. A., Leung, S. W., DeVries, T., and Deutsch, C.: Deep ocean nutrients imply large latitudinal variation in particle transfer efficiency, *Proc. Natl. Acad. Sci.*, 113, 8606–8611, <https://doi.org/10.1073/pnas.1604414113>, 2016.
- 1245 Wehrmann, L. M., Arndt, S., März, C., Ferdelman, T. G., and Brunner, B.: The evolution of early diagenetic signals in Bering Sea subseafloor sediments in response to varying organic carbon deposition over the last 4.3Ma, *Geochim. Cosmochim. Acta*, 109, 175–196, <https://doi.org/10.1016/j.gca.2013.01.025>, 2013.
- Westrich, J. T. and Berner, R. A.: The role of sedimentary organic matter in bacterial sulfate reduction: The *G* model tested, *Limnol. Oceanogr.*, 29, 236–249, <https://doi.org/10.4319/lo.1984.29.2.0236>, 1984.
- 1250 Wiltshire, K. H. and Manly, B. F. J.: The warming trend at Helgoland Roads, North Sea: phytoplankton response, *Helgol. Mar. Res.*, 58, 269–273, <https://doi.org/10.1007/s10152-004-0196-0>, 2004.
- Ye, H., Yang, T., Zhu, G., Jiang, S., and Wu, L.: Pore water geochemistry in shallow sediments from the northeastern continental slope of the South China sea, *Mar. Pet. Geol.*, 75, 68–82, <https://doi.org/10.1016/j.marpetgeo.2016.03.010>, 2016.

- 1255 Zebracki, M., Eyrolle-Boyer, F., Evrard, O., Claval, D., Mourier, B., Gairoard, S., Cagnat, X., and Antonelli, C.: Tracing the origin of suspended sediment in a large Mediterranean river by combining continuous river monitoring and measurement of artificial and natural radionuclides, *Sci. Total Environ.*, 502, 122–132, <https://doi.org/10.1016/j.scitotenv.2014.08.082>, 2015.
- 1260 Zonneveld, K. A. F., Versteegh, G. J. M., Kasten, S., Eglinton, T. I., Emeis, K.-C., Huguet, C., Koch, B. P., de Lange, G. J., de Leeuw, J. W., Middelburg, J. J., Mollenhauer, G., Prahl, F. G., Rethemeyer, J., and Wakeham, S. G.: Selective preservation of organic matter in marine environments; processes and impact on the sedimentary record, *Biogeosciences*, 7, 483–511, <https://doi.org/10.5194/bg-7-483-2010>, 2010.