

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/146177/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Couper, Keith, Murrells, Trevor, Sanders, Julie, Anderson, Janet E., Blake, Holly, Kelly, Daniel, Kent, Bridie, Maben, Jill, Rafferty, Anne Marie, Taylor, Rachel M. and Harris, Ruth 2022. The impact of COVID-19 on the wellbeing of the UK nursing and midwifery workforce during the first pandemic wave: a longitudinal survey study. *International Journal of Nursing Studies* 127, 104155. [10.1016/j.ijnurstu.2021.104155](https://doi.org/10.1016/j.ijnurstu.2021.104155)

Publishers page: <http://dx.doi.org/10.1016/j.ijnurstu.2021.104155>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



The impact of COVID-19 on the wellbeing of the UK nursing and midwifery workforce during the first pandemic wave: a longitudinal survey study

Keith COUPER ^{a,b}	k.couper@warwick.ac.uk	Twitter @keithcouper
Trevor MURRELLS ^c	trevor.murrells@kcl.ac.uk	Twitter @TJMurrells
Julie SANDERS ^{d,e}	j.sanders@qmul.ac.uk	Twitter @julesanders2
Janet E ANDERSON ^f	janet.anderson@city.ac.uk	Twitter @ProfAndersonHF
Holly BLAKE ^{g,h}	Holly.Blake@nottingham.ac.uk	Twitter @hollyblakenotts
Daniel KELLY ⁱ	KellyDM@cardiff.ac.uk	Twitter @profdkelly
Bridie KENT ^j	bridie.kent@plymouth.ac.uk	Twitter @BridieKent
Jill MABEN ^k	j.maben@surrey.ac.uk	Twitter @nursingpolicy
Anne Marie RAFFERTY ^c	anne_marie.rafferty@kcl.ac.uk	Twitter @annemarieraffer
Rachel M TAYLOR ^{l,m}	rtaylor13@nhs.net	Twitter @ctandrmt
Ruth HARRIS ^c	ruth.harris@kcl.ac.uk	Twitter @RuthHarris_

- a) Warwick Medical School, University of Warwick, Gibbet Hill Road, Coventry, UK
- b) Critical Care Unit, University Hospitals Birmingham NHS Foundation Trust, Birmingham Heartlands Hospital, Bordesley Green, Birmingham, B9 5SS, UK
- c) Florence Nightingale Faculty of Nursing, Midwifery & Palliative Care, King's College London, 57 Waterloo Road, London, SE1 8WA, UK
- d) St Bartholomew's Hospital, Barts Health NHS Trust, West Smithfield, London, EC1A 7BE, UK
- e) William Harvey Research Institute, Queen Mary University of London, Charterhouse Square, London, EC1M 6BQ
- f) School of Health Sciences, City University of London, Northampton Square, London, EC1V 0HB, UK
- g) School of Health Sciences, University of Nottingham, Queen's Medical Centre, Derby Road, Nottingham, NG7 2HA, UK
- h) NIHR Nottingham Biomedical Research Centre, Queen's Medical Centre, Derby Road, Nottingham, NG7 2UH, UK
- i) School of Healthcare Sciences, Cardiff University, Eastgate House, Cardiff, CF24 0AB, UK
- j) Faculty of Health, University of Plymouth, Rolle Building, Drake Circus, Plymouth, PL4 8AA, UK
- k) School of Health Sciences, University of Surrey, 30 Priestley Rd, Guildford, GU2 7YH
- l) Centre for Nurse, Midwife and AHP Led Research, University College London NHS Foundation Trust, 2nd Floor North, 250 Euston Rd, London NW1 2PG, UK
- m) EGA Institute for Women's Health, Faculty of Population Health Sciences, University College London, Medical School Building, 74 Huntley Street, London, WC1E 6AU, UK

Corresponding author:

Dr Keith Couper, Warwick Medical School, University of Warwick, Coventry, CV4 7AL, UK. Email: k.couper@warwick.ac.uk; Telephone: 02476151179

[Click here to view linked References](#)

ABSTRACT

Background

The specific challenges experienced by the nursing and midwifery workforce in previous pandemics have exacerbated pre-existing professional and personal challenges, and triggered new issues. We aimed to determine the psychological impact of the COVID-19 pandemic on the UK nursing and midwifery workforce and identify potential factors associated with signs of post-traumatic stress disorder.

Methods

A United Kingdom national online survey was conducted at three time-points during the first wave of the COVID-19 pandemic between April and August 2020 (T1 and T2 during initial wave; T3 at three-months following the first wave). All members of the UK registered and unregistered nursing and midwifery workforce were eligible to participate. The survey was promoted via social media and through organisational email and newsletters.

The primary outcome was an Impact of Events Scale-Revised score indicative of a post-traumatic stress disorder diagnosis (defined using the cut-off score ≥ 33). Multivariable logistic regression modelling was used to assess the association between explanatory variables and post-traumatic stress disorder.

Results

We received 7840 eligible responses (T1- 2040; T2- 3638; T3- 2162). Overall, 91.6% participants were female, 77.2% were adult registered nurses, and 28.7% were redeployed during the pandemic.

An Impact of Events Scale-Revised score ≥ 33 (probable post-traumatic stress disorder) was observed in 44.6%, 37.1%, and 29.3% participants at T1, T2, and T3 respectively. At all three time-points, both personal and workplace factors were associated with probable post-traumatic stress disorder, although some specific associations changed over the course of the pandemic. Increased age was associated with reduced probable post-traumatic stress disorder at T1 and T2 (e.g. 41-50 years at T1 odds ratio (OR) 0.60, 95% confidence interval (CI) 0.42-0.86), but not at T3. Similarly, redeployment with inadequate/ no training was associated with increased probable post-traumatic stress disorder at T1 and T2, but not at T3 (T1 OR 1.37, 95% CI 1.06-1.77; T3 OR 1.17, 95% CI 0.89-1.55). A lack of confidence in infection prevention and control training was associated with increased probable post-traumatic stress disorder at all three time-points (e.g. T1 OR 1.48, 95% CI 1.11-1.97).

Conclusion

A negative psychological impact was evident 3-months following the first wave of the pandemic. Both personal and workplace are associated with adverse psychological effects linked to the COVID-19 pandemic.

These findings will inform how healthcare organisations should respond to staff wellbeing needs both during the current pandemic, and in planning for future pandemics.

Keywords: COVID-19, nurses, midwives, pandemics, post-traumatic stress disorders, professional burnout, psychological distress, survey

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

What is already known about the topic?

- Nursing and midwifery workforce wellbeing is under strain due to staff shortages and high stress and burnout.
- The nursing and midwifery workforce are at the forefront of the COVID-19 response.
- In previous pandemics, the nursing and midwifery workforce has experienced worse psychological effects, compared with other healthcare professional groups.

What this paper adds

- Almost 30% of survey respondents reported experiences indicative of a probable post-traumatic stress disorder diagnosis three-months after the first pandemic peak.
- Key workplace-related factors were associated with adverse psychological effects, including redeployment to other clinical areas without adequate training and inadequate infection control training.
- Healthcare organisations should urgently implement evidence-based strategies to support the wellbeing of members of the nursing and midwifery workforce affected by the pandemic, and develop robust workforce plans in preparation for future pandemics.

Introduction

1 The year 2020 was designated by the World Health Organization as International Year of the Nurse
2 and Midwife (World Health Organization, 2020). This reflects the critical role nurses and midwives
3 play in international health and social care delivery, policy and research. Underpinning this
4 designation is a workforce already under significant pressure with high levels of attrition, ongoing
5 recruitment challenges, and an ageing workforce (World Health Organization, 2020). Across the
6 world, a shortfall of 5.7 million nurses has been predicted by 2030 (World Health Organization, 2020).
7 In the United Kingdom (UK), the National Health Service (NHS) long-term plan included a
8 commitment to improve staff recruitment and retention (National Health Service, 2019). However,
9 this is against a backdrop of over 40,000 whole-time equivalent nursing vacancies and rates of
10 dissatisfaction and plans to leave the profession in the UK that exceed many other European
11 countries (Aiken et al., 2012, National Audit Office, 2020).
12
13
14
15

16
17 At the end of 2019, just prior to the start of the International Year of the Nurse and Midwife, the
18 Severe Acute Respiratory Syndrome- Coronavirus 2 virus was first identified in China. The
19 subsequent COVID-19 pandemic has both highlighted nurses' and midwives' essential role in
20 healthcare delivery and served to exacerbate existing workforce challenges. The impact of COVID-
21 19 on society has been well-documented, both as a direct result of the virus and indirectly through
22 strategies (e.g., restrictions on travel and social gatherings; closure of workplaces and educational
23 institutions) implemented to limit infection transmission (Douglas et al., 2020). In addition to
24 these wider societal effects, nurses and midwives have faced the added challenge of healthcare
25 provision in the context of lack of personal protective equipment, the rapid implementation of
26 redeployment (move to a different clinical area to support the pandemic response), new ways of
27 working, and vast numbers of high acuity patients (Fernandez et al., 2020, Liu et al., 2020, Maben
28 and Bridges, 2020). Nurses and midwives have also faced the emotional challenge of high
29 mortality rates, with 30.8% patients hospitalised with COVID-19 dying (Navaratnam et al., 2021).
30 Observational studies highlight the increased risk of COVID-19 infection to frontline healthcare
31 workers, such as nurses and midwives (Chou et al., 2020, Mutambudzi et al., 2020, Nguyen et al.,
32 2020, Shah et al., 2020). The increased risk of infection may also extend to members of the
33 healthcare worker's family (Shah et al., 2020). Reports have also highlighted the tragic loss of life
34 due to COVID-19 infection amongst member of the nursing and midwifery
35 workforce (Bandyopadhyay et al., 2020).
36
37
38
39
40
41
42

43 Early COVID-19 data from China and studies of previous pandemics and disease outbreaks have
44 highlighted the potential effect of the pandemic on healthcare worker's mental health (Allan et al.,
45 2020, Cabarkapa et al., 2020, De Kock et al., 2021, Fernandez et al., 2020, Stuijtzand et al., 2020).
46 Some previous COVID-19 studies report worse psychological effects in nurses, compared with
47 other healthcare workers (Allan et al., 2020, Cabarkapa et al., 2020). A recent systematic review of
48 studies on mental health disorders in hospital-based healthcare workers in hospitals impacted by
49 pandemics included 19 studies, of which most described the impact of the Severe Acute
50 Respiratory Syndrome pandemic in Canada and South Asia (Allan et al., 2020). The review
51 identified that post-traumatic stress, anxiety, and depression were common in healthcare workers
52 during pandemics, with some concerning long-lasting effects.
53
54
55
56

57 This combination of pre-existing workforce challenges, rapid changes to professional life and ways
58 of working, and risk of illness to both the individual and their family may have important and
59 demonstrable psychological impacts on the UK nursing and midwifery workforce. On this basis, we
60 identified the need to rapidly design and undertake research to understand the psychological
61
62
63
64
65

1 impacts of the first wave of the COVID-19 pandemic on the UK nursing and midwifery workforce.
2 This has been recognised as a key research priority by members of the public, mental health
3 experts, and members of the nursing and midwifery workforce (Holmes et al., 2020, Manning et
4 al., 2021). The aims of the study were to assess the mental health effects of the pandemic over
5 time on the nursing workforce and identify the personal and workplace factors associated with
6 wellbeing to inform the response to the current pandemic and support preparation for future
7 pandemics.
8
9
10
11

12 **Methods**

13
14
15 A national longitudinal online survey study was conducted utilising a self-selecting sample of the
16 UK nursing and midwifery workforce across three time-points to explore the self-reported
17 personal psychological impact of the COVID-19 pandemic. The survey design and methodology
18 drew on that developed for the COVID-19 emergency response assessment study (Roberts et al.,
19 2021, Roberts et al., 2020). This paper is reported in accordance with the Checklist for Reporting
20 Results of Internet E-Surveys (Eysenbach, 2004).
21
22
23
24
25

26 **Overview of survey**

27 The survey was administered at three time-points (T1, T2, T3) through an online survey platform
28 (Qualtrics, Provo, Utah, USA) and could be completed in approximately 10-15 minutes.
29
30

31 Our T1, T2, and T3 surveys were open for responses between 2nd and 14th April 2020, 28th April
32 and 12th May 2020, and 30th July and 20th August 2020 respectively. During the first pandemic
33 wave the number of hospitalised cases increased more rapidly than expected and then decreased
34 more rapidly than expected. As such, both the highest number of hospitalised COVID-19 patients
35 (21,686 on 12th April 2020) and number of deaths where COVID-19 was recorded on the death
36 certificate (1,457 on 8th April 2020) occurred during our T1 survey (UK Government, 2021). Figure
37 one shows our survey distribution time points in the context of the number of hospitalised
38 patients across the UK. There was some regional variation in date of the peak and subsequent
39 recovery trajectory (supplementary material, figure e1). Our T3 survey, issued three months after
40 the T2 survey, was designed to reflect a COVID-19 recovery period.
41
42
43
44

45 Survey questions covered six broad domains: participant demographics, professional
46 characteristics, experience of work during the pandemic, personal impact of COVID-19, access to
47 psychological support, and mental health and wellbeing outcomes (supplementary material table
48 e1). Mental health and wellbeing outcomes were collected using validated tools, namely the
49 impact of events-revised scale, Depression Anxiety Stress scales- 21 (DASS-21), Maslach Burnout
50 Inventory (T3 survey only) and intention-to-leave scale (T3 survey only) (Antony et al., 1998, Chen
51 et al., 2019, Creamer et al., 2003, Lovibond and Lovibond, 1996, Poghosyan et al., 2009, Weiss,
52 2006).
53
54
55
56

57 Questions were consistent across each survey, although minor changes were made to reflect
58 emerging evidence as the pandemic progressed. In particular, we included ethnicity for our second
59 and third survey to reflect evidence highlighting the disproportionate impact of COVID-19 on
60 Black, Asian and minority ethnic populations (Ferrando-Vivas et al., 2021, Williamson et al., 2020).
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Eligibility criteria

Study participation was open to all members of the nursing and midwifery workforce working in the UK. This included registered nurses and midwives, student nurses and midwives, healthcare assistants, nursing associates, and trainee nursing associates. Inclusion was not limited by employer type.

Survey promotion and distribution

Information about the survey and an internet link were widely distributed through social media (Twitter, Facebook). High-profile individuals in the UK nursing and midwifery community actively encouraged survey promotion by participants. In addition, survey information was included in emails distributed by key nursing and midwifery organisations (e.g., Royal College of Nursing) and the UK nursing and midwifery regulator (Nursing and Midwifery Council). The survey was also reported as a news story in mainstream journals (Mitchell, 2020). Access to the survey was open to any individual with the study internet address link. At both T2 and T3, we also directly emailed individuals that had completed our survey at a previous time-point and who had provided their email address and consented to use of their personal data for this purpose. A reminder email was sent approximately half-way through the survey period. No incentive was offered for completion.

Following T1 and T2 surveys, a brief summary of aggregated data was rapidly distributed so that findings could inform public policy. These summaries were featured by various news outlets including BBC News, and national newspapers (BBC News, 2020, Hackett, 2020, Toynbee, 2020).

Survey design

Survey questions were based on the author’s expert subject knowledge, evidence from previous pandemics and disease outbreaks (e.g., Ebola, Severe Acute Respiratory Syndrome), other ongoing surveys, and emerging information about the COVID-19 pandemic (Roberts et al., 2021, Roberts et al., 2020). The first page of the survey provided key information, including its purpose and length, and how we would manage personal data. We mandated completion of only the first two questions, which were used to confirm eligibility and record consent for study participation. Completion of subsequent questions was not mandated.

Adaptive questioning was used to filter questions that were not relevant based on preceding responses (e.g., questions about experience of redeployment were shown only to those individuals that were redeployed). We did not randomise the order of any part of the survey. In designing the survey, text entry was minimised and categorical responses were favoured. A survey back button allowed participants to review their answers as they progressed through the survey. The survey design was reviewed at each time-point to reflect emerging data and, where necessary, alter question grammatical tense to reflect the stage of the pandemic. For each survey version, the research team completed a draft of the survey using different response variations to ensure that the survey functioned as expected.

1 In the survey, we invited participants to provide their email address, and explicit agreement was
2 sought to use this to: link information across surveys; inform the participant about surveys at
3 subsequent time-points, and invite them to participate in future research. We did not limit
4 responses by IP addresses as we anticipated that participants may complete the survey at their
5 place of work or using a shared home internet router, such that a restriction might prevent
6 completion by eligible colleagues or family members, who were also nurses.
7

8 **Outcomes**

9 The primary outcome was the prevalence of post-traumatic stress disorder, measured using the
10 impact of events scale-revised at each time point (Creamer et al., 2003, Weiss, 2006). For each of
11 the 22 items in the scale, participants were asked to rate the extent to which they had been
12 distressed in relation to COVID-19 over the preceding 7-days. Based on previous research, we
13 categorised a score of 33 or above as indicative of probable post-traumatic stress disorder
14 (Creamer et al., 2003).
15
16
17

18 Secondary outcomes were depression, anxiety, stress, job satisfaction, burnout and the
19 participant's intention to leave both their current role and healthcare, measured using validated
20 tools (Antony et al., 1998, Chen et al., 2019, Lovibond and Lovibond, 1996, Poghosyan et al.,
21 2009). A summary of outcome definitions and tools is included in the supplementary material
22 (table e2). Outcomes were measured at each of the three time points, except for burnout and
23 intention to leave which were captured at time-point three only to reflect the need to measure
24 these concepts in a period of relative stability.
25
26
27
28
29
30

31 **Statistical analysis**

32 Data were downloaded directly from the survey hosting platform (Qualtrics, Provo, Utah, USA).
33 Initial data processing and cleaning were undertaken using SPSS version 26 (IBM SPSS Statistics,
34 Armonk, NY, USA). Imputation and modelling were undertaken using Stata version 16.1 (College
35 Station, Texas, USA).
36
37
38

39 Our initial review of the three datasets identified 267 individuals that had completed the survey at
40 all three time-points and 898 (216 at T1/T2; 154 at T1/T3; 528 at T2/T3) individuals completed the
41 survey at two time-points. This is likely to be an underestimate of the true rate as we were only
42 able to track responses where an individual had both provided a matching email address and had
43 given permission to link data. Given the low frequency of completion at multiple time-points, we
44 decided to treat each survey time-point as an independent sample in our primary analysis.
45
46
47

48 To be included in the analysis, we required participants to have completed one or more questions
49 beyond the first two survey sections (professional characteristics and participant demographics),
50 and to have completed at least one outcome measure (one of: impact of events scale-revised,
51 Depression Anxiety Stress scales- 21 , Maslach Burnout Inventory (T3 only) and intention-to-leave
52 scale (T3 only)).
53
54

55 We identified duplicate responses through the recorded email address. Across T1, T2, and T3, we
56 identified 310 duplicate responses. In most cases (93% duplicates), the first record was selected
57 for inclusion. However, in a minority of cases (7%) where the first record contained little data or
58 key outcomes (impact of events scale-revised, Depression Anxiety Stress scales- 21, and Maslach
59 Burnout Inventory) were missing, we included the second response.
60
61
62

1 We observed data missingness across survey responses due to abandonment of the survey during
2 completion and, presumed unintentional, omission of individual questions within the survey.
3 Multiple imputation using the chained equations method was used to impute missing data. The
4 imputation model included all variables used in the subsequent multivariable modelling plus other
5 variables that were thought might predict either study outcomes or explanatory variables that we
6 planned to model (supplementary material table e1) (Royston and White, 2011). Twenty imputed
7 datasets were created using a method appropriate for data consisting of multiple psychometric
8 scales (Plumpton et al., 2016). Based on the fraction of missing information measure from the
9 multivariable models (T1- 0.098; T2- 0.133; T3- 0.121), 14 datasets were required to meet the
10 criterion proposed by the UCLA Statistical Consulting Group (UCLA: Statistical Consulting Group).
11
12

13 All data variables, except for some outcomes, were categorical and are described using numbers
14 and percentages. Our outcome measures included both those that provide an overall score and
15 those that categorise individuals based on score cut-off values. For the former, we summarise data
16 using the mean and 95% confidence interval or median and interquartile range, depending on data
17 normality. The latter, we present as number and percentage in each category. We calculated
18 Cronbach's α , mean scores and 95% confidence interval (CI) and Pearson correlations for the
19 impact of events scale-revised and Depression Anxiety Stress scales- 21 sub-scales across time-
20 points. For the Maslach Burnout Inventory, we calculated Cronbach's α for sub-scales. We also
21 explored the correlation between the impact of events scale-revised and Depression Anxiety
22 Stress scales- 21 sub-scales, and the impact of pattern of response on impact of events scale-
23 revised and Depression Anxiety Stress scales- 21.
24
25
26
27
28

29 We tested for equality across time. The type of test used depended on whether the variable had
30 been imputed or not, and its measurement scale. For variables with two categories a logistic
31 regression model was used and an ordinal regression model was used for three or more ordered
32 categories. For variables with three or more categories with no natural order a multilogit model
33 was used if the data had been imputed and the Pearson χ^2 test if not. The logistic, multilogit and
34 ordinal regression models fitted to imputed variables were tested for equality using a F statistic
35 and the likelihood ratio statistic (χ^2) for all other non-imputed variables. To improve readability,
36 we combine some of the ordinal categories in the text for reporting, although the statistical test
37 always relates to the original scale. A regression model was used to compute a F statistic for
38 continuous variables although the degrees of freedom for imputed and non-imputed variables
39 were calculated differently.
40
41
42
43
44

45 We fitted multivariable logistic regression models to individual imputed datasets, and then
46 combined the results to obtain estimates of the association between key explanatory variables
47 and an impact of events scale-revised score of 33 or more (probable post-traumatic stress
48 disorder) (Creamer et al., 2003). We included variables based on clinical plausibility
49 (supplementary material table e1), informed by previous literature, and developed individual
50 models for each time-point. We measured the association between the explanatory variables and
51 the dichotomous outcomes using odds ratios (OR) and 95% CI. For explanatory variables with
52 three or more categories, we tested the overall effect for statistical significance using a F statistic.
53 In a sensitivity analysis that was not defined *a priori*, we added a variable that categorised
54 participant's pattern of response (e.g. those that only completed at a single time-point and those
55 that completed at multiple time-points) to explore how this affected our model's findings.
56
57
58
59
60

61 Ethical approval

1 The study was approved by the University of Warwick Biomedical and Scientific Research Ethics
2 Committee on 27th March 2020 (reference 101/19-20). Participants gave informed consent for
3 participation through a checkbox at the start of the survey. All survey data were stored in
4 accordance with national legislation and institutional policies.
5
6

7 **Results**

8
9 Across the three survey time-points, the survey was accessed on 12,010 occasions (3299 T1, 5758
10 T2, 2953 T3), of which we finally included 7,840 responses (figure two). The highest number of
11 responses (n=3,638) was received at T2.
12
13

14 **Participant characteristics**

15 Across all survey responses, most participants were female (T1 91.6%; T2 91.6%; T3 91.9%, p=0.20,
16 and of white ethnicity (T2 89.3%; T3 93.1%, p<0.001). The most common age group was 51-60
17 years (T1 30.7%; T2 34.7%; T3 39.8%, p<0.001). Professionally, the majority were adult registered
18 nurses (T1 77.9%; T2 75.1%; T3 80.0%, p=0.001). Table one summarises key personal and
19 workplace variables, with the remainder summarised in table e3 (supplementary material).
20
21
22

23 **Personal and professional experiences of COVID-19**

24 Across time-points, 28.7% (n=2,251) participants reported being redeployed in response to the
25 pandemic (table one, table e3 supplementary material). Of these, 63.2% (n=1,422) reported that
26 training to prepare for redeployment either did not occur or was inadequate.
27
28
29

30 Participants often (n=3,133, 40.0%) reported that they lacked confidence in infection prevention
31 and control training, or that training had not been received, although this declined over time (T1
32 52.1%; T2 37.3%; T3 33.0%, p<0.001). In total, 22.6% (n=1772) disagreed or strongly disagreed that
33 the correct personal protective equipment was always available, similarly this decreased across
34 survey time-points (T1 35.5%; T2 18.3%; T3 17.8%, p<0.001). Table one summarises these data
35 across all time-points, with a breakdown by time-point included in the supplementary material.
36
37
38

39 At time-point three, 34.3% (n=741) participants reported that they had experienced a COVID-19
40 infection. The clinical workplace was the most commonly identified source of infection (n=477,
41 69.5%). Despite only 5.6% (n=38) requiring hospital care, 35.2% (n=241) reported that they had
42 not fully recovered.
43
44

45 **Primary outcome- Impact of events scale- -revised**

46 An impact of events scale-revised score of 33 or above (probable post-traumatic stress disorder)
47 was observed in 44.6% participants at time-point one, 37.1% participants at T2, and 29.3%
48 participants at T3 (p<0.001). The mean impact of events scale-revised score similarly decreased
49 across time-points (T1 32.16, 95% CI 31.25-33.07; T2 28.11, 95% CI 27.43-28.79; T3 23.81, 95% CI
50 22.91-24.71, p<0.001). Data are summarised in table two and the supplementary material (table
51 e4). Cronbach's α ranged from 0.89 to 0.94 across time-points and sub-scales (electronic supplement
52 table e5).
53
54
55
56

57 **Secondary outcomes**

58 We observed decreases over time in the prevalence of severe and extremely severe anxiety and
59 stress (e.g., severe/ extremely severe anxiety (T1 25.0%; T2 21.7%; T3 18.1%, p<0.001)). However,
60 no changes were observed in prevalence of depression over time (T1 19.7%; T2 20.0%; 19.6%,
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

p=0.25). Job satisfaction declined across time-points with the lowest proportion reporting being very satisfied or somewhat satisfied with their job at T3 (T1 69.5%; T2 67.8%; 57.2%, p<0.001). Cronbach's α ranged from 0.88 to 0.93 across time-points and sub-scales (electronic supplement table e5). There was a strong positive correlation between Depression Anxiety Stress scales- 21 sub-scales and impact of events scale-revised sub-scales across time-points (electronic supplement table e6).

At T3, mean scores for the Maslach Burnout Inventory scales of emotional exhaustion, depersonalization, and personal accomplishment were 26.48 (95% CI 25.89-27.07), 5.52 (95% CI 5.27-5.78), and 34.64 (95% CI 34.25-35.02) respectively. Cronbach's α ranged from 0.76 to 0.92 across sub-scales (electronic supplement table e5). The level of intention to leave was higher for an individual's current job (37.8, 95% CI 36.3-39.4) than it was for their intention to leave healthcare (28.6, 95% CI 27.2-30.0), although breakdown by response indicated polarised views with 26.1% having no plan to leave their current role, whilst 19.4% were giving high consideration to leaving with intermediate views having comparatively fewer responses (supplementary material (table e4).

Modelling of association between variables and probable post-traumatic stress disorder

In the multivariable logistic regression model (table three), factors were classified as personal (e.g., age, gender, caring responsibility) and workplace factors (e.g., clinical speciality, redeployment, personal protective equipment availability). For some factors, the low number of participants in a specific category means that our estimate of the association with the primary outcome is imprecise. Across time-points, we identified variability in the factors associated with our outcome of an impact of events scale-revised score of 33 or more.

Modelling- personal factors

For personal factors, compared with those aged 16-30, older participants were less likely to have an impact of events scale-revised score of 33 or more, although this was not statistically significant at T3 (e.g., age 51-60 T1 OR 0.53, 95% CI 0.37 – 0.75; T2 OR 0.67 95% CI 0.51 – 0.88; T3 OR 0.63 95% CI 0.38 – 1.06). Compared with white ethnicity, being of Black/ African/ Caribbean ethnicity was associated with impact of events scale-revised score of 33 or more at T2 (OR 1.72, 95% CI 1.24-2.38), but this association was not observed at T3 or for other ethnicities. A sole caring responsibility was associated with impact of events scale-revised score of 33 or more at T1 (OR 1.51, 95% CI 1.08-2.09) and T3 (OR 1.58, 95% CI 1.11-1.25), but not at T2 (OR 1.25, 95% CI 0.98-1.59). We did not find evidence of a clear association between gender, geographical region and professional qualification, and our outcome.

Modelling- workplace factors

For workplace factors, being employed by the NHS was associated with impact of events scale-revised score of 33 or more at T1 (OR 1.37, 95% CI 1.03-1.80), but not at subsequent time-points. Clinical speciality was not consistently associated with an impact of events scale-revised score of 33 or more, although at T2 working in critical/ operating theatres (OR 1.51, 95% CI 1.18-1.92) and working in a care/ nursing home (OR 1.64, 95% CI 1.03-2.60), compared with acute/ emergency hospital care, was associated with increased odds of impact of events scale-revised score of 33 or more.

Whilst redeployment with adequate training was not associated with impact of events scale-revised score of 33 or more, we identified an association between redeployment without training or inadequate training and impact of events scale-revised score of 33 or more at both T1 and T2 (e.g., T2 OR 1.50, 95% CI 1.22–1.84), but not at T3 (OR 1.17, 95% CI 0.89-1.55). A similar pattern

1 was observed in relation to participants who disagreed or strongly disagreed that the correct
2 personal protective equipment was always available (e.g., T2 OR 1.34, 95% CI 1.07-1.69; T3 OR
3 1.23, 95% CI 0.86-1.77). The only explanatory variable response associated with impact of events
4 scale-revised score of 33 or more at all three time-points was being not (or somewhat not)
5 confident in COVID-19 infection prevention and control training (T1 OR 1.48, 95% CI 1.11-1.97; T2
6 OR 1.29, 95% CI 1.03-1.61; T3 1.55, 95% CI 1.09-2.19). At T3, symptomatic COVID-19 infection was
7 associated with impact of events scale-revised score of 33 or more (OR 1.56, 95% CI 1.23–1.98).
8
9

10 **Modelling - participants pattern of response**

11
12 For the outcomes of impact of impact of events scale-revised score of 33 or more and Depression
13 Anxiety Stress scales- 21, results were broadly consistent with our primary analysis and across all
14 response patterns (electronic supplement table e7 and table e8). The inclusion of a pattern of
15 response variable in the multivariable models made little difference to the parameter estimates of
16 other variables (electronic supplement table e9).
17
18
19
20
21
22
23

24 **Discussion**

25
26 In this large national longitudinal survey study over the course of the first wave of the COVID-19
27 pandemic, we found that the UK nursing and midwifery workforce experienced a high prevalence
28 of negative psychological effects, including severe stress, severe anxiety and high impact of events
29 scale-revised scores, indicative of probable post-traumatic stress disorder. Prevalence of some
30 psychological issues decreased following the first pandemic peak, but a significant proportion of
31 participants continued to experience negative psychological effects during the pandemic recovery
32 phase. Factors associated with an impact of events scale-revised score of 33 or more, indicative of
33 probable post-traumatic stress disorder, varied across time-points. Personal factors including
34 younger age and caring responsibilities were associated with increased distress, as were
35 workplace factors such as availability of personal protective equipment and delivery of essential
36 training. Workplace factors are potentially modifiable. Our results provide important information
37 to support identification of at-risk staff and inform workforce support planning for future
38 pandemics.
39
40
41
42
43
44

45 Our methodology and survey design drew on that developed for the COVID-19 Emergency
46 Response Assessment study, which surveyed emergency medicine, intensive care, and anaesthetic
47 doctors during the pandemic acceleration phase, peak phase, and one-month following the peak
48 (Roberts et al., 2021, Roberts et al., 2020). Collecting data across three time-points allowed us to
49 explore changes in key psychological measures over the course the first wave of the pandemic,
50 and into the recovery period. This contrasts with the majority of other studies of mental health in
51 the context of a pandemic that have surveyed healthcare workers at only one single time-point,
52 often with a focus on hospital-based healthcare staff (Ali et al., 2020, Chew et al., 2020, Dykes et
53 al., Lai et al., 2020, Nickell et al., 2004, Wu et al., 2009, Zhu et al., 2020). Whilst such studies
54 provide some helpful information about the impact of the pandemic, follow-up assessments are
55 needed to understand the potential ongoing impact. A further limitation of many previous studies
56 in this area is their focus on the hospital-based workforce, thereby missing the impact on those
57 working in other areas, such as primary care and care homes.
58
59
60
61
62
63
64
65

1 As our experience shows, undertaking research in the context of a global pandemic, is challenging,
2 particularly in mapping survey distribution to the unpredictable impact of the pandemic on society
3 and the healthcare system. Despite rapidly developing the study protocol and survey tool, and
4 expedited ethical review, our T1 survey coincided with the rapid increase in COVID-19 hospitalised
5 cases in the UK and crossed the date on which the number of COVID-19 hospitalised patients
6 peaked during the first wave.(UK Government, 2021) This may, in part, explain the apparent
7 improvement in wellbeing between T1 and T2, as individuals became more normalised to
8 healthcare delivery in a pandemic, organisations began to improve their response (e.g., through
9 improved availability of personal protective equipment) and overall healthcare system demand
10 began to decline.
11
12

13
14 In our study, we reported a prevalence of probable post-traumatic stress disorder, based on an
15 impact of events scale-revised score of 33 or more, of 45%, 37%, and 29% at T1, T2, and T3
16 respectively. As such, our observed rate was higher than that reported in many studies prior to the
17 COVID-19 pandemic (Kinman et al., 2020). We identified similar patterns for the outcomes of
18 stress and anxiety, whilst prevalence of depression did not change across survey time-points.
19 Maslach burnout inventory scores indicated moderately high levels of emotional exhaustion, that
20 exceeded population norms, and levels of depersonalisation and personal accomplishment that
21 are in line with population norms (Maslach et al., 2018). Intention to leave showed polarised views
22 with some individuals expressing a clear desire to leave both their current role and healthcare.
23 There are likely important and complex relationships between these outcomes, such that the long-
24 term trajectory of these conditions remains unclear.
25
26
27
28

29 Recent studies of post-traumatic stress disorder prevalence, which also used the impact of events
30 scale-revised score and similar cut-off values, in healthcare workers in the context of COVID-19
31 have reported rates ranging from 12.6 to 29.8% during the pandemic peak (Dykes et al., Roberts et
32 al., 2021, Zhu et al., 2020). These data are consistent with a systematic review of 19 studies from
33 both the COVID-19 pandemic and previous pandemics that reported a pooled prevalence of
34 clinically important post-traumatic stress symptoms in healthcare workers in the acute pandemic
35 phase of 23.4% (95% CI 16.3-31.2%), although heterogeneity was high ($I^2=96.2%$) and seemingly
36 driven by differences in population, the nature of the pandemic, and method of measuring post-
37 traumatic stress (Allan et al., 2020).
38
39
40
41

42 The reason for the markedly higher prevalence of probable post-traumatic stress disorder in our
43 study, compared with previous studies, is unclear. Previous studies have reported that post-
44 traumatic stress disorder is more common in members of the nursing and midwifery workforce
45 than other healthcare professions, although this finding is not consistent across all studies (Allan
46 et al., 2020, Cabarkapa et al., 2020, Stuijtzand et al., 2020). Our study identified a variety of factors
47 associated with probable post-traumatic stress disorder, and these factors changed across time-
48 points. Key factors were linked to the organisational response to the pandemic (e.g., availability of
49 personal protective equipment and training adequacy) highlighting the important role of
50 employers in mitigating the impact of the pandemic on psychological wellbeing in the workforce.
51 These factors may, in part, reflect the risk of contracting COVID-19 or the perceived health impact
52 if the individual became infected. The factors identified are broadly consistent with those reported
53 in a recent systematic review of 59 studies from a range of countries across four continents and a
54 range of previous pandemics.(Kisely et al., 2020) This supports the potential generalisability of our
55 findings beyond the UK setting. However, some differences were noted. For example, we found
56 that redeployment with adequate training was not associated with probable post-traumatic stress
57
58
59
60
61
62
63
64
65

1 disorder, but an association was identified where redeployment occurred with inadequate or no
2 training. This indicates the importance of planning adequate workforce preparation for future
3 pandemics.

4 The observed decrease in negative psychological effects over the study period is consistent with
5 previous studies with a longer follow-up period, although these same studies highlight that an
6 important proportion of individuals will experience ongoing psychological effects (Allan et al.,
7 2020, McAlonan et al., 2007, Stuijzand et al., 2020, Wu et al., 2009). This has the potential to
8 increase levels of sickness and staff attrition, and negatively affect patient safety (Dyrbye et al.,
9 2019, Hall et al., 2016, Jun et al., 2021, Kinman et al., 2020). A further concern is that at T3, one-
10 third reported having had a COVID-19 infection, of whom 35.2% reported that they had not yet
11 fully recovered. Long-COVID is increasingly being recognised as an important health condition,
12 with both physical and psychological effects (Greenhalgh et al., 2020, Taquet et al., 2021). As such,
13 some individuals may require significant support to return to their previous work role, to avoid
14 their potential loss to the health and social care workforce.
15
16
17
18

19 The key strength of our study is its large sample size with participants from all members of the
20 nursing and midwifery workforce in a country that has been heavily impacted by COVID-19
21 (Bilinski and Emanuel, 2020, UK Government, 2021). In contrast to previous research, we chose
22 not to limit our sample to a specific clinical area or specific members of the nursing and midwifery
23 workforce (e.g. only registered nurses), thereby allowing us to produce findings that are relevant
24 to the whole workforce.
25
26
27

28 Our study does, however, have some important limitations. Firstly, whilst we collected nearly
29 8,000 eligible survey responses across the three time-points, our sample was self-selecting and
30 represented a relatively small proportion of the UK's total nursing and midwifery workforce. There
31 are around 400,000 members of the nursing and midwifery workforce directly employed by the
32 NHS in England, but these data do not include the number of individuals working in other UK
33 countries, or those not employed by the NHS, such as those working in social care. (NHS Digital,
34 2018) As such, it is not possible to estimate our survey response rate. Recent data from the
35 Nursing and Midwifery Council indicates that there 731,918 individuals on the professional
36 register, although not all of these individuals will be currently working and these data exclude
37 the many non-registered members of the workforce, such as healthcare assistants. (Nursing and
38 Midwifery Council, 2021) The proportion of nursing and midwifery registrants identifying as
39 female (89%) is similar to our sample (91.6%), whilst the proportion of non-white individuals is
40 markedly higher (21.7%) than our sample (9.3%). We also observed low numbers of responses
41 from healthcare assistants, who will typically receive lower salaries, despite our attempts to
42 specifically target both non-registered individuals and Black, Asian and minority ethnic groups
43 through social media at T2 and T3. Given the significant negative impact of COVID-19 on these
44 groups, we may have underestimated the true psychological impact on the total workforce
45 (Ferrando-Vivas et al., 2021, Williamson et al., 2020). Secondly, we observed some differences in
46 baseline characteristics across surveys. Whilst our modelling adjusted for known confounders, our
47 findings across surveys may have been influenced by unmeasured confounders. Thirdly, our data
48 linkage has some limitations. We allowed individuals to choose whether they supplied an email
49 address and how we used their data (including permission to link data across surveys). In addition,
50 at T2 and T3, we did not require participants to have completed one of the preceding surveys. As
51 such, for analysis purposes, we decided to treat the three surveys as independent samples.
52 However, where we explored the impact of pattern of response in sensitivity analyses, we found
53 that findings were consistent with our primary analysis. Fourthly, we developed our survey rapidly
54
55
56
57
58
59
60
61
62
63
64
65

1 and we sought to develop a tool that could be completed in 15-20 minutes. In developing our
2 survey, we drew on the wider literature and expert knowledge to identify and prioritise questions,
3 but it is possible that we may have missed additional factors associated with psychological
4 wellbeing. Fifthly, whilst we used validated tools to assess psychological impacts, these tools
5 cannot be used for the diagnosis of psychological conditions, but they may help identify individuals
6 that may be at risk and benefit from further assessment and support. In particular, our definition
7 of probable post-traumatic stress disorder did not fulfil all the necessary criteria (e.g., persistence
8 of symptoms for over one-month and evidence of functional impairment) for a clinical diagnosis
9 (Lancaster et al., 2016). Finally, our survey focussed on the first wave of the peak, such that the
10 enduring impact of subsequent pandemic waves on psychological wellbeing is yet to be
11 determined.
12

13 **Conclusion**

14
15
16
17
18 In this survey study, we observed a concerning prevalence of post-traumatic stress disorder,
19 stress, and anxiety in members of the UK nursing and midwifery workforce during the first wave of
20 the COVID-19 pandemic. Although prevalence decreased over time, even 3-months after the first
21 surge, almost three in ten participants reported probable post-traumatic stress disorder. Our
22 findings showed that both personal and workplace factors were associated with probable post-
23 traumatic stress disorder, and may have acted as triggers. Key similarities with previous studies
24 highlight the global relevance of our study findings.
25
26

27
28 Our findings should drive healthcare employers throughout the world to address shortcomings in
29 their organisational response to COVID-19, and future health emergencies, to both prevent
30 psychological issues developing and ensure that those affected receive timely high-quality
31 evidence-based support during the pandemic and as needed thereafter.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table and figure legends

Figure one: Survey time-points in context of number of hospitalised cases across UK

Figure two: study flow chart

Table one: baseline characteristics

Table two: Outcome data

Table three: multivariable model of association between explanatory variables and impact of events scale-revised score \geq 33

Supplementary materials

Table e1: summary of all survey questions, time-point of collection, question purpose, and inclusion in imputation/ modelling

Figure e1: Survey time-points in context of number of hospitalised cases across English regions, Scotland, Wales and Northern Ireland

Table e2: Summary of outcome measures

Table e3: Additional baseline characteristics

Table e4: additional outcome data

Table e5: Cronbach's α for outcome measures- page 14

Table e6: DASS-21 and Impact of Events Scale-revised subscale correlations (Pearson correlation coefficient)- page 15

Table e7: Impact of events scale-revised outcomes by response pattern- page 16

Table e8: DASS-21 outcomes by response pattern- page 17

Table e9: Multivariable models controlling for pattern of response- page 18

Funding

This work was supported by funding from The Burdett Trust for Nursing. The Burdett Trust for Nursing had no role in the design of the study, data collection, data analysis and interpretation, writing of the manuscript, or decision to publish.

RMT is a National Institute for Health Research (NIHR) Senior Nurse Research Leader, and receives funding through the NIHR PGfAR, HS&DR and UCLH Charity.

The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

Acknowledgements

We are grateful to the Royal College of Nursing, Nursing and Midwifery Council, Council of Deans for Health, Florence Nightingale Foundation, Cavell Nurses' Trust and all other individuals and organisations that supported survey dissemination.

We acknowledge the CERA (COVID-19 emergency response assessment) study team whose survey design informed our study.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Conflicts of interest:

None

Data availability:

The datasets generated during and/or analysed during the current study may be made available by the corresponding author on reasonable request.

References


- 1
2
3
4 Aiken, L.H., Sermeus, W., Van den Heede, K., Sloane, D.M., Busse, R., McKee, M., Bruyneel, L., Rafferty, A.M.,
5 Griffiths, P., Moreno-Casbas, M.T., Tishelman, C., Scott, A., Brzostek, T., Kinnunen, J., Schwendimann, R.,
6 Heinen, M., Zikos, D., Sjetne, I.S., Smith, H.L., Kutney-Lee, A., 2012. Patient safety, satisfaction, and quality of
7 hospital care: cross sectional surveys of nurses and patients in 12 countries in Europe and the United States.
8 *BMJ* 344, e1717.
- 9 Ali, S., Maguire, S., Marks, E., Doyle, M., Sheehy, C., 2020. Psychological impact of the COVID-19 pandemic on
10 healthcare workers at acute hospital settings in the South-East of Ireland: an observational cohort multicentre
11 study. *BMJ Open* 10 (12), e042930.
- 12 Allan, S.M., Bealey, R., Birch, J., Cushing, T., Parke, S., Sergi, G., Bloomfield, M., Meiser-Stedman, R., 2020. The
13 prevalence of common and stress-related mental health disorders in healthcare workers based in pandemic-
14 affected hospitals: a rapid systematic review and meta-analysis. *European Journal of Psychotraumatology* 11
15 (1), 1810903.
- 16 Antony, M.M., Bieling, P.J., Cox, B.J., Enns, M.W., Swinson, R.P., 1998. Psychometric properties of the 42-item and 21-
17 item versions of the Depression Anxiety Stress Scales in clinical groups and a community sample.
18 *Psychological Assessment* 10 (2), 176-181.
- 19 Bandyopadhyay, S., Baticulon, R.E., Kadhum, M., Alser, M., Ojuka, D.K., Badereddin, Y., Kamath, A., Parepalli, S.A.,
20 Brown, G., Iharchane, S., Gandino, S., Markovic-Obiago, Z., Scott, S., Manirambona, E., Machhada, A.,
21 Aggarwal, A., Benazaize, L., Ibrahim, M., Kim, D., Tol, I., Taylor, E.H., Knighton, A., Bbaale, D., Jasim, D.,
22 Alghoul, H., Reddy, H., Abuelgasim, H., Saini, K., Sigler, A., Abuelgasim, L., Moran-Romero, M., Kumarendran,
23 M., Jamie, N.A., Ali, O., Sudarshan, R., Dean, R., Kisyova, R., Kelzang, S., Roche, S., Ahsan, T., Mohamed, Y.,
24 Dube, A.M., Gwini, G.P., Gwokyaala, R., Brown, R., Papon, M.R.K.K., Li, Z., Ruzats, S.S., Charuvila, S., Peter,
25 N., Khalidy, K., Moyo, N., Alser, O., Solano, A., Robles-Perez, E., Tariq, A., Gaddah, M., Kolovos, S.,
26 Muchemwa, F.C., Saleh, A., Gosman, A., Pinedo-Villanueva, R., Jani, A., Khundkar, R., 2020. Infection and
27 mortality of healthcare workers worldwide from COVID-19: a systematic review. *BMJ Global Health* 5 (12),
28 e003097.
- 29 BBC News, 2020. Coronavirus: Nurses' leaders urge 'care for those who caring'. [Available from:
30 <https://www.bbc.co.uk/news/health-53101331>, accessed 8/02/2021]
- 31 Bilinski, A., Emanuel, E.J., 2020. COVID-19 and Excess All-Cause Mortality in the US and 18 Comparison Countries.
32 *JAMA* 324 (20), 2100-2102.
- 33 Cabarkapa, S., Nadjidai, S.E., Murgier, J., Ng, C.H., 2020. The psychological impact of COVID-19 and other viral
34 epidemics on frontline healthcare workers and ways to address it: A rapid systematic review. *Brain, behavior, &*
35 *immunity - health* 8, 100144-100144.
- 36 Chen, Y.-C., Guo, Y.-L.L., Chin, W.-S., Cheng, N.-Y., Ho, J.-J., Shiao, J.S.-C., 2019. Patient-Nurse Ratio is Related to
37 Nurses' Intention to Leave Their Job through Mediating Factors of Burnout and Job Dissatisfaction. *International*
38 *journal of environmental research and public health* 16 (23), 4801.
- 39 Chew, N.W.S., Lee, G.K.H., Tan, B.Y.Q., Jing, M., Goh, Y., Ngiam, N.J.H., Yeo, L.L.L., Ahmad, A., Ahmed Khan, F.,
40 Napoleon Shanmugam, G., Sharma, A.K., Komalkumar, R.N., Meenakshi, P.V., Shah, K., Patel, B., Chan,
41 B.P.L., Sunny, S., Chandra, B., Ong, J.J.Y., Paliwal, P.R., Wong, L.Y.H., Sagayanathan, R., Chen, J.T., Ying
42 Ng, A.Y., Teoh, H.L., Tsvigoulis, G., Ho, C.S., Ho, R.C., Sharma, V.K., 2020. A multinational, multicentre study
43 on the psychological outcomes and associated physical symptoms amongst healthcare workers during COVID-
44 19 outbreak. *Brain Behav Immun* 88, 559-565.
- 45 Chou, R., Dana, T., Buckley, D.I., Selph, S., Fu, R., Totten, A.M., 2020. Epidemiology of and Risk Factors for
46 Coronavirus Infection in Health Care Workers: A Living Rapid Review. *Annals of Internal Medicine* 173 (2), 120-
47 136.
- 48 Creamer, M., Bell, R., Failla, S., 2003. Psychometric properties of the Impact of Event Scale—Revised. *Behaviour*
49 *Research and Therapy* 41 (12), 1489-1496.
- 50 De Kock, J.H., Latham, H.A., Leslie, S.J., Grindle, M., Munoz, S.-A., Ellis, L., Polson, R., O'Malley, C.M., 2021. A rapid
51 review of the impact of COVID-19 on the mental health of healthcare workers: implications for supporting
52 psychological well-being. *BMC Public Health* 21 (1), 104.
- 53 Douglas, M., Katikireddi, S.V., Taulbut, M., McKee, M., McCartney, G., 2020. Mitigating the wider health effects of covid-
54 19 pandemic response. *BMJ* 369, m1557.
- 55 Dykes, N., Johnson, O., Bamford, P., 2021. Assessing the psychological impact of COVID-19 on intensive care workers:
56 A single-centre cross-sectional UK-based study. *Journal of the Intensive Care Society*. Epub ahead of print
57 January 5, 2021; <https://doi.org/10.1177/1751143720983182>.
- 58 Dyrbye, L.N., Shanafelt, T.D., Johnson, P.O., Johnson, L.A., Satele, D., West, C.P., 2019. A cross-sectional study
59 exploring the relationship between burnout, absenteeism, and job performance among American nurses. *BMC*
60 *Nursing* 18 (1), 57.
- 61 Eysenbach, G., 2004. Improving the Quality of Web Surveys: The Checklist for Reporting Results of Internet E-Surveys
62 (CHERRIES). *J Med Internet Res* 6 (3), e34.
- 63 Fernandez, R., Lord, H., Halcomb, E., Moxham, L., Middleton, R., Alananzeh, I., Ellwood, L., 2020. Implications for
64 COVID-19: A systematic review of nurses' experiences of working in acute care hospital settings during a
65 respiratory pandemic. *International Journal of Nursing Studies* 111, 103637.

- Ferrando-Vivas, P., Doidge, J., Thomas, K., Gould, D.W., Mouncey, P., Shankar-Hari, M., Young, J.D., Rowan, K.M., Harrison, D.A., on behalf of the, I.C.-T., 2021. Prognostic Factors for 30-Day Mortality in Critically Ill Patients With Coronavirus Disease 2019: An Observational Cohort Study. *Critical Care Medicine* 49 (1).
- Greenhalgh, T., Knight, M., A'Court, C., Buxton, M., Husain, L., 2020. Management of post-acute covid-19 in primary care. *BMJ* 370, m3026.
- Hackett, K., 2020. Nursing Standard: COVID-19: redeployment training is non-existent or inadequate, nurses reveal. [Available from: <https://rcni.com/nursing-standard/newsroom/news/covid-19-redeployment-training-non-existent-or-inadequate-nurses-reveal-160301> accessed 8/2/21]
- Hall, L.H., Johnson, J., Watt, I., Tsipa, A., O'Connor, D.B., 2016. Healthcare Staff Wellbeing, Burnout, and Patient Safety: A Systematic Review. *PLOS ONE* 11 (7), e0159015.
- Holmes, E.A., O'Connor, R.C., Perry, V.H., Tracey, I., Wessely, S., Arseneault, L., Ballard, C., Christensen, H., Cohen Silver, R., Everall, I., Ford, T., John, A., Kabir, T., King, K., Madan, I., Michie, S., Przybylski, A.K., Shafran, R., Sweeney, A., Worthman, C.M., Yardley, L., Cowan, K., Cope, C., Hotopf, M., Bullmore, E., 2020. Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science. *The Lancet Psychiatry* 7 (6), 547-560.
- Jun, J., Ojemeni, M.M., Kalamani, R., Tong, J., Crecelius, M.L., 2021. Relationship between nurse burnout, patient and organizational outcomes: Systematic review. *International Journal of Nursing Studies* 119, 103933.
- Kinman, G., Teoh, K., Harriss, A., 2020. The Mental Health and Wellbeing of Nurses and Midwives in the United Kingdom. [Available from: https://www.som.org.uk/sites/som.org.uk/files/The_Mental_Health_and_Wellbeing_of_Nurses_and_Midwives_in_the_United_Kingdom.pdf, accessed 19/04/2021]
- Kisely, S., Warren, N., McMahon, L., Dalais, C., Henry, I., Siskind, D., 2020. Occurrence, prevention, and management of the psychological effects of emerging virus outbreaks on healthcare workers: rapid review and meta-analysis. *BMJ* 369, m1642.
- Lai, J., Ma, S., Wang, Y., Cai, Z., Hu, J., Wei, N., Wu, J., Du, H., Chen, T., Li, R., Tan, H., Kang, L., Yao, L., Huang, M., Wang, H., Wang, G., Liu, Z., Hu, S., 2020. Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. *JAMA Network Open* 3 (3), e203976-e203976.
- Lancaster, C.L., Teeters, J.B., Gros, D.F., Back, S.E., 2016. Posttraumatic Stress Disorder: Overview of Evidence-Based Assessment and Treatment. *Journal of clinical medicine* 5 (11), 105.
- Liu, Q., Luo, D., Haase, J.E., Guo, Q., Wang, X.Q., Liu, S., Xia, L., Liu, Z., Yang, J., Yang, B.X., 2020. The experiences of health-care providers during the COVID-19 crisis in China: a qualitative study. *The Lancet Global Health* 8 (6), e790-e798.
- Lovibond, S.H., Lovibond, P.F., 1996. Manual for the depression anxiety stress scales. Psychology Foundation of Australia, Sydney.
- Maben, J., Bridges, J., 2020. Covid-19: Supporting nurses' psychological and mental health. *Journal of Clinical Nursing* 29 (15-16), 2742-2750.
- Manning, J.C., Bramley, L., Coad, J., Evans, C., Evans, K., Tinkler, L., Cooper, J., 2021. Priorities for research during the Coronavirus SARS-CoV-2 (COVID-19) pandemic and beyond: a survey of nurses, midwives and health visitors in the United Kingdom. *Journal of Research in Nursing* 26 (5), 442-454.
- Maslach, C., Jackson, S.E., Leiter, M.P., 2018. Maslach Burnout Inventory Manual. Mind Garden, Menlo Park, CA.
- McAlonan, G.M., Lee, A.M., Cheung, V., Cheung, C., Tsang, K.W., Sham, P.C., Chua, S.E., Wong, J.G., 2007. Immediate and sustained psychological impact of an emerging infectious disease outbreak on health care workers. *Can J Psychiatry* 52 (4), 241-247.
- Mitchell, G., 2020. Nursing Times: Nurses asked to share experiences of Covid-19 peak for major study. [Available from: <https://www.nursingtimes.net/news/workforce/nurses-asked-to-share-experiences-of-covid-19-peak-for-major-study-29-04-2020/> accessed 24/03/21].
- Mutambudzi, M., Niedwiedz, C., Macdonald, E.B., Leyland, A., Mair, F., Anderson, J., Celis-Morales, C., Cleland, J., Forbes, J., Gill, J., Hastie, C., Ho, F., Jani, B., Mackay, D.F., Nicholl, B., O'Donnell, C., Sattar, N., Welsh, P., Pell, J.P., Katikireddi, S.V., Demou, E., 2020. Occupation and risk of severe COVID-19: prospective cohort study of 120 075 UK Biobank participants. *Occupational and Environmental Medicine*, oemed-2020-106731.
- National Audit Office, 2020. The NHS nursing workforce. [Available from: [9/02/2021]. Available from: <https://www.nao.org.uk/wp-content/uploads/2020/03/The-NHS-nursing-workforce.pdf> accessed 9/02/21].
- National Health Service, 2019. The NHS Long Term Plan. [Available from: <https://www.longtermplan.nhs.uk/wp-content/uploads/2019/01/nhs-long-term-plan.pdf> accessed 31/01/2019].
- Navaratnam, A.V., Gray, W.K., Day, J., Wendon, J., Briggs, T.W.R., Patient factors and temporal trends associated with COVID-19 in-hospital mortality in England: an observational study using administrative data. *The Lancet Respiratory Medicine*. Epub ahead of print February 15, 2021; [https://doi.org/10.1016/S2213-2600\(20\)30579-8](https://doi.org/10.1016/S2213-2600(20)30579-8).
- Nguyen, L.H., Drew, D.A., Graham, M.S., Joshi, A.D., Guo, C.-G., Ma, W., Mehta, R.S., Warner, E.T., Sikavi, D.R., Lo, C.-H., Kwon, S., Song, M., Mucci, L.A., Stampfer, M.J., Willett, W.C., Eliassen, A.H., Hart, J.E., Chavarro, J.E., Rich-Edwards, J.W., Davies, R., Capdevila, J., Lee, K.A., Lochlainn, M.N., Varsavsky, T., Sudre, C.H., Cardoso, M.J., Wolf, J., Spector, T.D., Ourselin, S., Steves, C.J., Chan, A.T., Albert, C.M., Andreotti, G., Bala, B., Balasubramanian, B.A., Beane-Freeman, L.E., Brownstein, J.S., Bruinsma, F.J., Coresh, J., Costa, R., Cowan, A.N., Deka, A., Deming-Halverson, S.L., Elena Martinez, M., Ernst, M.E., Figueiredo, J.C., Fortuna, P., Franks, P.W., Freeman, L.B., Gardner, C.D., Ghobrial, I.M., Haiman, C.A., Hall, J.E., Kang, J.H., Kirpach, B., Koenen, K.C., Kubzansky, L.D., Lacey, Jr, J.V., Le Marchand, L., Lin, X., Lutsey, P., Marinac, C.R., Martinez, M.E., Milne, R.L., Murray, A.M., Nash, D., Palmer, J.R., Patel, A.V., Pierce, E., Robertson, M.M., Rosenberg, L., Sandler, D.P., Schurman, S.H., Sewalk, K., Sharma, S.V., Sidey-Gibbons, C.J., Slevin, L., Smoller, J.W., Steves, C.J., Tiirikainen, M.I., Weiss, S.T., Wilkens, L.R., Zhang, F., 2020. Risk of COVID-19 among front-line

health-care workers and the general community: a prospective cohort study. *The Lancet Public Health* 5 (9), e475-e483.

- 1 NHS Digital, 2018. Nurses, Midwives and Support staff by area, level, gender and age. [Available from:
2 [https://digital.nhs.uk/data-and-information/find-data-and-publications/supplementary-information/2018-](https://digital.nhs.uk/data-and-information/find-data-and-publications/supplementary-information/2018-supplementary-information-files/staff-numbers/nurses-midwives-and-support-staff-by-area-level-gender-and-age-january-18)
3 [supplementary-information-files/staff-numbers/nurses-midwives-and-support-staff-by-area-level-gender-and-](https://digital.nhs.uk/data-and-information/find-data-and-publications/supplementary-information/2018-supplementary-information-files/staff-numbers/nurses-midwives-and-support-staff-by-area-level-gender-and-age-january-18)
4 [age-january-18](https://digital.nhs.uk/data-and-information/find-data-and-publications/supplementary-information/2018-supplementary-information-files/staff-numbers/nurses-midwives-and-support-staff-by-area-level-gender-and-age-january-18) accessed 13/09/2021].
- 5 Nickell, L.A., Crighton, E.J., Tracy, C.S., Al-Enazy, H., Bolaji, Y., Hanjrah, S., Hussain, A., Makhlof, S., Upshur, R.E.G.,
6 2004. Psychosocial effects of SARS on hospital staff: survey of a large tertiary care institution. *Canadian*
7 *Medical Association Journal* 170 (5), 793-798.
- 8 Nursing and Midwifery Council, 2021. The NMC register: 1 April 2020 – 31 March 2021. [Available from:
9 [https://www.nmc.org.uk/globalassets/sitedocuments/data-reports/annual-2021/0005b-nmc-register-2021-](https://www.nmc.org.uk/globalassets/sitedocuments/data-reports/annual-2021/0005b-nmc-register-2021-web.pdf)
10 [web.pdf](https://www.nmc.org.uk/globalassets/sitedocuments/data-reports/annual-2021/0005b-nmc-register-2021-web.pdf) accessed 16/08/2021].
- 11 Plumpton, C.O., Morris, T., Hughes, D.A., White, I.R., 2016. Multiple imputation of multiple multi-item scales when a full
12 imputation model is infeasible. *BMC Research Notes* 9 (1), 45.
- 13 Poghosyan, L., Aiken, L.H., Sloane, D.M., 2009. Factor structure of the Maslach burnout inventory: An analysis of data
14 from large scale cross-sectional surveys of nurses from eight countries. *International Journal of Nursing Studies*
15 46 (7), 894-902.
- 16 Roberts, T., Daniels, J., Hulme, W., Hirst, R., Horner, D., Lyttle, M.D., Samuel, K., Graham, B., Reynard, C., Barrett, M.,
17 Foley, J., Cronin, J., Umana, E., Carlton, E., Vinagre, J., 2021. Psychological Distress and Trauma in Doctors
18 Providing Frontline Care During the COVID-19 Pandemic in the United Kingdom and Ireland: A Prospective
19 Longitudinal Survey Cohort Study. [Available from:
20 https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3760472 accessed 14th April 2021].
- 21 Roberts, T., Daniels, J., Hulme, W., Hirst, R., Horner, D., Lyttle, M.D., Samuel, K., Graham, B., Reynard, C., Barrett, M.,
22 Foley, J., Cronin, J., Umana, E., Vinagre, J., Carlton, E., On behalf of the collaborators of TERN, R., PERUKI,
23 ITERN, TRIC, and SATARN, 2021. Psychological distress during the acceleration phase of the COVID-19
24 pandemic: a survey of doctors practising in emergency medicine, anaesthesia and intensive care medicine in
25 the UK and Ireland. *Emerg Med J*. Epub ahead of print April 10, 2021; [https://doi.org/10.1136/emermed-2020-](https://doi.org/10.1136/emermed-2020-210438)
26 210438.
- 27 Roberts, T., Daniels, J., Hulme, W., Horner, D., Lyttle, M.D., Samuel, K., Graham, B., Hirst, R., Reynard, C., Barrett, M.,
28 Carlton, E., 2020. COVID-19 emergency response assessment study: a prospective longitudinal survey of
29 frontline doctors in the UK and Ireland: study protocol. *BMJ Open* 10 (8), e039851.
- 30 Royston, P., White, I.R., 2011. Multiple Imputation by Chained Equations (MICE): Implementation in Stata. 2011 45 (4),
31 20.
- 32 Shah, A.S.V., Wood, R., Gribben, C., Caldwell, D., Bishop, J., Weir, A., Kennedy, S., Reid, M., Smith-Palmer, A.,
33 Goldberg, D., McMenamin, J., Fischbacher, C., Robertson, C., Hutchinson, S., McKeigue, P., Colhoun, H.,
34 McAllister, D.A., 2020. Risk of hospital admission with coronavirus disease 2019 in healthcare workers and their
35 households: nationwide linkage cohort study. *BMJ* 371, m3582.
- 36 Stuijzand, S., Deforges, C., Sandoz, V., Sajin, C.-T., Jaques, C., Elmers, J., Horsch, A., 2020. Psychological impact of
37 an epidemic/pandemic on the mental health of healthcare professionals: a rapid review. *BMC Public Health* 20
38 (1), 1230.
- 39 Taquet, M., Geddes, J.R., Husain, M., Luciano, S., Harrison, P.J., 2021 6-month neurological and psychiatric outcomes
40 in 236 379 survivors of COVID-19: a retrospective cohort study using electronic health records. *The Lancet*
41 *Psychiatry* 8 (5), 416-27.
- 42 Toyne, P., 2020. The Guardian: It's officially the year of the nurse – what better time to reverse 10 years of bad care?
43 [Available from: [https://www.theguardian.com/commentisfree/2020/apr/24/year-nurse-tories-10-years-bad-care-](https://www.theguardian.com/commentisfree/2020/apr/24/year-nurse-tories-10-years-bad-care-nhs-crisis)
44 [nhs-crisis](https://www.theguardian.com/commentisfree/2020/apr/24/year-nurse-tories-10-years-bad-care-nhs-crisis) accessed 8/2/21]
- 45 UCLA: Statistical Consulting Group, Multiple imputation in stata. [Available from:
46 https://stats.idre.ucla.edu/stata/seminars/mi_in_stata_pt1_new/ accessed 5/3/21].
- 47 UK Government, 2021. Coronavirus (COVID-19) in the UK. [Available from: <https://coronavirus.data.gov.uk/> accessed
48 5/2/21].
- 49 Weiss, D.S., 2006. The Impact of Event Scale- Revised. In: Wilson, J.P., Keane, T.M., Guilford, P. (Eds.), *Assessing*
50 *psychological trauma and PTSD*. The Guilford Press, New York; London.
- 51 Williamson, E.J., Walker, A.J., Bhaskaran, K., Bacon, S., Bates, C., Morton, C.E., Curtis, H.J., Mehrkar, A., Evans, D.,
52 Inglesby, P., Cockburn, J., McDonald, H.I., MacKenna, B., Tomlinson, L., Douglas, I.J., Rentsch, C.T., Mathur,
53 R., Wong, A.Y.S., Grieve, R., Harrison, D., Forbes, H., Schultze, A., Croker, R., Parry, J., Hester, F., Harper, S.,
54 Perera, R., Evans, S.J.W., Smeeth, L., Goldacre, B., 2020. Factors associated with COVID-19-related death
55 using OpenSAFELY. *Nature* 584 (7821), 430-436.
- 56 World Health Organization, 2020. State of the world's nursing 2020: investing in education, jobs and leadership.
57 [Available from: <https://apps.who.int/iris/rest/bitstreams/1274201/retrieve> accessed 08/02/2020]
- 58 World Health Organization, 2020. Year of the Nurse and the Midwife 2020. [Available from:
59 <https://www.who.int/campaigns/year-of-the-nurse-and-the-midwife-2020> accessed 08/02/2021]
- 60 Wu, P., Fang, Y., Guan, Z., Fan, B., Kong, J., Yao, Z., Liu, X., Fuller, C.J., Susser, E., Lu, J., Hoven, C.W., 2009. The
61 Psychological Impact of the SARS Epidemic on Hospital Employees in China: Exposure, Risk Perception, and
62 Altruistic Acceptance of Risk. *The Canadian Journal of Psychiatry* 54 (5), 302-311.
- 63 Zhu, Z., Xu, S., Wang, H., Liu, Z., Wu, J., Li, G., Miao, J., Zhang, C., Yang, Y., Sun, W., Zhu, S., Fan, Y., Chen, Y., Hu,
64 J., Liu, J., Wang, W., 2020. COVID-19 in Wuhan: Sociodemographic characteristics and hospital support
65 measures associated with the immediate psychological impact on healthcare workers. *EClinicalMedicine* 24,
100443.

Figure one






Table one: baseline characteristics

	T1 (n=2040) n (%)	T2 (n=3638) n (%)	T3 (n=2162) n (%)	P-value
PERSONAL FACTORS†				
Gender (mlogit)	F(4, 4.5e+08) = 1.50			
Male	160 (7.9)	271 (7.5)	166 (7.7)	
Female	1868 (91.6)	3331 (91.6)	1986 (91.9)	
Other/prefer not to say	12 (0.6)	36 (1.0)	10 (0.5)	p = 0.20
Age (years) (oreg)	F(2, 1.6e+07) = 39.48			
16-30	265 (13.0)	381 (10.5)	158 (7.3)	
31-40	406 (19.9)	660 (18.1)	321 (14.9)	
41-50	600 (29.4)	1061 (29.2)	620 (28.7)	
51-60	626 (30.7)	1263 (34.7)	860 (39.8)	
61 and over	143 (7.0)	273 (7.5)	204 (9.4)	p<0.001
Ethnicity (mlogit)	F(3,336836.6) = 9.43			
White		3248 (89.3)	2012 (93.1)	
Mixed/Multiple ethnic		55 (1.5)	35 (1.6)	
Asian/Asian British		126 (3.5)	39 (1.8)	
Black/African/Caribbean		210 (5.8)	76 (3.5)	p<0.001
UK Region (mlogit)	F(20, 1.8e+08) = 4.58			
South East	322 (15.8)	562 (15.4)	326 (15.1)	
South West	220 (10.8)	383 (10.5)	234 (10.8)	
London	227 (11.1)	526 (14.4)	297 (13.7)	
East Midlands	214 (10.5)	267 (7.3)	173 (8.0)	
East of England	146 (7.2)	276 (7.6)	191 (8.9)	
North West	262 (12.9)	380 (10.4)	197 (9.1)	
North East (including Yorkshire)	126 (6.2)	340 (9.4)	227 (10.5)	
West Midlands	228 (11.2)	311 (8.6)	186 (8.6)	
Scotland	168 (8.3)	345 (9.5)	187 (8.7)	
Wales	86 (4.2)	154 (4.2)	108 (5.0)	
Northern Ireland	39 (1.9)	96 (2.6)	35 (1.6)	p<0.001
Professional qualification (mlogit)	F(12, 4.8e+07) = 2.84			
RN - adult	1589 (77.9)	2732 (75.1)	1730 (80.0)	
RN - child	109 (5.4)	211 (5.8)	111 (5.1)	
RN- learning disability	15 (0.7)	52 (1.4)	21 (1.0)	
RN- mental health	132 (6.5)	284 (7.8)	116 (5.4)	
Midwife/Health visitor	58 (2.9)	96 (2.6)	69 (3.2)	
HCA/ Nursing associate/ trainee	79 (3.9)	140 (3.8)	68 (3.2)	
Student nurse/ midwife	58 (2.8)	122 (3.4)	46 (2.1)	p<0.001
Caring responsibilities (mlogit)	F(4, 9.0e+07) = 2.99			
Yes- sole carer	223 (11.0)	407 (11.2)	205 (9.5)	
Yes, but not sole carer	774 (38.0)	1270 (34.9)	754 (34.9)	
No	1042 (51.1)	1961 (53.9)	1203 (55.7)	p = 0.018
WORKPLACE FACTORS†				
Health at risk due to clinical role (oreg)	F(1,15516.5) = 31.77			
Strongly disagree/disagree	247 (12.1)	568 (15.6)		
Neither agree nor disagree	279 (13.7)	642 (17.6)		
Strongly agree/agree	1514 (74.2)	2429 (66.8)		p<0.001
Personal COVID-19 infection	-			
Yes - I had symptoms			628 (29.0)	
Yes - did not have symptoms			113 (5.2)	
No			1421 (65.7)	-
Employed by NHS (lreg)	F(2, 2.3e+08) = 0.31			
No	470 (23.0)	830 (22.8)	512 (23.7)	
Yes	1570 (77.0)	2808 (77.2)	1650 (76.3)	p=0.74
NHS pay banding (or equivalent) (mlogit)	F(2, 1.5e+06) = 14.39			
Not banded	38 (1.9)	51 (1.4)	20 (0.9)	
HCA/ nursing associate	103 (5.1)	219 (6.0)	99 (4.6)	

	Staff nurse	567 (27.8)	1106 (30.4)	558 (25.8)	
	Senior staff nurse	555 (27.2)	1013 (27.9)	620 (28.7)	
	Charge nurse	446 (21.9)	780 (21.5)	547 (25.3)	
	Matron	187 (9.1)	281 (7.7)	184 (8.5)	
	Senior nurse	81 (4.0)	100 (2.7)	71 (3.3)	
	Very senior nurse	64 (3.1)	88 (2.4)	62 (2.9)	p<0.001
Clinical speciality during pandemic ^(mlogit)		F(22, 1.4e+07) = 2.44			
	Acute/emergency hospital care	547 (26.8)	935 (25.7)	511 (23.7)	
	Critical care/operating theatres	324 (15.9)	523 (14.4)	338 (15.6)	
	Learning disabilities	12 (0.6)	26 (0.7)	14 (0.7)	
	Mental health	102 (5.0)	253 (7.0)	95 (4.4)	
	Midwifery/Paediatrics	77 (3.8)	129 (3.5)	88 (4.1)	
	Palliative care	49 (2.4)	116 (3.2)	64 (3.0)	
	Outpatients	49 (2.4)	98 (2.7)	67 (3.1)	
	Care/nursing Home	49 (2.4)	119 (3.3)	60 (2.8)	
	Community or primary care	418 (20.5)	742 (20.4)	477 (22.1)	
	Education/ higher education	41 (2.0)	64 (1.8)	46 (2.1)	
	Research	79 (3.9)	131 (3.6)	123 (5.7)	
	Other	292 (14.3)	502 (13.8)	276 (12.8)	p<0.001
Redeployment ^(mlogit)		F(4, 5.7e+06) = 7.17			
	Not redeployed	1361 (66.8)	2643 (72.7)	1579 (73.1)	
	Redeployed - adequate training	259 (12.7)	368 (10.1)	202 (9.4)	
	Redeployed - inadequate or no training	419 (20.5)	624 (17.2)	380 (17.6)	p<0.001
Previously provided clinical care in pandemic ^(lreg)		F(2, 1.7e+07) = 1.92			
	Yes	520 (25.5)	963 (26.5)	608 (28.1)	
	No	1520 (74.5)	2675 (73.5)	1554 (71.9)	p=0.15
Main role during the pandemic		-			
	Direct clinical care			1533 (70.9)	
	Education			78 (3.6)	
	Research			140 (6.5)	
	Management/leadership			236 (10.9)	
	Unable to work			63 (2.9)	
	Other			112 (5.2)	-
Correct PPE always available ^(oreg)		F(2, 1.1e+06) = 199.97			
	Strongly agree/agree	795 (39.0)	2192 (60.3)	1468 (67.9)	
	Neither agree or disagree	522 (25.6)	782 (21.5)	311 (14.4)	
	Disagree/strongly disagree	724 (35.5)	664 (18.3)	384 (17.8)	p<0.001
Confidence in COVID-19 IPC training ^(mlogit)		F(2, 3.8e+06) = 72.76			
	Not confident at all/some not confident	858 (42.1)	1103 (30.3)	532 (24.6)	
	Neither not confident nor confident	346 (16.9)	634 (17.4)	295 (13.7)	
	Somewhat confident/very confident	632 (31.0)	1648 (45.3)	1152 (53.3)	
	Not received training	204 (10.0)	254 (7.0)	182 (8.4)	p<0.001
Prepared to provide care to COVID-19 patients ^(oreg)		F(2, 2.2e+06) = 75.79			
	Completely/somewhat unprepared	1096 (53.7)	1348 (37.1)	916 (42.4)	
	Neither prepared nor unprepared	237 (11.6)	474 (13.0)	292 (13.5)	
	Somewhat/very prepared	707 (34.7)	1816 (49.9)	954 (44.1)	p<0.001
Care quality for non-COVID-19 ^(oreg)		F(2,57974.0) = 35.09			
	Significantly/slightly worse than before	1007 (49.3)	1447 (39.8)	1078 (49.9)	
	Same as before	865 (42.4)	1777 (48.8)	867 (40.1)	
	Significantly/slightly better than before	169 (8.3)	414 (11.4)	217 (10.1)	p<0.001
Mental health first aid training ^(lreg)		F(2, 5.9e+07) = 0.47			
	Yes	396 (19.4)	729 (20.0)	446 (20.6)	
	No	1644 (80.6)	2909 (80.0)	1716 (79.4)	p=0.63

HCA- healthcare assistant; IPC- infection prevention and control; PPE- personal protective equipment; RN- registered nurse

All variables in table based on imputed dataset.

All variables collected at each of the three time-points except: ethnicity (not collected at time-point one); health at risk due to clinical role (not collected at time-point three); personal COVID-19 infection (not collected at time-points one/ two); main role during pandemic (not collected at time-points one/ two).

Statistical test obtained using: lreg = logistic regression, oreg = ordinal regression, mlogit = multinomial regression, PChi = Pearson Chi-squared †- The test statistic (where applicable) is shown adjacent to the variable name.

Table two: Outcome data

		T1 (n=2040)	T2 (n=3638)	T3 (n=2162)	Test statistic/ p-value
Impact of events scale-revised					
	Score cut-offs- n (%)				
	≥ 33 (probable PTSD)	909(44.6)	1350(37.1)	634(29.3)	<0.001†
	≥ 24 (PTSD clinical concern)	1227 (60.2)	1835 (50.4)	897 (41.5)	<0.001†
	Total score- mean (95% CI)	32.16 (31.25-33.07)	28.11 (27.43-28.79)	23.81 (22.91-24.71)	F(2, 7521.1) = 81.95 p<0.001‡
DASS-21- n(%)					
	Depression- severe/ extremely severe	402 (19.7)	727 (20.0)	424 (19.6)	F(2,124794.0) = 1.38, p = 0.25*,#
	Anxiety- severe/ extremely severe	509 (25.0)	792 (21.7)	392 (18.1)	F(2,303171.4) = 42.58, p<0.001*,#
	Stress- severe/ extremely severe	421 (20.6)	660 (18.1)	379 (17.5)	F(2,223990.2) = 9.30, p<0.001*, #
Satisfaction with job- n(%)					
	Very satisfied	598 (29.3)	991 (27.2)	395 (18.3)	
	Somewhat satisfied	819 (40.1)	1475 (40.6)	841 (38.9)	
	Average	349 (17.1)	667 (18.3)	431 (19.9)	
	Somewhat unsatisfied	176 (8.6)	333 (9.2)	323 (14.9)	
	Very unsatisfied	98 (4.8)	171 (4.7)	172 (8.0)	F(2, 4.3e+07) = 68.10, p<0.001
Maslach burnout inventory- mean (95% CI)					
	Emotional exhaustion			26.48 (25.89-27.07)	
	Depersonalization			5.52 (5.27-5.78)	
	Personal accomplishment			34.64 (34.25-35.02)	
Level of intention to leave					
	Current job- mean (95% CI)			37.8 (36.3-39.4)	
	Healthcare- mean (95% CI)			28.6 (27.2-30.0)	
<p>*- p-value based on comparison across five categories: Normal; Mild; Moderate; Severe; Extremely severe CI- confidence interval; PTSD- post-traumatic stress disorder Statistical test obtained from † logistic regression model ‡ regression model # ordinal regression model</p>					

Table three: multivariable model of association between explanatory variables and probable PTSD

	T1		T2		T3	
	Odds ratio (95% CI)	P- value	Odds ratio (95% CI)	P- value	Odds ratio (95% CI)	P- value
PERSONAL FACTORS						
Gender	F(2, 1.7e+07) = 2.09, p=0.12		F(2, 1.9e+06) = 1.83, p=0.16		F(2,498519.7)=1.38, p=0.25	
Male (i)	1.00		1.00		1.00	
Female	1.41 (0.98-2.04)	0.06	1.33 (0.99-1.77)	0.056	0.75 (0.50-1.06)	0.10
Other/prefer not to say	0.79 (0.21-2.96)	0.73	1.29 (0.59-2.82)	0.52	0.90 (0.21-3.79)	0.88
Age	F(4, 1.2e+06) = 6.91, p<0.001		F(4, 2.2e+06) = 6.12, p<0.001		F(4,67237.7)=0.89, p=0.47	
16-30 (i)	1.00		1.00		1.00	
31-40	0.83 (0.58-1.20)	0.33	0.89 (0.67-1.19)	0.43	0.69 (0.43-1.09)	0.11
41-50	0.60 (0.42-0.86)	0.006	0.71 (0.53-0.93)	0.015	0.75 (0.49-1.15)	0.19
51-60	0.53 (0.37-0.75)	<0.001	0.67 (0.51-0.88)	0.004	0.74 (0.49-1.12)	0.16
61 and over	0.32 (0.19-0.53)	<0.001	0.43 (0.29-0.63)	<0.001	0.63 (0.38-1.06)	0.083
Ethnicity			F(3, 2208925.4) = 3.98, p=0.008		F(3,37585.0)=0.24, p=0.87	
White (i)			1.00		1.00	
Mixed/Multiple ethnic			1.36 (0.76-2.44)	0.30	0.76 (0.35-1.65)	0.48
Asian/Asian British			1.30 (0.86-1.96)	0.21	1.07 (0.50-2.27)	0.87
Black/African/Caribbean			1.72 (1.24-2.38)	0.001	1.13 (0.64-1.98)	0.68
UK region	F(10, 998003.8) = 1.41, p=0.17		F(10, 2.5e+06) = 1.34, p=0.20		F(10,146571.6)=1.60, p=0.099	
South East (i)	1.00		1.00		1.00	
South West	0.98 (0.67-1.44)	0.91	0.92 (0.69-1.24)	0.60	0.57 (0.37-0.87)	0.010
London	1.28 (0.87-1.89)	0.21	0.85 (0.64-1.12)	0.24	0.89 (0.60-1.30)	0.54
East Midlands	1.56 (1.06-2.31)	0.025	0.80 (0.58-1.12)	0.19	0.94 (0.61-1.46)	0.78
East of England	1.02 (0.66-1.58)	0.92	0.74 (0.53-1.03)	0.077	0.99 (0.64-1.51)	0.95
North West	0.88 (0.61-1.26)	0.48	0.98 (0.73-1.31)	0.90	0.70 (0.46-1.08)	0.11
North East	1.05 (0.67-1.66)	0.82	0.91 (0.67-1.24)	0.55	0.61 (0.40-0.94)	0.024
West Midlands	1.39 (0.95-2.03)	0.09	1.00 (0.73-1.37)	0.99	0.88 (0.58-1.35)	0.57
Scotland	1.07 (0.70-1.63)	0.74	0.98 (0.72-1.33)	0.90	0.72 (0.46-1.13)	0.16
Wales	0.98 (0.58-1.66)	0.94	0.58 (0.38-0.88)	0.010	0.78 (0.47-1.31)	0.35
Northern Ireland	1.72 (0.82-3.62)	0.15	0.65 (0.40-1.07)	0.09	1.66 (0.73-3.77)	0.23
Professional qualification	F(3, 27417.4) = 2.12, p=0.096		F(3, 14277.6) = 0.23, p=0.87		F(3,27274.4)=0.29, p=0.84	
RN-Adult nurse (i)	1.00		1.00		1.00	
RN other/Midwives/HVs	0.65 (0.45-0.94)	0.023	1.07 (0.83-1.38)	0.62	0.99 (0.67-1.48)	0.98
HCA/ nursing associate/ trainee	0.36 (0.07-1.81)	0.21	1.35 (0.42-4.27)	0.61	0.47 (0.09-2.42)	0.37
Student nurse/ midwife	0.43 (0.09-1.99)	0.28	1.50 (0.46-4.88)	0.50	0.59 (0.12-2.93)	0.52

Caring responsibilities	F(2, 301685.9) = 3.84, p=0.021	F(2, 746013.4) = 1.58, p=0.21	F(2,23277.6)=3.97, p=0.019
Yes-sole carer	1.51 (1.08–2.09) 0.015	1.25 (0.98–1.59) 0.076	1.58 (1.11–2.25) 0.012
Yes-but not sole carer	1.26 (1.01–1.58) 0.041	1.05 (0.88–1.24) 0.60	1.26 (0.99–1.59) 0.056
No (i)	1.00	1.00	1.00
WORKPLACE FACTORS			
Health at risk due to clinical role	F(2, 10572.7) = 5.91, p=0.003	F(2, 7886.3) = 16.10, p<0.001	
Strongly disagree/disagree	1.29 (0.86–1.96) 0.22	1.34 (1.01–1.79) 0.040	
Neither agree/ disagree (i)	1.00	1.00	
Strongly agree/agree	1.68 (1.22–2.32) 0.001	1.84 (1.47–2.31) <0.001	
Personal COVID-19 infection			F(2,2434.4)=7.55, p<0.001
Yes – symptomatic			1.58 (1.25–2.00) <0.001
Yes- asymptomatic			1.40 (0.87–2.25) 0.17
No (i)			1.00
Employed by NHS†	1.37 (1.03–1.80) 0.028	1.16 (0.94–1.42) 0.17	0.89 (0.67–1.18) 0.40
NHS banding (or equivalent)	F(7, 108131.0) = 1.32, p=0.24	F(7, ,103269.3) = 1.01, p=0.42	F(7,75401.2)=1.28, p=0.25
Not banded	2.78 (0.57–13.65) 0.21	0.65 (0.19–2.23) 0.49	0.57 (0.10–3.35) 0.54
HCA/ nursing associate	2.80 (0.59–13.39) 0.20	0.89 (0.29–2.75) 0.84	1.60 (0.34–7.60) 0.56
Staff nurse (i)	1.00	1.00	1.00
Senior staff nurse	0.88 (0.67–1.15) 0.35	1.01 (0.83–1.22) 0.95	0.91 (0.69–1.20) 0.50
Charge nurse	0.73 (0.54–0.98) 0.038	0.89 (0.72–1.11) 0.31	0.75 (0.56–1.02) 0.067
Matron	0.70 (0.47–1.05) 0.087	0.72 (0.52–1.00) 0.053	0.71 (0.46–1.09) 0.12
Senior nurse	0.60 (0.34–1.06) 0.078	0.93 (0.56–1.54) 0.78	0.57 (0.29–1.14) 0.11
Very senior nurse	0.71 (0.36–1.39) 0.32	0.65 (0.36–1.17) 0.15	0.57 (0.27–1.19) 0.13
Clinical speciality*	F(11, 586562.0) = 1.31, p=0.021	F(11, 765173.0) = 4.67, p<0.001	F(11, 147539.3)=1.70, p=0.066
Acute/emergency hospital care(i)	1.00	1.00	1.00
Critical care/operating theatres	1.14 (0.79–1.64) 0.50	1.51 (1.18–1.92) 0.001	1.31 (0.94–1.82) 0.11
Learning disabilities	0.83 (0.20–3.43) 0.79	0.54 (0.20–1.41) 0.21	0.67 (0.16–2.85) 0.59
Mental health	0.78 (0.43–1.40) 0.41	0.71 (0.48–1.04) 0.075	0.85 (0.45–1.60) 0.61
Midwifery/Paediatrics	0.76 (0.40–1.44) 0.40	0.62 (0.39–1.01) 0.056	0.47 (0.22–0.99) 0.047
Palliative care	0.98 (0.48–2.01) 0.95	1.02 (0.64–1.61) 0.95	1.18 (0.62–2.22) 0.61
Outpatients	0.85 (0.54–1.36) 0.50	0.73 (0.45–1.18) 0.20	0.78 (0.39–1.53) 0.47
Care/nursing Home	1.12 (0.57–2.22) 0.74	1.64 (1.03–2.60) 0.036	1.39 (0.72–2.69) 0.33
Community or primary care	0.93 (0.68–1.27) 0.65	0.72 (0.57–0.91) 0.006	0.73 (0.52–1.02) 0.065
Education/ higher education	1.27 (0.69–2.32) 0.45	0.82 (0.42–1.57) 0.54	0.58 (0.26–1.31) 0.19
Research	0.48 (0.30–0.76) 0.002	0.71 (0.46–1.09) 0.12	0.73 (0.43–1.23) 0.23
Other	0.90 (0.65–1.26) 0.55	0.70 (0.54–0.90) 0.006	0.90 (0.62–1.30) 0.57
Redeployment	F(2, 379222.6) = 4.09, p=0.017	F(2, 3.1e+06) = 7.76, p<0.001	F(2,26653.7)=1.25, p=0.29
Not redeployed (i)	1.00	1.00	1.00

	Redeployed -adequate training	0.85 (0.62–1.16)	0.31	1.01 (0.78–1.31)	0.93	0.84 (0.56–1.25)	0.39
	Redeployed-inadequate/ no training	1.37 (1.06–1.77)	0.017	1.50 (1.22–1.84)	<0.001	1.17 (0.89–1.55)	0.26
	Previously provided clinical care in pandemic†	1.10 (0.87–1.39)	0.43	1.19 (1.00–1.41)	0.047	1.44 (1.14–1.82)	0.002
	Correct PPE always available	F(2, 148518.3) = 6.84, p=0.001		F(2, 327800.1) = 10.54, p<0.001		F(2,38969.5)=7.42, p<0.001	
	Strongly agree/agree	0.97 (0.74–1.28)	0.84	0.82 (0.67–1.00)	0.048	0.74 (0.54–1.01)	0.056
	Neither agree/ disagree (i)	1.00		1.00		1.00	
	Disagree/strongly disagree	1.49 (1.15–1.93)	<0.001	1.34 (1.07–1.69)	0.012	1.26 (0.88–1.80)	0.20
	Confidence in COVID-19 IPC training	F(3, 264019.8) = 3.98, p=0.008		F(3, 1.1e+06) = 11.45, p<0.001		F(3,63431.9)=3.35, p=0.018	
	Not confident at all/somewhat not confident	1.48 (1.11–1.97)	0.007	1.29 (1.03–1.61)	0.024	1.51 (1.07–2.13)	0.020
	Neither not confident/ confident (i)	1.00		1.00		1.00	
	Somewhat/ very confident	0.96 (0.70–1.32)	0.81	0.72 (0.57–0.90)	0.004	1.09 (0.78–1.52)	0.63
	Not received training	1.23 (0.83–1.82)	0.30	0.84 (0.60–1.17)	0.30	1.65 (1.07–2.53)	0.023
	Prepared to provide care to COVID-19 patients	F(2, 701907.5) = 8.03, p<0.001		F(2, 147563.2) = 13.47, p<0.001		F(2,21028.1)=19.80, p<0.001	
	Completely/somewhat unprepared	1.60 (1.16–2.22)	0.005	1.25 (0.98–1.59)	0.066	1.32 (0.95–1.84)	0.096
	Neither unprepared/ prepared (i)	1.00		1.00		1.00	
	Somewhat/very prepared	1.01 (0.70–1.44)	0.96	0.75 (0.59–0.96)	0.024	0.56 (0.40–0.80)	0.001
	Care quality for non-COVID-19 patients	F(2, 16636.2) = 7.11, p<0.001		F(2, 82544.4) = 0.95, p=0.39		F(2,29086.9)=3.37, p=0.034	
	Significantly/slightly worse	1.47 (1.19–1.82)	<0.001	1.12 (0.95–1.32)	0.17	1.34 (1.07–1.69)	0.012
	Same as before (i)	1.00		1.00		1.00	
	Significantly/slightly better	0.98 (0.67–1.42)	0.91	1.02 (0.79–1.31)	0.88	1.30 (0.90–1.89)	0.16
	Mental health first aid training†	0.86 (0.66–1.12)	0.26	0.87 (0.71–1.07)	0.19	0.73 (0.54–0.97)	0.031
	Pseudo R ²	0.11		0.11		0.13	
	C-Statistic (AUC)	0.72		0.72		0.74	

Note: the p-value for the overall effect of variables with three or more categories is shown in brackets in each row containing the variable name

i- Index category. Where index category is not specified, the index category is a negative response

† yes, compared with no

*- Clinical speciality on 1st February 2020 used for time-point one; Clinical speciality during pandemic used for time-points two and three.

AUC- Area Under Curve; CI- confidence interval; IPC- Infection prevention and control; HCA- Healthcare assistant; HV- Health visitor; NHS- National health service; PPE- personal protective equipment; RN- Registered nurse