

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/148986/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Aggleton, John P. and O'Mara, Shane M. 2022. The anterior thalamic nuclei: core components of a tripartite episodic memory system. *Nature Reviews Neuroscience* 23 , pp. 505-516. 10.1038/s41583-022-00591-8

Publishers page: <https://doi.org/10.1038/s41583-022-00591-8>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



The anterior thalamic nuclei: core components of a tripartite episodic memory system

John P. Aggleton^{1†} and Shane M. O'Mara^{2†}

¹School of Psychology, Cardiff University, Cardiff, United Kingdom.

²School of Psychology and Trinity College Institute of Neuroscience, Trinity College Dublin, The University of Dublin, Dublin, Ireland.

[†]Corresponding authors

[†]E-mail: aggleton@cardiff.ac.uk **Orcid** 0000-0002-5573-1308

[†]E-mail: smomara@tcd.ie **Orcid** 0000-0001-8087-8531

Abstract | Standard models of episodic memory focus on hippocampal–parahippocampal interactions, with the neocortex supplying sensory information and providing a final repository of mnemonic representations. However, recent advances have shown that other regions make distinct and equally critical contributions to memory. In particular, there is growing evidence that the anterior thalamic nuclei have a number of key cognitive functions that contribute to episodic memory. In this article, we describe these findings and argue for a core, tripartite memory system, comprising a ‘temporal lobe’ stream (hippocampal centred) and a ‘medial diencephalic’ stream (anterior thalamic centred) that together act on shared cortical areas. We demonstrate how these distributed brain regions form complementary and necessary partnerships in episodic memory formation.

[H1] Introduction

Many theoretical and computational models of episodic memory ignore the influence of medial diencephalic sites, while emphasising hippocampal interactions with the cortex¹⁻⁵. However, there is growing evidence that, by failing to account for the necessary and substantial contributions of medial diencephalic brain regions (most notably the anterior thalamic nuclei), such models are imbalanced and misleading.

The likely importance of the anterior thalamic nuclei for memory first emerged more than a century ago, when descriptions appeared⁶⁻⁸ of the devastating amnesia following medial diencephalic damage. The consistent pattern of pathology adjacent to the third ventricle helped to highlight the potential significance of the anterior thalamic nuclei for human episodic memory – one’s memory for day-to-day events. Subsequent animal studies have shown that the anterior thalamic nuclei play multiple, vital roles in aspects of cognition for

which their influence has frequently been underestimated, including for spatial processing, memory, and attention^{9,10,11,12}. It is proposed that (like the hippocampus) the contributions of the anterior thalamic nuclei to the processing of spatial information are a requirement for the contextual processing and binding that is integral to the encoding and retrieval of episodic information^{5,13-16}, thereby allowing the representation of experiences in cognitive space^{4,5}.

The anterior thalamic nuclei often function in partnership with the hippocampus, with the two structures acting in parallel to effect plasticity in widespread areas of medial and midline cortex¹⁷. In addition, some of the spatial functions of the hippocampus rely on its inputs from the anterior thalamic nuclei¹⁸⁻²¹. However, there is also evidence for memory-related functions of the anterior thalamic nuclei that are independent of the hippocampus^{9,10,22}. Consequently, the cognitive roles of these thalamic nuclei are not simply framed by the hippocampus, despite many overlapping areas of function.

In this article we outline a model that realigns the hippocampal formation, anterior thalamic nuclei, and cortex as necessary and co-equal partners within a tripartite mnemonic system, thereby highlighting vital, but relatively neglected, thalamic components of the brain's memory systems. The focus of the article is intentionally on the anterior thalamic nuclei in order to redress the balance of coverage given to each of the key elements of the model. Indeed, the hippocampal formation and its cortical interactions have been discussed in numerous reviews and models to which the reader is referred^{4,5,15,16}.

[H1] A brief history

One of the first formal descriptions of a persistent amnesic syndrome was published by Robert Lawson²³ in 1878. Summarising his observations of a series of patients with amnesia driven by alcohol abuse, Lawson wrote that the “feature of such cases which is sufficiently striking to give character to them is the almost absolute loss of memory for recent events”. Almost a decade later, the same alcohol-related amnesic syndrome was more famously described by Sergei Korsakoff²⁴, who noted that “especially characteristic is a derangement of memory and of the association of ideas”. Shortly thereafter, based on post-mortem studies, Hans Gudden⁶ reported that “atrophy of the mammillary bodies appears to be a frequent finding in cases of alcohol neuritis with amnesia (and perhaps generally in chronic alcoholism)”. By the early 20th century, further post-mortem studies⁷ confirmed the consistent mammillary body involvement in what had come to be known as Korsakoff syndrome,

alongside apparently more variable thalamic pathology (reviewed in REF⁸). The latter included changes in the anterior thalamic nuclei^{8,25} (which comprise the anterodorsal, anteromedial, and anteroventral nuclei; **Fig. 1**). Linking these diencephalic pathologies is how the anterior thalamic nuclei form the principal target of mammillary body neurons^{26,27}. It might, therefore, be supposed that the contributions of these two diencephalic sites (the anterior thalamic nuclei and the mammillary bodies) to episodic memory would have been intensively scrutinised. That is not what happened: instead, the hippocampus took centre stage.

In 1900, Vladimir Bekhterev²⁸ reported a possible link between hippocampal damage and memory loss in humans. However, it still took decades for this idea to become established (reviewed in REF²⁹), with the landmark case of amnesic patient H.M.³⁰ playing a central role. Meanwhile, research into diencephalic amnesia remained severely hampered by the lack of a similar landmark case with confirmed, discrete pathology. Indeed, identifying such patients has remained problematic, mainly because of the twin difficulty of distinguishing individual diencephalic nuclei and determining potential disruption to fibres of passage when relying on non-invasive imaging. In contrast, studies linking hippocampal pathology and memory took advantage of the sensitivity of this structure to conditions such as anoxia³¹⁻³³ while its size and location enabled imaging studies. Perhaps just as detrimental for diencephalic amnesia has been the pervasive view that the mammillary bodies and anterior thalamic nuclei function as downstream enablers for the hippocampus³⁴⁻³⁷ and are thereby of secondary importance. This perspective was first popularised in accounts of the ‘Papez circuit’, a loop of extended hippocampal connections that was previously thought to maintain emotions³⁴.

Hippocampal research was further galvanised by the parallel discoveries, in animal experiments, of place cells³⁸ (which fire selectively at certain locations), synaptic plasticity in the form of long-term potentiation³⁹, and confirmation that damage to the hippocampus severely disrupts performance in tests of spatial memory⁴⁰. More recent studies have demonstrated a key role of hippocampal oscillations (rhythmic patterns of neuronal firing) for the optimisation of plasticity^{41,42}, while findings from intracranial recordings in the medial temporal lobe of epileptic patients have shown that theta oscillations are associated with navigation⁴³ and episodic recall⁴⁴⁻⁴⁶. Inevitably, therefore, investigations into the mechanism of episodic memory gravitated towards the hippocampus and were later refined by considering hippocampo–cortical interactions^{15,41,47-49}. This hippocampal-centred approach

further reinforced the notion that the anterior thalamic nuclei merely serve as relays or have ill-specified roles in modulating or activating temporal lobe structures supporting memory^{35-37,50}.

[H1] Revisiting the anterior thalamic nuclei

To rebalance the situation outlined above, there is a pressing need to appreciate the distinct contributions of medial diencephalic sites to memory. Chief among these sites are the anterior thalamic nuclei. Over recent years, there has been a growing awareness that these nuclei can contribute to the cognitive changes associated with aging⁵¹, mild cognitive impairment^{52,53}, various dementias⁵³⁻⁵⁵, and, potentially, an array of other neurological and psychiatric disorders⁵⁶⁻⁵⁸. Meanwhile, the discovery of neurons within the rodent anterior thalamic nuclei that signal different types of spatial information (**Table 1**) has forced a further rethink regarding the significance of these nuclei in cognition⁵⁹.

[H2] Anterior thalamic connectivity

The anterior thalamic nuclei stand out among medial diencephalic structures as they receive dense, direct hippocampal and parahippocampal projections (**Fig. 2 ; Table 1**) from the subiculum, presubiculum, and postsubiculum, as well as providing direct return projections to the same sites⁶⁰⁻⁶³. Importantly, comparable sets of connections are present in rodents and nonhuman primates⁶⁴. In addition, the rodent anterior thalamic nuclei receive sparse inputs from hippocampal area CA1⁶⁵, while sparse inhibitory projections from CA3⁶⁶ target the anterodorsal nucleus. Meanwhile, almost every mammillary body neuron is thought to innervate the anterior thalamic nuclei through the mammillothalamic tract^{26,27}. Via this monosynaptic mammillary route, the subiculum, presubiculum, and postsubiculum provide additional, indirect influences on the anterior thalamus^{27,64}. The anterior thalamic nuclei also have extensive reciprocal connections with retrosplenial and anterior cingulate cortices⁶⁴. Consequently, the anterior thalamic nuclei interlink multiple cortical and subcortical areas strongly associated with memory, amnesia, and spatial processing^{35,37,50,64,67}.

[H2] Comparisons with other thalamic nuclei

Other medial diencephalic sites contribute to cognition⁶⁸. Foremost among these are the thalamic nucleus reuniens and the dorsomedial thalamic nucleus (reviewed in REFs⁶⁹⁻⁷¹). While both nucleus reuniens and the anterior thalamic nuclei support spatial learning and memory, anterior thalamic lesions have a wider impact in rats and are the more disruptive⁷¹⁻⁷⁵. Although nucleus reuniens is also reciprocally connected with the hippocampus, including

direct inputs to CA1, it only receives sparse inputs from the mammillary bodies^{76,77} and retrosplenial cortex⁶² in rats. Furthermore, although the anterior thalamic nuclei and nucleus reuniens appear to be connected to overlapping cortical sites, the respective inputs from these regions are largely separated by topography and lamina⁶², indicative of complementary functions. This difference can be seen in their respective importance for linking prefrontal (reuniens) and cingulate/retrosplenial regions (anterior thalamic nuclei) to the hippocampus^{62,78}. In association with its prefrontal connections, the rodent nucleus reuniens assists spatial working memory⁷⁹, gating information to the hippocampus⁸⁰.

Meanwhile, the dorsomedial thalamic nucleus, which lacks direct hippocampal connections⁸¹, is far less important for rodent spatial learning than the anterior thalamic nuclei^{69,70,82}. Instead, through its dense, reciprocal connections with the prefrontal cortex, the dorsomedial nucleus supports a range of functions that rely upon the prefrontal cortex, including executive actions (reviewed in REFs^{70,81}). The complementary actions of these adjacent thalamic nuclei (anterior, dorsomedial, reuniens) means that typical pathologies, which affect more than one thalamic nuclei, both increase the severity and breadth of cognitive impairments, while adding to the difficulty of discerning the roles of individual regions^{70,83}.

[H2] Hippocampal comparisons

To understand why the anterior thalamic nuclei deserve more attention it is helpful to make comparisons with the hippocampus. Numerous parallels emerge. Just as hippocampal tissue loss is principally responsible for temporal lobe amnesia³¹⁻³¹, so the anterior thalamic nuclei are at the core of diencephalic amnesia. The relevant clinical evidence for the latter statement largely derives from two sources; studies of Korsakoff syndrome^{8,25,84} and systematic analyses of diencephalic strokes^{8,85,86}. The second approach repeatedly shows that mammillothalamic tract involvement is the best predictor of memory loss following thalamic infarcts^{85,86}, highlighting the importance of the anterior thalamic nuclei, the principal target of the mammillothalamic tract (**Figs. 1C, 2**).

Meanwhile, behavioural studies in rodents repeatedly show that anterior thalamic lesions cause spatial memory deficits that parallel those following hippocampal lesions^{11,72,87-89}. These deficits are striking with respect to their range and severity¹¹. Indeed, the consequences of anterior thalamic lesions on spatial memory tasks are often greater than those of lesions to other sites beyond the hippocampal formation^{11,71,72,89-91}. A resulting question is, therefore, whether anterior thalamic lesions can be as disruptive as hippocampal damage. Unfortunately,

almost all evidence relating to this question is indirect, coming from comparisons of anterior thalamic lesions with lesions to the fornix, the principal tract containing extrinsic hippocampal fibres. For this reason, fornix lesions only partially replicate hippocampal damage^{92,93}.

While the effects of fornix lesions and anterior thalamic lesions on spatial memory tasks are sometimes similar^{11,94-98}, anterior thalamic damage can also be more disruptive^{73,99}. For example, a geometric deficit in the framing of space⁹⁹ is seen after hippocampal and anterior thalamic (but not fornix) lesions^{99,100}. Other relevant findings show that lesions of the fornix, hippocampal formation, and anterior thalamic nuclei all cause comparable deficits on tests of automated delayed nonmatching-to-sample in rats when using identical methodologies^{98,101}. This equivalence is notable as the same spatial paradigm has advanced our appreciation of hippocampal memory functions¹⁰². At the same time, complete hippocampal formation lesions (which include the subiculum) appear more disruptive than anterior thalamic lesions for reinforced spatial alternation¹¹. Interpretations of this difference should, however, be made with care as ‘hippocampus proper’ lesions, i.e., surgeries that spare the subiculum, are associated with more limited spatial deficits^{103,104}. These findings raise the possibility that hippocampus proper lesions and anterior thalamic lesions could have equivalent disruptive effects.

[H2] Not just head-direction

With hindsight, the failure of the severe spatial memory deficits after anterior thalamic lesions to spark wider interest probably stems from the misguided sense that the contribution of this site was already known. Within the anterior thalamic nuclei, the anterodorsal nucleus has long been recognised as a key component of the ‘head-direction system’^{18,105,106}, which provides compass-like directional signals to assist navigation¹⁸. Indeed, lesions involving the rat anterodorsal nucleus abolish parahippocampal head-direction signals¹⁹ and disrupt the activity of ‘grid’ cells²⁰, which normally fire at regular, discrete locations forming hexagonally spaced fields that create a metric for local space¹⁰⁷. In contrast, anterodorsal nucleus head-direction signals do not require the integrity of the hippocampus¹⁰⁸. However, more recent work reveals that, while the loss of head-direction signals undoubtedly contributes to the spatial deficits observed following anterior thalamic lesions²⁰, the loss of other anterior thalamic functions plays its part^{121,109}. Arguably, the clearest evidence comes from the demonstration in rats that disconnecting the ascending head-direction pathway (from the lateral mammillary nucleus to

the anterodorsal thalamic nucleus) only induces mild, transient spatial learning deficits^{110,111}. These mild impairments are in marked contrast to the severe, persistent, mnemonic deficits seen after lesions involving all three anterior thalamic nuclei^{11,12,89,112}. One important conclusion is that head-direction information is not a proxy for episodic memory processing¹¹¹.

It remains interesting to consider why attention remained firmly fixed on the hippocampus at the expense of the anterior thalamic nuclei given their demonstrable importance for spatial processing and memory. It seems most probable that the landmark discoveries of place cells and forms of synaptic plasticity, such as long-term potentiation (LTP), within the hippocampus^{38,39} opened the way for a myriad of investigations into memory at the neuronal level at a time when no counterparts existed for the thalamus¹¹³.

[H2] Spatial signalling and plasticity

The discovery of spatial cell-types beyond head-direction cells in the rat anterior and midline thalamus has more recently encouraged a broader conception of thalamic spatial signalling^{59,114-116}. These anterior and midline thalamic spatial cells resemble the grid, place, and border cells (which signal the perimeter of an environment) found in the medial temporal lobe^{114,115}. In addition, head-direction cells have been found more extensively across the anterior thalamic nuclei^{59,116}, i.e., beyond the anterodorsal nucleus. One feature of all of these thalamic spatial cells is the high-fidelity of the spatial information that they encode, undermining one previous suggestion that these nuclei merely enable some form of diffuse cortical arousal^{117,118}. These discoveries are additionally significant because spatial processing provides a framework not only for navigation but also for an array of other cognitive processes. In particular, contextual information plays an integral role in episodic memory encoding and retrieval, while being particularly important for the visualization of information^{4,14-16,47,119}.

The hippocampus understandably remains a favoured target for studies of neuronal plasticity, although we are also beginning to appreciate the plasticity that takes place within medial thalamic nuclei. A feature of this plasticity is that its underlying mechanisms may depend on the fibres of origin. One set of experiments¹²⁰ examined the plastic responses of rat anterior thalamic synapses to different types (high- and low-frequency) of stimulation. These investigations revealed that basal synaptic transmission mediated by the mammillothalamic

tract undergoes stimulation-dependent, BDNF-mediated potentiation¹²⁰. This pathway resulted in LTP of the thalamic field response, and was induced predominantly via high-frequency stimulation. By contrast, anterior thalamic long-term depression (LTD) of the field response could only be induced after low-frequency stimulation of the direct subiculum (hippocampal) projections to the anterior thalamic nuclei¹²⁰, which largely rely on the fornix. Hence, the two major tracts that mediate direct routes to the anterior thalamic nuclei (fornix vs mammillothalamic tract) exhibit opposing plasticity characteristics. This suggests the integration of hippocampal and mammillary body inputs depends on the pattern of prior activity in the fornix and mammillothalamic tract components of the circuit. Furthermore, evidence of the balancing effects of these parallel pathways upon anterior thalamic activity reinforces the notion that the anterior thalamic nuclei do not merely duplicate hippocampal functions, helping us to appreciate how they may make independent contributions to episodic memory consolidation, perhaps by offering complementary pathways to stabilise memory, allowing mnemonic streams that function in parallel and are partially independent of each other (**Fig 4**).

Deep-brain thalamic recordings in epileptic patients have added to our knowledge. Anterior thalamic activity has, for example, been linked to memory retrieval¹²¹. Meanwhile, a series of related studies describes how oscillatory activity in the anterior thalamic and dorsomedial thalamic nuclei correlates with memory performance¹²²⁻¹²⁴. For example, the presence of theta oscillation synchrony across neocortical and anterior thalamic areas, as well as cross-coupling with gamma oscillations, predicted subsequent memory for complex photographic scenes¹²². These findings have been incorporated into the hypothesis that the anterior thalamic nuclei promote the selection and coordination of task-relevant information during human memory formation¹²⁴. A further implication is that the role of the anterior thalamic nuclei in memory encoding involves accessing information from widespread neocortical sources, including frontal areas¹²⁴.

Imaging studies, using both functional and structural MRI, have repeatedly highlighted hippocampal associations with spatial^{125,126} and episodic¹²⁷⁻¹²⁹ memory. Comparable imaging information concerning the anterior thalamic nuclei had been largely lacking, partly due to the difficulty of localising these nuclei in the human brain. Evidence of progress comes from the relatively few but growing number of MRI-based studies of cognition reporting activity-related signals in anterior thalamic regions^{22,130-133}. fMRI studies of associative recognition¹³⁰,

as well as comparisons of familiarity with recollective based recognition¹³², implicate anterior thalamic nuclei activity in the encoding¹³⁰ and recall^{130,133} of episodic information. Moreover, advances in both structural and functional MRI¹³⁴⁻¹³⁶ make it increasingly feasible to interrogate individual anterior thalamic nuclei (**Fig. 1**).

[H1] Separating the anterior thalamic nuclei

Evidence that distinct functional contributions are made by the three anterior thalamic nuclei has come from multiple levels of animal research (**Table 1**). For example, electrophysiological recordings in rats show different distributions of both spatial^{59,114-116} and theta-modulated¹³⁷ cells across the three nuclei, while gene-transcription analyses show that each nucleus belongs to a different cluster of brain sites, each with individual gene expression profiles¹³⁸. Extending these differences, the anatomical inputs to each individual nucleus^{60,64} is distinct (**Fig. 2**). Meanwhile, lesion studies in rats repeatedly reveal that, while each nucleus contributes to spatial learning and memory, their combined impact is far greater than a lesion within any single nucleus^{21,66,109,112,140,141}.

The differences between the three principal anterior thalamic nuclei are most apparent for the anterodorsal nucleus, which — in addition to being the only nucleus to contain large numbers of head-direction cells — has distinct cytoarchitectural and connectomic properties (**Table 1**). The concentration of head-direction cells in this nucleus points to a principal role in navigation, allied to the growing appreciation that head-direction information can impact on other classes of spatial cells^{20,142}. In addition, the firing of mouse anterodorsal neurons immediately before hippocampal sharp-wave ripples during non-REM sleep may signal previously experienced directions of movement¹⁴³, potentially contributing to consolidation. The finding that ~60% of rat anterodorsal units can be classified as head-direction cells¹⁰⁵ does, however, raise questions about the functional properties of the remaining cells. It is, therefore, of interest that optogenetic silencing of the inhibitory projection from hippocampal area CA3 to the anterodorsal nucleus disrupts contextual fear memory retrieval after extended delays⁶⁶, again pointing to a role in mnemonic consolidation.

Functional differences between the anteromedial and anteroventral nuclei are, at present, often speculative because few behavioural studies have sought to isolate these nuclei (**Table 1**). Of the anterior thalamic nuclei, in both rodent and primates the anteromedial nucleus has the most links with the medial prefrontal and anterior cingulate cortices, alongside further

connections with perirhinal cortex, dysgranular retrosplenial (area 30) cortex, and related visual areas^{60,64,144-146}. The anteromedial nucleus also contains place cells and border cells, but few head-direction cells^{59,114,115}. Targeted lesions of this nucleus give modest deficits on spatial working memory tasks^{139,140}, with further evidence of a selective contribution to the retrieval of spatial information¹⁰⁹. Largely based on its frontal connectivity, it might be supposed that the anteromedial nucleus assists in the retrieval of high-interference information – that is, the separation of related target information^{147,148}. Other potential roles for this region include the integration of item and contextual information in parahippocampal areas, while its anterior cingulate connections suggest that it may support aspects of attention¹⁴⁹ (**Fig. 3**). The combination of functions implies dynamic relations between differing cognitive processes; spatial processing and attention are deployed during episodic memory encoding and consolidation (as they are contributory processes to memory), but might function independently of memory during other tasks.

Meanwhile, the anteroventral nucleus is well connected with dorsal hippocampal, presubicular and postsubicular sites, along with the granular retrosplenial cortex (area 29)^{60,64}. Its concentration of inputs from the rat distal subiculum⁵³ further underline its links with spatial processing^{150,151}. The anteroventral nucleus also contains most of the theta-modulated neurons within the anterior thalamic nuclei^{116,137}, which are jointly influenced by fibres in the fornix¹⁵² and the mammillothalamic tract¹⁵³. These activation patterns contribute to cortical (retrosplenial) and hippocampal oscillations, affecting plasticity^{124,153}. In animal studies, reversible lesions implicate the anteroventral nucleus in the encoding, consolidation, and retrieval of spatial information¹⁰⁹. These widespread effects are consistent with this region have the role of a hub nucleus with access to multiple sources of spatial codes, suggesting a potential role in on-line location monitoring, as well as gating and updating spatial information^{12,109,154}.

The proposed functions of the different anterior thalamic nuclei provide a working framework with which to generate testable hypotheses, with the caveat that the distinctions are heavily based on rodent studies. Other considerations include the degree of communication between these different nuclei and whether this might diminish their individual differences. While there is no evidence for local pathways between the three thalamic nuclei, Golgi stains show that dendritic fields may extend across the nuclei¹⁵⁵, potentially blurring distinctions. Likewise, while the connectivity of the three nuclei clearly differs, this is often by degree

rather than absolute differences⁶⁰. For example, the anteroventral nucleus may also contribute to attentional functions with the anterior cingulate cortex¹⁴⁶. Similarly, while the anterodorsal nucleus is the principal site for head-direction cells, they are also present in the anteroventral nucleus¹¹⁶. It is, therefore, likely that the functional differences between the three nuclei will prove to be nuanced.

[H1] Models of function

[H2] Interactions with the hippocampal formation

Historically, researchers have often conceptualised anterior thalamic function within the context of its dense hippocampal and parahippocampal inputs³⁴⁻³⁷, reinforced by evidence that damage to the fornix can cause anterograde amnesia in humans^{156,157}. However, the fornix makes numerous connections outside the medial diencephalon, highlighting the need to isolate hippocampal–anterior thalamic interactions to investigate their functions.

One experimental solution was to combine unilateral lesions in the anterior thalamic nuclei with unilateral lesions in the hippocampus of the opposite hemisphere. While this disconnection method helped to confirm the importance of anterior thalamic–hippocampal interactions for rat spatial learning^{158,159}, it could not determine the direction of effect. More targeted chemogenetic methods in rats have disrupted only those hippocampal projections that arise in the dorsal subiculum and terminate in the anteromedial and anteroventral nuclei²¹. This was sufficient to impair performance in a spatial working memory task (T-maze alternation), but only after maze rotation to further tax allocentric processing²¹. While this procedure, which spares anterodorsal nucleus function, further highlights the significance of anterior thalamic contributions to memory beyond head-direction information, the resulting deficit was not as severe as that seen after complete anterior thalamic lesions^{11,21}, again pointing to additive roles of the three nuclei^{139,140}. Meanwhile, optogenetic studies in rats have isolated subiculum cells with different targets, revealing that place-related firing is a feature of those subiculum neurons that innervate the anteroventral thalamus, as well as the medial mammillary bodies, retrosplenial cortex, and nucleus accumbens¹⁶⁰.

A complementary approach to study hippocampal–diencephalic connections in humans involves diffusion MRI, which has shown that changes in fornix properties are closely associated with levels of episodic memory¹⁶¹⁻¹⁶³, with further links to recollective-based

recognition¹⁶¹ and navigation¹⁶⁴. A valuable refinement has allowed the isolation of the postcommissural fornix (**Box 1**), which principally contains hippocampal projections to the anterior thalamic nuclei and mammillary bodies. Such studies have found correlations between the properties of this pathway and aspects of memory, including visual recall¹⁶⁵ and types of spatial learning¹⁶⁶, suggesting that the functions of the postcommissural fornix differ from those of the precommissural fornix (which links the hippocampus with septal, striatal, and frontal sites).

The reverse challenge is to understand the importance of anterior thalamic inputs to the hippocampal formation (**Table 1**). As described above, past analyses largely focussed on head-direction signals^{19,20}. More recently, however, the wider significance of anterior thalamic influences on the dorsal hippocampal formation was revealed by the discovery that in rats both permanent and transient anterior thalamic lesions cause a cessation of spatial-responsive firing (place, head-direction, border, and grid cells) within the subiculum¹⁵⁴, combined with impaired spatial alternation memory. In the same study, permanent anterior thalamic lesions spared hippocampal CA1 place cell fields (but see also¹⁰⁶). This is important because the very dense, plastic projections from CA1 to the subiculum¹⁶⁷ might otherwise have been assumed to be sufficient to ensure the effective function of spatial cells in the subiculum¹⁵⁶⁴. It remains to be determined whether the observed subiculum silencing arises from the loss of the direct or indirect inputs from the anterior thalamus. However, a recent chemogenetic study showed the importance of the direct anterior thalamic projections to the rat dorsal hippocampal formation for spatial working memory²¹. Other insights come from evidence that stimulation of the anterior thalamic nuclei in epileptic patients can modulate hippocampal gamma activity¹⁶⁸.

[H2] Communications beyond the hippocampus

To substantiate the case that the anterior thalamic nuclei are independently critical for memory it is necessary to highlight their connections and potential functions that are not part of traditional hippocampal circuitry. Sites such as the prelimbic, anterior cingulate, and retrosplenial cortices project to the anterior thalamic nuclei, but largely fail to directly innervate the hippocampal formation^{64,194-171}; trans-synaptic tracing in rats further confirms that the anterior thalamic nuclei form a major monosynaptic route from these cortical regions to the dorsal hippocampal formation⁷⁸. Consequently, along with the nucleus reuniens^{62,78}, the

anterior thalamic nuclei provide a potentially important (but rarely recognised) subcortical route, allowing frontal sites to influence hippocampal activity.

When comparing respective subcortical afferents to the anterior thalamic nuclei and hippocampal formation, the mammillary bodies provide the most striking difference. The lateral mammillary nucleus projects to the anterodorsal nucleus, whereas the medial mammillary nucleus projects to the anteromedial and anteroventral nuclei⁶⁴. In contrast, the mammillary bodies do not innervate the hippocampal formation. The finding that mammillothalamic tract lesions in rats, which disconnect the anterior thalamic nuclei, affect oscillatory activity and neuronal microstructure in the hippocampus¹⁵³ has been interpreted as indicating a suppression of learning-induced plasticity, set alongside changes in retrosplenial theta and immediate-early gene expression^{153,172}. The latter findings add to the discovery that anterior thalamic lesions cause a persistent retrosplenial hypoactivity, as measured by immediate-early genes, alongside a wider de-regulation of retrosplenial gene transcription^{173,174}. Additional attention has focused on the ascending tegmental inputs from Gudden's nuclei to the mammillary bodies, which contribute to spatial navigation and learning^{172,178}, but do not innervate the hippocampus. It is presumed that these same tegmental inputs have their principal actions via the anterior thalamic nuclei¹⁷⁵.

[H2] Widening our horizons

By moving from a hippocampal-centred view, a broader perspective emerges that is illustrated by the many convergent efferent projections from the anterior thalamic nuclei and hippocampal formation to cortical sites. Projections from both structures terminate in the retrosplenial cortex and parahippocampal region, as well as the anterior cingulate and prelimbic cortices^{17,63,64,176-178}. The need to look beyond reciprocal anterior thalamic–hippocampal interactions also stems from fMRI analyses of the human ‘default mode network’^{179,180}. This network, which was initially closely linked with self-related (introspective) tasks¹⁷⁸, is also activated during spatial, mnemonic, and social cognitive tasks¹⁸¹⁻¹⁸⁴. The default mode network is typically regarded as comprising several subsystems, including those in the medial prefrontal cortex, posteromedial parietal cortex, and medial temporal regions, with partial separation between different areas of cognitive engagement¹⁸⁻¹⁸³. One focus has been the apparent overlap between medial temporal and midline components of the default mode network and those areas that are active when remembering the past^{182,184,185}, including constructing mental scenes from memory as well as imagining the

future^{184,185}. This overlap extends to the distributed network of brain sites associated with anterograde amnesia⁶¹. Such findings have strongly influenced models of memory such as the ‘constructive episodic simulation hypothesis’^{185,186}. Rather than memory being a replay of the past, such models emphasise the predictive construction of relevant scene information^{15,186}.

Connectivity analyses show that the human anterior thalamic nuclei, along with some other thalamic nuclei^{187,188}, have close links with multiple components of the default mode network¹⁸⁷. Furthermore, damage to the anterior thalamic nuclei disrupts activity in the default mode network¹⁸⁹, while deep-brain stimulation of the anterior thalamic nuclei in epileptics can modulate the same network¹⁹⁰. Meanwhile, functional connectivity studies place the retrosplenial cortex as a key gateway between the medial temporal lobe and frontal default mode subsystems that support episodic memory¹⁹¹, suggesting that anterior thalamic damage may impact on landmark and scene construction via its disruptive effects on retrosplenial cortex^{68,173,174,192}. Interestingly, diffusion imaging suggests a relationship between precommissural fornix status and the episodic richness of past and future events, while a similar association was not found for the postcommissural fornix¹⁹³. One interpretation of this finding is that anterior thalamic–retrosplenial, rather than hippocampal–anterior thalamic interactions have particular significance for scene construction.

Arguably more striking evidence of the independence of the anterior thalamic nuclei from the hippocampus is seen in their contributions to specific aspects of attention^{10,22,149}. Selective lesions of the anterior thalamic nuclei impair the ability of rats to accelerate their learning over a series of closely related discriminations in which normal animals narrow their attention to the same reinforced stimulus dimension¹⁰, while ignoring other cue types. Quite remarkably, these same thalamic lesions facilitate performance when the discriminations require a dimension switch¹⁰. This profile of altered performance, not associated with hippocampal dysfunction^{194,195}, points to an independent role for the anterior thalamic nuclei in cognition that contributes to learning and memory.

Subsequent chemogenetic studies have found that disrupting anterior cingulate–anteromedial thalamic nucleus interactions cause the same profile of learning deficits and facilitation on the same test of attention assisted learning¹⁴⁹. One interpretation is that the anterior thalamic nuclei, along with the anterior cingulate cortex, normally help to maintain attention to previously rewarded stimulus categories, a role extending beyond spatial processing¹⁴⁹. The

paradoxical, facilitated switching occurs because the lesioned rats fail to initially disengage from a stimulus dimension that subsequently becomes the rewarded dimension. Support for this interpretation comes from functional imaging and from evidence from patients with rostral thalamic damage^{22,130}, though these human studies lack the anatomical precision of the rodent analyses. If correct, these rodent and human findings reveal an anterior thalamic attentional system that opposes¹⁹⁶ the well-established actions of those prefrontal areas that normally enable flexible responding¹⁹⁷⁻¹⁹⁹. Indeed, a prefrontal-like pattern of lesion effects was observed on these same attentional tasks in rats with hippocampal damage²⁰⁰, thereby contrasting with anterior thalamic lesion effects. These anterior thalamic actions on attention and memory may then facilitate a wide variety of learning tasks^{9,201-205}.

These same findings prompt renewed interest in earlier evidence that lesions to the anterior thalamic lead to a range of nonspatial problems^{9,147,148,201,202}. Some, such as deficits in recency memory^{147,148}, may align with alterations in hippocampal function²⁰³. Others, such as changes in anxiety and increased motor activity might again be linked to disrupted hippocampal function but could also reflect changes in anterior cingulate action²⁰⁴. Meanwhile, studies of classical conditioning and discrimination learning in rabbits, which employed both electrophysiological recordings and lesion analyses, highlighted the importance of delayed anterior thalamic interactions with the retrosplenial cortex that emerged after initial learning of these nonspatial associative tasks was established^{9,205}. These thalamo–cortical interactions, along with others involving the dorsomedial thalamic nucleus, were interpreted as supporting the associative attention to reward-related cues that emerges during different stages of discrimination learning^{9,205}. Consequently, these findings again point to the significance of the anterior thalamic nuclei for attention based on experience, contributions that may influence a variety of learning tasks, including those involving episodic memory.

[H1] A new tripartite memory model

As outlined above, the anterior thalamic nuclei make independent, wider contributions to cognitive function beyond those framed by their hippocampal connections⁶⁸ (**Fig. 3**). Through these contributions, the multiple actions of the anterior thalamic nuclei together affect episodic memory. Thus, we suggest there is a need to develop broader models of episodic memory that incorporate medial diencephalic contributions to encoding and consolidation. While this more inclusive approach appears in some existing models^{35,37,50}, the roles of the anterior thalamic nuclei remain poorly specified.

In the model we present here, it is suggested that there is a hippocampal–cortical memory stream, and a medial diencephalic–cortical memory stream (with the anterior thalamic nuclei at its core). The hippocampal memory stream courses from the entorhinal cortex to the hippocampus proper (the dentate gyrus, CA3 and CA1), thence to the subiculum, and onwards to cortical sites (such as parahippocampal, prefrontal, and retrosplenial cortical sites), along with parallel CA1 projections to parahippocampal and prefrontal areas¹⁷⁷. Meanwhile, the anterior thalamic nuclei project to the hippocampal formation (subiculum) and the parahippocampal and retrosplenial cortices, while receiving return inputs from the hippocampal formation (many via the mammillary bodies), retrosplenial, anterior cingulate, and prefrontal cortices. Both memory streams project to several cortical sites, including parahippocampal, prefrontal, and retrosplenial cortices, where synaptic plasticity supports memory consolidation (**Fig. 4**). These mnemonic streams are envisaged to operate in parallel, with activity in each stream being partially independent of the other stream. Critically, there are cortical zones where anatomical influences from both streams converge. Anterior thalamic nuclei- and hippocampus-dependent memory depends on these zones of interaction, where synchronous activity is required for mnemonic consolidation.

This tripartite model predicts that separate lesions to the hippocampal–cortical circuit or to the medial diencephalic–cortical circuit will cause approximately equivalent deficits in episodic memory, given the significance of synchronous cortical activity at the temporal lobe and medial diencephalic sites of convergence. Thus, damage to the specific cortical sites of convergence between these two streams will also cause memory impairments⁶⁷, providing further insights into memory consolidation^{15,48,49}.

While details of the respective synaptic actions of these two streams are beyond the scope of this review, recent reviews²⁰⁶ describe how cortical layer 1 might be critical for long-term plasticity, reflecting the convergence of thalamic inputs with those from sites such as the hippocampal formation (either directly or via parahippocampal structures)^{17,176,207}. The retrosplenial cortex is a likely candidate as it receives convergent projections from the anterior thalamic nuclei, hippocampal area CA1, and the dorsal subiculum^{17,176}. Within retrosplenial cortex layer 1, opposing excitatory anteroventral nucleus and inhibitory CA1 actions have contrasting, vital effects on mouse contextual fear conditioning¹⁷. Furthermore, superficial vesicular glutamate transporter 1-expressing (VGLUT1⁺) projections from the subiculum to

retrosplenial cortex may be involved in processing recent context memories, whereas corresponding VGLUT2⁺ projections to retrosplenial cortex contribute to their long-lasting storage¹⁷⁶. Such findings reinforce the value of initially exploring joint hippocampal and anterior thalamic influences on cortical sites, while underlining how the anterior thalamic nuclei make independent contributions. Intriguingly, the anterior thalamic and dorsal subiculum projections strongly target small low-rheobase pyramidal cells in the retrosplenial cortex, while neighbouring regular-spiking cells are preferentially controlled by claustral and anterior cingulate inputs, sources of mostly non-spatial information²⁰⁸. Such analyses begin to reveal the true complexity of many of these cortical interactions.

[H1] Future challenges

Many issues remain. These include the need to test the potential significance of the anterior thalamic nuclei for landmark and scene construction, while determining whether converging hippocampal formation and anterior thalamic projections within retrosplenial cortex support these functions. A striking feature of the hippocampal projections to the retrosplenial cortex in rats is that approximately 50% of them collateralise so that the same neurons also innervate the mammillary bodies²⁰⁹, creating a unique linkage across the anterior thalamic hub. Another issue arises from evidence pointing to an especially important role for the human anterior thalamic nuclei in recollective (associative), but not familiarity-based, episodic information^{37,83}. Currently, the absence of patients with truly selective anterior thalamic pathology has stalled the ability to test this prediction. Perhaps for this same reason, the findings from thalamic stroke patients remain mixed²¹⁰⁻²¹². There does, however, appear to be more agreement from initial fMRI studies^{132,133}, which find support for the recollection:familiarity distinction with respect to the anterior thalamic nuclei, but replications are required. Further issues concern the potential significance of the anterior thalamic nuclei for memory retrieval¹⁰⁹ and whether their roles are facilitated by their frontal connections^{41,78,169}.

Additional challenges involve making further comparisons with other components of the cognitive thalamus. For example, nucleus reuniens appears to be the principal thalamic relay to the rodent hippocampal formation from more rostral and ventral prefrontal cortical sites, while the anterior thalamic nuclei increasingly replace that role for cingulate and retrosplenial projections indirectly reaching the hippocampus^{62,78}. The functional consequences of these differences require added examination. Other issues stem from growing evidence that the

anterior thalamic nuclei contribute to specific aspects of attention^{22,149,201}, a function that is potentially independent of the hippocampus but can still affect learning and memory. This impact on learning can be seen in how animals with anterior thalamic lesions fail to benefit from the repeated experience of prior, similar discriminations¹⁰. Human neuropsychological studies also point to deficits in executive function following rostral thalamic damage¹³⁰ that stem from the attentional demand. Related fMRI studies also suggest anterior thalamic influences on working memory²² and point to contributions away from the medial temporal lobe^{130,201}. It may prove relevant that recent diffusion imaging studies²¹³ suggest that the anterior thalamic nuclei are far more richly connected with prefrontal areas than often suspected from animal tracer studies. While dense reciprocal anterior cingulate–anterior thalamic interconnections have long been recognised⁶⁰⁻⁶⁴, there are increasing reasons to believe that the anterior thalamic nuclei not only receive information from distributed areas of prefrontal cortex^{62,162}, but may prove to act back upon these same areas, both directly and indirectly, to support cognition¹²⁴. Determining these prefrontal contributions represents an important future challenge.

Our review prompts the question of whether the anterior thalamic nuclei have one paramount function that largely explains their significance for memory, or whether their importance arises from multiple contributions to memory, spatial processing, and attention. The present analysis strongly suggests the latter (**Fig. 3**). Furthermore, their widespread zones of prefrontal¹²⁴ and cingulate^{9,64,17} influence create broader opportunities for these thalamic nuclei to contribute to collective mnemonic schemas, affective states, and avoidance learning^{9,204,205,214}, offering a variety of clinical opportunities for cognitive investigations. One example arises from how deep brain stimulation of the anterior thalamic nuclei in epileptic patients can improve a colour-matching working memory task¹⁶⁸. Meanwhile optogenetic theta-burst stimulation of the rat anterior thalamic nuclei can rescue the spatial working memory deficit following mammillothalamic tract lesions²¹⁵ - findings that yet again underline the significance of these thalamic nuclei for cognition.

Acknowledgements

J.P.A. and S.M.O'M. are supported in the work described here by the Wellcome Trust (103722/Z14/Z). We thank S. Commins, S. Martin, A. Nelson, and C. Ranganath for very helpful feedback on a prior version of the manuscript. We also thank M.M. Jankowski for assistance with Figure 1.

References

1. Eichenbaum, H. A cortical–hippocampal system for declarative memory. *Nat. Rev. Neurosci.* **1**, 41-50 (2000).
2. Manns, J. R. & Eichenbaum, H. Evolution of declarative memory. *Hippocampus* **16**, 795-808 (2006).
3. Squire, L. R. Memory and brain systems: 1969–2009. *J. Neurosci.* **29**, 12711-12716 (2009).
4. Bellmund, J. L., Gärdenfors, P., Moser, E. I. & Doeller, C. F. Navigating cognition: Spatial codes for human thinking. *Science* **362**, 6415 (2018).
5. Ekstrom, A. D. & Ranganath, C. Space, time, and episodic memory: The hippocampus is all over the cognitive map. *Hippocampus* **28**, 680-687 (2018).
6. Gudden, H. Klinische und anatomische beitrage zur kenntnis der multiplen alkoholneuritis nebst bemerkungen uber die regenerationsvorgange im peripheren nervensystem *Arch. Psychiat.* **28**, 643-741 (1896).
7. Gamper, E. Zur Frage der Polioencephalitis hæmorrhagica der chronischen Alkoholiker: Anatomische Befund beim alkoholischen Korsakow und ihre Beziehungen zum klinischen Bild. *Deutsche Ztschr. Nervenhe.* **102**, 122 (1928).
8. Kopelman, M. D. What does a comparison of the alcoholic Korsakoff syndrome and thalamic infarction tell us about thalamic amnesia? *Neurosci. Biobehav. Rev.* **54**, 46-56 (2015).
9. Gabriel, M. In *Neurobiology of Cingulate Cortex and Limbic Thalamus* (eds. Vogt, B.A. & Gabriel, M.) 478-523 (Birkhäuser, Boston, MA 1993).
10. Wright, N. F., Vann, S. D., Aggleton, J. P. & Nelson, A. J. A critical role for the anterior thalamus in directing attention to task-relevant stimuli. *J. Neurosci.* **35**, 5480-5488 (2015).
11. Aggleton, J. P. & Nelson, A. J. Why do lesions in the rodent anterior thalamic nuclei cause such severe spatial deficits? *Neurosci. Biobehav. Rev.* **54**, 131-144. (2015).
12. Aggleton, J. P., O'Mara, S. M., Vann, S. D., Wright, N. F., Tsanov, M. & Erichsen, J. T. Hippocampal–anterior thalamic pathways for memory: uncovering a network of direct and indirect actions. *Eur. J. Neurosci.* **31**, 2292-2307. (2010).
13. Aggleton, J. P. & Pearce, J. M. Neural systems underlying episodic memory: insights from animal research. *Phil. Trans. R. Soc. Lond. Series B: Biol. Sci.* **356**, 1467-1482 (2001).
14. Bird, C. M. & Burgess, N. The hippocampus and memory: insights from spatial processing. *Nat. Rev. Neurosci.* **9**, 182-194 (2008).
15. Barry, D. N. & Maguire, E. A. Remote memory and the hippocampus: A constructive critique. *Trends Cogn. Sci.* **23**, 128-142 (2019).
16. Yonelinas, A. P., Ranganath, C., Ekstrom, A. D. & Wiltgen, B. J. A contextual binding theory of episodic memory: systems consolidation reconsidered. *Nat. Rev. Neurosci.* **20**, 364-375 (2019).

17. Yamawaki, N., Li, X., Lambot, L., Ren, L. Y., Radulovic, J. & Shepherd, G. M. Long-range inhibitory intersection of a retrosplenial thalamocortical circuit by apical tuft-targeting CA1 neurons. *Nat. Neurosci.* **22**, 618-626 (2019).
18. Taube, J. S. The head direction signal: origins and sensory-motor integration. *Annu. Rev. Neurosci.* **30**, 181-207 (2007).
19. Goodridge, J. P. & Taube, J. S. Interaction between the postsubiculum and anterior thalamus in the generation of head direction cell activity. *J. Neurosci.* **17**, 9315-9330 (1997).
20. Winter, S. S., Clark, B. J. & Taube, J. S. Disruption of the head direction cell network impairs the parahippocampal grid cell signal. *Science* **347**, 870-874 (2015).
21. Nelson, A. J., Kinnavane, L., Amin, E., O'Mara, S. M. & Aggleton, J. P. Deconstructing the direct reciprocal hippocampal-anterior thalamic pathways for spatial learning. *J. Neurosci.* **40**, 6978-6990 (2020).
22. Bourbon-Teles, J. et al. Thalamic control of human attention driven by memory and learning. *Curr. Biol.* **24**, 993-999 (2014).
23. Lawson, R. On the symptomatology of alcoholic brain disorders. *Brain* **1**, 182-194 (1878).
24. Korsakoff, S. S. Ob alkogol'nom paraliche (Of alcoholic paralysis: disturbance of psychic activity and its relation to the disturbance of the psychic sphere in multiple neuritis of nonalcoholic origin). *Vestn. Klin. Psychiat. Neurol.* **4**, 1-102. (1887).
25. Harding, A., Halliday, G., Caine, D. & Kril, J. Degeneration of anterior thalamic nuclei differentiates alcoholics with amnesia. *Brain* **123**, 141-154 (2000).
26. Guillery, R. W. A quantitative study of the mamillary bodies and their connexions. *J. Anat.* **89**, 19-32 (1955).
27. Takeuchi, Y., Allen, G. V. & Hopkins, D. A. Transnuclear transport and axon collateral projections of the mamillary nuclei in the rat. *Brain Res. Bull.* **14**, 453-468 (1985).
28. von Bekhterev, M. Demonstration eines Gehirns mit Zerstörung der vorderen und inneren Theile der Hirnrinde beider Schläfenlappen *Neurologisches Zeitblatt*, **19**, 990-991 (1900).
29. Clark, R. E. & Squire, L. R. An animal model of recognition memory and medial temporal lobe amnesia: history and current issues. *Neuropsychologia*, **48**, 2234-2244 (2010).
30. Scoville, W. B. & Milner, B. Loss of recent memory after bilateral hippocampal lesions. *J. Neurol. Neurosurg. Psychiat.* **20**, 11-21 (1957).
31. Zola-Morgan, S., Squire, L. R. & Amaral, D. G. Human amnesia and the medial temporal region: enduring memory impairment following a bilateral lesion limited to field CA1 of the hippocampus. *J. Neurosci.* **6**, 2950-2967 (1986).
32. Rempel-Clower, N. L., Zola, S. M., Squire, L. R. & Amaral, D. G. Three cases of enduring memory impairment after bilateral damage limited to the hippocampal formation. *J. Neurosci.* **16**, 5233-5255 (1996).
33. Spiers, H. J., Maguire, E. A. & Burgess, N. Hippocampal amnesia. *Neurocase* **7**, 357-382 (2001).
34. Papez, J. W. A proposed mechanism of emotion. *Arch. Neurol. Psychiat.* **38**, 725-743. (1937).
35. Bastin, C. et al. An integrative memory model of recollection and familiarity to understand memory deficits. *Behav. Brain Sci.* **42**, e281: 1–60. (2019).

36. Delay, J. & Brion, S. Le syndrome de Korsakoff. (Masson, 1969).
37. Aggleton, J. P. & Brown, M. W. Episodic memory, amnesia and the hippocampal-anterior thalamic axis. *Behav. Brain Sci.* **22**, 425-444. (1999).
38. O'Keefe, J. & Dostrovsky, J. The hippocampus as a spatial map: Preliminary evidence from unit activity in the freely-moving rat. *Brain Res.* **34**, 171-174 (1971).
39. Bliss, T. V. & Lømo, T. Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. *J. Physiol.* **232**, 331-356 (1973).
40. Morris, R. G., Garrud, P., Rawlins, J. A. & O'Keefe, J. Place navigation impaired in rats with hippocampal lesions. *Nature*, **297**, 681-683 (1982).
41. Eichenbaum, H. Prefrontal–hippocampal interactions in episodic memory. *Nat. Rev. Neurosci.* **18**, 547-558 (2017).
42. Korotkova, T. et al. Reconciling the different faces of hippocampal theta: The role of theta oscillations in cognitive, emotional and innate behaviors. *Neurosci. Biobehav. Rev.* **85**, 65-80 (2018).
43. Watrous, A. J. et al. A comparative study of human and rat hippocampal low-frequency oscillations during spatial navigation. *Hippocampus* **23**, 656-661 (2013).
44. Addante, R. J., Watrous, A. J., Yonelinas, A. P., Ekstrom, A. D. & Ranganath, C. Prestimulus theta activity predicts correct source memory retrieval. *Proc. Natl. Acad. Sci. USA* **108**, 10702-10707 (2011).
45. Horner, A. J. & Doeller, C. F. Plasticity of hippocampal memories in humans. *Curr. Opin. Neurobiol.* **43**, 102-109 (2017).
46. Kota, S., Rugg, M. D. & Lega, B. C. Hippocampal theta oscillations support successful associative memory formation. *J. Neurosci.* **40**, 9507-9518 (2020).
47. Zeidman, P. & Maguire, E. A. Anterior hippocampus: the anatomy of perception, imagination and episodic memory. *Nat. Rev. Neurosci.* **17**, 173-182 (2016).
48. Nadel, L., Samsonovich, A., Ryan, L. & Moscovitch, M. Multiple trace theory of human memory: computational, neuroimaging, and neuropsychological results. *Hippocampus*, **10**, 352-368 (2000).
49. Squire, L. R., Genzel, L., Wixted, J. T. & Morris, R. G. Memory consolidation. *Cold Spring Harbor Persp. Biol.* **7**, a021766 (2015).
50. Ritchey, M., Libby, L. A. & Ranganath, C. Cortico-hippocampal systems involved in memory and cognition: the PMAT framework. *Progr. Brain Res.* **219**, 45-64 (2015).
51. Fama, R. & Sullivan, E. V. Thalamic structures and associated cognitive functions: Relations with age and aging. *Neurosci. Biobehav. Rev.* **54**, 29-37 (2015).
52. Alderson, T., et al. Disrupted thalamus white matter anatomy and posterior default mode network effective connectivity in amnesic mild cognitive impairment. *Front. Aging Neurosci.* **9**, 370 (2017).

53. Aggleton, J. P., Pralus, A., Nelson, A. J. & Hornberger, M. Thalamic pathology and memory loss in early Alzheimer's disease: moving the focus from the medial temporal lobe to Papez circuit. *Brain*, **139**, 1877-1890 (2016).
54. Braak, H. & Braak, E. Alzheimer's disease affects limbic nuclei of the thalamus. *Acta Neuropath.* **81**, 261-268 (1991).
55. Forno, G., Lladó, A. & Hornberger, M. Going round in circles—The Papez circuit in Alzheimer's disease. *Eur. J. Neurosci.* **54**, 7668-7687 (2021).
56. Elvsåshagen, T. et al. The genetic architecture of the human thalamus and its overlap with ten common brain disorders. *Nature Commun.* **12**, 1-9 (2021).
57. Siniatchkin, M., Coropceanu, D., Moeller, F., Boor, R. & Stephani, U. EEG-fMRI reveals activation of brainstem and thalamus in patients with Lennox-Gastaut syndrome. *Epilepsia* **52**, 766-774 (2011).
58. Perry, J. C., Pakkenberg, B. & Vann, S. D. Striking reduction in neurons and glial cells in anterior thalamic nuclei of older patients with Down syndrome. *Neurobiol. Aging* **75**, 54-61 (2019).
59. O'Mara, S. M. & Aggleton, J. P. Space and memory (far) beyond the hippocampus: many subcortical structures also support cognitive mapping and mnemonic processing. *Front. Neur. Circ.* **13**, 52 (2019).
60. Wright, N. F., Vann, S. D., Erichsen, J. T., O'Mara, S. M. & Aggleton, J. P. Segregation of parallel inputs to the anteromedial and anteroventral thalamic nuclei of the rat. *J. Comp. Neurol.* **521**, 2966-2986 (2013).
61. Christiansen, K., Dillingham, C. M., Wright, N. F., Saunders, R. C., Vann, S. D., & Aggleton, J. P. Complementary subicular pathways to the anterior thalamic nuclei and mammillary bodies in the rat and macaque monkey brain. *Eur. J. Neurosci.* **43**, 1044-1061 (2016).
62. Mathiasen, M. L., Nelson, A. J., Amin, E., O'Mara, S. M. & Aggleton, J. P. A direct comparison of afferents to the rat anterior thalamic nuclei and nucleus reuniens: overlapping but different. *ENEURO*, **8**. ENEURO.0103-20.2021 (2021).
63. Shibata, H. Direct projections from the anterior thalamic nuclei to the retrohippocampal region in the rat. *J. Comp. Neurol.* **337**, 431-445 (1993).
64. Bubb, E. J., Kinnavane, L. & Aggleton, J. P. Hippocampal–diencephalic–cingulate networks for memory and emotion: An anatomical guide. *Brain Neurosci. Adv.* **1**, 2398212817723443 (2017).
65. Cenquizca, L. A. & Swanson, L. W. Analysis of direct hippocampal cortical field CA1 axonal projections to diencephalon in the rat. *J. Comp. Neurol.* **497**, 101-114 (2006).
66. Vetere, G., Xia, F., Ramsaran, A. I., Tran, L. M., Josselyn, S. A., & Frankland, P. W. An inhibitory hippocampal–thalamic pathway modulates remote memory retrieval. *Nat. Neurosci.* **24**, 685-693 (2021).
67. Ferguson, M. A. et al. A human memory circuit derived from brain lesions causing amnesia. *Nature Commun.* **10**, 1-9 (2019).
68. Wolff, M. & Vann, S.D. The cognitive thalamus as a gateway to mental representations. *J. Neurosci.* **39** 3-14 (2019).

69. Mitchell, A. S. & Chakraborty, S. What does the mediodorsal thalamus do? *Front. Syst. Neurosci.* **7**, 37 (2013).
70. Perry, B. A., Lomi, E. & Mitchell, A. S. Thalamocortical interactions in cognition and disease: the mediodorsal and anterior thalamic nuclei. *Neurosci. Biobehav. Rev.* **130**, 162-177 (2021).
71. Mathiasen, M. L., O'Mara, S. M. & Aggleton, J. P. The anterior thalamic nuclei and nucleus reuniens: So similar but so different. *Neurosci. Biobehav. Rev.* **119**, 268-280 (2020).
72. Moreau, P. H. et al. Lesions of the anterior thalamic nuclei and intralaminar thalamic nuclei: place and visual discrimination learning in the water maze. *Brain Struct. Funct.* **218**, 657-667 (2013).
73. Warburton, E. C. & Aggleton, J. P. Differential deficits in the Morris water maze following cytotoxic lesions of the anterior thalamus and fornix transection. *Behav. Brain Res.* **98**, 27-38 (1998).
74. Dolleman-van der Weel, M. J., Morris, R. G., & Witter, M. P. Neurotoxic lesions of the thalamic reuniens or mediodorsal nucleus in rats affect non-mnemonic aspects of watermaze learning. *Brain Struct. Funct.* **213**, 329-342 (2009).
75. Loureiro, M. et al. The ventral midline thalamus (reuniens and rhomboid nuclei) contributes to the persistence of spatial memory in rats. *J. Neurosci.* **32**, 9947-9959 (2012).
76. McKenna, J. T. & Vertes, R. P. Afferent projections to nucleus reuniens of the thalamus. *J. Comp. Neurol.* **480**, 115-142 (2004).
77. Mathiasen, M. L., Amin, E., Nelson, A. J., Dillingham, C. M., O'Mara, S. M. & Aggleton, J. P. Separate cortical and hippocampal cell populations target the rat nucleus reuniens and mammillary bodies. *Eur. J. Neurosci.* **49**, 1649-1672 (2019).
78. Prasad, J. A. & Chudasama, Y. Viral tracing identifies parallel disynaptic pathways to the hippocampus. *J. Neurosci.* **33**, 8494-8503 (2013).
79. Griffin, A. L. The nucleus reuniens orchestrates prefrontal-hippocampal synchrony during spatial working memory. *Neurosci. Biobehav. Rev.* **128**, 415-420 (2021).
80. Cassel, J. C. et al. The reuniens and rhomboid nuclei of the thalamus: a crossroads for cognition-relevant information processing? *Neurosci. Biobehav. Rev.* **126**, 338-360 (2021).
81. Mitchell, A. S. The mediodorsal thalamus as a higher order thalamic relay nucleus important for learning and decision-making. *Neurosci. Biobehav. Rev.* **54**, 76-88 (2015).
82. Hunt, P. R. & Aggleton, J. P. Medial dorsal thalamic lesions and working memory in the rat. *Behav. Neural Biol.* **55**, 227-246 (1991).
83. Aggleton, J. P., Dumont, J. R. & Warburton, E. C. Unraveling the contributions of the diencephalon to recognition memory: a review. *Learn. Mem.* **18**, 384-400 (2011).
84. Segobin, S. et al. Dissociating thalamic alterations in alcohol use disorder defines specificity of Korsakoff's syndrome. *Brain* **142**, 1458-1470 (2019).
85. Van der Werf, Y. D., Scheltens, P., Lindeboom, J., Witter, M. P., Uylings, H. B. & Jolles, J. Deficits of memory, executive functioning and attention following infarction in the thalamus; a study of 22 cases with localised lesions. *Neuropsychologia* **41**, 1330-1344 (2003).
86. Carlesimo, G. A., Lombardi, M. G. & Caltagirone, C. Vascular thalamic amnesia: a reappraisal. *Neuropsychologia* **49**, 777-789 (2011).

87. Mair, R. G., Burk, J. A. & Porter, M. C. Impairment of radial maze delayed nonmatching after lesions of anterior thalamus and parahippocampal cortex. *Behav. Neurosci.* **117**, 596-605 (2003).
88. Mitchell, A. S. & Dalrymple-Alford, J. C. Lateral and anterior thalamic lesions impair independent memory systems. *Learn. Mem.* **13**, 388-396 (2006).
89. Clark, B. J. & Harvey, R. E. Do the anterior and lateral thalamic nuclei make distinct contributions to spatial representation and memory? *Neurobiol. Learn. Mem.* **133**, 69-78 (2016).
90. Aggleton, J. P., Vann, S. D., Oswald, C. J. & Good, M. Identifying cortical inputs to the rat hippocampus that subserve allocentric spatial processes: a simple problem with a complex answer. *Hippocampus* **10**, 466-474 (2000).
91. Moran, J. P. & Dalrymple-Alford, J. C. Perirhinal cortex and anterior thalamic lesions: comparative effects on learning and memory. *Behav. Neurosci.* **117**, 1326-1341 (2003).
92. Cassel, J. C., Duconseille, E., Jeltsch, H. & Will, B. The fimbria-fornix/cingulate bundle pathways: a review of neurochemical and behavioural approaches using lesions and transplantation techniques. *Progr. Neurobiol.* **51**, 663-716 (1997).
93. Aggleton, J. P. & Brown, M. W. In *Neuropsychology of Memory* (eds. Squire, L.R. & Schacter, D.L) 377–394. (The Guilford Press. New York 2002).
94. Aggleton, J. P., Neave, N., Nagle, S. & Hunt, P. R. A comparison of the effects of anterior thalamic, mamillary body and fornix lesions on reinforced spatial alternation. *Behav. Brain Res.* **68**, 91-101 (1995).
95. Parker, A. & Gaffan, D. The effect of anterior thalamic and cingulate cortex lesions on object-in-place memory in monkeys. *Neuropsychologia* **35**, 1093-1102 (1997).
96. Gaffan, D. Scene-specific memory for objects: a model of episodic memory impairment in monkeys with fornix transection. *J. Cognitive Neurosci.* **6**, 305-320 (1994).
97. Sutherland, R. J. & Rodriguez, A. J. The role of the fornix/fimbria and some related subcortical structures in place learning and memory. *Behav. Brain Res.* **32**, 265-277 (1989).
98. Aggleton, J. P., Keith, A. B. & Sahgal, A. Both fornix and anterior thalamic, but not mammillary, lesions disrupt delayed non-matching-to-position memory in rats. *Behav. Brain Res.* **44**, 151-161(1991).
99. Dumont, J. R., Wright, N. F., Pearce, J. M. & Aggleton, J. P. The impact of anterior thalamic lesions on active and passive spatial learning in stimulus controlled environments: geometric cues and pattern arrangement. *Behav. Neurosci.* **128**, 161-177 (2014).
100. Pearce, J. M., Good, M. A., Jones, P. M. & McGregor, A. Transfer of spatial behavior between different environments: implications for theories of spatial learning and for the role of the hippocampus in spatial learning. *J. Exp. Psych: Anim. Behav. Proc.* **30**, 135 (2004).
101. Aggleton, J. P., Keith, A. B., Rawlins, J. N. P., Hunt, P. R. & Sahgal, A. Removal of the hippocampus and transection of the fornix produce comparable deficits on delayed non-matching to position by rats. *Behav. Brain Res.* **52**, 61-71 (1992).

102. Song, D., Harway, M., Marmarelis, V. Z., Hampson, R. E., Deadwyler, S. A. & Berger, T. W. Extraction and restoration of hippocampal spatial memories with non-linear dynamical modeling. *Front. Syst. Neurosci.*, **8**, 97 (2014).
103. Morris, R. G. M., Schenk, F., Tweedie, F. & Jarrard, L. E. Ibotenate lesions of hippocampus and/or subiculum: dissociating components of allocentric spatial learning. *Eur. J. Neurosci.* **2**, 1016-1028 (1990).
104. Jarrard, L. E. What does the hippocampus really do? *Behav. Brain Res.* **71**, 1-10 (1995).
105. Taube, J. S. Head direction cells recorded in the anterior thalamic nuclei of freely moving rats. *J. Neurosci.* **15**, 70-86 (1995).
106. Calton, J. L., Stackman, R. W., Goodridge, J. P., Archey, W. B., Dudchenko, P. A. & Taube, J. S. Hippocampal place cell instability after lesions of the head direction cell network. *J. Neurosci.* **23**, 9719-9731 (2003).
107. Moser, E. I., Roudi, Y., Witter, M. P., Kentros, C., Bonhoeffer, T. & Moser, M. B. Grid cells and cortical representation. *Nat. Rev. Neurosci.* **15**, 466-481 (2014).
108. Golob, E. J. & Taube, J. S. Head direction cells and episodic spatial information in rats without a hippocampus. *Proc. Natl Acad. Sci. USA* **94**, 7645-7650 (1997).
109. Safari, V. et al. Individual subnuclei of the rat anterior thalamic nuclei differently affect spatial memory and passive avoidance tasks. *Neurosci.* **444**, 19-32 (2020).
110. Vann, S. D. Transient spatial deficit associated with bilateral lesions of the lateral mammillary nuclei. *Eur. J. Neurosci.* **21**, 820-824 (2005).
111. Dillingham, C. M. & Vann, S.D. Why isn't the head direction system necessary for direction? lessons from the lateral mammillary nuclei. *Front. Neur. Circ.* **13**, 60 (2019).
112. van Groen, T., Kadish, I. & Wyss, J. M. Role of the anterodorsal and anteroventral nuclei of the thalamus in spatial memory in the rat. *Behav. Brain Res.* **132**, 19-28 (2002).
113. Moser, E. I., Moser, M. B. & McNaughton, B. L. Spatial representation in the hippocampal formation: a history. *Nat. Neurosci.* **20**, 1448-1464 (2017).
114. Matulewicz, P., Ulrich, K., Islam, M. N., Mathiasen, M. L., Aggleton, J. P. & O'Mara, S. M. Proximal perimeter encoding in the rat rostral thalamus. *Sci. Rep.* **9**, 1-12 (2019).
115. Jankowski, M. M., Passecker, J., Islam, M. N., Vann, S., Erichsen, J. T., Aggleton, J. P. & O'Mara, S. M. Evidence for spatially-responsive neurons in the rostral thalamus. *Front. Behav. Neurosci.* **9**, 256 (2015).
116. Tsanov, M. et al. Theta-modulated head direction cells in the rat anterior thalamus. *J. Neurosci.* **31**, 9489-9502 (2011).
117. Horikawa, K., Kinjo, N., Stanley, L. C. & Powell, E. W. Topographic organization and collateralization of the projections of the anterior and laterodorsal thalamic nuclei to cingulate areas 24 and 29 in the rat. *Neurosci. Res.* **6** 31-44 (1988).
118. Gibson, W. S., Ross, E. K., Han, S. R., Van Gompel, J. J., Min, H. K. & Lee, K. H. Anterior thalamic deep brain stimulation: functional activation patterns in a large animal model. *Brain Stim.* **9**, 770-773. (2016).

119. Horner, A. J., Bisby, J. A., Wang, A., Bogus, K. & Burgess, N. The role of spatial boundaries in shaping long-term event representations. *Cognition* **154**, 151-164 (2016).
120. Tsanov, M., Vann, S. D., Erichsen, J. T., Wright, N., Aggleton, J. P. & O'Mara, S. M. Differential regulation of synaptic plasticity of the hippocampal and the hypothalamic inputs to the anterior thalamus. *Hippocampus*, **21**, 1-8 (2011).
121. Bauch, E. M. et al. Theta oscillations underlie retrieval success effects in the nucleus accumbens and anterior thalamus: evidence from human intracranial recordings. *Neurobiol. Learn. Mem.* **155**, 104-112 (2018).
122. Sweeney-Reed, C. M. et al. Corticothalamic phase synchrony and cross-frequency coupling predict human memory formation. *Elife*, **3**, e05352 (2014).
123. Sweeney-Reed, C. M. et al. Thalamic theta phase alignment predicts human memory formation and anterior thalamic cross-frequency coupling. *Elife*, **4**, e07578 (2015).
124. Sweeney-Reed, C. M. et al. The role of the anterior nuclei of the thalamus in human memory processing. *Neurosci. Biobehav. Rev.* **126**, 146-158 (2021).
125. Li, A. W. & King, J. Spatial memory and navigation in ageing: A systematic review of MRI and fMRI studies in healthy participants. *Neurosci. Biobehav. Rev.* **103**, 33-49 (2019).
126. Fritch, H. A. et al. The anterior hippocampus is associated with spatial memory encoding. *Brain Res.* **1732**, 146696 (2020).
127. Petersen, R. C. et al. Memory and MRI-based hippocampal volumes in aging and AD. *Neurology* **54**, 581-581 (2000).
128. Rugg, M. D., Vilberg, K. L., Mattson, J. T., Sarah, S. Y., Johnson, J. D. & Suzuki, M. Item memory, context memory and the hippocampus: fMRI evidence. *Neuropsychologia* **50**, 3070-3079 (2012).
129. Horner, A. J. & Doeller, C. F. Plasticity of hippocampal memories in humans. *Curr. Opin. Neurobiol.* **43**, 102-109 (2017).
130. Geier, K. T., Buchsbaum, B. R., Parimoo, S. & Olsen, R. K. The role of anterior and medial dorsal thalamus in associative memory encoding and retrieval. *Neuropsychologia*, **148**, 107623 (2020).
131. Spets, D. S. & Slotnick, S. D. Thalamic functional connectivity during spatial long-term memory and the role of sex. *Brain Sci.* **10**, 898 (2020).
132. Kafkas, A., Mayes, A. R. & Montaldi, D. Thalamic-medial temporal lobe connectivity underpins familiarity memory. *Cereb. Cortex* **30**, 3827-3837 (2020).
133. Pergola, G., Ranft, A., Mathias, K. & Suchan, B. The role of the thalamic nuclei in recognition memory accompanied by recall during encoding and retrieval: an fMRI study. *Neuroimage* **74**, 195-208 (2013).
134. Su, J. H. et al. Thalamus optimized multi atlas segmentation (Thomas): fast, fully automated segmentation of thalamic nuclei from structural MRI. *Neuroimage* **194**, 272-282 (2019).
135. Choi, S. H., Kim, Y. B., Paek, S. H. & Cho, Z. H. Papez circuit observed by in vivo human brain with 7.0 T MRI super-resolution track density imaging and track tracing. *Front. Neuroanat.* **13**, 17. (2019).

136. Iglehart, C., Monti, M., Cain, J., Tourdias, T. & Saranathan, M. A systematic comparison of structural-, structural connectivity-, and functional connectivity-based thalamus parcellation techniques. *Brain Struct. Funct.* **225**, 1631-1642 (2020).
137. Albo, Z., Di Prisco, G. V., & Vertes, R. P. Anterior thalamic unit discharge profiles and coherence with hippocampal theta rhythm. *Thal. Related Syst.* **2**, 133-144 (2003).
138. Phillips, J. W. et al. A repeated molecular architecture across thalamic pathways. *Nat. Neurosci.* **22**, 1925-1935 (2019).
139. Aggleton, J. P., Hunt, P. R., Nagle, S. & Neave, N. The effects of selective lesions within the anterior thalamic nuclei on spatial memory in the rat. *Behav. Brain Res.* **81**, 189-198 (1996).
140. Byatt, G. & Dalrymple-Alford, J. C. Both anteromedial and anteroventral thalamic lesions impair radial-maze learning in rats. *Behav. Neurosci.* **110**, 1335-1348 (1996).
141. Wilton, L. A. K., Baird, A. L., Muir, J. L., Honey, R. C. & Aggleton, J. P. Loss of the thalamic nuclei for "head direction" impairs performance on spatial memory tasks in rats. *Behav. Neurosci.* **115**, 861-869 (2001).
142. Cullen, K. E. & Taube, J. S. Our sense of direction: progress, controversies and challenges. *Nat. Neurosci.* **20**, 1465 -1473 (2017).
143. Viejo, G. & Peyrache, A. Precise coupling of the thalamic head-direction system to hippocampal ripples. *Nature Commun.* **11**, 1-14 (2020).
144. van Groen, T., Kadish, I. & Wyss, J. M. Efferent connections of the anteromedial nucleus of the thalamus of the rat. *Brain Res. Rev.* **30**, 1-26 (1999).
145. Barbas, H., Henion, T. H. & Dermon, C. R. Diverse thalamic projections to the prefrontal cortex in the rhesus monkey. *J. Comp. Neurol.* **313**, 65-94 (1991).
146. Shibata, H. & Naito, J. Organization of anterior cingulate and frontal cortical projections to the anterior and laterodorsal thalamic nuclei in the rat. *Brain Res.* **1059**, 93-103 (2005).
147. Wolff, M., Gibb, S. J. & Dalrymple-Alford, J. C. Beyond spatial memory: the anterior thalamus and memory for the temporal order of a sequence of odor cues. *J. Neurosci.* **26**, 2907-2913 (2006).
148. Dumont, J. R. & Aggleton, J. P. Dissociation of recognition and recency memory judgments after anterior thalamic nuclei lesions in rats. *Behav. Neurosci.* **127**, 415-431. (2013).
149. Bubb, E. J., Aggleton, J. P., O'Mara, S. M. & Nelson, A. J. Chemogenetics reveal an anterior cingulate–thalamic pathway for attending to task-relevant information. *Cereb. Cortex* **31**, 2169-2186 (2021).
150. Kim, S. M., Ganguli, S. & Frank, L. M. Spatial information outflow from the hippocampal circuit: distributed spatial coding and phase precession in the subiculum. *J. Neurosci.* **32**, 11539-11558 (2012).
151. Poulter, S., Lee, S. A., Dachtler, J., Wills, T. J. & Lever, C. Vector trace cells in the subiculum of the hippocampal formation. *Nat. Neurosci.* **24**, 266-275 (2021).
152. Tsanov, M., Wright, N., Vann, S. D., Erichsen, J. T., Aggleton, J. P. & O'Mara, S. M. Hippocampal inputs mediate theta-related plasticity in anterior thalamus. *Neuroscience* **187**, 52-62 (2011).

153. Dillingham, C. M. et al. Mammillothalamic disconnection alters hippocampocortical oscillatory activity and microstructure: implications for diencephalic amnesia. *J. Neurosci.* **39**, 6696-6713 (2019).
154. Frost, B. E., Martin, S. K., Cafalchio, M., Islam, M. N., Aggleton, J. P. & O'Mara, S. M. Anterior thalamic function is required for spatial coding in the subiculum and is necessary for spatial memory. *J. Neurosci.* **41**, 6511-6525 (2021).
155. Hani, S. A. H. B., Al-Haidari, M. H. & Saboba, M. M. Neuronal types in the human anterior ventral thalamic nucleus: a Golgi study. *Cell. Mol. Neurobiol.* **27**, 745-755 (2007).
156. Gaffan, D. & Gaffan, E. A. Amnesia in man following transection of the fornix: a review. *Brain* **114**, 2611-2618 (1991).
157. Aggleton, J. P., et al. Differential cognitive effects of colloid cysts in the third ventricle that spare or compromise the fornix. *Brain* **123**, 800-815 (2000).
158. Warburton, E. C., Baird, A. L., Morgan, A., Muir, J. L. & Aggleton, J. P. Disconnecting hippocampal projections to the anterior thalamus produces deficits on tests of spatial memory in rats. *Eur. J. Neurosci.*, **12**, 1714-1726 (2000).
159. Henry, J., Petrides, M., St-Laurent, M. & Sziklas, V. Spatial conditional associative learning: effects of thalamo-hippocampal disconnection in rats. *Neuroreport* **15**, 2427-2431 (2004).
160. Kitanishi, T., Umaba, R. & Mizuseki, K. Robust information routing by dorsal subiculum neurons. *Sci. Adv.* **7**, eabf1913 (2021).
161. Rudebeck, S. R., Scholz, J., Millington, R., Rohenkohl, G., Johansen-Berg, H. & Lee, A. C. Fornix microstructure correlates with recollection but not familiarity memory. *J. Neurosci.* **29**, 14987-14992 (2009).
162. Metzler-Baddeley, C., Jones, D. K., Belaroussi, B., Aggleton, J. P. & O'Sullivan, M. J. Frontotemporal connections in episodic memory and aging: a diffusion MRI tractography study. *J. Neurosci.* **31**, 13236-13245 (2011).
163. Hartopp, N. et al. A key role for subiculum-fornix connectivity in recollection in older age. *Front. Syst. Neurosci.* **12**, 70 (2019).
164. Hodgetts, C. J. et al. The role of the fornix in human navigational learning. *Cortex*, **124**, 97-110 (2020).
165. Christiansen, K., Aggleton, J. P., Parker, G. D., O'Sullivan, M. J., Vann, S. D. & Metzler-Baddeley, C. The status of the precommissural and postcommissural fornix in normal ageing and mild cognitive impairment: An MRI tractography study. *NeuroImage* **130**, 35-47 (2016).
166. Coad, B. M., Craig, E., Louch, R., Aggleton, J. P., Vann, S. D. & Metzler-Baddeley, C. Precommissural and postcommissural fornix microstructure in healthy aging and cognition. *Brain Neurosci. Adv.* **4**, 2398212819899316 (2020).
167. Commins, S., Gigg, J., Anderson, M. & O'Mara, S. M. Interaction between paired-pulse facilitation and long-term potentiation in the projection from hippocampal area CA1 to the subiculum. *Neuroreport* **9**, 4109-4113 (1998).
188. Liu, J. et al. Anterior thalamic stimulation improves working memory precision judgments. *Brain Stim.* **14**, 1073-1080 (2021).

169. Xiao, D. & Barbas, H. Pathways for emotions and memory II. Afferent input to the anterior thalamic nuclei from prefrontal, temporal, hypothalamic areas and the basal ganglia in the rhesus monkey. *Thal. Related Syst.* **2**, 33-48 (2002).
170. Jones, B. F. & Witter, M. P. Cingulate cortex projections to the parahippocampal region and hippocampal formation in the rat. *Hippocampus* **17**, 957-976 (2007).
171. Sugar, J., Witter, M. P., van Strien, N. & Cappaert, N. The retrosplenial cortex: intrinsic connectivity and connections with the (para) hippocampal region in the rat. An interactive connectome. *Front. Neuroinf.* **5**, 7 (2011).
172. Dillingham, C. M., Milczarek, M. M., Perry, J. C. & Vann, S. D. Time to put the mammillothalamic pathway into context. *Neurosci. Biobehav. Rev.* **121**, 60-74 (2021).
173. Poirier, G.L. et al. Anterior thalamic lesions produce chronic and profuse transcriptional de-regulation in retrosplenial cortex: A model of retrosplenial hypoactivity and covert pathology. *Thal. Related Syst.* **4**, 59-77 (2008).
174. Poirier, G.L., Aggleton, J.P. Post-surgical interval and lesion location within the limbic thalamus determine extent of retrosplenial cortex hypoactivity. *Neurosci.* **160**, 452-469 (2009).
175. Vann, S. D. Dismantling the Papez circuit for memory in rats. *Elife*, **2**, e00736 (2013).
176. Yamawaki, N., Corcoran, K. A., Guedea, A. L., Shepherd, G. M. & Radulovic, J. Differential contributions of glutamatergic hippocampal→ retrosplenial cortical projections to the formation and persistence of context memories. *Cereb. Cortex* **29**, 2728-2736 (2019).
177. Aggleton, J. P. Multiple anatomical systems embedded within the primate medial temporal lobe: implications for hippocampal function. *Neurosci. Biobehav. Rev.* **36**, 1579-1596 (2012).
178. Jay, T. M. & Witter, M. P. Distribution of hippocampal CA1 and subicular efferents in the prefrontal cortex of the rat studied by means of anterograde transport of Phaseolus vulgaris-leucoagglutinin. *J. Comp. Neurol.* **313**, 574-586 (1991).
179. Raichle, M. E. The brain's default mode network. *Ann. Rev. Neurosci.* **38**, 433-447 (2015).
180. Mak, L. E., Minuzzi, L., MacQueen, G., Hall, G., Kennedy, S. H. & Milev, R. The default mode network in healthy individuals: a systematic review and meta-analysis. *Brain Conn.* **7**, 25-33 (2017).
181. Andrews-Hanna, J. R., Reidler, J. S., Sepulcre, J., Poulin, R. & Buckner, R. L. Functional-anatomic fractionation of the brain's default network. *Neuron* **65**, 550-562 (2010).
182. Wen, T., Mitchell, D. J. & Duncan, J. The functional convergence and heterogeneity of social, episodic, and self-referential thought in the default mode network. *Cereb. Cortex* **30**, 5915-5929 (2020).
183. Yeshurun, Y., Nguyen, M. & Hasson, U. The default mode network: where the idiosyncratic self meets the shared social world. *Nat. Rev. Neurosci.* **22**, 181-192 (2021).
184. Buckner, R. L., Andrews-Hanna, J. R. & Schacter, D. L. In *The Year in Cognitive Neuroscience* (eds. Kingstone, A. & Miller, M.B.) 1–38 (Blackwell Publishing, 2008).
185. Schacter, D. L., Addis, D. R. & Buckner, R. L. Remembering the past to imagine the future: the prospective brain. *Nat. Rev. Neurosci.* **8**, 657-661 (2007).

186. Schacter, D. L., Addis, D. R., Hassabis, D., Martin, V. C., Spreng, R. N. & Szpunar, K. K. The future of memory: remembering, imagining, and the brain. *Neuron* **76**, 677-694 (2012).
187. Alves, P. N., Foulon, C., Karolis, V., Bzdok, D., Margulies, D. S., Volle, E. & de Schotten, M. T. An improved neuroanatomical model of the default-mode network reconciles previous neuroimaging and neuropathological findings. *Comm. Biol.* **2**, 1-14 (2019).
188. Li, J., Curley et al. Mapping the subcortical connectivity of the human default mode network. *NeuroImage* **245**, 118758 (2021).
189. Jones, D. T., Mateen, F. J., Lucchinetti, C. F., Jack, C. R. & Welker, K. M. Default mode network disruption secondary to a lesion in the anterior thalamus. *Arch. Neurol.* **68**, 242-247 (2011).
190. Middlebrooks, E. H. et al. Functional activation patterns of deep brain stimulation of the anterior nucleus of the thalamus. *World Neurosurg.* **136**, 357-363 (2020).
191. Kaboodvand, N., Bäckman, L., Nyberg, L. & Salami, A. The retrosplenial cortex: A memory gateway between the cortical default mode network and the medial temporal lobe. *Hum. Brain Map.* **39**, 2020-2034 (2018).
192. Garden, D. L. et al. (2009). Anterior thalamic lesions stop synaptic plasticity in retrosplenial cortex slices: expanding the pathology of diencephalic amnesia. *Brain* **132**, 1847-1857.
193. Williams, A. N., Ridgeway, S., Postans, M., Graham, K. S., Lawrence, A. D. & Hodgetts, C. J. The role of the pre-commissural fornix in episodic autobiographical memory and simulation. *Neuropsychologia* **142**, 107457 (2020).
194. Corkin, S. Beware of frontal lobe deficits in hippocampal clothing. *Trends Cogn. Sci.* **5**, 321-323 (2001).
195. Hermann, B. & Seidenberg, M. Executive system dysfunction in temporal lobe epilepsy: effects of nociferous cortex versus hippocampal pathology. *J. Clin. Exp. Neuropsych.* **17**, 809-819 (1995).
196. Pearce, J. M. & Mackintosh, N. J. In *Attention and Associative Learning: From Brain to Behaviour* (eds. Mitchell, C. & Le Pelley, M.) 11-39 (Oxford University Press, Oxford, 2010).
197. Dias, R., Robbins, T. W. & Roberts, A. C. Dissociation in prefrontal cortex of affective and attentional shifts. *Nature* **380**, 69-72 (1996).
198. Birrell, J. M. & Brown, V. J. Medial frontal cortex mediates perceptual attentional set shifting in the rat. *J. Neurosci.* **20**, 4320-4324 (2000).
199. Barbas, H. & Zikopoulos, B. The prefrontal cortex and flexible behavior. *Neuroscientist* **13**, 532-545 (2007).
200. Marquis, J. P., Goulet, S. & Doré, F. Y. Neonatal ventral hippocampus lesions disrupt extra-dimensional shift and alter dendritic spine density in the medial prefrontal cortex of juvenile rats. *Neurobiol. Learn. Mem.* **90**, 339-346 (2008).
201. Nelson, A. J. The anterior thalamic nuclei and cognition: a role beyond space? *Neurosci. Biobehav. Rev.* **126**, 1-11 (2021).
202. Wolff, M., Alcaraz, F., Marchand, A. R. & Coutureau, E. Functional heterogeneity of the limbic thalamus: from hippocampal to cortical functions. *Neurosci. Biobehav. Rev.* **54**, 120-130. (2015).

203. Albasser, M. M., Amin, E., Lin, T. C. E., Iordanova, M. D. & Aggleton, J. P. Evidence that the rat hippocampus has contrasting roles in object recognition memory and object recency memory. *Behav. Neurosci.* **126**, 659-669 (2012).
204. Dupire, A. et al. A role for anterior thalamic nuclei in affective cognition: interaction with environmental conditions. *Hippocampus* **23**, 392-404 (2013).
205. Gabriel, M. & Talk, A. C. In *Model Systems and the Neurobiology of Associative Learning: A Festschrift in Honor of Richard F. Thompson* (eds. Steinmetz, J.E. Gluck, M.A. & Solomon, P.R.) 149-185. (Lawrence Erlbaum Associates Publishers, New Jersey, 2001).
206. Shin, J. N., Doron, G. & Larkum, M. E. Memories off the top of your head. *Science* **374**, 538-539 (2021).
207. Witter, M. P. & Groenewegen, H. J. Connections of the parahippocampal cortex in the cat. III. Cortical and thalamic efferents. *J. Comp. Neurol.* **252**, 1-31 (1986).
208. Brennan, E. K., Jedrasiak-Cape, I., Kailasa, S., Rice, S. P., Sudhakar, S. K. & Ahmed, O. J. Thalamus and claustrum control parallel layer 1 circuits in retrosplenial cortex. *eLife* **10**, e62207 (2021).
209. Kinnavane, L., Vann, S. D., Nelson, A. J., O'Mara, S. M. & Aggleton, J. P. Collateral projections innervate the mammillary bodies and retrosplenial cortex: a new category of hippocampal cells. *ENEURO*.0383-17.2018.
210. Zoppelt, D., Koch, B., Schwarz, M. & Daum, I. Involvement of the mediodorsal thalamic nucleus in mediating recollection and familiarity. *Neuropsychologia* **41**, 1160-1170 (2003).
211. Danet, L. et al. Medial thalamic stroke and its impact on familiarity and recollection. *Elife* **6**, e28141 (2017).
212. Carlesimo, G. A., Lombardi, M. G., Caltagirone, C. & Barban, F. Recollection and familiarity in the human thalamus. *Neurosci. Biobehav. Rev.* **54**, 18-28 (2015).
213. Grodd, W., Kumar, V. J., Schüz, A., Lindig, T. & Scheffler, K. The anterior and medial thalamic nuclei and the human limbic system: tracing the structural connectivity using diffusion-weighted imaging. *Sci. Rep.* **10**, 1-25 (2020).
214. Gagnepain, P. et al. Collective memory shapes the organization of individual memories in the medial prefrontal cortex. *Nat. Hum. Behav.* **4**, 189-200 (2020).
215. Barnett, S. C. et al. Anterior thalamic nuclei neurons sustain memory. *Curr. Res. Neurobiol.* **2**, 100022 (2021).
216. Lozsádi, D. A. Organization of connections between the thalamic reticular and the anterior thalamic nuclei in the rat. *J. Comp. Neurol.* **358**, 233-246 (1995).
217. Shibata, H. Topographic organization of subcortical projections to the anterior thalamic nuclei in the rat. *J. Comp. Neurol.* **323**, 117-127 (1992).
218. Swanson, L. W. & Cowan, W. M. An autoradiographic study of the organization of the efferent connections of the hippocampal formation in the rat. *J. Comp. Neurol.* **172**, 49-84 (1977).
219. van Groen, T. & Michael Wyss, J. Connections of the retrosplenial granular a cortex in the rat. *J. Comp. Neurol.* **300**, 593-606 (1990).

220. van Groen, T. & Wyss, J. M. Connections of the retrosplenial dysgranular cortex in the rat. *J. Comp. Neurol.* **315**, 200-216 (1992).
221. Poletti, C. E. & Creswell, G. Fornix system efferent projections in the squirrel monkey: an experimental degeneration study. *J. Comp. Neurol.* **175**, 101-127 (1977).
222. Shah, A., Jhavar, S. S. & Goel, A. Analysis of the anatomy of the Papez circuit and adjoining limbic system by fiber dissection techniques. *J. Clin. Neurosci.* **19**, 289-298 (2012).
223. Jankowski, M.M. et al. The anterior thalamus provides a subcortical circuit supporting memory and spatial navigation. *Front. Syst. Neurosci.* **7**, 45 (2013).

Acknowledgements

J.P.A. and S.M.O'M. are supported in the work described here by the Wellcome Trust (103722/Z14/Z). We thank S. Commins, S. Martin, A. Nelson, and C. Ranganath for very helpful feedback on a prior version of the manuscript.

Author contributions

The authors contributed equally to all aspects of the article.

Competing interests

The authors declare no competing interests.

Table 1 Contrasting properties of the three principal anterior thalamic nuclei

Property	Anterodorsal nucleus	Anteromedial nucleus	Anteroventral nucleus
Gene expression	Unique expression profile ¹³⁸	Expression profile corresponds to a cluster of 8 thalamic nuclei ¹³⁸	Expression profile corresponds to a cluster of 5 thalamic nuclei ¹³⁸
Electrophysiology	Head-direction cells predominate ¹⁰⁵ Activity associated with hippocampal sharp wave ripples ¹⁴³	Place cells ^{59,115} and perimeter cells ¹¹⁴ identified	Head-direction cells (modest numbers) ¹¹⁶ Theta oscillations ^{116,137}
Cortical and hippocampal inputs*‡	Presubiculum ^{64,218} Postsubiculum ^{64,218} Hippocampal area CA3 (sparse) ⁶⁶ Retrosplenial area 29 ^{§64,219}	Dorsal subiculum (proximal) ^{61,64,218} CA1 (sparse) ⁶⁵ Retrosplenial area 30 ^{64,220} Anterior cingulate cortex ^{60,146} Prelimbic cortex ^{60,146}	Dorsal subiculum (distal) ^{61,64,218} Retrosplenial area 29 ⁶⁴ Anterior cingulate cortex ^{60,146} Prelimbic cortex ^{60,146}
Subcortical inputs*‡	Lateral mammillary nucleus (bilateral) ^{27,217} Dorsal-most rostral reticular thalamic nucleus ²¹⁶	Medial mammillary nucleus (ipsilateral) ^{27,60,217} Upper half of the rostral reticular thalamic nucleus ²¹⁶	Medial mammillary nucleus (ipsilateral) ^{27,60,217} Dorsal rostral reticular thalamic nucleus ²¹⁶
Projection targets*	Presubiculum ⁶⁴ Postsubiculum ⁶⁴ Retrosplenial area 29 ^{64,117,219} (layers I, III and IV)	Subiculum (ventral) ⁶³ Area 29 ^{117,144, 219} (layers I and V; light input) Area 30 ^{117,144,220} (layers I, V and VI) Prelimbic cortex ¹⁴⁴ (layers I, III, IV and V) Anterior cingulate cortex ¹⁴⁴ (layers I and V)	Subiculum (dorsal) ⁶³ Area 29 ^{17,117, 219} (layers I and IV) Area 30 ^{117,220} (layer IV, light)
Transient lesion effects - passive avoidance	No effect ¹⁰⁹	Deficit in consolidation ¹⁰⁹	Deficits in consolidation and retrieval ¹⁰⁹
Transient lesion effects on spatial memory (Morris water maze)	Deficit in retrieval ¹⁰⁹	Deficit in retrieval ¹⁰⁹	Deficits in encoding, consolidation and retrieval ¹⁰⁹
Permanent lesions effects - spatial working memory	Deficit ¹⁴¹ (but included lateral dorsal nucleus damage)	Modest deficit ^{112,139,140}	Modest deficit ^{112,139,140}


The information in the table is derived from rodent data. *Some very light connections are not included. ‡Note that few individual neurons innervate more than one anterior thalamic nucleus⁵². §Note that retrosplenial cortex comprises areas 29 and 30.

Figure 1. Location of the anterior thalamic nuclei. **A, B.** Visualisation of the anterodorsal (red), anteromedial (yellow), and anteroventral (green) nuclei, along with adjacent thalamic nuclei from MRI images²¹³. Each column shows axial (upper), coronal (mid), sagittal (lower) sections. Those on the left are overlaid on the MNI brain template with corresponding X, Y, and Z coordinates. Those on the right are corresponding 3D rendered views of the selected nuclei overlaid on a thalamic template. **C.** Dissection of the human brain²²² showing midline sagittal view of the medial temporal lobe (lower) and medial diencephalon (upper). The major tracts appear white. **D.** Location and arrangement of thalamic nuclei in the rat brain, with the anterior thalamic nuclei highlighted in colour²²³. Abbreviations: 3V, third ventricle; AD, anterodorsal nucleus; AM, anteromedial nucleus; Amy, amygdala; ATN, anterior thalamic nuclei; AV, anteroventral nucleus; CeM, central medial nucleus; CL, centrolateral nucleus; F, fornix; IAM, interoanteromedial nucleus; ic, internal capsule; LD, laterodorsal nucleus; LGN lateral geniculate nucleus; LP, lateral posterior nucleus; Hb, habenula; HPC, hippocampus; MB, mammillary bodies; MD, mediodorsal nucleus; MGN, medial geniculate nucleus; MNI, Montreal Neurological Institute; MTT, mamillothalamic tract; PC, paracentral nucleus; PCF, postcommissural fornix; Pf, parafascicular nucleus; PT, paratenial nucleus; Pul, pulvinar; PuM, medial pulvinar; PV, paraventricular nucleus; RE, nucleus reuniens; RT, reticular thalamic nucleus; sm, stria medullaris; st, stria terminalis; VA, ventral anterior nucleus; VL, ventrolateral nucleus; VM, ventromedial nucleus; VPL, ventral posterior nucleus, pars lateralis. (A, B. With permission from W. Grodd²¹³; C, D permission from O'Mara²²³).

Fig. 2: Principal connections of the anterior thalamic nuclei. The main diagram depicts the connectivity of the rodent anteromedial nucleus (AM), the anteroventral nucleus (AV), and the anterodorsal (AD) thalamic nucleus^{26,27,60-64,144-146,217,218}. The thickness of the arrows reflects the relative density of the various connections. Dashed arrows indicate connections that do not directly target the anterior thalamic nuclei (ATN) but are potentially important for their contribution to episodic memory. The schematic to the right includes key pathways considered in the text. AC, anterior cingulate cortex; CB, cingulum bundle; F, fornix; HPC, hippocampal formation; LMB, lateral mammillary nucleus; M Bodies, mammillary bodies; MPF, medial prefrontal cortex; MMB, medial mammillary nucleus; MPF, medial prefrontal cortex; MTT, mamillothalamic tract; PHC, parahippocampal region; RSP, retrosplenial cortex.

Fig. 3: Proposed anterior thalamic participation in cognition. Hypothesised mapping of three interleaving, interconnected, 'cognitive zones' of anterior thalamic nuclear influence. Each zone maps cognitive functions — spatial processing^{18-21,37,50,59,87-89,94,105,109,124,154}, attention^{9,10,59,124,130,149}, and memory^{8,9,17,35,37,50,59,83-86,109,149} — to particular anterior nuclei (as confirmed by lesion, recording, anatomical tracing, or other evidence) illustrating their extensive contributions to multiple aspects of cognition. This mapping of functions implies fast-acting, dynamic relations between spatial processing, attentional, and mnemonic processes; spatial processing and attention are deployed during episodic memory encoding and consolidation (as they are contributory processes to memory), but equally they may well be independent of memory during other tasks. In sum, this figure illustrates that individual anterior thalamic nuclei participate in multiple, but particular, cognitive functions (namely, spatial processing, attention, and memory).

Fig. 4: A core, tripartite episodic memory system. An anatomically-simplified schematic of the parallel, interacting circuits supporting our model of a core, tripartite episodic memory system. According to the model, this system comprises a ‘temporal lobe’ stream (hippocampal-centred) and a ‘medial diencephalic’ stream (anterior thalamic-centred) that together act on independent and shared cortical areas (including parahippocampal, prefrontal, and retrosplenial cortices; note that more detailed connections are presented in **Fig. 2**). These mnemonic streams function in parallel and are partially independent of each other. There are cortical zones of anatomical overlap where projections from both streams converge; anterior thalamic nuclei-hippocampus dependent memory depends on these anatomical zones of interaction, where synchronous activity is required for mnemonic consolidation. The rules governing plasticity at many of the synapses in this circuit remain to be comprehensively investigated.



Box 1. Visualisation of fornix pathway and its subdivisions with diffusion imaging.

Diffusion imaging has delivered insights into the properties of fibre pathways associated with the anterior thalamic

nuclei and how they may change in disease states⁸⁴ or relate to individual variations in cognitive performance by healthy participants¹⁶¹⁻¹⁶⁶. Diffusion imaging uses the movement of water molecules to reconstruct white matter pathways and changes in the patterns of diffusion can provide indirect insights into the structural status of the pathway. This technique makes it possible to not only reconstruct the fornix (see the figure, part **A** green) but also to separate those fibres that pass in front (‘precommissural’ **B**) and behind (‘postcommissural’ **C**) the anterior commissure (from¹⁶⁶ with permission). This separation is informative as anatomical studies²²¹ show that the postcommissural fornix contains most of the direct hippocampal projections to the anterior thalamic nuclei along with the indirect projections via the mammillary bodies. To isolate the precommissural fornix (**B** – see also **Fig 1C**), a ‘NOT’ gate (red) is placed immediately behind the anterior commissure (centre of red box in **A** and pink pixel in **B** and **C**). This NOT gate precludes those fibre streams passing behind the commissure. Meanwhile, the AND gate in front of the commissure ensures that just fibre

streams in this region are visualised. The converse arrangement isolates the postcommissural fornix (C). The OR gate makes it possible to identify those fibres in the body of the fornix that pass in front or behind the anterior commissure. In this way it is possible to distinguish hippocampal – prefrontal (precommissural fornix) influences from hippocampal – diencephalic (postcommissural fornix) influences using non-invasive techniques.

