

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/157399/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Nagendrabau, Venkateshbabu, Vinothkumar, Thillar Sekar, Rossi-Fedele, Giampiero, Dogramaci, Esma J., Duncan, Henry F., Abbott, Paul V., Levin, Liran, Lin, Shaul and Dummer, Paul M.H. 2023. Dental patient-reported outcomes following traumatic dental injuries and treatment: a narrative review. *Dental Traumatology* 39 (4) , pp. 304-313. 10.1111/edt.12827

Publishers page: <https://doi.org/10.1111/edt.12827>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Dental patient-reported outcomes following traumatic dental injuries and treatment: a narrative review

Abstract

Dental patient-reported outcomes (dPROs) are self-reported descriptions of a patient's oral health status that are not modified or interpreted by a healthcare professional. Dental patient-reported outcome measures (dPROMs) are objective or subjective measurements used to assess dPROs. In oral healthcare settings, the emphasis on assessing treatment outcomes from the patient's perspective has increased and this is particularly important after traumatic dental injuries (TDIs), as this group of injuries represent the fifth most prevalent disease or condition worldwide. The purpose of this review is to summarise the current use of dPROs and dPROMs in the field of dental traumatology. Oral Health-Related Quality of Life, pain, swelling, aesthetics, function, adverse effects, patient satisfaction, number of clinical visits, and trauma-related dental anxiety are the key dPROs following TDIs. Clinicians and researchers should consider the well-being of patients as their top priority and conduct routine evaluations of dPROs using measures that are appropriate, accurate and reflect what is important to the patient. After a TDI, dPROs can assist clinicians and patients to choose the best management option(s) for each individual patient, and potentially improve the methodology, design and relevance of clinical studies.

Keywords

Dental patient-reported outcomes, Dental patient-reported outcome measures, dental traumatology, pain, Oral Health-Related Quality of Life.

1. AIM OF THE REVIEW

Dental practitioners who care for patients who have suffered traumatic dental injuries (TDIs) should routinely assess the sequelae of treatment using not only clinician-reported outcomes but also patient-reported outcomes (PROs) using reliable and valid measures. In order to not only promote a better understanding of the patient's experience but also encourage further research into dPROs, the objective of this review is to provide an overview of dental patient-reported outcomes (dPROs) and dental patient-reported outcome measures (dPROMs) in the field of dental traumatology. The regular use of dPROs and dPROMs by clinicians and the data they generate will over time add value to the experience of patients following TDIs and potentially have a positive impact on their subsequent management.

2. DENTAL PATIENT-REPORTED OUTCOMES (dPROs) and DENTAL PATIENT-REPORTED OUTCOME MEASURES (dPROMs)

In dentistry, the term "dental patient-reported outcome" (dPRO) refers to any report of a patient's oral health state that comes directly from the patient themselves, without interpretation by a clinician or another person.¹ Patient reported outcome measures (PROMs) "seek to ascertain patients' views of their symptoms, their functional status, and their health-related quality of life (HRQoL)"² and it is preceded with a "d" for dental, dPROMs, when focused within the field of dentistry. In oral health research, dPROs are gaining popularity not only because they capture what is important to the patient in order to determine the most effective treatment from the patient's perspective,³ but also form an essential feature of clinical guideline development.⁴ Within dentistry, pain is often measured by various tools and methods (e.g. numerical rating scales) that

makes comparison between and across different studies difficult.⁵ On the contrary, the Oral Health Impact Profile (OHIP) is a global instrument with established validity, reliability and precision that not only assesses pain, but also functional limitations, handicaps, and psychological discomfort, as well as physical, psychological and social disabilities in relation to an oral disorder, thus enabling measurement of an individual's Oral Health-Related Quality of Life (OHRQoL).⁶ Notably, OHRQoL is the most significant dPRO and a major contributor to HRQoL.⁷

3. TRAUMATIC DENTAL INJURIES

TDIs have been ranked as the fifth most prevalent disease or condition in the world after dental caries, tension-type headaches, iron-deficiency anaemia and hearing loss.⁸ The global prevalence of TDIs has been reported to be 22.7% for the deciduous dentition and 15.2% for the permanent dentition, with over one billion people having experienced TDIs.⁹ The incidence of TDIs is variable across different age groups, being highest amongst children under 12 years of age.¹⁰ TDIs can be caused by multiple events such as road traffic accidents, collisions during recreation and/or sporting activities, fights etc., that can have broader physical and psychosocial implications than just the TDI *per se*.

A range of sociodemographic, clinical and environmental factors are associated with a greater chance of TDI occurrence.¹¹ The role of gender and age as risks factors are well known, whereas others such as ethnicity and socio-economic status (SES) require further understanding.¹² Clinical dental factors include increased overjet, non-nutritive sucking behaviours, a soft tissue lip trap, Class II skeletal relationship¹³⁻¹⁵ and inadequate lip coverage of the anterior teeth.¹⁶ Thus, TDIs are the second highest dental

public health problem that impact children, affecting their masticatory function and quality of life, as well as creating ongoing economic consequences.^{9,17,18} Disease prevention is favoured over disease management, with interceptive orthodontic treatment and use of mouthguards advocated for reducing the incidence of TDIs within children and adolescents.¹⁵ If TDIs do occur, retention of the damaged deciduous or permanent teeth, where possible, and repair, regeneration or both of the surrounding tissue is encouraged in order to preserve function and aesthetics.¹⁹ TDIs can have long-term sequelae (e.g. pulp necrosis, apical periodontitis or invasive cervical resorption) requiring further clinical intervention to promote favourable outcomes, including those relevant to the present review such as root canal treatment, surgical intervention and decoronation.²⁰⁻²²

4. dPROs IN TRAUMATIC DENTAL INJURIES

It is evident that the currently reported outcomes of TDIs are numerous, varied, and heterogeneous, which results in difficulties when comparing studies and significant bias when evaluating reports in this area. In addition, it is not clear how and when these outcomes should be recorded.¹⁹ While dPROs are under-represented in published studies on TDIs,¹⁹ this, along with cross-study heterogeneity also impedes the conduct of meaningful meta-analyses.²³ Assessing treatment outcomes after TDIs has traditionally focused on clinical outcomes that rely on clinical and radiographic examination and special tests. Conversely, PROs such as quality of life (QoL) and trauma-related dental anxiety have been undervalued and not traditionally employed. A systematic review reported that PROs were poorly represented in the published research on TDIs.¹⁹ Consequently, there is likely to be a lack of evidence connecting the management of TDIs to PROs.

In recent years, the use of PROs in clinical trials has increased, with 6168 (45.1%) of the 13,666 trials documented in the Australian-New Zealand Clinical Trials Registry including at least one PRO.²⁴ Within endodontology, up to 2010, 19% of studies on root canal treatment and retreatment, and apexification procedures used dPROs, which has subsequently increased to 30%.²⁵ In a similar manner, researchers have more recently focused on PROs assessing the impact of TDIs and their management on OHRQoL compared with clinician reported outcomes in the field of dental trauma. For example, between 2017 and 2022, six systematic reviews were published assessing the impact of TDIs and their management on OHRQoL.^{18,26-30}

5. IMPORTANCE OF dPROS IN TRAUMATIC DENTAL INJURIES

Patient involvement in outcomes research “may improve the relevance of research questions”,³¹ with consideration of dPROs in conjunction with clinical outcomes being essential to ensuring “greater understanding of patient priorities, preferences, values, expectations, and experiences”.⁵ Otherwise, if a clinician's viewpoint is used to interpret a patient's perspective of treatment, valuable information from the patient's point of view on the effectiveness of treatment risks being ignored, leading to patient dissatisfaction due to unfulfilled expectations and a perception of being ignored.⁵

The advantages of using dPROs following treatment of TDIs includes:³²⁻³⁵

- Providing a more comprehensive view of treatment benefits and costs to aid clinicians and patients to select the most suitable treatment. For example, an uncomplicated crown fracture may be restored in various ways including re-

attachment of the fractured segment if the patient has kept it, a direct resin-based composite restoration, an indirect restoration (with various materials). Each has its own clinical and aesthetic advantages and disadvantages, and respective time requirements. There are also vast cost differences which must be discussed with the patient.

- Providing information to enhance clinical decision-making for the management of TDIs. For example, an avulsed tooth with an extended extra-oral dry time in a young patient will over time predictably develop ankylosis and external replacement resorption.³⁶ Management options include no treatment, decoronation, extraction or auto-transplantation. Each carries important consequences including aesthetics, loss of alveolar bone and soft tissue, the need for an interim prosthesis, and the long-term replacement of the tooth. In the case of long term extra-alveolar time, replantation is the recommended option as it will keep future treatment options open while restoring aesthetics and function in the interim, as well as maintaining alveolar bone in the edentulous space that may be required for future restorative options.³⁷
- PROs can contribute to the improvement of future clinical trial methodologies and be utilised as effective research outcomes for the development and evaluation of new drugs and therapies because they enable the estimation of treatment advantages and disadvantages separately from standard clinical measures of effectiveness.
- Improved translation of research studies to patients and lay stakeholders, for example understanding the management of TDIs in their own words.

6. dPROs USED IN TRAUMATIC DENTAL INJURIES

OHRQoL

The effect of TDIs on OHRQoL is complex and should take into consideration several clinical and non-clinical variables. OHRQoL is a global outcome measure encompassing functional limitations, oral symptoms, emotional and social impacts.⁵ TDIs may affect the OHRQoL of participants and also that of their families and/or carers.³⁸⁻⁴²

TDIs have a range of clinical presentations from an enamel fracture to tooth avulsion, with some studies reporting an association between the severity of the injury and its impact on the OHRQoL,^{43,44} but not others.⁴⁵ Injuries can involve variable numbers of teeth during different dentition and root maturation stages. These injuries can be managed with alternative protocols or can even remain untreated. The impact of TDIs on ORHQoL may be affected by other oral conditions such as dental caries and malocclusion.^{46,47} Injuries to multiple teeth are considered more common in traumatic events such as accidental falls and trips, acts of violence, sports, and road traffic accidents.¹²

TDIs can occur in conjunction with other oral and maxillofacial hard and soft tissue injuries as well as broader bodily injuries that can also affect the QoL of the patient. Bodily injuries may cause other non-dental disabilities and their management including long-term hospitalisation, which negatively influences the QoL of the subjects,⁴⁸ and requires interdisciplinary management. Several studies from Brazil have assessed TDIs in the primary dentition.^{49,50} When all types of TDIs were pooled, injuries in the primary dentition did not affect the OHRQoL, a finding that is likely to be due to a high incidence of enamel fractures in young children,^{44,49-52} which have limited impact

on OHRQoL.⁵¹ Conversely, severe TDIs were associated with poorer outcomes in a study that focussed on the primary dentition.⁵³ Amongst the different clinical presentations, tooth avulsion was associated with a negative impact on OHRQoL for the children and their families, although interestingly tooth discolouration had a negative effect solely on the parents.⁵² The latter is in agreement with a study where TDIs were associated with poorer outcomes for children, although only more severe presentations (i.e., avulsion and luxation) were reported by parents to have negative effects.⁴⁴ Mild dental trauma to the primary dentition appears to have a limited effect on OHRQoL.

The assessment of the association between TDIs and OHRQoL in the mixed and permanent dentitions has also attracted attention.⁵⁴⁻⁵⁹ When assessing the type of injury, crown fractures were associated with poorer OHRQoL amongst young children and adolescents in Brazil,^{56,59} and India.⁵⁸ Two Brazilian studies that studied trauma involving periodontal ligament damage (i.e. luxations and avulsion) reported negative outcomes in 12-year-olds⁵⁵ and 8-10 years old children.⁵⁷ Finally, tooth avulsion amongst children under-18 had an adverse effect on OHRQoL in an Italian population.⁵⁴ Thus, TDIs in the mixed and permanent dentitions appear to have a negative effect on OHRQoL.

The management of TDIs (e.g., treatment versus non-treatment and likely treatment modalities) may also impact OHRQoL. The absence of treatment may negatively impact OHRQoL due to potential functional and/or aesthetic limitations and depends on the severity of the injury. Untreated injuries in the permanent dentition commonly have a negative impact, as observed amongst 12-14 years old adolescents in Canada,⁶⁰ 12 years old Jordanians,⁴⁷ 12-14 years old Brazilians,⁶¹ and Albanians aged

16-19 years.⁶² Severe untreated injuries are also associated with a negative impact on the OHRQoL scores of Brazilian adolescents.⁶³ For the management of tooth avulsion, replantation, if possible, is the best treatment option.^{54,59} Conversely, a study in Brazil showed that the absence of treatment for crown fractures extending beyond enamel had no negative effect on OHRQoL.⁵⁹ In addition, a separate study from Brazil where children aged 8-14 years were treated for crown fractures reported enhanced OHRQoL scores for children, but no effect for their families.⁶⁴ Furthermore, even following treatment, adolescents are at risk of having poorer QoL when compared with their uninjured peers.⁶⁵ No studies are available comparing different treatment modalities and their effect on OHRQoL.

The role of dental caries and malocclusion as confounding factors when assessing the impact of TDI on OHRQoL may vary depending on clinical and participant-related factors. Studies from Brazil, Colombia, and Peru have reported a significant impact in OHRQoL amongst children aged 1 to 14 years for experience of dental caries and TDI when compared with occlusal discrepancies.^{41,44,66-69} Nonetheless, the additional impact of dentofacial anomalies plus TDIs on OHRQoL was highlighted in a study from India.⁷⁰ A study into 8-10 years old from Belo Horizonte, Brazil reported that dental caries had a higher impact on OHRQoL than TDIs and malocclusion.⁷¹ Similarly, another Brazilian study of 5-6-year-old children showed that severe caries, but not TDIs, was associated with poorer perception for children and carers with regards to OHRQoL.⁴⁹ These findings are in agreement with a comparable study also from Brazil.⁷² Therefore, TDIs should be considered alongside, not independent to, other oral presentations and diseases, when evaluating OHRQoL.

The role of SES on the association between TDIs and OHRQoL has been assessed with conflicting results, making definitive conclusions difficult. Amongst the studies reporting the negative impact of SES on OHRQoL, some criteria have been highlighted, such as: non-nuclear families (single parents or living with others),⁴¹ lower income,^{40,46,52,67,72} and low-level parental schooling/education.^{39,57} Conversely, other studies have reported no significant association between OHRQoL and SES.^{44,55,66,73}

The limitations of studies assessing the effect of TDIs on OHRQoL should be emphasised. Population studies rarely include radiographic examination, thus some types of injuries, such as root fractures, may elude detection during dental examinations in the absence of subsequent special tests. Grouping of TDIs of variable severity may mask negative effects on OHRQoL of more severe injuries, as less severe presentations may have a limited impact on this outcome measure and possibly be more detectable with clinical examination. Study designs in this area are often retrospective and/or cross-sectional in design and this can have a negative effect on the level of evidence on the topic and the ability to rigorously compare injuries. Experimental trials on the topic are not possible for ethical reasons, as participants cannot be intentionally subjected to TDIs. However, the role of various clinical and non-clinical factors, plus the management of the injury and the subsequent impact on OHRQoL is amenable to further assessment and does require further high-quality clinical studies.

Symptoms

Symptoms following treatment for TDIs are usually assessed on an individual basis as pain/swelling or indirectly as discomfort. Pain in endodontics has been defined as a “multifactorial noxious experience that involves a sensory response that can be

modified by cognitive, emotional and motivational influences related to past experience".⁷⁴ Pain can be spontaneous or it can be elicited by touching the tooth and/or the related soft tissues, with the latter often defined as 'tenderness'.⁵ In dentistry, pain is an important dPRO as well as having an influence on the Oral Health Impact Profile (OHIP) tool⁶ and on one of the four dimensions of the OHRQoL (orofacial pain).³ Pain and/or tenderness occur following most of the treatments provided after a TDI such as root canal treatment,⁴ dental implants,⁷⁵ intraoral bone grafting,⁷⁶ amongst others. The outcome assessment tools for postoperative pain following endodontic treatment generally involve a 11-point rating scale ranging from "no pain" to "worst possible pain"⁷⁶ or a numerical rating scale (NRS) ranging from 0 to 10.⁷⁵

The dental literature defines discomfort as a subjective experience resulting from oral or facial symptoms (pain, swelling, bleeding, and infection),⁷⁷ functional impairment (chewing, speech, and oral hygiene maintenance),⁷⁸ or general conditions (palpation, vomiting, and dizziness)⁷⁹ after dental treatment. Patients may become anxious about their treatment if they experience severe or ongoing discomfort.

Aesthetics

Aesthetics is an important characteristic of all dentitions. For example, it has been reported that fracture, displacement, and/or early loss of one or more primary maxillary incisor teeth will inevitably lead to impaired aesthetics in pre-school children.⁸⁰ When comparing intrusion injuries to other sub-types of luxation injuries, intrusion injuries result in higher mineralisation abnormalities in successional teeth. Mineralisation disturbances are typically linked to an aesthetic concern, which is crucial for the permanent incisors.⁸¹ Likewise, discolouration of the crown was commonly

encountered in the crowns of teeth following regenerative endodontic procedures.⁸² A systematic review reported that the most common unfavourable effects of trauma to permanent teeth are enamel hypoplasia and white or yellow brown discolouration.⁸¹ After trauma, pulp hyperaemia and pulpal haemorrhage release haemoglobin and erythrocyte products, resulting in crown discolouration. Due to the development of an apical tissue coagulum and subsequent odontoblastic stimulation, subluxated and intruded teeth exhibit a phenomenon known as amorphous calcification, which can also affect the subsequent colour of the crown.⁸³ The initial or emergency phase of treatment for TDIs includes care to relieve symptoms and to restore aesthetics and function.⁸⁴ Aesthetic impairment can be indirectly evaluated in pre-school children through questionnaires by assessing the parental demand for replantation of avulsed teeth.⁸⁵ Czochrowska et al.⁸⁶ evaluated the aesthetic outcome of autotransplanted premolars replacing maxillary incisors in adult patients by obtaining their opinions through the use of a questionnaire. The questions targeted the colour, length, width and position of the transplanted tooth and the patients' responses were categorised as satisfied, acceptable, and dissatisfied. Overall, 73% of the patients were satisfied with the aesthetic outcome of transplanted premolars.

Function

Speech is a function that may be affected as a consequence of sudden and premature loss of deciduous teeth following a TDI.⁸⁰ Gable et al.⁸⁷ conducted a non-randomised trial with 26 children following the natural exfoliation of their primary maxillary incisors and a further 26 children who had the same teeth extracted prematurely. Following the eruption of their permanent incisors, all subjects underwent testing for speech impairment. Intriguingly, there was no significant difference between the groups

for speech articulation impairments. Age-related reductions in articulation errors were seen in both groups, indicating a maturation effect. In addition to TDIs, post-trauma treatment could also have an impact on speech such as following autotransplantation⁸⁶ and implant placement.⁸⁸

Adverse effects

Adverse effects following TDIs often include pulpitis or pulp necrosis with infection and apical periodontitis which require some form of endodontic intervention.⁴ Discolouration due to the leaching of medicaments, root canal cements or endodontic materials into the dentine surrounding the pulp chamber is a known adverse effect following root canal treatment. Discolouration may also occur because of pulp canal calcification or pulp necrosis, as a response to trauma, following autotransplantation,⁸⁹ replantation⁹⁰ and luxation injuries.⁸¹ Pulp canal calcification, pulp necrosis and tooth loss are the most prevalent outcomes of luxation injuries.⁸¹ Discolouration can be measured subjectively by both the dentist and the patient or objectively using a spectrophotometer.⁹¹

Patient satisfaction

Patient satisfaction may include feedback specifically about the treatment delivered, which adds value during assessment of a patients' perception of a successful or unsuccessful treatment, usually measured through a patient questionnaire.^{86,92} In cases of auto-transplantation, aesthetic outcomes have been measured as a part of the patient's satisfaction by recording their responses in a 4-point Likert scale within a questionnaire.⁸⁶ Satisfaction can also be recorded qualitatively in the form of interview

with the patient that includes self-assessment as well as verbal evaluation of the procedure *per se* and long-term morbidity (discomfort, loss of sensation, and pain).⁷⁶ In this context, an 11-point rating scale has been used for self-assessment and a dichotomous scale for verbal evaluation. Patient satisfaction for implant-supported single crowns rated on a 4-point scale has been reported to be 93-98%.⁹³ In the context of satisfaction for paediatric patients, they should receive prompt, high-quality dental care that addresses symptoms and signs, as well as restores both tooth function and aesthetics. The prevalence of young people using social media, which potentially fuels bullying and taunting connected to appearance, is a growing concern.⁹⁴ Hakeberg et al.⁹⁵ assessed childrens' satisfaction towards the dentist using the Dental Visit Satisfaction Scale – Swedish version (DVSS-SV). In a study conducted to assess the psychosocial impact of TDIs on the self-esteem of adolescents using Rosenberg's self-esteem scale, children with TDIs reported poor self-esteem, which may lead to loss of an individual's self-confidence.⁹⁶

Number of clinical visits

The number of treatment and review visits is a secondary outcome associated with the nature and course of treatment offered for TDIs of both deciduous and permanent teeth.⁹⁷ The number of visits is usually measured by either verifying the patient's dental records or through retrospective multiple interviews of patients and their companions (relatives, neighbours, teachers and others) by telephone.^{98,99} The main factors that can influence the number of dental visits are clinically-related: complicated hard tissue injuries, complicated or uncomplicated periodontal injuries, the diagnosis and treatment of pulp necrosis, and treatment approaches of the clinician. Patient-related factors that can also determine the number of appointments include proximity of the

patient's residence to the clinic,^{5,97} availability of appointments,⁵ as well as patient preferences for the number of visits as there may be a desire to limit absence from work, particularly when this is related to loss of earnings.⁵ Thus, consideration of both clinical and patient-related factors is essential when deciding on the number of visits required in order to ensure that patients may experience high levels of satisfaction.⁵ Furthermore, the time involved for dental visits can be categorised as direct (involving professional healthcare providers) and indirect time (involving companions, e.g. relatives, neighbours, teachers and others).^{98,100}

Trauma related dental anxiety

TDIs involving children are likely to increase their levels of dental anxiety.^{23,101} Dental fear is a common emotional response to a particular external threat in a dental scenario. When a child experiences dental anxiety, they sense a generalised feeling of apprehension that is linked to aberrant circumstances.¹⁰² Dental fear is also characterised as a special type of anxiety that is accompanied with a propensity for having a negative dental experience.¹⁰³ Dental anxiety in children can be effectively measured by the clinician using various age-dependent tools such as: 5-8 years – Facial imaging scale;¹⁰⁴ 8-12 years – Faces version of the Modified Dental Anxiety Scale (MCDAS_f);¹⁰⁵ 12 years and above – Kleinknecht's Dental Fear Survey (DFS) and/or Corah's Dental Anxiety Scale (DAS).¹⁰⁶ Skaare et al.¹⁰⁷ distributed a questionnaire to care givers to assess the possible distress related to trauma of primary teeth during the follow-up examination of the permanent successors. Generally, girls, children who were more fearful of medical procedures,¹⁰⁸ and children with previous TDI were highly anxious.¹⁰⁸ Additionally, males and children who had experienced oral trauma had a generally more aggressive nature. There is a strong link between children's aggressive

behavior and dental anxiety and the likelihood of displaying violent behaviour increased marginally with the increase in anxiety level.¹⁰⁸

7. CORE OUTCOME SET (COS) IN TRAUMATIC DENTAL INJURIES

In an attempt to address heterogeneity and encourage the use of dPROs, the International Association of Dental Traumatology established a group of international experts to develop core outcome sets (COS) for TDIs in children and adults.²³ A COS is a collection of agreed-upon standard outcomes that must be included, measured, and reported at a minimum level in all clinical trials and outcome studies undertaken in a particular field. The protocol for developing the COS in dental traumatology followed an established methodological framework with prospective registration in the Core Outcome Measures in Effectiveness Trials (COMET). The process involved the identification of outcomes using literature searches and a web-based questionnaire that involved dentists globally, followed by an online Delphi process that determined the most significant outcomes for inclusion. Following the online Delphi process, conference calls were held to finalize the COS for severe TDIs. Finally, 23 outcomes were considered, 13 of which were "generic" and 10 of which were "injury-specific".²³ These COS can be used in future clinical and research studies, reviews, and the development of future clinical practice guidelines for TDIs with the hope that this will improve future research and allow meta-analyses to be performed.

8. FUTURE DIRECTIONS FOR RESEARCH

- Population level studies, ideally collecting outcomes consistent with COS from multiple centres, will be an ideal source of data when assessing dPROs in relation to TDIs.

- Future randomised clinical trials (when possible for management modalities) and observational studies should consider incorporating dPROs in addition to collecting data on clinical outcomes. Patients, patient representatives, or both, should be involved in deciding the appropriate patient-defined outcomes that are important to them.¹⁰⁹ This will eventually result in patient-centred, evidence-based care, enhancing the value of treatment following TDIs.
- Most PROMs used in dental traumatology are generic. Consequently, it is necessary to develop a set of validated dPROMs that are specific to the field of dental traumatology and then to map these to the four-dimensional framework (Oral function, Orofacial appearance, Orofacial pain, and Psychosocial impact). These steps will standardize the evaluation of various TDI treatment outcomes, which will facilitate synthesis of multiple research reports through systematic reviews and meta-analyses.
- Future research should identify the dPROMs applicable to TDIs, which will ultimately result in the identification of the most suitable outcome measures for patients undergoing treatment of TDIs moving forward.
- Clinicians and researchers have the opportunity to adopt electronic PROMs (ePROMs) for the routine monitoring of patients after a TDI and its management. ePROMs offer patients an "electronic" means of data submission, through any internet-connected device (e.g., PC, smartphone or tablet), providing clinicians and researchers with a flexible platform for viewing PROM data.^{110,111} Patients suffering from cancer and chronic kidney disease, for instance, have reported that the ePROM platform is a beneficial tool that improves their outcomes.^{112,113}

9. CONCLUSION

The important dPROs following TDIs are OHRQoL, pain, swelling, aesthetics, function, adverse effects, patient satisfaction, and trauma-related dental anxiety. The continued development of dPROs and an understanding of what matters to patients, clinicians and academics can improve therapies and research, which will ultimately benefit the management of DTIs, patient well-being, and wider society. To accurately represent the effects of TDIs and their treatment on patients, the proper use of dPROs is essential in dental traumatology. To more accurately capture the viewpoints of patients undergoing treatment following TDIs, a novel and unique assessment tool should be developed.

CONFLICT OF INTEREST

The authors declare there are no competing interests for the current manuscript. No funding was received for the current work.

ETHICAL STATEMENT

No ethics approval was required for this manuscript.

REFERENCES

1. John MT. Health outcomes reported by dental patients. *J Evid Based Dent Pract.* 2018;18:332–5.
2. Black N. Patient reported outcome measures could help transform healthcare. *BMJ.* 2013;346:f167.
3. John M, Omara M, Su N, List T, Sekulic S, Häggman-Henrikson B, et al. Recommendations for use and scoring of oral health impact profile versions. *J Evid Based Dent Pract.* 2022;22:101619.
4. Duncan HF, Nagendrababu V, El-Karim I, Dummer PMH. Outcome measures to

- assess the effectiveness of endodontic treatment for pulpitis and apical periodontitis for use in the development of European Society of Endodontology S3-level clinical practice guidelines: a consensus-based development. *Int Endod J.* 2021;54:2184–94.
5. Dođramacı EJ, Rossi-Fedele G. Patient-related outcomes and oral health-related quality of life in endodontics. *Int Endod J.* 2022;00:1–9.
 6. Slade GD, Spencer AJ. Development and evaluation of the oral health impact profile. *Community Dent Health.* 1994;11:3–11.
 7. John MT. Foundations of oral health-related quality of life. *J Oral Rehabil.* 2021;48:355–9.
 8. Petti S, Andreasen JO, Glendor U, Andersson L. The fifth most prevalent disease is being neglected by public health organisations. *Lancet Glob Heal.* 2018;6:e1070–1.
 9. Petti S, Glendor U, Andersson L. World traumatic dental injury prevalence and incidence, a meta-analysis-one billion living people have had traumatic dental injuries. *Dent Traumatol.* 2018;34:71–86.
 10. Andersson L. Epidemiology of traumatic dental injuries. *J Endod.* 2013;39:S2-5.
 11. Magno MB, Nadelman P, Leite KL de F, Ferreira DM, Pithon MM, Maia LC. Associations and risk factors for dental trauma: a systematic review of systematic reviews. *Community Dent Oral Epidemiol.* 2020;48:447–63.
 12. Glendor U. Epidemiology of traumatic dental injuries - a 12 year review of the literature. *Dent Traumatol.* 2008;24:603–11.
 13. Dođramacı EJ, Rossi-Fedele G. Establishing the association between nonnutritive sucking behavior and malocclusions. *J Am Dent Assoc.* 2016;147:926-934.e6.
 14. Dođramacı EJ, Rossi-Fedele G, Dreyer CW. Malocclusions in young children: does

- breast-feeding really reduce the risk? a systematic review and meta-analysis. *J Am Dent Assoc.* 2017;148:566-574.e6.
15. Arraj GP, Rossi-Fedele G, Dođramacı EJ. The association of overjet size and traumatic dental injuries—a systematic review and meta-analysis. *Dent Traumatol.* 2019;35:217–32.
 16. Dieguez-Perez M, de Nova-Garcia M, Mourelle-Martinez M, Bartolome-Villar B. Oral health in children with physical (cerebral palsy) and intellectual (down syndrome) disabilities: systematic review I. *J Clin Exp Dent.* 2016;0–0.
 17. Feldens CA, Kramer PF, Feldens EG, Pacheco LM, Vítolo MR. Socioeconomic, behavioral, and anthropometric risk factors for traumatic dental injuries in childhood: a cohort study. *Int J Paediatr Dent.* 2014;24:234–43.
 18. Zaror C, Martínez-Zapata MJ, Abarca J, Díaz J, Pardo Y, Pont À, et al. Impact of traumatic dental injuries on quality of life in preschoolers and schoolchildren: a systematic review and meta-analysis. *Community Dent Oral Epidemiol.* 2018;46:88–101.
 19. Sharif MO, Tejani-Sharif A, Kenny K, Day PF. A systematic review of outcome measures used in clinical trials of treatment interventions following traumatic dental injuries. *Dent Traumatol.* 2015;31:422–8.
 20. Malmgren B, Tsilingaridis G, Malmgren O. Long-term follow up of 103 ankylosed permanent incisors surgically treated with decoronation - a retrospective cohort study. *Dent Traumatol.* 2015;31:184–9.
 21. Lin S, Schwarz-Arad D, Ashkenazi M. Alveolar bone width preservation after decoronation of ankylosed anterior incisors. *J Endod.* 2013;39:1542–4.
 22. Levin L, Day PF, Hicks L, O’Connell A, Fouad AF, Bourguignon C, et al. International association of dental traumatology guidelines for the management

- of traumatic dental injuries: general introduction. *Dent Traumatol.* 2020;36:309–13.
23. Kenny KP, Day PF, Sharif MO, Parashos P, Lauridsen E, Feldens CA, et al. What are the important outcomes in traumatic dental injuries? An international approach to the development of a core outcome set. *Dent Traumatol.* 2018;34:4–11.
 24. Mercieca-Bebber R, Williams D, Tait MA, Roydhouse J, Busija L, Sundaram CS, et al. Trials with patient-reported outcomes registered on the Australian New Zealand Clinical Trials Registry (ANZCTR). *Qual Life Res.* 2018;27:2581–91.
 25. Azarpazhooh A, Sgro A, Cardoso E, Elbarbary M, Laghapour Lighvan N, Badewy R, et al. A scoping review of 4 decades of outcomes in nonsurgical root canal treatment, nonsurgical retreatment, and apexification studies—Part 2: outcome measures. *J Endod.* 2022;48:29–39.
 26. Borges TS, Vargas-Ferreira F, Kramer PF, Feldens CA. Impact of traumatic dental injuries on oral health-related quality of life of preschool children: a systematic review and meta-analysis. Milgrom PM, editor. *PLoS One.* 2017;12:e0172235.
 27. Lopez D, Waidyatillake N, Zaror C, Mariño R. Impact of uncomplicated traumatic dental injuries on the quality of life of children and adolescents: a systematic review and meta-analysis. *BMC Oral Health.* 2019;19:224.
 28. Antunes LAA, Lemos HM, Milani AJ, Guimarães LS, Küchler EC, Antunes LS. Does traumatic dental injury impact oral health-related to quality of life of children and adolescents? systematic review and meta-analysis. *Int J Dent Hyg.* 2020;18:142–62.
 29. Milani AJ, Castilho T, Assaf AV, Antunes LS, Antunes LAA. Impact of traumatic dental injury treatment on the oral health-related quality of life of children, adolescents, and their family: systematic review and meta-analysis. *Dent*

- Traumatol. 2021;37:735–48.
30. Das P, Mishra L, Jena D, Govind S, Panda S, Lapinska B. Oral health-related quality of life in children and adolescents with a traumatic injury of permanent teeth and the impact on their families: a systematic review. *Int J Environ Res Public Health*. 2022;19:3087.
 31. Concannon TW, Fuster M, Saunders T, Patel K, Wong JB, Leslie LK, et al. A systematic review of stakeholder engagement in comparative effectiveness and patient-centered outcomes research. *J Gen Intern Med*. 2014;29:1692–701.
 32. Gatten DL, Riedy CA, Hong SK, Johnson JD, Cohenca N. Quality of life of endodontically treated versus implant treated patients: a university-based qualitative research study. *J Endod*. 2011;37:903–9.
 33. Mercieca-Bebber R, King MT, Calvert MJ, Stockler MR, Friedlander M. The importance of patient-reported outcomes in clinical trials and strategies for future optimization. *Patient Relat Outcome Meas*. 2018;Volume 9:353–67.
 34. Liang M, Lian Q, Kotsakis GA, Michalowicz BS, John MT, Chu H. Bayesian network meta-analysis of multiple outcomes in dental research. *J Evid Based Dent Pract*. 2020;20:101403.
 35. Coleman RL, Beck JT, Baranda JC, Jacobs I, Smoyer KE, Lee LJ, et al. The use of patient-reported outcome measures in phase I oncology clinical trials. *Oncology*. 2021;99:444–53.
 36. Dođramaci EJ, Rossi-Fedele G, Jonest AG. Multi-disciplinary management of a patient with a post-traumatized incisor presenting concurrent replacement and inflammatory resorption: a case report. *Aust Orthod J*. 2015;31:216–25.
 37. Fouad AF, Abbott P V, Tsilingaridis G, Cohenca N, Lauridsen E, Bourguignon C, et al. International Association of Dental Traumatology guidelines for the

- management of traumatic dental injuries: 2. Avulsion of permanent teeth. *Dent Traumatol.* 2020;36:331–42.
38. Ramos-Jorge ML, Ramos-Jorge J, Mota-Veloso I, Oliva KJ, Zarzar PM, Marques LS. Parents' recognition of dental trauma in their children. *Dent Traumatol.* 2013;29:266–71.
 39. Barbosa Neves ÉT, Perazzo MF, Gomes MC, Martins CC, Paiva SM, Granville-Garcia AF. Perception of parents and self-reports of children regarding the impact of traumatic dental injury on quality of life. *Dent Traumatol.* 2017;33:444–50.
 40. Sardenberg F, Cavalcante-Leão BL, Todero SRB, Ferreira FM, Rebellato NLB, Fraiz FC. A population-based study on the impact of orofacial dysfunction on oral health-related quality of life among Brazilian schoolchildren. *Acta Odontol Scand.* 2017;75:173–8.
 41. Díaz S, Mondol M, Peñate A, Puerta G, Boneckér M, Martins Paiva S, et al. Parental perceptions of impact of oral disorders on Colombian preschoolers' oral health-related quality of life. *Acta Odontol Latinoam.* 2018;31:23–31.
 42. Ode W, Lopez V, Wong ML, Schou L, Yu VSH. Understanding patients' and dentists' perspectives in dental trauma management: A mixed-methods study. *Dent Traumatol.* 2018;34:320–8.
 43. Berger TD, Kenny DJ, Casas MJ, Barrett EJ, Lawrence HP. Effects of severe dentoalveolar trauma on the quality-of-life of children and parents. *Dent Traumatol.* 2009;25:462–9.
 44. Gomes MC, Pinto-Sarmiento TC de A, Costa EMM de B, Martins CC, Granville-Garcia AF, Paiva SM. Impact of oral health conditions on the quality of life of preschool children and their families: a cross-sectional study. *Health Qual Life Outcomes.* 2014;12:55.

45. Porritt JM, Rodd HD, Ruth Baker S. Quality of life impacts following childhood dento-alveolar trauma. *Dent Traumatol.* 2011;27:2–9.
46. Locker D. Disparities in oral health-related quality of life in a population of Canadian children. *Community Dent Oral Epidemiol.* 2007;35:348–56.
47. Rajab LD, Abu Al Huda D. Impact of treated and untreated traumatic dental injuries on oral health-related quality of life among 12-year-old schoolchildren in Amman. *Dent Traumatol.* 2019;35:153–62.
48. Hirobe Y, Koshinuma S, Nakamura M, Baba M, Yamamoto G, Hitosugi M. Factors influencing the long-term hospitalization of bicyclists and motorcyclists with oral and maxillofacial injuries. *Dent Traumatol.* 2021;37:234–9.
49. Abanto J, Tsakos G, Paiva SM, Carvalho TS, Raggio DP, Bönecker M. Impact of dental caries and trauma on quality of life among 5- to 6-year-old children: perceptions of parents and children. *Community Dent Oral Epidemiol.* 2014;42:385–94.
50. Vieira-Andrade RG, Siqueira MBL, Gomes GB, D'Avila S, Pordeus IA, Paiva SM, et al. Impact of traumatic dental injury on the quality of life of young children: a case–control study. *Int Dent J.* 2015;65:261–8.
51. Feldens CA, Day P, Borges TS, Feldens EG, Kramer PF. Enamel fracture in the primary dentition has no impact on children's quality of life: implications for clinicians and researchers. *Dent Traumatol.* 2016;32:103–9.
52. Viegas CM, Paiva SM, Carvalho AC, Scarpelli AC, Ferreira FM, Pordeus IA. Influence of traumatic dental injury on quality of life of Brazilian preschool children and their families. *Dent Traumatol.* 2014;30:338–47.
53. Abanto J, Tello G, Bonini GC, Oliveira LB, Murakami C, Bönecker M. Impact of traumatic dental injuries and malocclusions on quality of life of preschool

- children: a population-based study. *Int J Paediatr Dent*. 2015;25:18–28.
54. Giannetti L, Murri A, Vecci F, Gatto R. Dental avulsion: therapeutic protocols and oral health-related quality of life. *Eur J Paediatr Dent*. 2007;8:69–75.
 55. Silva-Oliveira F, Goursand D, Ferreira RC, Paiva PCP, Paiva HN, Ferreira EF, et al. Traumatic dental injuries in Brazilian children and oral health-related quality of life. *Dent Traumatol*. 2018;34:28–35.
 56. Damé-Teixeira N, Alves LS, Ardenghi TM, Susin C, Maltz M. Traumatic dental injury with treatment needs negatively affects the quality of life of Brazilian schoolchildren. *Int J Paediatr Dent*. 2013;23:266–73.
 57. Freire-Maia FB, Auad SM, Abreu MHNG de, Sardenberg F, Martins MT, Paiva SM, et al. Oral health-related quality of life and traumatic dental injuries in young permanent incisors in Brazilian schoolchildren: a multilevel approach. *PLoS One*. 2015;10:e0135369.
 58. Golai S, Nimbeni B, Patil SD, Baali P, Kumar H. Impact of untreated traumatic injuries to anterior teeth on the oral health related quality of life as assessed by video based smiling patterns in children. *J Clin Diagn Res*. 2015;9:ZC16-9.
 59. da Silva RLC, Dias Ribeiro AP, Almeida JCF, Sousa SJL, Garcia FCP. Impact of dental treatment and the severity of traumatic dental injuries on the quality of life of Brazilian schoolchildren. *Dent Traumatol*. 2021;37:562–7.
 60. Fakhruddin KS, Lawrence HP, Kenny DJ, Locker D. Impact of treated and untreated dental injuries on the quality of life of Ontario school children. *Dent Traumatol*. 2008;24:309–13.
 61. Ilma de Souza Cortes M, Marcenes W, Sheiham A. Impact of traumatic injuries to the permanent teeth on the oral health-related quality of life in 12-14-year-old children. *Community Dent Oral Epidemiol*. 2002;30:193–8.

62. Thelen DS, Trovik TA, Bårdsen A. Impact of traumatic dental injuries with unmet treatment need on daily life among Albanian adolescents: a case-control study. *Dent Traumatol.* 2011;27:88–94.
63. Bendo CB, Paiva SM, Varni JW, Vale MP. Oral health-related quality of life and traumatic dental injuries in Brazilian adolescents. *Community Dent Oral Epidemiol.* 2014;42:216–23.
64. Magno MB, Jural LA, Nogueira A do V, Lenzi MM, Pithon MM, Maia LC. Impact of crown fracture treatment on oral health-related quality of life of children, adolescents, and their families: a prospective clinical study. *Int J Paediatr Dent.* 2019;29:86–93.
65. Ramos-Jorge ML, Bosco VL, Peres MA, Nunes ACGP. The impact of treatment of dental trauma on the quality of life of adolescents? a case-control study in southern Brazil. *Dent Traumatol.* 2007;23:114–9.
66. Kramer PF, Feldens CA, Helena Ferreira S, Bervian J, Rodrigues PH, Peres MA. Exploring the impact of oral diseases and disorders on quality of life of preschool children. *Community Dent Oral Epidemiol.* 2013;41:327–35.
67. Schuch HS, dos Santos Costa F, Torriani DD, Demarco FF, Goettems ML. Oral health-related quality of life of schoolchildren: impact of clinical and psychosocial variables. *Int J Paediatr Dent.* 2015;25:358–65.
68. Firmino RT, Gomes MC, Clementino MA, Martins CC, Paiva SM, Granville-Garcia AF. Impact of oral health problems on the quality of life of preschool children: a case-control study. *Int J Paediatr Dent.* 2016;26:242–9.
69. Pulache J, Abanto J, Oliveira LB, Bönecker M, Porras JC. Exploring the association between oral health problems and oral health-related quality of life in Peruvian 11- to 14-year-old children. *Int J Paediatr Dent.* 2016;26:81–90.

70. Basavaraj P, Sunil MK, Nagarajappa R, Ashish S, Ramesh G. Correlation between oral health and child-OIDP Index in 12- and 15-year-old children from modinagar, India. *Asia Pacific J Public Heal.* 2014;26:390–400.
71. Martins MT, Sardenberg F, Bendo CB, Vale MP, Paiva SM, Pordeus IA. Dental caries are more likely to impact on children's quality of life than malocclusion or traumatic dental injuries. *Eur J Paediatr Dent.* 2018;19:194–8.
72. Scarpelli AC, Paiva SM, Viegas CM, Carvalho AC, Ferreira FM, Pordeus IA. Oral health-related quality of life among Brazilian preschool children. *Community Dent Oral Epidemiol.* 2013;41:336–44.
73. Viegas CM, Scarpelli AC, Carvalho AC, Ferreira F de M, Pordeus IA, Paiva SM. Impact of traumatic dental injury on quality of life among Brazilian preschool children and their families. *Pediatr Dent.* 2012;34:300–6.
74. American Association of Endodontics. *Glossary of Endodontic Terms.* 10th ed. Chicago: American Association of Endodontics; 2020. 36 p.
75. Wu X, Ye M, Sun J, Yan Q, Shi B, Xia H. Patient-reported outcome measures following surgeries in implant dentistry and associated factors: a cross-sectional study. *BMJ Open.* 2022;12:e059730.
76. Andersson L. Patient self-evaluation of intra-oral bone grafting treatment to the maxillary frontal region. *Dent Traumatol.* 2008;24:164–9.
77. Korsch M, Baum A, Bartols A. Postoperative discomfort after implant placement according to the all-on-4 concept with or without zygoma implants: a prospective clinical study. *Clin Oral Implants Res.* 2020;31:133–43.
78. Ribeiro AB, Della Vecchia MP, Cunha TR, Sorgini DB, dos Reis AC, Muglia VA, et al. Short-term post-operative pain and discomfort following insertion of mini-implants for retaining mandibular overdentures: a randomized controlled trial. *J*

- Oral Rehabil. 2015;42:605–14.
79. Getter L, Levin MP, Ayer W. Comparative effectiveness of some commonly prescribed drugs for the relief of post-surgical discomfort. *Anesth Prog.* 1973;20:137–40.
 80. Holan G, Needleman HL. Premature loss of primary anterior teeth due to trauma - potential short- and long-term sequelae. *Dent Traumatol.* 2014;30:100–6.
 81. Goswami M, Rahman B, Singh S. Outcomes of luxation injuries to primary teeth-a systematic review. *J Oral Biol Craniofacial Res.* 2020;10:227–32.
 82. Kahler B, Mistry S, Moule A, Ringsmuth AK, Case P, Thomson A, et al. Revascularization outcomes: a prospective analysis of 16 consecutive cases. *J Endod.* 2014;40:333–8.
 83. Qassem A, Martins N da M, da Costa VPP, Torriani DD, Pappen FG. Long-term clinical and radiographic follow up of subluxated and intruded maxillary primary anterior teeth. *Dent Traumatol.* 2015;31:57–61.
 84. Belmonte FM, Macedo CR, Day PF, Saconato H, Fernandes Moça Trevisani V. Interventions for treating traumatised permanent front teeth: luxated (dislodged) teeth. *Cochrane Database Syst Rev.* 2013;2014.
 85. Raphael SL, Gregory PJ. Parental awareness of the emergency management of avulsed teeth in children. *Aust Dent J.* 1990;35:130–3.
 86. Czochrowska EM, Stenvik A, Zachrisson BU. The esthetic outcome of autotransplanted premolars replacing maxillary incisors. *Dent Traumatol.* 2002;18:237–45.
 87. Gable TO, Kummer AW, Lee L, Creaghead NA, Moore LJ. Premature loss of the maxillary primary incisors: effect on speech production. *ASDC J Dent Child.* 1995;62:173–9.

88. Soo SY, Satterthwaite JD, Ashley M. Initial management and long-term follow up after the rehabilitation of a patient with severe dentoalveolar trauma: A case report. *Dent Traumatol.* 2020;36:84–8.
89. Rohof ECM, Kerdijk W, Jansma J, Livas C, Ren Y. Autotransplantation of teeth with incomplete root formation: a systematic review and meta-analysis. *Clin Oral Investig.* 2018;22:1613–24.
90. Abd-Elmeguid A, ElSalhy M, Yu DC. Pulp canal obliteration after replantation of avulsed immature teeth: a systematic review. *Dent Traumatol.* 2015;31:437–41.
91. Bosenbecker J, Barbon FJ, Souza Ferreira N, Morgental RD, Boscato N. Tooth discoloration caused by endodontic treatment: a cross-sectional study. *J Esthet Restor Dent.* 2020;32:569–74.
92. Sanz E, Azabal M, Arias A. Quality of life and satisfaction of patients two years after endodontic and dental implant treatments performed by experienced practitioners. *J Dent.* 2022;125:104280.
93. Andersson L, Emami-Kristiansen Z, Högström J. Single-tooth implant treatment in the anterior region of the maxilla for treatment of tooth loss after trauma: a retrospective clinical and interview study. *Dent Traumatol.* 2003;19:126–31.
94. Seehra J, Newton JT, DiBiase AT. Bullying in schoolchildren – its relationship to dental appearance and psychosocial implications: an update for GDPs. *Br Dent J.* 2011;210:411–5.
95. Hakeberg M, Heidari E, Norinder M, Berggren U. A Swedish version of the dental visit satisfaction scale. *Acta Odontol Scand.* 2000;58:19–24.
96. Goyal N, Singh S, Mathur A, Makkar DK, Aggarwal VP, Sharma A, et al. Traumatic dental injuries prevalence and their impact on self-esteem among adolescents in India: a comparative study. *J Clin Diagn Res.* 2017;11:ZC106–10.

97. Keasberry J, Munyombwe T, Duggal M, Day PF. A study of factors that influence the number of visits following traumatic dental injuries. *Br Dent J.* 2013;214:E28.
98. Glendor U, Halling A, Bodin L, Andersson L, Nygren Å, Karlsson G, et al. Direct and indirect time spent on care of dental trauma: a 2-year prospective study of children and adolescents. *Dent Traumatol.* 2000;16:16–23.
99. Al-Jundi SH. Type of treatment, prognosis, and estimation of time spent to manage dental trauma in late presentation cases at a dental teaching hospital: a longitudinal and retrospective study. *Dent Traumatol.* 2004;20:1–5.
100. Glendor U, Halling A, Andersson L, Andreasen JO, Klitz I. Type of treatment and estimation of time spent on dental trauma--a longitudinal and retrospective study. *Swed Dent J.* 1998;22:47–60.
101. Robertson A, Norén JG. Subjective aspects of patients with traumatized teeth a 15-year follow-up study. *Acta Odontol Scand.* 1997;55:142–7.
102. Gustafsson A, Arnrup K, Broberg AG, Bodin L, Berggren U. Child dental fear as measured with the dental subscale of the children's fear survey schedule: the impact of referral status and type of informant (child versus parent). *Community Dent Oral Epidemiol.* 2010;38:256–66.
103. Aartman IH, van Everdingen T, Hoogstraten J, Schuurs AH. Self-report measurements of dental anxiety and fear in children: a critical assessment. *ASDC J Dent Child.* 1998;65:252–8, 229–30.
104. Buchanan H, Niven N. Validation of a facial image scale to assess child dental anxiety. *Int J Paediatr Dent.* 2002;12:47–52.
105. Howard KE, Freeman R. Reliability and validity of a faces version of the modified child dental anxiety scale. *Int J Paediatr Dent.* 2007;17:281–8.
106. Kaakko T, Murtomaa H. Factors predictive of anxiety before oral surgery: efficacy

- of various subject screening measures. *Anesth Prog.* 1999;46:3–9.
107. Skaare AB, Aas ALM, Wang NJ. Enamel defects on permanent successors following luxation injuries to primary teeth and carers' experiences. *Int J Paediatr Dent.* 2015;25:221–8.
 108. Haliti F, Jurić H. The relationship between dental trauma, anxiety and aggression behavior in 7 to14 year old children in Kosovo. *Acta Stomatol Croat.* 2017;51:3–12.
 109. Kelson MC. Consumer collaboration, patient-defined outcomes and the preparation of cochrane reviews. *Health Expect.* 1999;2:129–35.
 110. Schick-Makaroff K, Molzahn A. Strategies to use tablet computers for collection of electronic patient-reported outcomes. *Health Qual Life Outcomes.* 2015;13.
 111. Kyte D, Bishop J, Brettell E, Calvert M, Cockwell P, Dutton M, et al. Use of an electronic patient-reported outcome measure in the management of patients with advanced chronic kidney disease: the RePROM pilot trial protocol. *BMJ Open.* 2018;8:e026080.
 112. Consolo L, Castellini G, Cilluffo S, Basile I, Lusignani M. Electronic patient-reported outcomes (e-PROMs) in palliative cancer care: a scoping review. *J patient-reported outcomes.* 2022;6.
 113. Aiyegbusi OL, Kyte D, Cockwell P, Marshall T, Dutton M, Walmsley-Allen N, et al. Patient and clinician perspectives on electronic patient-reported outcome measures in the management of advanced CKD: a qualitative study. *Am J Kidney Dis.* 2019;74:167–78.