



ON BOUNDEDNESS OF THE GENERALIZED RIESZ POTENTIAL IN LOCAL MORREY-TYPE SPACES

V. I. Burenkov^{1,2,3} · M. A. Senouci^{1,2}

Accepted: 20 October 2022 / Published online: 2 May 2023
© The Author(s) 2023

Abstract

For all admissible values of the numerical parameters sharp sufficient conditions on the functional parameters are obtained ensuring the boundedness of the generalized Riesz potential from one general local Morrey-type space to another one, which, for a certain range of the numerical parameters, coincide with the necessary ones.

Keywords Generalized Riesz potential operator · Boundedness · Local Morrey-type spaces

Introduction

In this paper, we study the boundedness from one general local Morrey-type space to another one of the generalized Riesz potential

$$(I_{\rho(\cdot)}f)(x) = \int_{\mathbb{R}^n} \rho(|x-y|)f(y)dy, \quad x \in \mathbb{R}^n,$$

under certain assumptions on the kernel ρ . Our aim is to generalize the results obtained in [6] for the case of the classical Riesz potential I_α , in which $\rho(t) = t^{\alpha-n}$, $t > 0$, $0 < \alpha < n$. Some of the presented results were stated without proofs in [7]. Let $F, G : A \times B \rightarrow [0, \infty]$. Throughout this paper we say that F is dominated by G uniformly in $x \in A$ and write

$$F \lesssim G \text{ uniformly in } x \in A$$

if there exists $c(B) > 0$ such that

$$F(x, y) \leq c(B)G(x, y) \text{ for all } x \in A.$$

✉ V. I. Burenkov
burenkov@cardiff.ac.uk

M. A. Senouci
senoucim@yandex.ru

¹ S.M. Nikol'skii Mathematical Institute, RUDN University, 6 Miklukho Maklay St, Moscow 117198, Russian Federation

² V.A. Steklov Mathematical Institute, Russian Academy of Sciences, 8 Gubkin St, Moscow 117966, Russian Federation

³ Cardiff School of Mathematics, Cardiff University, Senghennydd Rd, Cardiff CF24 4AG, United Kingdom

(So $c(B)$ is independent of $x \in A$, but may depend on $y \in B$.) Respectively, we say that F dominates G uniformly in $x \in A$ and write

$$F \gtrsim G \text{ uniformly in } x \in A$$

if there exists $c(B) > 0$ such that

$$F(x, y) \geq c(B)G(x, y) \text{ for all } x \in A.$$

We also say that F is equivalent to G uniformly in $x \in A$ and write

$$F \approx G \text{ uniformly in } x \in A$$

if F and G dominate each other uniformly in $x \in A$.

Definitions and basic properties of general Morrey-type spaces

In this section we recall basic facts of the theory of general Morrey-type spaces.

Let, for a Lebesgue measurable set $\Omega \subset \mathbb{R}^n$, $\mathfrak{M}(\Omega)$ denote the space of all functions $f : \Omega \rightarrow \mathbb{C}$ Lebesgue measurable on Ω , and $\mathfrak{M}^+(\Omega)$ denote the subset of $\mathfrak{M}(\Omega)$ of all non-negative functions.

Definition 2.1 Let $0 < p, \theta \leq \infty$ and let $w \in \mathfrak{M}^+((0, \infty))$ be not equivalent to 0. We denote by $LM_{p\theta, w(\cdot)}$ the local Morrey-type space, the space of all functions $f \in L_p^{\text{loc}}(\mathbb{R}^n)$ with finite quasinorm

$$\|f\|_{LM_{p\theta, w(\cdot)}} \equiv \|f\|_{LM_{p\theta, w(\cdot)}(\mathbb{R}^n)} = \|w(r)\|f\|_{L_p(B(0, r))}\|_{L_\theta(0, \infty)}.$$

Definition 2.2 Let $0 < p, \theta \leq \infty$. We denote by Ω_θ the set of all functions $w \in \mathfrak{M}^+((0, \infty))$ which are not equivalent to 0 and such that

$$\|w\|_{L_\theta(t, \infty)} < \infty \tag{2.1}$$

for some $t > 0$.

Lemma 2.1 [3–5]. Let $0 < p, \theta \leq \infty$ and let w be a non-negative Lebesgue measurable function on $(0, \infty)$, which is not equivalent to 0.

Then the space $LM_{p\theta, w(\cdot)}$ is nontrivial if and only if $w \in \Omega_\theta$.

In the sequel, keeping in mind Lemma 2.1, we always assume that $w \in \Omega_\theta$ for the local Morrey-type spaces $LM_{p\theta, w(\cdot)}$. It is well known that the spaces $LM_{p\theta, w(\cdot)}$ are Banach spaces if $1 \leq p, \theta \leq \infty$ and are quasi-Banach spaces if $0 < p < 1$, or $0 < \theta < 1$, or both $0 < p, \theta < 1$.

For further properties of general Morrey-type spaces, operators acting in such spaces and applications see, for example, survey papers [1, 2, 12–16] and references therein.

L_p and WL_p estimates of generalized Riesz potentials over balls

Let $n \in \mathbb{N}$, $0 < p < \infty$, $\Omega \subset \mathbb{R}^n$ be a Lebesgue measurable set. Recall that the weak L_p -space $WL_p(\Omega)$ is the space of all Lebesgue measurable functions $f : \Omega \rightarrow \mathbb{C}$ for which

$$\|f\|_{WL_p(\Omega)} = \sup_{t \geq 0} t |\{y \in \Omega : |f(y)| > t\}|^{\frac{1}{p}} < \infty.$$

Here $|G|$ denotes the Lebesgue measure of a set $G \subset \mathbb{R}^n$.

Remark 3.1 Let $M > 0$. It follows directly from the above definition that, if f is equivalent to M on Ω , then

$$\|M\|_{WL_p(\Omega)} = \|M\|_{L_p(\Omega)} = M|\Omega|^{\frac{1}{p}}$$

and if $f : \Omega \rightarrow \mathbb{R}$ is such that $f \geq M$ almost everywhere on Ω , then

$$\|f\|_{WL_p(\Omega)} \geq M|\Omega|^{\frac{1}{p}}.$$

Definition 3.1 Let $n \in \mathbb{N}$. We say that $\rho \in S_n$ if $\rho \in \mathfrak{M}^+((0, \infty))$ and

- 1) $\int_0^r \rho(t)t^{n-1} dt < \infty$ for all $r > 0$,
- 2) for some $c_1, c_2 > 0$,

$$c_1\rho(t) \leq \rho(s) \leq c_2\rho(t)$$

for all $s, t > 0$ satisfying the inequality $\frac{t}{2} \leq s \leq 2t$.

Remark 3.2 Let $x \in \mathbb{R}^n, r > 0, y \in B(x, r), z \in {}^cB(x, 2r)$. Then

$$|y - z| \leq |y - x| + |x - z| \leq r + |x - z| \leq \frac{1}{2}|x - z| + |x - z| < 2|x - z|$$

and

$$|y - z| \geq |x - z| - |y - x| \geq |x - z| - r \geq |x - z| - \frac{1}{2}|x - z| = \frac{1}{2}|x - z|.$$

Let $s = |y - z|$ and $t = |x - z|$. Then by Condition 2) of Definition 3.1 we have

- 3) $c_1\rho(|x - z|) \leq \rho(|y - z|) \leq c_2\rho(|x - z|)$
for all $x \in \mathbb{R}^n, r > 0, y \in B(x, r), z \in {}^cB(x, 2r)$.

Remark 3.3 If functions $\rho_1, \rho_2 : (0, \infty) \rightarrow (0, \infty)$ satisfy Condition 2) of Definition 3.1 with $c_{11}, c_{12} > 0, c_{21}, c_{22} > 0$, respectively, then the product $\rho_1\rho_2$ satisfies Condition 2) with $c_1 = c_{11}c_{21}$ and $c_2 = c_{12}c_{22}$. Indeed, it suffices to multiply the inequalities

$$c_{i1} \rho_i(t) \leq \rho_i(s) \leq c_{i2}\rho_i(t),$$

where $s, t > 0, \frac{t}{2} \leq s \leq 2t$ and $i = 1, 2$.

Remark 3.4 If a function $\rho_1 : (0, 1] \rightarrow (0, \infty)$ satisfies Condition 2) of Definition 3.1 with $c_{11}, c_{12} > 0$ and a function $\rho_2 : (1, \infty) \rightarrow (0, \infty)$ satisfies Condition 2) of Definition 3.1 with $c_{21}, c_{22} > 0$, then the function

$$\rho(t) = \begin{cases} \rho_1(t), & 0 < t \leq 1, \\ \rho_2(t), & 1 < t < \infty \end{cases}$$

satisfies Condition 2) of Definition 3.1 with certain $c_1, c_2 > 0$.

Indeed, let $s, t > 0, \frac{t}{2} \leq s \leq 2t$. If $t \leq \frac{1}{2}$, then $s \leq 1, \rho(s) = \rho_1(s), \rho(t) = \rho_1(t)$ and Condition 2) of Definition 3.1 is satisfied with $c_{11}, c_{12} > 0$. Respectively, if $t > 2$, then $s > 1, \rho(s) = \rho_2(s), \rho(t) = \rho_2(t)$ and Condition 2) of Definition 3.1 is satisfied with $c_{21}, c_{22} > 0$. Let $\frac{1}{2} < t \leq 2$, then $\frac{1}{4} < s \leq 4$,

$$\min\{c_{11}, c_{21}\} \leq \begin{cases} c_{11}, & \frac{1}{4} < s \leq 1, \\ c_{21}, & 1 < s < 4 \end{cases} \leq \rho(s) \leq \begin{cases} c_{12}, & \frac{1}{4} < s \leq 1, \\ c_{22}, & 1 < s < 4 \end{cases} \leq \max\{c_{12}, c_{22}\},$$

$$\min\{c_{11}, c_{21}\} \leq \begin{cases} c_{11}, & \frac{1}{2} < t \leq 1, \\ c_{21}, & 1 < t \leq 2 \end{cases} \leq \rho(t) \leq \begin{cases} c_{12}, & \frac{1}{2} < t \leq 1, \\ c_{22}, & 1 < t \leq 2 \end{cases} \leq \max\{c_{12}, c_{22}\},$$

hence

$$\min\{c_{11}, c_{21}\}(\max\{c_{12}, c_{22}\})^{-1}\rho(t) \leq \rho(s) \leq \max\{c_{12}, c_{22}\}(\min\{c_{11}, c_{21}\})^{-1}\rho(t).$$

So, for $\frac{1}{2} < t \leq 2, \frac{t}{2} \leq s \leq 2t$, Condition 2) of Definition 3.1 is satisfied with

$$c_1 = \min\{c_{11}, c_{21}\} \max\{c_{12}, c_{22}\}^{-1}, \quad c_2 = \max\{c_{12}, c_{22}\} \min\{c_{11}, c_{21}\}^{-1}.$$

The above inequality is satisfied for all $s, t > 0, \frac{t}{2} \leq s \leq 2t$, because $c_{11} \leq 1 \leq c_{12}, c_{21} \leq 1 \leq c_{22}$, hence $c_1 \leq c_{11}, c_1 \leq c_{21}, c_{12} \leq c_2, c_{22} \leq c_2$.

Remark 3.5 If a function $\rho : (0, \infty) \rightarrow (0, \infty)$ satisfies Condition 2) of Definition 3.1, then for any $\gamma > 0, c_{1\gamma}, c_{2\gamma} > 0$ such that for any $t > 0$

$$c_{1\gamma}\rho(t) \leq \rho(\gamma t) \leq c_{2\gamma}\rho(t).$$

Indeed, if $\frac{1}{2} \leq \gamma \leq 2$, then this inequality follows by Condition 2) with $s = \gamma t$. Let $2 \leq \gamma < 4$, then by Condition 2) with $t = 2\tau, \tau > 0$ and $s = \gamma\tau$ and the above inequality with $t = \tau$ and $\gamma = 2$

$$c_1^2\rho(\tau) \leq c_1\rho(2\tau) \leq \rho(\gamma\tau) \leq c_2\rho(2\tau) \leq c_2^2\rho(\tau), \quad \tau > 0,$$

because $\frac{1}{2}(2\tau) \leq \gamma\tau \leq 2(2\tau)$.

Next, let $4 \leq \gamma < 8$ by Condition 2) with $t = 4\tau, \tau > 0, s = \gamma\tau$,

$$\frac{1}{2}(4\tau) \leq \gamma\tau \leq 2(4\tau),$$

then

$$c_1^3\rho(\tau) \leq c_1^2\rho(2\tau) \leq c_1\rho(4\tau) \leq \rho(\gamma\tau) \leq c_2\rho(4\tau) \leq c_2^2\rho(2\tau) \leq c_2^3\rho(\tau), \quad \tau > 0.$$

Furthermore, if $2^{k-1} \leq \gamma < 2^k, k \in \mathbb{N}, k > 3 \Leftrightarrow k = \left\lceil \frac{\ln \gamma}{\ln 2} \right\rceil + 1$, then

$$c_1^k\rho(\tau) \leq c_1^{k-1}\rho(2\tau) \leq \dots \leq \rho(\gamma\tau) \leq c_2^2\rho(2\tau) \leq \dots \leq c_2^k\rho(\tau).$$

The argument for the case $0 < \gamma < \frac{1}{2}$ is similar.

Example 1 Let $\rho(t) = t^{\alpha-n}, t > 0, 0 < \alpha < n$. Then $\rho \in S_n$ because

$$\int_0^r t^{\alpha-n+n-1} dt < \infty \text{ for all } 0 < r < \infty \Leftrightarrow \alpha > 0$$

and, for any $s, t > 0$ for which $\frac{t}{2} \leq s \leq 2t$,

$$\begin{aligned} 2^{\alpha-n}\rho(s) &= 2^{\alpha-n}s^{\alpha-n} = (2s)^{\alpha-n} \\ &\leq \rho(t) = t^{\alpha-n} \leq \left(\frac{s}{2}\right)^{\alpha-n} = 2^{n-\alpha}s^{\alpha-n} = 2^{n-\alpha}\rho(s). \end{aligned}$$

Example 2 Let $0 < \alpha < n, -\infty < \beta_1, \beta_2 < \infty$,

$$\Psi_{\beta_1, \beta_2}(t) = \begin{cases} (1 + |\ln t|)^{\beta_1}, & 0 < t \leq 1, \\ (1 + |\ln t|)^{\beta_2}, & 1 < t < \infty \end{cases}$$

and $\rho(t) = t^{\alpha-n}\Psi_{\beta_1, \beta_2}(t), t > 0$. Then $\rho \in S_n$.

Since the function ρ is continuous on $(0, \infty)$ it suffices to prove Condition 1) of Definition 3.1 for $r = 1$. Recall that for any $\varepsilon > 0$ there exists $C_\varepsilon \geq 1$ such that

$$1 + |\ln t| \leq C_\varepsilon t^{-\varepsilon}$$

for all $t \in (0, 1]$. If $\beta_1 \leq 0$, then

$$\int_0^1 \rho(t)t^{n-1} dt = \int_0^1 t^{\alpha-1}(1 + |\ln t|)^{\beta_1} dt \leq \int_0^1 t^{\alpha-1} dt < \infty.$$

If $\beta_1 > 0$, then

$$\int_0^1 \rho(t)t^{n-1} dt = \int_0^1 t^{\alpha-1}(1 + |\ln t|)^{\beta_1} dt \leq C_\varepsilon^{\beta_1} \int_0^1 t^{\alpha-\varepsilon\beta_1-1} dt < \infty$$

if we choose $\varepsilon > 0$ to be such that $\varepsilon\beta_1 < \alpha$.

As for Condition 2) of Definition 3.1, by Remark 3.3 and Example 1, it suffices to prove that the function $\varphi_{\beta_1, \beta_2}$ satisfies this condition.

Let $s, t > 0$ and $\frac{t}{2} \leq s \leq 2t$. If $s > 1$, hence $t > \frac{1}{2}$, then

$$1 + |\ln s| \leq 1 + \ln 2t = 1 + \ln 2 + \ln t \leq 2(1 + |\ln t|)$$

and

$$1 + |\ln t| \leq 1 + \ln 2 < 2(1 + |\ln s|),$$

hence

$$\frac{1}{2}(1 + |\ln t|) \leq 1 + |\ln s| \leq 2(1 + |\ln t|).$$

For similar reasons this inequality also holds if $s \leq 1$, hence $t \leq 2$. Therefore, for $s \leq 1$

$$2^{-|\beta_1|}(1 + |\ln t|)^{\beta_1} \leq (1 + |\ln s|)^{\beta_1} \leq 2^{|\beta_1|}(1 + |\ln t|)^{\beta_1}.$$

If $t \leq 1$, this means that

$$2^{-|\beta_1|}\Psi_{\beta_1, \beta_2}(t) \leq \Psi_{\beta_1, \beta_2}(s) \leq 2^{|\beta_1|}\Psi_{\beta_1, \beta_2}(t).$$

If $1 < t \leq 2$, then

$$2^{-|\beta_1| - |\beta_2|} \leq (1 + |\ln t|)^{\beta_1} (1 + |\ln t|)^{-\beta_2} \leq 2^{|\beta_1| + |\beta_2|},$$

hence

$$2^{-2|\beta_1| - |\beta_2|} (1 + |\ln t|)^{\beta_2} \leq (1 + |\ln s|)^{\beta_1} \leq 2^{2|\beta_1| + |\beta_2|} (1 + |\ln t|)^{\beta_2}$$

which means that

$$2^{-2|\beta_1| - |\beta_2|}\Psi_{\beta_1, \beta_2}(t) \leq \Psi_{\beta_1, \beta_2}(s) \leq 2^{2|\beta_1| + |\beta_2|}\Psi_{\beta_1, \beta_2}(t).$$

So, this inequality holds for all $s \leq 1$ and all t satisfying the inequality $\frac{t}{2} \leq s \leq 2t$.

For similar reasons, for all $s > 1$ and all t satisfying the inequality $\frac{t}{2} \leq s \leq 2t$

$$2^{-|\beta_1| - 2|\beta_2|}\Psi_{\beta_1, \beta_2}(t) \leq \Psi_{\beta_1, \beta_2}(s) \leq 2^{|\beta_1| + 2|\beta_2|}\Psi_{\beta_1, \beta_2}(t).$$

Example 3 Let $0 < \alpha < n, -\infty < \beta < \infty$ and

$$\rho(t) = \begin{cases} t^{\alpha-n}, & 0 < t \leq 1, \\ t^\beta, & 1 < t < \infty. \end{cases}$$

Then $\rho \in S_n$. Indeed, Condition 1) is clearly satisfied. As for Condition 2), it is also satisfied by Remark 3.4, because $\rho_1(t) = t^{\alpha-n}$ satisfies Condition 2) on $(0, 1]$ by Example 1 and $\rho_2(t) = t^\beta$ satisfies Condition 2) on $(1, \infty)$ by an argument similar to that of the proof of Example 1.

Note that t^β can be replaced by any function $\rho(t)$ satisfying Condition 2) on $(1, \infty)$.

Definition 3.2 For $\rho \in S_n$ and $f \in \mathfrak{M}(\mathbb{R}^n)$

$$I_{\rho(\cdot)}f(x) = \int_{\mathbb{R}^n} \rho(|x - y|)f(y)dy,$$

$$\bar{I}_{\rho(\cdot),r}f(x) = \int_{cB(x,r)} \rho(|x - y|)f(y)dy$$

and

$$\underline{I}_{\rho(\cdot),r}f(x) = \int_{B(x,r)} \rho(|x - y|)f(y)dy.$$

Lemma 3.1 Let $n \in \mathbb{N}, \rho \in S_n, 0 < p \leq \infty, x \in \mathbb{R}^n, r > 0, f \in \mathfrak{M}(\mathbb{R}^n)$. Then¹ $G = r^{\frac{n}{p}} \left(\bar{I}_{\rho(\cdot),2r}|f| \right)(x)$.

$$\|I_{\rho(\cdot)}|f|\|_{WL_p(B(x,r))} \gtrsim r^{\frac{n}{p}} \left(\bar{I}_{\rho(\cdot),2r}|f| \right)(x) \tag{3.1}$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in \mathfrak{M}(\mathbb{R}^n)$.

Proof By Property 3) of Remark 3.2 for any $y \in B(x, r)$

$$\begin{aligned} I_{\rho(\cdot)}|f|(y) &\geq \int_{cB(x,2r)} \rho(|y - z|)|f(z)|dz \\ &\geq c_1 \int_{cB(x,2r)} \rho(|x - z|)|f(z)|dz, \end{aligned}$$

hence, by Remark 3.1, we get

$$\begin{aligned} \|I_{\rho(\cdot)}|f|\|_{WL_p(B(x,r))} &\geq c_1 (v_n r^n)^{\frac{1}{p}} \int_{cB(x,2r)} \rho(|x - z|)|f(z)|dz \\ &= c_1 v_n^{\frac{1}{p}} r^{\frac{n}{p}} \bar{I}_{\rho(\cdot),2r}|f|(x) \end{aligned}$$

¹ According to the definitions introduced in [Introduction](#), this means that for any $n \in \mathbb{N}, \rho \in S_n, 0 < p \leq \infty$ there exists $c(n, \rho, p) > 0$ such that

$$\|I_{\rho(\cdot)}|f|\|_{WL_p(B(x,r))} \gtrsim c(n, \rho, p) r^{\frac{n}{p}} \left(\bar{I}_{\rho(\cdot),2r}|f| \right)(x)$$

for all $x \in \mathbb{R}^n, r > 0$ and $f \in \mathfrak{M}(\mathbb{R}^n)$.

In this case $A = \{\mathbb{R}^n, (0, \infty), \mathfrak{M}(\mathbb{R}^n)\}, B = \{(n, \rho) : n \in \mathbb{N}, \rho \in S_n; [0, \infty], F = \|I_{\rho(\cdot)}|f|\|_{WL_p(B(x,r))}$

for all $x \in \mathbb{R}^n, r > 0$ and $f \in \mathfrak{M}(\mathbb{R}^n)$. Here v_n is the volume of the unit ball in \mathbb{R}^n .

Lemma 3.2 *Let $n \in \mathbb{N}, \rho \in S_n, 0 < p \leq \infty, x \in \mathbb{R}^n, r > 0, f \in \mathfrak{M}(\mathbb{R}^n)$. Then*

$$\|I_{\rho(\cdot)}|f|\|_{L_p(B(x,r))} \approx \|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{L_p(B(x,r))} + r^{\frac{n}{p}} \left(\bar{I}_{\rho(\cdot),2r}|f|\right)(x) \tag{3.2}$$

and

$$\|I_{\rho(\cdot)}|f|\|_{WL_p(B(x,r))} \approx \|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{WL_p(B(x,r))} + r^{\frac{n}{p}} \left(\bar{I}_{\rho(\cdot),2r}|f|\right)(x) \tag{3.3}$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in \mathfrak{M}(\mathbb{R}^n)$.

Proof According to the properties² of the spaces $L_p(\Omega)$ and $WL_p(\Omega)$

$$\|I_{\rho(\cdot)}|f|\|_{L_p(B(x,r))} \leq 2^{\left(\frac{1}{p}-1\right)_+} \left(\|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{L_p(B(x,r))} + \|I_{\rho(\cdot)}(|f|\chi_{cB(x,2r)})\|_{L_p(B(x,r))} \right), \tag{3.4}$$

$$\|I_{\rho(\cdot)}|f|\|_{WL_p(B(x,r))} \leq 2^{\frac{1}{p}} \left(\|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{WL_p(B(x,r))} + \|I_{\rho(\cdot)}(|f|\chi_{cB(x,2r)})\|_{WL_p(B(x,r))} \right). \tag{3.5}$$

$$\|I_{\rho(\cdot)}|f|\|_{L_p(B(x,r))} \geq \frac{1}{2} \left(\|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{L_p(B(x,r))} + \|I_{\rho(\cdot)}(|f|\chi_{cB(x,2r)})\|_{L_p(B(x,r))} \right), \tag{3.6}$$

and

$$\|I_{\rho(\cdot)}|f|\|_{L_p(B(x,r))} \|I_{\rho(\cdot)}|f|\|_{WL_p(B(x,r))} \geq \|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{WL_p(B(x,r))}. \tag{3.7}$$

Next,

$$\begin{aligned} \|I_{\rho(\cdot)}(|f|\chi_{cB(x,2r)})\|_{WL_p(B(x,r))} &\leq \|I_{\rho(\cdot)}(|f|\chi_{cB(x,2r)})\|_{L_p(B(x,r))} \\ &= \left[\int_{B(x,r)} \left(\int_{cB(x,2r)} \rho(|y-z|)|f(z)|dz \right)^p dy \right]^{\frac{1}{p}} = J. \end{aligned}$$

By Condition 3) of Remark 3.2

$$J \leq_{C2} \left[\int_{B(x,r)} \left(\int_{cB(x,2r)} \rho(|x-z|)|f(z)|dz \right)^p dy \right]^{\frac{1}{p}} =_{C2} v_n^{\frac{1}{p}} r^{\frac{n}{p}} \left(\bar{I}_{\rho(\cdot),2r}|f|\right)(x).$$

Thus, by (3.4) and (3.5)

$$\|I_{\rho(\cdot)}|f|\|_{L_p(B(x,r))} \lesssim \|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{L_p(B(x,r))} + r^{\frac{n}{p}} \left(\bar{I}_{\rho(\cdot),2r}|f|\right)(x),$$

and

$$\|I_{\rho(\cdot)}|f|\|_{WL_p(B(x,r))} \lesssim \|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{WL_p(B(x,r))} + r^{\frac{n}{p}} \left(\bar{I}_{\rho(\cdot),2r}|f|\right)(x),$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in \mathfrak{M}(\mathbb{R}^n)$.

² Below $a_+ = a$ if $a \geq 0$ and $a_+ = 0$ if $a < 0$.

Also, by Condition 3) of Remark 3.2

$$J \geq_{C1} \left[\int_{B(x,r)} \left(\int_{\epsilon B(x,2r)} \rho(|x-z|)|f(z)|dz \right)^p dy \right]^{\frac{1}{p}} =_{C1}^{-1} v_n^{\frac{1}{p}} r^{\frac{n}{p}} \left(\tilde{I}_{\rho(\cdot),2r}|f| \right)(x).$$

Hence, by (3.6)

$$\|I_{\rho(\cdot)}|f|\|_{L_p(B(x,r))} \gtrsim \|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{L_p(B(x,r))} + r^{\frac{n}{p}} \left(\tilde{I}_{\rho(\cdot),2r}|f| \right)(x),$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in \mathfrak{M}(\mathbb{R}^n)$.

Finally, by adding inequalities (3.1) and (3.7), we get that

$$\|I_{\rho(\cdot)}|f|\|_{WL_p(B(x,r))} \gtrsim \|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{WL_p(B(x,r))} + r^{\frac{n}{p}} \left(\tilde{I}_{\rho(\cdot),2r}|f| \right)(x),$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in \mathfrak{M}(\mathbb{R}^n)$.

Lemma 3.3 *Let, for $n \in \mathbb{N}, \rho \in S_n, 1 \leq p_1 < p_2 < \infty$, and*

$$\varphi_{n,\rho,p_1,p_2}(t) = \rho(t)t^{n\left(\frac{1}{p_1'} + \frac{1}{p_2}\right)}, \quad t > 0, \tag{3.8}$$

where p_1' is the conjugate number to p_1 .

1. Assume that ρ is such that the function φ_{n,ρ,p_1,p_2} is almost non-decreasing on $(0, \infty)$, that is for some $c > 0$

$$\varphi_{n,\rho,p_1,p_2}(t_1) \leq c\varphi_{n,\rho,p_1,p_2}(t_2) \text{ for all } 0 < t_1 < t_2 < \infty. \tag{3.9}$$

If $1 < p_1 < p_2 < \infty$, then

$$\|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{L_{p_2}(B(x,r))} \lesssim \varphi_{n,\rho,p_1,p_2}(r)\|f\|_{L_{p_1}(B(x,2r))} \tag{3.10}$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in L_{p_1}(B(x, 2r))$.

Also, if $p_1 = 1, 1 < p_2 < \infty$, then

$$\|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{WL_{p_2}(B(x,r))} \lesssim \varphi_{n,\rho,p_1,p_2}(r)\|f\|_{L_{p_1}(B(x,2r))} \tag{3.11}$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in L_1(B(x, 2r))$.

2. Inequality (3.10) also holds for $p_1 = 1$ and $1 \leq p_2 \leq \infty$ under the assumption

$$\|\rho(t)t^{\frac{n-1}{p_2}}\|_{L_{p_2}(0,r)} \lesssim \rho(r)r^{\frac{n}{p_2}} \tag{3.12}$$

uniformly in $r > 0$, replacing assumption (3.9).

3. Inequality (3.10) and hence inequality (3.11) also holds for $1 \leq p_1 < \infty$ and $0 < p_2 \leq p_1$ under the assumption

$$\|\rho(t)t^{n-1}\|_{L_1(0,r)} \lesssim \rho(r)r^n \tag{3.13}$$

uniformly in $r > 0$, replacing assumption (3.9).

4. Condition (3.13) is necessary for the validity of inequalities (3.10) and (3.11) for all $0 < p_1, p_2 \leq \infty$.

5. Condition (3.13) is necessary and sufficient for the validity of inequalities (3.10) and (3.11) for all $1 \leq p_1 < \infty$ and $0 < p_2 \leq p_1$.

Proof 1. Let $1 \leq p_1 < p_2 < \infty$, $x \in \mathbb{R}^n$, $r > 0$, $z \in B(x, r)$ and $f \in L_{p_1}(B(x, 2r))$. Then

$$\begin{aligned} \left(I_{\rho(\cdot)} \left(|f| \chi_{B(x, 2r)} \right) \right) (z) &= \int_{\mathbb{R}^n} \rho(|z - y|) |f(y)| \chi_{B(x, 2r)}(y) dy \\ &= \int_{B(x, 2r)} \rho(|z - y|) |z - y|^{n \left(1 - \frac{1}{p_1} + \frac{1}{p_2} \right)} \frac{|f(y)| \chi_{B(x, 2r)}(y)}{|z - y|^{n - n \left(\frac{1}{p_1} - \frac{1}{p_2} \right)}} dy \\ &\leq \left(\sup_{\substack{z \in B(x, r) \\ y \in B(x, 2r)}} \varphi_{n, \rho, p_1, p_2}(|z - y|) \right) \left(I_{n \left(\frac{1}{p_1} - \frac{1}{p_2} \right)} \left(|f| \chi_{B(x, 2r)} \right) \right) (z). \end{aligned}$$

Since the function $\varphi_{n, \rho, p_1, p_2}$ is almost non-decreasing on $(0, \infty)$ and, for $z \in B(x, r)$, $y \in B(x, 2r)$,

$$|z - y| \leq |z - x| + |x - y| \leq r + 2r = 3r,$$

we have by (3.9)

$$\sup_{\substack{z \in B(x, r) \\ y \in B(x, 2r)}} \varphi_{n, \rho, p_1, p_2}(|z - y|) \leq c \varphi_{n, \rho, p_1, p_2}(3r) = c 3^{n \left(1 - \frac{1}{p_1} + \frac{1}{p_2} \right)} \rho(3r) r^{n \left(1 - \frac{1}{p_1} + \frac{1}{p_2} \right)}.$$

By Remark 3.5 with $\gamma = 3$ it follows that

$$\rho(3r) \leq c_{23} \rho(r), \quad r > 0.$$

Hence,

$$\sup_{\substack{z \in B(x, r) \\ y \in B(x, 2r)}} \varphi_{n, \rho, p_1, p_2}(|z - y|) \leq c c_{23} 3^{n \left(1 - \frac{1}{p_1} + \frac{1}{p_2} \right)} \varphi_{n, \rho, p_1, p_2}(r)$$

and

$$\left(I_{\rho(\cdot)} \left(|f| \chi_{B(x, 2r)} \right) \right) (z) \lesssim \varphi_{n, \rho, p_1, p_2}(r) \left(I_{n \left(\frac{1}{p_1} - \frac{1}{p_2} \right)} \left(|f| \chi_{B(x, 2r)} \right) \right) (z) \tag{3.14}$$

uniformly in $x \in \mathbb{R}^n$, $r > 0$, $z \in B(x, r)$ and $f \in L_{p_1}(B(x, 2r))$.

Next, we apply the well-known inequalities for the Riesz potential. If $1 < p_1 < p_2 < \infty$, then

$$\| I_{n \left(\frac{1}{p_1} - \frac{1}{p_2} \right)} |f| \|_{L_{p_2}(\mathbb{R}^n)} \lesssim \| f \|_{L_{p_1}(\mathbb{R}^n)} \tag{3.15}$$

uniformly in $f \in L_{p_1}(\mathbb{R}^n)$. Also, if $1 < p_2 < \infty$, then

$$\| I_{n \left(1 - \frac{1}{p_2} \right)} |f| \|_{WL_q(\mathbb{R}^n)} \lesssim \| f \|_{L_1(\mathbb{R}^n)} \tag{3.16}$$

uniformly in $L_1(\mathbb{R}^n)$.

If $1 < p_1 < p_2 < \infty$, then, by (3.14) and (3.15), it follows that

$$\begin{aligned} \|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{L_{p_2}(\mathbb{R}^n)} &\lesssim \varphi_{n,\rho,p_1,p_2}(r) \| |f|\chi_{B(x,2r)} \|_{L_{p_1}(\mathbb{R}^n)} \\ &= \varphi_{n,\rho,p_1,p_2}(r) \|f\|_{L_{p_1}(B(x,2r))} \end{aligned}$$

uniformly in $x \in \mathbb{R}^n$, $r > 0$ and $f \in L_{p_1}(B(x, 2r))$, which is inequality (3.10).

Respectively, if $1 < p_2 < \infty$, then by (3.14) and (3.16) there follows inequality (3.11).

2. If $p_1 = 1$ and $1 \leq p_2 \leq \infty$, then, by applying Young's inequality for truncated convolutions, we get

$$\begin{aligned} \|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{L_{p_2}(B(x,r))} &= \left\| \int_{B(x,2r)} \rho(|-y|)|f(y)|dy \right\|_{L_{p_2}(B(x,r))} \\ &\leq \rho \| \chi_{L_{p_2}(B(x,r)-B(x,2r))} \| f \|_{L_1(B(x,2r))} \\ &= \rho \| \chi_{L_{p_2}(B(0,3r))} \| f \|_{L_1(B(x,2r))} \\ &= \sigma_n^{\frac{1}{p_2}} \| \rho(t)t^{\frac{n-1}{p_2}} \|_{L_{p_2}(0,3r)} \| f \|_{L_1(B(x,2r))}, \end{aligned}$$

where $\sigma_n = nv_n$ is the surface area of the unit ball in \mathbb{R}^n .

By inequality (3.12) and Remark 3.5 with $\gamma = 3$ it follows that

$$\| \rho(t)t^{\frac{n-1}{p_2}} \|_{L_{p_2}(0,3r)} \lesssim \rho(3r)r^{\frac{n}{p_2}} \lesssim \rho(r)r^{\frac{n}{p_2}} = \varphi_{n,\rho,1,p_2}(r),$$

which implies inequality (3.10).

3. If $1 \leq p_2 = p_1 < \infty$, then similarly to Step 2

$$\begin{aligned} \|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{L_{p_1}(B(x,r))} &\leq \rho \| \chi_{L_1(B(x,r)-B(x,2r))} \| f \|_{L_{p_1}(B(x,2r))} \\ &= \sigma_n \| \rho(t)t^{n-1} \|_{L_1(0,3r)} \| f \|_{L_{p_1}(B(x,2r))}. \end{aligned}$$

By inequality (3.13) and Remark 3.5 with $\gamma = 3$ it follows that

$$\| \rho(t)t^{n-1} \|_{L_1(0,3r)} \lesssim \rho(3r)r^n \lesssim \rho(r)r^n = \varphi_{n,\rho,p_1,p_1}(r),$$

which implies inequality (3.10).

If $0 < p_2 < p_1$, $1 \leq p_1 < \infty$, then, by applying Hölder's inequality and inequality (3.10) with $p_2 = p_1$ we get

$$\begin{aligned} \|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{L_{p_2}(B(x,r))} &\leq (v_n r^n)^{\frac{1}{p_2} - \frac{1}{p_1}} \|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{L_{p_1}(B(x,r))} \\ &\lesssim \rho(r)r^{n\left(\frac{1}{p_1} + \frac{1}{p_2}\right)} \|f\|_{L_{p_1}(B(x,2r))} = \varphi_{n,\rho,p_1,p_2}(r) \|f\|_{L_{p_1}(B(x,2r))}, \end{aligned}$$

which is inequality (3.10).

4. Assume that for some $0 < p_1, p_2 \leq \infty$ inequality (3.10) and (3.11) is satisfied. If inequality (3.10) is satisfied, then inequality (3.11) is also satisfied. Take in this inequality $x = 0$ and $f \equiv 1$. Then for any $y \in B(0, r)$ we have $B(y, 2r) \supset B(0, r)$ and

$$\begin{aligned}
 I_{\rho(\cdot)}(\mathcal{X}_{B(0,2r)})(y) &= \int_{B(0,2r)} \rho(y-z)dz = \int_{B(y,2r)} \rho(u)du \\
 &\geq \int_{B(0,r)} \rho(u)du = \sigma_n \int_0^r \rho(t)t^{n-1} dt.
 \end{aligned}$$

By Remark 3.1

$$\left\| I_{\rho(\cdot)}(\mathcal{X}_{B(0,2r)}) \right\|_{WL_{p_2}(B(0,r))} \gtrsim r^{\frac{n}{p_2}} \int_0^r \rho(t)t^{n-1} dt$$

and $\left\| \mathcal{X}_{B(0,2r)} \right\|_{L_{p_2}(B(0,r))} = (\sigma_n r^n)^{\frac{1}{p_1}}$, hence by (3.11)

$$r^{\frac{n}{p_2}} \int_0^r \rho(t)t^{n-1} dt \lesssim \rho(r)r^n \left(\frac{1}{p_1} + \frac{1}{p_2}\right) r^{\frac{n}{p_1}}$$

uniformly in $r > 0$, which implies inequality (3.13).

5. The last statement of Lemma 3.3 follows by Steps 3 and 4.

Remark 3.6 If $1 \leq p_1 < p_2 < \infty$ and

$$\lim_{t \rightarrow 0^+} \varphi_{n,\rho,p_1,p_2}(t) = 0, \tag{3.17}$$

then inequality (3.10) for $1 < p_1 < p_2 < \infty$ and inequality (3.11) for $1 < p_2 < \infty$ cannot hold for any $f \in L_{p_1}(\mathbb{R}^n)$, $f \in L_1(\mathbb{R}^n)$, respectively, not equivalent to 0 on \mathbb{R}^n . Indeed, by passing to the limit as $r \rightarrow \infty$, in (3.10) and (3.11), it follows that $\| I_{\rho(\cdot)}|f| \|_{L_{p_2}(\mathbb{R}^n)} = 0, \| I_{\rho(\cdot)}|f| \|_{WL_{p_2}(\mathbb{R}^n)} = 0$, hence, f is equivalent to 0 on \mathbb{R}^n .

If $\rho(t) = t^{\alpha-n}, 0 < \alpha < n$ and $1 \leq p_1 < p_2$, then Condition (3.9) reduces to the condition $\alpha \geq n(\frac{1}{p_1} - \frac{1}{p_2})$, Condition (3.12) reduces to the condition $\alpha > n(1 - 1/p_2)$ and Condition (3.13) is satisfied since $\alpha > 0$.

Corollary 3.1 *Let the assumptions of Lemma 3.3 for n, ρ, p_1, p_2 and the function φ_{n,ρ,p_1,p_2} be satisfied. Then, for $1 < p_1 < p_2 < \infty$, for $p_1 = 1, 1 < p_2 \leq \infty$, and for $1 \leq p_1 < \infty, 0 < p_2 \leq p_1$*

$$\| I_{\rho(\cdot)}|f| \|_{L_{p_2}(B(x,r))} \lesssim \varphi_{n,\rho,p_1,p_2}(r) \| f \|_{L_{p_1}(B(x,2r))} + r^{\frac{n}{p_2}} \left(\bar{I}_{\rho(\cdot),2r}|f| \right)(x) \tag{3.18}$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in \mathfrak{M}(\mathbb{R}^n) \cap L_{p_1}(B(x, 2r))$ and, for $1 < p_2 < \infty$

$$\| I_{\rho(\cdot)}|f| \|_{WL_{p_2}(B(x,r))} \lesssim \varphi_{n,\rho,1,p_2}(r) \| f \|_{L_1(B(x,2r))} + r^{\frac{n}{p_2}} \left(\bar{I}_{\rho(\cdot),2r}|f| \right)(x) \tag{3.19}$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in \mathfrak{M}(\mathbb{R}^n) \cap L_1(B(x, 2r))$.

Lemma 3.4 *Let $f \in \mathfrak{M}(\mathbb{R}^n)$ be a non-negative function and ρ be a non-negative, non-increasing, continuously differentiable function on $(0, \infty)$ such that $\lim_{t \rightarrow +\infty} \rho(t) = 0$. Then for any $x \in \mathbb{R}^n$ and $r > 0$*

$$\int_{c_{B(x,r)}} \rho(|x-y|)f(y)dy = \int_r^\infty \left(\int_{B(x,t) \setminus B(x,r)} f(y)dy \right) |\rho'(t)|dt. \tag{3.20}$$

For $\rho(t) = t^{-\beta}, t > 0, \beta > 0$ this lemma was proved in [3] (Lemma 3).

Proof By applying the Fubini theorem we get

$$\begin{aligned} \int_{c_{B(x,r)}} \rho(|x-y|)f(y)dy &= \int_{y:|x-y|>r} \rho(|x-y|)f(y)dy \\ &= \int_{y:|x-y|>r} \left(\int_{|x-y|}^{\infty} |\rho'(t)|dt \right) f(y)dy \\ &= \int_r^{\infty} \left(\int_{y:|x-y|>r,|x-y|<t} |\rho'(t)|f(y)dy \right) dt \\ &= \int_r^{\infty} \left(\int_{B(x,t)\setminus B(x,r)} f(y)dy \right) |\rho'(t)|dt. \end{aligned}$$

Definition 3.3 Let $n \in \mathbb{N}, 1 \leq p_1 < \infty, 0 < p_2 \leq \infty$. Then $\rho \in S_{n,p_1,p_2}$ if $\rho \in \mathfrak{M}^+((0, \infty))$ and there exists a positive non-increasing continuously differentiable function $\tilde{\rho} : (0, \infty) \rightarrow (0, \infty)$ such that

- 1) $\tilde{\rho}(t) \approx \rho(t)$ uniformly in $t > 0$,
- 2) $\lim_{t \rightarrow \infty} \tilde{\rho}(t) = 0$,
- 3) $\tilde{\rho} \in S_n$,
- 4) $\int_0^{\infty} \tilde{\rho}(t)t^{\frac{n}{p_1}-1} dt = \infty, \int_1^{\infty} \tilde{\rho}(t)t^{\frac{n}{p_1}-1} dt < \infty$,
- 5) $|\tilde{\rho}'(t)|t \gtrsim \tilde{\rho}(t)$ uniformly in $t > 0$,
- 6) if $0 < p_2 \leq p_1$ and $1 \leq p_1 < \infty$, then

$$\int_0^r \tilde{\rho}(t)t^{n-1} dt \lesssim \tilde{\rho}(r)r^n$$

uniformly in $r > 0$,

- 7) if $1 \leq p_1 < p_2 < \infty$, then the function $\phi_{n,\tilde{\rho},p_1,p_2}(t) = \tilde{\rho}(t)t^{n(\frac{1}{p_1} + \frac{1}{p_2})}$ is almost non-decreasing on $(0, \infty)$.

Moreover, if $p_1 = 1$ and $1 < p_2 < \infty$, then $\rho \in S_{n,1,p_2}$ if there exists a positive non-increasing continuously differentiable function $\tilde{\rho} : (0, \infty) \rightarrow (0, \infty)$ such that Conditions 1)–3) and 5) are satisfied, Conditions 4) and 6) are satisfied for $p_1 = 1$ and instead of Condition 7) the following condition is satisfied

- 8)

$$\|\tilde{\rho}(t)t^{\frac{n-1}{p_2}}\|_{L_{p_2}(0,r)} \lesssim \tilde{\rho}(r)r^{\frac{n}{p_2}}$$

uniformly in $r > 0$.

Remark 3.7 For the Riesz potential $\rho(t) = t^{\alpha-n}, 0 < \alpha < n, \tilde{\rho} = \rho$. Condition 2) is satisfied because $\alpha < n$, Condition 3) is satisfied because $\alpha > 0$, Condition 4) is satisfied if $p_1 < \infty$ and $\alpha < \frac{n}{p_1}$, Condition 5) is obviously satisfied, Condition 6) is satisfied because $\alpha > 0$, Condition 7) is satisfied if $\alpha \geq n(1/p_1 - 1/p_2)$, Condition 8) is satisfied if $\alpha > n(1 - 1/p_2)$.

Remark 3.8 Clearly, due to Condition 1), Conditions 2)–4), 6)–8) are equivalent to the conditions, obtained from them by replacing $\tilde{\rho}$ by ρ

Theorem 3.1 Let $n \in \mathbb{N}, 1 \leq p_1 < \infty, 0 < p_2 \leq \infty$.

- 1. If $1 < p_1 < p_2 < \infty$ or $1 \leq p_1 < \infty$ and $0 < p_2 \leq p_1$, and $\rho \in S_{n,p_1,p_2}$, then

$$\|I_{\rho(\cdot)}(|f|)\|_{L_{p_2}(B(x,r))} \lesssim r^{\frac{n}{p_2}} \int_r^{\infty} \rho(t)t^{\frac{n}{p_1}-1} \|f\|_{L_{p_1}(B(x,t))} dt \tag{3.21}$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in L_{p_1}^{\text{loc}}(\mathbb{R}^n)$.

2. If $p_1 = 1$ and $0 < p_2 < \infty$, and $\rho \in \tilde{S}_{n,1,p_2}$, then

$$\| I_{\rho(\cdot)}|f| \|_{WL_{p_2}(B(x,r))} \approx \| I_{\rho(\cdot)}|f| \|_{L_{p_2}(B(x,r))} \approx r^{\frac{n}{p_2}} \int_r^\infty \rho(t)t^{-1} \| f \|_{L_1(B(x,t))} dt \tag{3.22}$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in L_1^{\text{loc}}(\mathbb{R}^n)$.

3. If $p_1 = 1$ and $1 < p_2 < \infty$, and $\rho \in S_{n,1,p_2}$, then

$$\| I_{\rho(\cdot)}|f| \|_{WL_{p_2}(B(x,r))} \approx r^{\frac{n}{p_2}} \int_r^\infty \rho(t)t^{-1} \| f \|_{L_1(B(x,t))} dt \tag{3.23}$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in L_1^{\text{loc}}(\mathbb{R}^n)$.

For $\rho(t) = t^{\alpha-n}, 0 < \alpha < n$, hence for the Riesz potential I_α , this theorem is proved in [6] (Lemma 3.6 and Theorem 3.9).

Remark 3.9 If $p_1 = 1$ and $0 < p_2 < \infty$ and $\rho \in \tilde{S}_{n,1,p_2}$, equivalence (3.22) holds if Condition 6) of Definition 3.3 for $0 < p_2 \leq 1$ and Condition 7) of Definition 3.3 for $1 < p_2 < \infty$ are satisfied. If $\rho(t) = t^{\alpha-n}, 0 < \alpha < n$, then this means that $n(1 - \frac{1}{p_2})_+ < \alpha < n$.

If $p_1 = 1$ and $1 < p_2 < \infty$ and $\rho \in S_{n,1,p_2}$, equivalence (3.23) holds if Condition 8) of Definition 3.3 is satisfied. If $\rho(t) = t^{\alpha-n}, 0 < \alpha < n$, then this means that $n(1 - \frac{1}{p_2}) \leq \alpha < n$.

In particular, for $\alpha = n(1 - 1/p_2)$ Statement 3 of Theorem 3.1 holds, while, as proved in Lemma 3.3 of [6], Statement 2 does not hold.

Proof Step 1 (proof of (3.21)). We apply equivalence (3.2).

1.1a. To the first summand in the right-hand side of (3.2) we apply Condition 2) of Definition 3.1 with ρ replaced by $\tilde{\rho}, s$ by r, t by $2r$ and inequality (3.10) and get that

$$\begin{aligned} & r^{\frac{n}{p_2}} \int_r^\infty \rho(t)t^{\frac{n}{p_1}-1} \| f \|_{L_{p_1}(B(x,t))} dt \\ & \gtrsim r^{\frac{n}{p_2}} \int_r^\infty \tilde{\rho}(t)t^{\frac{n}{p_1}-1} \| f \|_{L_{p_1}(B(x,t))} dt \\ & \geq r^{\frac{n}{p_2}} \int_r^{2r} \tilde{\rho}(t)t^{\frac{n}{p_1}-1} \| f \|_{L_{p_1}(B(x,t))} dt \\ & \geq r^{\frac{n}{p_2}} \tilde{\rho}(2r) \left(\int_r^{2r} t^{\frac{n}{p_1}-1} dt \right) \| f \|_{L_{p_1}(B(x,r))} \\ & \gtrsim \tilde{\rho}(r)r^{\frac{n}{p_1} + \frac{n}{p_2}} \| f \|_{L_{p_1}(B(x,r))} = \varphi_{n,\tilde{\rho},p_1,p_2}(r) \| f \|_{L_{p_1}(B(x,r))} \\ & \gtrsim \| I_{\tilde{\rho}(\cdot)}(|f|\chi_{B(x,2r)}) \|_{L_{p_2}(B(x,r))} \gtrsim \| I_{\rho(\cdot)}(|f|\chi_{B(x,2r)}) \|_{L_{p_2}(B(x,r))} \end{aligned} \tag{3.24}$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in L_1^{\text{loc}}(\mathbb{R}^n)$.

1.1b (second proof of inequality (3.24)). To the first summand in the right-hand side of (3.2) we apply inequality (3.10) and Condition 2) of Definition 3.1 with ρ replaced by $\tilde{\rho}$, firstly, s replaced by r and t by $2r$, secondly, s replaced by $\frac{r}{2}$ and t by r , and we get that

$$\| I_{\rho(\cdot)}(|f|\chi_{B(x,2r)}) \|_{L_{p_2}(B(x,r))} \lesssim \| I_{\tilde{\rho}(\cdot)}(|f|\chi_{B(x,2r)}) \|_{L_{p_2}(B(x,r))}$$

$$\begin{aligned} &\lesssim \tilde{\rho}(r)r^{n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)} \|f\|_{L_{p_1}(B(x,2r))} \lesssim \tilde{\rho}(2r)r^{n\left(\frac{1}{p_1}+\frac{1}{p_2}\right)} \|f\|_{L_{p_1}(B(x,2r))} \\ &\lesssim r^{\frac{n}{p_2}} \left(\int_{2r}^{4r} \tilde{\rho}(t)t^{\frac{n}{p_1}-1} dt \right) \|f\|_{L_{p_1}(B(x,2r))} \lesssim r^{\frac{n}{p_2}} \int_{2r}^{\infty} \tilde{\rho}(t)t^{\frac{n}{p_1}-1} \|f\|_{L_{p_1}(B(x,2t))} dt \\ &= 2^{-\frac{n}{p_1}} r^{\frac{n}{p_2}} \int_r^{\infty} \tilde{\rho}\left(\frac{\tau}{2}\right) \tau^{\frac{n}{p_1}-1} \|f\|_{L_{p_1}(B(x,\tau))} d\tau \lesssim r^{\frac{n}{p_2}} \int_r^{\infty} \tilde{\rho}(\tau)^{\frac{n}{p_1}-1} \|f\|_{L_{p_1}(B(x,\tau))} d\tau \\ &\lesssim r^{\frac{n}{p_2}} \int_r^{\infty} \rho(\tau)^{\frac{n}{p_1}-1} \|f\|_{L_{p_1}(B(x,\tau))} d\tau \end{aligned}$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in L_{p_1}^{loc}(\mathbb{R}^n)$.

1.2a. In order to estimate the second summand in the right-hand side of (3.2) we obtain an estimate for $(\bar{I}_{\rho(\cdot),r}|f|)(x)$. First we note that

$$\begin{aligned} &\int_{2r}^{\infty} \tilde{\rho}(t)t^{-1} \|f\|_{L_1(B(x,t))} dt = \sum_{k=1}^{\infty} \int_{2^k r}^{2^{k+1}r} \tilde{\rho}(t)t^{-1} \|f\|_{L_1(B(x,t))} dt \\ &\geq \sum_{k=1}^{\infty} \tilde{\rho}(2^k r) \|f\|_{L_1(B(x,2^k r))} \int_{2^k r}^{2^{k+1}r} \frac{dt}{t} = \ln 2 \sum_{k=1}^{\infty} \tilde{\rho}(2^k r) \|f\|_{L_1(B(x,2^k r))} \\ &\geq \ln 2 \sum_{k=1}^{\infty} \tilde{\rho}(2^k r) \|f\|_{L_1(B(x,2^k r) \setminus B(x,2^{k-1}r))} \end{aligned}$$

Since by Condition 2) of Definition 3.1 $\tilde{\rho}(2^k r) \geq c_1 \tilde{\rho}(2^{k-1} r)$ and $|x - y| \geq 2^{k-1} r$ for all $y \in B(x, 2^k r) \setminus B(x, 2^{k-1} r)$, it follows that

$$\begin{aligned} &\int_{2r}^{\infty} \tilde{\rho}(t)t^{-1} \|f\|_{L_1(B(x,t))} dt \gtrsim \sum_{k=1}^{\infty} \int_{B(x,2^k r) \setminus B(x,2^{k-1} r)} \tilde{\rho}(2^{k-1} r)|f(y)| dy \\ &\geq \sum_{k=1}^{\infty} \int_{B(x,2^k r) \setminus B(x,2^{k-1} r)} \tilde{\rho}(|x - y|)|f(y)| dy = \int_{\mathbb{R}^n} \tilde{\rho}(|x - y|)|f(y)| dy = (\bar{I}_{\tilde{\rho}(\cdot),r}|f|)(x). \end{aligned}$$

So,

$$(\bar{I}_{\tilde{\rho}(\cdot),r}|f|)(x) \lesssim \int_{2r}^{\infty} \tilde{\rho}(t)t^{-1} \|f\|_{L_1(B(x,t))} dt \tag{3.25}$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in L_1^{loc}(\mathbb{R}^n)$.

Hence, by applying Hölder’s inequality, we get that

$$\begin{aligned} &r^{\frac{n}{p_2}} \left(\bar{I}_{\rho(\cdot),2r}|f| \right)(x) \lesssim r^{\frac{n}{p_2}} \left(\bar{I}_{\tilde{\rho}(\cdot),2r}|f| \right)(x) \lesssim r^{\frac{n}{p_2}} \int_{4r}^{\infty} \tilde{\rho}(t)t^{-1} \|f\|_{L_1(B(x,t))} dt \\ &\lesssim r^{\frac{n}{p_2}} \int_{4r}^{\infty} \tilde{\rho}(t)t^{\frac{n}{p_1}-1} \|f\|_{L_{p_1}(B(x,t))} dt \lesssim r^{\frac{n}{p_2}} \int_r^{\infty} \rho(t)t^{\frac{n}{p_1}-1} \|f\|_{L_{p_1}(B(x,t))} dt \end{aligned} \tag{3.26}$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in L_1^{loc}(\mathbb{R}^n)$.

Step 2 (proof of equivalences (3.22) and (3.23) for $p_1 = 1$). We apply equivalences (3.2) and (3.3).

2.1a. Under the assumption $\tilde{\rho} \in \tilde{S}_{n,1,p_2}$ inequality (3.12) holds for $p_1 = 1$ and $1 \leq p_2 \leq \infty$ and inequality (3.13) holds for $p_1 = 1$ and $0 < p_2 \leq 1$, therefore, Lemma 3.3 ensures the validity of inequality (3.10) with $p_1 = 1$ and we have

$$\begin{aligned} r^{\frac{n}{p_2}} \int_r^\infty \rho(t)t^{-1} \|f\|_{L_1(B(x,t))} dt &\gtrsim r^{\frac{n}{p_2}} \int_r^\infty \tilde{\rho}(t)t^{-1} \|f\|_{L_1(B(x,t))} dt \\ &\gtrsim \varphi_{n,\tilde{\rho},1,p_2}(r) \|f\|_{L_1(B(x,r))} \\ &\gtrsim \|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{L_{p_2}(B(x,r))} \geq \|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{WL_{p_2}(B(x,r))} \end{aligned} \tag{3.27}$$

uniformly in $x \in \mathbb{R}^n$, $r > 0$ and $f \in L_1^{\text{loc}}(\mathbb{R}^n)$ and we get the estimate above for the first summand in (3.22).

2.1b. If $\rho \in S_{n,1,p_2}$, then condition (3.9) holds for $p_1 = 1$ and $1 < p_2 < \infty$, therefore, Lemma 3.3 ensures the validity of inequality (3.11) and we, respectively, have

$$\begin{aligned} r^{\frac{n}{p_2}} \int_r^\infty \rho(t)t^{-1} \|f\|_{L_1(B(x,t))} dt \\ \gtrsim r^{\frac{n}{p_2}} \int_r^\infty \tilde{\rho}(t)t^{-1} \|f\|_{L_1(B(x,t))} dt \geq \varphi_{n,\tilde{\rho},1,p_2}(r) \|f\|_{L_1(B(x,r))} \\ \gtrsim \|I_{\rho(\cdot)}(|f|\chi_{B(x,2r)})\|_{WL_{p_2}(B(x,r))} \end{aligned} \tag{3.28}$$

uniformly in $x \in \mathbb{R}^n$, $r > 0$ and $f \in L_1^{\text{loc}}(\mathbb{R}^n)$ and we get the estimate above for the first summand in (3.23).

2.2a. If $\tilde{\rho} \in \tilde{S}_{n,1,p_2}$ and $1 \leq p_2 \leq \infty$, to the second summand in the right-hand side of (3.2) we apply inequality (3.25). Equivalence (3.2) and inequalities (3.25) and (3.27) imply that

$$\|I_{\rho(\cdot)}|f|\|_{WL_{p_2}(B(x,r))} \leq \|I_{\rho(\cdot)}|f|\|_{L_{p_2}(B(x,r))} \lesssim r^{\frac{n}{p_2}} \int_r^\infty \rho(t)t^{-1} \|f\|_{L_1(B(x,t))} dt \tag{3.29}$$

uniformly in $x \in \mathbb{R}^n$, $r > 0$ and $f \in L_1^{\text{loc}}(\mathbb{R}^n)$.

2.2b. Accordingly, if $\tilde{\rho} \in S_{n,1,p_2}$, $p_1 = 1$ and $1 < p_2 < \infty$, then equivalence (3.3) and inequalities (3.25) and (3.28) imply that

$$\|I_{\rho(\cdot)}|f|\|_{WL_{p_2}(B(x,r))} \lesssim r^{\frac{n}{p_2}} \int_r^\infty \rho(t)t^{-1} \|f\|_{L_1(B(x,t))} dt \tag{3.30}$$

uniformly in $x \in \mathbb{R}^n$, $r > 0$ and $f \in L_1^{\text{loc}}(\mathbb{R}^n)$.

2.3. For all $y \in B(x, r)$ we have $|y - z| \leq 2r$ if $z \in B(x, r)$ and $|y - z| \leq 2|x - z|$ if $z \in {}^c B(x, r)$ (since $|y - z| \leq |y - x| + |x - z| \leq r + |x - z| \leq 2|x - z|$). Therefore, since $\tilde{\rho}$ is non-increasing

$$\begin{aligned} I_{\tilde{\rho}(\cdot)}(|f|)(y) &= \int_{B(x,r)} \tilde{\rho}(|y - z|)|f(z)|dz + \int_{{}^c B(x,r)} \tilde{\rho}(|y - z|)|f(z)|dz \\ &\geq \tilde{\rho}(2r) \int_{B(x,r)} |f(z)|dz + \int_{{}^c B(x,r)} \tilde{\rho}(2|x - z|)|f(z)|dz. \end{aligned}$$

By Condition 2) of Definition 3.1 with $s = 2t$ we have $\tilde{\rho}(2t) \geq c_1 \tilde{\rho}(z)$ for any $t > 0$, hence, by equality (3.18) uniformly in $x \in \mathbb{R}^n$, $r > 0$, $y \in B(x, r)$ and $f \in L_1^{\text{loc}}(\mathbb{R}^n)$

$$I_{\rho(\cdot)}(|f|)(y) \gtrsim I_{\tilde{\rho}(\cdot)}(|f|)(y) \geq c_1 \left(\tilde{\rho}(r) \int_{B(x,r)} |f(z)|dz + \int_{{}^c B(x,r)} \tilde{\rho}(|x - z|)|f(z)|dz \right)$$

$$\begin{aligned}
 &= c_1 \left(\tilde{\rho}(r) \int_{B(x,r)} |f(z)| dz + \int_r^\infty \left(\int_{B(x,t) \setminus B(x,r)} |f(z)| dz \right) |(\tilde{\rho})'(t)| dt \right) \\
 &= c_1 \left(\tilde{\rho}(r) \int_{B(x,r)} |f(z)| dz + \int_r^\infty \left(\int_{B(x,t)} |f(z)| dz \right) |(\tilde{\rho})'(t)| dt - \left(\int_r^\infty |(\tilde{\rho})'(t)| dt \right) \int_{B(x,r)} |f(z)| dz \right) \\
 &= c_1 \int_r^\infty \left(\int_{B(x,t)} |f(z)| dz \right) |(\tilde{\rho})'(t)| dt.
 \end{aligned}$$

Consequently, by Remark 3.1 and by Condition 5) of Definition 3.3

$$\begin{aligned}
 \| I_{\rho(\cdot)} |f| \|_{L_{p_2}(B(x,r))} &\geq \| I_{\rho(\cdot)} |f| \|_{WL_{p_2}(B(x,r))} \gtrsim \| I_{\tilde{\rho}} |f| \|_{WL_{p_2}(B(x,r))} \\
 &\geq c_1 v_n^{\frac{n}{p_2}} r^{\frac{n}{p_2}} \int_r^\infty \left(\int_{B(x,t)} |f(z)| dz \right) |(\tilde{\rho})'(t)| dt \\
 &\gtrsim r^{\frac{n}{p_2}} \int_r^\infty \tilde{\rho}(t) t^{-1} \| f \|_{L_1(B(x,t))} dt \gtrsim r^{\frac{n}{p_2}} \int_r^\infty \rho(t) t^{-1} \| f \|_{L_1(B(x,t))} dt
 \end{aligned} \tag{3.31}$$

uniformly in $x \in \mathbb{R}^n, r > 0$ and $f \in L_1^{\text{loc}}(\mathbb{R}^n)$.

Step 3. Statement 1 of Theorem 3.1 follows by inequalities (3.24) and (3.26). Statement 2 follows by inequalities (3.25), (3.27) and (3.31). Statement 3 follows by inequalities (3.25), (3.28) and (3.31).

Generalized Riesz potential and Hardy operator

Let $\mathfrak{M}^+((0, \infty))$ be the subset of $\mathfrak{M}((0, \infty))$ consisting of all non-negative functions on $(0, \infty)$. We denote by $\mathfrak{M}^+((0, \infty); \downarrow)$ the cone of all functions in $\mathfrak{M}^+((0, \infty))$ which are non-increasing on $(0, \infty)$ and we set

$$\mathbb{A} = \left\{ \varphi \in \mathfrak{M}^+((0, \infty); \downarrow) : \lim_{t \rightarrow \infty} \varphi(t) = 0 \right\}.$$

Let H be the Hardy operator

$$(Hg)(t) := \int_0^t g(r) dr, \quad 0 < t < \infty$$

Moreover, let, for $0 < p \leq \infty$ and a Lebesgue measurable function $v : (0, \infty) \rightarrow [0, \infty)$, $L_{p, v(\cdot)}(0, \infty)$ denote the space of all Lebesgue measurable functions $f : (0, \infty) \rightarrow \mathbb{C}$ for which

$$\|f\|_{L_{p, v(\cdot)}(0, \infty)} = \|fv\|_{L_p(0, \infty)} < \infty.$$

Theorem 4.1 *Let $n \in \mathbb{N}, 1 \leq p_1 < \infty, 0 < p_2 \leq \infty, 0 < \theta_2 \leq \infty, w_2 \in \Omega_{\theta_2}, \rho$ be a positive continuous function on $(0, \infty)$ and*

$$\mu_{n, \rho, p_1}(r) = \frac{\int_r^\infty \rho(t) t^{\frac{n}{p_1} - 1} dt}{\int_1^\infty \rho(t) t^{\frac{n}{p_1} - 1} dt}, \quad r > 0, \tag{4.1}$$

$$v_2(r) = w_2 \left(\mu_{n,\rho,p_1}^{(-1)}(r) \right) \left(\mu_{n,\rho,p_1}^{(-1)}(r) \right)^{\frac{n}{p_2}} \left| \left(\mu_{n,\rho,p_1}^{(-1)}(r) \right)' \right|^{\frac{1}{\theta_2}}, \quad r > 0, \tag{4.2}$$

$$g_{n,\rho,p_1}(t) = \|f\|_{L_{p_1}(B(0,\mu_{n,\rho,p_1}^{(-1)}(t)))}, \quad t > 0. \tag{4.3}$$

1. If $1 < p_1 < p_2 < \infty$ or $1 \leq p_1 < \infty$ and $0 < p_2 \leq p_1$, and $\rho \in S_{n,p_1,p_2}$, then

$$\|I_{\rho(\cdot)}f\|_{LM_{p_2\theta_2,w_2(\cdot)}} \lesssim \|Hg_{n,\rho,p_1}\|_{L_{\theta_2,v_2(\cdot)}(0,\infty)} \tag{4.4}$$

uniformly in $f \in \mathfrak{M}(\mathbb{R}^n)$.

2. If $p_1 = 1$ and $0 < p_2 < \infty$, and $\rho \in \tilde{S}_{n,1,p_2}$, then

$$\|I_{\rho(\cdot)}f\|_{LM_{p_2\theta_2,w_2(\cdot)}} \approx \|Hg_{n,\rho,1}\|_{L_{\theta_2,v_2(\cdot)}(0,\infty)} \tag{4.5}$$

uniformly in all non-negative functions $f \in \mathfrak{M}(\mathbb{R}^n)$.

3. If $p_1 = 1$ and $1 < p_2 < \infty$, and $\rho \in S_{n,1,p_2}$, then

$$\|I_{\rho(\cdot)}f\|_{WLM_{p_2\theta_2,w_2(\cdot)}} \approx \|Hg_{n,\rho,1}\|_{L_{\theta_2,v_2(\cdot)}(0,\infty)} \tag{4.6}$$

uniformly in all non-negative functions $f \in \mathfrak{M}(\mathbb{R}^n)$.

Remark 4.1 If $\rho(t) = t^{\alpha-n}$, $1 \leq p_1 < \infty$, $0 < p_2 < \infty$, $0 < \alpha < \frac{n}{p_1}$, then $\mu_{n,\rho,p_1}(r) = r^{-\sigma}$, where $\sigma = \frac{n}{p_1} - \alpha$, $\mu_{n,\rho,p_1}^{(-1)}(r) = r^{-\frac{1}{\sigma}}$, $v_2(r) = \sigma^{-\frac{1}{\theta_2}} w_2(r^{-\frac{1}{\sigma}}) r^{-\frac{n}{\sigma p_2} - \frac{1}{\sigma \theta_2} - \frac{1}{\theta_2}}$, $g_{n,\rho,p_1}(t) = \|f\|_{L_{p_1}(B(0,t^{-\frac{1}{\sigma}}))}$ and Lemma 4.1 takes the form of Lemma 4.1 in [6].

Remark 4.2 Due to Condition 4) of Definition 3.3

$$\lim_{r \rightarrow 0^+} \mu_{n,\rho,p_1}(r) = \lim_{r \rightarrow 0^+} \mu_{n,\tilde{\rho},p_1}(r) = \infty, \quad \lim_{r \rightarrow \infty} \mu_{n,\rho,p_1}(r) = \lim_{r \rightarrow \infty} \mu_{n,\tilde{\rho},p_1}(r) = 0 \tag{4.7}$$

and μ_{n,ρ,p_1} is a strictly decreasing continuously differentiable function on $(0, \infty)$. Moreover,

$$\lim_{r \rightarrow 0^+} \mu_{n,\rho,p_1}^{(-1)}(r) = \infty, \quad \lim_{r \rightarrow \infty} \mu_{n,\rho,p_1}^{(-1)}(r) = 0. \tag{4.8}$$

Proof 1. Let $1 < p_1 < p_2 < \infty$ or $1 \leq p_1 \leq \infty$ and $0 < p_2 \leq p_1$, and $\rho \in S_{n,p_1,p_2}$. By inequality (3.21) we have

$$\begin{aligned} \left(c = \left(\int_1^\infty \rho(t) t^{\frac{n}{p_1}-1} dt \right)^{-1} \right) \\ \|I_{\rho(\cdot)}f\|_{LM_{p_2\theta_2,w_2(\cdot)}} &\lesssim \left\| w_2(r) r^{\frac{n}{p_2}} \int_r^\infty \rho(t) t^{\frac{n}{p_1}-1} \|f\|_{L_{p_1}(B(0,t))} dt \right\|_{L_{\theta_2}(0,\infty)} \\ &= c \left\| w_2(r) r^{\frac{n}{p_2}} \left(- \int_r^\infty \|f\|_{L_{p_1}(B(0,t))} d\mu_{n,\rho,p_1}(t) \right) \right\|_{L_{\theta_2}(0,\infty)} \\ &= c \left(\mu_{n,\rho,p_1}(t) = \tau, \quad t = \mu_{n,\rho,p_1}^{(-1)}(\tau), \quad \lim_{r \rightarrow \infty} \mu_{n,\rho,p_1}(t) = 0 \right) \\ &= c \left\| w_2(r) r^{\frac{n}{p_2}} \int_0^{\mu_{n,\rho,p_1}^{(-1)}(r)} \|f\|_{L_{p_1}(B(0,\mu_{n,\rho,p_1}^{(-1)}(\tau)))} d\tau \right\|_{L_{\theta_2}(0,\infty)} \end{aligned}$$

$$\begin{aligned}
 &= c \left(\mu_{n,\rho,p_1}(r) = u, r = \mu_{n,\rho,p_1}^{(-1)}(u), \lim_{r \rightarrow 0} \mu_{n,\rho,p_1}(r) = \infty, \lim_{r \rightarrow \infty} \mu_{n,\rho,p_1}(r) = 0 \right) \\
 &= c \left\| w_2 \left(\mu_{n,\rho,p_1}^{(-1)}(u) \right) \left(\mu_{n,\rho,p_1}^{(-1)}(u) \right)^{\frac{n}{p_2}} \left| \left(\mu_{n,\rho,p_1}^{(-1)}(u) \right)' \right|^{\frac{1}{\theta_2}} \int_0^u \|f\|_{L_{p_1}(B(0, \mu_{n,\rho,p_1}^{(-1)}(\tau)))} d\tau \right\|_{L_{\theta_2}(0, \infty)} \\
 &= c \left\| v_2(u) (Hg_{n,\rho,p_1})(u) \right\|_{L_{\theta_2}(0, \infty)} = c \|Hg_{n,\rho,p_1}\|_{L_{\theta_2, v_2(\cdot)}(0, \infty)}
 \end{aligned}$$

uniformly in $f \in \mathfrak{M}(\mathbb{R}^n)$.

2. Let $p_1 = 1$ and $0 < p_2 \leq 1$, and $\rho \in \tilde{S}_{n,p_1,p_2}$. By inequality (3.22) we have similarly to Step 1

$$\begin{aligned}
 \left\| I_{\rho(\cdot)} f \right\|_{LM_{p_2\theta_2, w_2(\cdot)}} &\approx \left\| w_2(r) r^{\frac{n}{p_2}} \int_r^\infty \rho(t) t^{-1} \|f\|_{L_1(B(0,t))} dt \right\|_{L_{\theta_2}(0, \infty)} \\
 &= c \left\| w_2(r) r^{\frac{n}{p_2}} \left(- \int_r^\infty \|f\|_{L_1(B(0,t))} d\mu_{n,\rho,1}(t) \right) \right\|_{L_{\theta_2}(0, \infty)} \\
 &= c \left\| v_2(u) (Hg_{n,1})(u) \right\|_{L_{\theta_2}(0, \infty)} = c \|Hg_{n,\rho,1}\|_{L_{\theta_2, v_2(\cdot)}(0, \infty)}
 \end{aligned}$$

uniformly in all non-negative functions $f \in \mathfrak{M}(\mathbb{R}^n)$.

3. Let $p_1 = 1$ and $1 < p_2 < \infty$, and $\rho \in S_{n,1,p_2}$. Equivalence (4.6) follows similarly to Step 2 from equivalence (3.23). □

Theorem 4.2 Assume that $n \in \mathbb{N}, 1 \leq p_1 < \infty, 0 < p_2 \leq \infty, 0 < \theta_1, \theta_2 \leq \infty, w_1 \in \Omega_{\theta_1}, w_2 \in \Omega_{\theta_2}, \mu_{n,\rho,p_1}$ is defined by formula (4.1),

$$v_1(r) = w_1 \left(\mu_{n,\rho,p_1}^{(-1)}(r) \right) \left| \left(\mu_{n,\rho,p_1}^{(-1)}(r) \right)' \right|^{\frac{1}{\theta_1}}, \quad r > 0, \tag{4.9}$$

and v_2 is defined by formula (4.2).

1. Let $1 < p_1 < p_2 < \infty$ or $1 \leq p_1 < \infty$ and $0 < p_2 \leq p_1$, and $\rho \in S_{n,p_1,p_2}$. If the operator H is bounded from $L_{\theta_1, v_1(\cdot)}(0, \infty)$ to $L_{\theta_2, v_2(\cdot)}(0, \infty)$ on the cone \mathbb{A} , that is

$$\|Hg\|_{L_{\theta_2, v_2(\cdot)}(0, \infty)} \lesssim \|g\|_{L_{\theta_1, v_1(\cdot)}(0, \infty)} \tag{4.10}$$

uniformly in $g \in \mathbb{A}$, then the operator $I_{\rho(\cdot)}$ is bounded from $LM_{p_1\theta_1, w_1(\cdot)}$ to $LM_{p_2\theta_2, w_2(\cdot)}$.

2. Let $p_1 = 1, 0 < p_2 < \infty$ and $\rho \in \tilde{S}_{n,1,p_2}$. Then the operator $I_{\rho(\cdot)}$ is bounded from $LM_{1\theta_1, w_1(\cdot)}$ to $LM_{p_2\theta_2, w_2(\cdot)}$ if and only if the operator H is bounded from $L_{\theta_1, v_1(\cdot)}(0, \infty)$ to $L_{\theta_2, v_2(\cdot)}(0, \infty)$ on the cone \mathbb{A} .

3. Let $p_1 = 1, 1 < p_2 < \infty$ and $\rho \in S_{n,1,p_2}$. Then the operator $I_{\rho(\cdot)}$ is bounded from $LM_{p_1\theta_1, w_1(\cdot)}$ to $WLM_{p_2\theta_2, w_2(\cdot)}$ if and only if the operator H is bounded from $L_{\theta_1, v_1(\cdot)}(0, \infty)$ to $L_{\theta_2, v_2(\cdot)}(0, \infty)$ on the cone \mathbb{A} .

Remark 4.3 If we put $r = \mu_{n_1,\rho,p_1}(t)$ in (4.9), then, taking into account that $\left(\mu_{n_1,\rho,p_1}^{(-1)}(r) \right)' = \left(\mu'_{n_1,\rho,p_1} \left(\mu_{n_1,\rho,p_1}^{(-1)}(r) \right) \right)^{-1}, r > 0$, we get

$$v_1(\mu_{n,\rho,p_1}(t)) = w_1(t) \left| \mu'_{n,\rho,p_1}(t) \right|^{-\frac{1}{\theta_1}}, \quad t > 0 \tag{4.11}$$

and

$$v_2(\mu_{n,\rho,p_1}(t)) = w_2(t) t^{\frac{n}{p_2}} \left| \mu'_{n,\rho,p_1}(t) \right|^{-\frac{1}{\theta_1}}, \quad t > 0, \tag{4.12}$$

Lemma 4.1 Assume that $0 < p_1, p_2, \theta_1, \theta_2 \leq \infty$, $w_1 \in \Omega_{\theta_1}$, ρ is a positive continuous function on $(0, \infty)$ such that

$$\int_0^1 \rho(t) t^{\frac{n}{p_1}-1} dt = \infty, \quad \int_1^\infty \rho(t) t^{\frac{n}{p_1}-1} dt < \infty, \tag{4.13}$$

and the functions μ_{n,ρ,p_1} , g_{n,ρ,p_1} , v_1 , v_2 are defined by formulas (4.1), (4.3), (4.9), (4.2), respectively. Then

$$\|g_{n,\rho,p_1}\|_{L_{\theta_1,v_1(\cdot)}} = \|f\|_{LM_{p_1,\theta_1,w_1(\cdot)}} \tag{4.14}$$

for all $f \in LM_{p_1,\theta_1,w_1(\cdot)}$, for any measurable function $\phi_1 : (0, \infty) \rightarrow (0, \infty)$ and $t > 0$

$$\|v_1 \phi_1\|_{L_{\theta_1}(0,t)} = \|w_1(r) \phi_1(\mu_{n,\rho,p_1}(r))\|_{L_{\theta_1}(\mu_{n,\rho,p_1}^{(-1)}(t), \infty)}, \tag{4.15}$$

$$\|v_1 \phi_1\|_{L_{\theta_1}(t, \infty)} = \|w_1(r) \phi_1(\mu_{n,\rho,p_1}(r))\|_{L_{\theta_1}(0, \mu_{n,\rho,p_1}^{(-1)}(t))}, \tag{4.16}$$

$$\|v_1 \phi_1\|_{L_{\theta_1}(0, \infty)} = \|w_1(r) \phi_1(\mu_{n,\rho,p_1}(r))\|_{L_{\theta_1}(0, \infty)}, \tag{4.16'}$$

$$\|v_1\|_{L_{\theta_1}(0, \infty)} = \|w_1\|_{L_{\theta_1}(0, \infty)}, \tag{4.17}$$

and for any measurable function $\phi_2 : (0, \infty) \rightarrow (0, \infty)$ and $t > 0$

$$\|v_2 \phi_2\|_{L_{\theta_2}(0,t)} = \|w_2(r) r^{\frac{n}{p_2}} \phi_2(\mu_{n,\rho,p_1}(r))\|_{L_{\theta_2}(\mu_{n,\rho,p_1}^{(-1)}(t), \infty)}, \tag{4.18}$$

$$\|v_2 \phi_2\|_{L_{\theta_2}(t, \infty)} = \|w_2(r) r^{\frac{n}{p_2}} \phi_2(\mu_{n,\rho,p_1}(r))\|_{L_{\theta_2}(0, \mu_{n,\rho,p_1}^{(-1)}(t))}, \tag{4.19}$$

$$\|v_2 \phi_2\|_{L_{\theta_2}(0, \infty)} = \|w_2(r) r^{\frac{n}{p_2}} \phi_2(\mu_{n,\rho,p_1}(r))\|_{L_{\theta_2}(0, \infty)}, \tag{4.19'}$$

$$\|v_2\|_{L_{\theta_2}(0, \infty)} = \left\| w_2(r) r^{\frac{n}{p_2}} \right\|_{L_{\theta_2}(0, \infty)}. \tag{4.20}$$

Proof 1. Indeed,

$$\begin{aligned} \|g_{n,\rho,p_1}\|_{L_{\theta_1,v_1(\cdot)}} &= \left\| v_1(t) \|f\|_{L_{p_1}(B(0, \mu_{n,\rho,p_1}^{(-1)}(t)))} \right\|_{L_{\theta_1}(0, \infty)} \\ &= \left(\mu_{n,\rho,p_1}^{(-1)}(t) = r, t = \mu_{n,\rho,p_1}(r), \text{ by (4.13)} \lim_{r \rightarrow 0^+} \mu_{n,\rho,p_1}^{(-1)}(t) = \infty, \lim_{n \rightarrow \infty} \mu_{n,\rho,p_1}^{(-1)}(t) = 0 \right) \\ &= \left\| v_1(\mu_{n,\rho,p_1}(r)) \left| \mu'_{n,\rho,p_1}(r) \right|^{\frac{1}{\theta_1}} \|f\|_{L_{p_1}(B(0,r))} \right\|_{L_{\theta_1}(0, \infty)} \end{aligned}$$

$$= \left\| w_1(r) \|f\|_{L_{p_1}(B(0,r))} \right\|_{L_{\theta_1}(0,\infty)} = \|f\|_{LM_{p_1,\theta_1,w_1(\cdot)}}.$$

(We have used equality (4.11)).

2. Note that, by (3.10) and (4.7), for $0 < \theta_1 < \infty, t > 0$

$$\begin{aligned} \|v_1 \phi_1\|_{L_{\theta_1}(0,t)} &= \left(- \int_0^t w_1 \left(\mu_{n,\rho,p_1}^{(-1)}(s) \right)^{\theta_1} \phi_1(s)^{\theta_1} \left(\mu_{n,\rho,p_1}^{(-1)}(s) \right)' ds \right)^{\frac{1}{\theta_1}} \\ &= \left(\mu_{n,\rho,p_1}^{(-1)}(s) = r \right) = \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(t)}^{\infty} w_1(r)^{\theta_1} \phi_1(\mu_{n,\rho,p_1}(r))^{\theta_1} dr \right)^{\frac{1}{\theta_1}} \\ &= \|w_1(r) \phi_1(\mu_{n,\rho,p_1}(r))\|_{L_{\theta_1}(\mu_{n,\rho,p_1}^{(-1)}(t),\infty)} \end{aligned}$$

and, similarly

$$\|v_1\|_{L_{\theta_1}(t,\infty)} = \|w_1(r) \phi_1(\mu_{n,\rho,p_1}(r))\|_{L_{\theta_1}(0,\mu_{n,\rho,p_1}^{(-1)}(t))}.$$

These equalities also hold for $\theta_1 = \infty$, because, for example, by (4.9)

$$\begin{aligned} \|v_1 \phi_1\|_{L_{\infty}(0,t)} &= \left\| w_1 \left(\mu_{n,\rho,p_1}^{(-1)}(s) \right) \phi_1(s) \right\|_{L_{\infty}(0,t)} \\ &= \operatorname{ess\,sup}_{0 < s < t} |w_1 \left(\mu_{n,\rho,p_1}^{(-1)}(s) \right) \phi_1(s)| \\ &= \left(\mu_{n,\rho,p_1}^{(-1)}(s) = r, \mu_{n,\rho,p_1}^{(-1)}((0,t)) = \left(\mu_{n,\rho,p_1}^{(-1)}, \infty \right) \right) \\ &= \operatorname{ess\,sup}_{\mu_{n,\rho,p_1}^{(-1)}(t) < r < \infty} |w_1(r) \phi_1(\mu_{n,\rho,p_1}(r))| = \|w_1(r) \phi_1(\mu_{n,\rho,p_1}(r))\|_{L_{\infty}(\mu_{n,\rho,p_1}^{(-1)}(t),\infty)}. \end{aligned}$$

Similarly, by (4.2), (4.7) and (4.12) for $0 < \theta_2 \leq \infty, t > 0$

$$\|v_2\|_{L_{\theta_2}(0,t)} = \left\| w_2(r) r^{\frac{n}{p_2}} \right\|_{L_{\theta_2}(\mu_{n,\rho,p_1}^{(-1)}(t),\infty)}$$

and

$$\|v_2\|_{L_{\theta_2}(t,\infty)} = \left\| w_2(r) r^{\frac{n}{p_2}} \right\|_{L_{\theta_2}(0,\mu_{n,\rho,p_1}^{(-1)}(t))}.$$

The proved equalities imply equalities (4.17) and (4.20) by passing to the limit as $t \rightarrow 0^+$ or $t \rightarrow +\infty$. □

Proof of Theorem 4.2 1. Let $1 < p_1 < p_2 < \infty$ or $1 \leq p_1 < \infty$ and $0 < p_2 \leq p_1$, and $\rho \in S_{n,p_1,p_2}$. Assume that the operator H is bounded from $L_{\theta_1,v_1(\cdot)}(0, \infty)$ to $L_{\theta_2,v_2(\cdot)}(0, \infty)$ on the cone \mathbb{A} . Since $g_{n,\rho,p_1} \in \mathbb{A}$, by Statement 1 of Theorem 4.1 and formula (4.14) we have

$$\left\| I_{\rho(\cdot)} f \right\|_{LM_{p_2,\theta_2,w_2(\cdot)}} \lesssim \left\| H g_{n,\rho,p_1} \right\|_{L_{\theta_2,v_2(\cdot)}(0,\infty)} \lesssim \left\| g_{n,\rho,p_1} \right\|_{L_{\theta_1,v_1(\cdot)}(0,\infty)} = \|f\|_{LM_{p_1,\theta_1,w_1(\cdot)}} \tag{4.21}$$

uniformly in $f \in LM_{p_1,\theta_1,w_1(\cdot)}$, hence the operator $I_{\rho(\cdot)}$ is bounded from $LM_{p_1,\theta_1,w_1(\cdot)}$ to $LM_{p_2,\theta_2,w_2(\cdot)}$.

2.1. Let $p_1 = 1, 0 < p_2 < \infty$ and $\rho \in \tilde{S}_{n,1,p_2}$. If the operator H is bounded from $L_{\theta_1, v_1(\cdot)}(0, \infty)$ to $L_{\theta_2, v_2(\cdot)}(0, \infty)$ on the cone \mathbb{A} , then by Statement 2 of Theorem 4.1 as in Step 1 it follows that the operator $I_{\rho(\cdot)}$ is bounded from $LM_{p_1, \theta_1, w_1(\cdot)}$ to $LM_{p_2, \theta_2, w_2(\cdot)}$.

2.2. Assume that $I_{\rho(\cdot)}$ is bounded from $LM_{1, \theta_1, w_1(\cdot)}$ to $LM_{p_2, \theta_2, w_2(\cdot)}$. Then by Statement 2 of Theorem 4.1 and formula (4.14) with $p_1 = 1$

$$\|H_{g_{n,\rho,1}}\|_{L_{\theta_2, v_2(\cdot)}(0, \infty)} \approx \|I_{\rho(\cdot)} f\|_{LM_{p_2, \theta_2, w_2(\cdot)}} \lesssim \|f\|_{LM_{1, \theta_1, w_1(\cdot)}} = \|g_{n,\rho,1}\|_{L_{\theta_1, v_1(\cdot)}} \tag{4.22}$$

uniformly in all non-negative functions $f \in L_1^{\text{loc}}(\mathbb{R}^n)$.

Let $g \in \mathbb{A}$ be continuously differentiable on $(0, \infty)$. Consider the positive Lebesgue measurable function h on $(0, \infty)$ defined uniquely up to equivalence by the equality

$$g(t) = \|h(|\cdot|)\|_{L_1(B(0, \mu_{n,\rho,1}^{(-1)}(t)))} = \sigma_n \int_0^{\mu_{n,\rho,1}^{(-1)}(t)} h(\tau) \tau^{n-1} d\tau, \quad t > 0$$

$$\Leftrightarrow g'(t) = \sigma_n (\mu_{n,\rho,1}^{(-1)}(t))' h(\mu_{n,\rho,1}^{(-1)}(t)) (\mu_{n,\rho,1}^{(-1)}(t))^{n-1}, \quad t > 0$$

$$\Leftrightarrow h(r) = \sigma_n^{-1} g'(\mu_{n,\rho,1}(r)) \mu_{n,\rho,1}'(r) r^{1-n}, \quad r > 0.$$

If we take in (4.22) $f(x) = h(|x|)$, then by (4.3) $\|f\|_{L_1(B(0, \mu_{n,\rho,1}^{(-1)}(t)))} = g(t)$ and by (4.21)

$$\|Hg\|_{L_{\theta_2, v_2(\cdot)}(0, \infty)} \lesssim \|g\|_{L_{\theta_1, v_1(\cdot)}} \tag{4.23}$$

uniformly in all $g \in \mathbb{A}$ which are continuously differentiable on $(0, \infty)$.

Finally, if g is an arbitrary function in \mathbb{A} , then there exist functions $g_k \in \mathbb{A}, k \in \mathbb{N}$, which are continuously differentiable on $(0, \infty)$ and such $g_k \leq g_{k+1}, k \in \mathbb{N}$, and $\lim_{k \rightarrow \infty} g_k = g$ on $(0, \infty)$. Therefore, by passing to the limit in (4.23), with g replaced by g_k , as $k \rightarrow \infty$, it follows that (4.23) holds for all $g \in \mathbb{A}$.

3. Let $p_1 = 1, 1 < p_2 < \infty$ and $\rho \in S_{n,1,p_2}$. In this case the argument is similar to that of Step 2, only Statement 3 of Theorem 4.1 should be used and the space $LM_{p_2, \theta_2, w_2(\cdot)}$ should be replaced by the space $WLM_{p_2, \theta_2, w_2(\cdot)}$. \square

Hardy’s inequality on the cone of monotonic functions

Necessary and sufficient conditions for the boundedness of the operator H from one weighted Lebesgue space $L_{\theta_1, v_1(\cdot)}(0, \infty)$ to another one $L_{\theta_2, v_2(\cdot)}(0, \infty)$ on the cone \mathbb{A} are known for all values of $0 < \theta_1, \theta_2 \leq \infty$. We present below these results which are corollaries of more general statements contained in the survey by A. Gogatishvili and V.D. Stepanov [10] (Theorems 2.5, 3.15 and 3.16). See also the survey by M.L. Goldman [11].

Theorem 5.1 *Let $0 < \theta_1, \theta_2 \leq \infty, v_1, v_2$ be non-negative Lebesgue measurable functions on $(0, \infty)$. Then the operator H is bounded from $L_{\theta_1, v_1(\cdot)}(0, \infty)$ to $L_{\theta_2, v_2(\cdot)}(0, \infty)$ on the cone \mathbb{A} if and only if*

(a) if $1 < \theta_1 \leq \theta_2 < \infty$, then

$$A_{11} := \sup_{t>0} \left(\int_0^t s^{\theta_2} v_2^{\theta_2}(s) ds \right)^{\frac{1}{\theta_2}} \left(\int_0^t v_1^{\theta_1}(s) ds \right)^{-\frac{1}{\theta_1}} < \infty,$$

and

$$A_{12} := \sup_{t>0} \left(\int_t^\infty v_2^{\theta_2}(\tau) d\tau \right)^{\frac{1}{\theta_2}} \left(\int_0^t z^{\theta_1'} \left(\int_0^z v_1^{\theta_1}(s) ds \right)^{-\theta_1'} v_1^{\theta_1}(z) dz \right)^{\frac{1}{\theta_1'}} < \infty,$$

(b) if $0 < \theta_1 \leq 1, \theta_1 \leq \theta_2 < \infty$, then

$$A_2 := \sup_{t>0} \left(\int_0^t v_1^{\theta_1}(s) ds \right)^{-\frac{1}{\theta_1}} \left(\int_0^\infty \min\{s, t\}^{\theta_2} v_2^{\theta_2}(s) ds \right)^{\frac{1}{\theta_2}} < \infty,$$

(c) if $1 < \theta_1 < \infty, 0 < \theta_2 < \theta_1 < \infty$, then

$$A_{31} := \left(\int_0^\infty \left(\int_0^t v_1^{\theta_1}(\tau) d\tau \right)^{-\frac{r}{\theta_1}} \left(\int_0^t s^{\theta_2} v_2^{\theta_2}(s) ds \right)^{\frac{r}{\theta_1}} t^{\theta_2} v_2^{\theta_2}(t) dt \right)^{\frac{1}{r}} < \infty,$$

and

$$A_{32} := \left(\int_0^\infty \left(\int_t^\infty v_2^{\theta_2}(s) ds \right)^{\frac{r}{\theta_1}} \left(\int_0^t s^{\theta_1'} \left(\int_0^s v_1^{\theta_1}(\tau) d\tau \right)^{-\theta_1'} v_1^{\theta_1}(s) ds \right)^{\frac{r}{\theta_1'}} v_2^{\theta_2}(t) dt \right)^{\frac{1}{r}} < \infty,$$

where $\frac{1}{r} = \frac{1}{\theta_2} - \frac{1}{\theta_1}$,

(d) if $0 < \theta_2 < \theta_1 \leq 1$, then $A_{41} := A_{31} < \infty$ and

$$A_{42} := \left(\int_0^\infty \left(\sup_{0 < s \leq t} s^{\theta_1} \left(\int_0^s v_1^{\theta_1}(\tau) d\tau \right)^{-1} \right)^{\frac{r}{\theta_1}} \left(\int_t^\infty v_2^{\theta_2}(z) dz \right)^{\frac{r}{\theta_1}} v_2^{\theta_2}(t) dt \right)^{\frac{1}{r}} < \infty,$$

(e) if $0 < \theta_1 \leq 1, \theta_2 = \infty$, then

$$A_5 := \sup_{t>0} \left(\int_0^t v_1^{\theta_1}(s) ds \right)^{-\frac{1}{\theta_1}} \left(\operatorname{ess\,sup}_{y>0} \min(t, y) v_2(y) \right) < \infty,$$

(f) if $1 < \theta_1 < \infty, \theta_2 = \infty$, then

$$A_6 := \operatorname{ess\,sup}_{s>0} v_2(s) \left(\int_0^s \left(\int_t^s \left(\int_0^y v_1^{\theta_1}(z) dz \right)^{-1} dy \right)^{\theta_1'} v_1^{\theta_1}(t) dt \right)^{\frac{1}{\theta_1'}} < \infty,$$

(g) if $\theta_1 = \infty, 0 < \theta_2 < \infty$, then

$$A_7 := \left(\int_0^\infty \left(v_2(x) \int_0^x \frac{dy}{\operatorname{ess\,sup}_{0 < z < y} v_1(z)} \right)^{\theta_2} dx \right)^{\frac{1}{\theta_2}} < \infty,$$

(h) if $\theta_1 = \theta_2 = \infty$, then

$$A_8 := \operatorname{ess\,sup}_{x>0} \left(v_2(x) \int_0^x \frac{dy}{\operatorname{ess\,sup}_{0 < z < y} v_1(z)} \right) < \infty.$$

Conditions, ensuring boundedness of the generalized Riesz potential

In order to obtain sufficient conditions and necessary and sufficient conditions for $p_1 = 1$ on the weight functions w_1, w_2 ensuring the boundedness of $I_{\rho(\cdot)}$ from $LM_{p_1, \theta_1, w_1(\cdot)}$ to $LM_{p_2, \theta_2, w_2(\cdot)}$ for $1 < p_1 < p_2 < \infty$ or $1 \leq p_1 < \infty$, it, clearly, suffices to apply Theorems 4.2 and 5.1.

In order to make these conditions have a simpler form we shall carry out certain changes of variables and apply equalities (4.15)–(4.20) proved in Lemma 4.1.

Theorem 6.1 *Assume that $0 < \theta_1, \theta_2 \leq \infty, w_1 \in \Omega_{\theta_1}, w_2 \in \Omega_{\theta_2}$.*

1. *Let $1 < p_1 < p_2 < \infty$ or $1 \leq p_1 < \infty$ and $0 < p_2 \leq \infty$, and $\rho \in S_{n, p_1, p_2}$. Then the operator $I_{\rho(\cdot)}$ is bounded from $LM_{p_1, \theta_1, w_1(\cdot)}$ to $LM_{p_2, \theta_2, w_2(\cdot)}$ if the following conditions are satisfied.*

(a) *If $1 < \theta_1 \leq \theta_2 < \infty$, then*

$$B_{11} := \sup_{t>0} \left(\int_t^\infty w_{2,n,p_2}^{\theta_2}(x) \mu_{n,\rho,p_1}^{\theta_2}(x) dx \right)^{\frac{1}{\theta_2}} \left(\int_t^\infty w_1^{\theta_1}(x) dx \right)^{-\frac{1}{\theta_1}} < \infty,$$

and

$$B_{12} := \sup_{t>0} \left(\int_0^t w_{2,n,p_2}^{\theta_2}(x) dx \right)^{\frac{1}{\theta_2}} \left(\int_t^\infty \frac{w_1^{\theta_1}(x) \mu_{n,\rho,p_1}^{\theta_1}(x)}{\left(\int_x^\infty w_1^{\theta_1}(s) ds \right)^{\theta_1}} dx \right)^{\frac{1}{\theta_1}} < \infty.$$

where $w_{2,n,p_2}(x) = w_2(x)x^{\frac{n}{p_2}}$.

(b) *If $0 < \theta_1 \leq 1, \theta_1 \leq \theta_2 < \infty$, then*

$$B_2 := \sup_{t>0} \left(\int_t^\infty w_1^{\theta_1}(x) dx \right)^{-\frac{1}{\theta_1}} \left(\int_0^\infty w_{2,n,p_2}^{\theta_2}(x) \min\{\mu_{n,\rho,p_1}(t), \mu_{n,\rho,p_1}(x)\}^{\theta_2} dx \right)^{\frac{1}{\theta_2}} < \infty.$$

(c) *If $1 < \theta_1 < \infty, 0 < \theta_2 < \theta_1 < \infty$, then*

$$B_{31} := \left(\int_0^\infty \left(\frac{\int_t^\infty w_{2,n,p_2}^{\theta_2}(x) \mu_{n,\rho,p_1}^{\theta_2}(x) dx}{\int_t^\infty w_1^{\theta_1}(x) dx} \right)^{\frac{r}{\theta_1}} w_{2,n,p_2}^{\theta_2}(t) \mu_{n,\rho,p_1}^{\theta_2}(t) dt \right)^{\frac{1}{r}} < \infty,$$

and

$$B_{32} := \left(\int_0^\infty \left(\int_z^\infty w_{2,n,p_2}^{\theta_2}(x) dx \right)^{\frac{r}{\theta_1}} \left(\int_z^\infty \frac{\mu_{n,\rho,p_1}^{\theta_1}(\tau) w_1^{\theta_1}(\tau)}{\left(\int_\tau^\infty w_1^{\theta_1}(u) du \right)^{\theta_1}} d\tau \right)^{\frac{r}{\theta_1}} w_{2,n,p_2}^{\theta_2}(z) dz \right)^{\frac{1}{r}} < \infty,$$

where $\frac{1}{r} = \frac{1}{\theta_2} - \frac{1}{\theta_1}$.

(d) *If $0 < \theta_2 < \theta_1 \leq 1$, then $B_{41} := B_{31} < \infty$ and*

$$B_{42} := \left(\int_0^\infty \sup_{y<z<\infty} \mu_{n,\rho,p_1}(z)^r \left(\int_z^\infty w_1^{\theta_1}(s) ds \right)^{-\frac{r}{\theta_1}} \left(\int_0^y w_{2,n,p_2}^{\theta_2}(x) dx \right)^{\frac{r}{\theta_1}} w_{2,n,p_2}^{\theta_2}(y) dy \right)^{\frac{1}{r}} < \infty.$$

(e) If $0 < \theta_1 \leq 1, \theta_2 = \infty$, then

$$B_5 := \sup_{t>0} \left(\int_0^\infty w_1^{\theta_1}(x) dx \right)^{-\frac{1}{\theta_1}} \operatorname{ess\,sup}_{x>0} w_{2,n,p_2}(x) \min\{\mu_{n,\rho,p_1}(t), \mu_{n,\rho,p_1}(x)\} < \infty.$$

(f) If $1 < \theta_1 < \infty, \theta_2 = \infty$, then

$$B_6 := \operatorname{ess\,sup}_{z>0} w_{2,n,p_2}(z) \left(\int_z^\infty \left(\int_x^z \left(\int_u^\infty w_1^{\theta_1}(s) ds \right)^{-1} \rho(u) u^{\frac{n}{p_1}-1} du \right)^{\theta_1'} w_1^{\theta_1}(x) dx \right)^{\frac{1}{\theta_1'}} < \infty.$$

(g) If $\theta_1 = \infty, 0 < \theta_2 < \infty$, then

$$B_7 := \left(\int_0^\infty \left(w_{2,n,p_2}(\tau) \int_\tau^\infty \frac{\rho(z) z^{\frac{n}{p_1}-1} dz}{\operatorname{ess\,sup}_{z<s<\infty} w_1(s)} \right)^{\theta_2} d\tau \right)^{\frac{1}{\theta_2}} < \infty.$$

(h) If $\theta_1 = \theta_2 = \infty$, then

$$B_8 := \operatorname{ess\,sup}_{\tau>0} \left(\int_\tau^\infty \frac{\rho(z) z^{\frac{n}{p_1}-1} dz}{\operatorname{ess\,sup}_{z<s<\infty} w_1(s)} \right) w_{2,n,p_2}(\tau) < \infty.$$

2. Let $p_1 = 1, 0 < p_2 < \infty$ and $\rho \in \tilde{S}_{n,1,p_2}$. Then the operator $I_{\rho(\cdot)}$ is bounded from $LM_{1,\theta_1,w_1(\cdot)}$ to $LM_{p_2,\theta_2,w_2(\cdot)}$ if and only if conditions (a) - (h) with $p_1 = 1$ are satisfied.

3. Let $p_1 = 1, 1 < p_2 < \infty$ and $\rho \in S_{n,1,p_2}$. Then the operator $I_{\rho(\cdot)}$ is bounded from $LM_{1,\theta_1,w_1(\cdot)}$ to $WLM_{p_2,\theta_2,w_2(\cdot)}$ if and only if conditions (a) - (h) with $p_1 = 1$ are satisfied.

Proof It suffices to prove by using Lemma 4.1 and the appropriate changes of variables that, for μ_{n,ρ,p_1}, ν_1 and ν_2 defined by formulas (4.1), (4.9), (4.2), respectively, $A_{11} = B_{11}, A_{12} = B_{12}, A_2 = B_2, A_{31} = B_{31}, A_{32} = B_{32}, A_{41} = B_{41}, A_5 = B_5$ and $A_6 = cB_6, A_7 = cB_7, A_8 = cB_8$, where $c = \left(\int_1^\infty \rho(t) t^{\frac{n}{p_1}-1} dt \right)^{-1}$.

(a) If $1 < \theta_1 \leq \theta_2 < \infty$, then according to Theorem 5.1 and by (4.15) with $\phi_1 \equiv 1$ and (4.18) with $\phi_2(s) \equiv s$, we get

$$\begin{aligned} A_{11} &= \sup_{t>0} \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(t)}^\infty w_{2,n,p_2}^{\theta_2}(r) \mu_{n,\rho,p_1}^{\theta_2}(r) dr \right)^{\frac{1}{\theta_2}} \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(t)}^\infty w_1^{\theta_1}(r) dr \right)^{-\frac{1}{\theta_1}} \\ &\quad \left(\mu_{n,\rho,p_1}^{(-1)}(t) = \tau \right) \\ &= \sup_{\tau>0} \left(\int_\tau^\infty w_{2,n,p_2}^{\theta_2}(r) \mu_{n,\rho,p_1}^{\theta_2}(r) dr \right)^{\frac{1}{\theta_2}} \left(\int_\tau^\infty w_1^{\theta_1}(r) dr \right)^{-\frac{1}{\theta_1}} = B_{11} \end{aligned}$$

and

$$A_{12} = \sup_{t>0} \left(\int_0^{\mu_{n,\rho,p_1}^{(-1)}(t)} w_{2,n,p_2}^{\theta_2}(x) dx \right)^{\frac{1}{\theta_2}} \left(\int_0^t z^{\theta_1'} \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(z)}^\infty w_1^{\theta_1}(r) dr \right)^{-\theta_1'} w_1^{\theta_1}(z) dz \right)^{\frac{1}{\theta_1'}}.$$

Next, by (4.16) with

$$\phi_1(z) = \left(z^{\theta'_1} \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(z)}^{\infty} w_1^{\theta_1}(r) dr \right)^{-\theta'_1} \right)^{\frac{1}{\theta'_1}},$$

we get

$$\begin{aligned} A_{12} &= \sup_{t>0} \left(\int_0^{\mu_{n,\rho,p_1}^{(-1)}(t)} w_{2,n,p_2}^{\theta_2}(x) dx \right)^{\frac{1}{\theta_2}} \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(t)}^{\infty} \mu_{n,\rho,p_1}^{\theta'_1}(x) \left(\int_x^{\infty} w_1^{\theta_1}(r) dr \right)^{-\theta'_1} w_1^{\theta_1}(x) dx \right)^{\frac{1}{\theta'_1}} \\ &\quad \left(\mu_{n,\rho,p_1}^{(-1)}(t) = \tau \right) \\ &= \sup_{\tau>0} \left(\int_0^{\tau} w_2^{\theta_2}(x) x^{\frac{n\theta_2}{p_2}} dx \right)^{\frac{1}{\theta_2}} \left(\int_{\tau}^{\infty} \mu_{n,\rho,p_1}^{\theta'_1}(x) \left(\int_x^{\infty} w_1^{\theta_1}(r) dr \right)^{-\theta'_1} w_1^{\theta_1}(x) dx \right)^{\frac{1}{\theta'_1}} = B_{12}. \end{aligned}$$

(b) If $0 < \theta_1 \leq 1, \theta_1 \leq \theta_2 < \infty$, then by (4.15) with $\phi_1 \equiv 1$ and by (4.19') with $\phi_2(s) = \min\{t, s\}, s > 0$, we get

$$\begin{aligned} A_2 &= \sup_{t>0} \left(\int_{\mu_{n,\rho,p_1}^{-1}(t)}^{\infty} w_1^{\theta_1}(r) dr \right)^{-\frac{1}{\theta_1}} \left(\int_0^{\infty} w_{2,n,p_2}^{\theta_2}(r) \min\{t, \mu_{n,\rho,p_1}(r)\}^{\theta_2} dr \right)^{\frac{1}{\theta_2}} \\ &\quad \left(\mu_{n,\rho,p_1}^{-1}(t) = \tau \right) \\ &= \sup_{\tau>0} \left(\int_{\tau}^{\infty} w_1^{\theta_1}(r) dr \right)^{-\frac{1}{\theta_1}} \left(\int_0^{\infty} w_{2,n,p_2}^{\theta_2}(r) \min\{\mu_{n,\rho,p_1}(\tau), \mu_{n,\rho,p_1}(r)\}^{\theta_2} dr \right)^{\frac{1}{\theta_2}} = B_2. \end{aligned}$$

(c) If $1 < \theta_1 < \infty, 0 < \theta_2 < \theta_1 < \infty$, then by (4.15) with $\phi_1 \equiv 1$, (4.18) with $\phi_s = s$ and by (4.19') we get

$$\begin{aligned} A_{31} &= \left(\int_0^{\infty} \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(t)}^{\infty} w_1^{\theta_1}(x) dx \right)^{\frac{r}{\theta_1}} \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(t)}^{\infty} w_{2,n,p_2}^{\theta_2}(x) \mu_{n,\rho,p_1}^{\theta_2}(x) dx \right)^{\frac{r}{\theta_1}} t^{\theta_2} v_2^{\theta_2}(t) dt \right)^{\frac{1}{r}} \\ &= \left(\int_0^{\infty} \left(\int_z^{\infty} w_1^{\theta_1}(x) dx \right)^{\frac{r}{\theta_1}} \left(\int_z^{\infty} w_{2,n,p_2}^{\theta_2}(x) \mu_{n,\rho,p_1}^{\theta_2}(x) dx \right)^{\frac{r}{\theta_1}} \mu_{n,\rho,p_1}^{\theta_2}(z) w_{2,n,p_2}^{\theta_2}(z) dz \right)^{\frac{1}{r}} = B_{31} \end{aligned}$$

and

$$A_{32} = \left(\int_0^{\infty} \left(\int_0^{\mu_{n,\rho,p_1}^{(-1)}(t)} w_{2,n,p_2}^{\theta_2}(x) dx \right)^{\frac{r}{\theta_1}} \left(\int_0^t \frac{s^{\theta'_1}}{\left(\int_{\mu_{n,\rho,p_1}^{(-1)}(s)}^{\infty} w_1^{\theta_1}(u) du \right)^{\theta'_1}} v_1^{\theta_1}(s) ds \right)^{\frac{r}{\theta'_1}} v_2^{\theta_2}(t) dt \right)^{\frac{1}{r}}$$

$$\begin{aligned}
&= \left(\int_0^\infty \left(\int_0^{\mu_{n,\rho,p_1}^{(-1)}(t)} w_{2,n,p_2}^{\theta_2}(x) dx \right)^{\frac{r}{\theta_1}} \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(\tau)}^\infty \frac{\mu_{n,\rho,p_1}^{\theta_1}(\tau) w_1^{\theta_1}(\tau)}{\left(\int_\tau^\infty w_1^{\theta_1}(u) du \right)^{\theta_1'}} d\tau \right)^{\frac{r}{\theta_1'}} v_2^{\theta_2}(t) dt \right)^{\frac{1}{r}} \\
&= \left(\int_0^\infty \left(\int_z^\infty w_{2,n,p_2}^{\theta_2}(x) dx \right)^{\frac{r}{\theta_1}} \left(\int_z^\infty \frac{\mu_{n,\rho,p_1}^{\theta_1}(\tau) w_1^{\theta_1}(\tau)}{\left(\int_\tau^\infty w_1^{\theta_1}(u) du \right)^{\theta_1'}} d\tau \right)^{\frac{r}{\theta_1'}} w_{2,n,p_2}^{\theta_2}(z) dz \right)^{\frac{1}{r}} = B_{32}.
\end{aligned}$$

(d) If $0 < \theta_2 < \theta_1 \leq 1$, then by (4.15) with $\phi_1 \equiv 1$, by (4.19) with $\phi_2 \equiv 1$, we get

$$\begin{aligned}
A_{41} &= \left(\int_0^\infty \sup_{0 < \tau < t} \tau^r \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(\tau)}^\infty w_1^{\theta_1}(u) du \right)^{\frac{r}{\theta_1}} \left(\int_0^{\mu_{n,\rho,p_1}^{(-1)}(t)} w_{2,n,p_2}^{\theta_2}(x) dx \right)^{\frac{r}{\theta_1}} v_2^{\theta_2}(t) dt \right)^{\frac{1}{r}} \\
&= \left(\int_0^\infty \sup_{0 < \tau < \mu_{n,\rho,p_1}^{(-1)}(y)} \tau^r \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(\tau)}^\infty w_1^{\theta_1}(s) ds \right)^{\frac{r}{\theta_1}} \left(\int_0^y w_{2,n,p_2}^{\theta_2}(x) dx \right)^{\frac{r}{\theta_1}} w_{2,n,p_2}^{\theta_2}(y) dy \right)^{\frac{1}{r}} \\
&\quad \left(\tau = \mu_{n,\rho,p_1}^{(-1)}(z) \right) \\
&= \left(\int_0^\infty \operatorname{ess\,sup}_{y < z < \infty} \mu_{n,\rho,p_1}(z)^r \left(\int_z^\infty w_1^{\theta_1}(s) ds \right)^{\frac{r}{\theta_1}} \left(\int_0^y w_{2,n,p_2}^{\theta_2}(x) dx \right)^{\frac{r}{\theta_1}} \mu_{n,\rho,p_1}(x)^{\theta_2}(y) dy \right)^{\frac{1}{r}} = B_{42}.
\end{aligned}$$

(e) If $0 < \theta_1 \leq 1, \theta_2 = \infty$, then by (4.15) with $\phi_1 \equiv 1$ and (4.19') with $\phi(y) = \min\{t, y\}, y > 0$, we get

$$\begin{aligned}
A_5 &= \sup_{t > 0} \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(t)}^\infty w_1^{\theta_1}(x) dx \right)^{-\frac{1}{\theta_1}} \operatorname{ess\,sup}_{x > 0} w_{2,n,p_2}(x) \min\{t, \mu_{n,\rho,p_1}(x)\} \\
&\quad \left(\mu_{n,\rho,p_1}^{(-1)}(t) = \tau \right) \\
&= B_{42} = \sup_{\tau > 0} \left(\int_\tau^\infty w_1^{\theta_1}(x) dx \right)^{-\frac{1}{\theta_1}} \operatorname{ess\,sup}_{x > 0} w_{2,n,p_2}(x) \min\{\mu_{n,\rho,p_1}(\tau), \mu_{n,\rho,p_1}(x)\} = B_5.
\end{aligned}$$

(f) If $1 < \theta_1 < \infty, \theta_2 = \infty$, then by (4.15) with $\phi_1 \equiv 1$ and by the same formula with

$$\phi_1(t) = \left(\int_t^s \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(y)}^\infty w_1^{\theta_1}(s) ds \right)^{-1} dy \right)^{\frac{\theta_1'}{\theta_1}}$$

and, finally, by (4.19') with $\theta_2 = \infty$ and

$$\phi_2(s) = \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(s)}^{\infty} \left(\int_{\mu_{n,\rho,p_1}(r)}^s \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(y)}^{\infty} w_1^{\theta_1}(s) ds \right)^{-1} dy \right)^{\theta_1'} w_1^{\theta_1}(r) dr \right)^{\frac{1}{\theta_1'}}$$

we get

$$\begin{aligned} A_6 &= \operatorname{ess\,sup}_{s>0} v_2(s) \left(\int_0^s \left(\int_r^s \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(y)}^{\infty} w_1^{\theta_1}(s) ds \right)^{-1} dy \right)^{\theta_1'} v_1^{\theta_1}(t) dt \right)^{\frac{1}{\theta_1'}} \\ &= \operatorname{ess\,sup}_{s>0} v_2(s) \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(s)}^{\infty} \left(\int_{\mu_{n,\rho,p_1}(r)}^s \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(y)}^{\infty} w_1^{\theta_1}(s) ds \right)^{-1} dy \right)^{\theta_1'} w_1^{\theta_1}(r) dr \right)^{\frac{1}{\theta_1'}} \\ &= \operatorname{ess\,sup}_{z>0} w_2(z) z^{\frac{n}{p_2}} \left(\int_z^{\infty} \left(\int_{\mu_{n,\rho,p_1}(r)}^z \left(\int_{\mu_{n,\rho,p_1}^{(-1)}(y)}^{\infty} w_1^{\theta_1}(s) ds \right)^{-1} dy \right)^{\theta_1'} w_1^{\theta_1}(r) dr \right)^{\frac{1}{\theta_1'}} \\ &\quad (y = \mu_{n,\rho,p_1}(u)) \\ &= cB_6 = \operatorname{ess\,sup}_{z>0} w_{2,n,p_2}(z) \left(\int_z^{\infty} \left(\int_r^z \left(\int_u^{\infty} w_1(s)^{\theta_1} ds \right)^{-1} |(\mu_{n,\rho,p_1}(u))'| du \right)^{\theta_1'} w_1^{\theta_1}(r) dr \right)^{\frac{1}{\theta_1'}} \end{aligned}$$

where

$$c = \left(\int_1^{\infty} \rho(t) t^{\frac{n}{p_1}-1} dt \right)^{-1}.$$

(g) If $\theta_1 = \infty, 0 < \theta_2 < \infty$, then by using formula (4.15) with $\phi_1 \equiv 1$ and formula (4.19') with $\phi_2(x) = \int_0^x \frac{dy}{\operatorname{ess\,sup}_{\substack{r>\mu_{n,\rho,p_1}^{(-1)}(y)}} w_1(r)}$

we get

$$\begin{aligned} A_7 &= \left(\int_0^{\infty} \left(v_2(x) \int_0^x \frac{dy}{\operatorname{ess\,sup}_{r>\mu_{n,\rho,p_1}^{(-1)}(y)} w_1(r)} \right)^{\theta_2} dx \right)^{\frac{1}{\theta_2}} \\ &= \left(\int_0^{\infty} \left(w_{2,n,p_2}(\tau) \int_0^{\mu_{n,\rho,p_1}(\tau)} \frac{dy}{\operatorname{ess\,sup}_{r>\mu_{n,\rho,p_1}^{(-1)}(y)} w_1(r)} \right)^{\theta_2} d\tau \right)^{\frac{1}{\theta_2}} \\ &\quad \left(\mu_{n,\rho,p_1}^{(-1)}(y) = z \right) \end{aligned}$$

$$= c \left(\int_0^\infty \left(w_{2,n,p_2}(\tau) \int_\tau^\infty \frac{\rho(z)z^{\frac{n}{p_1}-1}}{\operatorname{ess\,sup}_{z < r < \infty} w_1(r)} dz \right)^{\theta_2} d\tau \right)^{\frac{1}{\theta_2}} = cB_7.$$

(h) If $\theta_1 = \theta_2 = \infty$, then in A_8 according to (4.10) and (4.2)

$$v_1(x) = w_1 \left(\mu_{n,\rho,p_1}^{(-1)}(x) \right), \quad v_2(x) = w_2 \left(\mu_{n,\rho,p_1}^{(-1)}(x) \right) \left(\mu_{n,\rho,p_1}^{(-1)}(x) \right)^{\frac{n}{p_2}}, \quad x > 0.$$

By using formula (4.15) with $\psi_1 \equiv 1$ and by carrying out the following changes of variables: $y = \mu_{n,\rho,p_1}(z)$, $z > 0$, and, finally, $t = \mu_{n,\rho,p_1}(\tau)$, $\tau > 0$, we get

$$\begin{aligned} A_8 &= \operatorname{ess\,sup}_{t>0} \left(\int_0^t \frac{dy}{\operatorname{ess\,sup}_{s>\mu_{n,\rho,p_1}^{(-1)}(y)} w_1(s)} \right) w_2 \left(\mu_{n,\rho,p_1}^{(-1)}(t) \right) \left(\mu_{n,\rho,p_1}^{(-1)}(t) \right)^{\frac{n}{p_2}} \\ &= c \operatorname{ess\,sup}_{t>0} \left(\int_{\mu_{n,\rho,p_2}^{(-1)}(t)}^\infty \frac{\rho(z)z^{\frac{n}{p_1}-1}}{\operatorname{ess\,sup}_{s>z} w_1(s)} dz \right) w_2 \left(\mu_{n,\rho,p_2}^{(-1)}(t) \right) \left(\mu_{n,\rho,p_2}^{(-1)}(t) \right)^{\frac{n}{p_2}} \\ &= c \operatorname{ess\,sup}_{\tau>0} \left(\int_\tau^\infty \frac{\rho(z)z^{\frac{n}{p_1}-1}}{\operatorname{ess\,sup}_{z<s<\infty} w_1(s)} dz \right) w_{2,n,p_2}(\tau) = cB_8. \end{aligned}$$

□

Acknowledgements The authors are very grateful to the referees for careful reading of the paper and comments which allowed to improve the quality of the text.

Funding Sections 1–3 of this work were performed by V.I. Burenkov and M.A. Senouci at the RUDN University and supported by the Regional Mathematical Center of the Southern Federal University with the Agreement No. 075-02-2022-893 of the Ministry of Science and Higher Education of Russia. Sections 4–6 of this work were performed by them at the V.A. Steklov Mathematical Institute and supported by the Russian Science Foundation (project no. 19-11-00087, <https://rscf.ru/project/19-11-00087/>).

Associated data The manuscript has no associated data.

Declarations

Conflict of interest The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

REFERENCES

1. V.I. Burenkov, Recent progress in studying the boundedness of classical operators of real analysis in general Morrey-type spaces. I, *Eurasian Math. J.*, 3 (2012), no. 3, 11–32.
2. V.I. Burenkov, Recent progress in studying the boundedness of classical operators of real analysis in general Morrey-type spaces. II, *Eurasian Math. J.*, 4 (2013), no. 1, 21–45

3. V.I. Burenkov, H.V. Guliyev, Necessary and sufficient conditions for boundedness of the maximal operator in local Morrey-type spaces, *Studia Math*, 163 (2004), no. 2, 157-176.
4. V.I. Burenkov, H.V. Guliyev, V.S. Guliyev, Necessary and sufficient conditions for the boundedness of fractional maximal operators in local Morrey-type spaces, *J. Comput. Appl. Math.*, 208 (2007), 280-301.
5. V.I. Burenkov, P. Jain, T.V. Tararykova, On boundedness of the Hardy operator in Morrey-type spaces, *Eurasian Math. J.*, 2 (2011), no. 1, 52-80.
6. V.I. Burenkov, A. Gogatishvili, V.S. Guliyev, R.Ch. Mustafayev, Boundedness of the Riesz potential in local Morrey-type spaces, *Potential Analysis*, 35 (2011), no. 1, 67-87.
7. V.I. Burenkov, M.A. Senouci, Boundedness of the generalized Riesz potential in local Morrey type spaces, *Eurasian Math. J.*, 12 (2021), no. 4, 92–98.
8. M. Carro, L. Pick, J. Soria, V.D. Stepanov, On embeddings between classical Lorentz spaces, *Math. Ineq. Appl.*, 4 (2001), no. 3, 397-428.
9. M. Carro, A. Gogatishvili, J. Martin, L. Pick, Weighted inequalities involving two Hardy operators with applications to embeddings of function spaces, *J. Oper. Theory*, 59 (2008), no. 2, 309-332.
10. A. Gogatishvili, V.D. Stepanov, Reduction theorems for weighted integral inequalities on the cone of monotone functions, *Uspekhi Mat. Nauk* 68 (2013), no. 4 (412), 3–68 (in Russian); English transl., *Russian Math. Surveys* 68 (2013), no. 4, 597–664. MR3154814
11. M.L. Goldman, Hardy type inequalities on the cone of quasimonotone functions, Research Report No: 98/31. Khabarovsk: Computer Center, Far Eastern Branch of the Russian Academy of Sciences, 1998, 1-69.
12. V.S. Guliyev, Generalized weighted Morrey spaces and higher order commutators of sublinear operators, *Eurasian Math. J.*, 3 (2012), no. 3, 33–61.
13. P.-G. Lemarié-Rieusset, The role of Morrey spaces in the study of Navier–Stokes and Euler equations, *Eurasian Math. J.*, 3 (2012), no. 3, 62–93.
14. M.A. Ragusa, Operators in Morrey type spaces and applications, *Eurasian Math. J.*, 3 (2012), no. 3, 94–109.
15. W. Sickel, Smoothness spaces related to Morrey spaces – a survey. I, *Eurasian Math. J.*, 3 (2012), no. 3, 110–149.
16. W. Sickel, Smoothness spaces related to Morrey spaces — a survey. II, *Eurasian Math. J.*, 4 (2013), no. 1, 82–124.