

What works to support better access to mental health services (from primary care to inpatients) for minority groups to reduce inequalities? A rapid evidence summary.

Authors: Judit Csontos¹, Deborah Edwards¹, Elizabeth Gillen¹, Juliet Hounsome², Meg Kiseleva², Mala Mann², Abubakar Sha'aban³, Ruth Lewis⁴, Alison Cooper³, Adrian Edwards³

1 Wales Centre for Evidence Based Care (WCEBC), Cardiff University, United Kingdom

2 Specialist Unit for Review Evidence (SURE), Cardiff University, United Kingdom

3 Health and Care Research Wales Evidence Centre, Cardiff University, United Kingdom

4 Health and Care Research Wales Evidence Centre, Bangor University, United Kingdom

Abstract:

It is estimated that one in four people will experience poor mental health throughout their lifetimes. However, ethnic minority groups, refugees and asylum seekers experience more barriers accessing mental health services and have poorer mental health outcomes than those from non-ethnic minority groups. Evidence suggests that interventions that improve access and engagement with mental health services may help reduce disparities affecting ethnic minority groups, refugees and asylum seekers. Thus, the aim of this rapid evidence summary was to explore the literature on what works to support better access to mental health services for ethnic minority groups, refugees and asylum seekers to reduce inequalities. The review included interventions that were developed or assessed to improve equity in access, engagement, utilisation, or provision of mental health services.

Research Implications and Evidence Gaps:

There is limited review evidence regarding the effectiveness of interventions to improve access to mental healthcare across ethnic minority groups. Review evidence regarding interventions to support refugees and asylum seekers access to primary healthcare or specialised clinics (for example pregnancy and postpartum) is available, but the findings related to mental health care cannot be extracted.

Funding statement:

The Wales Centre for Evidence Based Care and the Specialist Unit for Review Evidence for Health Economics were funded for this work by the Health and Care Research Wales Evidence Centre, itself funded by Health and Care Research Wales on behalf of Welsh Government.



Health and Care
Research Wales
Evidence Centre
Canolfan Dystiolaeth
Ymchwil Iechyd a
Gofal Cymru

What works to support better access to mental health services (from primary care to inpatients) for minority groups to reduce inequalities? A rapid evidence summary.

February 2024

Review conducted by the Cardiff Evidence Synthesis Collaborative.

Report Contributors

Review Team

Judit Csontos, Deborah Edwards, Elizabeth Gillen, Juliet Hounsome, Meg Kiseleva, Mala Mann, Abubakar Sha'aban

Methodological Advice

Ruth Lewis

Evidence Centre Team

Ruth Lewis, Adrian Edwards, Alison Cooper, Elizabeth Doe involved in stakeholder engagement, review of report and editing

Public Partners

Olivia Gallen and Mel (Melanie) McAulay

Stakeholders

Beverly Morgan, Kim Swain, Hannah Bayfield

Evidence need submitted to the Evidence Centre: May 2023

Initial Stakeholder Consultation Meeting: June 2023

Final report issued: February 2024

This review should be cited as: Health and Care Research Wales Evidence Centre. What works to support better access to mental health services (from primary care to inpatients) for minority groups to reduce inequalities? A rapid evidence summary. (RES0024). February 2024.

Disclaimer: The views expressed in this publication are those of the authors, not necessarily Health and Care Research Wales. The Health and Care Research Wales Evidence Centre and authors of this work declare that they have no conflict of interest.

What works to support better access to mental health services (from primary care to inpatients) for minority groups to reduce inequalities? (Rapid Evidence Summary)

Report number RES0024 (February 2024)

EXECUTIVE SUMMARY

What is a Rapid Evidence Summary?

Our Rapid Evidence Summaries (RES) are designed to provide a rapid response product. They are based on a limited search of key resources and the assessment of abstracts. Priority is given to studies representing robust evidence synthesis. No quality appraisal or evidence synthesis are conducted, and the summary should be interpreted with caution.

This report is linked to a subsequent focused rapid review, to be published in Summer 2024, on the effectiveness of interventions to enhance equitable or overall access to mental health services by ethnic minority groups.

Who is this Rapid Evidence Summary for?

Health and Social Services Group - Mental Health & Vulnerable Groups (Policy).

Background / Aim of Rapid Evidence Summary

It is estimated that one in four people will experience poor mental health¹ throughout their lifetimes. However, ethnic minority groups¹, refugees and asylum seekers¹ experience more barriers accessing mental health services and have poorer mental health outcomes than those from non-ethnic minority groups. Evidence suggests that interventions that improve access and engagement with mental health services may help reduce disparities affecting ethnic minority groups, refugees and asylum seekers. Thus, the aim of this rapid evidence summary was to explore the literature on what works to support better access to mental health services for ethnic minority groups, refugees and asylum seekers to reduce inequalities. The review included interventions that were developed or assessed to improve equity¹ in access, engagement, utilisation, or provision of mental health services.

Results

Recency of the evidence base

- The included literature was published between 2006 and 2023.

Extent of the evidence base

- Bibliographic database searches identified systematic reviews¹ (n=19), scoping reviews¹ (n=7), a mapping review¹ (n=1), narrative reviews¹ (n=3), overviews of reviews¹ (n=2), a scoping review of reviews (n=1) and systematic review protocols (n=4).
- The stakeholders identified organisational reports (n=4), a rapid review¹ (n=1), a scoping review (n=1), and a systematic review (n=1).

^{1,1} Defined in the [Glossary](#)

Key findings

- The identified literature was summarised under the categories of: Access and pathways to mental health care (separately summarised for ethnic minority groups; Gypsy, Roma and Traveller communities; and refugees and asylum seekers); Mental health promotion, prevention, and treatment of mental health conditions including cultural adaptations to psychological interventions; Help-seeking behaviour; Treatment engagement; and Treatment initiation, participation or continuation.
- There is a **wealth of review evidence**, from across the UK and internationally, on the following topics: **barriers and facilitators to accessing mental health care** for ethnic minority groups, refugees and asylum seekers; **mental health promotion, prevention, and treatment** of mental health conditions (including treatment initiation, participation or continuation) for ethnic minority groups, refugees and asylum seekers; and **cultural adaptations** of psychological interventions whether it is for ethnic minority groups living in Western countries or majority populations living in non-Western countries.
- While evidence is available on improving mental health care access and experience of ethnic minority groups, refugees and asylum seekers, these often do not have an evaluative component and report their findings as suggestions / recommendations for interventions. Most common **recommendations to improve mental health care equity comprised** of: language and cultural adaptations; education of healthcare professionals; employing ethnically diverse staff; better information provision; collaborative working between different sectors; facilitating referral routes and improving pathways; specialist services for minority groups and outreach services; patient education and skill development; involvement of communities.
- Culturally adapted interventions (either mental health promotion, prevention or treatment) may lead to positive outcomes, including improved symptom severity, behaviours, self-efficacy, and wellness, although there is a lack of consensus in what components of adaptations work for which cultural groups. **It is unclear whether cultural adaptations truly result in improved outcomes compared to non-adapted interventions**, and there is a need for high quality well-designed randomised controlled trials.

Research Implications and Evidence Gaps

- There is limited review evidence regarding the effectiveness of interventions to improve access to mental healthcare across ethnic minority groups.
- Review evidence regarding interventions to support refugees' and asylum seekers' access to primary healthcare or specialised clinics (for example pregnancy and postpartum) is available, but the findings related to mental health care cannot be extracted.

Disclaimer: The views expressed in this publication are those of the authors, not necessarily Health and Care Research Wales. The Health and Care Research Wales Evidence Centre and authors of this work declare that they have no conflict of interest.

TABLE OF CONTENTS

1. CONTEXT / BACKGROUND	9
2. RESEARCH QUESTION(S).....	10
3. SUMMARY OF THE EVIDENCE BASE	11
3.1. Type and amount of evidence available	11
3.2. Key Findings	15
3.3. Areas of uncertainty	20
4. NEXT STEPS	20
5. REFERENCES.....	20
6. RAPID EVIDENCE SUMMARY METHODS	26
7. EVIDENCE	26
8. ADDITIONAL INFORMATION	55
8.1. Conflict of interest.....	55
8.2. Acknowledgments	55
9. APPENDIX.....	56

List of Tables

Table 1: Summary of review evidence identified	26
Table 2: Summary of included secondary evidence.....	27
Table 3: Summary of included secondary evidence identified by stakeholders	49
Table 4: Summary of included organisational reports identified by stakeholders.....	52

Abbreviations

Acronym	Full Description
BAME	Black, Asian and minority ethnic
NHS	National Health Service
OECD	Organisation for Economic Co-operation and Development
ONS	Office for National Statistics
PTSD	Post-traumatic stress disorder
RCT	Randomised Control Trial
VCSE	Voluntary Community and Social Enterprise

Glossary

Disparity: “Health disparity and health inequality are broad terms that include health inequity and signify more than just difference or variation: they signify a health difference that raises moral or ethical concerns.” (Braveman et al. 2018, p. 11)

Equity: Health equity means that “everyone has a fair and just opportunity to be as healthy as possible. Achieving health equity requires removing obstacles to health such as poverty, discrimination, and their consequences, which include powerlessness and lack of access to good jobs with fair pay; quality education, housing, and health care; and safe environments. For the purposes of measurement, health equity means reducing and ultimately eliminating disparities in health and health determinants that adversely affect excluded or marginalized groups. Health equity is the ethical and human rights principle motivating efforts to eliminate health disparities; health disparities are the metric for assessing progress toward health equity.” (Braveman et al. 2018, p. 11)

Ethnic minorities: The term refers to “all ethnic groups except the white British group. Ethnic minorities include white minorities, such as Gypsy, Roma and Irish Traveller groups.” (UK Government 2023b)

Mental health: Welsh Government defines mental health as a state of mental wellbeing that enables people to cope with the stresses of life, realise their abilities, learn well and work well, and contribute to their community. It is an integral component of health and wellbeing that underpins our individual and collective abilities to make decisions, build relationships and shape the world we live in. Mental health is a basic human right. And it is crucial to personal, community and socio-economic development. People with poor mental health can have a mental health condition but this is not always or necessarily the case.

Mental health condition: Welsh Government defines mental health condition as a broad term covering conditions that affect emotions, thinking and behaviour, and which substantially interfere with our life. Mental health conditions can significantly impact daily living, including our ability to work, care for ourselves and our family, and our ability to relate and interact with others. This is a term used to cover several conditions (e.g. depression, post-traumatic stress disorder, schizophrenia) with different symptoms and impacts for varying lengths of time, for each person. Mental health conditions can range from mild through to severe and enduring illness. People with mental health conditions are more likely to experience lower levels of physical and mental wellbeing, but this is not always or necessarily the case. Some mental health conditions like eating disorders and schizophrenia are associated with a higher risk of mortality.

Refugees and asylum seekers: “Refugees are people forced to flee their own country and seek safety in another country. They are unable to return to their own country because of feared persecution as a result of who they are, what they believe in or say, or because of

armed conflict, violence or serious public disorder” (UNHCR 2024b). “An asylum-seeker is someone whose request for sanctuary has yet to be processed” (UNHCR 2024a).

Different review types:

Mapping review: “is a transparent, rigorous, and systematic approach to identifying, describing, and cataloguing evidence and evidence gaps in a broader topic area.” (Campbell et al. 2023, p. 5)

Narrative review: Often referred to as literature review, can be defined as a “generic term for published materials that provide examination of recent or current literature, can cover wide range of subjects at various levels of completeness and comprehensiveness. It may include research findings.” (Grant and Booth 2009, p. 94)

Overview of reviews: “uses explicit and systematic methods to search for and identify multiple systematic reviews on related research questions in the same topic area for the purpose of extracting and analysing their results across important outcomes. Thus, the unit of searching, inclusion and data analysis is the systematic review.” (Pollock et al. 2023)

Rapid review: “a type of knowledge synthesis in which systematic review processes are accelerated and methods are streamlined to complete the review more quickly than is the case for typical systematic reviews.” (Tricco et al. 2017, p. 3)

Scoping review: “a type of evidence synthesis that aims to systematically identify and map the breadth of evidence available on a particular topic, field, concept, or issue, often irrespective of source (i.e., primary research, reviews, non-empirical evidence) within or across particular contexts. Scoping reviews can clarify key concepts/definitions in the literature and identify key characteristics or factors related to a concept, including those related to methodological research.” (Munn et al. 2022, p. 950)

Systematic review: “A systematic review attempts to identify, appraise and synthesize all the empirical evidence that meets pre-specified eligibility criteria to answer a specific research question. Researchers conducting systematic reviews use explicit, systematic methods that are selected with a view aimed at minimizing bias, to produce more reliable findings to inform decision making.” (Cochrane Library 2024)

1. CONTEXT / BACKGROUND

It is estimated that one in four people will experience poor mental health² throughout their lifetimes (Centre for Mental Health 2020). However, international research has documented that ethnic minority groups² experience more barriers to access to mental health services than non-ethnic minorities (Lowther-Payne et al. 2023). At the same time, these groups are at higher risk of mental health conditions² and this risk is often associated with being disproportionately impacted by detrimental social factors, such as racism and poverty (Bignall et al. 2019). Additionally, refugees and asylum seekers² experience higher rates of post-traumatic stress disorder (PTSD) and other mental health disorders linked to dangers associated with migration, poverty, and poor access to health care (Hynie et al. 2023, Iqbal et al. 2022).

Wales is home for diverse ethnic minority groups, including approximately 89,000 people who identify as Asian, Asian British or Asian Welsh), 28,000 Black, Black British, Black Welsh, Caribbean or African individuals, 49,000 mixed or multiple ethnic people, 26,000 members of other ethnic groups (ONS 2022), and 3,630 Gypsy and Irish Traveller residents (ONS 2023). Moreover, around 3,000 asylum seekers are supported in Wales (UK Government 2023a). Together for Mental Health, the Welsh Government's mental health strategy has focused on ensuring equality since 2012 (Welsh Government 2012). However, the equality impact assessment of this strategy found that since the publication of Together for mental health, stigma and discrimination was still more prevalent for people with protected characteristics, including ethnic minority groups (Welsh Government 2014). In addition, the COVID-19 pandemic highlighted systemic issues, and disproportionately affected ethnic minority groups, prompting the Welsh Government to focus more on reducing health inequalities in their Together for Mental Health updated delivery plan (Welsh Government 2020). A recently published research review of the Together for Mental Health Delivery Plan acknowledged that ethnic minority groups in Wales had poorer mental health outcomes than the wider public while identifying potential barriers to access and service provision (Lock et al. 2023). Improvement in the cultural competency of mental health services and providers in Wales was identified as necessary to help the engagement of ethnic minority groups (Lock et al. 2023). Additionally, issues with multi-lingual service provision in common international languages was also mentioned in the report, which could negatively impact on ethnic minority groups' help seeking who can discuss their condition better in a language other than English (Lock et al. 2023).

Evidence from the wider international literature also suggests that disparities² in ethnic minority groups' engagement with mental health services exist along the entire care pathway, with them being less likely to initiate mental health care and fill prescriptions and more likely to end treatment early (Aggarwal et al. 2016, Interian et al. 2013). These disparities are linked to various individual, organisational, and systemic barriers (Aggarwal et al. 2016). Individual-level barriers may include insufficient information to make treatment decisions, communication difficulties and linguistic issues, lack of trust in service providers, psychological distress, fear of stigma, and cultural beliefs resulting in feeling shame about seeking mental health support. Organisational-level barriers refer to unequal access to services and lack of cultural competence in service providers; and the systemic level includes poor funding of mental health services and inaccessibility of information about available services as well as broader issues such as lack of access to transportation or childcare necessary to attend mental health services (Aggarwal et al. 2016). Additionally, barriers to access to and underutilisation of health services, including mental health care, has also been noted among refugees and asylum seekers (Hynie et al. 2023, Iqbal et al. 2022).

² Defined in the [Glossary](#)

To improve ethnic minority groups' access to mental health services in Wales, a Mental Health Ethnic Minorities Task and Finish Group with the Wales Alliance for Mental Health was set up as part of the Anti-Racist Wales Action Plan (Welsh Government 2022). Moreover, suggestions were made that the new mental health strategy should be developed with the involvement of community organisations, third sector and NHS to make sure that the needs and experiences of ethnic minority groups are considered (Welsh Government 2022). To support these plans, it is also crucial to know what evidence is already available on interventions that could support ethnic minority groups' access to mental health services. It is thought that effective interventions to improve access and engagement with services may reduce disparities (Interian et al. 2013). Gask et al. (2012) described two conventional approaches to improving access to psychosocial therapies for common mental health problems in underserved groups. Firstly, the 'default' position of applying interventions developed for the general population e.g. collaborative care and secondly, modification of existing interventions to make them more acceptable to underserved populations such as developing culturally sensitive psychological therapies. This Rapid Evidence Summary will summarise the available evidence for these two approaches for ethnic minority groups.

2. RESEARCH QUESTION(S)

Review question	
What works to support better access to mental health services (from primary care to inpatients) for ethnic minority groups to reduce inequalities?	
Participants	Adults and children - Ethnic minorities (including Gypsy, Roma and Traveller communities) - Refugees - Immigrants - Asylum seekers
Condition/s	Mental wellbeing Psychological health Mental health - Anxiety - Depression - PTSD Severe mental illness - Bipolar - Schizophrenia - Psychosis Dementia
Context	Studies that address - Accessing services - Engaging with services - Utilising services - Provision of services
Other Study Considerations	
All types of reviews (systematic, narrative, scoping, rapid) No geographical restrictions No date restrictions	

3. SUMMARY OF THE EVIDENCE BASE

Secondary research evidence, such as systematic³, scoping³, and narrative³ reviews and overviews of reviews³, were identified via searches of various bibliographic databases and the stakeholders. The total number of reports identified is summarised in Table 1 and a summary of each report is provided in Tables 2 to 4 (Section 7). Further detail on the type of evidence identified, their focus and key findings are presented below.

While ethnic minority groups, refugees, asylum seekers, mental health and mental health conditions are defined in the glossary for this rapid evidence summary, included reviews referred to these concepts in different ways. Below the evidence is summarised by using the terms the original authors of the included reviews adopted.

3.1. Type and amount of evidence available

Nineteen systematic reviews (Aggarwal et al. 2016, Anderson et al. 2015, Arundell et al. 2021, Bhui et al. 2015, Castellanos et al. 2019, Degnan et al. 2018, Ellis et al. 2022, Giebel et al. 2015, Hernandez Robles et al. 2018, Interian et al. 2013, Iqbal et al. 2022, Lee-Tauler et al. 2018, Li et al. 2023, McFadden et al. 2018, Moffat et al. 2009, Moore 2018, Sass et al. 2009, Soltan et al. 2022, Taylor et al. 2023), **seven** scoping reviews (Apers et al. 2023, Handtke et al. 2019, Hynie et al. 2023, Place et al. 2021, Ratnayake et al. 2022, Rogers et al. 2020, Thomson et al. 2015), **one** mapping review³ (Arundell et al. 2020), **three** narrative review (Griner and Smith et al. 2006, Kalibatseva and Leong 2014, Mukadam et al. 2013), **two** overviews of reviews (Rathod et al. 2018, Uphoff et al. 2020), **one** scoping review of reviews (Chakanyuka et al. 2022) and **four** protocols (Aslam et al. 2018, Stockwell and Roche 2022, Wang and Kim 2019, Wood et al 2023) were identified across the literature searches of bibliographic databases. The detail from each of these reviews is reported in Table 2.

Four organisational reports (Roberts 2023, Traumatic Stress Wales 2023, VCSE Health and Wellbeing Alliance 2022, Welsh Government 2023), **one** rapid review (Kunorubwe et al. 2023), **one** scoping review (Pollard and Howard 2021), and **one** systematic review (Satinsky et al. 2019) were identified by the stakeholder group. The detail from each of these organisational reports and reviews are reported across Tables 3 to 4.

3.1.1 Summary of existing reviews identified via bibliographic databases

This section describes the included reviews and their research aims. Key findings of the included review evidence are summarised in Section 3.2.

The systematic reviews focused on:

- Assessing the relationship between clinician communication and the engagement of racial/ethnic minorities in mental health services across the UK, USA, Australia, Spain and Europe (Aggarwal et al. 2016).
- Assessing the effects of community coalition-driven interventions in improving health status or reducing health disparities among racial and ethnic minority populations of all ages (infant to elderly populations), across the USA, Australia, UK and the Netherlands (Anderson et al. 2015).
- Assessing the efficacy of adapted psychological interventions for people from ethnic minority groups. The specific questions asked were what types of adaptations have been implemented, are the types of culturally adapted interventions differentially effective and what effects of culturally adapted interventions are found across different ethnic minority groups (Arundell et al. 2021).

³ Defined in the [Glossary](#)

- Interventions designed to improve therapeutic communications for black and minority ethnic people (young people, adults and the elderly) using psychiatric services across the UK, USA, Canada and Germany (Bhui et al. 2015).
- The effectiveness of mindfulness-based interventions and their cultural adaptations for Hispanic populations from USA, Spain, Chile and Colombia (Castellanos et al. 2019).
- Cultural adaptations of psychosocial interventions for schizophrenia, their effectiveness, specific outcomes and how they were adapted to non-Western populations across 13 different countries (which included ethnic minorities) (Degnan et al. 2018).
- The acceptability and effectiveness of culturally adapted digital mental health interventions for ethnic / racial minorities across the world (Ellis et al. 2022).
- Barriers and facilitators of South Asian older adults' access to dementia care services in the UK, USA and Canada (Giebel et al. 2015).
- The effectiveness and characteristics of culturally adapted substance use interventions for Latino adolescents in the USA (Hernandez Robles et al. 2018).
- Interventions to improve mental health treatment engagement of underserved racial-ethnic adult groups in the USA (Interian et al. 2013).
- Primary health care interventions to improve the quality of health care (including mental health care) provided to refugees and asylum seekers across OECD countries (Iqbal et al. 2022).
- Interventions to improve the initiation of mental health care among racial-ethnic minority groups in the USA (Lee-Tauler et al. 2018).
- The effectiveness of culturally adapted treatments for common mental disorders in people of Chinese descent, including those living in mainland China and in other countries as ethnic minorities (Li et al. 2023).
- Addressing the question of how Gypsy, Roma and Traveller populations access healthcare and what are the best ways to enhance their engagement with health services (including mental health services), in various European countries and Canada, with about half of the evidence conducted in or including the UK (McFadden et al. 2018).
- Improving pathways to mental health care for adults and young people from black and ethnic minority groups in the UK (Moffat et al. 2009).
- Interventions to improve engagement with mental health services (initiation, participation, completion and service utilisation) among adolescents and young adults from underserved minority groups in the USA (Moore 2018).
- Initiatives to enhance pathways to mental health care for adults from black and ethnic minority populations across, UK, Australia and USA (Sass et al. 2009).
- The effectiveness and acceptability of community-based interventions based on randomised controlled trials (RCTs) only in comparison with controls (no treatment, waiting list, alternative treatment) for preventing and treating mental health problems (major depression, anxiety, PTSD, psychological distress) and improving mental health in refugee children and adolescents in high-income countries (Soltan et al 2022).

- Determining the efficacy of culturally adapted psychosocial interventions in comparison to generic treatment for young people with refugee or asylum-seeker status in the UK (Taylor et al. 2023)

The scoping reviews focused on:

- Health promotion, prevention, and non-medical treatment interventions targeting the mental health and mental wellbeing of migrants and ethnic minority groups in Europe (Apers et al. 2023).
- To identify and evaluate intervention components and strategies that create culturally competent healthcare (including mental health care) for culturally and linguistically diverse people and to develop a model of culturally competent healthcare provision (Handtke et al. 2019).
- The accessibility (affordability, availability/accommodation, appropriateness and acceptability) of virtual mental healthcare interventions and assessments for refugee and immigrant groups of all ages globally across multiple countries and settings (Hynie et al. 2023).
- Mapping the interventions to increase migrants' care-seeking behaviour in high-income countries for stigmatised conditions with a primary focus on mental health (Place et al. 2021).
- Mapping the types and characteristics of approaches and interventions that non-medical local immigrant settlement organisations undertake to support access to primary healthcare and mental health services for immigrants over 16 years of age in high-income countries (Canada, USA, Australia) (Ratnayake et al. 2022).
- Models of care in pregnancy and the first 12 months postpartum in migrant and refugee women living in high income countries, including but not limited to mental health care (Rogers et al. 2020).
- Barriers to immigrant populations utilisation of mental health services in Canada and analysis of policy and practice recommendations (Thomson et al. 2015).

The mapping review explored interventions that aim to reduce mental health inequalities, their cost-effectiveness and the barriers and facilitators affecting them (Arundell et al. 2020). This mapping review included populations from multiple protected characteristics, including race, religion and belief (Arundell et al. 2020).

The narrative reviews focused on:

- Determining the effectiveness of culturally adapted mental health interventions by using meta-analysis, although the methods of literature reviewing did not resemble a standard systematic review. The countries where the research was conducted were not reported (Griner and Smith 2006).
- Critically examining the literature on culturally sensitive treatments for depression. The countries where the research was conducted were not reported (Kalibatseva and Leong 2014).
- Reasons for underutilisation of dementia services by minority ethnic groups and the interventions that have been developed to address access issues. The countries where the research was conducted were not reported (Mukadam et al. 2013).

The overviews of reviews focused on:

- Summarising systematic and literature reviews with meta-analyses which aimed to determine the effectiveness of cultural adaptations of mental health interventions. The countries where the research was conducted were not reported (Rathod et al. 2018).
- Mapping the characteristics and methodological quality of existing systematic reviews and registered systematic review protocols on the promotion of mental health and prevention and treatment of common mental disorders among refugees, asylum seekers, and internally displaced persons. The countries where the research was conducted were not reported (Uphoff et al. 2020).

The objective of the scoping review of reviews was to appraise the existing literature to identify key elements, conceptualisations, and interventions of cultural safety to improve health services and dementia care for Indigenous people from Canada, Australia and New Zealand (Chakanyuka et al. 2022).

The systematic review protocols focused on:

- Barriers and facilitators to the uptake of psychosocial intervention that are delivered by lay therapists to improve asylum seekers' and migrants' mental health and wellbeing (Aslam et al. 2018).
- Culturally sensitive interventions for black men to improve access to mental health care and/or mental health outcomes in the community (Stockwell and Roche 2022).
- Community-based interventions used to address Asian Americans' mental health challenges (Wang and Kim 2019).
- Community-level interventions for people from minority ethnic groups to improve their access to primary care when they experience a first-episode psychosis (Wood et al. 2023).

3.1.2 Summary of organisational reports and existing reviews identified via stakeholder group

These explored the following:

- The rapid review aimed to provide up-to-date evidence on accessibility, appropriateness, acceptability and outcomes of psychological interventions for people from Black, Asian and minority ethnic communities and to provide evidence of both good practice and barriers to good practice for all ages groups in the UK (Kunorubwe et al. 2023).
- The scoping review explored barriers and enablers of mental health care for asylum seekers and refugees residing in the UK (Pollard and Howard 2021).
- The systematic review investigated the barriers to mental health care utilisation and access among refugees and asylum seekers in Europe (Satinsky et al. 2019).
- The scope of the evidence on the management of PTSD for refugees and asylum seekers, with a focus on psychological interventions (Roberts 2023).
- A scoping exercise identified good practice in Wales and across the UK with regard to access to psychological therapies and mental health services for Asylum Seekers, Refugees and Migrants in the UK (Traumatic Stress Wales 2023).

- A qualitative primary research report regarding the barriers and opportunities for improving access to mental health support for refugees and people seeking asylum in England (VCSE Health and Wellbeing Alliance 2022).
- A qualitative primary research report into “Good Access” in Community Pharmacy, NHS Dentistry and Allied Health Professional Services which included barriers and facilitators to access (Welsh Government 2023).

3.2. Key Findings

The identified literature was categorised based on the aspect of mental health care access and provision it investigated. The categories are: Access and pathways to mental health care; Mental health promotion and prevention, and treatment of common mental disorders; Cultural adaptations to psychological interventions; Help seeking behaviour; Treatment engagement; and Treatment initiation, participation or continuation. The key findings of the identified literature are briefly summarised below under each category. The findings relating to Access and pathways to mental health care for minority groups are reported separately for ethnic minority groups; Gypsy, Roma and travelling communities; and migrants, refugees and asylum seekers. Key findings from the literature identified by the stakeholders were narratively summarised if the reports focused on interventions aiming to improve access to services or treatment modalities.

Access and pathways to mental health care for ethnic minority groups

- Barriers to **South Asian older adults’** access to dementia care include insufficient knowledge about dementia, mental illness, and local services, stigma, culturally preferred coping strategies, and linguistic and cultural difficulties in communication. **Suggestions were provided that could help improve access** for South Asian older adults and these included better information provision, culturally tailored services, employment of healthcare staff with similar cultural background, and education of healthcare professionals about dementia and local services (Giebel et al. 2015).
- The key components of **effective pathway interventions to mental health care for black and ethnic minority groups** included specialist services for ethnic minority groups, collaboration between sectors, facilitating referral routes between services, outreach and facilitating access into care, and supporting access to rehabilitation and moving out of care (Moffat et al. 2009).
- Sass et al. (2009) reported evidence of interventions with **black and ethnic minority populations** that led to three types of pathway modification (accelerated transit through care pathways, removal of adverse pathways, the addition of beneficial pathways). It was found that ethnic matching promoted desired pathways in some ethnic groups, that managed care improved equity, that a pre-treatment service improved access to detoxification, and that an education leaflet increased recovery.

Access and pathways to mental health care for people from Gypsy, Roma and Traveller communities

- The barriers to health services (including mental health services) reported by people from **Gypsy, Roma and Traveller communities** included the organisation of health systems, discrimination, culture and language, health literacy, service-user attributes, and economic barriers. Strategies thought to improve engagement included specialist roles, outreach services, dedicated services, raising health awareness, handheld records, training for staff, and collaborative working (McFadden et al. 2018).

Access and pathways to mental health care for refugees and asylum seekers

- The scoping review by Hynie et al. (2023) described factors affecting accessibility of **virtual mental health interventions for refugee and immigrant groups** which were individual (e.g., literacy), program related (e.g., computer needed), and/or contextual/social (e.g., housing characteristics, internet bandwidth). Need for financial and technical support were identified as potential barriers to refugee and immigrant groups' access.
- The interventions provided to **refugees and asylum seekers** within the broad primary care setting to improve the quality of health care, spanning mental health (n=11) and general wellbeing (n=44) could be organized into four categories, for example those that focused on developing the skills of individual refugees/asylum seekers and their families (**including promoting access** and improving engagement with and adherence to health regimes); skills of primary health care workers; system and/or service integration models and structures; and lastly, interventions enhancing communication services (Iqbal et al. 2022).
- Findings from the scoping review by Ratnayake et al. (2022) showed that local non-medical **immigrant** settlement organisations had established approaches and interventions to support immigrants **access to primary healthcare services**, with most studies showing that mental health support was an important component. These included connecting to healthcare services and / or collaborating with health sector institutions; providing health promotion programs; undertaking community capacity-building and policy advocacy activities and providing on the ground assistance.
- The scoping review by Rogers et al. (2020) identified a range of **interventions that improved access to maternity and postpartum care for migrant and refugee women**, including bilingual/bicultural workers, group antenatal care, and specialised clinics for care in pregnancy or first 12 months postpartum. **Four** of the 17 included studies had a mental health focus but these did not report separate findings.
- Barriers to **immigrant populations'** utilisation of mental health services in Canada included the uptake of health information, the process of immigrant settlement, and service availability. **Recommendations to improve access** included linguistically and culturally sensitive services, provision of translators and training for mental health workers in cultural and language competencies, practitioners considering patients' circumstances and pre-immigration experiences, diverse and socially inclusive workforce, and involvement of the community among other suggestions (Thomson et al. 2015).

Mental health promotion, prevention and treatment of mental health conditions including cultural adaptations to psychological interventions

- The review by Anderson et al. (2015) found one high-quality study which was a cluster-randomised trial that implemented **depression care quality improvement** in a network of mental health and health and social care systems (primary care, substance abuse, social services, and homeless services) in the ethnically diverse South Los Angeles and Hollywood metropolitan area, USA. That study found added benefit of a community coalition-driven intervention⁴ for improved mental health among **African Americans**.

⁴ Community coalition-driven interventions involve using “coalitions that include representatives of target populations to plan and implement interventions for community level change.” (Anderson et al. 2015, p. 1)

- The scoping review by Apers et al. (2023) identified three main **successful mechanisms for intervention development and implementation** with regard to the **prevention and promotion** of mental health and wellbeing, secondary prevention and tertiary prevention for **migrants and ethnic minority** groups: a sound theory-base, systematic adaption to make interventions culturally sensitive and participatory approaches.
- The mapping review reported that 80% of their identified studies aimed at improving inequalities in people from lower socioeconomic groups, diverse age groups or **ethnic minority and indigenous/aboriginal communities**. The interventions identified included **psychological support, delivering education and training, engaging the community and other culturally adapted interventions** (Arundell et al. 2020).
- **Cultural adaptations** made to psychological interventions for **ethnic minority groups** with mental health problems appear to be efficacious relative to non-adapted or waitlist/no intervention comparators. Adaptions included adapting the form used to provide interventions, the time or length of the intervention, the location of treatment provision, and the method of access (Arundell et al. 2021).
- The review by Bhui et al. (2015) reports on beneficial interventions to improve **therapeutic communications** for **black and minority ethnic people** receiving specialist psychiatric care. The interventions with evidence of benefit were culturally adapted psychotherapies (cognitive-behavioural therapy and family therapies); ethnographic and motivational interviewing; communications skills training; community-based stepped care and case finding by including social venues in the care pathway; role induction and education for patients and telepsychiatry that included ethnic matching.
- Findings from a meta-analysis indicate that **cultural adaptation of mindfulness-based interventions** for **Hispanic populations** might improve depressive symptoms and stress and could help with managing a chronic illness (Castellanos et al. 2019). A moderate to large effect could also be detected on psychiatric distress. However, the authors conclude that more high quality research studies were needed.
- The scoping review of reviews found little evidence that reported on interventions that aimed to improve **cultural safety in health and dementia care** for **Indigenous people** (Chakanyuka et al. 2022). There was also a lack of research focusing on implementation and evaluation of cultural safety intervention, and using the concept of sex and gender (Chakanyuka et al. 2022).
- Psychosocial interventions for schizophrenia can be culturally adapted based on language, concepts and illness models, family, communication, content, cultural norms and practices, context and delivery, therapeutic alliance, and treatment goals. Findings from the meta-analysis of **culturally adapted psychosocial interventions** for **people from non-Western countries** imply reduced symptom severity, although change was proportionate to level of adaptation. Nevertheless, there is a lack of well-designed high quality studies comparing adapted interventions to non-adapted (Degnan et al. 2018).
- **Culturally adapted digital mental health interventions** for **ethnic / racial minorities** have been found effective and acceptable in outcomes such as, mental health symptomatology, behaviours, self-efficacy, or wellness, although there are a lack of studies focusing on feasibility, active comparison treatments or on Black and Indigenous groups (Ellis et al. 2022).
- A narrative review containing a meta-analysis indicates that **cultural adaptations of mental health interventions to specific populations** are four times more effective

than interventions aimed at mixed cultural or ethnic groups. In addition, intervention adapted to patients' mother tongue were twice as effective as English language treatments (Griner and Smith 2006).

- A scoping review focusing on **culturally competent healthcare** (including mental health care) for **culturally and linguistically diverse people** identified 20 categories of different strategies, which could be organised into four groups: Components of culturally competent healthcare–Individual level; Components of culturally competent healthcare–Organizational level; Strategies to implement culturally competent healthcare; and Strategies to provide access to culturally competent healthcare (Handtke et al. 2019). Both qualitative and quantitative findings indicate positive patient and service utilization outcomes as a result of culturally competent healthcare strategies, although the effect was often small or not statistically significant (Handtke et al. 2019).
- A meta-analysis of **culturally adapted substance use interventions** indicates that the substance use outcomes of **Latino adolescents** in the USA are slightly improved, although the quality of included studies was low (Hernandez Robles et al. 2018).
- **Culturally sensitive treatments for people with depression** can contain general or practical adaptations, including translation of written materials or incorporating cultural values (Kalibatseva and Leong 2014).
- A rapid review reports that both **culturally sensitive and culturally adapted psychological interventions** have shown positive results in terms of accessibility, appropriateness, acceptability and outcomes amongst **Black, Asian and minority ethnic communities**. It is of note however, that there is still limited research on the efficacy of such interventions across distinct cultural communities. The literature also supports the utilisation of more community based collaborative approaches to service development to remove access barriers and improve outcomes (Kunorubwe et al. 2023).
- **Culturally adapted interventions for Chinese descents with common mental disorder** have been found moderately effective in reducing symptom severity, although no difference was detected between culturally modified and culturally specific interventions. However, the quality of included studies could not be determined due to issues with reporting (Li et al. 2023).
- An overview of reviews focusing on **cultural adaptation of mental health interventions** highlighted that there is a lack of consensus regarding what components of adaptations work for which cultural groups. Additionally, it is unclear whether cultural adaptations truly result in improved outcomes compared to non-adapted interventions (Rathod et al. 2018).
- Roberts (2023) scoped the evidence on the **management of PTSD for refugees and asylum seekers**. The author reported on five RCTs, a network meta-analysis (n=23 studies) of psychosocial interventions for refugees and asylum seekers with PTSD (Turrini et al. 2021) and an individual-patient data meta-analysis of residual PTSD symptoms after provision of brief behavioural intervention in low- and middle-income countries (Akhtar et al. 2022). Cognitive behavioural therapy and Eye movement desensitisation and reprocessing were found significantly more effective in reducing PTSD symptoms than waitlist comparison (Turrini et al. 2021). Problem Management Plus⁵ intervention was also observed to be more effective for symptom reduction than

⁵ Problem Management Plus is “a transdiagnostic intervention that involves five sessions that teach people skills” (Akhtar et al. 2021, p. 72)

enhanced treatment as usual, although residual symptoms were reported for 30% of refugee and asylum seeker participants (Akhtar et al. 2022).

- There was no evidence of an effect of **community-based interventions** when compared with a waiting list for symptoms of PTSD, depression, and psychological distress in **refugee children and adolescents** in high-income countries (Soltan et al. 2022).
- A systematic review investigating **cultural adaptations** to psychosocial interventions for **young people with refugee or asylum-seeker status** found that these type of interventions may have some positive impact on symptom reduction, although results varied. However, the available evidence is insufficient to determine effectiveness of cultural adaptations compared to treatment as usual or to investigate which components of cultural adaptation work better, as study designs were mainly qualitative or observational (Taylor et al. 2023).
- The overview of reviews mapped the evidence-base and found that there were gaps in the literature regarding mental health interventions for refugees, asylum seekers and internally displaced persons. Existing systematic reviews focus more attention on the treatment of PTSD than mental health promotion or prevention, or the treatment of depression or anxiety. Studies of Cognitive Behavioural Therapy, Narrative Exposure Therapy, and integrative and interpersonal therapies were most likely to be included in reviews (Uphoff et al 2020).

Help seeking behaviour

- Educational campaigns to increase awareness and reduce stigma of dementia for **minority ethnic groups** have been described in the literature, although outcomes of these interventions have not been reported. However, other studies suggest that information targeted to specific ethnic groups has the potential to increase knowledge about dementia and consequently help-seeking (Mukadam et al. 2013).
- The scoping review by Place et al. (2021) identified three approaches that increased **migrants'** mental health care-seeking behaviour: health communication (n=9); support groups (n = 2); and primary care-based approaches (n=4), including a self-assessment tool for psychosocial risk, and engagement interventions seeking to increase collaboration between primary care and mental health services.

Treatment engagement

- The review by Interian et al. (2013) found that collaborative care for depression was efficacious as an engagement improvement intervention for **underserved racial-ethnic groups**.

Interventions that were family based or were culturally adapted for age group or race-ethnicity showed possible efficacy and promising results for improving treatment engagement among **underserved minority adolescents and young adults** (Moore 2018).

Treatment initiation, participation or continuation

- The review by Aggarwal et al. (2016) did not find any RCTs of interventions to target specific mechanisms of action for improving patient-clinical communication regarding treatment initiation, participation or continuation with **racial/ethnic minorities**. Although some studies were found that reported on clinician experiences to improve patient-clinician communication.
- Interventions to improve the initiation of mental health care among **racial-ethnic minority groups** included collaborative care (n=10), psychoeducation (n=7), case

management (n=5), co-location of mental health services within existing services (n=4), screening and referral (n=2) and a change of Medicare medication reimbursement policy (n=1). An increased uptake of psychotherapy or antidepressant use among members of racial-ethnic minority was observed across interventions involving either collaborative care, co-location of mental health services, and screening and referral interventions (Lee-Tauler et al. 2018).

3.3. Areas of uncertainty

- There is a wealth of review and overview of review evidence that explores the barriers and facilitators to accessing mental healthcare for ethnic minority groups, refugees and asylum seekers across the UK and internationally. However, these often do not have an evaluative interventions and report their findings as suggestions / recommendations for interventions.
- There is a wealth of review evidence that covers mental health promotion, prevention and treatment of mental health conditions (including treatment initiation, participation or continuation) for ethnic minority groups, refugees and asylum seekers across the UK and internationally.
- There is a wealth of review and overview of review evidence focusing on cultural adaptations of psychological interventions whether it is for ethnic minority groups living in Western countries or majority populations living in non-Western countries.
- There is limited review evidence regarding the effectiveness of interventions to improve access to mental health care across ethnic minority groups.
- There is wider review evidence regarding interventions to support refugees and asylum seekers access to primary healthcare services and specialised clinics for care in pregnancy and postpartum. However these reviews did not separate out the findings for the studies that focused on mental health.

4. NEXT STEPS

The findings of this Rapid Evidence Summary were shared with the stakeholders and used to inform our decision on a substantive focus for a subsequent Rapid Review (RR). It was decided that the RR would focus on interventions to support equitable access to mental health services for ethnic minority groups.

5. REFERENCES

Aggarwal NK, Pieh MC, Dixon L, et al. (2016). Clinician descriptions of communication strategies to improve treatment engagement by racial/ethnic minorities in mental health services: A systematic review. *Patient Education and Counselling*. 99(2): 198-209. <https://doi.org/10.1016/j.pec.2015.09.002>

Akhtar A, Koyiet P, Rahman A, et al. (2022). Residual posttraumatic stress disorder symptoms after provision of brief behavioral intervention in low- and middle-income countries: An individual-patient data meta-analysis. *Depression and Anxiety*. 39(1): 71-82. <https://doi.org/10.1002/da.23221>

Anderson LM, Adeney KL, Shinn C, et al. (2015). Community coalition-driven interventions to reduce health disparities among racial and ethnic minority populations. *Cochrane Database of Systematic Reviews*. 2015(6): CD009905. <https://doi.org/10.1002/14651858.CD009905.pub2>

Apers H, Van Praag L, Nostlinger C, et al. (2023). Interventions to improve the mental health or mental well-being of migrants and ethnic minority groups in Europe:

A scoping review. *Global Mental Health*. 10(2023): e23.

<https://doi.org/10.1017/gmh.2023.15>

Arundell L-L, Barnett P, Buckman JEJ, et al. (2021). The effectiveness of adapted psychological interventions for people from ethnic minority groups: A systematic review and conceptual typology. *Clinical Psychology Review*. 88: 102063.

<https://doi.org/10.1016/j.cpr.2021.102063>

Arundell L-L, Greenwood H, Baldwin H, et al. (2020). Advancing mental health equality: a mapping review of interventions, economic evaluations and barriers and facilitators. *Systematic reviews*. 9(1): 115. <https://doi.org/10.1186/s13643-020-01333-6>

Aslam RW, Dowrick C, Hagan R, et al. (2018). Barriers and facilitators to uptake of psychosocial interventions delivered by lay therapists to improve mental health and wellbeing of asylum seekers and migrants: a systematic review. 2018

CRD42018104453 Available at:

https://www.crd.york.ac.uk/PROSPERO/display_record.php?ID=CRD42018104453 [Accessed.

Bhui KS, Aslam RW, Palinski A, et al. (2015). Interventions to improve therapeutic communications between Black and minority ethnic patients and professionals in psychiatric services: systematic review. *The British Journal of Psychiatry* 207(2): 95-103. <https://doi.org/10.1192/bjp.bp.114.158899>

Bignall T, Jeraj S, Helsby E, et al. (2019). Racial disparities in mental health: Literature and evidence review. Race Equality Foundation. Available at:

<https://raceequalityfoundation.org.uk/wp-content/uploads/2022/10/mental-health-report-v5-2-2.pdf>.

Braveman P, Arkin E, Orleans T, et al. (2018). What is Health Equity? *Behavioral Science & Policy*. 4(1): 1-14. <https://doi.org/10.1177/237946151800400102>

Campbell F, Tricco AC, Munn Z, et al. (2023). Mapping reviews, scoping reviews, and evidence and gap maps (EGMs): the same but different— the “Big Picture” review family. *Systematic reviews*. 12(1): 45. <https://doi.org/10.1186/s13643-023-02178-5>

Castellanos R, Spinel MY, Phan V, et al. (2020). A systematic review and meta-analysis of cultural adaptations of mindfulness-based interventions for Hispanic populations. *Mindfulness*. 11(2): 317-32. <https://dx.doi.org/10.1007/s12671-019-01210-x>

Centre for Mental Health. (2020). Commission for Equality in Mental Health -

BRIEFING 1: Determinants of mental health. Centre for Mental Health. Available at:

<https://www.eatq.org/wp-content/uploads/2022/07/determinants-for-mental-health.pdf> [Accessed 16/01/2023].

Chakanyuka C, Bacsu JR, DesRoches A, et al. (2022). Indigenous-specific cultural safety within health and dementia care: a scoping review of reviews. *Social Science & Medicine*. 293: 114658. <https://doi.org/10.1016/j.socscimed.2021.114658>

Cochrane Library. (2024). What is a systematic review? Available at:

<https://www.cochranelibrary.com/about/about-cochrane-reviews> [Accessed 01/02/2024].

Degnan A, Baker S, Edge D, et al. (2018). The nature and efficacy of culturally-adapted psychosocial interventions for schizophrenia: A systematic review and meta-analysis. *Psychological medicine*. 48(5): 714-27.

<https://dx.doi.org/10.1017/S0033291717002264>

- Ellis DM, Draheim AA, Anderson PL. (2022). Culturally adapted digital mental health interventions for ethnic/racial minorities: A systematic review and meta-analysis. *Journal of Consulting and Clinical Psychology*. 90(10): 717-33. <https://dx.doi.org/10.1037/ccp0000759>
- Gask L, Bower P, Lamb J, et al. (2012). Improving access to psychosocial interventions for common mental health problems in the United Kingdom: narrative review and development of a conceptual model for complex interventions. *BMC Health Serv Res*. 12: 249. <https://doi.org/10.1186/1472-6963-12-249>
- Giebel CM, Zubair M, Jolley D, et al. (2015). South Asian older adults with memory impairment: improving assessment and access to dementia care. *International journal of geriatric psychiatry*. 30(4): 345-56. <https://doi.org/10.1002/gps.4242>
- Griner D, Smith TB. (2006). Culturally adapted mental health intervention: A meta-analytic review. *Psychotherapy: Theory, Research, Practice, Training*. 43(4): 531-48. <https://dx.doi.org/10.1037/0033-3204.43.4.531>
- Handtke O, Schilgen B, Mösko M. (2019). Culturally competent healthcare – A scoping review of strategies implemented in healthcare organizations and a model of culturally competent healthcare provision. *PloS one*. 14(7): e0219971. <https://doi.org/10.1371/journal.pone.0219971>
- Hernandez Robles E, Maynard BR, Salas-Wright CP, et al. (2018). Culturally adapted substance use interventions for Latino adolescents: A systematic review and meta-analysis. *Research on Social Work Practice*. 28(7): 789-801. <https://dx.doi.org/10.1177/1049731516676601>
- Hynie M, Oda A, Calaresu M, et al. (2023). Access to virtual mental healthcare and support for refugee and immigrant groups: a scoping review. *Journal of Immigrant and Minority Health*. 25(5): 1171-95. doi: <https://doi.org/10.1007/s10903-023-01521-1>
- Interian A, Lewis-Fernandez R, Dixon LB. (2013). Improving treatment engagement of underserved U.S. racial-ethnic groups: a review of recent interventions. *Psychiatric Services*. 64(3): 212-22. <https://doi.org/10.1176/appi.ps.201100136>
- Iqbal M, Walpola R, Harris-Roxas B, et al. (2022). Improving primary health care quality for refugees and asylum seekers: A systematic review of interventional approaches. *Health Expectations*. 25(5): 2065-94. <https://doi.org/10.1111/hex.13365>
- Johnson L, Clinton Frazee C, Peterson DC, et al. (2020). Chapter 15.1 - A death involving extremely high levels of 3,4-methylenedioxymethamphetamine. In: Ketha H & Garg U (eds.) *Toxicology Cases for the Clinical and Forensic Laboratory*. Academic Press.
- Kalibatseva Z, Leong FT. (2014). A critical review of culturally sensitive treatments for depression: recommendations for intervention and research. *Psychol Serv*. 11(4): 433-50. <https://doi.org/10.1037/a0036047>
- Kunorubwe T, Tyson PJ, Molina J, et al. (2023). Guidance and an associated action plan to improve access to, and provision of, psychological interventions to people from Black, Asian and minority ethnic communities Public Health Wales and University of South Wales.
- Lee-Tauler SY, Eun J, Corbett D, et al. (2018). A systematic review of interventions to improve initiation of mental health care among racial-ethnic minority groups. *Psychiatric Services*. 69(6): 628-47. <https://doi.org/10.1176/appi.ps.201700382>
- Li S, Xi Z, Barnett P, et al. (2023). Efficacy of culturally adapted interventions for common mental disorders in people of Chinese descent: A systematic review and meta-analysis. *The Lancet Psychiatry*. 10(6): 426-40. <https://dx.doi.org/10.1016/S2215-0366%2823%2900118-9>

- Lock K, Puntan L, Lewis-Richards M. (2023). Review of Together for Mental Health and Talk to Me 2 Strategies. Welsh Government. Available at: <https://www.gov.wales/sites/default/files/statistics-and-research/2023-03/review-of-together-for-mental-health-and-talk-to-me-2-strategies.pdf> [Accessed 07/12/2023].
- Lowther-Payne HJ, Ushakova A, Beckwith A, et al. (2023). Understanding inequalities in access to adult mental health services in the UK: a systematic mapping review. *BMC Health Serv Res.* 23(1): 1042. <https://doi.org/10.1186/s12913-023-10030-8>
- McFadden A, Siebelt L, Gavine A, et al. (2018). Gypsy, Roma and Traveller access to and engagement with health services: a systematic review. *European Journal of Public Health.* 28(1): 74-81. <https://doi.org/10.1093/eurpub/ckx226>
- Moffat J, Sass B, McKenzie K, et al. (2009). Improving pathways into mental health care for black and ethnic minority groups: a systematic review of the grey literature. *International Review of Psychiatry.* 21(5): 439-49. <https://doi.org/10.1080/09540260802204105>
- Moore KL. (2018). Mental health service engagement among underserved minority adolescents and young adults: a systematic review. *Journal of racial and ethnic health disparities.* 5(5): 1063-76. <https://doi.org/10.1007/s40615-017-0455-9>
- Mukadam N, Cooper C, Livingston G. (2013). Improving access to dementia services for people from minority ethnic groups. *Current Opinions in Psychiatry.* 26(4): 409-14. <https://doi.org/10.1097/YCO.0b013e32835ee668>
- Munn Z, Pollock D, Khalil H, et al. (2022). What are scoping reviews? Providing a formal definition of scoping reviews as a type of evidence synthesis. *JBI Evidence Synthesis.* 20(4).
- ONS. (2022). Ethnic group, England and Wales: Census 2021. London: Office for National Statistics. Available at: <https://www.ons.gov.uk/peoplepopulationandcommunity/culturalidentity/ethnicity/bulletins/ethnicgroupenglandandwales/census2021#main-points> [Accessed 16/01/2024].
- ONS. (2023). Gypsy or Irish Traveller populations, England and Wales: Census 2021. London: Office for National Statistics. Available at: <https://www.ons.gov.uk/peoplepopulationandcommunity/culturalidentity/ethnicity/articles/gypsyoririshtravellerpopulationsenglandandwales/census2021> [Accessed 04/12/2023].
- Place V, Nabb B, K. VA, et al. (2021). Interventions to increase migrants' care-seeking behaviour for stigmatised conditions: a scoping review. *Social psychiatry and psychiatric epidemiology.* 56(6): 913-30. <https://doi.org/10.1007/s00127-021-02065-1>
- Pollard T, Howard N. (2021). Mental healthcare for asylum-seekers and refugees residing in the United Kingdom: a scoping review of policies, barriers, and enablers. *International Journal of Mental Health Systems.* 15(1): 60. <https://doi.org/10.1186/s13033-021-00473-z>
- Pollock M, Fernandes RM, Becker LA, et al. (2023). Chapter V: Overviews of Reviews. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ & Welch VA (eds.) *Cochrane Handbook for Systematic Reviews of Interventions.* Version 6.4 ed.: Cochrane.
- Rathod S, Gega L, Degnan A, et al. (2018). The current status of culturally adapted mental health interventions: A practice-focused review of meta-analyses. *Neuropsychiatric Disease and Treatment* Vol 14, 2018, ArtID 165-178. 14. <https://dx.doi.org/10.2147/NDT.S138430>

- Ratnayake A, Sayfi S, Veronis L, et al. (2022). How are non-medical settlement service organizations supporting access to healthcare and mental health services for immigrants: a scoping review. *International journal of environmental research and public health*. 19(6). <https://doi.org/10.3390/ijerph19063616>
- Roberts N. (2023). Psychological interventions for refugees and asylum seekers: current evidence. Cardiff & Vale Univeristy Health Board & Cardiff University.
- Rogers HJ, Hogan L, Coates D, et al. (2020). Responding to the health needs of women from migrant and refugee background - models of maternity and postpartum care in high-income countries: a systematic scoping review. *Health & social care in the community*. 28(5): 1343-65. <https://doi.org/10.1111/hsc.12950>
- Sass B, Moffat J, Bhui K, et al. (2009). Enhancing pathways to care for black and minority ethnic populations: a systematic review. *International Review of Psychiatry*. 21(5): 430-8. <https://doi.org/10.1080/09540260802204121>
- Satinsky E, Fuhr DC, Woodward A, et al. (2019). Mental health care utilisation and access among refugees and asylum seekers in Europe: a systematic review. *Health Policy*. 123(9): 851-63. <https://doi.org/10.1016/j.healthpol.2019.02.007>
- Soltan F, Cristofalo D, Marshall D, et al. (2022). Community-based interventions for improving mental health in refugee children and adolescents in high-income countries. *Cochrane Database of Systematic Reviews*. 5(5): CD013657. <https://doi.org/10.1002/14651858.CD013657.pub2>
- Stockwell D, Roche M. (2022). A systematic review of interventions that help black men access timely and appropriate support for their mental health. PROSPERO 2022 CRD42022345323. Available at: https://www.crd.york.ac.uk/prospERO/display_record.php?ID=CRD42022345323 [Accessed 07/12/2023].
- Taylor A, Radford G, Calia C. (2023). Review: cultural adaptations to psychosocial interventions for families with refugee/asylum-seeker status in the United Kingdom - a systematic review. *Child and adolescent mental health*. 28(2): 241-57. <https://doi.org/10.1111/camh.12547>
- Thomson MS, Chaze F, George U, et al. (2015). Improving immigrant populations' access to mental health services in Canada: a review of barriers and recommendations. *Journal of Immigrant and Minority Health*. 17(6): 1895-905. <https://doi.org/10.1007/s10903-015-0175-3>
- Traumatic Stress Wales. (2023). Draft Asylum Seekers, Refugees and Migrants Mental Health Services Working Group Proposal. Traumatic Stress Wales & Wales Strategic Migration Partnership.
- Tricco AC, Langlois EV, Straus SE, et al. (2017). Rapid reviews to strengthen health policy and systems: a practical guide, Geneva, World Health Organization.
- Turrini G, Tedeschi F, Cuijpers P, et al. (2021). A network meta-analysis of psychosocial interventions for refugees and asylum seekers with PTSD. *BMJ Glob Health*. 6(6). <https://doi.org/10.1136/bmjgh-2021-005029>
- UK Government. (2023a). Asylum and resettlement datasets. London: UK Government. Available at: <https://www.gov.uk/government/statistical-data-sets/asylum-and-resettlement-datasets#local-authority-data> [Accessed 24/01/2024].
- UK Government. (2023b). Writing about ethnicity. London: UK Government. Available at: <https://www.ethnicity-facts-figures.service.gov.uk/style-guide/writing-about-ethnicity/#ethnic-minorities-and-ethnic-groups> [Accessed 24/01/2024].
- UNHCR. (2024a). Asylum-seeker. United Nations High Commissioner for Refugees. Available at: <https://www.unhcr.org/uk/asylum-seekers> [Accessed 25/01/2024].

UNHCR. (2024b). Refugees. United Nations High Commissioner for Refugees. Available at: <https://www.unhcr.org/about-unhcr/who-we-protect/refugees> [Accessed 25/01/2024].

Uphoff E, Robertson L, Cabieses B, et al. (2020). An overview of systematic reviews on mental health promotion, prevention, and treatment of common mental disorders for refugees, asylum seekers, and internally displaced persons. *Cochrane Database of Systematic Reviews*. 9(9): CD013458. doi: 10.1002/14651858.CD013458.pub2

VCSE Health and Wellbeing Alliance. (2022). Barriers and Opportunities: Improving access to mental health support for refugees and people seeking asylum. Voluntary Community and Social Enterprise Health and Wellbeing Alliance. Available at: <https://www.redcross.org.uk/about-us/what-we-do/we-speak-up-for-change/vcse-health-and-wellbeing-alliance> [Accessed 07/12/2023].

Wang C, Lim K. (2019). A systematic review of community-based mental health interventions for Asian Americans. PROSPERO 2019 CRD42019136832 Available at: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42019136832 [Accessed 07/12/2023].

Welsh Government. (2012). Together for Mental Health - A Strategy for Mental Health and Wellbeing in Wales Welsh Government. Available at: <https://www.gov.wales/sites/default/files/publications/2019-03/together-for-mental-health-a-strategy-for-mental-health-and-wellbeing-in-wales.pdf> [Accessed 07/12/2023].

Welsh Government. (2014). Together for Mental Health Strategy - Equality Impact Assessment. Welsh Government. Available at: <https://www.gov.wales/sites/default/files/publications/2019-03/together-for-mental-health-strategy-equality-impact-assessment.pdf> [Accessed 07/12/2023].

Welsh Government. (2020). Review of the Together for Mental Health Delivery Plan 2019-2022 in response to Covid-19 Welsh Government. Available at: https://www.gov.wales/sites/default/files/publications/2020-10/review-of-the-together-for-mental-health-delivery-plan-20192022-in-response-to-covid-19_0.pdf [Accessed 07/12/2023].

Welsh Government. (2022). Anti-racist Wales Action Plan. Welsh Government. Available at: https://www.gov.wales/sites/default/files/publications/2022-06/anti-racist-wales-action-plan_0.pdf [Accessed 07/12/2023].

Welsh Government. (2023). Research into “Good Access” in Community Pharmacy, NHS Dentistry and Allied Health Professional Services. Welsh Government. Available at: <https://www.gov.wales/research-good-access-community-pharmacy-nhs-dentistry-and-allied-health-professional-services> [Accessed 07/12/2023].

Wood N, Oduola S, Hodgekins J. (2023). The impact of community-level interventions on improving access to primary care for people from minority ethnic groups experiencing a first-episode psychosis. PROSPERO 2023 CRD42023398682. PROSPERO. Available at: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42023398682 [Accessed 07/12/2023]

6. RAPID EVIDENCE SUMMARY METHODS

A list of the resources searched during this Rapid Evidence Summary is provided within the Appendix. Relevant research was also identified via the stakeholders. Searches were limited to English-language publications and did not include searches for primary studies. Search hits were screened for relevance by a single reviewer.

Priority was given to robust evidence synthesis using minimum standards (systematic search, study selection, quality assessment, appropriate synthesis). The secondary research identified was not formally quality assessed. The included research may vary considerably in quality and the degree of such variation could be investigated during rapid review work which may follow-on. Citation, recency, evidence type, document status and key findings were tabulated for all relevant secondary research identified in this process.

Date of Search	November 2023
Search Concepts Used	<p>Mental health (synonyms and related terms: psychological health, mental illness, bipolar, schizophrenia, psychosis, depression, anxiety, post-traumatic stress disorder, dementia) AND</p> <p>Minority groups (synonyms and related terms: gypsy, roma, traveller, refugee, asylum, ethnic, minority, racial, immigrant) AND</p> <p>Intervention (synonyms and related terms: improve) AND</p> <p>Access (synonyms and related terms: provision, service, engage)</p>
Search Completed by	Cardiff Evidence Synthesis Collaborative

7. EVIDENCE

Table 1: Summary of review evidence identified

Evidence type	Total identified
Systematic reviews	20
Scoping reviews	8
Mapping review	1
Rapid reviews	1
Narrative reviews	3
Overview of reviews	2
Scoping review of reviews	1
Protocols for reviews that are underway	4
Stakeholder organisational reports	4

A more detailed summary of included evidence can be found across Table 2 to 4.

Table 2: Summary of included secondary evidence

Resource	Citation	Recency (Search dates)	Evidence Type*	Status**	Key findings from abstracts	Reviewer comments
Access and pathways to mental health care for ethnic minority groups						
Google	Giebel et al. 2015 South Asian older adults with memory impairment: improving assessment and access to dementia care. <i>International Journal of Geriatric Psychiatry</i> ; 30(4):345-356 https://doi.org/10.1002/gps.4242	1984 to 2012	SR	Published	Included studies: n=18 Participants: Older people Countries: UK (n=15), USA (n=1), Canada (n=2) South Asians and health professionals highlighted several difficulties which deterred help seeking and access to care: a lack of knowledge of dementia and mental illness, and of local services; stigma; culturally preferred coping strategies; and linguistic and cultural barriers in communication and decision making To improve access for these groups, service users and providers need to be better informed; services need to be more culturally tailored, sometimes employing staff with similar cultural backgrounds; and health professionals can benefit from dementia education and knowledge of local services. These factors are key to the delivery of the National Dementia Strategy in England	The review focuses on barriers and facilitators, and does not seem to evaluate interventions improving access
TripPro PsycINFO	Moffat et al. 2009 Improving pathways into mental health care for black and ethnic minority groups : A systematic review of the grey literature. <i>International Review of Psychiatry</i> ; 21(5):439-49	Inception to April 2006	SR of grey literature	Published	Included studies: n=8 Participants: Young people and adults Countries: UK The key components of effective pathway interventions to mental health care include specialist services for ethnic minority groups, collaboration between sectors, facilitating referral routes between services, outreach and facilitating access into care,	Details of services and type of evaluation data are presented in Table II All services were based in the UK; seven related to London-based services and one to a service in Birmingham

	https://doi.org/10.1080/09540260802204105				and supporting access to rehabilitation and moving out of care Services that support collaboration, referral between services, and improve access seem effective, but warrant further evaluation	Disadvantaged Arabic speaking women, Difficult to engage African and African Caribbean people (aged 16–25 years), Chinese clients, an underprivileged multi-ethnic group, Isolated Asian people, and Vietnamese people
PsycINFO	Sass et al. 2009 Enhancing pathways to care for black and minority ethnic populations: a systematic review. <i>International Review of Psychiatry</i> ; 21(5):430-438 https://doi.org/10.1080/09540260802204121	1982 to 2006	SR	Published	Included studies: n=6 Participants: Adults Countries: USA (n=4), Australia (n=1), UK (n=1) Reviewed initiatives to enhance pathways to mental health care for black and minority ethnic groups Interventions led to three types of pathways change: -accelerated transit through care pathways, -removed adverse pathways -added a beneficial pathway Ethnic matching promoted desired pathways in many groups but not African Americans Managed care improved equity A pre-treatment service improved access to detoxification An education leaflet increased recovery	
Access and pathways to mental health care for people from Gypsy, Roma and Traveller communities						
Google	McFadden et al. 2018 Gypsy, Roma and Traveller access to and engagement with health services: a systematic review. <i>European</i>	2000-2015	SR	Published	Included studies: n=99 (mental health n=20) Participants: Gypsy, Roma or Traveller Countries: 32 countries (31 European countries and Canada / UK: n=49)	Includes different healthcare services (not only mental health)

	<i>Journal of Public Health</i> ; 28(1):74-81 https://doi.org/10.1093/eurpub/ckx226				Reported barriers to health service usage related to organisation of health systems, discrimination, culture and language, health literacy, service-user attributes and economic barriers Promising engagement strategies included specialist roles, outreach services, dedicated services, raising health awareness, handheld records, training for staff and collaborative working	
Access and pathways to mental health care for refugees and asylum seekers						
PsycINFO	Hynie et al. 2023 Access to virtual mental healthcare and support for refugee and immigrant groups: a scoping review. <i>Journal of Immigrant and Minority Health</i> ; 25: 1171-1195 https://doi.org/10.1007/s10903-023-01521-1	Inception – (December 2020 through October 2021)	ScR	Published	Included studies: n=44 (41 unique interventions / assessment tools) Participants: All ages Countries: Australia (n=7), Canada (n=5), Denmark (n=2), Germany (n=2), Lebanon (n=1), USA (n=12), Mexico (n=1), the Netherlands (n=1), South Africa (n=1), Spain (n=2), Sweden (n=4), Switzerland (n=1), Turkey (n=1), mixed (n=4) Examined the accessibility (affordability, availability/accommodation, appropriateness and acceptability) of virtual mental health interventions and assessments Accessibility depended on individual (e.g., literacy) program (e.g., computer required) and contextual/social factors (e.g., housing characteristics, internet bandwidth). Participation often required financial and technical support , raising important questions about accessibility for immigrant/refugee populations	Only ran ti, ab, kw searches. Limited to English Language only. No other limits in terms of publication dates, geography.
PsycINFO	Iqbal et al. 2022 Improving primary health care quality for refugees and asylum seekers: A systematic	Inception to 2 September 2020	SR	Published	Included studies: n=55, Mental Health: n=11 Participants: Refugees and asylum seekers in OECD (Organization for Economic Co-Operation and Development) countries	11 studies that talked about interventions related to mental health References 27,35,47,55,59,68,74,54,79,49, and 69

	<p>review of interventional approaches. <i>Health Expectations</i>; 25(5):2065-2094</p> <p>https://doi.org/10.1111/hex.13365</p>				<p>Countries: North America (USA and Canada, n=35), European Countries (including United Kingdom, the Netherlands, Germany, Sweden and Italy n=14) and Australia (n=6)</p> <p>The interventions within the broad primary care setting could be organized into four categories</p> <ul style="list-style-type: none"> -Developing the skills of individual refugees/asylum seekers and their families; -Skills of primary health care workers -System and/or service integration models and structures -Interventions enhancing communication services. 	
PsycINFO	<p>Ratnayake et al. 2022</p> <p>How are non-medical settlement service organizations supporting access to healthcare and mental health services for immigrants: a scoping review. <i>International Journal of Environmental Research and Public Health</i>; 19: 3616-</p> <p>https://doi.org/10.3390/ijerph19063616</p>	<p>1st May 2013 to 31st May 2021</p>	ScR	Published	<p>Included studies: n=10</p> <p>Participants: 16 years and older</p> <p>Countries: Canada (n=8), USA (n=1), Australia (n=1)</p> <p>The scoping review aims to identify and map the types and characteristics of approaches and interventions that non-medical immigrant settlement organisations undertake to support access to primary healthcare for clients. Organisations support access by collaborating with health and sector partners in the community, connecting clients to health services and service providers, advocating for immigrant health, providing educational programming and initiating community development/ mobilization and advocacy activities</p>	<p>Study undertaken in Canada. Eligibility criteria state that they were interested in high income, industrialised countries with characteristics comparable to those of Canada</p>
PsycINFO	<p>Rogers et al. 2020</p> <p>Responding to the health needs of women from migrant and refugee backgrounds- Models of maternity and</p>	<p>2008 to 2019</p>	ScR	Published	<p>Included studies: n=17 (mental health-related: n=4)</p> <p>Participants: Pregnant women and women during the first 12 months postpartum</p> <p>Countries: USA (n=11), Australia (n=3), Canada (n=1), Sweden (n=1), UK (n=1)</p>	<p>Scoping review. 11 databases and hand-searching of references</p> <p>Four studies included mental health topics, such as</p>

	postpartum care in high-income countries: A systematic scoping review. <i>Health & Social Care in the Community</i> ; 28(5):1343-1365 https://doi.org/10.1111/hsc.12950				A diverse range of interventions were identified, including bilingual/bicultural workers, group antenatal care and specialised clinics. All identified interventions were acceptable to women, and improved access Synthesis of the included studies indicates the key elements of acceptable and accessible models, which were as follows: culturally responsive care, continuity of care, effective communication, psychosocial and practical support, support to navigate systems, flexible and accessible services	psychoeducation to reduce perinatal depression, maternal anxiety/stress and a mental health support group Most of the interventions were qualitative (n=12) Results not reported separately for mental health component
PsycINFO	Thomson et al. 2015 Improving immigrant populations' access to mental health services in Canada: a review of barriers and recommendations. <i>Journal of Immigrant and Minority Health</i> ; 17(1):1895-1905 https://doi.org/10.1007/s10903-015-0175-3	January 1990 to August 2013	ScR	Published	Included studies: n=131 Participants: All ages Countries: Canada (n=131) Immigrants and refugees came from diverse religious and cultural backgrounds and had complex mental health-related concerns that were not currently being adequately addressed by existing services The major barriers to the utilization of mental health services included: those related to the uptake of existing health information and services; those that were related to the process of immigrant settlement; and barriers related to availability of appropriate services Recommendations to improve access: - Linguistically and culturally sensitive service - Mental health workers to be trained in cultural and language competency - Practitioners to consider context/ circumstances including pre-migration experiences, etc - Time to establish safe relationship with clients - Provision of translators and interpreters - Diverse and socially inclusive workforce - Involvement of community - Public education and media campaigns	This ScR is derived from a larger body of work conducted in Canada. The paper focuses on barriers of access to mental health services, although it claims they conducted "thematic analysis" on recommendations The recommendations for practice section was extracted for this table.

					<ul style="list-style-type: none"> - Early intervention services to monitor the pathways to care for young people to address any disparities in accessing care - Efforts especially for refugees that facilitate their ability to cope with ongoing acculturation stressors from traumatic experiences - Preventative methods directed at the cultural evaluation of the psychiatric patient, the special needs of the individual and early intervention for these high-risk groups to prevent suicidal tendencies among immigrants 	
Mental health promotion, prevention and treatment of mental health conditions including cultural adaptations to psychological interventions						
Cochrane	Anderson et al. 2015 Community coalition-driven interventions to reduce health disparities among racial and ethnic minority populations . <i>Cochrane Database of Systematic Reviews</i> . 2015 (6):CD009905. https://doi.org/10.1002/14651858.CD009905.pub2	January 1990 to through 30 September 2013	SR	Published	<p>Included studies: n=58</p> <p>Participants: All age groups (infants, children young people, adults and the elderly)</p> <p>Countries: USA (n=2), Australia (n=2), Canada (n=2), England (n=2), the Netherlands (n=2)</p> <p>Interventions led by community coalitions may connect health and human service providers with ethnic and racial minority communities in ways that benefit individual health outcomes and behaviours, as well as care delivery systems. However, need a better understanding of how a program does or does not work</p> <p>This review found only one high-quality study which was a cluster-randomized trial that implemented depression care quality improvement in a network of mental health and health and social care systems (primary care, substance abuse, social services, and homeless services) in the ethnically diverse South Los Angeles and Hollywood metropolitan area, USA. That study found added benefit of a coalition-driven intervention for improved mental health among African Americans</p>	
PsycINFO	Apers et al. 2023	Inception to 1 July 2022	ScR	Published	Included studies: n=27	

	<p>Interventions to improve the mental health or mental well-being of migrants and ethnic minority groups in Europe: A scoping review. <i>Global Mental Health</i>. 2023 10:e23.</p> <p>https://doi.org/10.1017/gmh.2023.15</p>				<p>Participants: Adults aged +18 years old, migrants and ethnic minority groups</p> <p>Countries: UK (n=13), Scotland (n=1), the Netherlands (n=5), Sweden (n=3), Ireland (n=1), mixed countries (n=4)</p> <p>Interventions: Health promotion, prevention, and non-medical treatment</p> <p>Focus: Mental health and wellbeing</p> <p>The majority of interventions showed a positive effect on participants' mental health, indicating the importance of using a tailored approach</p> <p>Three main successful mechanisms for intervention development and implementation: a sound theory-base, systematic adaption to make interventions culturally sensitive and participatory approaches</p>	
Backchaining	<p>Arundell et al. 2020</p> <p>Advancing mental health equality: a mapping review of interventions, economic evaluations and barriers and facilitators. <i>Systematic reviews</i> 9:115</p> <p>https://doi.org/10.1186/s13643-020-01333-6</p>	January 2008 to December 2018	Mapping review	Published	<p>Included studies: n=128</p> <p>Participants: All age groups (including children and young people and older adults). All protected characteristics were considered for inclusion, including age, disability, race, religion and belief, pregnancy, sex, gender reassignment, sexual orientation.</p> <p>Countries: USA (n = 44), UK (n = 28), Australia (n=14), Ireland (n=9), The Netherlands (n=4), Iran (n=3), Canada (n=3), India (n=2), Colombia (n=2), Austria (n=1), Belgium (n=1), China (n=1), France (n=1), Germany (n=1), Israel (n=1), Norway (n=1), Pakistan (n=1), Portugal (n=1), Spain (n=1), Chile (n=1), Ethiopia (n=1), Kenya (n=1), Sweden (n=1)</p> <p>The majority (80%) of primary studies focused on targeting populations based on socioeconomic factors (n=65), age (children and young people as well as older adults; n=46) and race/ethnicity (ethnic minorities and indigenous/ aboriginal populations; n=29), indicating that these</p>	This mapping review focused on tabulating the type of evidence, but did not report on the effectiveness of the identified interventions.

					<p>populations are most frequently targeted in interventions to address mental health inequalities in the published literature</p> <p>Very few primary studies targeting populations on the basis of religious affiliation (n=2), sexual or gender identity and sexual orientation (LGBTQ+; n=2) and disability (n=3), were identified and none of the systematic reviews targeted these populations</p> <p>Intervention strategies used most frequently in primary studies were providing psychological support (n = 45), delivering education and training (n = 29), engaging the community (n = 26) and other—culturally adapted interventions (n = 26). The most frequently reported intervention strategy in systematic reviews was other—culturally adapted interventions (n=5) followed by providing psychological support (n=4)</p>	
PROSPERO	<p>Arundell et al. 2021</p> <p>The effectiveness of adapted psychological interventions for people from ethnic minority groups: A systematic review and conceptual typology. <i>Clinical Psychology Review</i>. 88:102063</p> <p>https://doi.org/10.1016/j.cpr.2021.102063</p>	1965 to December 2020	SR	Published	<p>Included studies: n=88 (67 were RCTs)</p> <p>Participants: Adults 18> Black, ethnic minority, migrant, refugee or asylum seeker communities, and people referred to as 'minorities' or defined as belonging to an identified racial or ethnic 'minority group'</p> <p>Countries: Not reported</p> <p>Included RCTs identified a significant effect on symptom reduction when adapted interventions were compared to non-adapted active treatments (K = 30, Hedge's g = -0.43 [95% CI: -0.61, -0.25], p < .001)</p> <p>Studies often incorporated multiple adaptations, limiting the exploration of the comparative effectiveness of different adaptation types, although inclusion of organisation-specific adaptations may be associated with greater benefits</p>	

<p>PubMed HTA PsycINFO</p>	<p>Bhui et al. 2015 Interventions designed to improve therapeutic communications between black and minority ethnic people and professionals working in psychiatric services: a systematic review of the evidence for their effectiveness. <i>Health Technology Assessment</i> 19(31) https://doi.org/10.3310/hta19310</p>	<p>Inception to 4 February 2012</p>	<p>SR</p>	<p>Published</p>	<p>Included studies: n=21 Participants: Young people, adults and the elderly Countries: UK (n=3), USA (n=13), Canada (n=4), Germany (n=1) The interventions with evidence of benefit were culturally adapted psychotherapies (cognitive-behavioural therapy and family therapies); ethnographic and motivational interviewing; communications skills training; community-based stepped care and case finding by including social venues in the care pathway; role induction and education for patients and telepsychiatry that included ethnic matching</p>	<p>Authors noted that studies tended to have small sample sizes or were pilot studies, not direct measures for therapeutic communications Derivative publications Bhui et al. 2013 <i>Systematic reviews</i>. 2(15) https://dx.doi.org/10.1186/2046-4053-2-15 Bhui et al. 2015 <i>The British Journal of Psychiatry</i>. 207(2):95-103 https://dx.doi.org/10.1192/bjp.bp.114.158899</p>
<p>PsycINFO</p>	<p>Castellanos et al. 2019 A systematic review and meta-analysis of cultural adaptations of mindfulness-based interventions for Hispanic populations. <i>Mindfulness</i> 11(2):317-332 https://dx.doi.org/10.1007/s12671-019-01210-x</p>	<p>Inception to August 2017</p>	<p>SR</p>	<p>Published</p>	<p>Included studies: n=20 Participants: All age groups, Latino or Hispanic populations (including college students) Countries: USA (n=6), Spain (n=11), Chile (n=2), Colombia (n=1) Culturally adapted mindfulness-based interventions are associated with depression symptom improvement, stress reduction, stress management, and chronic illness management. Results from meta-analysis suggest a moderate to large effect of the interventions on psychiatric distress relative to scores in the comparison group Findings from this study suggest there is clear evidence that cultural adaptations can improve evidence-based treatment implementation among</p>	<p>None of the studies reported on ethnicity despite considerable ethnic heterogeneity among Hispanic populations in the USA.</p>

					Hispanics, but more methodologically rigorous studies are needed	
PsycINFO	Chakanyuka et al. 2022 Indigenous-specific cultural safety within health and dementia care: A scoping review of reviews. <i>Social Science & Medicine</i> 293:114658. https://doi.org/10.1016/j.socscimed.2021.114658	Jan 2010 to Dec 2020	ScR of reviews	Published	Included reviews: n=17 Participants: Indigenous people with dementia Countries: Canada (n=10), Australia (n=6), and New Zealand (n=1) Conclude that few reviews described specific interventions, implementation strategies, evaluation methods, or the concept of sex and gender to improve cultural safety in healthcare delivery	
PsycINFO	Degnan et al. 2018 The nature and efficacy of culturally-adapted psychosocial interventions for schizophrenia: A systematic review and meta-analysis. <i>Psychological Medicine</i> 48(5):714-727 https://dx.doi.org/10.1017/S0033291717002264	Inception to 3 March 201	SR	Published	Included studies: n=46 Participants: 18+ years with schizophrenia, interventions were adopted for majority populations (n=39), and minority populations (n=7) Countries: China (n=25), Pakistan (n=2), Taiwan (n=2), India (n=2), Iran (n=1), Saudi Arabia (n=1), Malaysia (n=1), USA (n=6), Mexico (n=2), Brazil (n=1), Italy (n=1), Australia (n=1), Egypt (n=1) Cultural adaptations were grouped into nine themes: language, concepts and illness models, family, communication, content, cultural norms and practices, context and delivery, therapeutic alliance, and treatment goals. The adaptation process can be described within a framework that serves as a benchmark for development or assessment of future adaptations Meta-analyses showed significant post-treatment effects in favour of adapted interventions for total symptom severity (n = 2345, g: -0.23, 95% confidence interval (CI) -0.36 to -0.09), positive (n = 1152, g: -0.56, 95% CI -0.86 to -0.26), negative (n = 855, g: -0.39, 95% CI -0.63 to -0.15), and general (n = 525, g: -0.75, CI -1.21 to -0.29) symptoms. Culturally adapted interventions were	Some of the included studies focused on adapting the format, delivery or content of an existing Western intervention (i.e. influenced by European culture, including Europe, USA, Canada and Australia) to meet the cultural needs of majority populations in countries other than Western countries.

					<p>more efficacious than usual treatment in proportion to the degree of adaptation.</p> <p>There is insufficient evidence to show that adapted interventions are better than non-adapted interventions. Features of context, intervention and design influenced efficacy. Investigating whether adaptation improves efficacy, most importantly amongst ethnic minorities, requires better designed trials with comparisons against unadapted interventions</p>	
PsycINFO	<p>Ellis et al. 2022</p> <p>Culturally adapted digital mental health interventions for ethnic/racial minorities: A systematic review and meta-analysis. <i>Journal of Consulting and Clinical Psychology</i> 90(10):717-733</p> <p>https://dx.doi.org/10.1037/ccp0000759</p>	<p>January 2000 and July 2021</p>	SR	Published	<p>Included studies: n=32</p> <p>Participants: Adults; Minority status was defined as a group with distinct racial, ethnic, or linguistic traditions that differ from that of the majority of the national population where the research was based or where the original evidence-based treatment was developed</p> <p>Countries: USA (n=7), Brazil (n=1), New Zealand (n=1), Dominican Republic (n=1), Australia (n=1), Chile (n=1), China (n=4), Lebanon (n=1), Germany (n=1), Mexico (n=2), Sweden (n=2), American Samoa (n=1), Colombia (n=1), The Netherlands (n=1), Mixed (n=7)</p> <p>Among eligible randomized controlled studies (n=12) comprising 653 participants, results indicated that culturally adapted digital mental health interventions produced a large, positive, significant effect ($g=0.90$) across a range of outcomes when compared to wait-list and treatment as usual control conditions. The average attrition rate per study was 42%, and most participants did not complete all modules despite reporting high satisfaction</p> <p>Culturally adapted digital mental health interventions are efficacious and acceptable. Such interventions represent a powerful opportunity to circumvent barriers to mental health treatment and improve mental health equity among racially and ethnically</p>	

					minoritized communities. However, the prevalence of feasibility studies, lack of active comparison treatments—and limited research for Black and Indigenous populations—indicate that more research is needed to achieve this purpose	
PsycINFO	Griner and Smith 2006 Culturally adapted mental health intervention: A meta-analytic review. <i>Psychotherapy: Theory, Research, Practice, Training</i> 43(4):531-548 https://dx.doi.org/10.1037/0033-3204.43.4.531	Inception to August 2004	NR	Published	Included studies: n=76 Participants: Age not reported, racial and ethnic groups Countries: Not reported Across 76 studies the resulting random effects weighted average effect size was $d=.45$, indicating a moderately strong benefit of culturally adapted interventions. Interventions targeted to a specific cultural group were four times more effective than interventions provided to groups consisting of clients from a variety of cultural backgrounds. Interventions conducted in clients' native language (if other than English) were twice as effective as interventions conducted in English	While searches were conducted across multiple databases and meta-analysis was performed, reporting does not represent a standard systematic review. Reference is missing to study selection, and critical appraisal.
Citation tracking	Handtke et al. 2019 Culturally competent healthcare – A scoping review of strategies implemented in healthcare organizations and a model of culturally competent healthcare provision. <i>PLoS ONE</i> 14(7): e0219971 https://doi.org/10.1371/journal.pone.0219971	Inception to August 2016 Update January 2017	ScR	Published	Included studies: n=67 (mental health n=19) Participants: All age groups. Migrants, culturally and linguistically diverse patients, ethnic minorities (e.g., Latino population, Native Americans, South Asian Americans) or refugees Countries: USA (n=51), Europe (n=6), Canada (n=5), Australia / New Zealand (n=3), Others (n=2) Identified strategies and components of culturally competent healthcare extracted from these studies were clustered into twenty categories, which were organized in four groups: Components of culturally competent healthcare—Individual level; Components of culturally competent healthcare—Organizational level; Strategies to implement culturally competent healthcare and Strategies to provide access to culturally competent healthcare	

					<p>The overall effects on patient outcomes and utilization rates of identified components or strategies were positive but often small or not significant</p> <p>Reported effects of single components or strategies are limited because most studies implemented a combination of different components and strategies simultaneously</p> <p>Qualitative data suggest that components and strategies of culturally competent healthcare were appreciated by patients and providers</p>	
PsycINFO	<p>Hernandez Robles et al. 2018</p> <p>Culturally adapted substance use interventions for Latino adolescents: A systematic review and meta-analysis. <i>Research on Social Work Practice</i> 28(7):789-801</p> <p>https://dx.doi.org/10.1177/1049731516676601</p>	January 1990 and March 2014	SR	Published	<p>Included studies: n=10 (across 14 reports)</p> <p>Participants: Adolescents (Latino) (age 11 to 18)</p> <p>Countries: USA</p> <p>Culturally adapted substance use interventions with Latino adolescents may be slightly more effective than other active interventions. Meta-analytic results suggest positive, yet small effects on substance use outcomes at posttest and slightly larger effects at follow-up. A moderate amount of heterogeneity was observed; however, no variables tested explained the variance. The risk of bias assessment revealed that most studies were at high risk for performance and selection bias.</p> <p>The authors also uncovered important gaps and deficiencies in this body of research, including the need to examine potential secondary benefits of culturally adapted interventions</p>	
PsycINFO	<p>Kalibatseva and Leong 2014</p> <p>A critical review of culturally sensitive treatments for depression: recommendations for</p>	Inception to December 2012	NR	Published	<p>Included studies: n=16</p> <p>Participants: Not reported</p> <p>Countries: Not reported</p> <p>The majority of culturally sensitive treatments for depression employed an evidence-based bottom-up approach, which involved general and practical</p>	<p>Only one database searched (PsycINFO) and only one author screened records</p>

	intervention and research. <i>Psychological Services</i> 11(4):433–450 https://dx.doi.org/10.1037/a0036047				adaptations, such as translating materials or infusing specific cultural values. Most studies used cognitive– behavioural strategies and included Latinos and African Americans	
PsycINFO	Li et al. 2023 Efficacy of culturally adapted interventions for common mental disorders in people of Chinese descent: A systematic review and meta-analysis. <i>The Lancet Psychiatry</i> 10(6):426-440 https://dx.doi.org/10.1016/S2215-0366%2823%2900118-9	Inception to 10 March 2023	SR	Published	Included studies: n=67 Participants: Aged 15 years or older with common mental health disorders (Chinese descents) Countries: China (n=60), Hong Kong (n=4), Taiwan (n=1), Australia (n=1), USA (n=1) Culturally adapted interventions had medium effect sizes in terms of reducing both self-reported (Hedges' g 0.77 [95% CI 0.61–0.94]; I ² 84%) and clinician-rated (0.75 [0.54–0.96]; I ² 86%) symptom severity across all disorders at end of treatment, irrespective of adaptation types. We noted no difference in efficacy between culturally modified interventions and culturally specific interventions. Subgroup analyses showed considerable heterogeneity. Inadequate reporting in included studies largely restricted risk-of-bias appraisals across all domains Psychological interventions can be transported across cultures with appropriate modifications. Adaptations to interventions can be made by modifying evidence-based interventions, or in culturally specific ways that are rooted in the sociocultural context. However, findings are limited by the insufficient reporting of interventions and cultural adaptations	Some of the included studies focused on adapting the format, delivery or content of an existing Western intervention (i.e. influenced by European culture, including Europe, USA, Canada and Australia) to meet the cultural needs of Chinese descents in China.
PsycINFO	Rathod et al. 2018 The current status of culturally adapted mental health interventions: A	2006–2016	OoR	Published	Included studies: n=12 Participants: All age groups; populations who belonged to a non-western cultural background, whether living in the west or outside the west	

	practice-focused review of meta-analyses. <i>Neuropsychiatric Disease and Treatment</i> 14:165-178 https://dx.doi.org/10.2147/NDT.S138430				Countries: Not reported There is value in cultural adaptation. Current evidence does not offer a solution to the issue of which components of cultural adaptation are effective, for what population, and whether cultural adaptation works better than noncultural adaptation. Well-defined and standardized frameworks for adaptation are needed, and further research in testing the application of these frameworks to evaluate the effectiveness of adapted interventions is essential. There is also a need to further refine the criteria and process of RCTs and meta-analyses in this field	
Cochrane	Soltan et al. 2022 Community-based interventions for improving mental health in refugee children and adolescents in high-income countries Cochrane Database of Systematic Reviews 5(5):CD013657 https://doi.org/10.1002/14651858.cd013657.pub2	Inception to Feb 2021	SR	Published	Included studies: n=62 Participants: Refugee children and adults Countries: High income Interventions: preventing and treating mental health problems (major depression, anxiety, post-traumatic stress disorder, psychological distress) and improving mental health There was no evidence of an effect of community-based interventions when compared with a waiting list for symptoms of post-traumatic stress (mean difference (MD) -1.46, 95% confidence interval (CI) -6.78 to 3.86; 1 study; low-certainty evidence), symptoms of depression (MD 0.26, 95% CI -2.15 to 2.67; 1 study; low-certainty evidence), and psychological distress (MD -10.5, 95% CI -47.94 to 26.94; 1 study; very low-certainty evidence)	There is insufficient evidence to determine the efficacy and acceptability of community-based mental health interventions for refugee children and adolescents
PsycINFO	Taylor et al. 2023 Review: Cultural adaptations to psychosocial interventions for families with refugee/asylum	Inception to Dec 2020	SR	Published	Included studies: n=11 Participants: Young people with refugee or asylum-seeker status Countries: UK	Whilst there is evidence for the use of CAIs with R/AS young people, the heterogeneity between studies limits the generalisability of these results. The available

	<p>seeker status in the United Kingdom – a systematic review. <i>Children & Adolescent Mental Health</i> 28(2):241-257.</p> <p>https://doi.org/10.1111/camh.12547</p>				<p>Interventions: Cultural adaptations to psychosocial interventions</p> <p>Studies used a variety of cultural adaptations including surface-level and deep-level adaptations. Studies showed some support for the use of CAIs with young people with R/AS, with varying degrees of symptom reduction. It was not possible to compare the effectiveness of CAIs against 'treatment-as-usual', nor to determine the effectiveness of different CAI components.</p>	<p>research is not sufficient to provide conclusive evidence of the use of CAIs over 'treatment-as-usual'</p>
Cochrane	<p>Uphoff et al. 2020</p> <p>An overview of systematic reviews on mental health promotion, prevention, and treatment of common mental disorders for refugees, asylum seekers, and internally displaced persons. Cochrane Database of Systematic Reviews. 9(9):CD013458.</p> <p>https://doi.org/10.1002/14651858.CD013458.pub2</p>	Inception to Sept 2019	OoR	Published	<p>Included reviews: (n=23)</p> <p>Included protocols: (n=15)</p> <p>Participants: Adults and children who are refugees, asylum seekers, and internally displaced</p> <p>Countries: No geographical restrictions</p> <p>Interventions: Interventions must have been aimed at mental health promotion (for example, classroom-based well-being interventions for children), prevention of mental health problems (for example, trauma-focussed Cognitive Behavioural Therapy to prevent post-traumatic stress disorder), or treatment of common mental disorders and symptoms (for example, narrative exposure therapy to treat symptoms of trauma)</p> <p>Reviews more commonly included refugees and asylum seekers than internally displaced persons, and were more frequently focused on adults than children</p> <p>There was more attention on the treatment of post-traumatic stress disorder than there was for mental health promotion or prevention, or for the treatment of depression or anxiety</p>	<p>This overview of systematic reviews summarises the characteristics of reviews available on this topic, to help determine which research questions are the most important to address in future Cochrane reviews</p>

					<p>Studies of Cognitive Behavioural Therapy, Narrative Exposure Therapy, and integrative and interpersonal therapies were most likely to be included in reviews</p> <p>Gaps exist in the evidence on mental health interventions for refugees, asylum seekers, and internally displaced persons. Most reviews do not specify that internally displaced persons are included in the selection criteria, even though they make up the majority of involuntary migrants worldwide. Reviews specific to mental health promotion and prevention of common mental disorders are missing, and there is more evidence available for adults or mixed populations than for children. The literature is focused on post-traumatic stress disorder and trauma-related symptoms, with less attention for depression and anxiety disorders. Better quality systematic reviews and better report of review design and methods would help those who may use these reviews to inform implementation of mental health interventions.</p>	
Help seeking behaviour						
Google PsycINFO	<p>Mukadam et al. 2013</p> <p>Improving access to dementia services for people from minority ethnic groups. <i>Current Opinion in Psychiatry</i>; 26(4):409-414</p> <p>https://doi.org/10.1097/YCO.0b013e32835ee668</p>	Not reported	NR	Published	<p>Included studies: Not reported</p> <p>Participants: Not reported</p> <p>Countries: Not reported</p> <p>Many countries are carrying out educational campaigns in an effort to increase awareness about dementia and reduce stigma, but none of these have reported any measurable outcomes of their interventions</p> <p>Studies show that knowledge about dementia has the potential to increase help-seeking, but information should be targeted to the recipient audience</p>	<p>No methods section that describes the search strategy, inclusion and exclusion criteria, study selection process, or whether quality assessment was conducted. Thus, there are a potential for bias in the review</p>

PsycINFO	Place et al. 2021 Interventions to increase migrants' care-seeking behaviour for stigmatised conditions: A scoping review. <i>Social Psychiatry and Psychiatric Epidemiology</i> ; 56:913–930 https://doi.org/10.1007/s00127-021-02065-1	Inception to 5 July 2019	ScR	Published	Included studies: n=16. Mental Health: n=15 Participants: migrants in high-income countries. Countries: USA (n=15), Canada (n=1) Three approaches were identified to increase migrants' mental health care-seeking behaviour: health communication (n=9), support groups (n=2), and primary care-based approaches (n=4), including a self-assessment tool for psychosocial risk, and engagement interventions seeking to increase collaboration between primary care and mental health services.	N.B: One study (1 out of 16) sought to increase migrants' care-seeking behaviour for Hepatitis B Virus (HBV) infection using health communication approach All quantitative studies
Treatment engagement						
Epistemonikos	Interian et al. 2013 Improving treatment engagement of underserved U.S. racial-ethnic groups: a review of recent interventions. <i>Psychiatric Services</i> ; 64(3):212-222 https://doi.org/10.1176/aipi.ps.201100136	2001-2011	SR (stated – only one database searched)	Published	Included studies: n=10 Participants: Adults Countries: USA (n=10) Focused on mental health treatment engagement among underserved racial-ethnic minority populations Evidence supported the efficacy of collaborative care for depression as an engagement enhancement intervention among underserved racial-ethnic populations Several other interventions demonstrated possible efficacy The effect of the interventions on clinical outcomes, such as symptom improvement and rehospitalization, was mixed	Only Medline was searched Limited to US adults and randomised studies
Epistemonikos TRIP PsycINFO	Moore 2018 Mental health service engagement among underserved minority adolescents and young	January 1995 to November 2016	SR	Published	Included studies: n=10 Participants: Adolescents and young adults Countries: USA (n=10)	Interventions were classified into 3 groups according to their Level of Evidence as Probably Efficacious,

	adults: a systematic review. <i>Journal of Racial and Ethnic Health Disparities</i> ; 5:1063-1076 https://doi.org/10.1007/s40615-017-0455-9				Four interventions were deemed probably efficacious Interventions that were family based or were culturally adapted for age group or race-ethnicity also showed possible efficacy and promising results	Possibly Efficacious, and Promising Interventions The lack of studies focused on this population limited findings
Treatment initiation, participation or continuation						
Epistemonikos TripPro PsycINFO	Aggarwal et al. 2016 Clinician descriptions of communication strategies to improve treatment engagement by racial/ethnic minorities in mental health services: a systematic review. <i>Patient Education and Counseling</i> ; 99(2):198-209 https://doi.org/10.1016/j.pec.2015.09.002	Inception to November 2014	SR	Published	Included studies: n=23 Participants: Healthcare professionals and patients Countries: USA (n=12), Australia (n=3), UK (n=1), Spain (n=1) and other European countries (n=1) Reasons for patient delays in treatment initiation Five studies noted clinician attempts to improve treatment participation by changing the context of communication Reasons for decisions to continue or discontinue treatment Some studies reported on clinician experiences to improve patient-clinician communication, though none reported randomized controlled interventions to target specific mechanisms of action	Mainly qualitative data (n=21)
Google TRIP Epistemonikos TripPro PsycINFO	Lee-Tauler et al. 2018 A systematic review of interventions to improve initiation of mental health care among racial-ethnic minority groups. <i>Psychiatric Services</i> ; 69(6):628-647. https://doi.org/10.1176/aipi.ps.201700382	Inception to February 2016	SR	Published	Included studies: n=29 Participants: All ages Countries: USA (n=29) Focused on mental health care among racial-ethnic minority groups Identified interventions: collaborative care (n=10), psychoeducation (n=7), case management (n=5), colocation of mental health services within existing services (n=4), screening and referral (n=2), and a	Limited to US residents.

					change in Medicare medication reimbursement policy that served as a natural experiment (n=1) Reduction of disparities in the initiation of antidepressants or psychotherapy was noted in four involving collaborative care, two involving colocation of mental health services, and one involving screening and referral interventions	
Review protocols						
PROSPERO	Aslam et al. 2018 Barriers and facilitators to uptake of psychosocial interventions delivered by lay therapists to improve mental health and wellbeing of asylum seekers and migrants: a systematic review https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42018104453	January 2007 to July 2018	Protocol	Published	Participants: Lay health workers, refugees, asylum seekers and their families, policy makers, programme managers, other health workers, or any others involved in or affected by the programmes. Countries: Not restricted by country To conduct a systematic review to identify the barriers and facilitators to uptake of psychosocial interventions delivered by lay therapists to improve mental health and well-being of asylum seekers and migrants - What are the psychosocial interventions delivered by lay therapists to improve mental health and wellbeing of asylum seekers and migrants? This will be a broad mapping exercise - What are the barriers and facilitators to uptake of these interventions using lay therapists? Full review - What evidence is there on the relative cost-effectiveness of psychosocial interventions delivered by lay therapists to improve mental health and wellbeing of asylum seekers and migrants?	Studies involving migrants were excluded
PROSPERO	Stockwell and Roche 2022 A systematic review of interventions that help black men access timely	No date restrictions	Protocol	Published	Participants: Black men who have a mental health need requiring support in a community setting. 18-70 years old Countries: Not restricted by country	English only People without a settled status i.e. asylum seekers or refugees. were excluded

	and appropriate support for their mental health https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42022345323				What culturally sensitive interventions are there for black men that improve access and/or treatment in the community for their mental health care - How effective are these interventions? - What is the experience of black men utilizing community mental health interventions specifically designed to support them in the community? - What are the components and mechanisms of community mental health interventions designed for black men which help them in accessing timely and appropriate care?	Diagnosis does not need to be established if mental health symptoms are described i.e. stress, low mood, anxiety etc.
PROSPERO	Wang and Kim 2019 A systematic review of community-based mental health interventions for Asian Americans https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42019136832	Not reported	Protocol	Published	Participants: Asian Americans and/or first-generation Asian immigrants to the United States Countries: US only What community-based interventions are being used to address mental health challenges facing Asian American communities? How do interventions address barriers to mental health resource utilization? How do interventions address the barrier of cultural stigma?	English only Studies that do not include a community-based component (i.e. only take place in clinical settings) will be excluded
PROSPERO	Wood et al. 2023 The impact of community-level interventions on improving access to primary care for people from minority ethnic groups experiencing a first-episode psychosis https://www.crd.york.ac.uk/prospero/display_rec	Not reported	Protocol	Published	Participants: Adult participants with first episode psychosis from minority ethnic groups Countries: Not restricted by country What is the impact of community-level interventions focused on improving access to primary care for people from minority ethnic groups experiencing a first-episode psychosis	Excluded -Studies that do not include community-level interventions -Studies that do not include primary care as a pathway to care or the duration of untreated psychosis

	ord.php?ID=CRD42023398682					
--	---	--	--	--	--	--

Key: CAI - Culturally adapted interventions; NR - narrative review; OoR – overview of reviews; R/AS - refugee or asylum-seeker status; SR - systematic review; ScR - scoping review

Table 3: Summary of included secondary evidence identified by stakeholders

Citation	Recency (Search dates)	Evidence Type*	Status**	Key findings from abstracts	Reviewer comments
<p>Kunorubwe et al. 2023 (Public Health Wales and University of South Wales)</p> <p>Guidance and an associated action plan to improve access to, and provision of, psychological interventions to people from Black, Asian and minority ethnic communities - Summary Report of a Rapid Review</p> <p>No link is available</p>	<p>Inception to 14th March 2023</p>	<p>RR</p>	<p>Unpublished</p>	<p>Included studies: n=19 (culturally adapted therapies (n=3); culturally adapted cognitive therapy (n=2))</p> <p>Participants: All ages from Black, Asian and minority communities</p> <p>Countries: UK (n=19)</p> <p>The available research indicates both culturally sensitive and culturally adapted psychological therapies have shown positive results in terms of accessibility, appropriateness, acceptability and outcomes. It is of note however, that there is still limited research on the efficacy of such interventions across distinct cultural communities. The literature also supports the utilisation of more community based collaborative approaches to service development to remove access barriers and improve outcomes</p> <ol style="list-style-type: none"> 1. Therapists and service leads should consider adopting existing models for developing culturally appropriate interventions. For example, Bernal et al's ecological model for the cultural adaptation of psychotherapeutic interventions (Bernal, Bonilla, & Bellido, 1995) 2. Therapists should adopt a collaborative, 'knowledge exchange' approach to intervention when a client's cultural background is unfamiliar to them 3. Imagery, proverbs metaphor, stories, and other narratives can be utilised as important tools for clients to vocalise and understand their experiences and distress, and somatisation of distress is also more likely in some ethnic groups 4. Therapists should establish the client's perspective on their safety, both physical (e.g., fear of harm, repatriation) and psychosocial (e.g., stigma, confidentiality) as a key feature of therapy 	<p>This RR aimed to provide up-to-date evidence on accessibility, appropriateness, acceptability and outcomes of psychological therapies for those from Black, Asian and minority ethnic communities and to provide evidence of both good practice and barriers to good practice</p> <p>Majority of the included evidence was qualitative</p>

				<p>5. Community stakeholders (e.g., faith leaders, third sector organisations) should be involved in discussions about developing and delivering interventions</p> <p>6. There should be utilisation of tools to facilitate a culturally-minded approach to practice and research. For example, the template to guide the development of a culturally adapted or culturally sensitive psychological therapy (Fig 2, p38)</p> <p>7. A culture of evidence-based practice and practice-based evidence should be embedded across the full scope of mental health services</p> <p>8. Guidelines such as the Medical Research Council framework for developing and evaluating complex interventions for diverse cultural groups, should be utilised when evaluating service provision</p>	
<p>Pollard and Howard 2021</p> <p>Mental healthcare for asylum-seekers and refugees residing in the United Kingdom: a scoping review of policies, barriers, and enablers. <i>International Journal of Mental Health Systems</i>; 15(60)</p> <p>https://doi.org/10.1186/s13033-021-00473-z</p>	2011-2020	ScR	Published	<p>Included studies: n=39</p> <p>Participants: All ages (included studies with specific groups (n=12): child and adolescent refugees or unaccompanied minors (n = 8), refugee men (n = 3), and refugee women (n = 1))</p> <p>Countries: UK (n=39)</p> <p>Synthesised themes from literature and interviews included existing barriers (i.e. communication difficulties and lack of funding, resources, and political will) and potential enablers (i.e. proposed provision practices, social needs of ASR, and policy changes)</p> <p>There is a gap in the literature regarding UK-wide assessment of access and delivery of mental healthcare for ASR in the UK</p> <p>Time sensitive and culturally appropriate approaches are needed, with greater funding and resource support from the UK Government</p> <p>This study provides justification for a call to relax hostile environment policies, and for ASR-specific mental health</p>	Data from stakeholder interviews (n=10) were also included

				services and support to be considered within the UK. Further research is needed to assess implementation of guidelines across the UK	
Satinsky et al. 2019 Mental health care utilisation and access among refugees and asylum seekers in Europe: A systematic review. <i>Health Policy</i> ; 123(9):851-863 https://doi.org/10.1016/j.healthpol.2019.02.007	January 2007 to August 2017	SR	Published	<p>Included studies: n=27</p> <p>Participants: All ages (including children and adolescents or refugees and asylum seekers who arrived as teenagers (n=6))</p> <p>Countries: Some studies were conducted in multiple countries, which included UK (n=9), Germany (n=5), the Netherlands (n=4), Norway (n=4), Ireland (n=4), Denmark (n=3), Sweden (n=3), Austria (n=2), Belgium (n=2), France (n=2), Italy (n=2), Poland (n=2), Switzerland (n=2), Croatia (n=1), the Czech Republic (n=1), Hungary (n=1), Portugal (n=1), and Spain (n=1)</p> <p>The findings suggest inadequate MHPSS utilisation. Major barriers to accessing care included language, help-seeking behaviours, lack of awareness, stigma, and negative attitudes towards and by providers</p> <p>Refugees and asylum seekers have high mental health needs but under-utilise services in European host countries</p> <p>This underutilisation may be explained by cultural-specific barriers which need to be tackled to increase treatment demand</p> <p>Training health providers on cultural models of mental illness may facilitate appropriate identification, referral, and care</p> <p>Based on these findings, it is crucial to review policies regarding MHPSS provision across the EU</p>	

Key: ASR – asylum-seekers and refugees; MHPSS - Mental health and psychosocial support; RR – rapid review; SR - systematic review; ScR - scoping review

Table 4: Summary of included organisational reports identified by stakeholders

Citation Citation retrieval source	Details
<p>Roberts 2023</p> <p>Psychological interventions for refugees and asylum seekers: current evidence</p> <p>No link is available</p> <p>Identified by stakeholder group</p>	<p><u>Type of publication</u> Presentation (unpublished)</p> <p>Overview of evidence that included: -A network meta-analysis of psychosocial interventions for refugees and asylum seekers with PTSD https://gh.bmj.com/content/6/6/e005029.long n=23 studies -Residual posttraumatic stress disorder symptoms after provision of brief behavioral intervention in low- and middle-income countries: An individual-patient data meta-analysis https://doi.org/10.1002/da.23221 -And 5 RCTs</p> <p><u>Focus</u> To scope the evidence on the management of post-traumatic stress disorder</p> <p>Cognitive behavioural therapy (SMD=-1.41; 95% CI -2.43 to -0.38) and eye movement desensitisation and reprocessing (SMD=-1.30; 95% CI -2.40 to -0.20) were significantly more effective than waitlist for refugees and asylum seekers.</p> <p>There was greater symptom reduction for Problem Management Plus than for enhanced treatment as usual for most symptoms for refugees and asylum seekers. Hyperarousal symptoms were the most common residual symptoms after PM+, with more than 30% of participants reporting persistent sleep disturbance, concentration difficulties, and anger.</p>
<p>Traumatic Stress Wales 2023</p> <p>Draft Asylum Seekers, Refugees and Migrants Mental Health Services Working Group Proposal</p> <p>No link is available</p> <p>Identified by stakeholder group</p>	<p><u>Type of publication</u> Working group proposal (unpublished)</p> <p><u>Focus</u> The proposal aims to result in a whole system approach that is integrated into current structures and initiatives, facilitates access to existing services and helps services to operate in a fully trauma-informed manner.</p> <p>A scoping exercise identified good practice in Wales and across the UK:</p> <p>Gwent Health Inclusion Service -To address health inequalities within the Aneurin University Health Board catchment area. The service is a nurse-led specialist NHS community team which provides primary health care level services for vulnerable groups of all ages, with a specific focus on people who have difficulty in accessing standard primary care services</p> <p>-A Social Determinants of Mental Health & Community Approach</p>

	<p>Based in Aneurin Bevan University Health Board Child & Family Community Psychology team, places an emphasis not on “what is wrong” with someone but “what has happened” to them</p> <p>-Cardiff and Vale Health Inclusion Service (CAVHIS) CAVHIS provides health screening and urgent care for newly arrived asylum seekers and trafficked individuals moved to Cardiff by the Home Office</p> <p>-Guy’s & St Thomas’ Health Inclusion Team, London This team provides a primary health care level service for vulnerable groups (refugees, asylum seekers, homeless populations), and the refugee and asylum seeker services are across Southwark and Lambeth</p> <p>-Solace (Yorkshire & Humber) Solace is a third sector charitable organisation delivering psychotherapy and mental health support to refugees and asylum seekers across the Yorkshire and Humber region</p>
<p>VCSE Health and Wellbeing Alliance 2022</p> <p>Barriers and Opportunities: Improving access to mental health support for refugees and people seeking asylum</p> <p>https://www.redcross.org.uk/about-us/what-we-do/we-speak-up-for-change/vcse-health-and-wellbeing-alliance</p> <p>Identified by stakeholder group</p>	<p><u>Type of publication</u> Qualitative primary research report</p> <p><u>Focus</u> To collaborate with people with lived experience of seeking asylum in England to design research which explores the barriers faced by refugees and people seeking asylum when trying to access support for their mental health</p> <p>To explore what professionals working within organisations providing mental health and related support to refugees and people seeking asylum, including within the NHS, identify as barriers to accessing mental health support among this group</p> <p>To enable people with lived experience of seeking asylum to identify opportunities for improving the access to, and provision of, mental health support for refugees and people seeking asylum in England</p>
<p>Welsh Government 2023</p> <p>Research into “Good Access” in Community Pharmacy, NHS Dentistry and Allied Health Professional Services</p> <p>https://www.gov.wales/research-good-access-community-pharmacy-nhs-dentistry-and-allied-health-professional-services</p> <p>Identified by stakeholder group</p>	<p><u>Type of publication</u> Qualitative primary research report</p> <p><u>Focus</u> To identify key barriers and facilitators to accessing primary care across Community Pharmacy, NHS Dentistry and Allied Health Professionals</p> <p>To explore what good access means to people (general public) and the expectations of what good access looks like in Community Pharmacy, NHS Dentistry and Allied Health Professionals</p> <p>To explore perceptions of phone first and telephone consultations / triage</p>

	To explore perceptions of face-to-face appointments vs use of technology To identify awareness of available options such as; NHS 111, Common Ailment Scheme; My health online and the NHS Wales App
--	--

Key: SMD – Standardised Mean Difference; VCSE – Voluntary Community and Social Enterprise

8. ADDITIONAL INFORMATION

8.1. Conflict of interest

The authors declare they have no conflicts of interest to report.

8.2. Acknowledgments

The authors would like to thank Beverly Morgan, Kim Swain, Hannah Bayfield, Thomas Hoare, Olivia Gallen and Mel (Melanie) McAulay for their contributions during stakeholder meetings in guiding the focus of the review and interpretation of findings.

9. APPENDIX

APPENDIX 1: – Resources searched during Rapid Evidence Summary

A single list of resources has been developed for guiding and documenting the sources searched as part of a Rapid Evidence Summary. Not all resources will be searched, depending on relevancy. Some sources will be searched as part of the subsequent Rapid Review (or Rapid Evidence Map).

Secondary research resources	Success or relevancy of the retrieval
Medical and health	
Cochrane Database of Systematic Reviews (CDSR) https://www.cochranelibrary.com/cdsr/reviews	Searched, results found
JBI (via OVID) (Subscription based service – WCEBC has a subscription)	Searched, nothing found
NIHR Journals Library https://www.journalslibrary.nihr.ac.uk/#/	Searched, results found
Trip https://www.tripdatabase.com/	Searched, results found
PROSPERO https://www.crd.york.ac.uk/prospéro/	Searched, results found
Additional resources searched	
Google Advanced Search https://www.google.co.uk/advanced_search	Searched, results found
PsycINFO (Subscription based service – CU has a subscription)	Searched, results found

APA PsycInfo <1806 to November Week 1 2023>

Mental health conditions

#	Query	Results from 17 Nov 2023
1	exp "Racial and Ethnic Groups"/	159,436
2	exp Ethnic Identity/	19,369
3	exp Minority Groups/	21,400
4	exp Human Migration/	43,522
5	exp Refugees/	8,799
6	ethnic*.tw.	110,866
7	(migrant* or immigrant* or emigrant*).tw.	43,736
8	(minorit* adj2 group*).tw.	11,478
9	(racial or race equity).tw.	59,534
10	(refugee* or asylum*).tw.	15,159
11	(black or BAME).tw.	74,926
12	((Afro or African or Asian or Latin or Indian) adj1 american).tw.	61,249
13	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12	348,531
14	exp Mental Health/	93,486
15	exp Mental Health Services/	58,487
16	*Mental Disorders/	77,498
17	(mental adj (health or ill* or well* or disease* or disorder* or condition*)).tw.	338,885
18	psychological health.tw.	8,102
19	(depression or anxiety).tw.	447,748

20	exp Posttraumatic Stress Disorder/	40,922
21	(post-traumatic stress disorder* or ptsd).tw.	46,663
22	exp Bipolar Disorder/	35,005
23	exp Schizophrenia/	99,998
24	(schizo* or "mood disorder*" or "personality disorder*" or psychotic* or psychosis or psychoses).tw.	241,878
25	(bipolar or mania).tw.	51,644
26	*Personality Disorders/	11,478
27	*Psychosis/	28,356
28	*Paranoid Psychosis/	1,089
29	14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23 or 24 or 25 or 26 or 27 or 28	956,706
30	exp Health Care Access/	9,814
31	(access* or utilis* or utiliz* or uptake* or availab* or service* or pathway* or refer*).tw.	1,131,604
32	engag*.ti.	27,370
33	30 or 31 or 32	1,153,171
34	(intervention* or innovation* or improv* or trial*).tw.	1,081,567
35	exp "Systematic Review"/	830
36	exp Meta Analysis/	5,399
37	(systematic review* or scoping review* or rapid review* or narrative review* or meta-analysis or evidence synthes#s).ti.	57,656
38	35 or 36 or 37	60,322
39	13 and 29 and 33 and 34 and 38	266

APA PsycInfo <1806 to 28 November 2023>

Dementia

#	Query	Results from 28 Nov 2023
1	exp "Racial and Ethnic Groups"/	159,567
2	exp Ethnic Identity/	19,386
3	exp Minority Groups/	21,426
4	exp Human Migration/	43,555
5	exp Refugees/	8,814
6	ethnic*.tw.	110,964
7	(migrant* or immigrant* or emigrant*).tw.	43,764
8	(minorit* adj2 group*).tw.	11,486
9	(racial or race equity).tw.	59,594
10	(refugee* or asylum*).tw.	15,178
11	(black or BAME).tw.	75,006
12	((Afro or African or Asian or Latin or Indian) adj1 american).tw.	61,282
13	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12	348,828
14	exp dementia/	92,970
15	exp Alzheimer's Disease/	56,227
16	dementia.tw.	79,129
17	alzheimer.tw.	16,210
18	14 or 15 or 16 or 17	113,897
19	exp Health Care Access/	9,838
20	(access* or utilis* or utiliz* or uptake* or availab* or service* or pathway* or refer*).tw.	1,132,640

21	engag*.ti.	27,399
22	19 or 20 or 21	1,154,227
23	(intervention* or innovation* or improv* or trial*).tw.	1,082,671
24	exp "Systematic Review"/	832
25	exp Meta Analysis/	5,401
26	(systematic review* or scoping review* or rapid review* or narrative review* or meta-analysis or evidence synthes#s).ti.	57,864
27	23 or 24 or 25 or 26	60,531
28	13 and 18 and 22 and 23 and 27	19

APA PsycInfo <1806 to November Week 3 2023>

Gypsy, Roma or Travellers

#	Search	Results from 29 Nov 2023
1	exp Romanies/	419
2	roma.mp.	865
3	romany.mp.	25
4	romani.mp.	167
5	gypsy.mp.	325
6	gypsies.mp.	293
7	gipsies.mp.	6
8	travel?er*.mp.	2,640
9	1 or 2 or 3 or 4 or 5 or 6 or 7 or 8	3,899
10	exp Mental Health/	93,835
11	exp Mental Health Services/	58,584
12	*Mental Disorders/	77,635
13	(mental adj (health or ill* or well* or disease* or disorder* or condition*)).tw.	339,778
14	psychological health.tw.	8,123
15	(depression or anxiety).tw.	448,503
16	exp Posttraumatic Stress Disorder/	41,013
17	(post-traumatic stress disorder* or ptsd).tw.	46,752
18	exp Bipolar Disorder/	35,034
19	exp Schizophrenia/	100,063
20	(schizo* or "mood disorder*" or "personality disorder*" or psychotic* or psychosis or psychoses).tw.	242,071
21	(bipolar or mania).tw.	51,694
22	*Personality Disorders/	11,489
23	*Psychosis/	28,409
24	*Paranoid Psychosis/	1,089
25	exp dementia/	93,028
26	exp Alzheimer's Disease/	56,265
27	dementia.tw.	79,166
28	alzheimer.tw.	16,222
29	10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22 or 23 or 24 or 25 or 26 or 27 or 28	1,050,286
30	exp Health Care Access/	9,861
31	(access* or utilis* or utiliz* or uptake* or availab* or service* or pathway* or refer*).tw.	1,133,781
32	engag*.ti.	27,464
33	30 or 31 or 32	1,155,416
34	(intervention* or innovation* or improv* or trial*).tw.	1,083,963
35	exp "Systematic Review"/	836
36	exp Meta Analysis/	5,407

37	(systematic review* or scoping review* or rapid review* or narrative review* or meta-analysis or evidence synthes#s).ti.	58,033
38	35 or 36 or 37	60,704
39	9 and 29 and 33 and 34 and 38	0

An additional search focused search was conducted in PsycINFO using the following terms (cultur* or adapt* or modif* or target* or inform* or specific* or tailor* or sensitive).ti and 61 citations were retrieved and screened.



The Health and Care Research Wales Evidence Centre

Our dedicated team works together with Welsh Government, the NHS, social care, research institutions and the public to deliver vital research to tackle health and social care challenges facing Wales.

Funded by Welsh Government, through Health and Care Research Wales, the Evidence Centre answers key questions to improve health and social care policy and provision across Wales.

Along with our collaborating partners, we conduct reviews of existing evidence and new research, to inform policy and practice needs, with a focus on ensuring real-world impact and public benefit that reaches everyone.

Director: Professor Adrian Edwards

Associate Directors: Dr Alison Cooper, Dr Natalie Joseph-Williams, Dr Ruth Lewis



@EvidenceWales @tystiolaethcym



healthandcareevidence@cardiff.ac.uk



www.researchwalesevidencecentre.co.uk