



Review/Comparative article

Methodological approaches to measuring mental health in a cost-of-living crisis: A rapid review

Clare England^{a,*}, David Jarrom^a, Jennifer Washington^a, Elise Hasler^a, Leona Batten^a, Adrian Edwards^b, Ruth Lewis^c

^a Health Technology Wales, United Kingdom

^b Health and Care Research Wales Evidence Centre, Cardiff University, United Kingdom

^c Health and Care Research Wales Evidence Centre, Bangor University, United Kingdom



ARTICLE INFO

Keywords:

Cost-of-living
Financial crisis
Mental health
Health policy

ABSTRACT

Background: Cost-of-living crises are damaging to population mental health and require a public health response. It is important to assess whether public health interventions are effective. We aimed to identify population-level methods and measures and the appropriateness of the measures for vulnerable populations.

Methods: A rapid evidence review was undertaken. Nineteen databases, including grey literature, were searched for evidence published between 1970 and April 2023.

Results: Seven reviews, nine primary studies and two reports from grey literature were identified. Methods consisted of analyses of existing data from national or regional cohort studies, household panel surveys, repeated cross-sectional surveys, routine medical data, or data on suicide death rates. Twelve brief validated mental health measurement tools, embedded in population-level surveys, were identified. Two quasi-experimental studies used data from a UK household panel survey to examine the impact of the introduction of specific welfare policies on mental health. Studies identified socio-economic vulnerabilities, but it was not possible to determine whether data were effectively captured from people from minority ethnic groups.

Conclusion: Population-level surveys can be used in quasi-experimental studies to measure the effects of a public health initiative with specific roll out dates to tackle cost-of-living impacts. It is unclear as to whether the identified methods and tools are suitable for use with people from minority ethnic groups.

1. Background

During 2021 and 2022 the UK experienced a sharp rise in inflation. For many people, wages and welfare payments did not keep up with the rate of inflation, leading to a cost-of-living crisis. Energy, food, fuel, and housing became less affordable leading to adverse short and long-term health outcomes [1]. There is evidence from past economic crises, such as the 2008/2009 global financial crisis (the Great Recession) that economic shocks and increases in the cost-of-living have a detrimental effect on mental health across the population and that vulnerable populations are disproportionately affected [2]. Furthermore, people with poorer mental health are more likely to experience further reductions in income. The negative impacts on mental health and well-being induced

by the cost-of-living crisis could lead to future financial and health problems, potentially creating a downward spiral that can persist even if economic conditions improve [3].

In response, national and regional public health interventions may be put into place to mitigate the effects of the cost-of-living crisis. It is important for policymakers and practitioners to be able to evaluate the impact of any public health measure on mental health at a population level, but designing the studies, recruiting participants, and collecting the data can be expensive and time consuming. However, there are ongoing surveys and cohort studies undertaken by governments and national and regional organisations that include mental health measurement methods and which could potentially be used to evaluate the effect of public health interventions. An overview of mental health

Abbreviations: Crpd, clinical practice research datalink; Ghq-12, 12-item general health questionnaire; Gdp, gross domestic product; Ons, office for national statistics; Phq-8, population health survey; Sf-12, short form 12 mental health component summary; Thin, The Health Improvement Network; Ukhls, Understanding Society: The UK Household Longitudinal Study; Wemwbs, Warwick-Edinburgh Mental Well-being Scale.

* Corresponding author at: Health Technology Wales (HTW), Second Floor, The Life Science Hub, 3 Assembly Square, Cardiff, CF10 4PL, United Kingdom.

E-mail address: clare.england2@wales.nhs.uk (C. England).

<https://doi.org/10.1016/j.healthpol.2024.105062>

Received 1 November 2023; Received in revised form 26 March 2024; Accepted 1 April 2024

Available online 4 April 2024

0168-8510/© 2024 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

measures could assist researchers and policy makers seeking to understand the impact of public health interventions on mental health.

We conducted a rapid review of articles that evaluated the impact of economic crises on population mental health. The aims of the review were to 1) identify and appraise available population-level measures for assessing the impact on mental health of any public health response to the cost-of-living crisis and 2) identify whether the measures are appropriate for use in vulnerable populations.

The purpose of the review was not to summarise the findings of included articles (i.e., impact of the economic crises on mental health or outcome of any public health interventions), or to identify how often different methods of measurement were used to evaluate the effect of economic crises on mental health, but explore what methods and tools are available and appropriate for monitoring the impact of the cost-of-living crisis on mental health and determine if these methods and mental health measurement tools suitable for specific vulnerable populations?.

2. Methods

We conducted a rapid review, which is a form of knowledge synthesis that accelerates the process of conducting a conventional systematic review by abbreviating or omitting specific methods to generate the evidence for stakeholders in a resource-efficient manner [4]. The review was conducted in line with best practice guidance for rapid reviews [5], and was based on existing reviews plus more recent primary studies that may not be covered by reviews.

2.1. Study selection and searches

The study selection criteria are shown in Table 1. The search strategy was developed and carried out in several parts (JW and EH). As per the study selection criteria, we searched for secondary evidence published between 1970 and 2023 and for primary evidence from 2021 to 2023 relating to cost-of-living and mental health. Further ‘mop-up’ searches were conducted to ensure we had identified all relevant mental health tools used to measure mental health during an economic crisis (1970 to 2023) and, because this study was of particular interest to Welsh policy-makers we conducted a search to identify research on cost-of-living and Wales (1970 to 2023). Search results were downloaded to Endnote 20.3 (Clarivate) and duplicates removed.

Searches ran to April 2023 in the following databases: KSR Evidence; Ovid Medline, Embase, PsycInfo, SSP and HMIC; EbscoHOST CINAHL; Scopus; Wiley Cochrane; ProQuest ASSIA, SSA, SA and SD; Social Care Online; TripPro; PROSPERO; Google and Google Scholar. The MEDLINE search strategies are available in the supplementary materials.

2.2. Study selection process

All references identified by the literature searches were screened for eligibility using the criteria listed in Table 1 by title and abstract. Potentially relevant full texts were assessed for inclusion using a structured approach. Firstly, existing reviews describing the effect of the cost-of-living crisis on mental health were assessed for relevance using their inclusion/exclusion criteria and characteristics of included studies. Reviews that did not clearly define their inclusion/exclusion criteria or where the majority of included studies did not fit our eligibility criteria (e.g., mainly cross-sectional studies) or that did not include a description of measurement tools were excluded. We made a pragmatic decision to include reviews with a minority of studies that did not meet our inclusion criteria because we were aware that many individual studies in this field were cross-sectional, and few reviews existed that only included longitudinal studies. However, due to the number of relevant articles identified it was not possible to include all reviews and studies. As multiple reviews were found that could potentially answer the research questions, the largest or most recent reviews that included details of the

Table 1
Study selection criteria.

| | Inclusion criteria | Exclusion criteria |
|--------------------------------|---|---|
| Population | Adults or households | Organisations and systems (e.g., exploring how an organisation such as a health / educational organisation or system responds) Children/adolescents only |
| Exposure | Cost-of-living crisis or economic / financial crisis | Related to financial stress caused by Covid-19 and/or measures to tackle Covid-19 or other infectious disease or disaster (natural or man-made) |
| Outcome | Mental health e.g., <ul style="list-style-type: none"> • Depressive/affective disorders • Anxiety disorders • Disruptive behaviour and dissociative disorders • Stress • Wellbeing • Suicide attempts/ideation • Suicide completion rates | Physical health Behavioural outcomes (such as healthy eating, physical activity) Sleep outcomes Psychosis Schizophrenia Substance misuse Eating disorders |
| Evaluation | Appraisal of the following: <ul style="list-style-type: none"> • Validated population-level surveys • Population-level mental healthcare use (including, but not limited to primary and secondary care records, hospital admissions, medication use) | Appraisal of questionnaires used only for individual level data. Non-validated tools developed specifically for a study (bespoke tools) |
| Study design | <ul style="list-style-type: none"> • Systematic reviews; scoping reviews; rapid reviews, organisational reports; clinical guidance Primary research, limited to: <ul style="list-style-type: none"> • Longitudinal observational studies (panel surveys, time series, interrupted time series and qualitative studies) • Repeated, population-level cross-sectional studies • Population-level intervention studies (randomised, non-randomised and single arm) | Single timepoint cross-sectional studies Individual-level intervention studies |
| Countries | Conducted in UK, Europe, Australia, New Zealand, English | All other countries |
| Language of publication | English | |
| Publication date | Secondary evidence from 1970 Primary studies 2021- present | |
| Publication type | Published, including grey literature | Conference abstracts Preprints |
| Other factors | Prior to 2021 relevant measures will be identified from secondary evidence (systematic reviews; scoping reviews; rapid reviews; rapid evidence assessments; organisational reports; clinical guidance). This date limit was arrived at since preparatory work had identified a large, eligible, scoping review that included articles published up to the end of Dec 2020 | |

available tools and methods were chosen for inclusion.

Relevant full texts of primary studies from 2021 to April 2023 were screened and included according to the following criteria 1) used evaluation methods not previously identified from reviews 2) with an experimental/quasi-experimental design 3) studies of high relevance to UK populations 4) included detailed exploration of strengths and limitations of evaluation methods.

Initial screening was conducted by a single reviewer (CE). A second reviewer dual screened 20 % of the primary studies for quality assurance purposes (LB).

2.3. Data extraction

The following data were extracted: study details (author, year, country, purpose, design), number of included studies and key characteristics from secondary evidence, mental health measurement methods and tools, economic measures, sources of data, strengths and limitations of the measurement tools as identified by study authors and vulnerable groups as identified by study authors. Where necessary further, non-systematic, searches were conducted to obtain details of the characteristics of the identified measurement tools and methods, prioritising articles describing the original development and validation of the tool. A full review and quality assessment of the psychometric properties of each tool was beyond the scope of this rapid review, instead a brief description of each tool, including domain, number of items, time taken to administration and availability was extracted. Data were extracted by

the lead author, with 20 % of records dual extracted for quality assurance by a second reviewer (DJ).

2.4. Quality appraisal

Formal quality appraisal of the evidence presented by the reviews and studies that used the methods was beyond the scope of the rapid review. We provide a narrative description of the search strategy and inclusion criteria of included reviews and studies and discuss pros and cons of potential study designs using the methods identified.

3. Results

After duplicates were removed, 6775 titles and abstracts of records were screened, of which 190 were identified for full text screening. We




Fig. 1. Search results and study selection.

identified 18 papers for inclusion (Fig. 1).

Four systematic reviews [6–9], one scoping review [2] one clinical guideline [10] and one rapid review from grey literature [11] were included. The reviews included a total of 322 different identifiable studies, of which 52 were included in more than one review (although Martin-Carrasco reported using evidence from a total of 354 studies, detailed data were only extracted from 69). In addition, we identified nine further peer-reviewed primary studies [3,12–19] and two reports from grey literature [20,21].

A detailed summary of study characteristics, mental health and economic measurement methods, author assessments of the methods used, and groups identified as particularly vulnerable are available in Tables 2, 3 and 4 for peer-reviewed secondary evidence, peer-reviewed primary studies, and grey literature respectively.

3.1. Identified study designs and sources of data

Table 5 summarises the different study designs, characteristics, and examples of sources of data from the reviews and primary studies. Most studies included in the reviews were large observational studies involving sample sizes in the thousands. They were secondary analyses of existing data drawn from national or regional cohort studies, household panel surveys, repeated cross-sectional surveys [2,6–10], or used medical records or prescribing data [2,9,10] or were ecological time-series studies using national or regional suicide death rates [2,6,8,10]. One case-control study that was included compared mental health outcomes from a country that experienced a recession with one that did not [6]. The reviews were, in general, not specific about which datasets were used, although one review [10] named two UK surveys: “The Health and Activity Lifestyle Survey” and “The English Longitudinal Study of aging” as sources of data.

Many studies included in the reviews compared mental health in the years pre- and post-onset of an economic crisis, some described treating the Great Recession of 2008/2009 as a ‘natural experiment’. It is not clear from the reviews exactly how the date of onset of the economic crisis was determined and whether this was comparable across studies. Only two reviews [6,9] gave details of specific economic measures. The macro-economic indicators that were used included national and regional rates of unemployment, gross domestic product (GDP) and home foreclosure rates. Individual or household-level indicators included employment status, job security, household income, perceived financial stress, indebtedness, and household tenure.

The additional primary studies and grey literature provided more details about study design and data sources. Of the UK-based studies, seven used data from “Understanding Society: The UK Household Longitudinal Study” (UKHLS) [3,11,15–17,19,20], one used data from the “Scottish Longitudinal Study” [14], one from the “Welsh Health Survey” (now “National Survey for Wales”) [18] and one from the Office for National Statistics (ONS) “Opinions and Lifestyle Survey” [21]. The former two are panel surveys that follow the same households or individuals over time, replacing participants who leave the sample and include data linkage with external datasets, including health data. The latter two are repeated cross-sectional surveys and include a different representative sample at each wave. Economic indicators were those also identified in the reviews. In addition, primary studies in the UK measured deprivation using the Index of Multiple Deprivation, other regional income indicators and aggregated data from other household surveys such as the “Wealth and Assets Survey” and “Household Labour Survey”.

Seven of the primary studies were observational [3,12–16,18] and two used a quasi-experimental design [17,19]. Both quasi-experimental studies used data from UKHLS to examine the effect of the introduction of specific welfare policies (the “Bedroom Tax” and the introduction of Universal Credit) on mental health.

3.2. Mental health measures

Mental health outcomes were frequently assessed using validated mental health measurement tools, which are embedded in many of the cohort and panel surveys. Twelve brief validated mental health or mental wellness measurement tools, self-administered, or administered by an interviewer, were identified from the included articles [22–33]. These are listed in Table 6. The 12-item General Health Questionnaire (GHQ-12) [22], the Short Form 12 Mental Health Component Summary (SF-12) [23] and the Warwick-Edinburgh Mental Well-being Scale (WEMWBS) [24] are all embedded into UKHLS, the WEMWBS [24] is used in the “National Survey for Wales” and the 8-item Population Health Survey (PHQ-8) [25] is used in the “Opinions and Lifestyle Survey”. Additional measures of mental health included data from medical and hospital records for incidence of mental illness, prescription of medication, suicide attempts and suicide rates. In the UK, aggregated data from GP recorded diagnoses from the Clinical Practice Research Datalink (CRPD) and primary care data from The Health Improvement Network (THIN) are available and panel surveys such as the Scottish Longitudinal Study are linked with medical records on an individual level.

3.3. Potential biases in study design

All reviews and studies included some discussion on strengths and limitations of study methodology (Tables 2, 3 and 4).

We identified one systematic review [8] that had the aim of evaluating biases in studies assessing the effects of the Great Recession on health in Spain. This review included 53 studies and evaluated bias using an adapted tool. Four main biases were identified: problems with evaluation, time bias, failure to adjust for confounders such as seasonal effects, and inconsistencies in defining the date of onset of the Great Recession. The biases identified are summarised below.

Problems with evaluation existed with a repeated-cross sectional design where the sample surveyed before recession onset did not consist of the same individuals as the sample surveyed after onset, and therefore were subject to confounding. Studies judged at lower risk of bias in this domain attempted to control for confounding by either matching subjects from the samples before and after the crisis; stratifying the samples by common characteristics, often into groups considered most vulnerable, or using multilevel modelling analyses. Glonti et al. (2015) also recommended other analyses such as dynamic modelling or structural equation modelling. Other reviews also point out that it is not possible to infer causality from studies with a repeated cross-sectional design, although a strength of these studies is that they are frequently national surveys that are representative of the population. However, if a longitudinal panel survey with a repeated measures design is used, the evaluation problems may be reduced and there is stronger evidence for causality.

Time bias existed in time series studies where there were a very few data points after crisis onset, or where there was no consideration given to any potential lag between exposure and outcome. Time series studies were mainly those where the health outcome was suicide. Studies into suicide rates are ecological studies using aggregated data of unknown quality and causality cannot be inferred.

Other reviewers also discussed failure or inability to adjust for confounding. A limitation of using existing datasets for secondary analysis is that data of relevance to the secondary analysis may not have been collected. For example, it was observed that UK population panel and repeated cross-sectional surveys do not always ask about both mental health and household economics in detail, and not all income- or employment-related factors of interest were included in all survey waves.

Surveys use self-reported data which may not be accurate. However, potential biases in the use of medical records were discussed in several other reviews and studies. Data in hospital and medical records were

Table 2
Summary of peer-reviewed secondary evidence.

| Author, date (Country of interest) | Review details (type, purpose, search period, economic crisis, outcomes, review quality) | Included studies (number, key characteristics) | Mental health measurement tools and methods | Specific economic measures | Author assessment of methods/other observations | Groups identified as particularly vulnerable |
|------------------------------------|---|--|---|--|---|---|
| Frasquilho, 2016 [6] (Any) | Systematic review Examined evidence on population mental health during recession and identified vulnerable groups Articles published between 2004 and 2014 Any economic crisis/recession Outcomes: Psychological well-being; mental health distress; common mental disorders; depression; suicide (including suicide behaviours and ideation) Search strategy, inclusion and exclusion criteria clearly described; narrative summary; no formal quality assessment but strengths and weaknesses for study methodology described | 101 studies 66 studies used national population samples; 16 cross-national population samples; 19 regional/community samples 79 studies were in a general population, 22 in different specific populations (e.g., working population, unemployed, older adults, hospital patients) 61 European, of which 9 are in the UK; 7 from Australasia; 33 from other countries. 2 case-control / quasi-experimental studies; 30 cohort; 17 repeated cross-sectional; 28 ecological; 24 cross-sectional All included studies had ≥ 1000 participants | Mental health measurement tools: <ul style="list-style-type: none"> • SF-36 MCS (General mental health) • CES-D scale (Psychological well-being and mental health) • EURO-D (Depressive symptoms) • HSCL-25 (Mental health distress caseness) • GHQ-12 (Mental health distress) • PSS-4 (Psychological stress) • Kessler-10 (Mental health distress) • MHI-5 (Mental health distress) • SCID-I (Depression) • WHO—CIDI (Depression) • Other methods: • GP / hospital records on incidence and prevalence of mental illness • Suicide rates • Hospital records on suicide attempts Population surveys: <ul style="list-style-type: none"> • Unspecified national and international cohort, panel and repeated cross-sectional surveys | Time variables (pre- and post-economic recession changes) Macroeconomic indicators: <ul style="list-style-type: none"> • rates of unemployment • GDP • home foreclosure rates Individual-level indicators: <ul style="list-style-type: none"> • employment status • psychosocial job quality and security • household income • perceived financial strain or security • perceived economy/recession stress • deprivation • indebtedness • housing payment problems • socioeconomic status | Representative data can be obtained from national and regional population samples. In longitudinal cohort studies, the chronological sequence of exposure, outcomes and confounders affect all participants at the same time, giving stronger evidence for causality. Repeated cross-sectional designs, studies using aggregated data and ecological studies cannot give evidence for causality. Many studies had a limited time-period, which meant long-term effects could not be measured. Important to measure and adjust for confounding factors such as seasonal variation and mental health prior to exposure. | People unemployed (either due to job loss during recession, or prior unemployment) People in insecure work Living with debt Low socioeconomic status At risk of home foreclosure / eviction People with pre-existing mental health conditions |
| Glonti 2015 [7] (Any) | Systematic review Examined evidence from longitudinal studies on factors influencing resilience among the general population living in countries exposed to financial crises Searched up to October 2013 Any economic crisis/recession Outcomes: Mental health; depression; perceived stress; psychological distress; mental well-being; happiness; attempted suicide. Search strategy, inclusion and exclusion criteria clearly described; narrative summary; quality of included studies assessed using an adapted version of the Quality Assessment Tool for Quantitative Studies | 22 studies in total, of which 11 included mental health outcomes in countries of interest. 11 European 5 cohort studies; 6 repeated cross-sectional | <ul style="list-style-type: none"> • GHQ-12 (Mental health) • PSS-4 (Stress levels) • Population surveys: • Unspecified national and international cohort, panel and repeated cross-sectional surveys | Not reported | Different analytical methods could be used to explore causality and suggest dynamic modelling or structural equation modelling. Multilevel modelling could be used to explore the effect of environmental influences on mental health. | People who are unemployed or in precarious employment People on lower incomes Women's mental health was found to be worse than men's mental health pre-crises. Mental health worsened during economic crises for both men and women, some studies found the change in mental health was greater in men, but women still tended to experience worse mental health overall. |

(continued on next page)

Table 2 (continued)

| Author, date (Country of interest) | Review details (type, purpose, search period, economic crisis, outcomes, review quality) | Included studies (number, key characteristics) | Mental health measurement tools and methods | Specific economic measures | Author assessment of methods/other observations | Groups identified as particularly vulnerable |
|------------------------------------|---|--|--|----------------------------|---|--|
| Guerra, 2021 [2] (Any) | Scoping review Summaries available evidence of the impact of economic recession on mental health Articles published between 2008 and end-December 2020 Any economic crisis/recession Outcomes: depressive symptoms; anxiety disorders; suicide mortality; suicide attempts; suicide ideation Search strategy, inclusion and exclusion criteria clearly described; narrative summary; no quality assessment | 127 studies 11 prospective cohort; 3 retrospective cohort; 1 case-control; 84 time-series; 28 cross-sectional. 72 studies in all adults; 36 in working age adults; 9 in older adults (definitions ranged from adults >50 to >75 years old); 2 young adults; 3 middle-aged adults. Also 5 in children and adolescents. 80 European; 7 Australasia; 40 other countries | Mental health measurement tools: • CES-D (Depressive symptoms) • EURO-D (Depressive symptoms) • Goldberg Depression and Anxiety Scales • SCID-I (Major depressive disorder) • WHO CIDI (Anxiety and mood disorders) Other measurements: • Rates of psychotropic medication use • Mental health service use (community or hospital) • Length of hospital admissions • Hospital records on suicide attempt rates • Suicide rates • Population surveys: • National and international cohort, panel and repeated cross-sectional surveys e.g., European Social Survey; Portuguese National Mental Health Survey | Not reported | There is a large quantity of available data, but it is very varied. Studies may not be generalisable since basis of each economic crisis is variable and many studies are based on 2008/2009 financial crash in housing or stock markets. The population affected by cost-of-living crisis may be different and rates of mental illness may vary. Populations at specific risk may be different in different countries. Local welfare policies may also be different and produce different impacts | Men are more vulnerable to suicide / suicide attempts Unemployed men People in insecure work At risk of home foreclosure / eviction People on low incomes People with low education People with pre-existing mental health conditions |
| Martin-Carrasco, 2016 [10] (Any) | Clinical Guidance Identify the impact on mental health in Europe of the economic downturn and the measures that may be taken to respond to it. Search to end 2014 Focus on economic crisis of 2007–2014 Search strategy, inclusion and exclusion criteria clearly described; narrative review; no quality assessment | Evidence is drawn from 354 articles, but only 69 European studies are summarised in detail. Of these: 17 studies relate to psychiatric morbidity (9 repeated cross-sectional; 4 cumulative registers; 1 prospective cohort; 1 longitudinal cross-cohort; 1 clinical descriptive; 1 cross-sectional) 20 relate to suicide behaviour (13 time-trends analysis) | Mental health measurement tools: • GHQ-12 (Mental health distress) • WHO CIDI (Anxiety and mood disorders) • Other measurements • Data extracted from calls made to mental health helpline (Greece) • Suicide rates • Population surveys: • National and international cohort, panel, and repeated cross-sectional surveys, including Health and Activity Lifestyle Survey (HALS); English Longitudinal Study of Ageing (ELSA) | Unemployment rate | The advantage of aggregate-level data is that they reflect the environmental effects of changes in the economy beyond an individual's situation. Aggregate level data cannot allow for variation in services e.g., An economic crisis may increase suicide rates for some population sub-groups but if suicide prevention strategies are provided at the same time that benefit other population sub-groups the net result may be a zero increase. | Men of working age (may be mediated by socioeconomic status) People who are unemployed or in precarious employment People at risk of foreclosure / eviction or living in neighbourhoods with high rates of foreclosure People living with debt People with existing mental health problems People with low social capital |
| Saez, 2019 [8] (Spain) | Systematic review Evaluate bias in studies assessing the effect of the Great Recession (between 2008 – 2013) on health in Spain. Search up to June 2018 Great Recession Outcomes: General mental health outcomes; suicide Search strategy, inclusion and exclusion criteria | 53 studies in total, of which 32 included mental health outcomes. Study designs and populations not clearly reported, although most studies reporting mental health outcomes used data from health surveys with repeated cross-sectional design and there were fewer cohort studies; time series studies were generally used where | • GHQ-12 (Mental health distress) • Suicide rates • Population surveys: • National and international cohort, panel, and repeated cross-sectional surveys | Not reported | 60 % of the included studies (all health conditions) were assessed as having a high risk of bias. Four main biases identified: evaluation problems, time-bias, lack of control of observed and unobserved confounders and non-exogenous definition of the onset of the Great Recession. A repeated cross-sectional design means different samples at each timepoint. | People who are unemployed Migrants People with low education People at risk of foreclosure / eviction |

(continued on next page)

Table 2 (continued)

| Author, date (Country of interest) | Review details (type, purpose, search period, economic crisis, outcomes, review quality) | Included studies (number, key characteristics) | Mental health measurement tools and methods | Specific economic measures | Author assessment of methods/other observations | Groups identified as particularly vulnerable |
|------------------------------------|--|---|---|---|--|--|
| | clearly described; narrative summary; assessment of study bias using 8 domains was adapted from a previously published tool (Parmer 2016) | the outcome was suicide All conducted in Spain | | | Authors suggest controlling by matching subjects across surveys, stratification by analysing data from selected vulnerable groups or adjustment for confounders in a multivariate model. A better study design would be use of longitudinal data from population-based cohort studies, with repeated measures on the same individual. Many time series studies were assessed as having insufficient number of periods after the crisis and did not consider lag effects. Not all studies control for observed or unobserved confounding (e.g., long-term temporal effects, seasonal variation) or control for the variability in mental health outcomes that is not explained by the crisis or the confounders (which could be done by introducing random effects). Varying definitions as to the onset of the Great Recession can affect results. | |
| Silva, 2018 [9] (Any) | Systematic review Examine impact of economic crises on use of mental health care Search to June 2018 Outcomes: Use of mental health services Early 1990s Post-Communist Depression; Late 1990s East Asian financial crisis; 2008 Great Recession Search strategy, inclusion and exclusion criteria clearly described; narrative summary; quality of studies assessed using Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies | 17 studies 5 repeated cross-sectional studies; 3 cohort; 1 retrospective cohort; 1 panel study; 4 time series; 3 ecological 10 European; 1 Australia; 6 other countries | Hospital records for: <ul style="list-style-type: none"> Incidence of attendance due to suicidal behaviour Use of emergency department due to mental health issues / suicidal behaviour Discharge from psychiatric hospitals Medical records for: <ul style="list-style-type: none"> Use of mental healthcare Use of prescription drugs National population surveys involving structured interviews that include questions on: Use of mental health care Prescription drugs | Change in rates of unemployment Local unemployment | Hospital / medical records are dependant on the structure of services and record keeping. Most studies reported that there was no reorganisation of mental health services during the period studies, but any changes (e.g., due to austerity measures) change availability of services and may alter availability disproportionately for more vulnerable people. | Women are higher users of mental health services Adults aged 35–54yrs increased use of care due to suicidal behaviour Men’s use of care due to suicidal behaviour was influenced by socioeconomic factors. Authors note that people seeking care were not necessarily those most impacted by economic crises |

CES-D=Centre for Epidemiological Study Depression Scale; GDP = Gross Domestic Product; GHQ-12 = 12-item General Health Questionnaire; GP = General Practitioner; HSCL-25 = Hopkins Symptom Checklist; MHI-5 = 5-item Mental Health Inventory; PSS-4 = Short form Perceived Stress Scale; SCID-I = Structured Interview to Identify Major Depressive Disorders; SF-36 MCS = 36-item Short Form Health Survey Questionnaire Mental Component Summary; WHO CIDI = World Health Organisation Structured Interview to Identify Anxiety and Mood Disorder.

Table 3
Summary of peer-reviewed primary studies.

| Citation (Country) | Study Details (purpose, design, mental health outcomes) | Study characteristics (source of data, details of sample, dates of data collection, analysis) | Mental health measurement tools and methods | Specific economic measures | Covariates / confounders | Author assessment of strengths and limitations of measurement | Groups identified as particularly vulnerable |
|--|---|--|--|--|--|--|---|
| Alvarez-Galvez, 2021 [12] (Spain) | Assess association between recent economic downturns and age-standardised suicide rates adjusting for social cohesion and community values (interpersonal trust, social capital, post materialist values obtained from World Values Survey) Time-series study Outcomes: Suicide | Population-level data Age-standardised suicide rates from: (1) 1980 to 1991; (2) from 1992 until 2007; (3) from 2008 to 2010; (4) from 2011 to 2017 Interrupted time-series analysis | Suicide rates per 100,000 obtained from the National Statistics Institute of Spain | From Eurostat data: • Unemployment rate • GDP growth • Social expenditure as a percentage of GDP | Social relationships data from the World Values Survey: • Interpersonal trust • Social capital • Postmaterialist values Men and women analysed separately Age standardised results presented | Methodological variations in suicide data from registries Aggregate national data used which means individual behaviour cannot be inferred | Men more at risk of suicide at all timepoints |
| Aretz, 2022 [13] (Netherlands/Germany) | Investigate whether the Great Recession (2007 to 2009) increased the risk of depression in older people in Europe. Panel study Outcomes: late-life depression | Data from Survey of Health, Ageing and Retirement in Europe (waves 1 – 7), conducted in Austria, Belgium, Denmark, France, Germany, Italy, Spain, Sweden, Switzerland Data from 2004 to 2017 Eligible participants were ≥50yrs in wave 1; had valid depression data in waves 1 and 2 (prior to Great Recession) and for at least one wave after; depression-free at baseline; did not relocate during study period 6866 participants eligible at baseline (out of 95,534); 3144 included in fixed effects model (53 % female) Analysis: Binary logistic individual fixed effect (FE) models using within-estimators. | EURO-D depression symptom scale | Country-level GDP data used to identify start of recession by country Perceived area deprivation (from SHARE) | Individual level confounders: • Age • Job status • Physical activity • Cognitive functioning • Frailty • Activity of daily living • Number of chronic diseases Household level confounders: • Living together with a partner • Household size • Urban/rural | EURO-D was developed to specifically to measure late-life depression Use of a mental health questionnaire identifies undiagnosed depression Survey only conducted every 2 years, which limits ability to identify very short-term effects. Individual-level confounders available (Note: This study is useful to show how a subset from cohort data can be used/derived) | People living in more deprived areas |
| Cherrie, 2021 [14] (Scotland) | Examine how the trend in 'new' antidepressant prescription rates during the Great Recession varied by regional economic conditions Longitudinal data linkage study Outcomes: Anti-depressant prescriptions | Data from the Scottish Longitudinal Study, linked to NHS service use and prescriptions for mental health conditions Eligible participants supplied data in 2001 and 2011 censuses (pre- and post-recession onset); were aged between 16 and 60 in 2011; were economically active and employed in the labour market in 2011. 86,500 participants eligible Linked to NHS data from | New anti-depressant prescriptions from linked NHS data | Labour market trajectories derived from annual trends in full-time employment (derived from NOMIS Official Labour Market Statistics, ONS) from 2004 to 2014 Average regional income lost per working adult due to welfare reforms | • Age • Gender • Ethnicity • Marital status • Living alone • Social grade • Carstairs neighbourhood deprivation • Health selective migration prior to study period controlled for with data on Local Authority of residence in 2001 and 2011 | Population representative sample used Linked data on prescribing No data on diagnosis of depression so some prescriptions may not be for mental illness Unable to control for differences between prescribers in treating depression with anti-depressants | People living in areas with low or declining levels of full-time employment |

(continued on next page)

Table 3 (continued)

| Citation (Country) | Study Details (purpose, design, mental health outcomes) | Study characteristics (source of data, details of sample, dates of data collection, analysis) | Mental health measurement tools and methods | Specific economic measures | Covariates / confounders | Author assessment of strengths and limitations of measurement | Groups identified as particularly vulnerable |
|-----------------------------|--|--|---|--|---|---|---|
| Clair, 2022 [15] (UK) | Explore the impact of cold homes on mental health, the role of gender as a mediator and the effect of transitioning into a cold home Household panel study Outcomes: mental distress Note: During the study period, energy prices and housing costs were rising | 2009 to 2014 Analysis: Multilevel logistic regression and mediation analysis Data from UKHLS 2009 to Dec 2021 Eligible participants entered the study at wave 1; ≥ 16 yrs; had data for at least 3 waves Sample stratified into: Good mental health and lived in warm home at wave 1 ($n = 21,281$) Borderline mental health and lived in warm home at wave 1 ($n = 2258$) Analysis: multilevel discrete-time event models | GHQ-12 (Mental health distress) (Checked with SF-12 MCS) | Cold home measured by responses to question: “In winter are you able to keep this accommodation warm enough” | <ul style="list-style-type: none"> • Age • Gender • Ethnicity • Region • Employment status • Diagnosis of longstanding condition or disability • Adjusted income quartile • Highest educational qualification • Household composition • Financial situation • Household tenure • Building type • Housing payment arrears | Observed that it is possible that mental distress perceptions of temperature; that different people may experience household temperature differently | People transitioning into living in cold homes |
| Curtis, 2021 [16] (England) | Determine if the risk of worsening mental health was greater for people living in the most deprived areas in terms of unemployment and impact of austerity policies Household panel survey Outcomes: Trends in self-reported mental health | Data from UKHLS waves 3 to 7, 2011 to 2017 Eligible participants were ≥ 16 yrs; living in England; had full data on the variables of interest. 17,212 participants Maximum likelihood multinomial logit models | SF-12 MCS (version 2) | Geographical variables to categorise place of residence according to socio-geographic aspects (employment domain of the Index of Multiple Deprivation; average income loss per person of working age due to welfare reforms; rural / urban; West Midlands/ other) Indicator showing whether someone had moved to a location in a different category | <ul style="list-style-type: none"> • Gender • Age group • Ethnicity • Living with a partner • Occupational social class • Income in the month prior to interview • Employment status • Housing tenure at wave 4 • In receipt of welfare benefits at wave 4 (other than child benefits and state pensions) • Change in socio-economic status | Using a sub-sample from the panel survey means it may not no longer be representative of England as a whole | People who were more deprived Living in economically disadvantaged neighborhoods Living in areas most impacted by austerity policies Becoming unemployed Moving onto welfare benefits |
| Kim, 2022 [17] (England) | Examine the impact of the announcement (April 2012) and implementation (April 2013) of the spare room subsidy (“bedroom tax”) on psychological distress in social housing renters Quasi-experimental study with | Data from UKHLS 2010 to 2014 for main analysis and up to 2017 for sensitivity tests. Eligible participants were living in social housing; had pre- and post- policy data; were over 16 and under state pension age; lived in England | GHQ-12 (Mental health distress) | None | <ul style="list-style-type: none"> • Observed and unobserved individual-level time-invariant characteristics (e.g., immigration status and family history of illnesses) • Regional differences | Pre- and post- design and use of matching and appropriate analysis enhances ability to determine causality Other covariates not captured may confound the treatment effects of an intervention GHQ-12 is limited compared | People of lower socioeconomic status |

(continued on next page)

Table 3 (continued)

| Citation (Country) | Study Details (purpose, design, mental health outcomes) | Study characteristics (source of data, details of sample, dates of data collection, analysis) | Mental health measurement tools and methods | Specific economic measures | Covariates / confounders | Author assessment of strengths and limitations of measurement | Groups identified as particularly vulnerable |
|----------------------------|---|---|---|--|---|--|---|
| | matched controls Outcomes: psychological distress | 2078 eligible participants, of which 422 were living in underoccupied housing and were affected by the policy prior to April 2013 ('treatment group') and 1926 were not affected. Final matched sample = 412 treatment group and 412 controls. Difference-in-differences analysis | | | | with formal mental health assessment from a trained health care professional Loss to follow-up is common in panel surveys, which may bias estimates if different between control and treatment arms. Matching may reduce representativeness of the sample Treatment status was assumed based on household characteristics prior to the intervention period; household characteristics may be vulnerable to misreporting or may change over time Other studies examine associations between welfare reforms and mental health during economic crises but not the impact of a very specific policy using a quasi-experimental design | |
| Saville, 2021 [18] (Wales) | Examine whether pre-economic crisis aggregated social capital within a region (ecological social capital) was protective against post-economic crisis depression Repeated cross-sectional survey Outcomes: self-reported treatment for depression | Data from the Welsh Health Survey (now superseded by the National Survey for Wales) from all 12 waves (2003/4–2015) Overall sample size 180,462 (minimum 12,689, maximum 15,699 per year) Analysis: multilevel regression with post-stratification estimates | Self-reported treatment for depression | Data from Living in Wales survey used to estimate pre-crisis ecological social capital for each Middle Super Output Area of Wales Classification of occupation from National Statistics Socio-economic classification | <ul style="list-style-type: none"> • Age band • Gender • Area population density • Occupation type | Welsh Health Survey represents a relatively large proportion of the Welsh population Self-reported treatment for depression may not be accurate | Not reported, although women had higher levels of self-reported treatment for depression at all timepoints Rates of treatment for depression increased with age up to the age of 50yrs |
| Thomson, 2022 [3] (UK) | Compare total and direct effects of unemployment on mental health, examining the impact of income Household panel survey Outcomes: prevalence of poor mental health | Data from UKHLS 2019 Eligible participants were all working aged adults aged 25 to 64. 45,497 participants Analysis: double-robust marginal structural modelling | GHQ-12 (Mental health distress) | Binary measure of whether someone was employed or unemployed Continuous income Binary measure of poverty (living above or below household equivalised income <60 % median after housing costs) | <ul style="list-style-type: none"> • Age • Gender • Ethnicity • Education • Socio-economic status • Marital status • Number of dependents • Physical health • Location | Unable to include measures for other income-related factors such as wealth, savings, or debt since these were not included in all survey waves | Not reported |

(continued on next page)

Table 3 (continued)

| Citation (Country) | Study Details (purpose, design, mental health outcomes) | Study characteristics (source of data, details of sample, dates of data collection, analysis) | Mental health measurement tools and methods | Specific economic measures | Covariates / confounders | Author assessment of strengths and limitations of measurement | Groups identified as particularly vulnerable |
|-------------------------|---|--|--|---|---|---|--|
| Wickham, 2020 [19] (UK) | Examine the effect of the introduction of Universal Credit on mental health Quasi-experimental study Outcomes: psychological distress (NB: Included due to study design despite date. Not included in reviews, identified via grey literature) | Data from UKHLS between 2009 and 2018 Eligible participants were aged between (16 to 64yrs), had data on variables of interest, did not live in Northern Ireland, were not on disability benefit 52,187 participants Intervention group for each survey wave were participants who answered that they were unemployed; control participants were those who answered other than unemployed Analysis: Difference-in-differences analysis | GHQ-12 (Mental health distress) (Checked using SF-12 MCS) | Self-reported unemployment Data on when Universal Credit was introduced into an area where each respondent lived | Country of residence Age Gender Education status Marital status | Natural policy experiment approach | Not reported |

GDP = Gross Domestic Product; GHQ-12 = 12-item General Health Questionnaire; NHS = National Health Service; ONS = Office for National Statistics; SF-12 MCS = Short Form 12 Mental Health Component Summary; SHARE = Survey of Health, aging and Retirement in Europe; UKHLS = Understanding Society; The UK Household Longitudinal Study.

ved to be of variable quality and dependent on the structure of services and record keeping. Any reorganisation of mental health care services post-onset of an economic crisis can introduce bias by potentially altering who can access services and may disproportionately affect the most vulnerable. The use of prescribing data can introduce bias since the availability of drugs may vary, different prescribers may have different prescribing thresholds which may affect different population sub-groups differently, and the same drug may be given for conditions that are not mental-health related. One review [9] commented that people seeking medical care were not necessarily those most impacted by economic crises. Where data from medical records have been aggregated, individual behaviour cannot be inferred, and the data cannot capture variation in services between population sub-groups. For example, an economic crisis may increase suicide rates for some population sub-groups but if suicide prevention strategies are provided at the same time, these may benefit different population sub-groups and the net result may be a zero increase. However, it was observed that an advantage of aggregated data was that they reflect the environmental effects of changes in the economy beyond an individual’s personal circumstances.

3.4. Vulnerable groups

All the reviews and grey literature and some primary studies identified population groups whose mental health was most likely to be affected by an economic crisis. All the mental health measurement tools were used for this purpose, except for the WEMWBS. All the reviews presented the data narratively and neither reviews nor primary studies commented on the suitability of the methods and tools to measure mental health in vulnerable groups. However, data were presented on people living on lower incomes, unemployed or at risk of unemployment, in financial or housing insecurity or living in more deprived areas prior to the crisis. The reviews also examined data stratified by gender and age. Other groups identified as being vulnerable were people with pre-existing mental health problems, people living in households with dependent children, and migrants. One primary study [21] also identified adults with a disability or long-term illness as being at risk. No included study or review presented data from people from minority ethnic groups.

4. Discussion

4.1. Summary of the findings

This rapid review aimed to answer two questions:

4.2. What methods and tools are available and appropriate for monitoring the impact of the cost-of-living crisis on mental health?

We identified multiple studies that examined the impact of economic crises on mental health over time using national surveys from many different countries using different methods. Data from population-level panel surveys may be most useful. These large surveys incorporate validated mental health tools and questions to determine financial security at household or individual level and may be linked to medical records (e.g. Clinical Practice Research Datalink (CPRD) data in UK). Many studies are observational in design, but we also identified two relevant quasi-experimental case-control studies to examine the impact of specific welfare policies in the UK (the “Bedroom tax” and the introduction of Universal Credit). A quasi-experimental study design could also be suitable for measuring the impact of a specific public health initiative that has a clear roll-out date.

Table 4
Summary of grey literature.

| Citation (country) | Details (aim, study type, organisation) | Mental health measurement tools and methods | Specific economic measures | Author assessment of methods | Groups identified as particularly vulnerable |
|---|--|--|---|--|---|
| Clark, 2022 [20] (UK) | Report examining associations between insecurity and anxiety in the UK Uses data from GP records, national surveys (panel surveys and repeated cross-sectional), although some data used is cross-sectional being taken from a single year. Joseph Rowntree Foundation | <ul style="list-style-type: none"> • GHQ –12 • SF-12 • Warwick-Edinburgh mental well-being scale (WEMWBS)¹ • Single anxiety score • Medical records: • GP recording diagnosis of anxiety from: Clinical Practice Research Datalink • The Health Improvement Network • Population survey: • UKHLS • Annual Population Survey | Employment data: <ul style="list-style-type: none"> • Office for National Statistics • UKHLS • Annual Population Survey Housing data: <ul style="list-style-type: none"> • Labour Force Survey • English Housing Survey Household savings: <ul style="list-style-type: none"> • Wealth and Assets Survey • Family Resources Survey Social capital: <ul style="list-style-type: none"> • Office for National Statistics Deprivation: <ul style="list-style-type: none"> • Index of Multiple Deprivation | Prescribing data needs care with interpretation since the availability of drugs may vary and different prescribers may have different prescribing thresholds and the same drug may be given for different conditions Noted that UK official surveys do not always ask about both mental health and household economics in detail, may miss specific economic details and data from different surveys cannot be linked The authors report that UKHLS has 10,000 s of respondents and has information on both health and economics, although there are difficulties in identifying people in insecure work or forced freelancers UKHLS allows individuals to be tracked over time to determine how different events affect mental health Overall, the authors report that official statistics and secondary sources on mental health and financial stress are easily available in the UK | Living in debt Living in rental accommodation (especially social renters) Less than £1000 in savings or expecting less than enough as a pension |
| Preece 2019 [11] (Wales) | Policy report / rapid review Examine the relationship between housing insecurity and mental health, identify key lessons and relate to Wales Post 2008 economic crisis Rapid review: literature searched from 2008 – 2019 39 studies identified from searches, plus an unspecified number from handsearching UK Collaborative Centre for Housing Evidence | Mental health measurement tools: <ul style="list-style-type: none"> • GHQ-12 (Mental health distress) • Population survey: • UKHLS | Landlord possession rates and orders Mortgage repossession rates | Although UKHLS data has been used there is limited data on housing characteristics, and it was not possible to control for housing condition Authors note that data gaps around housing and mental health is not drawn specifically from Wales (evidence is mainly from Australia and England) and may not be generalisable | Households with dependant children People receiving means tested benefits Migrant groups People experiencing socio-economic deprivation People living with existing mental health problems |
| Office for National Statistics, 2022 [21] (Great Britain) | To examine depression in the context of rising cost-of-living Brief report describing repeated cross-sectional data collected fortnightly Office for National Statistics | Mental health measurement tool: <ul style="list-style-type: none"> • PHQ-8 (Depression) • Population survey: • The Opinions and Lifestyle Survey | Specific questions asked during survey on e.g., housing tenure, ease of paying bills, savings etc | Authors make a point of stating that no causal link between rising cost-of-living and depression is suggested | Economically inactive because of long-term sickness Unpaid carers for 35 or more hours a week Disabled adults Living in the most deprived areas of England Young adults aged 16 to 29 years Single person household Women Living in rental accommodation |

GHQ-12 = 12-item General Health Questionnaire; PHQ-8 = 8-item Public Health Questionnaire; SF-12 = Short Form 12; UKHLS = Understanding Society: The UK Household Longitudinal Study.

¹ Also used in UKHLS, although data from this questionnaire was not utilised in this study.

4.3. Are methods and mental health measurement tools suitable for specific, vulnerable populations?

All the reviews and most primary studies found people living in financial insecurity were at higher risk of poor or worsening mental health and most examined gender differences. It therefore appears that the methods and tools are suitable for use in men and women, across the socio-economic spectrum, living in different housing tenures and with different employment status. However, it is possible that some

potentially vulnerable groups are not well represented when using the methods and data sources identified. Few studies presented data from other groups that are often marginalised such as migrants and people living with disabilities or long-term poor health. We also identified no studies that examined whether people from minority ethnic groups were more at risk of deteriorating mental health during economic crises. It is therefore not clear as to whether the methods identified are suitable for all marginalised populations.

Table 5
Study designs and main characteristics.

| Study designs | Study characteristics | Examples of data sources of relevance to Wales ¹ |
|---|---|---|
| Cohort studies | Not necessarily representative of the population Self-reported data is collected but may include data-linked with other records, including medical records Longitudinal, repeated measures, following the same individuals over time, strengthening the evidence for causality Loss to follow-up is common Not all known confounders may be available in the dataset | None identified in the evidence presented, but an example is UK Biobank |
| Household or individual panel surveys | Representative of the population Self-reported data is collected but may include data-linked with other records, including medical records Longitudinal, repeated measures, following the same households or individuals over time, strengthening the evidence for causality Loss to follow-up is common Not all confounders may be available in the dataset | Understanding Society: the UK Household Longitudinal Study (UKHLS) Scottish Longitudinal Study (SLS) |
| National and regional repeated cross-sectional surveys | Representative of the population Different sample is used for each wave Self-reported measures are typically used Cannot infer causality, although sophisticated analysis techniques may strengthen inference Not all confounders may be available in the dataset | National Survey for Wales ONS Opinions and Lifestyle Survey |
| Quasi-experimental / case-control using panel survey data | Design reduces confounding, especially if matched samples are derived, and allows causality to be inferred Use of a sub-set of the main sample reduces overall representativeness It may not be straightforward to derive robust exposure variables Loss to follow-up may introduce bias if losses are different between arms | Data used from UKHLS |
| Studies using real-world longitudinal data (e.g., from medical records) | Does not depend on self-reported data Quality depends on standards of record keeping Reorganisation of services after an event such as an economic crisis or introduction of public health intervention may introduce bias since availability may change independent of need Use of aggregate data does not capture individual behaviour or variation in service provision between population sub-groups | CRPD Hospital Episode Statistics THIN |

Table 5 (continued)

| Study designs | Study characteristics | Examples of data sources of relevance to Wales ¹ |
|--------------------------------|---|---|
| Ecological time-series studies | Aggregated data are often easily accessible but may be of unknown quality Causality cannot be inferred | Hospital Episode Statistics ONS suicide rates |

CPRD = Clinical Practice Research Datalink; ONS = Office for National Statistics; THIN = The Health Improvement Network; UKHLS = Understanding Society; UK Household Longitudinal Study.

¹ The Catalogue of Mental Health Measures provides a comprehensive summary of mental health and wellbeing data available from population-level longitudinal studies conducted in Britain (<https://www.catalogumentalhealth.ac.uk/>).

4.4. Strengths and limitations of this rapid review

Evidence for this rapid review has been drawn from reviews that included over 300 studies, and from primary studies and grey literature. We identified appropriate available population level datasets that include both validated mental health tools and collect socioeconomic outcomes and identified appropriate study designs and provided information about their strengths and limitations.

However, our rapid review has limitations. All the included reviews summarised data from individual studies narratively and varied in the detail they provided on study designs and strengths and limitations. The identification of vulnerable groups appeared not to be systematic in most reviews and it is possible that data on vulnerable groups were not captured. There is a strong overlap in the literature with studies describing the impact of economic crises on mental health and the impact of any austerity policies that followed the Great Recession on mental health. We did not include austerity as a search term, and it is possible that relevant studies were excluded. However, the purpose of this review was not to examine the evidence that either economic crises or austerity policies affect mental health, but to describe available methods of measuring changes to population mental health in response to public health intervention. In addition, although the search strategy was comprehensive and included evidence from peer-reviewed articles and from grey literature, we had to exclude much of the identified grey literature due to lack of methodological detail and lack of information on data sources and measurement tools and, due to challenges in searching grey literature, it is likely that we did not identify all relevant documents. However, our overall aim was to identify the range of relevant methodological approaches (or population-level measures) used rather than identify all papers reporting the use of each approach. It is possible that there are datasets and tools used by charities and institutions that we did not identify.

4.5. Implications for policy and research

Our rapid review has identified existing methods and tools likely to be suitable for measuring the impact of public health initiatives on mental health in people from different socio-economic groups during a cost-of-living crisis. However, it is unclear as to whether the identified methods and tools adequately capture data from people from minority ethnic groups, who already experience disparities in mental health care [34]. The reported evidence did not allow us to determine whether the methods used to capture ethnicity of participants are inadequate, or whether recruitment of people from minority ethnic groups was low. In the short-term, reports and studies using panel surveys such as UKHLS or that use medical records should present data on ethnicity, or highlight where it is missing, and plan to stratify analyses by ethnicity if possible. In the longer-term it may be necessary to develop new methods to better capture data from people from minority ethnic groups.

Table 6
Mental health measurement scales.

| Name of measurement tool (abbreviation) | (Author(s), year) | Domains | Number of items | Target population | Time to complete | Comments |
|---|------------------------------|--|-----------------|----------------------------------|------------------|--|
| Used in UK surveys | | | | | | |
| 12-item General Health Questionnaire GHQ-12 | (Goldberg et al. 1997) [23] | Presence of mental health disorders (depression, anxiety, somatic symptoms, and social withdrawal) | 12 | General and clinical populations | 5–10mins | Used in UKHLS Survey |
| Short form 12 Mental Health Component Summary (SF-12) (Derived from SF-36) | (Ware et al. 1996) [24] | Psychological distress | 12 | General and clinical populations | 2–5mins | Used in UKHLS Survey Licenced by QualityMetric |
| Warwick-Edinburgh Mental Well-being Scale (WEMWBS) | (Tennant et al. 2007) [25] | Mental Wellbeing | 14 | General and clinical populations | 5–10mins | Used in UKHLS Survey and National Survey for Wales |
| Patient Health Questionnaire depression scale (PHQ-8) | (Kroenke et al. 2009) [26] | Presence of depression | 8 | General and clinical populations | | Used in ONS Opinions and Lifestyle Survey Based on PHQ-9 with item asking about suicidal or self-injurious thoughts removed |
| Other measurement tools | | | | | | |
| 5-item Mental Health Inventory (MHI-5) (Derived from SF-36) | (Berwick et al. 1991) [27] | Anxiety and depression | 5 | General population | 5mins | |
| 36-item Short Form Health Survey Questionnaire Mental Component Summary (SF-36 MCS) | (Brazier et al. 1992) [28] | Psychological distress | 35 | General and clinical populations | 5–10mins | Developed by the RAND corporation. SF-36 v1 is in the public domain SF-36v2 is licenced by QualityMetric |
| Centre for Epidemiological Study Depression Scale (CES-D) | (Radloff 1977) [29] | Depressive symptoms | 20 | General and clinical populations | Up to 10mins | In public domain |
| EURO-D | (Prince et al. 1999) [30] | Depressive symptoms | 12 | General population ≥65yrs | 5mins | Used in the Survey of Health, ageing and Retirement in Europe (SHARE) |
| Goldberg Depression and Anxiety Score | (Goldberg et al. 1988) [31] | Presence of anxiety and depression | 18 | General and clinical populations | 5–10mins | In public domain |
| Hopkins Symptom Checklist (HSCL-25) | (Hesbacher et al. 1980) [32] | Presence of anxiety and depression | 25 | General practice patients | 5–10mins | Widely translated and validated in multiple languages |
| Kessler-10 | (Furukawa et al. 2003) [33] | Anxiety and depression over the past 4 weeks | 10 | General population | Unclear | Developed for an Australian population, validated in other populations |
| Short form perceived stress scale (PSS-4) (Goldberg & Hillier 1979) | (Cohen et al. 1983) [34] | Psychological stress | 4 | General population | 5mins | Not a diagnostic instrument so there are no cut-offs. |

UKHLS = Understanding Society: The UK Household Longitudinal Study.

5. Conclusion

Internationally a wide variety of data sources and datasets has been used to explore the impact of economic crises on mental health. Both medical records and longitudinal individual and household surveys that include validated brief mental health measurement tools are available. The surveys aim to be representative of the population and collect information on other health-related and socio-economic outcomes and using them to measure the effects of public health policy is efficient. The choice of study design is likely to be a pragmatic decision based on the exact research question and availability of data, although care should be taken if data on important confounders are missing.

Funding

Health Technology Wales was funded for this work by the Health and Care Research Wales Evidence Centre, itself funded by Health and Care Research Wales on behalf of Welsh Government.

CRedit authorship contribution statement

Clare England: Conceptualization, Formal analysis, Investigation, Methodology, Writing – original draft, Writing – review & editing. **David Jarrom:** Conceptualization, Methodology, Supervision, Writing –

review & editing. **Jennifer Washington:** Conceptualization, Investigation, Methodology. **Elise Hasler:** Conceptualization, Investigation, Methodology. **Leona Batten:** Investigation, Writing – review & editing. **Adrian Edwards:** Conceptualization, Methodology, Writing – review & editing. **Ruth Lewis:** Conceptualization, Methodology, Supervision, Writing – review & editing.

Declaration of competing interest

The authors declare they have no conflicts of interest to report.

Acknowledgements

The authors would like to thank James Burgess, Brendan Collins, Alison Cooper, Jacob Davies, Rhiannon Tudor Edwards, Rebecca Masters, and Nigel Pearson for their contributions during stakeholder meetings in guiding the focus of the review and interpretation of findings.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.healthpol.2024.105062](https://doi.org/10.1016/j.healthpol.2024.105062).

References

- [1] Roberts M., Petchey L., Challenger A., Azam S., Masters R., Peden J. The cost to life: how soaring living costs affect people's health and wellbeing 2022 [Available from: <https://research.senedd.wales/research-articles/the-cost-to-life-how-soaring-living-costs-affect-people-s-health-and-wellbeing/>].
- [2] Guerra O, Eboeime E. The impact of economic recessions on depression, anxiety, and trauma-related disorders and illness outcomes-a scoping review. *Behav Sci (Basel)* 2021;11(9):31.
- [3] Thomson RM, Kopasker D, Leyland A, Pearce A, Katikireddi SV. To what extent does income explain the effect of unemployment on mental health? Mediation analysis in the UK Household Longitudinal Study. *Psychol Med* 2022;1–9.
- [4] Hamel C, Michaud A, Thuku M, Skidmore B, Stevens A, Nussbaumer-Streit B, et al. Defining rapid reviews: a systematic scoping review and thematic analysis of definitions and defining characteristics of rapid reviews. *J Clin Epidemiol* 2021; 129:74–85.
- [5] Garrity C, Gartlehner G, Nussbaumer-Streit B, King VJ, Hamel C, Kamel C, et al. Cochrane Rapid Reviews Methods Group offers evidence-informed guidance to conduct rapid reviews. *J Clin Epidemiol* 2021;130:13–22.
- [6] Frاسquilho D, Matos MG, Salonna F, Guerreiro D, Storti CC, Gaspar T, et al. Mental health outcomes in times of economic recession: a systematic literature review. *BMC Public Health* 2016;16(1):115.
- [7] Glonti K, Gordeev VS, Goryakin Y, Reeves A, Stuckler D, McKee M, et al. A systematic review on health resilience to economic crises. *PLoS ONE* 2015;10(4): e0123117.
- [8] Saez M, Barcelo MA, Saurina C, Cabrera A, Daponte A. Evaluation of the biases in the studies that assess the effects of the great recession on health. a systematic review. *Int J Environ Res Public Health* 2019;16(14):11.
- [9] Silva M, Resurreccion DM, Antunes A, Frاسquilho D, Cardoso G. Impact of economic crises on mental health care: a systematic review. *Epidemiol Psychiatr Sci* 2018;29:e7.
- [10] Martin-Carrasco M, Evans-Lacko S, Dom G, Christodoulou NG, Samochowicz J, Gonzalez-Fraile E, et al. EPA guidance on mental health and economic crises in Europe. *Eur Arch Psychiatry Clin Neurosci* 2016;266(2):89–124.
- [11] Preece DJ, Bimpson DE. Housing insecurity and mental health in wales: an evidence review. European: UK Collaborative Centre for Housing Evidence (CaCHE); 2019.
- [12] Alvarez-Galvez J, Suarez-Lledo V, Salvador-Carulla L, Almenara-Barrrios J. Structural determinants of suicide during the global financial crisis in Spain: integrating explanations to understand a complex public health problem. *PLoS ONE* 2021;16(3):e0247759.
- [13] Aretz B. The short- and long-term effects of the Great Recession on late-life depression in Europe: the role of area deprivation. *Soc Sci Med* 2022;294:114697.
- [14] Cherrie M, Curtis S, Baranyi G, Cunningham N, Dibben C, Bamba C, et al. A data linkage study of the effects of the Great Recession and austerity on antidepressant prescription usage. *Eur J Public Health* 2021;31(2):297–303.
- [15] Clair A, Baker E. Cold homes and mental health harm: evidence from the UK Household Longitudinal Study. *Soc Sci Med* 2022;314(pagination).
- [16] Curtis S, Cunningham N, Pearce J, Congdon P, Cherrie M, Atkinson S. Trajectories in mental health and socio-spatial conditions in a time of economic recovery and austerity: a longitudinal study in England 2011–17. *Soc Sci Med* 2021;270:113654.
- [17] Kim C, Teo C, Nielsen A, Chum A. What are the mental health consequences of austerity measures in public housing? A quasi-experimental study. *J Epidemiol Commun Health* 2022;15:15.
- [18] Saville CWN. Ecological social capital does not predict geographical variance in increases in depression following the 2008 financial crisis. *Br J Psychol* 2021;112(1):163–79.
- [19] Wickham S, Bentley L, Rose T, Whitehead M, Taylor-Robinson D, Barr B. Effects on mental health of a UK welfare reform, Universal Credit: a longitudinal controlled study. *Lancet Public Health* 2020;5(3):e157–ee64.
- [20] Clark T, Wenham A. Anxiety nation? economic insecurity and mental distress in 2020s Britain. Joseph Rowntree Foundation; 2022.
- [21] Office for National Statistics. Cost of living and depression in adults, Great Britain: 29 September to 23 October 2022 2022 [Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/mentalhealth/articles/costoflivinganddepressioninadultsgreatbritain/29septemberto23october2022>].
- [22] Goldberg DP, Gater R, Sartorius N, Ustun TB, Piccinelli M, Gureje O, et al. The validity of two versions of the GHQ in the WHO study of mental illness in general health care. *Psychol Med* 1997;27(1):191–7.
- [23] Ware Jr J, Kosinski M, Keller SD. A 12-Item Short-Form Health Survey: construction of scales and preliminary tests of reliability and validity. *Med Care* 1996;34(3):220–33.
- [24] Tennant R, Hiller L, Fishwick R, Platt S, Joseph S, Weich S, et al. The warwick-edinburgh mental well-being scale (WEMWBS): development and UK validation. *Health Qual Life Outcomes* 2007;5(1):63.
- [25] Kroenke K, Strine TW, Spitzer RL, Williams JB, Berry JT, Mokdad AH. The PHQ-8 as a measure of current depression in the general population. *J Affect Disord* 2009; 114(1–3):163–73.
- [26] Berwick DM, Murphy JM, Goldman PA, Ware JE, Barsky AJ, Weinstein MC. Performance of a five-item mental health screening test. *Med Care* 1991;29(2): 169–76.
- [27] Brazier JE, Harper R, Jones NM, O'Cathain A, Thomas KJ, Usherwood T, et al. Validating the SF-36 health survey questionnaire: new outcome measure for primary care. *Br Med J* 1992;305(6846):160–4.
- [28] Radloff LS. The CES-D Scale: a Self-Report Depression Scale for Research in the General Population. *Appl Psychol Meas* 1977;1(3):385–401.
- [29] Prince MJ, Reischies F, Beekman ATF, Fuhrer R, Jonker C, Kivela SL, et al. Development of the EURO-D scale – a European Union initiative to compare symptoms of depression in 14 European centres. *Br J Psychiatry* 1999;174(4): 330–8.
- [30] Goldberg D, Bridges K, Duncan-Jones P, Grayson D. Detecting anxiety and depression in general medical settings. *Br Med J* 1988;297(6653):897–9.
- [31] Hesbacher PT, Rickels K, Morris RJ, Newman H, Rosenfeld H. Psychiatric illness in family practice. *J Clin Psychiatry* 1980;41(1):6–10.
- [32] Furukawa TA, Kessler RC, Slade T, Andrews G. The performance of the K6 and K10 screening scales for psychological distress in the Australian National Survey of Mental Health and Well-Being. *Psychol Med* 2003;33(2):357–62.
- [33] Cohen S. Perceived stress in a probability sample of the United States. The social psychology of health. The claremont symposium on applied social psychology. Thousand Oaks, CA, US: Sage Publications, Inc; 1988. p. 31–67.
- [34] Ahmad G, McManus S, Cooper C, Hatch SL, Das-Munshi J. Prevalence of common mental disorders and treatment receipt for people from ethnic minority backgrounds in England: repeated cross-sectional surveys of the general population in 2007 and 2014. *Br J Psychiatry* 2022;221(3):520–7.