

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/176083/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Cooke, Steven J., Lynch, Abigail J., Abell, Robin, Piczak, Morgan L., Arthington, Angela H., Thieme, Michele, Perry, Denielle, Britton, J. Robert, Dalu, Tatenda, Birnie-Gauvin, Kim, Ormerod, Steve J. , Matuk, Fernanda Ayaviri, Raghavan, Rajeev and Smol, John P. 2025. A roadmap for implementing the emergency recovery plan for freshwater biodiversity. *Environmental Reviews*

Publishers page:

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



A Roadmap for Implementing the Emergency Recovery Plan for Freshwater Biodiversity

Preface for Special Issue

Steven J. Cooke¹, Abigail J. Lynch², David Tickner³, Robin Abell⁴, Morgan L. Piczak¹, Angela H. Arthington⁵, Michele Thieme⁶, Denielle Perry⁷, J. Robert Britton⁸, Tatenda Dalu⁹, Kim Birnie-Gauvin¹⁰, Steve J. Ormerod¹¹, Fernanda Ayaviri Matuk¹², Rajeev Raghavan¹³, and John P. Smol¹⁴

¹Department of Biology and Institute of Environmental and Interdisciplinary Science, Carleton University, 1125 Colonel By Dr., Ottawa, ON, K1S 5B6, Canada; <https://orcid.org/0000-0002-5407-0659>

²United States Geological Survey, National Climate Adaptation Science Center, 12201 Sunrise Valley Drive MS 516, Reston, VA, 20192, USA, <https://orcid.org/0000-0001-8449-8392>

³WWF-UK, Living Planet Centre, Woking, GU21 4LL, UK, <https://orcid.org/0000-0001-5928-0869>

⁴The Nature Conservancy, Arlington, VA, USA

⁵Australian Rivers Institute, Griffith University, Nathan, Queensland, 4111, Australia; <https://orcid.org/0000-0001-5967-7954>

⁶WWF, Washington, DC, USA

⁷Free-flowing Rivers Lab, School of Earth and Sustainability, Northern Arizona University, 624 Knoles Dr, Flagstaff, AZ, 86011, USA; <https://orcid.org/0000-0002-3603-3731>

⁸Fish Ecology and Conservation Research Cluster, Bournemouth University, Poole, BH12 5BB, United Kingdom

⁹School of Biology and Environmental Sciences, University of Mpumalanga, Nelspruit 1200, South Africa

¹⁰Section for Freshwater Fisheries and Ecology, Technical University of Denmark, Vejlssøvej 39, 8600 Silkeborg, Denmark; <https://orcid.org/0000-0001-9242-0560>

¹¹Water Research Institute, Cardiff School of Biosciences, Cardiff University, CF10 3AX, UK; <https://orcid.org/0000-0002-8174-302X>

¹²Federal Institute of Minas Gerais, Geography, Av. Primeiro de Junho, 1043 - Centro, São João Evangelista, MG, 39705-000, Brazil; <https://orcid.org/0000-0002-5152-5566>

¹³Kerala University of Fisheries and Ocean Studies, Panangad, Kerala, India

¹⁴Paleoecological Environmental Assessment and Research Laboratory (PEARL), Department of Biology, Queen's University, Kingston, ON, K7L 3N6, Canada

Abstract - PREFACE PAPERS IN ENV REVIEWS DO NOT HAVE ABSTRACTS

Context

Although hyperbole abounds in the context of environmental issues (Katz 2011), it is entirely appropriate to use the word “crisis” when referring to the current state of freshwater biodiversity (Harrison et al. 2018). There is a growing body of studies (at the regional, national, and international scale) illustrating negative trends in freshwater biodiversity (e.g., He et al. 2019, IUCN 2019, Jahnig et al 2021; Isbell et al. 2023; Sayer et al. 2025), indicative of a global decline in essential ecosystem services (Harrison et al. 2018). The numbers are considerable with the World Wide Fund for Nature (WWF) Living Planet Index (LPI) revealing that average freshwater vertebrate populations are down 83% since 1970, a level of decline more than double that observed in marine and terrestrial systems (WWF 2022). The problems span taxa, ecosystems, regions, and issues. A recent International Union for the Conservation of Nature (IUCN) analysis found that ~1/4 of all freshwater fauna are threatened with extinction (Sayer et al. 2025). For example, 224 (44%) of the 511 freshwater molluscs in Europe are classified by the IUCN as ‘near threatened’ or ‘threatened’ species (Lopez-Lima et al. 2017). In Borneo, the majority of native mussel populations are now imperiled (Zieritz et al. 2018). Such declines are ubiquitous on a global

scale (Strayer et al. 2004). The extinction rate for freshwater fishes in North America is estimated to be 877 times greater than the background extinction rate, with an acceleration in loss since the 1950s (Burkhead 2012). Amphibians (the vast majority of which are freshwater-dependent) are the most imperiled taxonomic class on the planet, with a major extinction underway (Ceballos et al. 2020). —For some taxa, such as freshwater plants, we simply have no knowledge about their status (Lacoul and Freedman 2006). However, in Canada, for example, 11.7% and 17.9% of all freshwater species of plants and animals assessed were found to be ‘at risk’ or identified as being of ‘special concern’, respectively (Desforges et al. 2022). Beyond specific taxa, these declines are paralleled in freshwater habitats as well. Wetland loss from human activities exceeds at least 30% (Hu et al. 2017) and may possibly be higher than 50% (Davidson 2014), with at least 21% loss over the last three centuries alone (Fluet-Chouinard et al. 2023). Due to damming, only 37% of rivers longer than 1,000 kilometres remain free-flowing over their entire length (Grill et al. 2019). We will not belabour the point; these example statistics are representative of the overall state of freshwater ecosystems and their biodiversity, which is rather grim (Albert et al. 2021).

The threats facing freshwater biodiversity and ecosystems are numerous and often interacting in complex and unexpected ways (Ormerod et al. 2010; Dudgeon 2019) across spatial scales (Birk et al. 2020). Pollution, fragmentation, water withdrawal, invasive species, and over exploitation are among the most relevant, long-standing and emerging threats (Dudgeon et al. 2006; Reid et al. 2019). Climate change also serves as a “threat multiplier” (Smol 2010), whose interactions with other stressors can be synergistic (Dudgeon 2019; Dodson et al. 2020; Barbarossa et al. 2021). Inadequate governance and poor management of freshwater ecosystems and water resources either causes or compounds these threats.

Collectively, these threats do not just impact freshwater biodiversity and ecosystems, but also the people who depend upon them (Basurto et al. 2013). We acknowledge that essential human activities such as drinking water, electricity generation, agriculture and transportation require freshwater resources. Yet some freshwater resource development and exploitation achieve short-term benefits at the expense of long-term sustainability, and those benefits are often distributed unequally, with collateral impacts disproportionately affecting disadvantaged groups in society (e.g., Sherman et al. 2015). Freshwater biodiversity and ecosystems yield numerous ecosystem services, spanning from nutrition for some of the world’s most food insecure peoples to supporting livelihoods (Postel and Carpenter 1997; Lynch et al. 2016; Parmesan et al. 2022; Lynch et al. 2023). In fact, freshwater resources are the basis for many cultural practices and religions (Lynch et al. 2023). There is clear evidence that the impacts (direct and indirect) of humans and associated threats on freshwater ecosystem services are diminishing the values that can be derived in the future (Naiman and Dudgeon 2011; Dodds et al. 2013).

For years, the scientific community has been calling the attention to the threatened state of freshwater ecosystems and of their biodiversity loss (e.g., Strayer and Dudgeon 2010; Carpenter et al. 2011; Gleick 2018; Albert et al. 2021; Kuiper 2023). While there is little evidence that these calls have promoted substantial changes in trajectories for freshwater ecosystems and biodiversity (Dudgeon and Strayer 2025), a few initiatives show promise. For instance, WWF has spearheaded the development of the ‘Emergency Recovery Plan for Freshwater Biodiversity’ (see Tickner et al. 2020; herein, the “FW Emergency Recovery Plan”). The term ‘emergency’ was used intentionally to acknowledge the urgency needed in addressing the crisis. Overall, the FW Emergency Recovery Plan consists of six actions that, if embraced, would “bend the curve” (*sensu* Mace et al. 2018) of freshwater biodiversity loss globally (Figure 1). The actions are as follows:

Commented [KAT1]: Maybe (?) clearer if changed to:
These declines in specific taxa are paralleled in freshwater habitats as well.

1. Accelerating the implementation of environmental flows;
2. Improving water quality;
3. Protecting and restoring critical habitats;
4. Managing the exploitation of freshwater ecosystem resources, especially species and riverine aggregates;
5. Preventing and controlling non-native species invasions;
6. Safeguarding and restoring river connectivity.

The FW Emergency Recovery Plan has served as an effective framework for broad policy influence, such as in the lead up to the 15th Conference of the Parties of the Convention on Biological Diversity (CBD) (UN Water 2020). However, the original article by Tickner et al. (2020) contains insufficient information for those intending to use it to operationalize each of the six actions that would lead to the recovery of freshwater biodiversity. Accordingly, there is a need to take a deeper dive into each of these six actions to synthesize relevant evidence required to implement the recovery plan. Also, case studies can further exemplify how these actions are being implemented in both lower and higher income countries and thus inform delivery of international agreements and initiatives such as the CBD Kunming-Montreal Global Biodiversity Framework (see <https://www.cbd.int/doc/c/e6d3/cd1d/daf663719a03902a9b116c34/cop-15-l-25-en.pdf> and Cooke et al. 2023) and the Freshwater Challenge, launched by eight governments during the 2023 UN Water Summit (https://wwf.panda.org/wwf_news/?7942466/Launch-of-Freshwater-Challenge).

To that end, we assembled a collection of papers focused on a deeper examination of each of the six FW Emergency Recovery Plan actions. In addition to having a paper exploring barriers to and opportunities for implementation of each of those actions, we also included a paper focused on understanding and influencing the wider policy context for such efforts (Tickner et al. This Issue) as well as a paper that considers how to future-proof such actions (Lynch et al. This Issue). This preface provides the context and briefly summarizes key messages arising from each paper in the special issue. Underpinning all papers in this collection is the goal of ensuring that we include and amplify voices from around the globe that represent different regions, identities, knowledge systems, career stages, sectors, and roles. The core papers have accordingly been authored by a diverse group of experts and include case studies that attempt to highlight how the various high-level actions (and associated response measures) relate to different regional contexts.

Synopsis of Content in the Special Issue

Accelerating implementation of environmental flows

Arthington et al. (This Issue) outline threats arising from hydrological alterations caused by dams, freshwater overuse and climate change, and identify critical factors that enable recovery of freshwater biodiversity in flow-stressed rivers. These enabling factors include overarching environmental flows (e-flows) legislation and participatory governance, sustained financial and human resources, diverse stakeholder engagement and co-production of western and Indigenous knowledge, capacity training and research, and collaborative monitoring of ecological and social-economic outcomes. Biodiversity and societal outcomes from e-flow implementations can be strengthened by exploring trade-offs among water users, removing or retrofitting water infrastructure to facilitate e-flows and connectivity, and adaptations to address climate-change scenarios.

Improving water quality

This paper was not complete at time of writing this editorial but some key points were emerging from ongoing writing efforts. There are a range of response options that can reduce diverse types and sources of pollution and deliver multiple ~~other~~ benefits to progress ~~other~~ aspects of the FW Emergency Recovery Plan including (1) improvements in local to global policy; (2) better governance, enforcement, and regulation; (3) market instruments and innovative finance to recognise and incentivise more sensitive water use; (4) education and cultural encouragement to raise awareness of water quality issues; (5) more holistic management of water resource use from supply to waste disposal; and (6) nature-based solutions in the management of urban and rural catchments. Issues of pollution are shared between human well-being and freshwater biodiversity, so solutions will have wide-ranging benefits.

Commented [KAT2]: Mark Mallory commented:
It seems weird to start the "Improving Water Quality" section on p 3 with the caveat of something not being complete ... not sure that's necessary, I'd jettison the first line.

Protecting and restoring critical habitats

Piczak et al. (This Issue) synthesize the threats associated with habitat fragmentation, degradation, and loss, and outline specific response options to save freshwater biodiversity. These response options include legislating the protection of healthy and productive freshwater ecosystems; prioritizing habitats for protection and restoration; enacting durable protection; conserving habitat in a coordinated and integrated manner; engaging in evidence-based restoration using an adaptive management approach; ensuring that potential freshwater habitat alterations are mitigated or off-set; and enabling future-proofing of protection and restoration actions. Expected challenges involve: (1) institutional and management practices and laws (e.g., limited resources), (2) scientific uncertainties (e.g., regarding climate change), and (3) social and political goals (e.g., shifting political priorities).

Managing the exploitation of freshwater ecosystem resources, especially species and riverine aggregates

Cooke et al. (This Issue) outline the threats arising from the exploitation of freshwater biota and aggregate resources (e.g., sand, gravel, boulders) and identify response options to ensure that methods and levels of extraction are sustainable and allow recovery of over-exploited freshwater biodiversity and ecosystems. Response options for managing species exploitation include strengthening assessment and reporting, using science-based approaches to reduce overexploitation and support recovery, embracing community engagement, and building or tightening legislation. Response options for managing exploitation of freshwater aggregate resources include reducing demand for extraction, strengthening governance, reporting, and monitoring of environmental impacts, and promoting the restoration of degraded ecosystems or compensating for losses.

Preventing and controlling non-native species invasions

Britton et al. (This Issue) emphasise that the most effective management measure to protect freshwater biodiversity from non-native species is to prevent their introduction from occurring (e.g., inspecting and quarantining imports, decontaminating possible vectors). If this is unsuccessful, the early detection of non-native species enables the implementation of rapid response measures to prevent population establishment, dispersal, and impact (e.g., intense eradication efforts). Where these steps fail to prevent the invasion from progressing, management response options are then usually limited to methods that aim to control and/or contain the non-native species (e.g., sterile male release techniques). Developments in invasion risk assessment enable non-native species to be prioritised according to their invasion risk; for species that are already invasive, these assessments help ensure that the management responses are commensurate with the risks to freshwater biodiversity. Issues associated with non-native species in freshwaters include their presence often being a symptom of a degraded freshwater, rather than the species being the main driver of biodiversity loss, and so management requires more holistic interventions that aim to more generally restore the freshwater environment.

Safeguarding and restoring river connectivity

Thieme et al. (This Issue) summarize measures for maintaining and restoring river connectivity, including system-scale planning for energy and water resources to limit the loss of freshwater connectivity; putting in place protections for keeping critically important freshwater habitats connected; mitigating impacts on freshwater ecosystems via barrier design, fish passage, or implementation of environmental flows; and restoring freshwaters via barrier removal and reconnection of rivers, wetlands, and floodplains and via active management of groundwater recharge.

On Future Proofing Freshwater Biodiversity Protections

Lynch et al. (This Issue) warn that the FW Emergency Recovery Plan will not bend the curve of biodiversity loss if it only considers response options to address historic and current conditions. Uncertainty related to climatic, demographic, geopolitical, socio-economic, and cultural contexts make historical baselines somewhat less relevant and conventional conservation and restoration objective targets potentially maladaptive (Barnett and O'Neill 2010). 'Future-proofing' is a concept that explicitly addresses future uncertainties and surprises. It refers to the process of anticipating future events and developing methods to mitigate or minimize plausible stressors and shocks to a system (Rehman et al. 2017). Future-proofing the FW Emergency Recovery Plan can safeguard against future known uncertainties (e.g., climate change, emerging pollutants) and build resilience to withstand future unknown uncertainties (e.g., unexpected non-native species, algal blooms, diseases) and unanticipated surprises (see Table 1 in Lynch et al., This Issue for examples).

Commented [KAT3]: To make this clearer to reader, should this section be introduced as Lynch et al. (This issue) - rather than at end?

On Policy

Tickner et al. (This Issue) describe the key sectors that the international freshwater conservation and science communities should seek to influence to address underlying drivers of freshwater biodiversity loss, such as water resource management, agriculture, and energy. They describe major risks and opportunities emerging from current sector-specific policy discourses (e.g., public and private sector funding) and suggest priorities for research and advocacy interventions. Freshwater biodiversity issues cannot be addressed without significant inter-sectoral collaboration and coordinated policy development and action. There is much that can be done to embrace a more systemic, driver-focused approach to freshwater conservation research to ensure that freshwater biodiversity is considered in decisions related to both policy and business.

From Roadmap to Implementation

The evidence-base to guide management responses is ever-growing and, although some research gaps remain (Harper et al. 2021; Arthington 2021; Maasri et al. 2022), the bigger challenge lies in decision-making and implementation. We need, for example, to determine the most effective means of communicating the need for, and benefits of, action to different audiences including governments (from local to regional to national to international bodies), other decision makers, and diverse publics around the globe drawing on insights from behavioural and cognitive sciences (Toomey, 2023). There is also immediate need to embrace different knowledge systems (e.g., Indigenous ways of knowing, stakeholder knowledge) to ensure that all relevant forms of evidence are considered when identifying the best management measures for a given scenario. All relevant actor groups (including rights holders and their rights to self determination as outlined in the UN Declaration for the Rights of Indigenous Peoples; https://www.un.org/development/desa/indigenouspeoples/wp-content/uploads/sites/19/2018/11/UNDRIP_E_web.pdf) need to be included in decision-making processes (Birnle-Gauvin et al. 2023).

Twardek et al. (2021) and Cooke and Birnie-Gauvin (2022) emphasized the important role that will be played by practitioners who are front-line workers as it relates to freshwater protection, management, and restoration. While large-scale policy changes at the level of regional and global governance (e.g., UN, global agreements such as Conference of Parties (COP) Biodiversity; see Gonçalves and Hermoso 2022) can surely affect change, those governance-level efforts can only be as impactful as the on-the-ground actions that can be undertaken. Throughout the papers in the special issue, there are case studies from high-income and low-income regions that range from localized community-level initiatives to multi-national policy initiatives.

The freshwater biodiversity crisis is often overshadowed by other more visible issues. However, demonstrating the value of freshwater biodiversity (see Lynch et al. 2023) yields many stories that can be shared with the public and politicians, on the path to generating political will for action. We also need to capture and share success stories where governments, communities, and other organizations work alone or together to implement aspects of the FW Emergency Recovery Plan. Communication activities can use a combination of hope and fear to best elicit engagement (Kidd et al. 2019) and can be framed intelligently and adaptively to resonate with the values and priorities of target audiences in specific contexts. There is need for a coherent approach to planning effective communication efforts that can be applied in different contexts and with different audiences in mind. There is also an opportunity to use change models (see Reddy et al. 2017) to ensure that all elements of the pathway (e.g., from any communication or public engagement efforts through to behaviour change and associated actions related to the FW Emergency Recovery Plan) yield tangible improvements in freshwater biodiversity.

Conclusion

In this special issue, we have provided insights on how the world can implement the FW Emergency Recovery Plan (summarized in Figure 1). Notably, we have also provided examples (case studies) where actions have been implemented with some success. Given that we are facing a critical environmental state with freshwater ecosystems and species, all constructive efforts, no matter the scale, are helpful. However, actions that are conducted in a systematic manner (Linke et al. 2011) and across relevant scales (e.g., using a watershed approach; Brumm et al. 2022) are particularly relevant. We also acknowledge that resources for implementing the plan will be limited such that it will be necessary to prioritize efforts (see Linke et al. 2019). Keeping an eye on the future (i.e., future-proofing) in all efforts can help ensure that actions are durable (*sensu* Higgins et al. 2021) in a changing world.

The policies related to freshwater ecosystems likely benefit from adopting a deeper recognition of the connections between water, biodiversity, and people such that relevant policies, ranging from global to local scales (Dudgeon and Strayer 2025) and involving social actors of different sectors, are improved, created, or strengthened. Fortunately, freshwater biodiversity has recently received more attention in global policy arenas (Gonçalves and Hermoso 2022; Cooke et al. 2023) but more work is needed. Governance structures that support freshwater biodiversity protection and restoration are not sufficiently present in many global regions, making it difficult or impossible to effectively implement the FW Emergency Recovery Plan actions. Building capacity for informed action and strengthening governance remain priorities, particularly in the vulnerable regions where much of the freshwater biodiversity exists and is used by locals, and where the consequences of that loss will be even more profound for people and the planet.

Finally, we acknowledge that implementing the plan will require the work of many actors including scientists, practitioners, rights holders, stakeholders, politicians, business leaders, investors and voters. There are many ways for these individuals and groups to become involved in enacting the FW

Emergency Recovery Plan and bending the curve for freshwater biodiversity. We now know how to operationalize the plan; the next step is putting those strategies into action!

Positionality Statement

The special issue was coordinated by Cooke, Lynch, Tickner and Abell. Smol served as Editor for the papers in the collection. The other authors represent leads of various papers within the special issue or individuals that were particularly engaged in the process. The first author acknowledges the stewards of the lands and waters where we live, work, and play recognizing that many of us are settlers and are guests on the traditional territories of Indigenous peoples. We also acknowledge the varied conditions and contexts in which various contributors to our special issue engage with freshwaters - having different professional roles, active in varied socio-cultural and geo-political contexts, and operating in systems with varied capacity, governance stability, and freedoms. The freshwater biodiversity crisis impacts people around the globe and similarly to implement the Emergency Recovery Plan requires broad engagement across all continents, biomes, countries, and regions. We hope that the papers presented here serve as a roadmap for bending the curve for freshwater biodiversity.

References

- Albert, J. S., Destouni, G., Duke-Sylvester, S. M., Magurran, A. E., Oberdorff, T., Reis, R. E., ... & Ripple, W. J. (2021). Scientists' warning to humanity on the freshwater biodiversity crisis. *Ambio*, 50(1), 85-94.
- Allen, C.R., Garmestani, A.S. (Eds.), 2015. *Adaptive Management of Social-Ecological Systems*. Springer, New York, pp. 271.
- Arthington, A. H. (2021). Grand challenges to support the freshwater biodiversity emergency recovery plan. *Frontiers in Environmental Science*, 9, 118.
- Arthington, A. H., Tickner, D., McClain, M. E., Acreman, M. C., Anderson, E. P., Babu, S., et al. (This Issue). Accelerating environmental flows implementation to bend the curve of global freshwater biodiversity loss. *Environmental Reviews*. 00:000-000.
- Barbarossa, V., Bosmans, J., Wanders, N., King, H., Bierkens, M. F., Huijbregts, M. A., & Schipper, A. M. (2021). Threats of global warming to the world's freshwater fishes. *Nature Communications*, 12(1), 1701.
- Barnett J, O'Neill S (2010) Maladaptation. *Glob Environ Chang* 20:211–213.
<https://doi.org/10.1016/j.gloenvcha.2009.11.004>
- Basurto, X., Gelcich, S., Ostrom, E. The social-ecological system framework as a knowledge classificatory system for benthic small-scale fisheries. *Global Environmental Change*, 23 (6): 1366-1380, 2013.
<https://doi.org/10.1016/j.gloenvcha.2013.08.001>
- Birk, S., Chapman, D., Carvalho, L., Spears, B. M., Andersen, H. E., Argillier, C., et al. (2020). Impacts of multiple stressors on freshwater biota across spatial scales and ecosystems. *Nature Ecology and Evolution*, 4(8), 1060-1068.
- Birnie-Gauvin, K., Lynch, A. J., Franklin, P. A., Reid, A. J., Landsman, S. J., Tickner, D., et al. (2023). The RACE for freshwater biodiversity: Essential actions to create the social context for meaningful conservation. *Conservation Science and Practice*, 5(4), e12911.

- Britton, J. R., Lynch, A. J., Bardal, H., Bradbeer, S. J., Coetzee, J. A., Coughlan, N. E., et al. (2023). Preventing and controlling nonnative species invasions to bend the curve of global freshwater biodiversity loss. *Environmental Reviews*, 31(2), 310-326.
- Brumm, K. J., Hanks, R. D., Baldwin, R. F., & Peoples, B. K. (2022). A scale-linked conservation planning framework for freshwater ecosystems. *Landscape Ecology*, 37(10), 2589-2605.
- Bunn, S. E. (2016). Grand challenge for the future of freshwater ecosystems. *Frontiers in Environmental Science*, 4, 21. Research gaps remain but we also do know what works in some scenarios and need to take action.
- Burkhead, N. M. (2012). Extinction rates in North American freshwater fishes, 1900–2010. *BioScience*, 62(9), 798-808.
- Carpenter, S. R., Stanley, E. H., & Vander Zanden, M. J. (2011). State of the world's freshwater ecosystems: physical, chemical, and biological changes. *Annual review of Environment and Resources*, 36, 75-99.
- Ceballos, G., Ehrlich, P. R., & Raven, P. H. (2020). Vertebrates on the brink as indicators of biological annihilation and the sixth mass extinction. *Proceedings of the National Academy of Sciences of the United States of America*, 117(24), 13596–13602. <https://doi.org/10.1073/pnas.1922686117>
- Cooke, S. J., Lapointe, N. W. R., Martins, E. G., Thiem, J. D., Raby, G. D., Taylor, M. K., et al. (2013). Failure to engage the public in issues related to inland fishes and fisheries: strategies for building public and political will to promote meaningful conservation. *Journal of Fish Biology*, 83(4), 997-1018.
- Cooke, S.J., I. Harrison, M.L. Thieme, S.J. Landsman, K. Birnie-Gauvin, R. Raghavan, I.F. Creed, G. Pritchard, A. Ricciardi and D.E.L. Hanna. 2023. Is it a new day for freshwater biodiversity?: Reflections on outcomes of the Kunming-Montreal Global Biodiversity Framework. *PLoS Sustainability and Transformation*. 2(5), e0000065.
- Cooke, S. J., Piczak, M., Nyboer, E. A., Michalski, F., Bennett, A., Koning, A. A., et al. (This Issue). Managing exploitation of freshwater species and aggregates to protect and restore freshwater biodiversity. *Environmental Reviews*. 00:000-000.
- Cooke, S.J., and Birnie-Gauvin, K. (2022). The conservation and restoration of freshwater ecosystems and biodiversity can be enhanced with ecopracticology. *Socio-Ecological Practice Research*, 4, 409-416.
- Darwall W, Smith K, Allen D, Seddon M, McGregor Reid G, Clausnitzer V, Kalkman V. 2008. Freshwater biodiversity – a hidden resource under threat. In *The 2008 Review of the IUCN Red List of Threatened Species*, Vié J-C, Hilton-Taylor C, Stuard SN (eds). IUCN: Gland, Switzerland.
- Darwall, W. R., & Freyhof, J. (2016). Lost fishes, who is counting? The extent of the threat to freshwater fish biodiversity. *Conservation of Freshwater Fishes*, 1-36.
- Davidson, N. C. (2014). How much wetland has the world lost? Long-term and recent trends in global wetland area. *Marine and Freshwater Research*, 65(10), 934-941.
- Desforges, J.E., Clarke, J., Harmsen, E.J., Jardine, A.M., Robichaud, J.A., Serré, S., Chakrabarty, P., Bennett, J.R., Hanna, D.E., Smol, J.P., et al. 2022. The alarming state of freshwater biodiversity in Canada. *Canadian Journal of Fisheries and Aquatic Sciences*, 79(2), pp.352-365.

- Dodds, W. K., Perkin, J. S., & Gerken, J. E. (2013). Human impact on freshwater ecosystem services: a global perspective. *Environmental Science and Technology*, 47(16), 9061-9068.
- Dodson, J. C., Dérer, P., Cafaro, P., & Götmark, F. (2020). Population growth and climate change: Addressing the overlooked threat multiplier. *Science of the Total Environment*, 748, 141346.
- Dudgeon, D. (2019). Multiple threats imperil freshwater biodiversity in the Anthropocene. *Current Biology*, 29(19), R960-R967.
- Dudgeon, D., and Strayer, D. L. (2024). Bending the curve of global freshwater biodiversity loss: what are the prospects?. *Biological Reviews*. 100, 205–226.
- Fluet-Chouinard, E., Stocker, B.D., Zhang, Z., Malhotra, A., Melton, J.R., Poulter, B., Kaplan, J.O., Goldewijk, K.K., Siebert, S., Minayeva, T. and Hugelius, G., 2023. Extensive global wetland loss over the past three centuries. *Nature*, 614(7947), pp.281-286.
- Gleick, P. H. (2018). Transitions to freshwater sustainability. *Proceedings of the National Academy of Sciences*, 115(36), 8863-8871.
- Gonçalves, D. V., & Hermoso, V. (2022). Global goals overlook freshwater conservation. *Science*, 377(6604), 380-380.
- Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., et al. (2019). Mapping the world's free-flowing rivers. *Nature*, 569(7755), 215-221.
- Harrison, I., Abell, R., Darwall, W., Thieme, M. L., Tickner, D., & Timboe, I. (2018). The freshwater biodiversity crisis. *Science*, 362(6421), 1369-1369.
- Higgins, J., Zablocki, J., Newsock, A., Krolopp, A., Tabas, P., & Salama, M. (2021). Durable freshwater protection: A framework for establishing and maintaining long-term protection for freshwater ecosystems and the values they sustain. *Sustainability*, 13(4), 1950.
- Hu, S., Niu, Z., Chen, Y., Li, L., & Zhang, H. (2017). Global wetlands: Potential distribution, wetland loss, and status. *Science of the Total Environment*, 586, 319-327.
- IPBES. 2018. The IPBES assessment report on land degradation and restoration. Montanarella L, Scholes R, Brainich A, editors. Bonn (Germany): Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services; p. 744. doi:10.5281/zenodo.3237392
- Isbell, F., Balvanera, P., Mori, A. S., He, J. S., Bullock, J. M., Regmi, G. R., et al. (2023). Expert perspectives on global biodiversity loss and its drivers and impacts on people. *Frontiers in Ecology and Environment*, 21(2), 94-103.
- Katz, D. (2011). Hydro-political hyperbole: Examining incentives for overemphasizing the risks of water wars. *Global Environmental Politics*, 11(1), 12-35.
- Kidd, L. R., Bekessy, S. A., & Garrard, G. E. (2019). Neither hope nor fear: Empirical evidence should drive biodiversity conservation strategies. *Trends in Ecology and Evolution*, 34(4), 278-282.
- Kuiper, J.J. 2023. A plea for more freshwater experts in IPBES. *Aquat. Conserv. Mar. Freshw. Ecosyst.* (December 2022): 1–9. doi:10.1002/aqc.3956.
- Lacoul, P., & Freedman, B. (2006). Environmental influences on aquatic plants in freshwater ecosystems. *Environmental Reviews*, 14(2), 89-136.

- Linke, S., Hermoso, V., & Januchowski-Hartley, S. (2019). Toward process-based conservation prioritizations for freshwater ecosystems. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 29(7), 1149-1160.
- Linke, S., Turak, E., & Nel, J. (2011). Freshwater conservation planning: the case for systematic approaches. *Freshwater Biology*, 56(1), 6-20.
- Lopes-Lima, M., Sousa, R., Geist, J., Aldridge, D. C., Araujo, R., Bergengren, J., et al. (2017). Conservation status of freshwater mussels in Europe: state of the art and future challenges. *Biological Reviews*, 92(1), 572-607.
- Lynch, A.J., Cooke, S.J., Arthington, A.H., Baigun, C., Bossenbroek, L., Dickens, C., et al. (2023). People need freshwater biodiversity. *WIREs Water* 10(3): e1633. doi:10.1002/wat2.1633.
- Lynch, A.J., Cooke, S.J., Deines, A.M., Bower, S.D., Bunnell, D.B., Cowx, I.G., Nguyen, V.M., Nohner, J., Phouthavong, K., Riley, B., Rogers, M.W., Taylor, W.W., Woelmer, W., Youn, S.-J., and Beard, T.D. 2016. The social, economic, and environmental importance of inland fish and fisheries. *Environmental Reviews*. 24(2): 1–7. doi:10.1139/er-2015-0064.
- Lynch, A. J., Hyman, A. A., Cooke, S. J., Capon, S. J., Franklin, P. A., Jähnig, S. C., et al. (This Issue). Future-proofing the Emergency Recovery Plan for freshwater biodiversity. *Environmental Reviews*. 00:000-000.
- Naiman, R. J., & Dudgeon, D. (2011). Global alteration of freshwaters: influences on human and environmental well-being. *Ecological Research*, 26, 865-873.
- Ormerod, S. J., Dobson, M., Hildrew, A. G., & Townsend, C. (2010). Multiple stressors in freshwater ecosystems. *Freshwater Biology*, 55, 1-4.
- Parmesan, C., Morecroft, M. D., Trisurat, Y., Adrian, R., Anshari, G. Z., Arneith, A., ... & Young, K. (2022). *Terrestrial and freshwater ecosystems and their services*. Cambridge University Press.
- Piczak, M. L., Perry, D., Cooke, S. J., Harrison, I., Benitez, S., Koning, A., et al. (This Issue). Protecting and restoring habitats to benefit freshwater biodiversity. *Environmental Reviews*. 00:000-000.
- Postel, S., & Carpenter, S. (1997). Freshwater ecosystem services. *Nature's services: Societal dependence on natural ecosystems*, 195.
- Reddy, S. M., Montambault, J., Masuda, Y. J., Keenan, E., Butler, W., Fisher, J. R., et al. (2017). Advancing conservation by understanding and influencing human behavior. *Conservation Letters*, 10(2), 248-256.
- Rehman OU, Ryan MJ, Efatmaneshnik M (2017) Future Proofing Process. *INCOSE Int Symp* 27:921–934. <https://doi.org/https://doi.org/10.1002/j.2334-5837.2017.00403.x>
- Sayer, C. A., Fernando, E., Jimenez, R. R., Macfarlane, N. B., Rapacciuolo, G., Böhm, M., et al. (2025). One-quarter of freshwater fauna threatened with extinction. *Nature*. <https://doi.org/10.1038/s41586-024-08375-z>
- Sherman, M., Ford, J., Llanos-Cuentas, A. et al. Vulnerability and adaptive capacity of community food systems in the Peruvian Amazon: a case study from Panaillo. *Natural Hazards* 77, 2049–2079 (2015). <https://doi.org/10.1007/s11069-015-1690-1>
- Smol, J.P. 2010. The power of the past: Using sediments to track the effects of multiple stressors on lake ecosystems. *Freshwater Biology* 55 (Suppl. 1): 43-59.

Strayer, D. L., & Dudgeon, D. (2010). Freshwater biodiversity conservation: recent progress and future challenges. *Journal of the North American Benthological Society*, 29(1), 344-358.

Strayer, D. L., Downing, J. A., Haag, W. R., King, T. L., Layzer, J. B., Newton, T. J., & Nichols, J. S. (2004). Changing perspectives on pearly mussels, North America's most imperiled animals. *BioScience*, 54(5), 429-439.

Thieme, M., Birnie-Gauvin, K., Opperman, J. J., Franklin, P. A., Richter, H., Baumgartner, L., et al. (This Issue). Measures to Safeguard and Restore River Connectivity. *Environmental Reviews*. 00:000-000.

Tickner, D., Opperman, J.J., Abell, R., Acreman, M., Arthington, A.H., Bunn, S.E., Cooke, S.J., Dalton, J., Darwall, W., et al. 2020. Bending the curve of global freshwater biodiversity loss: an emergency recovery plan. *BioScience*, 70(4), 330-342. <https://doi.org/10.1093/biosci/biaa002>

Tickner, D., Schiller, L., Cooke, S.J., Barrios Ordóñez, E., Collier, U., Dalton, J., Harrison, I., Lifeng, L., Phang, S.C., Young, W.J., and Lynch, A.J. This Issue. Addressing the policy and business drivers of global freshwater biodiversity loss. *Environmental Reviews*. 00:000-000.

Toomey, A. H. (2023). Why facts don't change minds: Insights from cognitive science for the improved communication of conservation research. *Biological Conservation*, 278, 109886.

Twardek, W. M., Nyboer, E. A., Tickner, D., O'Connor, C. M., Lapointe, N. W., Taylor, M. K., et al. (2021). Mobilizing practitioners to support the Emergency Recovery Plan for freshwater biodiversity. *Conservation Science and Practice*, 3(8), e467.

WWF (2022) Living Planet Report 2022 – Building a nature-positive society. Almond, R.E.A., Grooten, M., Juffe Bignoli, D. & Petersen, T. (Eds). WWF, Gland, Switzerland.

Zieritz, A., Bogan, A. E., Rahim, K. A. A., Sousa, R., Jainih, L., Harun, S., et al. (2018). Changes and drivers of freshwater mussel diversity and distribution in northern Borneo. *Biological Conservation*, 219, 126-137.

Implementing the Emergency Recovery Plan for Freshwater Biodiversity




Figure 1. The status quo will presumably lead to the further loss of freshwater biodiversity. By implementing the FW Emergency Recovery Plan it is possible to begin to “bend the curve” and restore FW biodiversity. We know what to do (as outlined in the various papers in this special issue) so the biggest challenge is implementation. There is dire need to focus on implementation priorities (indicated on the figure) to ensure that the FW Emergency Recovery Plan is implemented at relevant scales (regional, national, international) and in a way that engages all relevant actors.