

RESEARCH

Open Access



# Avoidant/restrictive food intake disorder (ARFID) symptoms in gender diverse adults and their relation to autistic traits, ADHD traits, and sensory sensitivities

Kai S. Thomas<sup>1\*</sup>, Jennifer Keating<sup>2</sup>, Avalon A. Ross<sup>1</sup>, Kate Cooper<sup>3,4</sup> and Catherine R. G. Jones<sup>1</sup>

## Abstract

**Background** There is emerging evidence to suggest gender diverse people are overrepresented in avoidant/restrictive food intake disorder (ARFID) samples. However, the mechanisms underlying elevated risk for ARFID in this group are currently unknown. Gender diversity and neurodivergence commonly co-occur, with elevated sensory sensitivities reported to be a shared experience common across autism, attention deficit hyperactivity disorder (ADHD), and ARFID. We aimed to examine the unique contribution of sensory sensitivities, including hypo- and hyper-sensitivity, in predicting ARFID symptoms in gender diverse adults, whilst controlling for autistic and ADHD traits.

**Methods** Gender diverse adults ( $N=182$ ; 142 assigned female at birth;  $M$  age = 28.6 years) in the UK participated in an online survey. We examined correlations between their self-reported ARFID symptoms, sensory sensitivities, autistic traits (not including sensory sensitivities), and ADHD traits whilst controlling for weight and shape concerns. We then used hierarchical multiple regression to investigate the unique contribution of sensory sensitivities to ARFID symptoms whilst controlling for the other neurodivergent traits.

**Results** In our gender diverse sample, higher levels of ARFID symptoms were associated with higher levels of sensory sensitivities, autistic traits, and ADHD traits, after controlling for weight and shape concerns. Furthermore, sensory sensitivities, specifically hyper-sensitivity, uniquely predicted levels of ARFID symptoms once we accounted for autistic and ADHD traits.

**Conclusions** When considering neurodivergence, sensory hyper-sensitivities may be particularly relevant to ARFID symptomatology in gender diverse adults. Future research should explore associations between ARFID presentations and sensory sensitivities in large samples of gender diverse adults, to enable separate analyses by gender identity.

## Plain english summary

Gender diverse people, whose gender identity does not align with their sex assigned at birth, are at increased risk for eating disorders such as avoidant/restrictive food intake disorder (ARFID). Sensory sensitivities, autism and attention deficit hyperactivity disorder (ADHD) may increase a person's risk of developing ARFID. Gender diverse

\*Correspondence:

Kai S. Thomas  
ThomasK30@cardiff.ac.uk

Full list of author information is available at the end of the article



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

people commonly report co-occurring autistic and ADHD traits, making it challenging to determine whether ARFID symptoms are linked to the autistic and ADHD traits they report, or whether ARFID symptoms are specifically associated with sensory sensitivities. We asked 182 gender diverse adults to complete an online questionnaire about their levels of ARFID symptoms, autistic traits, ADHD traits, and sensory sensitivities. We found gender diverse people with higher levels of ARFID symptoms also reported higher levels of autistic traits, ADHD traits, and sensory sensitivities. When we accounted for autistic and ADHD traits in our gender diverse sample, sensory hypersensitivity (i.e., an over-response to sensory information) had the strongest association with ARFID symptoms. Our findings suggest sensory sensitivities may be important to the development of ARFID symptoms in gender diverse people, over and above the influence of autistic and ADHD traits.

**Keywords** Avoidant/restrictive food intake disorder (ARFID), Gender diversity, Transgender, Autistic traits, ADHD traits, Sensory sensitivity

## Introduction

Gender diverse people, whose gender identity does not align with their sex assigned at birth, are at elevated risk of developing eating disorders [1–10]. Eating disorders are prevalent across the gender diverse spectrum, including in binary trans individuals, such as trans men and trans women, and individuals who identify with an identity between or outside of the gender binary, such as non-binary, gender fluid, or genderqueer [11–13]. A recent meta-analysis reported estimated prevalence rates of eating disorders to be 17.7% in binary trans individuals [14]. There is some evidence to suggest these rates may be even higher for individuals who identify with a non-binary identity [13]. These estimated rates are considerably elevated compared to rates of 1.8% and 0.2% reported in cisgender women and men, respectively [15]. Eating disorder symptomatology in gender diverse people is likely to serve a number of functions, such as attaining body characteristics that are more aligned with their gender identity [2, 4, 7], controlling pubertal development, such as menstruation cessation for individuals assigned female at birth [6, 11, 16, 17], and/or managing heightened levels of emotional distress due to interpersonal and intrapersonal stressors [18, 19]. The majority of research with gender diverse people to-date has focused on eating pathology related to weight and shape concerns [1, 11, 20, 21], with little attention given to other types of feeding and eating disorders.

Avoidant/restrictive food intake disorder (ARFID) is a feeding and eating disorder characterised by an inadequate intake of nutrition and/or energy that is associated with significant physical and/or psychological difficulties, such as nutritional deficiencies, weight loss, and a marked interference with psychosocial functioning [22]. ARFID was introduced in the Diagnostic and Statistical Manual of Mental Disorders, fifth edition (DSM-5; 22) in 2013, where it replaced Feeding Disorder of Infancy and Early Childhood, recognising that ARFID symptoms can occur at any age. ARFID is distinguished from other eating disorders by an absence of weight or shape concerns [22]. Restrictive and avoidant eating behaviour presented

in ARFID can be related to aversions to sensory properties (i.e., related to the texture, taste, smell, or appearance of the food), low appetite or limited interest in eating, and/or fear of negative consequences from eating (e.g., choking; 23). According to the three-dimensional model of ARFID proposed by Thomas et al. [23], these underpinning drivers are not mutually exclusive and can vary in severity, resulting in heterogeneous presentations of ARFID.

The true prevalence of ARFID is currently unknown and estimates have primarily been focussed on child samples. Prevalence estimates vary from 1.5 to 64% among clinical eating disorder populations and <1–15.5% in non-clinical cohorts [24]. Two of the only studies to examine ARFID prevalence in older adolescents and adults from the general population reported rates of 0.3% and 4.7%, assessing ARFID using interviews and self-report questionnaires, respectively [25, 26]. Research examining the developmental trajectories of ARFID is lacking, but there is some evidence to suggest that ARFID typically presents before adulthood and can remain stable across the lifespan [27]. Therefore, ARFID symptomatology is not confined to childhood, and further research focussed specifically on adults is necessary to increase understanding of ARFID symptoms across the lifespan [24, 28, 29].

There is emerging evidence to suggest gender diverse people are overrepresented in ARFID samples. In a retrospective chart review of adults treated for ARFID at an eating disorder clinic, MacDonald, Liebman [30] found 16.7% of their sample identified as transgender, non-binary, or another gender diverse identity. This figure far exceeds the prevalence that would be expected, given gender diverse people make up less than 2% of the general population [31, 32]. The only other studies exploring ARFID in gender diverse people have focussed on screening for ARFID using self-report measures [33–35]. All three studies used the Nine-Item ARFID Screen (NIAS; 36), a well-established screening measure for ARFID. The NIAS has three subscales that capture symptoms associated with the three main overlapping

ARFID presentations: sensory sensitivity, lack of interest, and fear of aversive consequences [36]. The measure has recently been validated in gender diverse youth and young adults attending a US gender clinic [35]. In this study, Zickgraf, Garwood [35] reported 22% of their sample screened positive for ARFID, with gender diverse people assigned female at birth scoring significantly higher on the NIAS compared to those assigned male at birth. Despite growing evidence suggesting a link between ARFID and gender diversity, the underlying mechanisms are currently unclear.

Neurodivergence may play a significant role in the co-occurrence of gender diversity and ARFID. Research has found an overlap between autism, attention deficit hyperactivity disorder (ADHD), and gender diversity [37–41]. Both autism and ADHD are around 2–6 times more common in gender diverse individuals compared to cisgender individuals, whose gender identity aligns with their sex assigned at birth [41]. In addition, both autistic and non-autistic gender diverse people endorse more autistic traits on self-report measures compared to cisgender people [41–45]. Autistic people and people with ADHD are found to be at elevated risk of developing ARFID [30, 46]. For example, in a cohort of adults attending treatment for ARFID, MacDonald, Liebman [30] found 16.7% of the sample had an autism diagnosis whilst 26.2% had an ADHD diagnosis. Previous research has not yet explored the relations between autistic traits, ADHD traits, and ARFID symptoms in a gender diverse sample.

Based on the three maintaining mechanisms typically seen in ARFID [23, 24], autistic people and people with ADHD more commonly exhibit presentations driven by food-related sensory sensitivities and/or a lack of appetite/interest in eating, compared to those driven by a fear of aversive consequences of eating [30, 46]. The heightened occurrence of these ARFID presentations may be reflective of behavioural and cognitive patterns characteristic of and/or associated with both autism and ADHD. These include sensory sensitivities, encompassing both hyper- and hypo-sensitivity, emotion regulation differences, hyperactivity and impulsivity, and behavioural and cognitive rigidity, such as stimming, monotropic interests, and difficulties coping with novelty or uncertainty [47–55]. These characteristics can lead to eating behaviours typically seen in ARFID, such as rigid rules around food preparation, a limited food repertoire, insistence on specific brands of food only, as well as increased consumption of foods with pleasant sensory properties. Taken together, these findings suggest the elevated rates of ARFID in gender diverse people may be driven by the high rates of co-occurring neurodivergence.

When considering neurodivergent traits that may be relevant to ARFID, sensory sensitivities may be particularly relevant as a transdiagnostic feature. They are

included in the diagnostic criteria for both autism and ARFID [22] and are also observed at elevated levels in people with ADHD [53], suggesting they are a transdiagnostic experience. Sensory sensitivities are also present in samples independent of neurodivergent traits, such as individuals with migraine [56], anxiety and depressive disorders [57], and the entire spectrum of eating disorders [58, 59]. Sensory hyper-sensitivities, defined as an over-response (i.e., speed, intensity, or duration of response) to sensory stimuli [60], and sensory hypo-sensitivities, defined as an under-response to sensory stimuli [61], have both been reported in autistic children and adults [62–64], as well as children and adults with ADHD [65–67].

Although limited, there is evidence to suggest both autistic and non-autistic gender diverse people experience elevated levels of sensory sensitivities compared to cisgender autistic and non-autistic people, respectively [41]. These sensory sensitivities may contribute to increased gender dysphoria and elevated distress resulting in a heightened risk of developing ARFID symptoms. Qualitative research with autistic gender diverse adults suggests sensory sensitivities may exacerbate feelings of gender dysphoria by increasing sensitivity to negative experiences of one's body, known as 'sensory dysphoria' [63]. These sensory experiences can include facial hair growth, menstruation, and wearing clothing that is particularly uncomfortable due to the fabric or shape [63]. It is possible that sensory sensitivities are a key driver of ARFID symptoms in gender diverse people, alongside co-occurring neurodivergent traits.

In the current study, we aimed to examine the unique contribution of sensory sensitivities to levels of ARFID symptoms in gender diverse adults, whilst controlling for autistic traits (not including sensory sensitivities) and ADHD traits. Given that commonly used self-report measures of ARFID, such as the NIAS, are not able to reliably differentiate between restrictive eating behaviours specific to ARFID and those related to other eating disorders [68], we also controlled for weight and shape concerns. We hypothesised that: (1) ARFID symptoms would be positively correlated with levels of autistic traits, ADHD traits, and sensory sensitivities after controlling for weight and shape concerns and (2) sensory sensitivities would uniquely contribute to levels of ARFID symptoms in our gender diverse sample.

## Methods

### Participants

A total of 281 participants consented to participate in the study. All participants were recruited through social media advertisements and relevant support organisations. Advertisements specified that the aims of the research were to understand how eating behaviours were

**Table 1** Sample characteristics ( $N = 182$ )

|  | Frequency | % of the sample |
|--|-----------|-----------------|
| <b>Sex assigned at birth</b>                 |           |                 |
| Female                                       | 142       | 78.0            |
| Male   | 31        | 17.0            |
| Prefer not to say/missing                    | 9         | 4.9             |
| <b>Gender identity<sup>a</sup></b>           |           |                 |
| Trans man                                    | 44        | 24.2            |
| Trans woman                                  | 21        | 11.5            |
| Gender expansive <sup>b</sup>                | 28        | 15.4            |
| Non-binary <sup>b</sup>                      | 64        | 35.2            |
| Trans masculine non-binary <sup>b</sup>      | 18        | 9.9             |
| Trans feminine non-binary <sup>b</sup>       | 7         | 3.8             |
| <b>Eating disorder diagnosis<sup>c</sup></b> |           |                 |
| Yes  | 28        | 15.4            |
| Anorexia Nervosa                             | 15        |                 |
| Avoidant/Restrictive Food Intake Disorder    | 3         |                 |
| Bulimia Nervosa                              | 2         |                 |
| Eating Disorder Not Otherwise Specified      | 8         |                 |
| No   | 153       | 84.1            |
| Prefer not to say                            | 1         | 0.5             |
| <b>Status of Eating Disorder</b>             |           |                 |
| Current                                      | 19        | 10.4            |
| Historical                                   | 8         | 4.4             |
| Unsure                                       | 1         | 0.5             |
| <b>Autism diagnosis<sup>c</sup></b>          |           |                 |
| Yes  | 54        | 29.7            |
| No   | 126       | 69.2            |
| Prefer not to say                            | 2         | 1.1             |
| <b>ADHD diagnosis<sup>c</sup></b>            |           |                 |
| Yes  | 47        | 25.8            |
| No   | 131       | 72.0            |
| Prefer not to say                            | 4         | 2.2             |
| <b>Mental health diagnoses<sup>c</sup></b>   |           |                 |
| Yes  | 135       | 74.2            |
| Depressive Disorders                         | 111       |                 |
| Anxiety Disorders                            | 93        |                 |
| Post-Traumatic Stress Disorder               | 27        |                 |
| Obsessive-Compulsive Disorder                | 16        |                 |
| Borderline Personality Disorder              | 11        |                 |
| No   | 46        | 25.3            |
| Prefer not to say                            | 1         | 0.5             |

Note.<sup>a</sup>To verify that participants identified as gender diverse, they were asked to indicate whether their gender identity aligned with their sex assigned at birth. They were then asked to specify their gender identity using an open-text box. These free-text responses were coded by the authors into one of the six categories displayed

<sup>b</sup>Sex assigned at birth for participants in each of these categories: Gender expansive = 2 people assigned male at birth (AMAB), 23 people assigned female at birth (AFAB); Non-binary = 7 people AMAB, 55 people AFAB; Transmasculine non-binary = 1 person AMAB, 17 people AFAB; Transfeminine non-binary = 5 people AMAB, 1 person AFAB

<sup>c</sup>Participants were asked whether they had received a formal diagnosis of an eating disorder of any type, autism, ADHD, or diagnosis of a mental health condition. If participants responded 'Yes' to either an eating disorder or mental health diagnosis, they were given the option to specify one or more diagnoses using a free text response

$N = 24$  (13.2% of the sample) reported both autism and ADHD diagnoses

related to gender identity and neurodivergent characteristics. Participants were required to identify as gender diverse, be aged 18 years or older, be currently based in the UK, and be fluent in English. Eighty participants were excluded due to incomplete responses (>10% missing items on individual questionnaires), one participant was excluded due to failure of attention checks, and 18 participants were excluded as they reported identifying as cisgender or did not provide their gender identity. This resulted in a final sample of 182 participants ( $M$  age = 28.60 years,  $SD = 8.57$ , ranged from 18 to 70 years). An a priori power analysis performed using G\*Power indicated a minimum sample size of 178 was needed for a linear multiple regression with eleven predictors, assuming a medium effect size ( $f^2 = 0.15$ ), alpha of 0.05, and power of 0.95 [69].

The project received ethical approval from the University's School of Psychology Ethics Committee (EC.23.10.10.6846GA3). Informed consent was obtained online from all participants before completing the questionnaires. Demographic information and diagnostic history are presented in Table 1, with additional demographics (ethnicity, highest level of education and sexual orientation) included in the Supplementary Materials (Table S1).

### Procedure

The study was hosted on an online platform [70]. Participants first answered a screening question to allow us to identify invalid low-quality responses [71]. In this screening question, participants were required to answer an identification question involving processing, understanding, and responding. Participants were presented with a snow sculpture and asked to briefly describe what they predicted would happen to this sculpture on a warm, sunny day. We screened individual free-text responses from each participant to ensure these were consistent with a description of melting snow. Participants were then asked to provide demographic information and complete a series of self-report questionnaires examining eating disorder behaviours and cognitions, sensory sensitivities, and neurodivergent traits. An additional measure of internalising symptoms was also included but was not relevant to the current research questions. An attention check was presented twice during the series of questionnaires, at the start and half-way through completion. This read: "Please ignore the question below and select both 'yes' and 'unsure'. This way, we can be more confident that you are reading the questions carefully and will pay attention throughout the study. Do you frequently get less than 7 hours of sleep?" Response options were 'Yes', 'No', or 'Unsure'. After completing the questionnaires, participants were offered the option to enter a prize draw to win a monetary voucher as compensation for their time.



Participants' contact details were collected via a separate link to their data.

## Measures

### **Nine-item ARFID screen (NIAS; 36)**

The NIAS is a nine-item self-report measure of ARFID symptoms. It is comprised of three subscales: the picky/selective eating subscale measuring sensory aversion to food (e.g., "I dislike most foods that other people eat"), the appetite subscale measures a lack of interest in food or eating (e.g., "I am not very interested in eating; I seem to have a smaller appetite than other people"), and the fear subscale (e.g., "I eat small portions because I am afraid of GI discomfort, choking, or vomiting"). Participants respond on a 6-point scale from 0 (Strongly disagree) to 5 (Strongly agree). The three subscales (ranging from 0 to 15) can be summed to create a total score (ranging from 0 to 45). Higher scores indicate higher levels of ARFID symptoms. The NIAS has recently been validated in gender diverse youth and young adults attending a US gender clinic [35]. In the current study, we only used the NIAS total score in our analyses (Cronbach's  $\alpha = 0.866$ ).

### **Autism spectrum quotient short (AQ-S)**

The AQ-S [72] is an abridged version of the AQ [73], a 50-item self-report measure of autistic traits in adults. The AQ-S contains a subset of 28 items from the AQ which form five subscales: social skills, routine, switching, imagination, and numbers/patterns. Participants respond on a 4-point scale from 1 (Definitely agree) to 4 (Definitely disagree) according to their endorsement of each statement. The scoring is then reversed for items where an 'agree' response is characteristic of autism (e.g., 'I frequently get strongly absorbed in one thing'). Higher scores indicate higher levels of autistic traits. Items are summed to calculate scores, including the total score which ranges from 28 to 112. Subscales ranged from 8 to 32 (Social skills and Imagination), 4–16 (Routine and Switching), or 5–20 (Numbers/patterns), depending on their number of items. Hoekstra, Vinkhuyzen [72] recommended including one item ('New situations make me anxious') in the scoring of both the social skills and routine subscales for English samples. As our sample were all based in the UK and fluent in English, we followed this recommendation. Cronbach's  $\alpha$  for the AQ-S were as follows: total score = 0.866; social skills subscale score = 0.795; routine subscale score = 0.645; switching subscale score = 0.563; imagination subscale score = 0.786; numbers and patterns subscale score = 0.784. We used all five subscales in our analyses.

### **Adult ADHD self-report scale (ASRS)**

The ASRS [74] is an 18-item self-report measure of ADHD traits. For each item (e.g., how often do you have

problems remembering appointments or obligations?), the participant is asked to respond on a 5-point Likert scale ('Never' to 'Very often'), based on how they have felt/behaved over the past 6 months. Depending on the item, responses are either scored as 0 or 1. The total score is calculated through summing the items and ranges from 0 to 18. Higher scores indicate higher levels of current ADHD traits. Three subscales can be calculated from the items based on the factors identified by Stanton, Forbes [75]: inattentive (ranges from 0 to 9), hyperactive/impulsive motor (0–5) and hyperactive/impulsive verbal (0–4). Cronbach's  $\alpha$  for the ASRS was as follows: total score = 0.848, inattentive subscale score = 0.754, hyperactive/impulsive motor subscale score = 0.698, hyperactive/impulsive verbal subscale score = 0.695. We used all three subscales in our analyses.

### **Glasgow sensory questionnaire (GSQ)**

The GSQ [76] is a self-report measure of sensory sensitivities across seven sensory domains (visual; auditory; gustatory; olfactory; tactile; vestibular and proprioceptive). The 42 items are evenly split to measure hyper- and hypo-sensitivity. The measure was initially constructed based on sensory sensitivities associated with autism, but has been found to be reliable in both autistic and non-autistic adults [77]. Participants respond based on the frequency of each type of sensory experience (e.g., Do you react very strongly when you hear an unexpected sound?) on a 5-point scale from 0 (Never) to 4 (Always). Given sensory sensitivities are a key component of ARFID, we removed overlapping eating-related items (items 2, 27, 23, 26, 28, and 40) from the GSQ before scoring. Hypo- and hyper-sensitivity subscales are calculated by summing corresponding items (using the adapted scoring method, scores ranged from 0 to 72), whilst total scores are calculated by summing all items (0–144). Higher scores indicate higher levels of sensory sensitivities. Cronbach's  $\alpha$  was calculated for total score ( $\alpha = 0.923$ ) and both hyper-sensitivity ( $\alpha = 0.888$ ) and hypo-sensitivity ( $\alpha = 0.841$ ) subscales. We used both hyper- and hypo-sensitivity subscales in our analyses.

### **Eating disorder examination questionnaire (EDE-Q)**

The EDE-Q [78] assesses eating disorder behaviours and cognitions over the previous 28 days (e.g., on how many of the past 28 days have you had a definite fear that you might gain weight? ), with responses captured on a 7-point scale ranging from 0 ('no days') to 6 ('every day'). Averaging the scores of relevant items is used to capture four subscales: restraint, eating concern, shape concern, and weight concern. These subscale scores are averaged to create a global EDE-Q score (ranging from 0 to 6), with higher scores indicating a greater endorsement of eating disorder symptoms. Only the shape concern and weight

concern subscales were used in the current analyses. The EDE-Q is a well-established measure that has been validated in gender diverse people [12, 79, 80]. Cronbach's  $\alpha$  for the shape concern ( $\alpha=0.896$ ) and weight concern ( $\alpha=0.838$ ) subscales in the current study were acceptable.

### Statistical analyses

All statistical analyses were conducted using R Statistical Software (version 4.3.2; R Core Team, 2021), accessed through R Studio (version 23.09.1). Missing values were low across all measures (<0.35% of all datapoints) and mean imputation was used to replace these. The mean, standard deviation, and range were calculated for each subscale of the NIAS, AQ-S, ASRS, and GSQ, in addition to the shape concern and weight concern subscales of the EDE-Q. To examine our first hypothesis that ARFID symptoms would be positively correlated with levels of autistic traits, ADHD traits, and sensory sensitivities, Spearman's correlations were conducted between ARFID symptoms (NIAS total score) and the subscales of the AQ-S (autistic traits), ASRS (ADHD traits), and GSQ (sensory sensitivities). These associations were further examined using partial correlations to control for weight and shape concerns (EDE-Q). To address our second hypothesis that sensory sensitivities would uniquely contribute to levels of ARFID symptoms, multiple linear regressions were conducted to examine the contributions

of each independent variable (autistic traits, ADHD traits, and sensory sensitivities) on ARFID symptoms. In the multiple linear regression model, our control variables, sex at birth and EDE-Q Shape and Weight Concern subscales, were entered at Step 1, followed by GSQ hyper- and hypo-sensitivity subscales at Step 2, and AS-Q and ASRS subscales at Step 3. For the linear regression model, there was independence of residuals, no evidence of multicollinearity, homoscedasticity was present, and the assumption of normality was met, as assessed by a P-P Plot. The regression model was repeated with individuals with and without an eating disorder diagnosis (current and/or historical). Lastly, we conducted descriptive statistics and exploratory correlation analyses for each gender diverse group (Supplementary Materials).

### Results

The mean, standard deviation, and range for each subscale of the NIAS, AQ-S, ASRS, and GSQ, in addition to the shape concern and weight concern subscales of the EDE-Q for the whole sample are presented in Table 2. Descriptive statistics for each gender identity category (trans masculine, trans feminine, non-binary, gender expansive) are presented in Supplementary Materials.

Our sample reported a moderate level of ARFID symptoms, with mean scores on each NIAS subscale below the proposed screening cutoffs of  $\geq 10$  (picky eating),  $\geq 9$  (appetite), and  $\geq 10$  (fear) by Burton Murray, Dreier [81]. Picky eating was the most endorsed presentation of ARFID across the sample, with 53 participants (29.1%) scoring above the NIAS-picky eating subscale cutoff. This was closely followed by the 49 participants (26.9%) who scored above the NIAS-appetite cutoff, while only 23 participants (12.6%) met the cutoff score for the NIAS-fear subscale. Autistic traits were elevated in our sample, with 88.5% of the overall sample exceeding the screening cutoff score ( $> 65$ ) suggested by Hoekstra, Vinkhuyzen [72]. Eating concern and weight concern mean scores, measured using the EDE-Q, were slightly higher than previously reported in both binary transgender [80] and gender-expansive populations [12].

In relation to the first hypothesis, correlational analyses revealed significant positive associations between ARFID symptoms (NIAS total score) and autistic traits (AQ-S), ADHD traits (ASRS), and sensory sensitivities (GSQ) (Table 3). The only associations that were not significant were between the switching subscale (AQ-S) and ARFID symptoms (NIAS total score) and the verbal subscale (ASRS) and ARFID symptoms (NIAS total score). Furthermore, there were significant positive correlations between all autistic and ADHD traits and both hyper- and hypo-sensory sensitivities.

Partial correlations were conducted with the variables above, while controlling for shape concern and weight

**Table 2** Descriptive statistics for the questionnaires ( $n=182$ )

|                                     | Min-Max | Available scores | Mean (SD)     |
|-------------------------------------|---------|------------------|---------------|
| <b>ARFID symptoms (NIAS)</b>        | 0–44    | 0–45             | 15.91 (10.49) |
| Picky eating                        | 0–15    | 0–15             | 6.71 (4.85)   |
| Appetite                            | 0–15    | 0–15             | 5.69 (4.44)   |
| Fear                                | 0–15    | 0–15             | 3.50 (4.33)   |
| <b>Autistic traits (AQ-S)</b>       | 38–107  | 28–112           | 80.02 (12.25) |
| Social skills                       | 8–32    | 8–32             | 23.63 (4.53)  |
| Routine                             | 4–16    | 4–16             | 12.99 (2.26)  |
| Switching                           | 6–16    | 4–16             | 13.00 (2.25)  |
| Imagination                         | 8–32    | 8–32             | 19.81 (5.26)  |
| Numbers and patterns                | 5–20    | 5–20             | 14.28 (3.54)  |
| <b>ADHD traits (ASRS)</b>           | 1–18    | 0–18             | 12.36 (4.01)  |
| Inattentive                         | 0–8     | 0–9              | 6.04 (1.98)   |
| Hyperactivity/impulsivity motor     | 0–5     | 0–5              | 3.41 (1.48)   |
| Hyperactivity/impulsivity verbal    | 0–4     | 0–4              | 2.21 (1.35)   |
| <b>Sensory sensitivities (GSQ)*</b> | 12–115  | 0–144            | 69.28 (21.61) |
| Hyper-sensitivity                   | 6–66    | 0–72             | 37.95 (12.50) |
| Hypo-sensitivity                    | 5–55    | 0–72             | 31.33 (10.55) |
| <b>Shape concern (EDE-Q)</b>        | 0–6     | 0–6              | 3.54 (1.56)   |
| <b>Weight concern (EDE-Q)</b>       | 0–6     | 0–6              | 3.12 (1.61)   |

Note. NIAS: Nine-Item ARFID Scale, ARFID: Avoidant/Restrictive Food Intake Disorder; AQ-S: Autism Spectrum Quotient - Short, ADHD: Attention-deficit Hyperactivity Disorder, ASRS: Adult ADHD Self-Report Scale, GSQ: Glasgow Sensory Questionnaire, EDE-Q: Eating Disorder Examination Questionnaire

\*Adapted GSQ scales were used (removing any eating-related items)

**Table 3** Spearman’s correlations between neurodivergent trait subscales, sensory sensitivities, shape and weight concerns, and NIAS total score

|                                    | NIAS Total Score | GSQ Hyper-sensitivity | GSQ Hypo-sensitivity |
|------------------------------------|------------------|-----------------------|----------------------|
| <b>Autistic traits (AQ-S)</b>      | 0.405***         | 0.629***              | 0.399***             |
| Social skills                      | 0.282***         | 0.456***              | 0.331***             |
| Routine                            | 0.271***         | 0.406***              | 0.284***             |
| Switching                          | 0.129            | 0.324***              | 0.244***             |
| Imagination                        | 0.306***         | 0.362***              | 0.284***             |
| Numbers and patterns               | 0.323***         | 0.538***              | 0.549***             |
| <b>ADHD traits (ASRS)</b>          | 0.186*           | 0.395***              | 0.605***             |
| Inattentive                        | 0.163*           | 0.301***              | 0.420***             |
| Hyperactivity/impulsivity motor    | 0.253***         | 0.430***              | 0.568***             |
| Hyperactivity/impulsivity verbal   | 0.096            | 0.250***              | 0.383***             |
| <b>Sensory sensitivities (GSQ)</b> |                  |                       |                      |
| Hyper-sensitivity                  | 0.557***         |                       |                      |
| Hypo-sensitivity                   | 0.403***         |                       |                      |
| <b>Shape concern (EDE-Q)</b>       | 0.100            |                       |                      |
| <b>Weight concern (EDE-Q)</b>      | 0.122            |                       |                      |

\* $p < .05$ , \*\* $p < .01$ , \*\*\* $p < .001$ .

Note: NIAS: Nine-Item ARFID Scale; ARFID: Avoidant/Restrictive Food Intake Disorder; AQ-S: Autism Spectrum Quotient– Short; ASRS: Adult ADHD Self-Report Scale; GSQ: Glasgow Sensory Questionnaire; EDE-Q: Eating Disorder Examination Questionnaire

**Table 4** Partial correlations between neurodivergent trait subscales, sensory sensitivities and ARFID symptoms (NIAS) while controlling for EDE-Q weight and shape concern subscales

|                                    | NIAS Total Score |
|------------------------------------|------------------|
| <b>Autistic traits (AQ-S)</b>      | 0.389***         |
| Social skills                      | 0.280***         |
| Routine                            | 0.284***         |
| Switching                          | 0.160*           |
| Imagination                        | 0.288***         |
| Numbers and patterns               | 0.304***         |
| <b>ADHD traits (ASRS)</b>          | 0.179*           |
| Inattentive                        | 0.169*           |
| Hyperactivity/impulsivity motor    | 0.259***         |
| Hyperactivity/impulsivity verbal   | 0.044            |
| <b>Sensory sensitivities (GSQ)</b> |                  |
| Hyper-sensitivity                  | 0.558***         |
| Hypo-sensitivity                   | 0.417***         |

\* $p < .05$ , \*\* $p < .01$ , \*\*\* $p < .001$

Note: NIAS: Nine-Item ARFID Scale; ARFID: Avoidant/Restrictive Food Intake Disorder; AQ-S: Autism Spectrum Quotient– Short; ASRS: Adult ADHD Self-Report Scale; GSQ: Glasgow Sensory Questionnaire; EDE-Q: Eating Disorder Examination Questionnaire

**Table 5** Regression of sensory, attention, and autistic traits predicting ARFID scores

|                                      | NIAS Total Score |
|--------------------------------------|------------------|
| <b>Step 1</b>                        |                  |
| Sex assigned at birth                | 0.035            |
| EDE-Q Shape concerns                 | −0.151           |
| EDE-Q Weight concerns                | 0.240            |
| $R^2$                                | 0.016            |
| F                                    | 0.954            |
| <b>Step 2</b>                        |                  |
| Sex assigned at birth                | −0.060           |
| EDE-Q Shape concerns                 | −0.051           |
| EDE-Q Weight concerns                | −0.041           |
| GSQ Hyper-sensitivity                | 0.575***         |
| GSQ Hypo-sensitivity                 | 0.022            |
| $R^2$                                | 0.320            |
| F                                    | 16.450***        |
| <b>Step 3</b>                        |                  |
| Sex assigned at birth                | −0.065           |
| EDE-Q Shape concerns                 | −0.023           |
| EDE-Q Weight concerns                | −0.072           |
| GSQ Hyper-sensitivity                | 0.499***         |
| GSQ Hypo-sensitivity                 | 0.034            |
| AQ-S Social skills                   | 0.019            |
| AQ-S Routine                         | 0.053            |
| AQ-S Imagination                     | 0.061            |
| AQ-S Numbers and patterns            | 0.000            |
| ASRS Inattentive                     | −0.034           |
| ASRS Hyperactivity/impulsivity motor | 0.050            |
| $R^2$                                | 0.329            |
| F                                    | 7.521***         |

\* $p < .05$ , \*\* $p < .01$ , \*\*\* $p < .001$

Note: NIAS: Nine-Item ARFID Scale; ARFID: Avoidant/Restrictive Food Intake Disorder; AQ-S: Autism Spectrum Quotient– Short; ASRS: Adult ADHD Self-Report Scale; GSQ: Glasgow Sensory Questionnaire; EDE-Q: Eating Disorder Examination Questionnaire

concern subscales (EDE-Q) (Table 4). Similar significant correlations remained for all subscales; however, the correlation between the AQ-S switching subscale and NIAS total score was now also significant ( $p = .032$ ). Correlations conducted for each of the gender diverse groups (see Supplementary Materials) were broadly consistent with the whole sample analyses. However, participants in the gender expansive group displayed additional weak positive correlations between ARFID symptoms and both sensory hypo-sensitivities and some autistic traits.

With regard to the second hypothesis, multiple linear regression analysis investigated the unique contribution of sensory sensitivities, autistic traits, and ADHD traits on ARFID symptoms (Table 5). Sex assigned at birth and shape and weight concerns (EDE-Q) were controlled for at Step 1 and this model was not statistically significant. The addition of the hyper-sensitivity and hypo-sensitivity subscales at Step 2 was statistically significant, with an increase in model fit of 0.304. In Step 3, autistic

traits (AQ-S) and ADHD traits (ASRS) that were significant correlates of ARFID symptoms were added into the model, resulted in a small increase in model fit. In this final model, hyper-sensitivity was the only significant predictor of NIAS scores ( $\beta = 0.499, p < .001$ ). Higher hyper-sensitivity scores were associated with higher scores on the NIAS. Additional exploratory analysis confirmed that the pattern of data were the same when weight and shape concerns were not controlled for in the model.

## Discussion

The present study is the first to examine the unique contribution of sensory sensitivities in predicting ARFID symptoms in a sample of gender diverse adults. Importantly, we controlled for autistic and ADHD traits, both of which are associated with sensory processing differences [62–67], as well as overrepresented in the gender diverse population [41, 63]. In line with our first hypothesis, we found higher levels of ARFID symptoms were associated with higher levels of sensory sensitivities, autistic traits, and ADHD traits after controlling for weight and shape concerns. In line with our second hypothesis, we found sensory hyper-sensitivity to be the only unique predictor of ARFID symptoms when controlling for autistic and ADHD traits. This pattern of findings suggests sensory hyper-sensitivities may be central to the experiences of ARFID symptomatology in gender diverse adults, over and above autistic and ADHD traits.

We found ARFID symptoms were positively correlated with autistic and ADHD traits, as well as sensory hyper- and hypo-sensitivities in gender diverse adults. These findings are consistent with research from cisgender populations, demonstrating a high co-occurrence between ARFID symptomatology and sensory sensitivities, autism, and ADHD [82–91]. Our findings suggest gender diverse people with higher levels of sensory sensitivities, autistic traits, and ADHD traits are at elevated risk for developing ARFID symptoms, which is broadly in line with findings from cisgender populations [88]. However, compared to cisgender samples using the same measures, gender diverse people in our study reported elevated levels of sensory sensitivities [76], ARFID symptoms [68], and neurodivergent traits [92, 93]. The level of ARFID symptoms in our sample was similar to previously published values for gender diverse youth and young adults [35] and autistic traits were closely aligned with previously published AQ-S subscale scores for autistic gender diverse adults [94]. In a recent study measuring sensory sensitivities in gender diverse adults, Warrier, Greenberg [41] found higher levels of sensory hypersensitivity in gender diverse individuals compared to cisgender individuals when using the Sensory Perception Quotient (SPQ; 95). Therefore, our study findings provide support for a body of literature demonstrating ARFID symptoms,

sensory sensitivities and neurodivergent traits are elevated in gender diverse adults.

It is also important to highlight that weight and shape concern, which were controlled for in our analyses, were elevated in our gender diverse sample compared to previous literature with largely cisgender samples in the UK [93, 96]. These findings suggest gender diverse adults are likely to report a range of eating pathologies, but the individual's specific pattern of neurodivergent or related characteristics may contribute to the type of eating pathology they experience. For example, ARFID symptoms may be particularly prevalent in individuals with higher levels of sensory sensitivities, compared to characteristics such as intense interests and preoccupations with food and weight, which are more likely to present with anorexia nervosa symptoms. Further, the trajectories of these symptoms and co-occurring factors may change across time. For example, sensory sensitivities may drive the initial development of restrictive eating behaviours, but symptoms may be later maintained by neurodivergent traits like cognitive rigidity, insistence on sameness, and intense interests [86]. Future research examining associations between ARFID, sensory sensitivities, and neurodivergent traits should also consider differences between individuals who develop ARFID in childhood compared to those with adult-onset ARFID. Research suggests individuals with adult-onset ARFID most commonly endorse the driver 'fear of aversive consequences' [27], which differs from the mixed presentations of ARFID reported in children [97].

In our correlational analysis, we explored the associations between ARFID symptoms and categories of autistic and ADHD traits. Both before and after controlling for weight and shape concerns, we found ARFID symptoms were positively associated with all autistic and ADHD characteristics captured in the measures except for verbal hyperactivity/impulsivity. Switching, captured in the measure of autistic traits, was not significantly associated with ARFID symptoms before controlling for weight and shape concerns. Our findings are consistent with research demonstrating individuals with ARFID display repetitive and routinised behaviours around food and eating [24, 86, 98], as well as differences in social responsiveness [90]. Indeed, the diagnostic criteria for ARFID can include difficulties in social functioning, such as inability to sustain social relationships and participate in social activities, as a consequence of these avoidant and restrictive eating behaviours [22]. Previous research has also reported associations between ARFID symptoms and rates of ADHD diagnoses/traits in children and adolescents [99, 100]. However, our study is the first to demonstrate this association in adults. Specifically, we found inattentiveness and hyperactivity/impulsivity to be positively associated with ARFID symptoms, although this



association was no longer significant when controlling for sensory sensitivities. This may suggest that sensory sensitivities are driving the association between neurodivergent traits and ARFID symptoms in our sample of gender diverse adults.

The results of our hierarchical regression analyses suggest there is unique variance associated with sensory sensitivities that is important above and beyond any contributions of autistic or ADHD traits to ARFID symptoms. This is consistent with previous research demonstrating that both autistic and non-autistic gender diverse people experience elevated levels of sensory sensitivities compared to cisgender autistic and non-autistic people, respectively [41]. Although sensory sensitivities may be heightened in gender diverse adults as a result of co-occurring autistic and ADHD traits, this transdiagnostic experience may also uniquely contribute to gender dysphoria. Sensory sensitivities may exacerbate feelings of gender dysphoria by increasing sensitivity to negative sensory experiences of one's body, such as facial hair growth and menstruation [63, 101]. This heightened distress may in turn facilitate the development and/or maintenance of ARFID symptoms, as a way of managing the negative sensory experiences of one's body. Given we controlled for shape and weight concerns in our analysis, it is important to consider the potential for multiple pathways from the experience of heightened distress, such as increased drive for muscularity/thinness, as well as to avoid unpleasant sensory experiences. In conclusion, sensory sensitivities may increase the risk for ARFID symptoms in gender diverse adults, over and above the contribution of co-occurring neurodivergent traits.

### Strengths and limitations

By using a dimensional approach, we were able to study associations between ARFID symptoms and sensory sensitivities, autistic, and ADHD traits. This allowed us to account for different types of neurodivergent traits in our model, which is important given the well-established links between sensory sensitivities, autism, and ADHD [76, 89, 102, 103]. In addition, we accounted for weight and shape concerns in our analyses to ensure we were examining restrictive eating behaviours specific to ARFID, rather than those associated with other eating disorders, such as anorexia nervosa [68]. However, it is important to acknowledge the potential limitations of our measures, specifically the AQ-S [72]. The measure includes a social skills subscale which frames impairments in social skills to be characteristic of autism. However, this assumption has been criticised by the autistic community, and many have proposed the double empathy problem as an alternative perspective [104–107]. This framework challenges the deficit model that assumes autistic people are exclusively contributing to difficult

or unusual cross-neurotype social interactions. Instead, it emphasises the bidirectionality of social communication and expectations, with both autistic and non-autistic social partners contributing to the outcome. Indeed, research has found autistic people communicate with other autistic people just as effectively as non-autistic people communicate with other non-autistic people [108]. Therefore, it is important to consider the use of current self-report measures of autistic traits and explore alternative tools or adaptations that are informed by the voices of autistic people, encompassing the range of the autistic experience and using accessible and strength-based language.

It is also important to highlight the potential limitations of the GSQ. The GSQ captures sensory processing across seven sensory domains (with six items representing each domain), including items on exteroceptive processing, such as visual and auditory processing, as well as items on proprioception and vestibular processing, which are thought to play a role in interoceptive processing [109]. As sensory sensitivity represents a wide range of distinct sensory domains across both exteroceptive and interoceptive processing, it can be difficult for researchers to precisely classify the type of sensory processing they are investigating, particularly in the context of challenges around classifying sensory processing as purely exteroceptive or interoceptive [110]. The GSQ does not explore distinct subtypes of the sensory experience that might be relevant for research on eating pathology, such as misophonia [111, 112]. Therefore, although the GSQ enables us to capture a number of sensory domains, it does not capture all sensory experiences, and a domain-specific measure would provide greater specificity and internal reliability. In turn, this is likely to result in more nuanced understanding of the role of each sensory processing domain in relation to eating pathology. Lastly, it is important to note that the psychometric properties of the measure may have been impacted by our decision to remove eating-related items due to the overlap between sensory sensitivities and ARFID symptoms.

A limitation of our study is that we did not explore associations between ARFID symptoms and different sensory modalities in our analyses. We chose to focus on the contribution of sensory hyper- and hypo-sensitivities only, as our sample size was moderate, and we did not have the power to explore all sensory domains. It may be the case that certain domains of sensory processing are more relevant to ARFID presentations, as well as co-occurring neurodivergent traits, than others. For example, oral texture sensitivity is the most common sensory difference reported by autistic people [82] and has been found to be an independent predictor of ARFID symptoms in children and adolescents [86]. Hyper-sensitivity to visual stimuli is associated with higher levels of

ED symptoms and autistic eating behaviours in autistic adults, while hypo-sensitivity to taste is linked to higher levels of ED symptoms only [113]. Furthermore, we did not explore associations between sensory sensitivities, autistic and ADHD traits, and types of ARFID presentations i.e., aversions to sensory properties, low appetite or limited interest in eating, and/or fear of negative consequences from eating [23]. This is important as drivers of ARFID symptoms may differ depending on the individuals' neurodivergent and sensory profile, and has implications for clinical support [114]. Further research is needed to explore these relations with larger samples to enable a more fine-grained and nuanced approach. Lastly, our cross-sectional design does not allow us to establish causality or directionality with respect to the associations between ARFID symptoms, sensory sensitivities, and neurodivergent traits.

A strength of our study is the recruitment of a moderate sample of gender diverse people who identified with a range of diverse gender identities. For example, previous research has reported challenges recruiting non-binary participants [35], whereas just under 50% of our sample identified as non-binary (including trans masculine and trans feminine non-binary identities). However, the small sample sizes for each gender diverse identity in our study limited our ability to conduct regression analyses for each group separately. This is an important limitation given previous research has found ED prevalence rates and presentations to differ across gender diverse groups [11, 13]. For example, transmasculine people assigned female at birth often report engaging in weight control behaviours and excessive exercise to achieve a more lean and muscular body and appear less feminine [115–117]. For transfeminine people assigned male at birth, engagement in weight control behaviours may be used to appear more traditionally feminine, such as altering body shape to align with feminine body ideals and reducing facial and body hair growth [7, 118–120]. Furthermore, studies exploring eating pathology in non-binary people have found thinness to be driven in part by a desire for androgynous gender expression [121–123]. Although these findings are specific to eating pathology related to weight and shape concerns rather than ARFID, further support for exploring gender diverse groups separately is reported in autism research. For example, higher levels of autistic traits have been found in non-binary and gender expansive individuals compared to binary transgender adults [42]. Therefore, neurodivergent traits may interact with gender identity to influence ARFID symptom development and presentation.

Although we controlled for sex assigned at birth in our regression analyses, gender diverse people assigned female at birth were overrepresented in our sample, even when considering recent gender clinic referral data

[124]. This appears to be a challenge shared by other ARFID studies with gender diverse people, as Zickgraf, Garwood [35] report just over 18% of their sample were assigned male at birth. In their study, gender diverse people assigned female at birth were reported to score significantly higher on the NIAS compared to those assigned male at birth [35]. However, given the underrepresentation of gender diverse people assigned male at birth, future research is needed to examine the intersection of sex assigned at birth, gender identity, and ARFID symptoms. Further, the current sample is predominantly White and educated, highlighting the need to consider a range of recruitment strategies to reach communities who may be less familiar with or comfortable engaging in research.

It is important to acknowledge the high levels of neurodivergence and mental health diagnoses reported by our sample. Although previous research has reported gender diverse people to be more likely to be autistic and/or have ADHD, just under 30% of our sample reported receiving an autism diagnosis, whilst 25% reported an ADHD diagnosis. These figures are elevated compared to rates reported by other studies with gender diverse samples and may have influenced the relations between autistic traits, ADHD traits, sensory sensitivity, and ARFID symptoms in our study. In a recent meta-analysis, Kallitsounaki and Williams [45] found autism diagnoses to have a pooled estimate of 11% in gender diverse people. A similar rate of 10% has been reported for ADHD diagnoses in gender diverse people [125]. However, the authors acknowledge large variations in prevalence rates across studies, potentially due to differences in study designs (e.g., clinical studies vs. population studies) and participant type (e.g., individuals who have received a gender dysphoria/gender incongruence diagnosis vs. individuals who report gender incongruence). Prevalence estimates of autism diagnoses are lower in studies recruiting individuals from gender clinics who have received a diagnosis of gender dysphoria compared to individuals from the general population who identify as gender diverse [45]. Although this could suggest overestimation of self-reported neurodivergent diagnoses in individuals recruited from the community, it is also possible that neurodivergent diagnoses are under-diagnosed in gender diverse people attending gender clinics [45], or may not be explored or identified until later on in their gender transition. In addition, our recruitment materials explicitly stated that we were investigating neurodivergent characteristics alongside eating behaviours and gender diversity. This may have attracted more people with neurodivergent diagnoses or self-identified neurodivergent characteristics than would typically be expected in a gender diverse sample. In addition to high rates of neurodivergent diagnoses in our sample, around 75% of our

sample reported a co-occurring mental health diagnosis, excluding eating disorders. Although previous studies with gender diverse people also report high levels of mental health diagnoses [125–127], this may be a potential confounding factor due to the high co-occurrence between eating disorders and mental health difficulties, including anxiety and depression [15, 128–130].

### Clinical implications

The findings from our study have implications for clinicians' understanding of ARFID presentations in gender diverse people, as well as considerations for treatment in this group. Firstly, clinicians supporting gender diverse people should be aware of the strong positive association between sensory sensitivities and ARFID symptoms. This may indicate a transdiagnostic feature that is elevated in neurodivergent individuals, and so clinicians should carefully assess this when working with individuals with ARFID, using standardised questionnaires where appropriate. Clinicians should aim to establish a shared understanding of sensory needs with the gender diverse individual they are supporting, as well as how these needs may intersect with eating difficulties. Gender diverse people may experience a complex intersection of gender-related distress, sensory sensitivities, and eating behaviours. The individual's experience of neurodivergence, whether this relates to traits, identity or a formal diagnosis, are likely to factor into their experience of sensory sensitivities and ARFID and should also be carefully considered. It is important that clinicians aim to disentangle and understand these experiences to reach an accurate shared formulation of eating problems.

In conclusion, our findings demonstrate higher levels of ARFID symptoms are associated with higher levels of autistic traits, ADHD traits, and sensory sensitivities in a gender diverse sample. Furthermore, sensory sensitivities, specifically hyper-sensitivity, uniquely predicted levels of ARFID symptoms once we accounted for autistic and ADHD traits. This pattern of findings suggests sensory hyper-sensitivities may be particularly relevant to ARFID symptomatology in gender diverse adults, compared to other traits that are relevant to neurodivergence. Future research is needed with larger and more diverse samples to corroborate and extend these findings.

### Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s40337-025-01215-z>.

Supplementary Material 1

### Acknowledgements

We would like to thank the participants who contributed their time to the study, as well as Grace Williams, Grace Clark, Lowri Evans, Klaudia Zwawiak, and Holly Young for their help with data collection. For the purpose of Open

Access, the author has applied a CC BY public copyright license to any Author Accepted Manuscript (AAM) version arising from this submission.

### Author contributions

All authors were involved in the conceptualisation and design of the study. JK, KT, and AR conducted the formal analysis and data curation. KT led the project administration and writing of the original draft, which was reviewed and edited by all co-authors. CJ and KC provided mentorship to the first author.

### Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

### Data availability

Data are available at <https://osf.io/s4eua/>.

### Declarations

#### Ethics approval and consent to participate

The project received ethical approval from Cardiff University's School of Psychology Ethics Committee (EC.23.10.10.6846GA3). Informed consent was obtained online from all participants before completing the questionnaires.

#### Consent for publication

Not applicable.

#### Competing interests

The authors declare no competing interests.

#### Author details

<sup>1</sup>School of Psychology, Cardiff University, Cardiff, Wales, UK

<sup>2</sup>Wales Institute of Social and Economic Research and Data, School of Social Sciences, Cardiff University, Cardiff, Wales, UK

<sup>3</sup>Department of Clinical, Educational and Health Psychology, University College London, London, England, UK

<sup>4</sup>Centre for Applied Autism Research, University of Bath, Bath, England, UK

Received: 19 December 2024 / Accepted: 6 February 2025

Published online: 17 February 2025

### References

- Diemer EW, Grant JD, Munn-Chernoff MA, Patterson DA, Duncan AE. Gender identity, sexual orientation, and Eating-Related Pathology in a National Sample of College Students. *J Adolesc Health*. 2015;57(2):144–9.
- Jones BA, Haycraft E, Murjan S, Arcelus J. Body dissatisfaction and disordered eating in trans people: a systematic review of the literature. *Int Rev Psychiatry*. 2016;28(1):81–94.
- Nagata JM, McGuiire FH, Lavender JM, Brown TA, Murray SB, Greene RE, et al. Appearance and performance-enhancing drugs and supplements, eating disorders, and muscle dysmorphia among gender minority people. *Int J Eat Disord*. 2022;55(5):678–87.
- Ålgars M, Alanko K, Santtila P, Sandnabba NK. Disordered eating and gender identity disorder: a qualitative study. *Eat Disord*. 2012;20(4):300–11.
- Brewer G, Hanson L, Caswell N. Body image and eating behavior in transgender men and women: the importance of stage of gender affirmation. *Bull Appl Transgender Stud*. 2022;1(1):71–95.
- Coelho JS, Suen J, Clark BA, Marshall SK, Geller J, Lam P-Y. Eating disorder diagnoses and Symptom Presentation in Transgender Youth: a scoping review. *Curr Psychiatry Rep*. 2019;21(11):107.
- Gordon AR, Austin S, Krieger N, White Hughto JM, Reisner SL. I have to constantly prove to myself, to people, that I fit the bill: perspectives on weight and shape control behaviors among low-income, ethnically diverse young transgender women. *Soc Sci Med*. 2016;165:141–9.
- Joy P, White M, Jones S. Exploring the influence of gender dysphoria in eating disorders among gender diverse individuals. *Nutr Dietetics*. 2022;79(3):390–9.
- Sedgewick F, Leppanen J, Tchanturia K. Gender differences in mental health prevalence in autism. *Adv Autism*. 2021;7(3):208–24.
- Strauss P, Cook A, Watson V, Winter S, Whitehouse A, Albrecht N, et al. Mental health difficulties among trans and gender diverse young people with an

- autism spectrum disorder (ASD): findings from Trans pathways. *J Psychiatr Res.* 2021;137:360–7.
11. Diemer EW, White Hughto JM, Gordon AR, Guss C, Austin SB, Reisner SL. Beyond the Binary: differences in eating disorder prevalence by gender identity in a Transgender Sample. *Transgender Health.* 2018;3(1):17–23.
  12. Nagata JM, Compte EJ, Cattle CJ, Flentje A, Capriotti MR, Lubensky ME, et al. Community norms for the eating disorder examination Questionnaire (EDE-Q) among gender-expansive populations. *J Eat Disorders.* 2020;8(1):74.
  13. Romano KA, Lipson SK. Weight misperception and thin-ideal overvaluation relative to the positive functioning and eating disorder pathology of transgender and nonbinary young adults. *Psychol Sex Orientat Gend Divers.* 2022;9(4):446–53.
  14. Rasmussen SM, Dalgaard MK, Roloff M, Pinholt M, Skrubbeltrang C, Clausen L, et al. Eating disorder symptomatology among transgender individuals: a systematic review and meta-analysis. *J Eat Disorders.* 2023;11(1):84.
  15. Duffy ME, Henkel KE, Joiner TE. Prevalence of self-injurious thoughts and behaviors in Transgender individuals with eating disorders: a National Study. *J Adolesc Health.* 2019;64(4):461–6.
  16. Kamody RC, Yonkers K, Pluhar EI, Olezeski CL. Disordered eating among Trans-Masculine Youth: considerations through a Developmental Lens. *LGBT Health.* 2020;7(4):170–3.
  17. Romito M, Salk RH, Roberts SR, Thoma BC, Levine MD, Choukas-Bradley S. Exploring transgender adolescents' body image concerns and disordered eating: semi-structured interviews with nine gender minority youth. *Body Image.* 2021;37:50–62.
  18. Hendricks ML, Testa RJ. A conceptual framework for clinical work with transgender and gender nonconforming clients: an adaptation of the minority stress model. *Prof Psychology: Res Pract.* 2012;43(5):460–7.
  19. Testa RJ, Habarth J, Peta J, Balsam K, Bockting W. Development of the gender minority stress and resilience measure. *Psychol Sex Orientat Gend Divers.* 2015;2(1):65–77.
  20. Hallward L, Nagata JM, Rodgers RF, Ganson KT. Examination of eating disorder psychopathology across sexual and gender identities among a Canadian sample. *Int J Eat Disord.* 2023;56(3):604–15.
  21. Nagata JM, Stuart E, Hur JO, Panchal S, Low P, Chaphekar AV, et al. Eating disorders in sexual and gender minority adolescents. *Curr Psychiatry Rep.* 2024;26(7):340–50.
  22. American Psychiatric Association D-TR. Diagnostic and statistical manual of mental disorders: DSM-5™. 5th ed. Arlington, VA, US: American Psychiatric Publishing, Inc.; 2013. pp. 947–xlv.
  23. Thomas JJ, Lawson EA, Micali N, Misra M, Deckersbach T, Eddy KT. Avoidant/Restrictive food intake disorder: a three-Dimensional Model of Neurobiology with implications for etiology and treatment. *Curr Psychiatry Rep.* 2017;19(8):54.
  24. Bourne L, Bryant-Waugh R, Cook J, Mandy W. Avoidant/restrictive food intake disorder: a systematic scoping review of the current literature. *Psychiatry Res.* 2020;288:112961.
  25. Hay P, Mitchison D, Collado AEL, González-Chica DA, Stocks N, Touyz S. Burden and health-related quality of life of eating disorders, including Avoidant/Restrictive Food Intake Disorder (ARFID), in the Australian population. *J Eat Disorders.* 2017;5(1):21.
  26. D'Adamo L, Smolar L, Balantekin KN, Taylor CB, Wilfley DE, Fitzsimmons-Craft EE. Prevalence, characteristics, and correlates of probable avoidant/restrictive food intake disorder among adult respondents to the National Eating Disorders Association online screen: a cross-sectional study. *J Eat Disorders.* 2023;11(1):214.
  27. Manwaring JL, Blalock DV, Rienecke RD, Le Grange D, Mehler PS. A descriptive study of treatment-seeking adults with avoidant/restrictive food intake disorder at residential and inpatient levels of care. *Eat Disord.* 2024;32(1):13–28.
  28. Archibald T, Bryant-Waugh R. Current evidence for avoidant restrictive food intake disorder: implications for clinical practice and future directions. *JCPP Adv.* 2023;3(2):e12160.
  29. Willmott E, Dickinson R, Hall C, Sadikovic K, Wadhwa E, Micali N, et al. A scoping review of psychological interventions and outcomes for avoidant and restrictive food intake disorder (ARFID). *Int J Eat Disord.* 2024;57(1):27–61.
  30. MacDonald DE, Liebman R, Trotter K. Clinical characteristics, treatment course and outcome of adults treated for avoidant/restrictive food intake disorder (ARFID) at a tertiary care eating disorders program. *J Eat Disorders.* 2024;12(1):15.
  31. Goodman M, Adams N, Corneil T, Kreukels B, Motmans J, Coleman E. Size and distribution of transgender and gender nonconforming populations: a narrative review. *Endocrinol Metab Clin North Am.* 2019;48(2):303–21.
  32. Herman JL, Flores AR, O'Neill KK. How many adults and Youth identify as Transgender in the United States? The UCLA School of Law; 2022.
  33. Linsenmeyer W, Stiles D, Garwood S, Giedinghagen A, Lewis C, Strand G. The Sick, Control, One Stone, Fat, Food (SCOFF) is a valid eating disorder questionnaire to Use with Transgender Youth. *Clin Pediatr.* 2023;63(7):971–6.
  34. Linsenmeyer WR, Katz IM, Reed JL, Giedinghagen AM, Lewis CB, Garwood SK. Disordered Eating, Food Insecurity, and weight status among transgender and gender nonbinary youth and young adults: a cross-sectional study using a Nutrition Screening Protocol. *LGBT Health.* 2021;8(5):359–66.
  35. Zickgraf HF, Garwood SK, Lewis CB, Giedinghagen AM, Reed JL, Linsenmeyer WR. Validation of the nine-item Avoidant/Restrictive Food Intake Disorder Screen among Transgender and Nonbinary Youth and Young adults. *Transgender Health.* 2023;8(2):159–67.
  36. Zickgraf HF, Ellis JM. Initial validation of the Nine Item Avoidant/Restrictive Food Intake disorder screen (NIAS): a measure of three restrictive eating patterns. *Appetite.* 2018;123:32–42.
  37. n der Miesen ALR, Hurley H, Bal AM, de Vries ALC. Prevalence of the Wish to be of the Opposite gender in adolescents and adults with Autism Spectrum Disorder. *Arch Sex Behav.* 2018;47(8):2307–17.
  38. Goetz TG, Adams N. The transgender and gender diverse and attention deficit hyperactivity disorder nexus: a systematic review. *J Gay Lesbian Mental Health.* 2022:1–18.
  39. Antshel KM, Zhang-James Y, Faraone SV. The comorbidity of ADHD and autism spectrum disorder. *Expert Rev Neurother.* 2013;13(10):1117–28.
  40. Cooper K, Smith LGE, Russell AJ. Gender identity in Autism: sex differences in Social Affiliation with gender groups. *J Autism Dev Disord.* 2018;48(12):3995–4006.
  41. Warrier V, Greenberg DM, Weir E, Buckingham C, Smith P, Lai M-C, et al. Elevated rates of autism, other neurodevelopmental and psychiatric diagnoses, and autistic traits in transgender and gender-diverse individuals. *Nat Commun.* 2020;11(1):3959.
  42. Huisman B, Noens I, Steensma TD, van der Kreukels BPC. Autism traits in transgender and gender-diverse adults seeking gender-affirming medical treatment. *Int J Transgender Health.* 2024:1–17.
  43. Kallitsounaki A, Williams DM, Implicit EG-R, Cognition G, Dysphoria. Autistic-like traits, and Mentalizing: differences between autistic and non-autistic Cisgender and Transgender adults. *Arch Sex Behav.* 2022;51(7):3583–600.
  44. Kung KTF. Autistic traits, systemising, empathising, and theory of mind in transgender and non-binary adults. *Mol Autism.* 2020;11(1):73.
  45. Kallitsounaki A, Williams DM. Autism spectrum disorder and Gender Dysphoria/Incongruence. A systematic literature review and Meta-analysis. *J Autism Dev Disord.* 2023;53(8):3103–17.
  46. Watts R, Archibald T, Hembry P, Howard M, Kelly C, Loomes R, et al. The clinical presentation of avoidant restrictive food intake disorder in children and adolescents is largely independent of sex, autism spectrum disorder and anxiety traits. *eClinicalMedicine.* 2023;63:102190.
  47. Brede J, Babb C, Jones C, Elliott M, Zanker C, Tchanturia K, et al. For me, the Anorexia is just a Symptom, and the cause is the autism: investigating restrictive eating disorders in Autistic Women. *J Autism Dev Disord.* 2020;50(12):4280–96.
  48. Kinnaird E, Norton C, Stewart C, Tchanturia K. Same behaviours, different reasons: what do patients with co-occurring anorexia and autism want from treatment? *Int Rev Psychiatry.* 2019;31(4):308–17.
  49. Mansour S, Rozenblat V, Fuller-Tyszkiewicz M, Paganini C, Treasure J, Krug I. Emotions mediate the relationship between autistic traits and disordered eating: a new autistic-emotional model for eating pathology. *Psychiatry Res.* 2016;245:119–26.
  50. Kaisari P, Dourish CT, Higgs S. Attention deficit hyperactivity disorder (ADHD) and disordered eating behaviour: a systematic review and a framework for future research. *Clin Psychol Rev.* 2017;53:109–21.
  51. Hanč T, Cortese S. Attention deficit/hyperactivity-disorder and obesity: a review and model of current hypotheses explaining their comorbidity. *Neurosci Biobehavioral Reviews.* 2018;92:16–28.
  52. Nimbley E, Golds L, Sharpe H, Gillespie-Smith K, Duffy F. Sensory processing and eating behaviours in autism: a systematic review. *Eur Eat Disorders Rev.* 2022;30(5):538–59.
  53. Scheerer NE, Pourtousi A, Yang C, Ding Z, Stojanoski B, Anagnostou E, et al. Transdiagnostic Patterns of Sensory Processing in Autism and ADHD. *J Autism Dev Disord.* 2024;54(1):280–92.
  54. Dwyer P, Williams ZJ, Lawson WB, Rivera SM. A trans-diagnostic investigation of attention, hyper-focus, and monotropism in autism, attention



- dysregulation hyperactivity development, and the general population. *Neurodiversity*. 2024;2:27546330241237883.
55. Charlton RA, Entecott T, Belova E, Nwaordu G. It feels like holding back something you need to say: autistic and non-autistic adults accounts of sensory experiences and stimming. *Res Autism Spectr Disorders*. 2021;89:101864.
  56. Price A, Sumner P, Powell G. Subjective sensory sensitivity and its relationship with anxiety in people with probable migraine. *Headache: J Head Face Pain*. 2021;61(9):1342–50.
  57. Engel-Yeger B, Muzio C, Rinosi G, Solano P, Geoffroy PA, Pompili M, et al. Extreme sensory processing patterns and their relation with clinical conditions among individuals with major affective disorders. *Psychiatry Res*. 2016;236:112–8.
  58. Saure E, Laasonen M, Raevuori A. Anorexia nervosa and comorbid autism spectrum disorders. *Curr Opin Psychiatry*. 2021;34(6).
  59. Cobbaert L, Hay P, Mitchell PB, Roza SJ, Perkes I. Sensory processing across eating disorders: a systematic review and meta-analysis of self-report inventories. *Int J Eat Disord*. 2024;57(7):1465–88.
  60. Miller LJ, Anzalone ME, Lane SJ, Cermak SA, Osten ET. Concept evolution in sensory integration: a proposed nosology for diagnosis. *Am J Occup Ther*. 2007;61(2):135–40.
  61. Ward J, Hoadley C, Hughes JEA, Smith P, Allison C, Baron-Cohen S, et al. Atypical sensory sensitivity as a shared feature between synaesthesia and autism. *Sci Rep*. 2017;7(1):41155.
  62. Ben-Sasson A, Gal E, Fluss R, Katz-Zetler N, Cermak SA. Update of a Meta-analysis of sensory symptoms in ASD: a New Decade of Research. *J Autism Dev Disord*. 2019;49(12):4974–96.
  63. Cooper K, Mandy W, Butler C, Russell A. The lived experience of gender dysphoria in autistic adults: an interpretative phenomenological analysis. *Autism*. 2022;26(4):963–74.
  64. Dellapiazza F, Michelon C, Vernhet C, Muratori F, Blanc N, Picot M-C, et al. Sensory processing related to attention in children with ASD, ADHD, or typical development: results from the ELENA cohort. *Eur Child Adolesc Psychiatry*. 2021;30(2):283–91.
  65. Bijlenga D, Tjon-Ka-Jie JYM, Schuijers F, Kooij JJS. Atypical sensory profiles as core features of adult ADHD, irrespective of autistic symptoms. *Eur Psychiatry*. 2017;43:51–7.
  66. Kamath MS, Dahm CR, Tucker JR, Huang-Pollock CL, Etter NM, Neely KA. Sensory profiles in adults with and without ADHD. *Res Dev Disabil*. 2020;104:103696.
  67. Engel-Yeger B, Ziv-On D. The relationship between sensory processing difficulties and leisure activity preference of children with different types of ADHD. *Res Dev Disabil*. 2011;32(3):1154–62.
  68. Presseller EK, Cooper GE, Thornton LM, Birgegård A, Abbaspour A, Bulik CM et al. Assessing Avoidant/Restrictive food intake disorder (ARFID) symptoms using the Nine Item ARFID screen in > 9000 Swedish adults with and without eating disorders. *Int J Eat Disord*. 2024.
  69. Faul F, Erdfelder E, Lang A-G, Buchner A. G\*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods*. 2007;39(2):175–91.
  70. Qualtrics. Qualtrics. December 2023 ed. Provo, Utah, USA2020.
  71. Rodriguez C, Oppenheimer DM. Creating a bot-tleneck for malicious AI: psychological methods for bot detection. *Behav Res Methods*. 2024;56(6):6258–75.
  72. Hoekstra RA, Vinkhuysen AAE, Wheelwright S, Bartels M, Boomsma DI, Baron-Cohen S, et al. The construction and validation of an abridged version of the autism-spectrum quotient (AQ-Short). *J Autism Dev Disord*. 2011;41(5):589–96.
  73. Baron-Cohen S, Wheelwright S, Skinner R, Martin J, Clubley E. The autism-spectrum quotient (AQ): evidence from Asperger Syndrome/High-Functioning autism, males and females, scientists and mathematicians. *J Autism Dev Disord*. 2001;31(1):5–17.
  74. Kessler RC, Adler L, Ames M, Demler O, Faraone S, Hiripi EVA, et al. The World Health Organization adult ADHD self-report scale (ASRS): a short screening scale for use in the general population. *Psychol Med*. 2005;35(2):245–56.
  75. Stanton K, Forbes MK, Zimmerman M. Distinct dimensions defining the adult ADHD self-report scale: implications for assessing inattentive and hyperactive/impulsive symptoms. *Psychol Assess*. 2018;30(12):1549–59.
  76. Robertson AE, Simmons DR. The relationship between sensory sensitivity and autistic traits in the General Population. *J Autism Dev Disord*. 2013;43(4):775–84.
  77. Hwang YI, Arnold S, Srasuebku P, Trollor J. Understanding anxiety in adults on the autism spectrum: an investigation of its relationship with intolerance of uncertainty, sensory sensitivities and repetitive behaviours. *Autism*. 2019;24(2):411–22.
  78. Fairburn CG, Beglin SJ. Assessment of eating disorders: interview or self-report questionnaire? *Int J Eat Disord*. 1994;16(4):363–70.
  79. Nagata JM, Compte EJ, McGuire FH, Brown TA, Lavender JM, Murray SB, et al. Investigating the factor structure and measurement invariance of the eating disorder examination-questionnaire (EDE-Q) in a community sample of gender minority adults from the United States. *Int J Eat Disord*. 2023;56(8):1570–80.
  80. Nagata JM, Murray SB, Compte EJ, Pak EH, Schauer R, Flentje A, et al. Community norms for the eating disorder examination Questionnaire (EDE-Q) among transgender men and women. *Eat Behav*. 2020;37:101381.
  81. Burton Murray H, Dreier MJ, Zickgraf HF, Becker KR, Breithaupt L, Eddy KT, et al. Validation of the nine item ARFID screen (NIAS) subscales for distinguishing ARFID presentations and screening for ARFID. *Int J Eat Disord*. 2021;54(10):1782–92.
  82. Bourne L, Mandy W, Bryant-Waugh R. Avoidant/restrictive food intake disorder and severe food selectivity in children and young people with autism: a scoping review. *Dev Med Child Neurol*. 2022;64(6):691–700.
  83. Fonseca NKO, Curtarelli VD, Bertolotti J, Azevedo K, Cardinal TM, Moreira JD, et al. Avoidant restrictive food intake disorder: recent advances in neurobiology and treatment. *J Eat Disord*. 2024;12(1):74.
  84. Kuschner ES, Eisenberg IW, Orionzi B, Simmons WK, Kenworthy L, Martin A et al. A preliminary study of self-reported food selectivity in adolescents and young adults with autism spectrum disorder. *Res Autism Spectr Disorders*. 2015;15–6:53–9.
  85. Koomar T, Thomas TR, Pottschmidt NR, Lutter M, Michaelson JJ. Estimating the prevalence and genetic risk mechanisms of ARFID in a large autism cohort. *Front Psychiatry*. 2021;12.
  86. Zickgraf HF, Richard E, Zucker NL, Wallace GL. Rigidity and sensory sensitivity: independent contributions to selective eating in children, adolescents, and young adults. *J Clin Child Adolesc Psychol*. 2022;51(5):675–87.
  87. Xiao Y, Han J, Zickgraf HF, Barnhart WR, Zhao Y, He J. Questionnaire-assessed atypical sensory sensitivity is prospectively related to higher ARFID psychopathology, higher food neophobia, and lower satisfaction with food-related life in an online sample of Chinese older adults. *Food Qual Prefer*. 2025;122:105303.
  88. Young RL, Smith P, Lim A, Short M. The relationship between Autism Diagnostic Criteria and Problematic Eating behaviors. *Adv Neurodevelopmental Disorders*. 2022;6(1):53–64.
  89. Panagiotidi M, Overton PG, Stafford T. The relationship between ADHD traits and sensory sensitivity in the general population. *Compr Psychiatry*. 2018;80:179–85.
  90. Calisan Kinter R, Ozbaran B, Inal Kaleli I, Kose S, Bildik T, Ghaziuddin M. The sensory profiles, eating behaviors, and quality of life of children with Autism Spectrum disorder and Avoidant/Restrictive food intake disorder. *Psychiatr Q*. 2024;95(1):85–106.
  91. Zhang S, Hu R, Zhao S. Autistic traits and ARFID-associated eating behaviors in preschoolers: mediating effects of sensory processing patterns. *Appetite*. 2024;196:107237.
  92. Hargitai LD, Livingston LA, Waldren LH, Robinson R, Jarrold C, Shah P. Attention-deficit hyperactivity disorder traits are a more important predictor of internalising problems than autistic traits. *Sci Rep*. 2023;13(1):31.
  93. Thomas KS, Cooper K, Jones CRG. The intersection of autistic traits, ADHD traits, and gender diversity in disordered eating and drive for muscularity within the general population. *In press*.
  94. Walsh RJ, Krabbendam L, Dewinter J, Beeger S. Brief report: gender identity differences in autistic adults: associations with Perceptual and Socio-cognitive profiles. *J Autism Dev Disord*. 2018;48(12):4070–8.
  95. Tavassoli T, Hoekstra RA, Baron-Cohen S. The sensory perception quotient (SPQ): development and validation of a new sensory questionnaire for adults with and without autism. *Mol Autism*. 2014;5(1):29.
  96. Carey M, Kupeli N, Knight R, Troop NA, Jenkinson PM, Preston C. Eating disorder examination Questionnaire (EDE-Q): norms and psychometric properties in U.K. females and males. *Psychol Assess*. 2019;31(7):839–50.
  97. Norris ML, Spettigue W, Hammond NG, Katzman DK, Zucker N, Yelle K, et al. Building evidence for the use of descriptive subtypes in youth with avoidant restrictive food intake disorder. *Int J Eat Disord*. 2018;51(2):170–3.
  98. Coglan L, Otasowie J. Avoidant/restrictive food intake disorder: what do we know so far? *BJPsych Adv*. 2019;25(2):90–8.

99. Keery H, LeMay-Russell S, Barnes TL, Eckhardt S, Peterson CB, Lesser J, et al. Attributes of children and adolescents with avoidant/restrictive food intake disorder. *J Eat Disorders*. 2019;7(1):31.
100. Zanna V, Criscuolo M, Mereu A, Cinelli G, Marchetto C, Pasqualetti P, et al. Restrictive eating disorders in children and adolescents: a comparison between clinical and psychopathological profiles. *Eating and Weight disorders - studies on Anorexia. Bulimia Obes*. 2021;26(5):1491–501.
101. Cooper K, Mandy W, Russell A, Butler C. Healthcare clinician perspectives on the intersection of autism and gender dysphoria. *Autism*. 2023;27(1):31–42.
102. Mayer JL. The relationship between autistic traits and atypical sensory functioning in Neurotypical and ASD adults: a Spectrum Approach. *J Autism Dev Disord*. 2017;47(2):316–27.
103. Horder J, Wilson CE, Mendez MA, Murphy DG. Autistic traits and abnormal sensory experiences in adults. *J Autism Dev Disord*. 2014;44(6):1461–9.
104. Jones DR, Botha M, Ackerman RA, King K, Sasson NJ. Non-autistic observers both detect and demonstrate the double empathy problem when evaluating interactions between autistic and non-autistic adults. *Autism*. 2024;28(8):2053–65.
105. Cheang RT, Skjervling M, Blakemore AI, Kumari V, Puzzo I. Do you feel me? Autism, empathic accuracy and the double empathy problem. *Autism*. 2020;24(10):13623613241252320.
106. Milton DEM. On the ontological status of autism: the 'double empathy problem'. *Disabil Soc*. 2012;27(6):883–7.
107. Milton D, Gurbuz E, López B. The 'double empathy problem': ten years on. *Autism*. 2022;26(8):1901–3.
108. Crompton CJ, Ropar D, Evans-Williams CV, Flynn EG, Fletcher-Watson S. Autistic peer-to-peer information transfer is highly effective. *Autism*. 2020;24(7):1704–12.
109. Candia-Rivera D, Engelen T, Babo-Rebello M, Salamone PC. Interoception, network physiology and the emergence of bodily self-awareness. *Neurosci Biobehavioral Reviews*. 2024;165:105864.
110. Chen WG, Schloesser D, Arensdorf AM, Simmons JM, Cui C, Valentino R, et al. The Emerging Science of Interoception: sensing, integrating, interpreting, and regulating signals within the self. *Trends Neurosci*. 2021;44(1):3–16.
111. Kluckow H, Telfer J, Abraham S. Should we screen for misophonia in patients with eating disorders? A report of three cases. *Int J Eat Disord*. 2014;47(5):558–61.
112. Erfanian M, Kartsonaki C, Keshavarz A. Misophonia and comorbid psychiatric symptoms: a preliminary study of clinical findings. *Nord J Psychiatry*. 2019;73(4–5):219–28.
113. Nisticò V, Faggioli R, Tedesco R, Giordano B, Priori A, Gambini O, et al. Brief report: sensory sensitivity is associated with disturbed eating in adults with Autism Spectrum disorders without Intellectual Disabilities. *J Autism Dev Disord*. 2023;53(8):3295–300.
114. Duffy F, Willmott E, Nimbley E, Lawton A, Sharpe H, Buchan K, et al. Avoidant restrictive food intake disorder (ARFID)—Looking beyond the eating disorder lens? *Eur Eat Disorders Rev*. 2024;32(4):824–7.
115. Nagata JM, Ganson KT, Austin SB. Emerging trends in eating disorders among sexual and gender minorities. *Curr Opin Psychiatry*. 2020;33(6).
116. Amodeo AL, Esposito C, Antuoni S, Saracco G, Bacchini D. Muscle dysmorphia: what about transgender people? *Culture. Health Sexuality*. 2022;24(1):63–78.
117. Duffy ME, Henkel KE, Earnshaw VA. Transgender clients' experiences of eating disorder treatment. *J LGBT Issues Couns*. 2016;10(3):136–49.
118. Murray SB, Boon E, Touyz SW. Diverging eating psychopathology in Transgendered Eating Disorder patients: a report of two cases. *Eat Disord*. 2013;21(1):70–4.
119. Ewan LA, Middleman AB, Feldmann J. Treatment of anorexia nervosa in the context of transsexuality: a case report. *Int J Eat Disord*. 2014;47(1):112–5.
120. Winston AP, Acharya S, Chaudhuri S, Fellowes L. Anorexia nervosa and gender identity disorder in biologic males: a report of two cases. *Int J Eat Disord*. 2004;36(1):109–13.
121. Cusack CE, Galupo MP. Body checking behaviors and eating disorder pathology among nonbinary individuals with androgynous appearance ideals. *Eating and Weight disorders - studies on Anorexia. Bulimia Obes*. 2021;26(6):1915–25.
122. Cusack CE, Lampieri AO, Galupo M. I'm still not sure if the eating disorder is a result of gender dysphoria: Trans and nonbinary individuals' descriptions of their eating and body concerns in relation to their gender. *Psychol Sex Orientat Gen Divers*. 2022;9(4):422–33.
123. Galupo MP, Cusack CE, Morris ER. Having a non-normative body for me is about survival: androgynous body ideal among trans and nonbinary individuals. *Body Image*. 2021;39:68–76.
124. Masala B, Love A, Carmichael P, Masic U. Demographics of referrals to a specialist gender identity service in the UK between 2017 and 2020. *Clin Child Psychol Psychiatry*. 2024;29(2):624–36.
125. Wanta JW, Niforatos JD, Durbak E, Viguera A, Altinay M. Mental Health diagnoses among Transgender patients in the clinical setting: an All-Payer Electronic Health Record Study. *Transgender Health*. 2019;4(1):313–5.
126. Wittlin NM, Kuper LE, Olson KR. Mental Health of transgender and gender diverse youth. *Ann Rev Clin Psychol*. 2023;19:207–32.
127. Kidd JD, Tettamanti NA, Kaczmarkiewicz R, Corbeil TE, Dworkin JD, Jackman KB, et al. Prevalence of substance use and mental health problems among transgender and cisgender U.S. adults: results from a national probability sample. *Psychiatry Res*. 2023;326:115339.
128. Kerr-Gaffney J, Harrison A, Tchanturia K. Social anxiety in the eating disorders: a systematic review and meta-analysis. *Psychol Med*. 2018;48(15):2477–91.
129. Pallister E, Waller G. Anxiety in the eating disorders: understanding the overlap. *Clin Psychol Rev*. 2008;28(3):366–86.
130. Puccio F, Fuller-Tyszkiewicz M, Ong D, Krug I. A systematic review and meta-analysis on the longitudinal relationship between eating pathology and depression. *Int J Eat Disord*. 2016;49(5):439–54.

## Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.