Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/53944/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Krüger, F., Clare, E. L., Symondson, William Oliver Christian ORCID: <https://orcid.org/0000-0002-3343-4679>, Keišs, O. and Pēteršons, G. 2014. Diet of the insectivorous bat 'Pipistrellus nathusii' during autumn migration and summer residence. *Molecular Ecology* 23 (15) , pp. 3672-3683. 10.1111/mec.12547 file

Publishers page: <http://dx.doi.org/10.1111/mec.12547>
<<http://dx.doi.org/10.1111/mec.12547>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.

MOLECULAR ECOLOGY

Diet of the insectivorous bat *Pipistrellus nathusii* during autumn migration and summer residence

Journal:	<i>Molecular Ecology</i>
Manuscript ID:	MEC-13-0708.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Krüger, Frauke; Institute for Natural Resource Conservation, Landscape Ecology Clare, Elizabeth; School of Biological and Chemical Sciences Queen Mary University London, Symondson, William; Cardiff University, Cardiff School of Biosciences; Keiss, Oskars; Institute of Biology, University of Latvia, Laboratory of Ornithology Petersons, Gunars; Faculty of Veterinary Medicine, Latvia University of Agriculture,
Keywords:	Migration, Diet Analysis, Chiroptera, <i>Pipistrellus nathusii</i>

Diet of the insectivorous bat *Pipistrellus nathusii* during autumn migration and summer residence

Frauke Krüger^{*1}, Elizabeth L. Clare², William O.C. Symondson³, Oskars Keišk⁴, Gunārs Pētersons⁵

¹ Institute of Natural Resource Conservation, University of Kiel, Germany

² School of Biological and Chemical Sciences Queen Mary University of London, United Kingdom

³ Cardiff School of Biosciences, Sir Martin Evans Building, Cardiff University, United Kingdom.

⁴ Institute of Biology, Laboratory of Ornithology, University of Latvia, Salaspils, Latvia

⁵ Faculty of Veterinary Medicine, Latvia University of Agriculture, Jelgava, Latvia

Corresponding author:

*Frauke Krüger, Institute of Natural Resource Conservation, University of Kiel,,

fkrueger@ecology.uni-kiel.de

Key words: Chiroptera, Diet Analysis, Migration, *Pipistrellus nathusii*

Running title: Diet of the migrating bat *Pipistrellus nathusii*

Abstract

Migration is widespread among vertebrates, yet bat migration has received little attention and only in the recent decades has a better understanding of it been gained. Migration can cause significant changes in behaviour and physiology, due to increasing energy demands and aerodynamic constraints. Dietary shifts, for example, have been shown to occur in birds before onset of migration. For bats it is not known if a change in diet occurs during migration, although breeding season related dietary preference has been documented. It is known that a diet rich in fats and the accumulation of fat deposits do increase the flight range of migratory bats. Some bat species can be regarded as long-distance migrants, covering up to 2,000 km between summer and winter roosting areas. *Pipistrellus nathusii* (Vespertilionidae), a European long-distant migrant, travels each year along the Baltic Sea from north-eastern Europe to hibernate in central and southern Europe. This study presents

29 data on the dietary habits of migrating *Pipistrellus nathusii* compared with those during the
30 breeding season. We analysed faecal samples from bats on fall migration caught at the
31 Ornithological Field Station in Pape, Latvia and from samples collected in North-Latvian
32 summer roosts. We applied both morphological identification and molecular methods, as
33 morphological methods also recognize life stages of prey and can contribute frequency data.
34 The diets of bats on migration and breeding bats were similar, with Diptera and Lepidoptera
35 comprising the major prey categories. However, certain prey groups could be explained by the
36 different hunting habitats exploited during migration vs. summer residence.

37

38

39 Introduction

40 Across the animal kingdom numerous species make annual movements of both short
41 and long duration. In particular, bird migration has been intensively studied since the late 19th
42 century. In the last decades these studies have been sophisticated both in methods and
43 explanations and still are being developed further (Berthold 2001; Robinson *et al.* 2007;
44 Wikelski *et al.* 2007; Fiedler 2009). Many species of bats, the only volant mammals, are also
45 known to migrate. Although the first interest in bat migration arose as early as the end of the
46 19th century (Merriam 1887), bat migration has been largely ignored until recently. Unlike
47 birds, the elusive life strategies of bats, difficulties regarding visual observations, and low
48 success in mark recapture programs, have made these species difficult to study. However,
49 substantial recent advances have been made, which increase our understanding of orientation
50 and physiology (Holland *et al.* 2006; Cryan & Brown 2007; McGuire & Guglielmo 2009;
51 Voigt *et al.* 2010, 2012b). New techniques have contributed to our ability to track and infer
52 actual range of movement, such as satellite tracking and stable isotope analysis (Cryan *et al.*
53 2004; Richter & Cumming 2008; Popa-Lisseanu & Voigt 2009; Voigt *et al.* 2012a; Tsoar *et*
54 *al.* 2012)

55 Studies of bat migration can profit from previous work on bird migration (McGuire &
56 Guglielmo 2009). Both birds and bats need to maintain a sufficient nutrient intake to meet the
57 increased energy demand during migration over distances of sometimes several thousand
58 kilometers between summer and winter habitats (Griffin 1970; Petersons 2004). As in birds,
59 the scale of bat migration can vary considerably between short-distance, regional migrants
60 (e.g., *Myotis daubentonii*, *M. lucifugus*) and long-distance migrants (e.g., *Pipistrellus nathusii*,
61 *Lasioncyteris noctivagans*) (Fleming & Eby 2003; Dzal *et al.* 2009; Dzal *et al.* 2011). On
62 their journeys birds and bats face similar tradeoffs between acquiring sufficient fat deposits
63 (energy reserves) to fuel flight and maintaining body conditions (weight, size) optimal for
64 flight with low energetic costs.

Birds are known to start to build up fat reserves before migratory flight and during stopovers at resting sites (e.g., Wadden Sea; McWilliams *et al.* 2004). Before the onset of migration birds show different adaptations to increase fat stores: they may become hyperphagous, their digestive and biosynthetic systems may alter, for example increase liver mass and liver activity (Egeler *et al.* 2000; Guglielmo & Williams 2003), and they may increase or reduce the size of their digestive systems (Piersma 1998; Piersma *et al.* 1999; McWilliams & Karasov 2004). Additionally birds are able to fly during the night and forage and refuel during the day.

Bats have to accomplish the dual task of both flying and refueling at night. Recent studies show that bats also become hyperphagous and increase their body fat and catabolic enzyme activity during pre-migration (Ewing *et al.* 1970; Bairlain 2001; McGuire *et al.* 2009, 2013a, b; Šuba *et al.* 2010). Furthermore, they are able to fuel their migration both directly from insects caught during flight and from stored fatty acid reserves to maintain both steady state and refill reserves (Voigt *et al.* 2010; Voigt *et al.* 2012b). The fly-and-forage hypothesis, which states that bats forage on the wing during migration, is supported by acoustical observations along migration routes (Ahlén *et al.* 2009; Valdez & Cryan 2009, Šuba *et al.* 2012). Yet, it is not clear to what extent bats are segregating foraging and migratory flight during these periods or whether they can truly hunt while migrating.

Another adaptation, the shift in diet towards different food items (e.g., from insects to fruit), helps some birds to gain sufficient energy during the pre-migration period (Bairlein 1990; Bairlein & Gwinner 1994; Bairlein 2001; McWilliams & Karasov 2005). It is not known if bats show similar behaviour. While most insectivorous bats use a generalist strategy, consuming prey in relation to their abundance (Anthony & Kunz 1977; Swift *et al.* 1985) within a given habitat (Clare *et al.* 2013a/ b in review, Special Issue), selective feeding and the ability to discriminate between food items have been demonstrated in some bat species (Von der Emde & Schnitzler 1990; Koselj *et al.* 2011). Dietary shifts over time have been

described in bats (Agosta 2002) and may be related to physiological state (pregnancy, lactation, preparation for hibernation) or changes in insect abundance (Clare *et al.* 2009, 2011, 2013a/b in press, Special Issue).

Here we tested the hypothesis that bat diet differs between summer roosting and fall migration. We used high throughput sequencing which yields detailed species-level data on prey in predator diets (Symondson 2002; King *et al.* 2008), and has been particularly successful in insectivores such as bats, (Razgour *et al.* 2011; Bohmann *et al.* 2011; Clare *et al.* 2013a/b in review) and shrews (Brown *et al.* in press, Special Issue). From species-level data (DNA sequences) we can draw conclusion on differences in prey items, apparent energy values or fat content and on putative foraging area differences between summer and migration habitats. We focused on a long-distance migrating bat, *Pipistrellus nathusii* (Keyserling & Blasius 1839), a generalist pipistrelle bat, which feeds to a large extent on insects connected to aquatic habitats, mainly on Diptera, particular Chironomidae (Beck 1994-1995; Vaughan 1997; Arnold *et al.* 2000; Flaquer *et al.* 2006). This species is known to travel up to 2000 km between the summer roosting grounds and hibernacula (Petersons 2004).

Methods

Study sites and sample collection

We collected samples for the purpose of dietary analysis at the Pape Ornithological Research Station at the southern Baltic coast of Latvia (56.165° N, 21.017° E) during the fall migration between 11 August and 12 September 2012 (Fig. 1). The station has been a central site for intensive research on bird and bat migration, particularly during the past twenty years (Petersons 2004). The surrounding areas are dominated by low sand dunes, partly covered by unmanaged pine woodlands (*Pinus sylvestris*) and grasslands. In the vicinity of the station is Pape Nature Park with Lake Pape, a 12km² shallow coastal lagoon lake, and a mosaic of marshland, reeds, meadows, forests and peat bogs. We caught bats from dusk until 01:00am using a Helgoland funnel trap following Petersons (2004). Bats were placed in a clean soft cotton bags and held for approximately 1h to collect faecal samples. Samples of *P. nathusii* faeces from summer colonies were collected from nursery colonies situated in buildings, at Vecpiebalga (57.058° N; 25.815° E) and artificial roosts with male groups at Garkalne (57.048° N; 24.382° E), Latvia during June 2013 (Fig. 1). Both sites are located in a mosaic landscape of forests, pasture and in proximity to large lakes.

Molecular diet analysis from faecal samples

For the molecular analysis we extracted DNA from faecal samples from individual migrating bats (n= 35 faecal samples) and from summer colony bulk samples, non-individually collected from under the roosting bats (n = 21 faecal samples) using the QIAamp DNA Stool Mini Kit (Qiagen, UK) following Zeale *et al.* (2011). To amplify the arthropod prey DNA we used modified primers based on the universal COI primer ZBJ-ArtF1c and ZBJ-ArtR2c. PCR (following the protocol of Zeale *et al.* 2011) produce a 157bp amplicon at the 5' end of the 658bp COI barcode region (Hebert *et al.* 2004). DNA was sequenced via a high throughput Ion Torrent sequencing platform (Life Technology) at the University of Bristol Genomics facility (School of Biological Sciences, Bristol, UK). For the adjustment,

trimming and organisation of sequences by MIDs after sequencing we used the Galaxy V platform (<https://main.g2.bx.psu.edu/root>; Giardine *et al.* 2005; Blankenberg *et al.* 2005; Blankenberg *et al.* 2010; Goecks *et al.* 2010). To allow niche analysis procedure for all sequences, we clustered the sequences into molecular operational taxonomic units (MOTU) using the program jMOTU (Jones *et al.* 2011). We tested grouping thresholds from 1-10bp and selected a 4bp threshold as the most appropriate for this data set (see Razgour *et al.* 2011). We extracted representative sequences for each MOTU and compared sequences against references within the Barcode of Life Data System (Ratnasingham & Hebert 2007; Clare *et al.* 2009). If sequences matched completely to a reference sequence without matching any other arthropod, we regarded the sequence as belonging to the same species. However, the short amplicon length also constrains some species assignments. We used a modified version of the criteria used by Razgour *et al.* (2011) as follows:

- 1a. True species match (>99 % similarity)
- 1b. Likely species match (>98% similarity)
2. Match (>98%) to more than one species, only one of which belongs to local assemblage
3. Match (>98%) to several species or genera – genus or family level assignment considered provisional.

Morphological identification from faecal samples

For morphological faecal analysis, we dried samples after DNA extraction (see Zeale *et al.* 2011) at room temperature and stored them at -20°C to avoid coprophagous insects. Before analysis, we soaked the pellets for 48 h in 70% ethanol and then dissected them under a binocular vision microscope (x 40 – 60). We separated characteristic fragments and mounted them in Euparal for further examination. We identification taxa to class, order, family, or genus level (where possible), by comparison of fragments with whole collected insects, arthropod identification keys from the literature (Medvedev 1989; Savage 1990; Shiel

et al. 1997; Osterbroek *et al.* 2005) and fragment photos from earlier studies (Krüger *et al.* 2012).

For every sample we calculated the frequency of each prey group relative to all samples, to estimate relative importance of prey groups (Shiel *et al.* 1997; Vaughn 1997; Krüger *et al.* 2012).

Statistical analysis

As molecular and morphological analysis produce presence-absence data and frequency data, respectively, we analysed the two data sets in different ways.

We used the Hamming distance and Bray-Curtis index (Equation 1) for similarity to analyze the dietary differences between the migratory group and the summer group as measured by molecular data. Both indices use binomial data. The Hamming distance is related to the number of changes needed to adjust two strings of same length to each other (Hamming 1950). A smaller value for Hamming distances reflects high overlap in dietary choices. The Bray-Curtis index (Bray & Curtis 1957) measures the dissimilarity between the dietary data sets, where C_{ij} is the sum of the lesser value for only those items which both data sets have in common. S_i and S_j are the total number of items counted in both data sets. If the data sets are identical, then both predators feed on the same prey and the Bray-Curtis index is 0. If the two data sets do not share any prey items then the index is 1 (Bloom 1981).

$$BC_{ij} = \frac{2C_{ij}}{S_i + S_j} \quad (\text{Equation 1})$$

$$D = 1 - \sum_{i=1}^S \frac{n_i(n_i-1)}{n(n-1)} \quad (\text{Equation 2})$$

$$O_{jk} = \frac{\sum p_{ij}p_{ik}}{(\sum p_{ij}^2 \sum p_{ik}^2)^{1/2}} \quad (\text{Equation 3})$$

To assess dietary niche breadth based on the morphological diet data, we used the Simpson's index for diversity and heterogeneity (Equation 2), where n_i is the relative proportion of a prey item i (with $i = 1 \dots n$) of a total of n prey items. Thus, D is 0, if all eaten prey belongs to one prey group. The higher the diversity, the closer D gets to 1. As an additional niche parameter we calculated Pianka's index of niche overlap (Equation 3), where p_i is the frequency of occurrence of prey item i in the diets of species j and k (Pianka 1973):

We used a non-metric multi-dimensional scaling (NMDS) with Jaccard distance to visualize the degree of similarity or dissimilarity of the diet. The resulting two-dimensional ordination plot shows the samples sorted relative to their dissimilarity, with similar samples in close proximity and dissimilar samples further apart. We used a threshold (=stress value) of < 0.2 for ecological interpretation of the NMDS plot (Clark & Warwick 2001).

We conducted indices calculation, Adonis, and NMDS using the *vegan* R library (Oksanen *et al.* 2011). We applied generalized linear models (GLM) with a binomial distribution and a logit link function (Zuur *et al.* 2007), to assess level of significance of differences between the two data sets regarding the presence or absence of prey groups, using *multcomp* R library (Hothorn *et al.* 2008).

Results

Molecular analysis

We found 220 MOTUs, of which 148 could be assigned to insect species (Table 1). For 72 MOTUs we found no matches in the BOLD System. We rejected 1 MOTU, because it contained only very short reads. Of the MOTUs 32% could be assigned to species level, 28% to genus level, 23% to family and 17% only to order level. We found 108 MOTUs in samples from migrating bats, whereas 58 MOTUs were assigned to samples from summer roosts. 19 MOTUs were found in both groups.

Hamming Distances between migratory bats and bats at summer colonies was 197. Additionally overall Bray-Curtis similarity between migratory bats and bats at summer colonies was 0.84. Both suggest low dietary similarity.

Morphological analysis

We found that the diet was significantly different between bats from summer roosts and on migration, indicated by the conducted permutational analysis of variance (ADONIS: F = 4.371, df = 1, $p < 0.001$). Comparing diversity and species richness in the diet of *P. nathusii* between the two sites, we found no differences (Table 2). The trophic niche overlap, indicated by Pianka index, was relatively high (Table 2). The ordination plot (NMDS) shows samples spread out evenly along the two dimensions, overlapping to a great extent. The slight clustering along the first dimension has to be interpreted cautiously, as a stress value of 0.2 was reached (Fig. 2).

Based on GLMs we found significant difference between certain prey groups. *P. nathusii* from summer roosts appear to feed more often on Chironomidae than migrating *P. nathusii*. In contrast Tipulidae occurred more often in the diet of migrating bats (Table 3).

Discussion

The high values for Hamming distance and Bray-Curtis similarity we found for the molecular diet data indicate that diets of *P. nathusii* during migration and at summer colonies show low similarity. Despite this we found similar diversity indices for both groups, based on the morphological analysis, and a relatively high niche overlap. Yet, if compared to the niche overlap between different species foraging in similar habitats, for example *Myotis dasycneme* and *Myotis daubentonii*, the niche overlap between *P. nathusii* appears less strong (Krüger *et al.* 2012; Krüger *et al.* in press). Subtle but significant differences appear regarding Chironomidae and Tipulidae occurrence in the diet of migrating and summer bats, respectively. Chironomid species, especially in areas between latitude of 50° and 60°, can have two or more generations per year with diverging peaks from April to October. Several species (up to 15) can form groups which emerge in synchrony and cause an increase in potential prey biomass. As different groups follow different emergence patterns, alternating peaks of different Chironomid groups exist, replacing each other during the season and forming a more or less constant food resource (Oliver 1971, Pinder 1986, Berg and Hellenthal 1992, Tokeshi 1995, Benke 1998). The higher occurrence of Chironomidae in summer roosting bats might be explained by the reproductive state of bats and hence their needs for easily accessible prey, such as swarming Chironomidae. The higher occurrence of Tipulidae in migrating bats could be related to a peak in Tipulidae during that time at Pape, coinciding with migration paths. Diptera too are thought to migrate (Hogsette & Ruff 1985), and the tracks of migrating *P. nathusii* and tipulids may have coincided. Alternatively the bats may have been hunting more often in terrestrial habitats during this period.

A significant issue here is whether the difference is due to “migration” or “location”. Clare *et al.* (In press) demonstrated the degree of location, season and inter-annual variation in bat diet across landscapes. Since most bats forage among prey in their immediate vicinity it is not clear to what degree shifts in diet observed here are caused by changes in insect

phenology, bat physiological demands and habitat-insect relationships. We report here that there is variation between summer colonies and migrating individuals, though the underlying cause is very likely a mix of these competing factors. We suggest that the next logical step is to expand this type of study to include multiple summer and migration sites so that comparisons can be made across locations.

The higher resolution of molecular diet analyses of prey species, compared with morphological analyses, provides valuable information on associations between prey, habitat and predator (Clare *et al.* 2010; Razgour *et al.* 2011, Clare *et al.* 2013a in press; Krüger *et al.* in press). While our morphological observations suggest that diet was strongly overlapping between summer and migratory groups, we did observe a higher species richness in the migratory diet based on the molecular data. There are two potential drivers of increased diversity in the diet of migrating *P. nathusii*. First, migrating bats cover more space and more potential habitat types. This may expose them to a higher diversity of potential prey as a consequence. Second, insect diversity is general reduced later in the summer. At this point a reduced availability of prey may force the *P. nathusii* to become more flexible in the prey they consume (Clare *et al.* 2013a/b in press).

We also found that the diet of migrating *P. nathusii* contained higher occurrence of insect species inhabiting aquatic habitats like the beetles *Cyphon phragmiteticola* and *Agonum piceum*. This probably reflects the fact that *P. nathusii* forage in the adjacent bog and marsh lands of Lake Pape. The moths *Epinotia immundana*, *Epinotia nisella* and *Phyllonorycter apparella* are associated with riverine forests and trees in marshes, supporting the inference that bats forage in the vicinity of aquatic habitats. Further indications for aquatic foraging habitats are the occurrences of Trichoptera and Megaloptera. By contrast the moth *Malacosoma castrensis* indicates foraging over dunes, as this is the major habitat of this moth. The dunes at Pape spread out parallel to the coastline, and are also used by *P. nathusii* as a

major flight corridor during migration (Šuba *et al.* 2012) suggesting prey habitat and predator habitat overlap at this point.

In the diet of *P. nathusii* from summer colonies we found prey species which are typically associated with forested areas, like *Bupalus pinaria*, a pine pest species, or *Promethes sulcator*, an ichneumon wasp. These species were not identified in samples from migrating *P. nathusii*. As the colony sites are also within a few kilometres of lakes, we also found prey species associated with aquatic habitats, like Chironomidae or Ephemeroptera. Overall we can observe how the foraging habitat of *P. nathusii* determines the diet and thus differences between migrating and summering bats might be explained. In birds it has been shown that during migration sedge warblers (*Acrocephalus schoenobaenus*) select stop-over sites with high abundance of aphids (Bibby & Green 1981). Insectivorous bats, like *P. nathusii*, are known to forage particularly in habitats with high insect abundance like riverine and semi-aquatic habitats.

Many insect species are also known to migrate (e.g. Hummingbird Hawk-moth *Macroglossum stellatarum*, Monarch butterfly *Danaus plexippus*). The beet army worm, *Spodoptera exigua*, originally distributed in the Americas, now occurring globally, is also a known long-distance migrant (Westbrook 2008). In Europe this species has been observed to travel long distances, from Russia over Fennoscandia towards Denmark and the British Isles (Mikkola 1970). The occurrence of a migrating insect in the diets of migrating bats may be a coincidental overlap of migration routes and the opportunistic foraging behaviour of pipistrelles, which has been also observed in other species such as *Tadarida brasiliensis* which feeds opportunistically on migrating moths (Lee & McCracken 2005). Hoary bats are believed to time migration with the mass emergence of moths, its major prey (Valdez & Cryan 2009). In Europe the noctula bat *Nyctalus lasiopterus* has been found to exploit migrating songbirds during spring and autumn migration (Ibáñez *et al.* 2001; Popa-Lisseanu

et al. 2007). Similar behavior has been also reported for the birdlike noctula, *Nyctalus aviator*, in Japan (Fukui et al. 2013).

Energetic values of insect prey might influence foraging behaviour and diet preference in migrating bats. Due to the high costs of migration flights, bats might prefer prey with high fat content and high nutritional value, to maintain energy flow and fat deposits. The variation in insect nutritional values is high, with large moths or beetles showing relatively higher fat content than many other groups (Verkerk et al. 2007). In addition, some migrating insect also accumulate fat to survive long dispersal flights, for example noctuid moths (Beall 1948; Angelo & Slansky 1984; Kevan & Kendall 1997). The occurrence of *Spodoptera exigua* (Noctuidae), and other Lepidoptera and Coleoptera in the diet of migrating *P. nathusii*, suggest that these bats feed on prey with high fat content. Voigt et al. (2010, 2012b) showed how *P. nathusii* and also other bat species fuel their flight during migration with endogenous fatty acids from adipocytes in combination with proteins and carbohydrates directly metabolised from exogenous sources, such as insects. This process is determined by a limited capacity for energy storage and primarily saves energy during the costly process of converting macronutrients to lipids for storage. Nevertheless, it can be regarded as a beneficial digestive adaptation to flying and migration. Hence, the selective exploitation of prey with high fat content would enable *P. nathusii* to balance its fat reserves despite their high energy demands from long-distance and foraging flights. Birds are known to alter their dietary preference and select different food sources shortly before or during migration. Geese have been shown to select for certain plants species during migration (Gwinner 1990). Insectivorous migratory birds like garden warbler (*Sylvia borin*) switch from arthropod based diet to fruit based diets (Bairlein 1990) and furthermore select for fruit with certain fatty acid compositions prior to migration (McWilliams et al. 2004). Other birds like willow warblers switch to insect prey high in sugars, like aphids (Berthold 2001). Adaptive alteration of diet selectivity during migration seems to be a valuable trait in migrants. .

This kind of selectivity would require the bats to discriminate between prey of different energy values. In bats selective behaviour and prey discrimination based on size has previously been demonstrated only for horseshoe bats (Koselj *et al.* 2011). Whether the same ability exists for other bats, like *Pipistrellus*, which, in contrast to horseshoe bats, use short frequency-modulated (FM) calls and mainly feed on Diptera on the wing, is not clear. Also like for frugivory in insectivorous birds during migration, the differences in diet between summer and migratory *P. nathusii* may result from the seasonal changes in availability of certain food items, insects and fruit, respectively.

The fat stores of migrating bats have higher proportions of polyunsaturated fatty acids (PUFAs) (McGuire *et al.* 2013b). Thus PUFAs may be an important resource during pre-migration and migration itself. Naturally, the diet of *Pipistrellus nathusii* is often dominated by Diptera, particularly Chironomidae, which are rich in highly unsaturated fatty acids (Thompson 1982; Hanson *et al.* 1985). Thus increased lipid biosynthesis capacity and additional intake of bigger, or fatter prey, may not be required during migration (McGuire & Guglielmo 2009).

In general, our results demonstrate differences in the diet of *P. nathusii* in summer roosts and on migration. *P. nathusii* is a generalist predator and feeds on prey groups thought to be rich in important fatty acids (e.g. Chironomidae) thus the need to select for prey with particularly high fat content during migration might be low. Yet, there is no evidence that endogenous triggered selectivity can be observed in insectivorous bats as is the case with some insectivorous and grazing birds.

Additionally, the ability of pipistrelle bats to discriminate between prey of differing energetic values might be poor and hamper shifts in prey selection. Diet of migrating bats like *P. nathusii* might rather depend on the availability of prey at the respective stop-over site and the differences between migrating and summering individuals found in prey groups are likely to be related to habitat differences along migratory routes and in the summering grounds. For

355 the future it would be beneficial to find and add more migratory stop-over sites, where species
356 can be studied.

357

358

For Review Only

Acknowledgments

We are grateful to the Baltic-German University Liaison Office / DAAD for financial support. Many thanks to Alma Vītola, Ilze Brila, Jurgis Šuba for help in the field. We thank the two anonymous reviewers for valuable comments on an earlier version of this manuscript.

Data Accessibility

Molecular (sequences) and morphological (binary presence-absence) dietary data will be provided on a DRYAD account (doi: 10.5061/dryad.2d38f).

References

- Agosta SJ (2002) Habitat use, diet and roost selection by the Big Brown Bat (*Eptesicus fuscus*) in North America: a case for conserving an abundant species. *Mammal Review* **32**, 179-198.
- Ahlén I, Baagø H, BachL(2009)Behaviour of Scandinavian bats during migration and foraging at sea. *Journal of Mammalogy*, **90**, 1318-1323.
- Angelo MJ, SlanskyJr F (1984) Body building by insects: trade-offs in resource allocation with particular reference to migratory species. *Florida Entomologist*, 22-41
- Anthony, E.L.P., Kunz, T. H., 1977. Feeding Strategies of the Little Brown Bat, *Myotis lucifugus*, in Southern New Hampshire. *Ecology*, **58**, 775-786.
- Bairlein F (1990) Nutrition and food selection in migratory birds. In: *Bird migration: physiology and ecophysiology* (ed Gwinner E), pp 198–213. Springer, Berlin Heidelberg New York.
- Bairlein F (2001). Results of bird ringing in the study of migration routes and behaviour.*Ardea*, **89**, 7-19.
- Bairlein F,GwinnerE (1994) Nutritional mechanisms and temporal control of migratory energy accumulation in birds. *Annual Review of Nutrition*, **14**,187-215
- Beall F (1948) The fat content of a butterfly, *Danaus plexippus* Linn., as affected by migration. *Ecology*, **29**, 80-94.
- Beck A (1994-95). Fecal analyses of European bat species. *Myotis*, **32-33**, 109-119.

- 388 Benke AC (1998) Production Dynamics of Riverine Chironomids: Extremely High Biomass
389 Turnover Rates of Primary Consumers. *Ecology*, **79**, 899-910.
- 390 Berg MB, Hellenthal RA (1992) The role of Chironomidae in energy flow of a lotic
391 ecosystème. *Netherlands Journal of Aquatic Ecology*, **26**, 471–476.
- 392 Berthold P (2001). *Bird migration: a general survey*. Oxford University Press.
- 393 Blankenberg D, Taylor J, Schenck I, He J, Zhang Y, Ghent M, Veeraraghavan N, Albert I,
394 Miller W, Makova KD, Hardison RC, Nekrutenko A (2007) A framework for
395 collaborative analysis of ENCODE data: Making large-scale analyses biologist-
396 friendly. *Genome Research*, **17**, 960-964.
- 397 Blankenberg D, Gordon A, Von Kuster G, Coraor N, Taylor J, Nekrutenko A, The Galaxy
398 Team (2010) Manipulation of FASTQ data with Galaxy. *Bioinformatics Application*
399 *Note*, **26**, 1783-1785.
- 400 Bloom SA (1982) Similarity indices in community studies: Potential pitfalls. *Marine Ecology*
401 *– Progress Series*, **5**, 125-128.
- 402 Bohmann K, Monadjem A, Lehmkuhl-Noer C, Rasmussen M, Zeale MRK, Clare EL, Jones G,
403 Willerslev E, Gilbert MTP (2011) Molecular Diet Analysis of Two African Free-
404 Tailed Bats (Molossidae) Using High Throughput Sequencing. *PLOS one*, **6**, 1-11.
- 405 Brown DS, Burger R, Cole N, Vencatasamy D, Clare EL, Montazam A, Symondson WOC (in
406 press) Dietary competition between the alien Asian Musk Shrew (*Suncus murinus*) and
407 a reintroduced population of Telfair's Skink (*Leiolopisma telfairii*). *Molecular*
408 *Ecology*.
- 409 Clare EL, Fraser EE, Braid HE, Fenton MB, Hebert PD (2009) Species on the menu of a
410 generalist predator, the eastern red bat (*Lasiurus borealis*): using a molecular approach
411 to detect arthropod prey. *Molecular Ecology*, **18**, 2532–2542.
- 412 Clare EL, Barber BR, Sweeney BW, Hebert PDN, Fenton MB (2011) Eating local: influences
413 of habitat on the diet of little brown bats (*Myotis lucifugus*). *Molecular Ecology*, **20**,
414 1772-1780.
- 415 Clare EL, Symondson WOC, Broders H et al. (THIS ISSUEa) The diet of *Myotis lucifugus*
416 across Canada: assessing foraging quality and diet variability. *Molecular Ecology*,
417 (XXXX-XXXX)

418 Clare EL, Symondson WOC, Fenton MB (THIS ISSUEb) An inordinate fondness for beetles?
 419 Variation in seasonal dietary preferences of night roosting big brown bats (*Eptesicus*
 420 *fuscus*). *Molecular Ecology*, (XXXX-XXXX)

421 Clarke KR, Warwick RM (2001) Change in marine communities: an approach to statistical
 422 analysis and interpretation, 2nd ed. PRIMER-E, Plymouth.

423 Cryan PM, Bogan MA, Rye RO, Landis GP, Kester CL (2004) Stable hydrogen isotope
 424 analysis of bat hair as evidence for seasonal molt and long-distance migration. *Journal*
 425 *of Mammalogy*, **85**, 995-1001.

426 Cryan PM, Brown AC (2007) Migration of bats past a remote island offers clues toward the
 427 problem of bat fatalities at wind turbines. *Biological Conservation*, **139**, 1-11.

428 Dzal Y, Hooton LA, Clare EL, Fenton MB (2009). Bat activity and genetic diversity at Long
 429 Point, Ontario, an important bird stopover site. *Acta Chiropterologica*, **11**, 307-315.

430 Dzal Y, McGuire LP, Veselka N, Fenton MB (2011). Going, going, gone: the impact of
 431 white-nose syndrome on the summer activity of the little brown bat (*Myotis lucifugus*).
 432 *Biology letters*, **7**, 392-394.

433 Egeler O, Williams TD, Guglielmo CG (2000) Modulation of lipogenic enzymes, fatty acid
 434 synthase and $\Delta 9$ -desaturase, in relation to migration in the western sandpiper (*Calidris*
 435 *mauri*). *Journal of Comparative Physiology B*, **170**, 169-174

436 Ewing WG, Studier EH, O'Farrell MJ (1970) Autumn fat deposition and gross body
 437 composition in three species of *Myotis*. *Comparative Biochemistry and Physiology*,
 438 **36**, 119-129.

439 Fiedler W (2009) New technologies for monitoring bird migration and behaviour. *Ringling &*
 440 *Migration*, **24**, 175-179.

441 Fleming TH, Eby P (2003) Ecology of bat migration. In: *Bat Ecology* (eds Kunz TH &Fenton
 442 MB), pp. 156-208. University of Chicago Press, Chicago, USA.

443 Fukui D, Dewa H, Katsuta S, Sato A (2013) Bird predation by the birdlike noctula in Japan.
 444 *Journal of Mammalogy*, **94**, 657-661.

445 Giardine B, Riemer C, Ross CH *et al.* (2005) Galaxy: A platform for interactive large-scale
 446 genome analysis. *Genome Research*, **15**, 1451-1455.

447 Goecks J, Nekrutenko A, Taylor J, The Galax Team (2010) Galaxy: a comprehensive
 448 approach for supporting accessible, reproducible, and transparent computational
 449 research in the life sciences. *Genome Biology*, doi:10.1186.

- Griffin DR (1970) Griffin, D. R. (1970). Migration and homing of bats. In: *Biology of bats*, Vol 2, (ed Wimsatt WA), pp. 233-264. Academic Press, New York.
- Guglielmo CG, Williams TD (2003) Phenotypic flexibility of body composition in relation to migratory state, age, and sex in the western sandpiper (*Calidris mauri*). *Physiological and Biochemical Zoology*, **76**, 84-98.
- Hamming RW (1950) Error detecting and error correcting codes. *The Bell System Technical Journal*, **29**, 147-160.
- Hanson BJ, Cummins KW, Cargill AS, Lowry RR (1985) Lipid content, fatty acid composition, and the effect of diet on fats of aquatic insects. *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry*, **80**, 257-276.
- Hebert PDN, Stoeckle MY, Zemplak TS, Francis CM (2004) Identification of birds through DNA barcodes. *PLoS Biology*, **2**, e312.
- Hogsette JA, Ruff JP (1985) Stable fly (Diptera: Muscidae) migration in northwest Florida. *Environmental entomology*, **14**, 170-175.
- Holland RA, Thorup K, Vonhof MJ, Cochran WW, Wikleski M (2006) Bat orientation using Earth's magnetic fields. *Nature*, **44**, 702.
- Holland RA, Borissov I, Siemers BM (2010) A nocturnal mammal, the greater mouse-eared bat, calibrates a magnetic compass by the sun. *Proceedings of the National Academy of Sciences*, **107**, 6941-6945.
- Hothorn T, Bretz F, Westfall P (2008) Simultaneous Inference in General Parametric Models. *Biometrical Journal*, **50**, 346 - 363.
- Jones M, Ghoorah A, Blaxter M (2011) jMOTU and Taxonator: turning DNA Barcode sequences into annotated operational taxonomic units. *PLoS one*, **6**, e19259.
- Kevan PG, Kendall DM (1997) Liquid assets for fat bankers: Summer nectarivory by migratory moths in the Rocky Mountains, Colorado, U.S.A. *Arctic and Alpine Research*, **29**, 478-482.
- Ibáñez C, Juste J, García-Mudarra JL, Agirre-Mendi PT (2001) Bat predation on nocturnally migrating birds. *Proceedings of the National Academy of Sciences*, **98**, 9700-9702.
- King RA, Read DS, Traugott M, Symondson WOC (2008) Molecular analysis of predation: a review of best practice for DNA-based approaches. *Molecular Ecology*, **17**, 947 - 963.
- Koselj K, Schnitzler H-U, Siemers BM (2011) Horseshoe bats make adaptive prey-selection decisions, informed by echo cues. *Proceedings of the Royal Society B*, **278**, 3034-3041.

- 483 Krüger F, Harms I, Fichtner A, Wolz I, Sommer RS (2013) High trophic similarity in the
 484 sympatric North European trawling bat species *Myotis daubentonii* and *Myotis*
 485 *dasycneme*. *Acta Chiropterologica*, **14**, 347–356.
- 486 Krüger F, Clare EL, Greif S, Siemers BM, Symondson WOC, Sommer RS (THIS ISSUE) An
 487 integrative approach to detect subtle trophic niche differentiation in the sympatric
 488 trawling bat species *Myotis dasycneme* and *Myotis daubentonii*. *Molecular Ecology*,
 489 (XXXX-XXXX)
- 490 Lee Y, McCracken GF (2009) Dietary variation of Brazilian free-tailed bats links to migratory
 491 populations of pest insects. *Journal of Mammalogy*, **86**, 67-76.
- 492 McGuire LP, Fenton MB, Guglielmo CG (2009) Effect of age on energy storage during
 493 prehibernation swarming in little brown bats (*Myotis lucifugus*). *Canadian Journal of*
 494 *Zoology*, **87**, 515-519.
- 495 McGuire L, Guglielmo CG (2009) What can birds tell us about the migration physiology of
 496 bats? *Journal of Mammalogy*, **90**, 1290-129.
- 497 McGuire LP, Fenton MB, Guglielmo CG (2013a). Phenotypic flexibility in migrating bats:
 498 seasonal variation in body composition, organ sizes and fatty acid profiles. *The*
 499 *Journal of experimental biology*, **216**, 800-808.
- 500 McGuire LP, Fenton MB, Guglielmo CG (2013b) Seasonal upregulation of catabolic enzymes
 501 and fatty acid transporters in the flight muscle of migrating hoary bats,
 502 *Lasiurus cinereus*. *Comparative Biochemistry and Physiology B*, **165**, 138-143.
- 503 McWilliams SR, Guglielmo C, Pierce B, Klaassen M (2004) Flying, fasting and feeding in
 504 birds during migration: a nutritional and physiological ecology perspective. *Journal of*
 505 *Avian Biology*, **35**, 377–393.
- 506 McWilliams SR, Karasov WH (2004) Migration takes guts: digestive physiology of migratory
 507 birds and its ecological significance. In: *Birds of Two Worlds* (eds Marra P Greenberg
 508 R). Johns Hopkins University Press, Baltimore, M. D.
- 509 Medvedev GS (Ed.) (1989) Keys to the Insects of the European Part of the USSR.:
 510 Lepidoptera. IV, 1 (Vol. 117). Brill Archive.
- 511 Merriam CH (1887) Do any Canadian bats migrate? Evidence in the affirmative. *Transaction*
 512 *of the Royal Society of Canada*, **5**, 85-87.
- 513 Mikkola K (1970) The interpretation of long-range migrations of *Spodoptera exigua*
 514 Hb.(Lepidoptera: Noctuidae). *The Journal of Animal Ecology*, **39**, 593-598.

- 515 Norberg UM, Rayner JMV (1987) Ecological morphology and flight in bats (Mammalia;
516 Chiroptera): wing adaptations, flight performance, foraging strategy and echolocation.
517 *Philosophical Transaction of the Royal Society B Biological Sciences*, **316**, 335-427.
- 518 Oksanen JR, Kindt R, Legendre P, O'Hara B, Simpson GL, Solymos P, Stevens MHH,
519 Wagner H (2010) *Vegan: community ecology package*. Available from [http://cran.r-](http://cran.r-project.org/)
520 [project.org/](http://vegan.r-forge.r-project.org) and <http://vegan.r-forge.r-project.org>.
- 521 Oliver DR (1971) Life history of the Chironomidae. *Annual Review of Entomology*, **16**, 211–
522 230.
- 523 Oosterbroek P, de Jong H, Sijstermans L (2005) De Europese families van muggen en vliegen
524 (Diptera). Determinatie, diagnose, biologie. KNNV Uitgeverij, Utrecht.
- 525 Petersons G (2004) Seasonal migrations of north-eastern populations of Nathusius' bat
526 *Pipistrellus nathusii* (Chiroptera). *Myotis*, **41**, 29–56.
- 527 Pinder, LCV (1986) Biology of freshwater Chironomidae. *Annual Review of Entomology*, **31**,
528 1-23.
- 529 Pianka ER (1974) Niche overlap and diffuse competition. *Proceedings of the National*
530 *Academy of Sciences, USA*, **71**, 2141-2145.
- 531 Piersma T (1998) Phenotypic flexibility during migration: optimization of organ size
532 contingent on the risks and rewards of fueling and flight. *Journal of Avian Biology*, **29**,
533 511-520.
- 534 Piersma T, Gudmundsson GA, Lilliendahl K (1999) Rapid changes in the size of different
535 functional organ and muscle groups during refueling in a long-distance migrating
536 shorebird. *Physiol. Biochem. Zool.*, **72**, 405-415.
- 537 Popa-Lisseanu AG, Delgado-Huertas A, Forero MG, Rodríguez A, Arlettaz R, Ibáñez C
538 (2007) Bats' conquest of a formidable foraging niche: The myriads of nocturnally
539 migrating songbirds. *PLoS One*, **2**, e205.
- 540 Popa-Lisseanu AG, Voigt CC (2009) Bats on the move. *Journal of Mammalogy*, **90**, 1283-
541 1289.
- 542 Razgour O, Clare EL, Zeale MR, Hanmer J, Schnell IB, Rasmussen M, Jones G (2011)
543 High-throughput sequencing offers insight into mechanisms of resource partitioning in
544 cryptic bat species. *Ecology and evolution*, **1**, 556-570.
- 545 Ratnasingham S, Hebert PD (2007) BOLD: The Barcode of Life Data System ([http://www.](http://www.barcodinglife.org)
546 [barcodinglife.org](http://www.barcodinglife.org)). *Molecular Ecology Notes*, **7**, 355-364.
- 547 Richter HV, Cumming GS (2008) First application of satellite telemetry to track African
548 straw-coloured fruit bat migration. *Journal of Zoology*, **275**, 172-176.

- Robinson WD, Bowlin MS, Bisson I, Shamoun-Baranes J, Thorup K, Diehl RH, Winkler DW (2009) Integrating concepts and technologies to advance the study of bird migration. *Frontiers in Ecology and the Environment*, **8**, 354-361.
- Savage AA (1990) A key to the adults of British lesser water boatmen (Corixidae). *Field Studies*, **7**, 485-515.
- Shiel CB, McAney CM, Sullivan C, Fairley JS (1997) *Identification of arthropod fragments in bat droppings*. Occasional Publication 17, The Mammal Society, London.
- Suarez RK, Welch KC (2011) The sugar oxidation cascade: aerial refueling in hummingbirds and nectar bats. *The Journal of Experimental Biology*, **214**, 172-178.
- Šuba J, Vintulis V, Petersons G (2010) Body weight provides insights into the feeding strategy of swarming bats. *Hystrix, the Italian Journal of Mammalogy*, **22**, 179-187.
- Šuba J, Petersons G, Rydell J (2012) Fly-and-forage strategy in the bat *Pipistrellus nathusii* during autumn migration. *Acta Chiropterologica*, **14**, 379-385.
- Swift SM, Racey PA, Avery MI (1985) Feeding ecology of *Pipistrellus pipistrellus* (Chiroptera: Vespertilionidae) during pregnancy and lactation. II. Diet. *The Journal of Animal Ecology*, **54**, 217-225.
- Symondson WOC (2002) Molecular identification of prey in predator diets. *Molecular Ecology*, **11**, 627 – 641.
- Thompson SN (1973) A review and comparative characterization of the fatty acid compositions of seven insect orders. *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry*, **45**, 467-482.
- Tokeshi M (1995) Species interaction and community structure. In *The Chironomidae. Biology and ecology of non-biting midges* (P.D. Armitage, P.S. Cranston, and L.C.V. Pinder, eds.). Chapman and Hall, London.
- Tsoar A, Nathan R, Bartan Y, Vyssotski A, Dell'Omo G, Ulanovsky N (2011) Large-scale navigational map in a mammal. *Proceedings of the National Academy of Sciences*, **108**, E718-E724.
- Vaughan N (1997) The diets of British bats (Chiroptera). *Mammal Review*, **27**, 77-94.
- Valdez EW, Cryan PM (2009) Food habits of the hoary bat (*Lasiurus cinereus*) during spring migration through New Mexico. *The Southwestern Naturalist*, **54**, 195-200.
- Verkerk MC, Tramper J, van Trijp JCM, Martens DE (2007) Insect cells for human food. *Biotechnology Advances*, **25**, 198-202.

- 581 von der Emde G, Schnitzler HU (1990) Classification of insects by echolocating greater
582 horseshoe bats. *Journal of Comparative Physiology A*, **167**, 423-430.
- 583 Voigt CC, Sörgel K, Dechmann DKN (2010) Refueling while flying: Foraging bats combust
584 food rapidly and directly to power flight. *Ecology*, **91**, 2908-2917
- 585 Voigt CC, Popa-Lisseanu AG, Niermann I, Kramer-Schadt S (2012a) The catchment area of
586 wind farms for European bats: A plea for international regulations. *Biological
587 Conservation*, **153**, 80-86.
- 588 Voigt CC, Sörgel K, Suba J, Keiss O, Petersons G (2012b) The insectivorous bat *Pipistrellus
589 nathusii* uses a mixed-fuel strategy to power autumn migration. *Proceedings of the
590 Royal Society B Biological Sciences*, **279**, 3772-3778.
- 591 Westbrook JK (2008) Noctuid migration in Texas within the nocturnal aeroecological
592 boundary layer. *Integrative and Comparative biology*, **48**, 99-106.
- 593 Wikelski M, Kays RW, Kasdin NK, Thorup K, Smith JA, Swenson GW (2007) Going wild:
594 what a global small-animal tracking system could do for experimental biologists.
595 *Journal of Experimental Biology*, **210**, 181-186.
- 596 Zeale MRK, Butlin RK, Barker GLA, Lees DC, Jones G (2011) Taxon-specific PCR for DNA
597 barcoding arthropod prey in bat faeces. *Molecular Ecology Resources*, **11**, 236 – 244.
- 598
- 599

Table 1 Taxa which were assigned to MOTU utilising the BOLD search system (V.3). The confidence levels signify 1a) perfect match to one genus or species (>99%), 1b) match to one genus or species (>98%), 2) match to more than one species, of which only one was a local species, 3) match >98% to several species of different genera or to reference sequences only identified to family level. Presence and absence of prey items in the diet of bat groups is indicated by 1 and 0, respectively.

Order	Family	Species	Conf	Migration	Summer
Diptera	Asilidae	unknown	3	1	0
	Chaoboridae	unknown	3	1	0
	Chironomidae	unknown	3	1	1
		<i>Glyptotendipes sp.</i>	3	0	1
		<i>Microtendipes sp.</i>	3	1	0
		<i>Microtendipes brevitarsis</i>	1b	1	0
		<i>Parachironomus tenuicaudatu</i>	1b	1	0
		<i>Paracladopelma winnelli</i>	1b	1	1
		<i>Procladius sp.</i>	1b	1	1
		<i>Synenotendipes impar</i>	1b	0	1
		<i>Tanytarsus mendax</i>	1a	1	0
		<i>Xenochironomus xenolabis</i>	1a	1	0
		unknown	2	1	0
	Culicidae	<i>Aedes sp.</i>	2	1	0
		<i>Anopheles sp.</i>	3	0	1
		<i>Culiseta sp.</i>	2	1	0
		<i>Ochlerotatus annulipes</i>	1a	1	0
	Dolichopodidae	unknown	3	1	0
	Empididae	unknown	3	1	0
	Limoniidae	<i>Dicranomyia frontalis</i>	1a	1	1
		<i>Dicranomyia sp.</i>	1b	1	1
		<i>Erioptera sp.</i>	2	1	0
		<i>Helius flavus</i>	1a	1	0
		<i>Limonia nubeculosa</i>	1b	1	0
		<i>Metalimnobia sp.</i>	1a	1	1
		<i>Molophilus sp.</i>	1a	0	0
		<i>Phylidorea ferruginea</i>	3	1	0
		<i>Phylidorea fulvonervosus</i>	1b	0	1
		<i>Rhipidia maculata</i>	1a	1	0
		<i>Helina impunctata</i>	1a	0	1
	Muscidae	unknown	3	0	1
	Mycetophilidae	<i>Mycetophila luctuosa</i>	1a	0	1

		unknown	3	1	0
	Pedicidae	<i>Psychoda phalaenoides</i>	1a	0	1
	Psychodidae	unknown	3	1	0
	Sciaridae	<i>Anticheta sp.</i>	1b	0	1
	Sciomyzidae	<i>Haematopota pluvialis</i>	1a	0	1
	Tabanidae	<i>Hybomitra lurida</i>	1a	1	0
		<i>Nephrotoma sp.</i>	1b	1	0
	Tachinidae	unknown	3	1	0
	Tipulidae	<i>Tipula sp.</i>	1a	1	1
Lepidoptera	Amphisbatidae	<i>Pseudatemelia josephinae</i>	1a	0	1
	Argyresthiidae	<i>Argyresthiago edartella</i>	1a	0	1
	Blastobasidae	unknown	3	1	0
	Coleophoridae	<i>Coleophora glitzella</i>	1a	0	1
		<i>Coleophora limosipennella</i>	1a	1	0
	Elachistidae	<i>Agonopterix sp.</i>	1b	1	0
		<i>Semioscopis sp.</i>	1a	1	0
	Gelechiidae	<i>Exoteleia dodocella</i>	1a	1	1
		<i>Coleotechnite spiceaella</i>	1b	1	0
	Geometridae	<i>Hydriomena sp.</i>	1a	1	0
		<i>Bupalus pinaria</i>	1a	0	1
	Gracillariidae	<i>Phyllonorycter apparella</i>	1a	1	0
	Lasiocampidae	<i>Malacosoma castrensis</i>	1b	1	0
		<i>Dendrolimus pini</i>	1a	0	1
	Noctuidae	<i>Spodoptera exigua</i>	1b	1	0
	Tortricidae	<i>Acleris emargana</i>	1a	1	0
		<i>Adoxo phyesorana</i>	1a	0	1
		<i>Cnephasia sp.</i>	1a	1	1
		<i>Epinotia immunda</i>	1a	1	0
		<i>Epinotia nisella</i>	1a	1	0
		<i>Eudemis porphyra</i>	1a	0	1
		<i>Lozotaenia forsterana</i>	1a	0	1
		<i>Rhopobota naevana</i>	1a	1	0
		<i>Sparganothis sp.</i>	1b	1	0
Ephemeroptera	Baetidae	<i>Cloeon dipterum</i>	1a	1	0
	Caenidae	<i>Caenis sp.</i>	1a	1	1
	Ephemerellidae	<i>Eurylophella temporalis</i>	1b	1	1
	Ephemeridae	unknown	3	1	0

	Heptageniidae	unknown	1a	1	0
	Isonychiidae	<i>Heptagenia sp.</i>	1a	1	1
		<i>Isonychia sp.</i>	1b	1	0
Trichoptera	Leptoceridae	unknown	3	1	0
	Glossosomatidae	<i>Glossosoma intermedium</i>	1a	1	0
Neuroptera	Chrysopidae	<i>Chrysoperla sp.</i>	1a	1	0
	Hemerobiidae	<i>Hemerobius sp.</i>	1a	1	1
Hemiptera	Notonectidae	<i>Notonecta sp.</i>	1b	1	0
Coleoptera	Carabidae	<i>Agonom sp.</i>	1b	1	0
		<i>Agonom piceum</i>	1a	1	0
		<i>Dromius sp.</i>	1b	1	0
	Scirtidae	<i>Cyphon sp.</i>	1b	1	0
		<i>Cyphon phragmiteticola</i>	1a	1	0
Megaloptera	Sialidae	<i>Sialis sp.</i>	1b	1	0
Hymenoptera	Ichneumonidae	<i>Promethes sulcator</i>	1a	0	1

Table 2 Indices for richness, diversity and niche overlap for diet of bats from summer colonies and migration, derived from morphological analysis of faecal samples.

	Summer	Migration
Simpson's index	0.88	0.87
Species richness	14	13
Pianka's index	0.76	

Table 3 The frequency of prey groups in the diet of *P. nathusii* from summer colonies and migration based on morphological presence / absence data. We tested data with generalized liner model (GLM) and Tukey post-hoc test. Significant differences are indicated in bold.

	Migration	Summer	z	p<
Nematocera	73.3%	47.8%	-1.531	0.126 -
Anisopodidae	0.0%	21.7%	0.007	0.995
Chironomidae	40.0%	91.3%	3.034	0.00241
Culicidae	0.0%	17.4%	0.006	0.995
Tipulidae	86.7%	30.4%	-3.051	0.00228
Brachycera	40.0%	52.2%	0.732	0.464
Hemiptera	6.7%	0.0%	-0.003	0.998
Corixidae	0.0%	8.7%	0.004	0.997
Cicada	0.0%	4.3%	0.004	0.997
Aphidoidea	13.3%	26.1%	0.927	0.354
Trichoptera	26.7%	17.4%	-0.681	0.496
Lepidoptera	53.3%	30.4%	-1.396	0.163-
Ephemeroptera	26.7%	0.0%	-0.005	0.996
Neuroptera	46.7%	34.8%	-0.73	0.465
Coleoptera	13.3%	0.0%	-0.005	0.996
Hymenoptera	6.7%	4.3%	-0.311	0.756
Araneae	13.3%	4.3%	-0.957	0.338
Simuliidae	0.0%	4.3%	0.004	0.997
Formicidae	26.7%	0.0%	-0.005	0.996

637 Figure 1 Overview of the sampling locations for faecal samples of *Pipistrellus nathusii* (Picture) in
638 Latvia. Summer colonies were sampled in Garkalne and Vecpiebalga. Migrating bats were caught and
639 sampled in Pape, Ornithological Station, situated within the migration route of *P. nathusii* (Species
640 photo by Viesturs Vintulis).

For Review Only

641

Figure 2 Plot of a non-metric two-dimensional ordination scale (NMDS) based on the presence-absence prey data derived from the morphological diet analysis on migrating *P. nathusii* (circle) and *P. nathusii* from summer colonies (cross) (n= 50, stress = 0.20).

For Review Only

Diet of the insectivorous bat *Pipistrellus nathusii* during autumn migration and summer residence

Frauke Krüger^{*1}, Elizabeth L. Clare², William O.C. Symondson³, Oskars KeiĶš⁴, Gunārs Pētersons⁵

¹ Institute of Natural Resource Conservation, University of Kiel, Germany

² School of Biological and Chemical Sciences Queen Mary University of London, ~~United Kingdom~~

³ Cardiff School of Biosciences, Sir Martin Evans Building, Cardiff University, United Kingdom.

⁴ Institute of Biology, Laboratory of Ornithology, University of Latvia, ~~Miera Street 3, 2169~~ Salaspils, Latvia

⁵ Faculty of Veterinary Medicine, Latvia University of Agriculture, ~~K. Helmana Street 8, 3004~~ Jelgava, Latvia

Corresponding author:

Frauke Krüger, fkrueger@ecology.uni-kiel.de

Formatted: German (Germany)

Key words: ~~Chiroptera, Diet Analysis, Migration, *Pipistrellus nathusii*, Migration, Chiroptera, *Pipistrellus nathusii*, Diet Analysis~~

Running title: Diet of the migrating bat *Pipistrellus nathusii*

Abstract

Migration is widespread among vertebrates. Yet bat migration has received little attention and only in the recent decades ~~knowledge has a better understanding~~ of it has been gained. Migration can cause significant changes in behaviour and physiology, due to increasing energy demands and aerodynamic constraints. Dietary shifts, for examples, have been shown to occur in birds before onset of migration. For bats it is not known if a change in diet occurs during migration, although ~~especially~~ breeding season related dietary preference has been documented. It is known that ~~a diet rich in fats and the fat rich diets, and subsequent~~ accumulation of ~~high~~ fat deposits, do increase the flight range of migratory bats. Some bat species can be regarded as long-distance migrants, covering up to 2,000 km ~~on their way~~ between summer and winter roosting areas. *Pipistrellus nathusii* (Vespertilionidae), a European long-distant migrant, travels each year along the Baltic Sea from north-eastern

29 Europe to hibernate in central and southern Europe. This study presents data on the dietary
30 habits of migrating *Pipistrellus nathusii* compared with ~~dietary habits~~ those during the
31 breeding season. We analysed faecal samples from bats on fall migration caught at the
32 Ornithological Field Station in Pape, Latvia and from samples collected in North-Latvian
33 summer roosts. We applied both morphological identification and molecular methods, as
34 morphological methods also recognize life stages of prey and can contribute frequency data.
35 The diets of bats on migration and breeding bats were similar, with Diptera and Lepidoptera
36 comprising the major prey categories. However, certain prey groups could be explained by the
37 different hunting habitats ~~used~~ exploited during migration vs. summer residence.

40 **Introduction**

41 ~~Across the animal kingdom numerous species~~ ~~Thousands of insects, fish, birds and~~
42 ~~mammals~~ make annual movements of both short and long duration. In particular, bird
43 migration ~~between hemispheres~~ has been ~~observed~~ intensively ~~and~~ studied since the late 19th
44 century. In the last decades these studies have been sophisticated both in methods and
45 explanations ~~and still are being developed further~~ ([Berthold 2001](#); [Robinson *et al.* 2007](#);
46 ~~Cryan *et al.* 2004~~; [Wikelski *et al.* 2007](#); [Fiedler 2009](#)). Many species of bats, the only volant
47 mammals, are also known to migrate. Although the first interests in bat migration arose as
48 early as the end of the 19th century ~~by~~ ~~(Merriam (1887))~~, bat migration has been largely
49 ignored until recently. Unlike birds, the elusive life strategies of bats, difficulties regarding
50 visual observations, and low success in mark recapture programs, have made these species
51 difficult to study. However, substantial recent advances have been made, which increase our
52 understanding of orientation and physiology ([Holland *et al.* 2006](#); [Cryan & Brown 2007](#);
53 ~~Richter & Cumming 2008~~; [McGuire & Guglielmo 2009](#); [Voigt *et al.* 2010, 2012b](#)). New
54 techniques have contributed to our ability to track and infer actual range of movement, such
55 as satellite tracking and stable isotope analysis ([Cryan *et al.* 2004](#); [Richter & Cumming 2008](#);
56 [Popa-Lisseanu & Voigt 2009](#); [Voigt *et al.* 2012a](#); [Tsoar *et al.* 2012](#))

57 Studies of bat migration can profit from previous work on bird migration ([McGuire &](#)
58 [Guglielmo 2009](#)). Both birds and bats need to maintain a ~~steady-sufficient nutrient intake to~~
59 ~~meet the increased energy demand during migration state (energy in -- energy out)~~ over
60 distances of sometimes several thousand kilometers between summer and winter habitats
61 ([Griffin 1970](#); [Petersons 2004](#)). As in birds, the scale of bat migration can vary considerably
62 between short-~~distance~~, regional migrants (e.g., *Myotis daubentonii*, *M. lucifugus*) and long-
63 distance migrants (e.g., *Pipistrellus nathusii*, *Lasioncyteris noctivagans*) ([Fleming & Eby](#)
64 [2003](#); [Dzal *et al.* 2009](#); [Dzal *et al.* 2011](#)). On their journeys birds and bats face similar
65 tradeoffs between acquiring sufficient fat deposits (energy reserves) to fuel flight and

Formatted: Font: Italic

~~maintaining optimal body conditions (weight, size) optimal for aerodynamic constitution~~
~~flight with low energetic costs.~~

Birds are known to start to build up fat reserves before migratory flight and during stopovers at resting sites (e.g., Wadden Sea; McWilliams *et al.* 2004). Before the onset of migration birds show different adaptations to increase fat ~~storages stores and decrease mass:~~ they may become hyperphagous, their digestive and biosynthetic systems may alter, ~~e.g., for example~~ increase ~~in~~ liver mass and liver activity (Egeler *et al.* 2000; Guglielmo & Williams 2003), and they may increase or reduce the size of their digestive systems (Piersma 1998; Piersma *et al.* 1999; McWilliams & Karasov 2004). Additionally birds are able to fly during the night and forage and refuel during the day.

Bats have to accomplish the dual task of both flying and refueling at night. Recent studies show that bats also become hyperphagous and increase their body fat and catabolic enzyme activity during pre-migration (Ewing *et al.* 1970; Bairlain 2001; McGuire *et al.* 2009, 2013a, b; Šuba *et al.* 2010). Furthermore, they are able to fuel their migration both directly from insects caught during flight and from stored fatty acid reserves to maintain both steady state and refill reserves (Voigt *et al.* 2010; ~~Suarez & Welch 2011;~~ Voigt *et al.* 2012b). The fly-and-forage ~~strategy hypothesis, which states that bats forage on the wing during migration,~~ is ~~also~~ supported by acoustical observations along migration routes (Ahlén *et al.* 2009; Valdez & Cryan 2009, Šuba *et al.* 2012). ~~Yet, it is not for certain clear to which what extent or if at all bats are segregating foraging and migratory flight during these periods or whether they can truly hunt while commuting migrating. -~~

Another adaptation, the shift in diet towards different food items (e.g., from insects to fruit), helps some birds to gain sufficient energy during the pre-migration period (Bairlein 1990; Bairlein & Gwinner 1994; Bairlein 2001; McWilliams & Karasov 2005). It is not known if bats show similar behaviour. While most insectivorous bats use a generalist strategy, consuming prey in relation to their abundance (Anthony & Kunz 1977; Swift *et al.* 1985)

within a given habitat (Clare *et al.* 2013a/ b in review, Special Issue), selective feeding and the ability to discriminate between food items have been demonstrated in some bat species (Von der Emde & Schnitzler 1990; Koselj *et al.* 2011). Dietary shifts over time have been described in bats (Agosta 2002) and may be related to physiological state (pregnancy, lactation, preparation for hibernation) or changes in insect abundance (Clare *et al.* 2009, 2011, 2013a/b in reviewpress, Special Issue).

Here we tested the hypothesis that bat diet differs between summer roosting and fall migration. We used high throughput sequencing which yields detailed species-level data on prey in predator diets (Symondson 2002; King *et al.* 2008), and has been particularly successful in insectivores such as bats, (Razgour *et al.* 2011; Bohmann *et al.* 2011; Clare *et al.* 2013a/b in review) and shrews (Brown *et al.* in press, Special Issue). From species-level data (DNA sequences) we can draw conclusion on differences in prey items, apparent energy values or fat content and on putative foraging area differences between summer and migration habitats. We focused on a long-distance migrating bat, *Pipistrellus nathusii* (Keyserling & Blasius 1839), a generalist pipistrelle bat, which feeds to a large extent on insects connected to aquatic habitats, mainly on Diptera, particular Chironomidae (Beck 1994-1995; Vaughan 1997; Arnold *et al.* 2000; Flaquer *et al.* 2006). This species is known to travel up to 2000 km between the summer roosting grounds and hibernacula (Petersons 2004).

112 **Methods**

113 *Study sites and sample collection*

114 We collected samples for the purpose of dietary analysis at the Pape Ornithological
115 Research Station at the southern Baltic coast of Latvia (56.165° N, 21.017° E) during the fall
116 migration between 11 August and 12 September 2012 (Fig. 1). The station has been a central
117 site for intensive research on bird and bat migration, particularly during the past twenty years
118 (Petersons 2004). The surrounding areas are dominated by low sand dunes, partly covered by
119 unmanaged pine woodlands (*Pinus sylvestris*) and grasslands. In the vicinity of the station is
120 Pape Nature Park with Lake Pape, a 12km² shallow coastal lagoon lake, and a mosaic of
121 marshland, reeds, meadows, forests and peat bogs. We caught bats from dusk until 01:00am
122 using a Helgoland funnel trap following Petersons (2004). Bats were placed in a clean soft
123 cotton bags and held for approximately 1h to collect faecal samples. Samples of *P. nathusii*
124 faeces from summer colonies were collected from nursery colonies situated in buildings, at
125 Vecpiebalga (57.058° N; 25.815° E) and artificial roosts with male groups at Garkalne
126 (57.048° N; 24.382° E), Latvia during June 2013 (Fig. 1). Both sites are located in a mosaic
127 landscape of forests, pasture and in proximity to large lakes.

128 *Molecular diet analysis from faecal samples*

129 For the molecular analysis we extracted DNA from faecal samples from individual
130 migrating bats (n= 35 faecal samples) and from summer colony bulk samples, non-
131 individually collected from under the roosting bats (n = 21 faecal samples) using the QIAamp
132 DNA Stool Mini Kit (Qiagen, UK) following Zeale *et al.* (2011). To amplify the arthropod
133 prey DNA we used modified primers based on the universal COI primer ZBJ-ArtF1c and
134 ZBJ-ArtR2c. PCR (following the protocol of Zeale *et al.* 2011) produce a 157bp amplicon at
135 the 5' end of the 658bp COI barcode region (Hebert *et al.* 2004). DNA was sequenced via a
136 high throughput Ion Torrent sequencing platform (Life Technology) at the University of
137 Bristol Genomics facility (School of Biological Sciences, Bristol, UK). For the adjustment,

trimming and organisation of sequences by MIDs after sequencing we used the Galaxy V platform (<https://main.g2.bx.psu.edu/root>; Giardine *et al.* 2005; Blankenberg *et al.* 2005; Blankenberg *et al.* 2010; Goecks *et al.* 2010). To allow niche analysis procedure for all sequences, we clustered the sequences into molecular operational taxonomic units (MOTU) using the program jMOTU (Jones *et al.* 2011). We tested grouping thresholds from 1-10bp and selected a 4bp threshold as the most appropriate for this data set (see Razgour *et al.* 2011). We extracted representative sequences for each MOTU and compared sequences against references within the Barcode of Life Data System (Ratnasingham & Hebert 2007; Clare *et al.* 2009). If sequences matched completely to a reference sequence without matching any other arthropod, we regarded the sequence as belonging to the same species. However, the short amplicon length also constrains some species assignments. We used a modified version of the criteria used by Razgour *et al.* (2011) as follows:

- 1a. True species match (>99 % similarity)
 - 1b. Likely species match (>98% similarity)
 - 2. Match (>98%) to more than one species, only one of which belongs to local assemblage
 - 3. Match (>98%) to several species or genera – genus or family level assignment
- made by considered provisional.

Morphological identification from faecal samples

For morphological faecal analysis, we dried samples after DNA extraction (see Zeale *et al.* 2011-~~notes~~) at room temperature and stored them at -20°C to avoid coprophagous insects. Before analysis, we soaked the pellets for 48 h in 70% ~~Ethanol~~-ethanol and then dissected them under a binocular vision microscope (x 40 – 60). We separated characteristic fragments and mounted them in Euparal for further examination. We identification taxa to class, order, family, or genus level (where possible), by comparison of fragments with whole collected insects, arthropod identification keys from the literature ([Medvedev 1989](#); [Savage](#)

1990; Shiel *et al.* 1997; Osterbroek *et al.* 2005) and fragment photos from earlier studies (McAney *et al.* 1991; Krüger *et al.* 2012).

For every sample we calculated the frequency of each prey group relative to all samples, to estimate relative importance of prey groups (McAney-Shiel *et al.* 1991; Vaughn 1997; Krüger *et al.* 2012).

Statistical analysis

As molecular and morphological analysis produce presence-absence data and frequency data, respectively, we analysed the two data sets in different ways.

We used the Hamming distance and Bray-Curtis index (Equation 1) for similarity to analyze the dietary differences between the migratory group and the summer group as measured by molecular data. Both indices use binomial data. The Hamming distance is related to the number of changes needed to adjust two strings of same length to each other (Hamming 1950). A smaller value for Hamming distances reflects high overlap in dietary choices. The Bray-Curtis index (Bray & Curtis 1957) measures the dissimilarity between the dietary data sets, where C_{ij} is the sum of the lesser value for only those items which both data sets have in common. S_i and S_j are the total number of items counted in both data sets. If the data sets are identical, then both predators feed on the same prey and the Bray-Curtis index is 0. If the two data sets do not share any prey items then the index is 1 (Bloom 1981).

$$BC_{ij} = \frac{2C_{ij}}{S_i + S_j} \quad (\text{Equation 1})$$

$$D = 1 - \sum_{i=1}^s \frac{n_i(n_i-1)}{n(n-1)} \quad (\text{Equation 2})$$

$$O_{jk} = \frac{\sum p_{ij}p_{ik}}{(\sum p_{ij}^2 \sum p_{ik}^2)^{1/2}} \quad (\text{Equation 3})$$

188 To assess dietary niche breadth based on the morphological diet data, we used the
189 Simpson's index for diversity and heterogeneity (Equation 2), where n_i is the relative
190 proportion of a prey item i (with $i = 1 \dots n$) of a total of n prey items. Thus, D is 0, if all eaten
191 prey belongs to one prey group. The higher the diversity, the closer D gets to 1. As an
192 additional niche parameter we calculated Pianka's index of niche overlap (Equation 3), where
193 p_i is the frequency of occurrence of prey item i in the diets of species j and k (Pianka 1973):

194 We used a non-metric multi-dimensional scaling (NMDS) with Jaccard distance to
195 visualize the degree of similarity or dissimilarity of the diet. The resulting two-dimensional
196 ordination plot shows the samples sorted relative to their dissimilarity, with similar samples in
197 close proximity and dissimilar samples further apart. We used a threshold (=stress value) of <
198 0.2 for ecological interpretation of the NMDS plot (Clark & Warwick 2001).

199 We conducted indices calculation, Adonis, and NMDS using the *vegan* R library
200 (Oksanen *et al.* 2011). We applied generalized linear models (GLM) with a binomial
201 distribution and a ~~and a~~ logit link function (Zuur *et al.*, 2007) ~~general linear hypotheses (glht)~~
202 ~~with Tukey's post-hoc test~~, to assess level of significance of differences between the two data
203 sets regarding the presence or absence of prey groups, using *multcomp* R library (Hothorn *et*
204 *al.* 2008).

205

Formatted: English (U.K.)

Formatted: Font: (Default) Times New Roman, English (U.K.)

Formatted: English (U.K.)

206 Results

207 *Molecular analysis*

208 We found 220 MOTUs, of which 148 could be assigned to insect species (Table 1).
 209 For 72 MOTUs we found no matches in the BOLD System. We rejected 1 MOTU, because it
 210 contained only very short reads. Of the MOTUs 32% could be assigned to species level, 28%
 211 to genus level, 23% to family and 17% only to order level. We found 108 MOTUs in samples
 212 from migrating bats, whereas 58 MOTUs were assigned to samples from summer roosts. 19
 213 MOTUs were found in both groups.

214 Hamming Distances between migratory bats and bats at summer colonies was 197.
 215 Additionally overall Bray-Curtis similarity between migratory bats and bats at summer
 216 colonies was 0.84. Both suggest low dietary similarity.

217 *Morphological analysis*

218 We found that the diet was significantly different between bats from summer roosts
 219 and on migration, indicated by the conducted permutational analysis of variance (ADONIS: F
 220 = 4.371, df = 1, $p < 0.001$). Comparing diversity and species richness in the diet of *P. nathusii*
 221 between the two sites, we found no differences (Table 2). The trophic niche overlap, indicated
 222 by Pianka's index, was relatively high (Table 2). The ordination plot (NMDS) shows samples
 223 spread out evenly along the two dimensions, overlapping to a great extent. The slight
 224 clustering along the first dimension has to be interpreted cautiously, as a stress value of 0.2
 225 was reached (Fig. 4 2).

226 Based on GLMs we found significant difference between certain prey groups. *P.*
 227 *nathusii* from summer roosts appear to feed more often on Chironomidae than migrating *P.*
 228 *nathusii*. In contrast Tipulidae occurred more often in the diet of migrating bats (Table 3).

Formatted: Indent: First line: 0"

Discussion

The high values for Hamming distance and Bray-Curtis similarity we found for the molecular diet data indicate that diets of ~~migrating *P. nathusii* bats during migration~~ and ~~*P. nathusii* bats~~ at summer colonies show ~~high-low~~ similarity. ~~We-Despite this, we also~~ found ~~very~~ similar diversity indices for both groups, based on the morphological analysis and a relatively high niche overlap. ~~Yet, if compared to the niche overlap between different species foraging in similar habitats, for example *Myotis dasycneme* and *Myotis daubentonii*, the niche overlap between *P. nathusii* appears less strong (Krüger *et al.* 2012; Krüger *et al.* in press). In comparison to other data we found higher niche overlap between different species foraging in similar habitats, e.g. *Myotis dasycneme* and *Myotis daubentonii* (Krüger *et al.* 2012; Krüger *et al.* 2013 in review press). However, s~~Subtle but significant differences appear regarding Chironomidae and Tipulidae occurrence in the diet of migrating and summer bats, respectively. Chironomid species, especially in areas between latitude of 50° and 60°, can have two or more generations per year with diverging peaks from April to October. Several species (up to 15) can form groups which emerge in synchrony and cause an increase in potential prey biomass. As different groups follow different emergence patterns, alternating peaks of different Chironomid groups exist, replacing each other during the season and forming a more or less constant food resource (Oliver 1971; Pinder 1986; Berg and Hellenthal 1992; Tokeshi 1995; Benke 1998). The higher occurrence of Chironomidae in summer roosting bats might be explained by the reproductive state of bats and hence their needs for easily accessible prey, ~~like such as~~ swarming Chironomidae. The higher occurrence of Tipulidae in migrating bats could be ~~either~~ related to a peak in Tipulidae during that time at Pape, coinciding with migration paths. Diptera too are thought to migrate (Hogsette & Ruff 1985), and the tracks of migrating ~~*P. nathusii* bats~~ and ~~Diptera like~~ tipulids may have coincided. Alternatively the bats may have been hunting more often in terrestrial habitats during this period. (Hogsette & Ruff 1985).

A significant issue here is whether the difference is due to “migration” or “location”.

Clare *et al.* (In press) demonstrated the degree of location, season and inter-annual variation in bat diet across landscapes. Since most bats forage among prey in their immediate vicinity it is not clear to what degree shifts in diet observed here are caused by changes in insect phenology, bat physiological demands and habitat-insect relationships. We report here that there is variation between summer colonies and migrating individuals, though the underlying cause is very likely a mix of these competing influences. We suggest that the next logical step is to expand this type of study to include multiple summer and migration sites so that comparisons can be made across location.

Formatted: Font: Italic

The higher resolution of molecular diet analyses of prey species, compared contrasting with morphological analysis, provides valuable information on associations between prey, habitat and predator (Clare *et al.* 2010; Razgour *et al.* 2011, Clare *et al.* 2013a in press; Krüger *et al.* in press). While our morphological observations suggest that diet was strongly overlapping between summer and migratory groups, we did observe a higher species richness in the migratory diet based on the molecular data. There are two potential drivers of increased diversity in the diet of migrating *P. nathusii* bats. First, migrating individuals bats migrating are covering cover more space and more potential habitat types. This may expose them to a higher diversity of potential prey as a consequence. Second, insect diversity, in general is general reduced later, falls later in the summer. At this point a reduced availability of prey may force the *P. nathusii* bats to become more flexible in the prey they consume (Clare *et al.* 2013a/b in review press).

Formatted: Font: Italic

We also found that the diet of migrating *Pipistrellus* *P. nathusii* contained higher occurrence of insect species inhabiting aquatic habitats like the beetles *Cyphon phragmiteticola* and *Agonum piceum*. This probably reflects the fact that *P. nathusii* bats forage in the adjacent bog and marsh lands of Lake Pape. The moths *Epinotia immundana*, *Epinotia nisella* and *Phyllonorycter apparella* are associated with riverine forests and trees in

Formatted: Font: Italic

Formatted: English (U.S.)

Formatted: Font: Italic

marshes, supporting the inference that bats forage in the vicinity of aquatic habitats. Further indications for aquatic foraging habitats are the occurrences of Trichoptera and Megaloptera. By contrast the moth *Malacosoma castrensis* indicates foraging over dunes, as this is the major habitat of this moth. The dunes at Pape spread out parallel to the coastline, and are also used by *P. nathusii* bats as a major flight corridor during migration (Šubař et al. 2012) suggesting prey habitat and predator habitat overlap at this point.

In the diet of *P. nathusii* bats from summer colonies we found prey species which are typically associated with forested areas, like *Bupalus pinaria*, a pine pest species, or *Promethes sulcator*, an ichneumon wasp. These species were not identified in samples from migrating *P. nathusii* bats. As the colony sites are also within a few kilometres of lakes, we also found prey species associated with aquatic habitats, like Chironomidae or Ephemeroptera. Overall we can observe how the foraging habitat of *P. nathusii* determines the diet and thus differences between migrating and summering bats might be triggered explained. In birds it has been shown that during migration sedge warblers (*Acrocephalus schoenobaenus*) select for stop-over sites with high abundance of aphids (Bibby & Green 1981). Insectivorous bats, like *P. nathusii*, are known to forage particularly in habitats with high insect abundance like riverine and semi-aquatic habitats.

Formatted: Font: Italic

Many insect species are also known to migrate (e.g. ~~Hummingbird Hawk-moth~~ *Macroglossum stellatarum*, Monarch ~~butterflies~~ *butterfly* *Danaus plexippus*). The beet army worm, *Spodoptera exigua*, originally distributed in the Americas, now occurring globally, is also a known long-distance migrant (Westbrook 2008). In Europe this species has been observed to travel long distances, from Russia over Fennoscandia towards Denmark and the British Isles (Mikkola 1970). The occurrence of a migrating insect in the diets of migrating bats may be a coincidental overlap of migration routes and the opportunistic foraging behaviour of pipistrelles, which has been also observed in other species such as *Tadarida brasiliensis* which feeds opportunistically on migrating moths (Lee & McCracken 2005). ~~Also~~

for Hoary bats it is believed that it times its migration with the mass emergence of moths, its major prey (Valdez & Cryan 2009). In Europe the noctula bat, *Nyctalus lasiopterus*, has been found to a bat that predate bird, has adapted to exploit the occasional food source of migrating songbirds during spring and autumn migration (Ibáñez *et al.* 2001; Popa-Lisseanu *et al.* 2007). Similar behavior has been also reported for the case for the birdlike noctula, *Nyctalus aviator*, in Japan (Fukui *et al.* 2013).

Energetic values of insect prey might influence foraging behaviour and diet preference in migrating bats. Due to the high costs of migration flights, bats might prefer prey with high fat content and high nutritional value, to maintain energy flow and fat deposits. The variation in insect nutritional values is high, with large moths or beetles showing relatively higher fat content than many other groups (Verkerk *et al.* 2007). In addition, some migrating insect also accumulate fat to survive long dispersal flights, e.g. moths of the Noctuidae (Beall 1948; Angelo & Slansky 1984; Kevan & Kendall 1997). The occurrence of *Spodoptera exigua* (Noctuidae), and other Lepidoptera and Coleoptera in the diet of migrating *P. nathusii* suggest that these bats feed on prey with high fat content. Voigt *et al.* (2010, 2012b) proposed how that *P. nathusii* and also other bat species fuel their migration-flight during migration with endogenous fatty acids from adipocytes in combination with proteins and carbohydrates from directly metabolised from exogenous sources, such as insects. This process is determined by a limited capacity for energy storage and primarily saves energy during the costly process of converting macronutrients to lipids for storage. Nevertheless, it can be regarded as beneficial digestive adaptation to flying and hence migration. Similarly other bats are able to fuel flight with energy oxidized from both insect prey and fat deposits and are able to directly reload fat deposits with fatty acids from insect prey (Voigt *et al.* 2010). Hence, the selective exploitation of prey with high fat content would enable *P. nathusii* to balance its fat and energy reserves despite their high energy demands from long-distance and foraging flights. Birds are known to alter their dietary preference and select different food

Formatted: Font: Italic

Formatted: Font: Italic

336 ~~sources shortly before or during migration. Geese are known~~ have been shown to select for
337 ~~certain plants species during migration (Bairlein 1990). Insectivorous migratory birds like~~
338 ~~garden warbler (*Sylvia borin*) switch from arthropod based diet to fruit based diets (Bairlein~~
339 ~~1990) and furthermore select for fruit with certain fatty acid compositions prior to migration~~
340 ~~(McWilliams *et al.* 2004). Other birds like willow warblers switch to insect prey high in~~
341 ~~sugars, like aphids (Berthold 2001). Adaptive alteration of diet selectivity during migration~~
342 ~~seems to be a valuable trait in migrants. The latter is comparable to possible shifts in diets of~~
343 ~~migratory bats.~~

Formatted: Font: Italic

344 This kind of selectivity would require the bats to discriminate between prey of
345 different energy ~~fat~~ values. In ~~bats selective behaviour and prey discrimination based on size~~
346 ~~has previously been demonstrated only for~~ horseshoe bats ~~selective behaviour and prey~~
347 ~~discrimination based on size has been demonstrated previously~~ (Koselj *et al.* 2011). Whether
348 the same ability exists for other bats, like *Pipistrellus*, which ~~in contrast to horseshoe bats,~~
349 use short frequency-modulated (FM) calls and mainly feed on Diptera ~~on the wing,~~ is not
350 clear. ~~Also like for frugivory in insectivorous birds during migration, the differences in diet~~
351 ~~between summer and migratory *P. nathusii* may result from the seasonal changes in~~
352 ~~availability of certain food items, insects and fruit, respectively.~~

Formatted: Font: Italic

Formatted: English (U.S.)

353 The fat stores of migrating bats have higher proportions of polyunsaturated fatty acids
354 (PUFAs) (McGuire *et al.* 2013b). Thus PUFAs ~~are seem to may be~~ an important resource
355 during pre-migration and migration itself. Naturally, the diet of *Pipistrellus nathusii* is often
356 dominated by Diptera, particularly Chironomidae, which are rich in highly unsaturated fatty
357 acids (Thompson 1982; Hanson *et al.* 1985). ~~Overall aquatic insects have higher PUFA~~
358 ~~content than do terrestrial insects, though this varies depending on life stage (Hanson *et al.*~~
359 ~~1985). The high occurrence of Chironomidae in the summer diet of *P. nathusii* demonstrates~~
360 ~~that bats already have good supply of fat resources, needed for building up reserves.~~
361 ~~Migrating bats still feed to large extent on Nematocera, often associated with high PUFA~~

362 | ~~content~~. Thus increased lipid biosynthesis capacity and additional intake of bigger, or fatter
363 | prey, may not be required during migration (McGuire & Guglielmo 2009).

364 | In general, our results demonstrate **differences** in the diet of *P. nathusii* from summer
365 | roosts and *P. nathusii* on migration. *P. nathusii* is a generalist predator and feeds on prey
366 | groups thought to be rich in important fatty acids (e.g. Chironomidae) thus the need to select
367 | for prey with particularly high fat content during migration might be low. Yet, there is no
368 | evidence that endogenous triggered selectivity can be observed in insectivorous bats like it is
369 | the case for some insectivorous and grazing birds.

370 | Additionally, ~~their~~ the ability of pipistrelle bats to discriminate between prey of
371 | differing energetic values might be poor and hamper shifts in prey selection. Diet of migrating
372 | bats like *P. nathusii* might rather depend on the availability of prey at the respective stop-over
373 | site and the differences between migrating and summering individuals found in prey groups
374 | ~~can~~ are likely to be related to habitat differences along migratory routes and in the summering
375 | grounds. For the future it would be beneficial to find and add more migratory stop-over sites,
376 | where species can be studied.

Formatted: Normal

Formatted

Acknowledgments

We are grateful to the Baltic-German University Liaison Office / DAAD for financial support. Many thanks to Alma Vītola, Ilze Brila, Jurgis Šuba for help in the field. We thank the two anonymous reviewers for valuable comments on an earlier version of this manuscript.

Data Accessibility

Molecular (sequences) and morphological (binary presence-absence) dietary data will be provided on a DRYAD account (doi: 10.5061/dryad.2d38f).

References

Agosta SJ (2002) Habitat use, diet and roost selection by the Big Brown Bat (*Eptesicus fuscus*) in North America: a case for conserving an abundant species. *Mammal Review* **32**, 179-198.

Ahlén I, Baagø H, Bach L (2009) Behaviour of Scandinavian bats during migration and foraging at sea. *Journal of Mammalogy*, **90**, 1318-1323.

Angelo MJ, Slansky Jr F (1984) Body building by insects: trade-offs in resource allocation with particular reference to migratory species. *Florida Entomologist*, 22-41

Anthony, E.L.P., Kunz, T. H., 1977. Feeding Strategies of the Little Brown Bat, *Myotis lucifugus*, in Southern New Hampshire. *Ecology*, **58**, 775-786.

Bairlein F (1990) Nutrition and food selection in migratory birds. In: *Bird migration: physiology and ecophysiology* (ed Gwinner E), pp 198–213. Springer, Berlin Heidelberg New York.

Bairlein F (2001). Results of bird ringing in the study of migration routes and behaviour. *Ardea*, **89**, 7-19.

Bairlein F, Gwinner E (1994) Nutritional mechanisms and temporal control of migratory energy accumulation in birds. *Annual Review of Nutrition*, **14**, 187-215

Beall F (1948) The fat content of a butterfly, *Danaus plexippus* Linn., as affected by migration. *Ecology*, **29**, 80-94.

Beck A (1994-95). Fecal analyses of European bat species. *Myotis*, **32-33**, 109-119.

- 408 [Benke AC \(1998\) Production Dynamics of Riverine Chironomids: Extremely High Biomass](#)
 409 [Turnover Rates of Primary Consumers. *Ecology*, **79**, 899-910.](#)
- 410 ~~Berg, M-B, and R-A. Hellenthal-RA (1992). The role of Chironomidae in energy flow of a~~
 411 ~~lotic ecosystem. *Netherlands Journal of Aquatic Ecology*, **26**, 471-476.~~
- 412 ~~[Benke, A.C. 1998. Production Dynamics of Riverine Chironomids: Extremely High Biomass](#)~~
 413 ~~[Turnover Rates of Primary Consumers. *Ecology*, **79**, 899-910.](#)~~
- 414 [Berthold P \(2001\). *Bird migration: a general survey*. Oxford University Press.](#)
- 415 Blankenberg D, Taylor J, Schenck I, He J, Zhang Y, Ghent M, Veeraraghavan N, Albert I,
 416 Miller W, Makova KD, Hardison RC, Nekrutenko A (2007) A framework for
 417 collaborative analysis of ENCODE data: Making large-scale analyses biologist-
 418 friendly. *Genome Research*, **17**, 960-964.
- 419 Blankenberg D, Gordon A, Von Kuster G, Coraor N, Taylor J, Nekrutenko A, The Galaxy
 420 Team (2010) Manipulation of FASTQ data with Galaxy. *Bioinformatics Application*
 421 *Note*, **26**, 1783-1785.
- 422 Bloom SA (1982) Similarity indices in community studies: Potential pitfalls. *Marine Ecology*
 423 *– Progress Series*, **5**, 125-128.
- 424 Bohmann K, Monadjem A, LehmkuhlNoer C, Rasmussen M, Zeale MRK, Clare EL, Jones G,
 425 Willerslev E, Gilbert MTP (2011) Molecular Diet Analysis of Two African Free-
 426 Tailed Bats (Molossidae) Using High Throughput Sequencing. *PLOS one*, **6**, 1-11.
- 427 Brown DS, Burger R, Cole N, Vencatasamy D, Clare EL, Montazam A, Symondson WOC (in
 428 press) Dietary competition between the alien Asian Musk Shrew (*Suncus murinus*) and
 429 a reintroduced population of Telfair's Skink (*Leiopisma telfairii*). *Molecular*
 430 *Ecology*.
- 431 ~~Clare EL, Fraser EE, Braid HE, Fenton MB, Hebert PD (2009) Species on the menu of a~~
 432 ~~generalist predator, the eastern red bat (*Lasiurus borealis*): using a molecular approach~~
 433 ~~to detect arthropod prey. *Molecular Ecology*, **18**, 2532-2542.~~
- 434 Clare EL, Barber BR, Sweeney BW, Hebert PDN, Fenton MB (2011) Eating local: influences
 435 of habitat on the diet of little brown bats (*Myotis lucifugus*). *Molecular Ecology*, **20**,
 436 1772-1780.
- 437 [Clare EL, Symondson WOC, Broders H et al. \(THIS ISSUEa\) The diet of *Myotis lucifugus*](#)
 438 [across Canada: assessing foraging quality and diet variability. *Molecular Ecology*.](#)
 439 [\(XXXX-XXXX\)](#)

Formatted: English (U.S.), Not Small caps

Formatted: English (U.S.)

Formatted: Font: Italic, English (U.S.)

Formatted: English (U.S.)

Formatted: Font: Bold, English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.S.), Not Small caps

Formatted: English (U.S.)

Formatted: Font: Italic, English (U.S.)

Formatted: English (U.S.)

Formatted: Font: Bold, English (U.S.)

Formatted: English (U.S.)

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Auto, English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.K.)

Formatted: English (U.K.)

Formatted: Normal, Indent: Left: 0", Hanging: 0.49", Keep with next

440 | [Clare EL, Symondson WOC, Fenton MB \(THIS ISSUEb\) An inordinate fondness for beetles?](#)
441 | [Variation in seasonal dietary preferences of night roosting big brown bats \(*Eptesicus*](#)
442 | [fuscus\). *Molecular Ecology*. \(XXXX-XXXX\)](#) Formatted: English (U.K.)

443 | Clarke KR, Warwick RM (2001) Change in marine communities: an approach to statistical
444 | analysis and interpretation, 2nd ed. PRIMER-E, Plymouth. Formatted: English (U.S.)

445 | Cryan PM, Bogan MA, Rye RO, Landis GP, Kester CL (2004) Stable hydrogen isotope
446 | analysis of bat hair as evidence for seasonal molt and long-distance migration. *Journal*
447 | *of Mammalogy*, **85**, 995-1001.

448 | Cryan PM, Brown AC (2007) Migration of bats past a remote island offers clues toward the
449 | problem of bat fatalities at wind turbines. *Biological Conservation*, **139**, 1-11. Formatted

450 | Dzal Y, Hooton LA, Clare EL, Fenton MB (2009). Bat activity and genetic diversity at Long
451 | Point, Ontario, an important bird stopover site. *Acta Chiropterologica*, **11**, 307-315. Formatted
Formatted

452 | Dzal Y, McGuire LP, Veselka N, Fenton MB (2011). Going, going, gone: the impact of
453 | white-nose syndrome on the summer activity of the little brown bat (*Myotis lucifugus*).
454 | *Biology letters*, **7**, 392-394. Formatted: Spanish (International Sort)

455 | Egeler O, Williams TD, Guglielmo CG (2000) Modulation of lipogenic enzymes, fatty acid
456 | synthase and $\Delta 9$ -desaturase, in relation to migration in the western sandpiper (*Calidris*
457 | *mauri*). *Journal of Comparative Physiology B*, **170**, 169-174

458 | Ewing WG, Studier EH, O'Farrell MJ (1970) Autumn fat deposition and gross body
459 | composition in three species of *Myotis*. *Comparative Biochemistry and Physiology*,
460 | **36**, 119-129.

461 | [Fiedler W \(2009\) New technologies for monitoring bird migration and behaviour. *Ringling &*](#)
462 | [Migration. **24**, 175-179.](#) Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Auto, English (U.S.)
Formatted: English (U.S.)

463 | Fleming TH, Eby P (2003) Ecology of bat migration. In: *Bat Ecology* (eds Kunz TH &Fenton
464 | MB), pp. 156-208. University of Chicago Press, Chicago, USA. Formatted: Font: Bold, Not Italic, English (U.S.)
Formatted: English (U.S.)

465 | [Fukui D, Dewa H, Katsuta S, Sato A \(2013\) Bird predation by the birdlike noctula in Japan.](#)
466 | [Journal of Mammalogy. **94**, 657-661.](#) Formatted: Font: Italic
Formatted: Font: Bold

467 | Giardine B, Riemer C, Ross CH *et al.* (2005) Galaxy: A platform for interactive large-scale
468 | genome analysis. *Genome Research*, **15**, 1451-1455.

469 | Goecks J, Nekrutenko A, Taylor J, The Galax Team (2010) Galaxy: a comprehensive
470 | approach for supporting accessible, reproducible, and transparent computational
471 | research in the life sciences. *Genome Biology*, doi:10.1186. Formatted: English (U.S.)

- Griffin DR(1970) Griffin, D. R. (1970). Migration and homing of bats. In: *Biology of bats*, Vol 2, (ed Wimsatt WA), pp. 233-264. Academic Press, New York.
- Guglielmo CG, Williams TD (2003) Phenotypic flexibility of body composition in relation to migratory state, age, and sex in the western sandpiper (*Calidris mauri*). *Physiological and Biochemical Zoology*, **76**, 84-98.
- Hamming RW (1950) Error detecting and error correcting codes. *The Bell System Technical Journal*, **29**, 147-160.
- Hanson BJ, Cummins KW, Cargill AS, Lowry RR (1985) Lipid content, fatty acid composition, and the effect of diet on fats of aquatic insects. *Comparative Biochemistry and Physiology Part B: Comparative Biochemistry*, **80**, 257-276.
- Hebert PDN, Stoeckle MY, Zemlak TS, Francis CM (2004) Identification of birds through DNA barcodes. *PLoS Biology*, **2**, e312.
- Hogsette JA, Ruff JP (1985) Stable fly (Diptera: Muscidae) migration in northwest Florida. *Environmental entomology*, **14**, 170-175.
- Holland RA, Thorup K, Vonhof MJ, Cochran WW, Wikleski M (2006) Bat orientation using Earth's magnetic fields. *Nature*, **44**, 702.
- Holland RA, Borissov I, Siemers BM (2010) A nocturnal mammal, the greater mouse-eared bat, calibrates a magnetic compass by the sun. *Proceedings of the National Academy of Sciences*, **107**, 6941-6945.
- Hothorn T, Bretz F, Westfall P (2008) Simultaneous Inference in General Parametric Models. *Biometrical Journal*, **50**, 346 - 363.
- Jones M, Ghoorah A, Blaxter M (2011) jMOTU and Taxonator: turning DNA Barcode sequences into annotated operational taxonomic units. *PLoS one*, **6**, e19259.
- Kevan PG, Kendall DM (1997) Liquid assets for fat bankers: Summer nectarivory by migratory moths in the Rocky Mountains, Colorado, U.S.A. *Arctic and Alpine Research*, **29**, 478-482.
- Ibáñez C, Juste J, García-Mudarra JL, Agirre-Mendi PT (2001) Bat predation on nocturnally migrating birds. *Proceedings of the National Academy of Sciences*, **98**, 9700-9702.
- King RA, Read DS, Traugott M, Symondson WOC (2008) Molecular analysis of predation: a review of best practice for DNA-based approaches. *Molecular Ecology*, **17**, 947 - 963.
- Koselj K, Schnitzler H-U, Siemers BM (2011) Horseshoe bats make adaptive prey-selection decisions, informed by echo cues. *Proceedings of the Royal Society B*, **278**, 3034-3041.

Formatted: German (Germany)

505 Krüger F, Harms I, Fichtner A, Wolz I, Sommer RS (2013) High trophic similarity in the
506 sympatric North European trawling bat species *Myotis daubentonii* and *Myotis*
507 *dasycneme*. *Acta Chiropterologica*, **14**, 347–356.

508 [Krüger F, Clare EL, Greif S, Siemers BM, Symondson WOC, Sommer RS \(THIS ISSUE\) An](#)
509 [integrative approach to detect subtle trophic niche differentiation in the sympatric](#)
510 [trawling bat species *Myotis dasycneme* and *Myotis daubentonii*. *Molecular Ecology*,](#)
511 [\(XXXX-XXXX\)](#)

512 Lee Y, McCracken GF (2009) Dietary variation of Brazilian free-tailed bats links to migratory
513 populations of pest insects. *Journal of Mammalogy*, **86**, 67–76.

514 McGuire LP, Fenton MB, Guglielmo CG (2009) Effect of age on energy storage during
515 prehibernation swarming in little brown bats (*Myotis lucifugus*). *Canadian Journal of*
516 *Zoology*, **87**, 515–519.

517 McGuire L, Guglielmo CG (2009) What can birds tell us about the migration physiology of
518 bats? *Journal of Mammalogy*, **90**, 1290–129.

519 McGuire LP, Fenton MB, Guglielmo CG (2013a). Phenotypic flexibility in migrating bats:
520 seasonal variation in body composition, organ sizes and fatty acid profiles. *The*
521 *Journal of experimental biology*, **216**, 800–808.

522 McGuire LP, Fenton MB, Guglielmo CG (2013b) Seasonal upregulation of catabolic enzymes
523 and fatty acid transporters in the flight muscle of migrating hoary bats,
524 *Lasiurus cinereus*. *Comparative Biochemistry and Physiology B*, **165**, 138–143.

525 McWilliams SR, Guglielmo C, Pierce B, Klaassen M (2004) Flying, fasting and feeding in
526 birds during migration: a nutritional and physiological ecology perspective. *Journal of*
527 *Avian Biology*, **35**, 377–393.

528 McWilliams SR, Karasov WH (2004) Migration takes guts: digestive physiology of migratory
529 birds and its ecological significance. In: *Birds of Two Worlds* (eds Marra P Greenberg
530 R). Johns Hopkins University Press, Baltimore, M. D.

531 ~~Medvedev G-S (Ed.) (1989) Keys to the Insects of the European Part of the USSR:~~
532 ~~Lepidoptera. IV, 1 (Vol. 117). Brill Archive.~~

533 Merriam CH (1887) Do any Canadian bats migrate? Evidence in the affirmative. *Transaction*
534 *of the Royal Society of Canada*, **5**, 85–87.

535 Mikkola K (1970) The interpretation of long-range migrations of *Spodoptera exigua*
536 Hb.(Lepidoptera: Noctuidae). *The Journal of Animal Ecology*, **39**, 593–598.

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.K.)

Formatted: English (U.K.)

Formatted: English (U.S.)

- 537 Norberg UM, Rayner JMV (1987) Ecological morphology and flight in bats (Mammalia;
538 Chiroptera): wing adaptations, flight performance, foraging strategy and echolocation.
539 *Philosophical Transaction of the Royal Society B Biological Sciences*, **316**, 335-427.
- 540 Oksanen JR, Kindt R, Legendre P, O'Hara B, Simpson GL, Solymos P, Stevens MHH,
541 Wagner H (2010) *Vegan: community ecology package*. Available from [http://cran.r-](http://cran.r-project.org/)
542 [project.org/](http://cran.r-project.org/) and <http://vegan.r-forge.r-project.org>.
- 543 Oliver D-R (1971) Life history of the Chironomidae. *Annual Review of Entomology*, **16**,
544 211-230.
- 545 Oosterbroek P, de Jong H, Sijstermans L (2005) *De Europese families van muggen en vliegen*
546 *(Diptera). Determinatie, diagnose, biologie*. KNNV Uitgeverij, Utrecht.
- 547 Petersons G (2004) Seasonal migrations of north-eastern populations of Nathusius' bat
548 *Pipistrellus nathusii* (Chiroptera). *Myotis*, **41**, 29-56.
- 549 Pinder LCV (1986) Biology of freshwater Chironomidae. *Annual Review of Entomology*, **31**,
550 1-23.
- 551 Pianka ER (1974) Niche overlap and diffuse competition. *Proceedings of the National*
552 *Academy of Sciences, USA*, **71**, 2141-2145.
- 553 Piersma T (1998) Phenotypic flexibility during migration: optimization of organ size
554 contingent on the risks and rewards of fueling and flight. *Journal of Avian Biology*, **29**,
555 511-520.
- 556 Piersma T, Gudmundsson GA, Lilliendahl K (1999) Rapid changes in the size of different
557 functional organ and muscle groups during refueling in a long-distance migrating
558 shorebird. *Physiol. Biochem. Zool.*, **72**, 405-415.
- 559 Popa-Lisseanu AG, Delgado-Huertas A, Forero MG, Rodríguez A, Arlettaz R, Ibáñez C
560 (2007) Bats' conquest of a formidable foraging niche: The myriads of nocturnally
561 migrating songbirds. *PLoS One*, **2**, e205.
- 562 Popa-Lisseanu AG, Voigt CC (2009) Bats on the move. *Journal of Mammalogy*, **90**, 1283-
563 1289.
- 564 Razgour O, Clare EL, Zeale MR, Hanmer J, Schnell IB, Rasmussen M, Jones G (2011)
565 High-throughput sequencing offers insight into mechanisms of resource partitioning in
566 cryptic bat species. *Ecology and evolution*, **1**, 556-570.
- 567 Ratnasingham S, Hebert PD (2007) BOLD: The Barcode of Life Data System ([http://www.](http://www.barcodinglife.org)
568 [barcodinglife.org](http://www.barcodinglife.org)). *Molecular Ecology Notes*, **7**, 355-364.
- 569 Richter HV, Cumming GS (2008) First application of satellite telemetry to track African
570 straw-coloured fruit bat migration. *Journal of Zoology*, **275**, 172-176.

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: Not Small caps

Formatted: Font: Italic

Formatted: Font: Bold

Formatted: English (U.S.)

Formatted: Not Small caps

Formatted: Not Small caps

Formatted: Font: Italic, English (U.S.)

Formatted: Font: Bold, English (U.S.)

Formatted: English (U.S.)

571 [Robinson WD, Bowlin MS, Bisson I, Shamoun-Baranes J, Thorup K, Diehl RH, Winkler DW](#)
572 [\(2009\) Integrating concepts and technologies to advance the study of bird migration.](#)
573 [Frontiers in Ecology and the Environment, 8, 354-361.](#)
574 [Savage AA \(1990\) A key to the adults of British lesser water boatmen \(Corixidae\). Field](#)
575 [Studies, 7, 485-515.](#)
576 Shiel CB, McAney CM, Sullivan C, Fairley JS (1997) *Identification of arthropod fragments*
577 *in bat droppings*. Occasional Publication 17, The Mammal Society, London.
578 Suarez RK, Welch KC (2011). The sugar oxidation cascade: aerial refueling in hummingbirds
579 and nectar bats. *The Journal of Experimental Biology*, **214**, 172-178.
580 Šuba J, Vintulis V, Petersons G (2010) Body weight provides insights into the feeding
581 strategy of swarming bats. *Hystrix, the Italian Journal of Mammalogy*, **22**, 179-187.
582 Šuba J, Petersons G, Rydell J (2012) Fly-and-forage strategy in the bat *Pipistrellus nathusii*
583 during autumn migration. *Acta Chiropterologica*, **14**, 379-385.
584 Swift SM, Racey PA, Avery MI (1985) Feeding ecology of *Pipistrellus pipistrellus*
585 (Chiroptera: Vespertilionidae) during pregnancy and lactation. II. Diet. *The Journal of*
586 *Animal Ecology*, **54**, 217-225.
587 Symondson WOC (2002) Molecular identification of prey in predator diets. *Molecular*
588 *Ecology*, **11**, 627 – 641.
589 Thompson SN (1973) A review and comparative characterization of the fatty acid
590 compositions of seven insect orders. *Comparative Biochemistry and Physiology Part*
591 *B: Comparative Biochemistry*, **45**, 467-482
592 Tokeshi M (1995) Species interaction and community structure. In *The Chironomidae.*
593 *Biology and ecology of non-biting midges* (P.D. Armitage, P.S. Cranston, and L.C.V.
594 Pinder, eds.). Chapman and Hall, London.
595 Tsoar A, Nathan R, Bartan Y, Vyssotski A, Dell'Omo G, Ulanovsky N (2011) Large-scale
596 navigational map in a mammal. *Proceedings of the National Academy of Sciences*, **108**,
597 E718-E724.
598 [Vaughan N \(1997\) The diets of British bats \(Chiroptera\). Mammal Review, 27, 77-94.](#)
599 Valdez EW, Cryan PM (2009) Food habits of the hoary bat (*Lasiurus cinereus*) during spring
600 migration through New Mexico. *The Southwestern Naturalist*, **54**, 195-200.
601 Verkerk MC, Tramper J, van Trijp JCM, Martens DE (2007) Insect cells for human food.
602 *Biotechnology Advances*, **25**, 198-202.

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Auto, English (U.S.)

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Auto, English (U.S.)

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Auto, English (U.S.)

Formatted: English (U.S.)

Formatted: Font: Bold, Not Italic, English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: Not Small caps

Formatted: Heading 7, References, Indent: Left: 0", First line: 0", Line spacing: single, Tab stops: Not at 0.1" + 0.49"

Formatted: English (U.S.)

Formatted: Font: Italic, English (U.S.)

Formatted: English (U.S.)

Formatted: Font: Bold, English (U.S.)

Formatted: English (U.S.)

Formatted: Font: Italic

- 603 von der Emde G, Schnitzler HU (1990) Classification of insects by echolocating greater
604 horseshoe bats. *Journal of Comparative Physiology A*, **167**, 423-430.
- 605 Voigt CC, Sörgel K, Dechmann DKN (2010) Refueling while flying: Foraging bats combust
606 food rapidly and directly to power flight. *Ecology*, **91**, 2908-2917
- 607 Voigt CC, Popa-Lisseanu AG, Niermann I, Kramer-Schadt S (2012a) The catchment area of
608 wind farms for European bats: A plea for international regulations. *Biological*
609 *Conservation*, **153**, 80-86.
- 610 Voigt CC, Sörgel K, Suba J, Keiss O, Petersons G (2012b) The insectivorous bat *Pipistrellus*
611 *nathusii* uses a mixed-fuel strategy to power autumn migration. *Proceedings of the*
612 *Royal Society B Biological Sciences*, **279**, 3772-3778.
- 613 Westbrook JK (2008) Noctuid migration in Texas within the nocturnal aeroecological
614 boundary layer. *Integrative and Comparative biology*, **48**, 99-106.
- 615 Wikelski M, Kays RW, Kasdin NK, Thorup K, Smith JA, Swenson GW (2007) Going wild:
616 what a global small-animal tracking system could do for experimental biologists.
617 *Journal of Experimental Biology*, **210**, 181-186.
- 618 Zeale MRK, Butlin RK, Barker GLA, Lees DC, Jones G (2011) Taxon-specific PCR for DNA
619 barcoding arthropod prey in bat faeces. *Molecular Ecology Resources*, **11**, 236 – 244.
- 620
- 621 |

622 Table 1 Taxa which were assigned to MOTU utilising the BOLD search system (V.3). The confidence
623 levels signify 1a) perfect match to one genus or species (>99%), 1b) match to one genus or species
624 (>98%), 2) match to more than one species, of which only one was a local species, 3) match >98% to
625 several species of different genera or to reference sequences only identified to family level. Presence
626 and absence of prey items in the diet of bat groups is indicated by 1 and 0, respectively.

Order	Family	Species	Conf	Migration	Summer
Diptera	Asilidae	unknown	3	1	0
	Chaoboridae	unknown	3	1	0
	Chironomidae	unknown	3	1	1
		<i>Glyptotendipes</i> sp.	3	0	1
		<i>Microtendipes</i> sp.	3	1	0
		<i>Microtendipes brevitarsis</i>	1b	1	0
		<i>Parachironomus tenuicaudatus</i>	1b	1	0
		<i>Paracladopelma winnelli</i>	1b	1	1
		<i>Procladius</i> sp.	1b	1	1
		<i>Synenotendipes impar</i>	1b	0	1
		<i>Tanytarsus mendax</i>	1a	1	0
		<i>Xenochironomus xenolabis</i>	1a	1	0
		unknown	2	1	0
	Culicidae	<i>Aedes</i> sp.	2	1	0
		<i>Anopheles</i> sp.	3	0	1
		<i>Culiseta</i> sp.	2	1	0
		<i>Ochlerotatus annulipes</i>	1a	1	0
	Dolichopodidae	unknown	3	1	0
	Empididae	unknown	3	1	0
	Limoniidae	<i>Dicranomyia frontalis</i>	1a	1	1
		<i>Dicranomyia</i> sp.	1b	1	1
		<i>Erioptera</i> sp.	2	1	0
		<i>Helius flavus</i>	1a	1	0
		<i>Limonia nubeculosa</i>	1b	1	0
		<i>Metalimnobia</i> sp.	1a	1	1
		<i>Molophilus</i> sp.	1a	0	0
		<i>Phylidorea ferruginea</i>	3	1	0
		<i>Phylidorea fulvonervosus</i>	1b	0	1
		<i>Rhipidia maculata</i>	1a	1	0
		<i>Helina impunctata</i>	1a	0	1
	Muscidae	unknown	3	0	1
	Mycetophilidae	<i>Mycetophila luctuosa</i>	1a	0	1

Formatted	... [28]
Formatted	... [29]
Formatted	... [30]
Formatted	... [31]
Formatted	... [32]
Formatted	... [33]
Formatted	... [34]
Formatted	... [35]
Formatted	... [36]
Formatted	... [37]
Formatted	... [38]
Formatted	... [39]
Formatted	... [1]
Formatted	... [40]
Formatted	... [41]
Formatted	... [42]
Formatted	... [43]
Formatted	... [44]
Formatted	... [45]
Formatted	... [46]
Formatted	... [47]
Formatted	... [48]
Formatted	... [49]
Formatted	... [50]
Formatted	... [51]
Formatted	... [52]
Formatted	... [53]
Formatted	... [54]
Formatted	... [55]
Formatted	... [56]
Formatted	... [57]
Formatted	... [58]
Formatted	... [59]
Formatted	... [60]
Formatted	... [61]
Formatted	... [62]
Formatted	... [63]
Formatted	... [64]
Formatted	... [65]
Formatted	... [2]
Formatted	... [3]
Formatted	... [66]
Formatted	... [67]
Formatted	... [4]
Formatted	... [5]
Formatted	... [68]
Formatted	... [69]
Formatted	... [6]
Formatted	... [7]
Formatted	... [70]
Formatted	... [71]
Formatted	... [8]
Formatted	... [9]
Formatted	... [10]
Formatted	... [11]
Formatted	... [12]
Formatted	... [13]
Formatted	... [14]
Formatted	... [15]

		unknown	3	1	0	Formatted	... [72]
	Pediciidae	<i>Psychoda phalaenoides</i>	1a	0	1	Formatted	... [73]
	Psychodidae	unknown	3	1	0	Formatted	... [74]
	Sciaridae	<i>Anticheta sp.</i>	1b	0	1	Formatted	... [75]
	Sciomyzidae	<i>Haematopota pluvialis</i>	1a	0	1	Formatted	... [76]
	Tabanidae	<i>Hybomitra lurida</i>	1a	1	0	Formatted	... [77]
		<i>Nephrotoma sp.</i>	1b	1	0	Formatted	... [78]
	Tachinidae	unknown	3	1	0	Formatted	... [79]
	Tipulidae	<i>Tipula sp.</i>	1a	1	1	Formatted	... [80]
Lepidoptera	Amphisbatidae	<i>Pseudatemelia josephinae</i>	1a	0	1	Formatted	... [119]
	Argyresthiidae	<i>Argyresthiago edartella</i>	1a	0	1	Formatted	... [120]
	Blastobasidae	unknown	3	1	0	Formatted	... [121]
	Coleophoridae	<i>Coleophora glitzella</i>	1a	0	1	Formatted	... [122]
		<i>Coleophora limosipennella</i>	1a	1	0	Formatted	... [123]
	Elachistidae	<i>Agonopterix sp.</i>	1b	1	0	Formatted	... [124]
		<i>Semioscopis sp.</i>	1a	1	0	Formatted	... [125]
	Gelechiidae	<i>Exoteleia dodocella</i>	1a	1	1	Formatted	... [126]
		<i>Coleotechnite spiceaella</i>	1b	1	0	Formatted	... [81]
	Geometridae	<i>Hydriomena sp.</i>	1a	1	0	Formatted	... [82]
		<i>Bupalus pinaria</i>	1a	0	1	Formatted	... [83]
	Gracillariidae	<i>Phyllonorycter apparella</i>	1a	1	0	Formatted	... [84]
	Lasiocampidae	<i>Malacosoma castrensis</i>	1b	1	0	Formatted	... [129]
		<i>Dendrolimus pini</i>	1a	0	1	Formatted	... [130]
	Noctuidae	<i>Spodoptera exigua</i>	1b	1	0	Formatted	... [85]
	Tortricidae	<i>Acleris emargana</i>	1a	1	0	Formatted	... [86]
		<i>Adoxo phyesorana</i>	1a	0	1	Formatted	... [131]
		<i>Cnephasia sp.</i>	1a	1	1	Formatted	... [132]
		<i>Epinotia immunda</i>	1a	1	0	Formatted	... [87]
		<i>Epinotia nisella</i>	1a	1	0	Formatted	... [88]
		<i>Eudemis porphyra</i>	1a	0	1	Formatted	... [133]
		<i>Lozotaenia forsterana</i>	1a	0	1	Formatted	... [93]
		<i>Rhopobota naevana</i>	1a	1	0	Formatted	... [94]
		<i>Sparganothis sp.</i>	1b	1	0	Formatted	... [91]
						Formatted	... [92]
						Formatted	... [137]
						Formatted	... [138]
Ephemeroptera	Baetidae	<i>Cloeon dipterum</i>	1a	1	0	Formatted	... [93]
	Caenidae	<i>Caenis sp.</i>	1a	1	1	Formatted	... [94]
	Ephemerellidae	<i>Eurylophella temporalis</i>	1b	1	1	Formatted	... [95]
	Ephemeridae	unknown	3	1	0	Formatted	... [96]
						Formatted	... [97]
						Formatted	... [98]
						Formatted	... [99]
						Formatted	... [100]
						Formatted	... [101]
						Formatted	... [102]
						Formatted	... [103]
						Formatted	... [104]
						Formatted	... [105]
						Formatted	... [106]
						Formatted	... [107]
						Formatted	... [108]
						Formatted	... [109]
						Formatted	... [110]
						Formatted	... [111]
						Formatted	... [112]
						Formatted	... [113]
						Formatted	... [114]
						Formatted	... [115]
						Formatted	... [116]
						Formatted	... [117]
						Formatted	... [118]
						Formatted	... [119]
						Formatted	... [120]
						Formatted	... [121]
						Formatted	... [122]
						Formatted	... [123]
						Formatted	... [124]
						Formatted	... [125]
						Formatted	... [126]
						Formatted	... [127]
						Formatted	... [128]
						Formatted	... [129]
						Formatted	... [130]
						Formatted	... [131]
						Formatted	... [132]
						Formatted	... [133]
						Formatted	... [134]
						Formatted	... [135]
						Formatted	... [136]
						Formatted	... [137]
						Formatted	... [138]
						Formatted	... [139]
						Formatted	... [140]
						Formatted	... [141]
						Formatted	... [142]
						Formatted	... [143]
						Formatted	... [144]
						Formatted	... [145]
						Formatted	... [146]
						Formatted	... [147]
						Formatted	... [148]
						Formatted	... [149]
						Formatted	... [150]
						Formatted	... [151]
						Formatted	... [152]
						Formatted	... [153]
						Formatted	... [154]
						Formatted	... [155]
						Formatted	... [156]
						Formatted	... [157]
						Formatted	... [158]
						Formatted	... [159]
						Formatted	... [160]
						Formatted	... [161]
						Formatted	... [162]
						Formatted	... [163]
						Formatted	... [164]
						Formatted	... [165]
						Formatted	... [166]
						Formatted	... [167]
						Formatted	... [168]
						Formatted	... [169]
						Formatted	... [170]
						Formatted	... [171]
						Formatted	... [172]
						Formatted	... [173]
						Formatted	... [174]
						Formatted	... [175]
						Formatted	... [176]
						Formatted	... [177]
						Formatted	... [178]
						Formatted	... [179]
						Formatted	... [180]
						Formatted	... [181]
						Formatted	... [182]
						Formatted	... [183]
						Formatted	... [184]
						Formatted	... [185]
						Formatted	... [186]
						Formatted	... [187]
						Formatted	... [188]
						Formatted	... [189]
						Formatted	... [190]
						Formatted	... [191]
						Formatted	... [192]
						Formatted	... [193]
						Formatted	... [194]
						Formatted	... [195]
						Formatted	... [196]
						Formatted	... [197]
						Formatted	... [198]
						Formatted	... [199]
						Formatted	... [200]

	Heptageniidae	unknown	1a	1	0
	Isonychiidae	<i>Heptagenia sp.</i>	1a	1	1
		<i>Isonychia sp.</i>	1b	1	0
Trichoptera	Leptoceridae	unknown	3	1	0
	Glossosomatidae	<i>Glossosoma intermedium</i>	1a	1	0
Neuroptera	Chrysopidae	<i>Chrysoperla sp.</i>	1a	1	0
	Hemerobiidae	<i>Hemerobius sp.</i>	1a	1	1
Hemiptera	Notonectidae	<i>Notonecta sp.</i>	1b	1	0
Coleoptera	Carabidae	<i>Agonom sp.</i>	1b	1	0
		<i>Agonom piceum</i>	1a	1	0
		<i>Dromius sp.</i>	1b	1	0
	Scirtidae	<i>Cyphon sp.</i>	1b	1	0
		<i>Cyphon phragmiteticola</i>	1a	1	0
Megaloptera	Sialidae	<i>Sialis sp.</i>	1b	1	0
Hymenoptera	Ichneumonidae	<i>Promethes sulcator</i>	1a	0	1

Table 2 Indices for richness, diversity and niche overlap for diet of bats from summer colonies and migration, derived from morphological analysis of faecal samples..

	Summer	Migration
Simpson's index	0.88	0.87
Species richness	14	13
Pianka's index	0.76	

Formatted: Indent: First line: 0"

Table 3 The frequency of prey groups in the diet of *P. nathusii* from summer colonies and migration based on morphological presence / absence data. We tested data with generalized liner model (GLM) and Tukey post hoc test. Significant differences are indicated with p values (in bold, -).

Formatted: Font: 11 pt, English (U.S.)

	Migration	Summer	z	p<
Nematocera	73.3%	47.8%	-1.531 -	-0.126 -
Anisopodidae	0.0%	21.7%	-0.007	0.995-
Chironomidae	40.0%	91.3%	3.034	0.00241
Culicidae	0.0%	17.4%	0.006- -	0.995-
Tipulidae	86.7%	30.4%	-3.051	0.00228
Brachycera	40.0%	52.2%	0.732 -	0.464-
Hemiptera	6.7%	0.0%	-0.003	-0.998
Corixidae	0.0%	8.7%	0.004-	0.997-
Cicada	0.0%	4.3%	0.004-	0.997-
Aphidoidea	13.3%	26.1%	0.927-	-0.354
Trichoptera	26.7%	17.4%	-0.681-	0.496
Lepidoptera	53.3%	30.4%	-1.396 -	0.163-
Ephemeroptera	26.7%	0.0%	-0.005-	0.996-
Neuroptera	46.7%	34.8%	-0.73	0.465-
Coleoptera	13.3%	0.0%	-0.005	0.996-
Hymenoptera	6.7%	4.3%	-0.311	-0.756
Araneae	13.3%	4.3%	-0.957	-0.338
Simuliidae	0.0%	4.3%	0.004-	-0.997
Formicidae	26.7%	0.0%	-0.005	0.996-

664 Figure 1 Overview of the sampling locations for faecal samples of *Pipistrellus nathusii* (Picture) in
665 Latvia. Summer colonies were sampled in Garkalne and Vecpiebalga. Migrating bats were caught and
666 sampld in Pape, Ornithological Station, situated within the migration route of *P. nathusii* (Species
667 photo by Viesturs Vintulis).

For Review Only

668 Figure 1 Overview of the sampling locations for faecal samples of *Pipistrellus nathusii* (Picture) in
669 Latvia. Summer colonies were sampled in Garkalne and Vecpiebalga. Migrating bats were caught and
670 sampld in Pape, Ornithological Station, situated within the migration route of *P. nathusii* (Species
671 photo by Viesturs Vintulis).

672

Figure 2 Plot of a non-metric two-dimensional ordination scale (NMDS) based on the presence-absence prey data derived from the morphological diet analysis on migrating *P. nathusii* (circle) and *P. nathusii* from summer colonies (corss) Non-metric two-dimensional ordination scale (NMDS) of morphological prey data (n= 50, stress = 0.20).

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

For Review Only

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

For Review Only

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

For Review Only

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

For Review Only

For Review Only

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

For Review Only

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

For Review Only

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

For Review Only

For Review Only

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

For Review Only

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

For Review Only

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

6/24/2013 4:58:00 PM

For Review Only

For Review Only

For Review Only

237x236mm (96 x 96 DPI)

65x50mm (300 x 300 DPI)