Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/57102/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Seric Jelaska, Lucija, Jurasovic, Jasna, Brown, David S., Vaughan, Ian P.
ORCID: <https://orcid.org/0000-0002-7263-3822> and Symondson, William O. C.
ORCID: <https://orcid.org/0000-0002-3343-4679> 2014. Molecular field analysis of trophic relationships in soil-dwelling invertebrates to identify mercury, lead and cadmium transmission through forest ecosystems. *Molecular Ecology* 23 (15) , pp. 3755-3766. 10.1111/mec.12566 file

Publishers page: <http://dx.doi.org/10.1111/mec.12566>
<<http://dx.doi.org/10.1111/mec.12566>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.

Molecular field analysis of trophic relationships in soil-dwelling invertebrates to identify mercury, lead and cadmium transmission through forest ecosystems

^{1,3}Lucija Šerić Jelaska, ²Jasna Jurasović, ³David S. Brown, ³Ian P. Vaughan, ³William O. C. Symondson

¹Faculty of Science, University of Zagreb, Rooseveltov trg 6, 10000 Zagreb, Croatia

²Institute for Medical Research and Occupational Health, Ksaverska cesta 2, 10000 Zagreb, Croatia

³Cardiff School of Biosciences, Cardiff University, Sir Martin Evans Building, Museum Avenue, Cardiff, CF10 3AX, UK

Key words: carabid beetles, earthworms, metal bioaccumulation, molecular gut content analyses, prey choice, slugs

Correspondence:

Lucija Šerić Jelaska, Fax: + 385 (0)1 4826260;

E-mail: slucija@biol.pmf.hr

Running title:

Molecular diet analyses and metal transfer

22 **Abstract**

23 Contamination pathways in complex food chains in soil ecosystems can be difficult to
24 elucidate. Molecular analysis of predator gut content can, however, rapidly reveal previously
25 unidentified trophic interactions between invertebrates and thereby uncover pathways of
26 pollutant spread. Here we measured concentrations of the toxic metals lead, cadmium and
27 mercury in carabid beetle predators and their prey. Invertebrates were sampled at one control
28 and four heavy metal polluted sites in order to reveal the impact of diet composition and
29 seasonal variation in prey availability on metal burden in carabids and metal transfer
30 pathways through forest ecosystems. This is the first report, to our knowledge, of carabid
31 diet composition based on PCR analysis of gut contents at the forest community level, rather
32 than in cultivated fields. Extensive screening using group- and species-specific primers
33 revealed that carabids ate primarily earthworms and slugs, as well as smaller numbers of
34 woodlice and springtails. Metal concentrations in carabids correlated with seasonal changes
35 in diet. Mercury accumulated in beetle predators more than in their slug prey. Since
36 earthworms, slugs and carabid beetles are the major prey of many birds and mammals, prey-
37 predator transfer and associated toxicity are major risks at mercury-contaminated sites.
38 Carabids may be useful bioindicators for assessing the impact of pollutants on soil
39 ecosystems, as long as species and seasonal factors are taken into account.

40

Introduction

Lead, cadmium and mercury are highly toxic metals without any biological function. While they do occur naturally in the soil at low levels, more than 200 years of industrialisation have led to elevated levels in the soil, degrading its quality and threatening soil biota (European Environment Agency & Joint Research Centre 2010). Eco-toxicity of metals correlates with their bioavailability and accumulation rate in soil biota. Bioavailability of metals depends, in turn, on the physical and chemical properties of the soil, and bioaccumulation depends on factors such as the metal tolerance of organisms, species-specific physiology and exposure history (e.g. van Straalen & van Wensem 1986; Hopkin 1989).

The main biological factors controlling metal accumulation in terrestrial invertebrates are diet and the structure and physiology of the digestive system (Hopkin 1989). After ingesting toxic metals, some organisms may avoid harmful effects by sequestering them in certain tissues and immobilising them in a form that cannot interfere with vital biological processes (Hopkin 1989). When organisms assimilate large amounts of heavy metals and do not excrete them, biomagnification up the food chain can occur. By the process of biomagnification, even low concentrations of some toxins (e.g. DDT, PCBs, organic mercury) can end up in living organisms at higher concentrations and can be disseminated to distant places through migration. Biomagnification of some toxic compounds occurs in both aquatic and terrestrial ecosystems affecting many organisms, such as raptors and otters (e.g. Castro *et al.* 2011; Mayack 2012), and can be harmful to human health. After mercury discharge in Japan in 1956, mercury contamination of fish and shellfish severely affected humans and animal populations for decades (e.g. Harada 1995). Although biomagnification of metals is less common in terrestrial food chains (reviewed in Janssen *et al.* 1993), it does occur in some

invertebrate species such as predatory lycosid spiders in the case of cadmium (Hendrickx *et al.* 2003), as well as along some species-specific trophic chains (van Straalen & Ernst 1991).

The large diversity of soil invertebrates makes it difficult to identify general patterns in metal distribution in the soil and in consequence bioaccumulation. Some organisms, such as earthworms, slugs and isopods accumulate metals in their tissue (e.g. Hopkin 1989; Dallinger 1993), while some hexapods efficiently excrete them (Hopkin 1989; Janssen *et al.* 1991; Lindquist *et al.* 1995). Taxonomic grouping by itself is not a sufficient basis for predicting accumulation rates in the environment. Furthermore, different species at the same trophic level have been found to accumulate different lead and cadmium concentrations (Georgii 1986; van Straalen & van Wensem 1986).

Seasonal variations are another complicating factor. The slug *Arion ater*, after feeding in the laboratory, takes up more metal in July than in September (Ireland 1981). The carabid *Calathus melanocephalus* takes up more cadmium in autumn than during other times of year (Janssen *et al.* 1991). Some carabid species active in autumn accumulate more cadmium than do species active in late spring (Šerić Jelaska *et al.* 2007). In addition to these apparently species-specific differences, seasonal changes in abiotic factors like temperature, day length and moisture, as well as seasonal changes in species composition, may cause variations in metal concentration in soil ecosystems during the year (Ireland 1981; Hopkin 1989; Šerić Jelaska *et al.* 2007).

In addition, to reveal heavy metal distribution and bioaccumulation in soil ecosystems, the pathways by which metals move from the soil into the biota need to be identified. Soil ecosystems feature complex food webs and highly biodiverse communities comprising different developmental stages from eggs and juveniles to adults, as well as numerous

microscopic and even cryptic species. These factors make it difficult to identify the many different trophic interactions involved and hence metal transfer pathways.

Carabids beetles have been used extensively for studying trophic interactions (Symondson *et al.* 1996; Holland 2002; Symondson *et al.* 2000, 2006; Eskelson *et al.* 2011; Hatteland *et al.* 2011; Boreau de Roince *et al.* 2012; Traugott *et al.* 2012; Davey *et al.* 2013) and the dynamics of heavy metal bioaccumulation and decontamination (Kramarz 1999; Stone *et al.* 2002; Butovsky 2011). They thus provide ecologically highly-connected indicator taxa for assessing metal contamination in many soil environments. Assessments based on single predator species may not capture effects on other species in the community, since even species within the same taxonomic group can show different behavioural, physiological and morphological responses to environmental stresses.

As hard-bodied chitinous insects, carabid adults are exposed to metal contamination mostly through food ingestion. Previous studies based on molecular and microscopic gut content analysis, as well as field observations, show that predatory carabids feed on a wide range of other soil invertebrates including earthworms, molluscs, woodlice, springtails and many insect taxa (e.g. Sunderland 1975; Hengeveld 1980; Sunderland & Sutton 1980; Symondson *et al.* 1996, 2000; Harper *et al.* 2005; King *et al.* 2010). Thus, their exposure to heavy metals may depend more on the concentration of the metal in prey tissue than its bioavailable concentration in the soil. Carabids, earthworms and slugs are in turn important food sources for birds and mammals.

Quantifying predation by carabids, through direct observation or microscopic gut screening, is difficult because they are mainly nocturnal and many of their prey are soft-bodied

organisms. As a result, knowledge of their food preferences is limited. Precise information on trophic interactions between beetles and other parts of the ecosystem is critical for understanding how different pollutants move from the soil into the biota. Recent advances in molecular analysis of gut contents allows more precise, *in situ* predation analyses that can reveal trophic interactions in greater detail (Harper *et al.* 2005; King *et al.* 2010).

Group- and species-specific PCR primers were used to detect prey choice by carabids among four dominant prey groups (earthworms, slugs, woodlice and springtails). 25 carabid species screened were collected in the field in late spring-early summer and in autumn. We quantified carabid predation in both seasons and we analysed lead, cadmium and mercury concentrations in carabids and in their main prey (earthworms and slugs) sampled from the same sites. We tested the hypothesis that prey choice, and lead, cadmium and mercury concentrations in prey, would explain the heavy metal concentrations in the carabids, providing a trophic pathway for transport of contaminants from soil into wildlife. We also assessed the extent to which each carabid species, and their community assemblage as a whole, might prove useful as indicators for metal pollution assessment.

Materials and methods

1. Field collection

Soil and animals were collected from four heavy metal polluted locations (L1-L4) and one control location (L5) in Croatia and the UK: Rudnik Zrinski – Medvednica, Croatia (L1); Tusti vrh – Medvednica, Croatia (L2); Llantrisant, Wales, UK (L3); Rudry 1, Wales, UK (L4); and Rudry 2, Wales, UK (L5). All sites were located in deciduous woodlands, and L1 - L4 were located near old coal mines and smelters, and consequently had high concentrations of mercury, lead and cadmium in the soil (Morgan & Morgan 1998; Nahmani *et al.* 2007; Šerić Jelaska *et al.* 2007). Site L5 was not located near pollution sources and therefore served as a control site. All sites had acid soil with average pH values ranging from 3.75 at site L1 to 5.85 at site L3 (Table S3).

Soil and invertebrate samples were collected from the end of May to the end of July and from mid-September to the end of October in 2007 in Croatia, and from mid-June to the end of July and in October in 2010 in Wales, UK.

For metal analyses, 8-16 top-soil samples (3L) per site were taken twice, in late spring and in autumn. Samples were taken randomly from the top 10 cm across the area at each site. Earthworms, slugs and carabids were collected from the same layer of soil at the same sites by digging and hand sorting, during three visits per season. Adult carabids were also collected by pitfall trapping to get substantial number of individuals for PCR analysis of their gut contents and heavy metal analyses. Five empty traps (0.5 L plastic cups) were left at each site for two weeks during each season. Traps were emptied every morning and beetles used for PCR analysis of gut content were placed individually in plastic tubes, transported to the laboratory in a cooler, killed immediately at -80 °C and stored at that temperature until

extracted for molecular analysis. Animals used in metal analyses were left in separate Petri dishes containing moist filter paper to empty their gut for 48 h and the filter paper was changed daily. They were killed at -80 °C and stored in the freezer before metal analysis. Woodlice found by digging and trapped in pitfalls were not stored but their densities were recorded.

All carabid beetles, earthworms and adult slugs were identified by morphology to species level using Cameron *et al.* (1983), Sims & Gerard (1985), Mršić (1997), Freude *et al.* (2004) and Luff (2007).

2. Molecular analyses

DNA extractions

DNA was extracted from the beetle foregut for diet analyses. DNA from earthworm and slug species was also extracted to serve as positive controls during PCR.

All primers, including newly designed ones for *Limax cinereoniger*, were tested for cross-amplification against DNA extracted from 35 soil invertebrate species representing potential non-target prey in the field, including predator DNA (Table S1). The non-target organisms were tested individually and no cross-amplifications were found.

Beetles were thawed to room temperature, the foreguts were removed as described in Symondson *et al.* (2000), and DNA was extracted using the DNeasy Blood & Tissue Kit (Qiagen), following the manufacturer's instructions. Negative controls with no added animal tissue were included in each batch of samples to check for potential DNA carry-over contamination during extraction.

182

183 To check for the presence of DNA after extraction and to avoid false negatives, extractions
184 were tested by PCR using general invertebrate primers for a 710-bp fragment of the
185 mitochondrial cytochrome oxidase I (COI) gene (Folmer *et al.* 1994). Each PCR (10 µL)
186 contained the following: 1 µL of template DNA, 0.625 U *Taq* polymerase (Invitrogen), 10
187 µM of each primer, 0.8 µL 2.5 mM dNTPs, 10x PCR buffer, 50 mM MgCl₂ (Invitrogen), 10
188 µg bovine serum albumin (New England Biolabs) and dH₂O (Sigma Aldrich). PCRs were
189 carried out in a GeneAmp 9700 thermal cycler (Applied Biosystems) using the following
190 conditions: 94 °C for 3 min; 45 cycles of 94 °C for 30 sec, 47 °C for 1 min, 72 °C for 1 min;
191 and finally 72 °C for 10 min.

192

193 *Selection of primers to identify prey in predator gut*

194 The primers were used to screen the gut contents of 317 field-caught carabid beetles for
195 earthworms, slugs, woodlice and springtails (Table 1).

196

197 Predation on earthworms was detected using general earthworm primers (Harper *et al.* 2005).
198 To screen for slugs, we used general *Arion* primers (Dodd 2004; Harper *et al.* 2005) to
199 amplify DNA of four *Arion* species (*A. hortensis*, *A. distinctus*, *A. silvaticus*, *A. subfuscus*)
200 and species-specific primers for *Deroceras reticulatum* (Dodd 2004; Harper *et al.* 2005). All
201 of these species were abundant at the study sites. We also designed species-specific primers
202 to amplify DNA of the slug *Limax cinereoniger* (Limacidae) (see below), which were
203 abundant at the two study sites in Croatia. The slugs *A. flagellus*, *A. ater*, *Malacolimax*
204 *tenellus*, *Testacella maugei* and *T. scutulum* were found at low abundance at the study sites,
205 and DNA of *A. flagellus* and *A. ater* was not amplified after testing with the general *Arion*
206 primers. No other adult slug species were recorded in the field surveys, except for *Tandonia*

207 *budapestensis* present at site L3. Since this species is highly toxic for carabid beetles
 208 (Symondson 1997) we assumed that it was not consumed by the beetles.
 209

210 *Primer design for Limax cinereoniger*

211 *Limax cinereoniger* COI sequences (accession number FJ606460) were aligned together with
 212 other sequences of slug species from the GenBank (*Limax maximus*, FJ606471; *L. sarnensis*,
 213 FJ606493; *L. wohlberedti*, FJ606481; *Deroceras reticulatum*, FJ481179; *Arion subfuscus*,
 214 AM259721; *A. hortensis*, AM259726) using Clustal W (Thompson *et al.* 1994) as
 215 implemented in BioEdit (Hall 1999). Species specific primers were then designed for *L.*
 216 *cinereoniger*, COI-Lcin-F1 and COI-Lcin-R1 (Table 1), that amplified a 198 bp DNA
 217 fragment. After evaluating their compatibility with NetPrimer (Premier Biosoft), annealing
 218 temperature was optimised in PCR.
 219

220 DNA from field-collected *L. cinereoniger* and negative control samples were amplified using
 221 the species-specific primers in a 10 µL reaction containing ca. 200 ng of total genomic DNA,
 222 0.625 U *Taq* polymerase (Invitrogen), 10 µM of each primer, 0.8 µL 2.5 mM dNTPs, 10x
 223 PCR buffer, 50 mM MgCl₂ (Invitrogen), 10 µg bovine serum albumin (New England
 224 Biolabs) diluted up to volume with sterile water (Sigma Aldrich). Thermal cycling conditions
 225 were as follows: 95 °C for 3 min; 35 cycles of 94 °C for 30 s, 61 °C for 45 s, 72 °C for 45 s;
 226 and finally 72 °C for 7 min. This novel primer pair proved to be species-specific under the
 227 optimised PCR cycling conditions with no evidence of cross-amplification, even when tested
 228 with related slug species such as *Limax maximus*.
 229

230 *Screening of field-caught predators*

Singleplex and multiplex PCRs were used to screen the gut contents of each field-caught beetle for the presence of earthworm and slug species (Table 1).

Singleplex PCR reactions (10 µL) contained 1 µL of template DNA, 0.625 U *Taq* polymerase (Invitrogen), 10 µM of each primer, 0.8 µL 2.5 mM dNTPs, 10x PCR buffer, 50 mM MgCl₂ (Invitrogen), 10 µg bovine serum albumin (New England Biolabs) and sterile water (Sigma Aldrich). PCRs were carried out in an AB Veriti 96-well thermocycler (Applied Biosystems). Cycling conditions for earthworm DNA samples were as follows: 94 °C for 3 min; 35 cycles of 94 °C for 30 s, 65 °C for 45 s, 72 °C for 45 s and finally 72 °C for 10 min. Cycling conditions for *L. cinereoniger* DNA samples were as follows: 95 °C for 3 min; 35 cycles of 94 °C for 30 s, 61 °C for 45 s, 72 °C for 45 s; and finally 72 °C for 7 min. Cycling conditions for amplifying woodlice and springtail DNA were as described in Jarman *et al.* (2006) and Kuusk & Agustí (2008).

Multiplex PCR reaction (10 µL), for amplifying *D. reticulatum* and *Arion sp.* DNA, contained 1.2 µL of extracted DNA, 5 µL of Multiplex PCR Master Mix (Qiagen), 0.2 µM each primer, 10 µg bovine serum albumin (New England Biolabs), and sterile distilled water (Qiagen). After initial denaturing step at 95 °C for 15 min, amplification proceeded for 35 cycles at 94 °C for 30s, 53 °C for 1 min 30 s, 72 °C for 1 min 30 s and a final extension at 72 °C for 10 min.

All PCRs included a positive control (target prey) and a negative control (sterile water instead of DNA). PCR products were separated on a 2% agarose gel for 40 min at 120 V and visualised with ethidium bromide (0.075 µg/mL).

All samples were screened for target prey twice using the same PCR conditions and those that came up positive only once were also accepted as positive and included in further analyses.

3. Metal analyses

Metal analyses were carried out on 206 beetles, 135 earthworms, 96 slugs and 111 soil samples using the ICP-MS technique. The procedure is described in Supporting Information (Metal analyses). Accumulation factors (AFs) for each metal were calculated by dividing mean metal concentrations in predator tissue per season per plot by the mean metal concentrations in prey.

4. Statistical analyses

Statistical analyses and figures were prepared using Statistica 9.1 (Statsoft, Inc. 2010), R (version 2.11.1, R Development Core Team, 2011) and Gephi 0.8.2 beta [Common Development and Distribution License (CDDL) & GNU General Public License 2008-2012]. Details of statistical analyses used in the paper are provided in Supporting Information (Statistical analyses).

Results

317 carabids were submitted for diet analyses and 206 for metal analyses. A total of 135 earthworms and 96 slugs were analysed for metal concentrations in tissue to investigate whether they can serve as metal transfer vectors through the food web.

Among the samples collected there were 25 carabid species, with *Nebria brevicollis*, *Abax parallelepipedus* and *Abax parallelus* accounting for 70% of captured animals (Table S2). *N. brevicollis* and *A. parallelepipedus* account for 77.3% of the carabid population sampled in the UK, while *A. parallelepipedus* and *A. parallelus* account for 61.4% of the total population in Croatian samples.

Prey consumption by carabids in the field

Carabids were shown to be consuming all four prey groups (earthworms, slugs, woodlice and springtails). Of 317 tested carabids, 213 (67.2%) contained at least one prey from these groups in their foregut. The foreguts of most beetles (138) contained only one prey group, 66 contained two, while eight contained three and one was positive for all four. Earthworms were detected most frequently, with 148 (46.7%) testing positive, while 98 beetles (30.9%) were positive for slugs, of which 52 beetles were positive for both earthworms and slugs. Woodlice and springtails were consumed by 35 (11.0%) and 17 (5.36%) beetles respectively. Prey consumption rates in the UK and Croatia were very similar with only woodlice showing a significant difference (chi-squared = 7.76, df = 1, $P = 0.005$), being consumed at higher rates in the UK.

Seasonal differences in prey consumption

Percentage of carabids testing positive for slugs or woodlice at the same study site differed between the two seasons (Fig. 1). Within sites L1 and L2, more slugs were preyed on in autumn and within sites L4 and L5 more woodlice were preyed on in spring.

There was little difference in predation on *Arion* slugs between seasons at all study sites. 28 individual beetles in the overall sample were positive for *Arion* sp. in spring and 26 in

autumn. However, carabid foregut analysis showed higher predation of *L. cinereoniger* and *D. reticulatum* in autumn than in spring. 80% and 60% of carabid beetles at L1 (Cro) and L2 (Cro) sites tested positive for *L. cinereoniger* in autumn compared to no positives and 6% of positives in spring, respectively. 7% and 10% of carabids tested positive for *D. reticulatum* in autumn at sites L4 (UK) and L5 (UK) respectively, with no positives in the spring-summer season.

Of the 25 carabid species recorded on the five study sites, eight were observed during both seasons. Foregut analysis of these eight species revealed significant seasonal differences in how much they preyed on slugs (T-test, $t=3.3$, $df=7$, $P=0.013$). The overall proportions of carabid species tested for the DNA of the four prey groups are listed in Table S2.

Proportions of earthworms, slugs and woodlice in the environment and in predators testing positive for these prey (Fig. 2) showed that isopods were taken in clearly lower proportions than would be expected if the carabids were predating randomly. All other prey were taken in approximately the ratios expected. Croatian sites were not included in these graphs because of low numbers of recorded animals (slugs, earthworms and woodlice) in the upper 10 mm of the soil, after extremely dry weather conditions throughout the year.

324 *Metal concentrations in invertebrates*

325 Metal concentrations in earthworms, slugs and carabids correlated positively with those in
326 soil samples from the same sites within the seasons ($r=0.97$, 0.84 and 0.83 for lead
327 respectively; $r=0.86$, 0.81 and 0.79 for cadmium respectively and $r=0.87$, 0.75 and 0.58 for
328 mercury respectively). Earthworm tissue showed the highest mean concentrations of lead,
329 cadmium and mercury of all predator and prey samples tested, while beetle tissue showed the
330 lowest concentrations of lead and cadmium but not of mercury. Slug tissue showed the lowest
331 mean concentrations of mercury, both in control and highly polluted sites (Fig. 3). Metal
332 concentrations were compared using the Kruskal–Wallis test. The median concentrations of
333 lead, cadmium and mercury differed significantly among invertebrate groups (lead, $N=437$,
334 $df=2$, $H=220.5$, chi squared= 184.3 ; cadmium, $H=208.9$, chi square= 173.2 ; mercury,
335 $H=73.63$, chi square= 73.6 ; $P<0.001$). All groups exhibited significantly higher lead,
336 cadmium and mercury concentrations on highly polluted sites (L1-L4) compared with the
337 control one (L5), (Mann-Whitney U Test, $P<0.05$).

338
339 Lead and cadmium concentrations in carabid tissue correlated strongly with those in
340 earthworms ($r=0.94$ for Cd, $r=0.92$ for Pb, $P<0.05$) and slugs ($r=0.85$ for Cd and Pb, $P<0.05$),
341 but not mercury concentration ($r=0.28$, $P=0.43$ with earthworms; $r=0.04$, $P=0.92$ with slugs).
342 The highest AF (>1), as the measure of metal transmission from prey to predator, was found
343 for mercury in the beetle-slug trophic pathway [Friedman ANOVA, chi squared ($N=9$, $df=5$)
344 = 33.125 , $P<0.01$], followed by mercury in the beetle-earthworm pathway [Friedman
345 ANOVA, chi squared ($N=9$, $df=4$) = 23.933 , $P<0.001$], (Fig. S1, Fig. 4 c). The lowest AF
346 was recorded for lead in the beetle-earthworm pathway (Fig. S1, Fig. 4 a).

347

348 *Seasonal and species variations in metal concentrations*

Seasonal comparisons of metal concentrations in animals within each study site revealed significant differences. Carabids contained higher lead and mercury concentrations in spring than in autumn, and higher cadmium concentrations in autumn (Mann-Whitney U Test for lead: $U = 95.000$, $P = 0.032$ on location L3 and $U = 94.000$, $P = 0.036$ on location L5; for cadmium: $U = 136.000$, $P = 0.009$ on location L4), (Table S3). Cadmium concentration in slugs was also higher in autumn, even higher than in the earthworms (Fig. 3). In general, lead, cadmium and mercury concentrations in animal tissue at highly polluted sites showed greater seasonal differences in carabids and slugs than in earthworms (Fig. 3).

Node diagrams (Fig. 4) show the links between mean metal concentrations in carabid species and the main prey, presented by the size of nodes and predation events in both seasons. Significant differences were found in metal concentrations between carabid species (Kruskal-Wallis ANOVA, $P < 0.05$), (Table S4). *N. brevicollis* accumulated more lead and cadmium (Fig. 4, Table S4) than did the two *Abax* species (lead, $U = 718.000$, $Z = 5.31836$; cadmium, $U = 996.000$, $Z = 3.83615$; $P < 0.001$). *A. parallelepipedus* accumulated more mercury than did *Nebria* species ($U = 393.000$, $Z = -7.05115$, $P < 0.001$). *Abax* species were dominant at the highly mercury-polluted sites.

Discussion

Molecular gut content analyses provided detailed insight into carabid diet under field conditions, allowing the construction of metal transfer pathways and providing possible explanations for seasonal variations in metal tissue concentrations. We revealed earthworms and slugs as the main prey within carabid communities with more slugs being consumed in autumn than in late spring and early summer. We confirmed our main hypothesis that

seasonal shift in prey consumed, and lead, cadmium and mercury concentrations in those prey, influenced metal concentrations in the carabids.

We observed a decrease in mercury and lead concentrations in predators in autumn, which coincided with a doubling in the proportion of beetles testing positive for slugs, allowing us to correlate the decrease in metal concentrations in carabids with the fact that slugs accumulated lower levels of these metals than did earthworms. In the case of cadmium, the tissue burden in carabids was higher in autumn than spring, coinciding with higher concentrations in slugs.

Diet composition and heavy metal tissue burden

Although many studies have used molecular methods to study natural regulation of pests (Symondson *et al.* 1996; Chen *et al.* 2000; Nash *et al.* 2008; Eskelson *et al.* 2011; Hatteland *et al.* 2011; Boreau de Roincé *et al.* 2012) they can also elucidate complex food webs and thereby track pathways of pollutants between trophic levels in ecosystems. For example, data on metal transmission through trophic interactions have been obtained from feeding experiments (Janssen *et al.* 1991, Kramarz 1999; Hendrickx *et al.* 2003) and from analysis of stable isotope ratios (e.g. Cabana & Rasmussen 1994).

Metal transfer has been measured *in situ* indirectly by measuring metal concentrations in different food chain compartments (e.g. Notten *et al.* 2005, Roodbergen *et al.* 2008). We used a similar approach in our study to measure metal concentrations in different food web compartments (soil, earthworms that ingest soil substrate, slugs as herbivores and carabids as carnivorous), in order to explore the relationship between the relative proportion of each prey group in the beetle diet and metal accumulation in beetle tissue. Our results confirmed the

expected positive relationship between metal concentrations in predator and in prey. Earthworms as the main prey with the highest metal concentrations in the tissue represent the most important link in trophic pathways of contaminants from the soil to predators. In addition, we observed that carabid beetles had lower concentrations of lead and cadmium than slugs and earthworms, but had more mercury than slugs ($AF > 1$), raising the possibility of mercury biomagnification (Fig. S1). Unlike spiders, which lack mechanisms to excrete cadmium (Hendrickx *et al.* 2003), carabids showed lower Pb and Cd concentrations, with seasonal and species differences detected.

Taking into account that each prey contributes differently to metal accumulation in the predator, the AF could be calculated by multiplying mean metal concentrations in earthworms and slugs by the percentage of beetles testing positive and adding up the concentrations of both prey to yield the total metal concentration from prey. Then, the mean metal concentration in the predator (carabids) would be divided by this total concentration from both prey. This approach could be useful for assessing metal transfer from multiple prey combinations to predators and thus calculating more accurate AF values, but predation rates need to be adjusted for all predator-prey combinations. As it was not possible within this study to correct for the multiple combinations of primers and predator species in the analyses, simple AF calculations have been done.

Although carabid tissue showed lower Pb and Cd concentrations than did their prey, carabids from highly polluted sites exhibited much higher metal concentrations than did animals from control sites (Fig. 3). Some differences between sites may be influenced by metal availability due to chemical and physical properties of the soil (i.e. pH values, Table S3). Still some clear trends in metal concentrations in invertebrates were evident.

423

424 *Carabid beetles as bioindicators for metal pollution assessment*

425 Carabid species differ in lifespan, as well as in feeding and breeding behaviours (autumn or
426 spring breeders). Moreover, the same species may even change its diet during the year:

427 *Carabus violaceus* consume more slugs in early spring and again in autumn (Paill 2000). All
428 these differences may affect metal concentrations and pollution assessment. Species and
429 seasonal variations in heavy metal tissue burden have previously been described in carabids
430 collected from the field in other studies (i.e. Janssen *et al.* 1991; Purchart & Kula 2007; Šerić
431 Jelaska *et al.* 2007; Butovsky 2011).

432

433 *Seasonal differences in metal burden*

434 Carabids had higher lead and mercury concentrations in spring than in autumn, but higher
435 cadmium concentrations in autumn. In an attempt to explain these differences, we
436 superimposed changes in heavy metal concentrations in carabid tissue with numbers of
437 carabids testing positive for each prey group. Since earthworms showed the greatest lead and
438 mercury accumulation and were also the main carabid prey, we speculate that trophic
439 pathways involving earthworms should be the main determinant of metal concentration in
440 carabids at highly polluted sites. Slugs accumulate lead and mercury to a much smaller extent
441 than do earthworms, and carabids consume a higher proportion of slugs in the autumn than in
442 spring and early summer (Table S2). This shift in consumed prey may explain why lead and
443 mercury concentrations in carabid predators were lower in autumn. This same shift may also
444 explain seasonal variations in cadmium concentrations. Not only carabid predators but also
445 the slugs showed higher cadmium tissue concentrations on polluted sites in autumn (Fig. 3).
446 During this time, slugs showed maximal levels of this metal (Table S3), coinciding with the
447 time when carabids consumed an increasing proportion of them as prey. Although Ireland

(1981) found greater cadmium uptake by *A. ater* slugs after feeding trials in July than in September, here we measured higher concentrations in slugs in autumn on the two cadmium-rich locations L3 and L4; concentrations at the other sites were higher in spring.

Species differences in metal burden

Species differences in metal concentrations in carabid tissue were observed, including among the most abundant *Abax* and *Nebria* species. For example, the highest cadmium concentrations were measured in *Nebria brevicollis* tissue with 2.2 and 2.4 times more cadmium on L3 and L4 locations (with higher cadmium concentrations in the soil) compared with the control L5 location. On the other hand, *A. parallelepipedus* accumulated 1.2 times more lead at lead-contaminated site L3 compared with control site. In both species, lead concentrations were higher in spring. *Nebria* accumulated more cadmium than did *Abax*, especially in autumn when more *Nebria* were trapped, predating more on slugs than did species collected in spring-summer (Fig. 4). Links in food webs (Fig. 4) gave a good indication of interaction strengths. Still, some adjustments based upon modeling the decay curves for each predator-prey combination would probably be needed to obtain more precise results, although given the large numbers of carabid-primer combinations involved this may not be practical. At least 100 individual predators would be needed for each trial to model the decay curve for each predator-prey combination shown in Fig. 4.

Similar seasonal trends were observed in previous studies (reviewed in Butovsky 2011). These findings suggest that season and carabid community structure should be taken into account when using these beetles to assess metal pollution in the field.

473 *Carabid beetles diet*

474 Earthworms and slugs were the primary components of the carabid diet, with more slugs
475 consumed in autumn than in late spring or early summer. Some carabid species, such as
476 *Pterostichus melanarius*, may rely on their earthworm diet to improve their fitness
477 parameters (Symondson *et al.* 2006). Also, some species of carabids have been shown to
478 prefer slug eggs and juveniles (Paill 2000; Hatteland *et al.* 2010) because adult slugs have
479 effective defence mechanisms (McKemey *et al.* 2003). A study by Foltan (2004) showed that
480 arionid slugs deterred ground beetle attacks more effectively than did limacid slugs. Our data
481 strongly suggest that the observed number of isopod positive beetles was less than expected
482 in both seasons (Fig. 2). Again, we have conducted no statistical analyses on these results for
483 the reasons explained. As earthworms were clearly the most abundant prey available in the
484 autumn, they represent the most important trophic pathway available at that time.

485

486 Although *Nebria brevicollis* has been described as preying primarily on springtails (Thiele
487 1977), we detected all four prey groups in its foregut, with earthworms present in more than
488 40% of individuals. The proportion of carabids in our study positive for woodlice (18.8% in
489 UK locations) was similar to the 17% reported by Sunderland & Sutton (1980) in grasslands
490 in UK. The number of carabids testing positive for each prey group could be changed to some
491 degree by different detection periods following ingestion. Decay rates might be different for
492 each predator and primer combination, and therefore have to be calculated to get more
493 precise results and interaction strengths.

494

495 This study has provided insights not only at the level of individual species, but also at the
496 community level. Molecular gut content analyses allowed us to screen the entire community

rapidly for a broad range of prey, elucidate the position of carabids within the ecosystem and identify nutrient flows while taking into account species composition and abundance. These findings suggest the possibility of using carabids as bioindicators of heavy metal contamination in soil-based ecosystems, as long as species and seasonal variations are taken into account, since these factors affect metal concentrations in carabids. The high accumulation factor for mercury (>1) on mercury polluted sites highlights the need to analyse the potential for mercury transmission in other predators.

Acknowledgements

This work was supported by postdoctoral grant of the Croatian Science Foundation (Project No. 02.03/88), awarded to LSJ and by the Croatian Ministry of Science, Education and Sports (Grants No. 022-0222148-2135 and 119-1193080-1206). The Project was hosted by WOCS at Cardiff University. We would like to thank numerous colleagues (S. Jelaska, I. Russo, J.K. Bluemel, S. Mondol, M. Temunović, V. San Andrés, A. Sekovanić, A. Sulimanec Grgec, T. Orct, S. Mataušić, K. Vlahoviček, D. Franjević) for all their help in the field and in the laboratory and valuable advices on data analyses. Great thank to J. Morgan and R. Donnelly for their help in choosing the study sites and for providing valuable data on soil conditions.

References

- Bureau de Roince C, Lavigne C, Ricard J, Franck P, Bouvier J, Garcin A, Symondson WOC (2012) Predation by generalist predators on the codling moth versus a closely-related emerging pest the oriental fruit moth: a molecular analysis. *Agricultural and Forest Entomology*, **14**, 260–269.
- Butovsky RO (2011) Heavy metals in carabids (Coleoptera, Carabidae). *Zookeys*, **100**, 215–222.
- Cabana G, Rasmussen JB (1994) Modelling food chain structure and contaminant bioaccumulation using stable nitrogen isotopes. *Nature (London)*, **372**, 255–257.
- Cameron RAD, Jackson N, Eversham B (1983) A field key to the slugs of the British Isles Taunton: Field Studies Council.
- Castro I, Aboal JR, Fernandez JA, Carballeira A (2011) Use of raptors for biomonitoring of heavy metals: gender, age and tissue selection. *Bulletin of Environmental Contamination and Toxicology*, **86**, 347–351.
- Chen Y, Giles KL, Payton ME, Greenstone MH (2000) Identifying key cereal aphid predators by molecular gut analysis. *Molecular Ecology*, **9**, 1887–1898.

539 Dallinger R (1993) Strategies of metal detoxification in terrestrial invertebrates. In: Dallinger,
540 R., Rainbow, P.S. (Eds.), *Ecotoxicology of Metals in Invertebrates*. Lewis Publishers,
541 London, pp. 245–280.

542

543 Davey JS, Vaughan IP, King RA, Bell JR, Bohan DA, Bruford MW, Holland JM,
544 Symondson WOC (2013) Intraguild predation in winter wheat: prey choice by a common
545 epigeal carabid consuming spiders. *Journal of Applied Ecology*, **50**, 271–279.

546

547 Dodd, CS (2004) Development and optimisation of PCR-based techniques in predator gut
548 analysis. PhD thesis, Cardiff University, Cardiff, UK.

549

550 European Environment Agency, Joint Research Centre (2010) The European environment —
551 state and outlook 2010, Soil. EEA and JCR, Luxemburg, Publications Office of the European
552 Union.

553

554 Eskelson MJ, Chapman EG, Archbold DD, Obrycki JJ, Harwood JD (2011) Molecular
555 identification of predation by carabid beetles on exotic and native slugs in a strawberry
556 agroecosystem, *Biological Control*, **56**, 245-253.

557

558 Freude H, Harde KW, Lohse GA, Klausnitzer B (2004) Die Käfer Mitteleuropas, Band 2,
559 Spectrum, Elsevier, München, 521 pp.

560

561 Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of
562 mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates.

563 *Molecular Marine Biology and Biotechnology*, **3**, 294–299.

564

565 Foltan P (2004) Influence of slug defence mechanisms on the prey preferences of the carabid
 566 predator *Pterostichus melanarius* (Coleoptera: Carabidae). *European Journal of Entomology*,
 567 **101**, 359–364.

568

569 Georgii HW (1986) Atmospheric Pollutants in Forest Areas: Their Deposition and
 570 Interception. Kluwer Academic Publishers 289 pp.

571

572 Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis
 573 program for Windows 95/98/NT. *Nucleic Acids Symposium Series*, **41**, 95–98.

574

575 Harada (1995) Minamata disease: methylmercury poisoning in Japan caused by
 576 environmental pollution. *Critical Reviews in Toxicology*, 25/1: 1-24.

577

578 Harper GL, King RA, Dodd CS, Harwood JD, Glen DM, Bruford MW, Symondson WOC
 579 (2005) Rapid screening of invertebrate predators for multiple prey DNA targets. *Molecular*
 580 *Ecology*, **14**, 819–827.

581

582 Hatteland BA, Symondson WOC, King RA, Skage M, Schander C, Solhøy T (2011)
 583 Molecular analysis of predation by carabid beetles (Carabidae) on the invasive Iberian slug
 584 *Arion lusitanicus*. *Bulletin of Entomological Research* 101, 675-686.

585

586 Hendrickx F, Maelfait J, Langenbick F (2003) Absence of cadmium excretion and high
 587 assimilation result in cadmium biomagnification in a wolf spider. *Ecotoxicology and*
 588 *Environmental Safety*, **55**, 287–292.

589

590 Hengeveld R (1980) Polyphagy, oligophagy and food specialization in ground beetles
591 (Coleoptera, Carabidae). *Netherlands Journal of Zoology*, **30**, 564-584.

592

593 Holland J (ed.) (2002) *The Agroecology of Carabid Beetles*. Intercept Ltd., 356 pp.

594

595 Hopkin SP (1989) *Ecophysiology of Metals in Terrestrial Invertebrates*. Elsevier Applied
596 Science, London, 366pp.

597

598 Ireland MP (1981) Uptake and distribution of cadmium in the terrestrial slug *Arion ater* (L.).
599 *Comparative Biochemistry and Physiology*, **68A**, 37 – 41.

600

601 Janssen MPM, Bruins A, De Vries TH, Van Straalen NM (1991) Comparison of cadmium
602 kinetics in four soil arthropod species. *Archives of Environmental Contamination Toxicology*
603 **20**, 305-312.

604

605 Janssen MPM, Ma WC, Van Straalen NM (1993) Biomagnification of metals in terrestrial
606 ecosystem. *Science of Total Environment Suppl.*, 511-524.

607

608 Jarman SN, Redd KS, Gales NJ (2006) Group-specific primers for amplifying DNA
609 sequences that identify Amphipoda, Cephalopoda, Echinodermata, Gastropoda, Isopoda,
610 Ostracoda and Thoracica. *Molecular Ecology Notes*, **6**, 268–271.

611

612 King RA, Vaughan IP, Bell JR, Bohan DA, Symondson WOC (2010) Prey choice by carabid
 613 beetles feeding on an earthworm community analysed using species- and lineage-specific
 614 PCR primers. *Molecular Ecology*, **19**, 1721–1732.
 615
 616 Kramarz P (1999) Dynamics of accumulation and decontamination of cadmium and zinc in
 617 carnivorous invertebrates. 1. The Ground Beetle, *Poecilus cupreus* L. *Bulletin of*
 618 *Environmental Contamination and Toxicology*, **63**, 531–537.
 619
 620 Kuusk AK, Agustí N (2008) Group-specific primers for DNA-based detection of springtails
 621 (Hexapoda: Collembola) within predator gut contents. *Molecular Ecology Resources*, **8**, 678–
 622 681.
 623
 624 Lindquist L, Block M, Tjälve H (1995) Distribution and excretion of Cd, Hg, methyl-Hg and
 625 Zn in the predatory beetle *Pterostichus niger* (Coleoptera: Carabidae). *Environmental*
 626 *Toxicology and Chemistry*, **14**, 1195–1201.
 627
 628 Luff ML (2007) The Carabidae (ground beetles) of Britain and Ireland. Carl Hildebrand
 629 Lindroth, Royal Entomological Society of London, 247 pp.
 630
 631 Mayack DT (2012) Hepatic mercury, cadmium, and lead in mink and otter from New York
 632 State: monitoring environmental contamination. *Environmental Monitoring and Assessment*,
 633 **184**, 2497–2516.
 634

635 McKemey AR, Symondson WOC, Glen DM (2003) Predation and prey size choice by the
636 carabid beetle *Pterostichus melanarius* (Coleoptera: Carabidae): the dangers of extrapolating
637 from laboratory to field. *Bulletin of Entomological Research*, **93**, 227-234.

638

639 Morgan JE, Morgan AJ (1998) The distribution and intracellular compartmentation of metals
640 in the endogeic earthworm *Aporrectodea caliginosa* sampled from an unpolluted and a metal-
641 contaminated site. *Environmental pollution*, **99/2**, 67-175.

642

643 Mršić N (1997) Živali naših tal. Uvod v pedozoologijo—sistematika in ekologija s splošnim
644 pregledom talnih živali. Tehniška založba Slovenije, Ljubljana, 416 pp.

645

646 Nahmani J, Hodson ME, Black S (2007) A review of studies performed to assess metal
647 uptake by earthworms. *Environmental Pollution*, **145**, 402-424.

648

649 Nash MA, Thomson LJ, Horne PA, Hoffman AA (2008) *Notomus gravis* (Chaudoir)
650 (Coleoptera: Carabidae) predation of *Deroceras reticulatum* Müller (Gastropoda:
651 Agrolimacidae), an example of fortuitous biological control. *Biological Control* **47**, 328–334.

652

653 Notten M, Oosthoek A, Rozema J, Aerts R (2005) Heavy metal concentrations in a soil–
654 plant–snail food chain along a terrestrial soil pollution gradient. *Environmental Pollution*,
655 **138**, 178–190.

656

657 Paill W (2000) Slugs as a prey for larvae and imagines of *Carabus violaceus* L. (Coleoptera,
658 Carabidae) pp. 221-222. In Brandmayr P, Lovei GL, Brandmayr TZ, Casale A and Vigna

659 Taglianti A (Eds.), Natural history and applied ecology of carabid beetles. Sofia, Bulgaria,
660 Pensoft Publishers.
661

662 Purchart L, Kula E (2007) Content of heavy metals in bodies of field ground beetles
663 (Coleoptera, Carabidae) with respect to selected ecological factor. *Polish Journal of Ecology*
664 **55**, 305-314.
665

666 Roodbergen M, Klok C, van der Hout A (2008) Transfer of heavy metals in the food chain
667 earthworm Black-tailed godwit (*Limosa limosa*): comparison of a polluted and a reference
668 site in The Netherlands. *Science of Total Environment*, **406**, 407-412.
669

670 Sims RW, Gerard BM (1985) Earthworms: Keys and Notes for the Identification and Study
671 of the Species. Linnean Society of London, Estuarine and Brackish-Water Sciences
672 Association, Brill Archive, 171 pp.
673

674 Stone D, Jepson P, Laskowski R (2002) Trends in detoxification enzymes and heavy metal
675 accumulation in ground beetles (Coleoptera: Carabidae) inhabiting a gradient of pollution.
676 *Comparative Biochemistry and Physiology. Part C* **132**, 105-112.
677

678 Sunderland KD (1975) The diet of some predatory arthropods in cereal crops. *Journal of*
679 *Applied Ecology*, **12**, 389–396.
680

681 Sunderland, KD, Sutton SL (1980) A serological study of arthropod predation on terrestrial
682 isopods in a dune grassland ecosystem. *Journal of Animal Ecology*. **49/3**, 987-1004.
683

684 Symondson WOC (1997) Does *Tandonia budapestensis* (Mollusca: Pulmonata) contain
 685 toxins? Evidence from feeding trials with the slug predator *Pterostichus melanarius*
 686 (Coleoptera: Carabidae). *Journal of Molluscan Studies* **63**: 541–545.
 687
 688 Symondson WOC, Cesarini S, Dodd PW, Harper GL, Bruford MW, Glen DM, Wiltshire
 689 CW, Harwood JD (2006) Biodiversity vs. biocontrol: positive and negative effects of
 690 alternative prey on control of slugs by carabid beetles. *Bulletin of Entomological Research*,
 691 **96**, 637–645.
 692
 693 Symondson WOC, Glen DM, Erickson ML, Liddell JE, Langdon CJ (2000) Do earthworms
 694 help to sustain the slug predator *Pterostichus melanarius* (Coleoptera: Carabidae) within
 695 crops? Investigations using monoclonal antibodies. *Molecular Ecology*, **9**, 1279–1292.
 696
 697 Symondson, W.O.C., Glen, D.M., Wiltshire, C.W, Langdon, C.J. & Liddell, J.E. (1996)
 698 Effects of cultivation techniques and methods of straw disposal on predation by *Pterostichus*
 699 *melanarius* (Coleoptera: Carabidae) upon slugs (Gastropoda: Pulmonata) in an arable field.
 700 *Journal of Applied Ecology* 33, 741-753.
 701
 702 Šerić Jelaska L, Blanusa M, Durbesic P, Jelaska SD (2007) Heavy metal concentrations in
 703 ground beetles, leaf litter, and soil of a forest ecosystem. *Ecotoxicology and Environmenal*
 704 *Safety*, **66**, 74-81.
 705
 706 Thiele HU (1977) Carabid beetles in their environments. Springer Verlag, Berlin, Heidelberg,
 707 New York, 369 pp.
 708

Thompson JD, Higgins DG, Gibson TJ (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research*, **22**, 4673-4680.

Traugott M, Bell JR, Raso L, Sint D, Symondson WOC (2012) Generalist predators disrupt parasitoid aphid control by direct and coincidental intraguild predation. *Bulletin of Entomological Research*, **102**, 239-247.

Van Straalen NM, Ernst WHO (1991) Metal biomagnification may endanger species in critical pathways. *Oikos*, **62**, 255–256.

Van Straalen NM, van Wensem J. (1986) Heavy metal content of forest litter arthropods as related to body-size and trophic level. *Environmental Pollution*, **42**, 209-221.

Data Accessibility

Lead, cadmium and mercury concentrations (dry weight) in soil and invertebrates: DRYAD entry doi:10.5061/dryad.5n02f/1. Raw data of molecular gut content analyses (MGCA) in carabid beetles: DRYAD entry: doi:10.5061/dryad.5n02f/2 .

Author Contributions Box

LSJ designed the study, collected the samples, performed molecular diagnostics of sampled material, data analyses and wrote the manuscript; DSB conducted part of molecular gut content analyses; JJ performed metal analyses; LSJ and IPV performed statistical analysis;

733 WOCS supervised this work from the very beginning, adding valuable suggestions and
734 revising the manuscript.
735
736
737

Figure legends:

Figure 1. Beetles testing positive for each prey group in spring-early summer (in 2007 and 2010 together) and autumn (in both years together). Box represents mean $\pm$ SE and whisker $\pm$ 0.95 confidence interval.

Figure 2. Number of PCR positive beetles for three prey groups (shown in open bars) and abundance of each prey group in the environment (shown in black bars). Upper graph showed data across the three locations in the UK (L3-5) for the spring-summer period, and lower graph showed data across three locations in the UK for autumn.

Figure 3. Lead, cadmium and mercury concentrations (mean, $m \pm SE$, $m \pm 1.96*SE$) in earthworms, slugs and carabids collected in spring-summer 2007, 2010 (open boxes) and in autumn 2007, 2010 (filled boxes) in control (L5) and polluted locations (L1-4). Mean metal concentration values from the control site are marked as squares, and those from the polluted sites are marked as circles.

Figure 4. Node diagrams of observed trophic interactions between prey groups and predator species with measured metal concentrations in the tissue. Size of the nodes reflects mean concentrations of lead (a), cadmium (b), and mercury (c) in earthworms, slugs and carabids. Yellow circles represent beetles caught in spring-summer, and purple circles represent those caught in autumn. Arrow thickness represents the percentage of carabid species testing positive for the given trophic interaction. Black circle represents metal concentration of 0.1 mg/kg, with an arrow thickness representing 10% of carabid species being positive for a given interaction. Force Atlas model was chosen as a layout. Node sizes range from 20-150

763 for Pb (0.08-623.1 mg/kg), Cd (0.1-56.5 mg/kg) and Hg (0.035-0.666 mg/kg) concentrations
764 measured in carabids. The same range is chosen for arrow thickness, representing the
765 percentage (from 0.1 to 1) of PCR positive carabid species tested for each prey within the
766 season.
767
768 Tables

769 Table 1. Sequences of primers (5'-3') used for multiplex (M) and singleplex (S) PCR analyses.

Species	PCR	Primer name	Primer sequence	Amplicon size (bp)	Reference
Earthworms	S	185F	TGTGTACTGCCGTCGTAAGCA	225-236	Harper <i>et al.</i> 2005
		14233R	AAGAGCGACGGGCGATGTGT		
<i>Deroceras reticulatum</i>	M	Dr11F	CTATACACAATTTTAAATAAG	109	Dodd 2004, Harper <i>et al.</i> 2005
		DRF29RC	GCTTCTGGTTTATCTATTATTGGT		
<i>Arion sp.</i>	M	Ai1F	CACATAAATGATAGTCACC	208-221	Dodd 2004, Harper <i>et al.</i> 2005
		AR2R	ATACTTACAAGTCCATCTTT		
<i>Limax cinereoniger</i>	S	COI-Lcin-F1	TGAACTGTATACCCGCCTTTG	198	This study
		COI-Lcin-R1	CCTGCCAATACAGGAAGCG		
Springtails	S	Col3F	GGACGATYTTRTTRGTTCGT	177&272	Kuusk & Agustí 2008
		Col4F	GCTACAGCCTGAACAWTWG		
		Col5R	TCTTGGCAAATGCTTTCGCAGTA		
Woodlice	S	IsopodNSSf1	TCATGATTYATGGGATGT	201-278	Jarman <i>et al.</i> 2006
		IsopodNSSr1	AAGACCTCAGCGCTCGGC

Supporting information

Additional supporting information may be found in the online version of this article.

Metal analyses

Lead, cadmium and mercury were analysed by inductively coupled plasma – mass spectrometry (ICP-MS) using the Agilent 7500cx (Agilent Technologies, Waldbronn, Germany).

Whole animals were washed with ultrapure water (18 MΩ cm; obtained with a GenPure system, TKA, Germany), weighed in a quartz digestion vessels, dried, weighed again and digested in 50% (v/v) nitric acid (3-5 mL, depending on the weight of the samples) in an UltraCLAVE IV microwave digestion system (Milestone, Italy). Soil samples were air-dried and sieved through 2 mm meshes. Dried powdered soil samples (approximately 0.1 g) were weighed into PTFE vessels and digested in 2 mL conc. HNO₃ and 0.4 mL conc. HF, according to UltraCLAVE IV application instructions for the digestion of soil (*UltraCLAVE Application Note N. UC-17. In: UltraCLAVE IV user manual Rev. 02/2008. Leutkirch: MLS-Milestone GmbH Mikrowellen-Laborsysteme*).

Digested samples were adjusted to 30 mL and just before the analysis by ICP-MS those samples were additionally diluted with 1% nitric acid in ultrapure water by a factor of 10-50, depending on the type and starting dry mass of the sample. All standard solutions (for external calibration and internal standards) were prepared from 1000 mg/L PlasmaCAL (SCP Science, Canada) single element standards. Isotopes ¹¹⁴Cd, ²⁰²Hg and ²⁰⁸Pb were used for element quantification, while ¹⁰³Rh, ¹⁵³Tb and ¹⁹³Ir were used as internal standards to correct for instrument drift and matrix effects.

Certified standard reference materials were used in duplicate with each sample digestion series for quality assurance. BCR185R Bovine Liver and BCR186 Pig Kidney (Institute for Reference Materials and Measurements, Geel, Belgium) were used to control measurement accuracy of earthworm and slug samples, while San Joaquin Soil 2709 (National Institute of Standards and Technology, USA) was used to control for soil sample measurement accuracy. The results agreed well with the certified values - recoveries ranged from 92% to 110%, with the exception of Hg in San Joaquin Soil NIST 2709 sample (84-92%).

Statistical analyses

Prior to statistical analysis, all data were checked using Shapiro-Wilk's test to see whether they followed a normal distribution; if not, they were transformed [$\log(x+1)$]. The Levene post-hoc test was used to test the assumption of equal variances. If the transformed data followed a normal distribution based on Shapiro-Wilk's test, they were analysed using parametric statistical tests. If not, they were analysed using the following nonparametric tests: Friedman ANOVA and the Mann-Whitney U test, to compare two independent samples; the Kruskal-Wallis test, to compare multiple independent samples followed by multiple comparisons of mean ranks for all groups.

Friedman ANOVA with a significance threshold of $P < 0.05$ was used to compare lead, cadmium and mercury concentrations factors between carabids and earthworms and between carabids and slugs at the five study sites in the two seasons. The Kruskal-Wallis test was used to test the null hypothesis of no difference in group median metal concentrations among earthworms, slugs and carabids among the sites. The Mann-Whitney U Test was used to compare the metal concentrations in each group of animals (earthworms, slugs and carabids) between the two seasons. Two tailed T-test was used to check for differences in predation

events within the same predator-prey combinations between two seasons. Correlations between prey consumption and metal concentration in carabids at different study sites and during both seasons were calculated using the Pearson correlation test.

Figure S1. Mean ($\pm$ SE and $1.96 \times$ SE) accumulation factors (AFs) for lead, cadmium and mercury in carabids (c) and their prey (e, earthworms; s, slugs).

Table S1. List of invertebrates used for non-target primer tests.

Carabid beetles

Carabus nemoralis

Abax parallelepipedus

Calathus fuscipes

Notiophilus biguttatus

Bembidion sp.

Other beetles

Rove beetles - *Ocypus olens*

Dor beetle - *Geotrupes* sp.

Hymenoptera

Ant (sp 1)

Ant (sp 2)

Ant (sp3)

Diptera

Crane fly larvae (sp 1)

Butterflies and Moths

Pieris brassicae

Cydia pomonella

Grapholita molesta

Earwings (sp 1)

Springtails (sp 1)

Spiders

Pardosa sp.

Erigone atra

Woodlice

Philoscia muscorum

Oniscus asellus

Porcellio scaber

Earthworms

Aporrectodea caliginosa

Aporrectodea smaragdina

Eisenia foetida

Lumbricus castanea

Lumbricus rubellus

Lumbricus terrestris

Deandrobaena octaedra

Slugs

Deroceras reticulatum

Limax maximus

Lehmannia marginata

Tandonia budapestensis

Snails

Cepea nemoralis

Helix aspersa

Nematodes (sp 1)

Table S2. Screening of field-caught carabid species for four prey groups. Below are the numbers of individuals of each species tested and percentage of beetles testing positive for each prey group.

Species	Locations where species occurred	No. ind. tested for diet	Beetles testing positive (%)			
		(overall	Earthworms	Slugs	Woodlice	Springtails
		abundances)				
<i>Nebria brevicollis</i>	3,4,5	87 (165)	40.23	26.44	10.3	6.9
<i>Abax parallelus</i>	1,2	68 (92)	78.95	84.21	1.5	10.3
<i>A. parallelepipedus</i>	1,2,3,4,5	63 (107)	39.68	33.33	34.9	1.6
<i>Carabus nemoralis</i>	1,2	18 (28)	72.22	38.89	0.0	0.0
<i>C. ullrichi</i>	2	13 (19)	53.85	7.69	0.0	0.0
<i>C. violaceus</i>	1,2,4,5	9 (19)	44.44	22.22	0.0	0.0
<i>Cychrus attenuates</i>	1,2	7 (10)	71.43	14.29	0.0	0.0
<i>Pterostichus madidus</i>	3,4,5	7 (16)	71.43	28.57	14.3	0.0
<i>Agonum sp</i>	5	6 (9)	16.67	33.33	0.0	0.0

<i>C. convexus</i>	2	6 (6)	83.33	0.0	0.0	0.0
<i>C. coriaceus</i>	2	6 (8)	83.33	33.33	0.0	0.0
<i>C. intricatus</i>	1,2	6 (7)	50.00	0.0	0.0	0.0
<i>Leistus fulvibarbis</i>	3,4	3 (6)	66.67	0.0	0.0	66.7
<i>Bembidion nigricorne</i>	3	2 (5)	100	0.0	0.0	0.0
<i>C. problematicus</i>	4	2 (2)	0.0	50.00	50.0	0.0
<i>Molops piceus</i>	2	2 (4)	100	50.00	0.0	0.0
<i>P. melanarius</i>	3,4	2 (2)	50.00	50.00	0.0	0.0
<i>P. transversalis</i>	1,2	2 (2)	50.00	50.00	0.0	0.0
<i>Synuchus vivalis</i>	3,4	2 (7)	100	0.0	50.0	0.0
<i>Aptinus bombardia</i>	1	1 (1)	100	0.0	0.0	100
<i>Notiophilus rufipes</i>	2	1 (2)	0.0	100	0.0	0.0

Table S3. Metal concentrations and pH values (mean±SD) in soil, earthworms (e), slugs (s) and carabids (c) at five locations. Concentrations are reported as mg/kg dry weight for lead and cadmium, and as µg/kg for mercury. Significant seasonal differences in metal concentrations at each study site are indicated with different superscripts; the same number with different letters denotes significantly different concentrations at the 0.05 level (Mann-Whitney U test). Abbreviations: Spring-summer (s), autumn (a)

Location	Groups	Season	N ind.	pH	Pb (mg/kg)	Cd (mg/kg)	Hg (µg/kg)
L1 (Cro)	c	s	10		0.455±0.40	0.784±0.81	105.1±76.97
	e	a	9		180.2±120.9	7.889±4.97	413.3±335
	e	s	22		168.9±92.52	8.182±3.54	361.5±169
	s	a	13		0.883±0.93	7.841±7.45	217.6 ^{1a} ±244.9
	s	s	6		0.748±1.38	6.662±7.74	57.88 ^{1b} ±103.6
	soil	a	11	3.75±0.14	67.91±15.3	0.565±0.208	310.9±152.4
	soil	s	11	3.80± 0.16	73.54±12.6	0.465±0.15	234.2±74.6
L2 (Cro)	c	a	13		0.406±0.41	1.194±0.79	491.7±264.6
	c	s	53		0.289±0.20	1.321±1.32	538.7±688.3
	e	a	16		75.45±74.98	15.13±9.20	1515±1420
	e	s	13		116±48.92	24.58±13.91	1075±686

	s	a	2		1.689±2.27	6.42±4.54	314.7±27.14
	s	s	3		16.78±28.07	33.75±22.91	115±35.30
	soil	a	11	4.96±0.17	127.2±71.9	1.557±0.62	442.2±422.9
	soil	s	11	4.96±0.13	139.7±60.06	1.369±0.53	617.8±415.2
L3 (UK)	c	a	22		26.17 ^{2a} ±25.17	9.480±18.21	123.2±59.48
	c	s	15		114.8 ^{2b} ±240.7	4.692±4.21	146.1±126.9
	e	a	17		2441±1154	174.3±132.4	708.5±681.9
	e	s	15		2446±1025	162.7±146.9	1790±2310
	s	a	25		227.5±138.2	122.3±79.71	193.8 ^{3a} ±149.7
	s	s	9		940.7±1092	84.98±38.11	478.3 ^{3b} ±490.9
	soil	a	12	5.73±0.10	8204±2406	24.27±13.25	590.2±311.2
	soil	s	14	5.86±0.15	8829±2830	24.84±15.2	669±290.7
L4 (UK)	c	a	45		8.020±7.18	8.989±9.97 ^{4a}	58.40±26.91
	c	s	12		4.628±3.15	2.752±3.27 ^{4b}	44.69±40.47
	e	a	9		294.1±132.2	101.7±76.13	414.1±511.4
	e	s	7		367.3±344.6	46.57±41.00	203.5±187

	s	a	8		36.31 ^{5a} ±17.52	53.27±29.60	118.3±61.92
	s	s	10		73.33 ^{5b} ±40.71	30.95±15.29	82.99±34.77
	soil	a	15	5.64±0.18	2525±2426	21.48±9.18	191.8±46.7
	soil	s	14	5.55±0.19	2630±1054	23.57±6.15	217.9±53
L5 (UK)	c	a	13		0.210 ^{6a} ±0.09	1.346±1.78	82.72±37.21
	c	s	25		0.457 ^{6b} ±0.56	0.776±0.64	129.7±87.91
	e	a	18		107.9 ^{7a} ±60.10	12.62±12.46	109.8±146.9
	e	s	9		33.63 ^{7b} ±22.6	11.08±7.73	70.46±47.06
	s	a	6		4.375±1.85	4.202±1.70	21.55±14.26
	s	s	14		5.092±2.92	5.524±3.03	30.15±21.61
	soil	a	8	4.52±0.10	69.11±18.8	0.713 ^{8a} ±0.28	76.2 ^{9a} ±17.6
	soil	s	16	4.12±0.10	56.05±12.7	0.466 ^{8b} ±0.15	170.2 ^{9b} ±43.9

Means at different study sites bearing a superscript with the same number but different letters differ significantly from each other at the 0.05 level. For example: 1a differs significantly from 1b. Values carrying different superscripted numbers do not differ significantly.

Table S4. Mean metal concentrations in ground beetles at five locations. Concentrations are expressed as mg/kg dry weight for lead and cadmium, and as µg/kg dry weight for mercury. *N*, the number of beetles used in the metal analysis.

Species	Site	<i>N</i>	Body weight (g dry weight)	Pb mg/kg	Cd mg/kg	Hg µg/kg
<i>Abax ater</i>	L2 (Cro)	23	0.132	0.379	1.382	718.1
	L3 (UK)	6	0.118	35.02	4.317	91.62
	L4 (UK)	3	0.123	5.517	3.559	84.27
	L5 (UK)	12	0.110	0.249	0.998	136.9
<i>A. parallelus</i>	L1 (Cro)	3	0.133	0.385	1.146	155.6
	L2 (Cro)	21	0.069	0.305	0.825	639.3
<i>Agonum sp.</i>	L5 (UK)	3	0.019	0.226	0.164	35.17
<i>Bembidion quadrimaculatum</i>	L3 (UK)	2	0.003	10.28	0.474	119.7
	L4 (UK)	1	0.004	15.55	0.144	16.43
	L5 (UK)	1	0.002	0.660	0.205	417.2

<i>Carabus cancellatus</i>	L4 (UK)	1	0.144	6.783	0.168	34.35
<i>C. coriaceus</i>	L2 (Cro)	2	0.128	1.780	0.823	0.252
<i>C. intricatus</i>	L1 (Cro)	1	0.336	0.399	0.591	95.15
<i>C. nemoralis</i>	L1 (Cro)	1	0.288	1.459	0.474	131.9
	L2 (Cro)	9	0.275	0.266	2.335	143.8
<i>C. ullrichi</i>	L2 (Cro)	6	0.293	0.084	1.245	370.9
<i>C. violaceus</i>	L1 (Cro)	3	0.247	0.351	1.016	76.65
	L2 (Cro)	1	0.299	0.053	1.831	590.2
	L4 (UK)	1	0.314	1.708	2.786	11.87
	L5 (UK)	5	0.256	0.279	0.862	54.03
<i>Cychrus attenuatus</i>	L1 (Cro)	1	0.075	0.200	0.054	83.67
	L2 (Cro)	2	0.088	0.486	1.531	159.0
<i>Cy. caraboides</i>	L5 (UK)	1	0.151	0.166	0.084	105.9
<i>Leistus fulvibarbis</i>	L4 (UK)	1	0.006	7.852	2.883	153.6
<i>Molops piceus</i>	L2 (Cro)	2	0.040	0.122	0.402	157.6
<i>Nebria brevicollis</i>	L3 (UK)	22	0.022	28.39	9.505	114.1

	L4 (UK)	46	0.019	7.455	8.672	55.62
	L5 (UK)	11	0.023	0.236	1.598	95.18
<i>Pterostichus</i>						
<i>fasciatopunctatus</i>	L1 (Cro)	1	0.064	0.284	0.238	43.42
<i>P. madidus</i>	L3 (UK)	4	0.041	79.69	7.510	162.6
	L5 (UK)	5	0.044	1.138	0.448	145.6
<i>Synuchus vivalis</i>	L3 (UK)	3	0.004	374.5	4.306	317.5
	L4 (UK)	2	0.006	5.823	1.652	18.02