Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/59379/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Crunelli, Vincenzo ORCID: <https://orcid.org/0000-0001-7154-9752>, David, Francois, Leresche, Nathalie and Lambert, Régis C. 2014. Role for T-type Ca^{2+} channels in sleep waves. Pflugers Archiv European Journal of Physiology 466 (4) , pp. 735-745. 10.1007/s00424-014-1477-3 file

Publishers page: <http://dx.doi.org/10.1007/s00424-014-1477-3>
<<http://dx.doi.org/10.1007/s00424-014-1477-3>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.

Role for T-type Ca²⁺ channels in sleep waves

▲ Vincenzo Crunelli^{1*}, Francois David¹, Nathalie Leresche^{2,3,4} and Régis C. Lambert^{2,3,4*}

Formatted: French (France)

¹School of Biosciences, Cardiff University, Museum Avenue, Cardiff CF10 3US, UK,
²Sorbonne Universités, UPMC Univ Paris 06, UM CR18, Neuroscience Paris Seine
(NPS), F-75005, Paris, France, ³CNRS, UMR 8246, NPS, F-75005, Paris, France,
⁴INSERM, U1130, NPS, F-75005, Paris, France

Short title: T-channels and sleep

Key words: cortex, thalamus, sleep slow oscillation, sleep spindles, delta waves, slow waves, theta waves, I_{Twindow}, I_h, I_{CAN}

▲ *Correspondence:
V. Crunelli (crunelli@cardiff.ac.uk), R. C. Lambert (regis.lambert@upmc.fr)

Formatted: French (France)

Formatted: French (France)

Formatted: French (France)

Field Code Changed

Formatted: French (France)

This article is published as part of the Special Issue on T-type channels

Abstract

Since their discovery more than thirty years ago, low-threshold T-type Ca^{2+} channels (T-channels) have been suggested to play a key role in many EEG waves of non-REM sleep, which has remained exclusively linked to the ability of these channels to generate low-threshold Ca^{2+} potentials and associated high-frequency bursts of action potentials. Our present understanding of the biophysics and physiology of T-channels, however, highlights a much more diverse and complex picture of the pivotal contributions that they make to different sleep rhythms. In particular, recent experimental evidence has conclusively demonstrated the essential contribution of thalamic T-channels to the expression of slow waves of natural sleep and the key role played by Ca^{2+} entry through these channels in the activation or modulation of other voltage-dependent channels that are important for the generation of both slow waves and sleep spindles. However, the precise contribution to sleep rhythms of T-channels in cortical neurons and other sleep-controlling neuronal networks remains unknown, and a full understanding of the cellular and network mechanisms of sleep delta waves is still lacking.

Introduction

As this review is part of a Special Issue on T-type Ca^{2+} channels (T-channels), it feels appropriate to firstly provide the reader with an overview of the stereotyped sequences of electrical waves that are recorded in the EEG during natural non-REM sleep. Importantly, while the source(s) of the electrical waves observed in scalp EEG recordings are located in the upper layers of the neocortex, their generator(s) are the dynamical interactions between the different neuronal network activities that are expressed by various component neurons of the corticothalamic loop. In humans, the occurrence of theta waves (3-7 Hz) over a generally desynchronized EEG characterizes the first stage of non-REM sleep, while in stage 2 occasional K-complexes and slow waves start to appear. Sleep spindles are also present in stage 2, either in isolation or associated with a K-complex. The EEG in stage 3 sleep still presents spindle episodes but also shows clearly defined periods of delta waves (0.5-4 Hz) that together with slow waves (<1-2 Hz) become the predominant activity as sleep deepens into stage 4. This smooth and progressive transition from stage 1 to stage 4 non-REM sleep is invariably accompanied at the neuronal level by a reduction in the depolarizing tone exerted by cholinergic, monoaminergic and histaminergic afferents from brain stem and mammillary body onto both cortical and thalamic neurones [60] (with some exceptions, see ref. [60]), leading to a progressive hyperpolarization of the majority of cortical and thalamic neurones [42, 78, 89].

The first insight into the role of T-channels in sleep waves came from their discovery in thalamic neurons, and in particular the finding that activation of these channels, following a period of membrane hyperpolarization, leads to a voltage

waveform known as the low threshold Ca^{2+} potential (LTCP) or low threshold spike [30, 32, 46, 47, 55]. Since then, the main and only widely recognized function of thalamic T-channels in sleep waves has been that of providing the rhythmic LTCP-mediated sequences of high frequency bursts of action potentials that characterize the cellular activity of these neurons during sleep spindles and delta waves as well as at the start of an Up state of sleep slow waves [7, 8, 45, 53, 54, 61, 76, 81, 82, 84, 92]. However, the role of the T-channels in sleep and non-REM EEG oscillations can no longer be restricted to the stereotypical LTCPs of thalamic neurons, since i) it involves other physiological voltage waveforms that are dependent on the “window current” of these channels (i.e. I_{Twindow}) [10, 43, 44, 91, 93], and ii) because non-thalamic neuronal populations, e.g. those in the neocortex and in sleep-controlling brain regions, show a marked expression of T-channels [28, 35, 38, 40, 41, 48, 71, 87]. This short review will address these issues after presenting a brief overview on the biophysics of T-channels, and in particular on I_{Twindow} and its physiological consequences for neuronal excitability (for detailed descriptions of the molecular genetics, biophysics and neuronal cell type distribution of T-channels, see other contributions to this Special Issue). In addition, as few experiments have so far analysed the role of T currents in naturally sleeping animals, we will also discuss how the current view of T-channel function in sleep may be clouded by the speculative extrapolations of data obtained either in brain slices or in anesthetized preparations where EEG waves similar, though not identical, to those observed during the various stages of natural non-REM sleep can be recorded (see [21]). Finally, we will highlight the current difficulty in correctly identifying the cellular and network

mechanisms of delta waves because of the partial overlap of their frequency band with that of slow waves.

Biophysics and physiological impact of the “window current” generated by T-channels

Since their original characterization in primary sensory neurons [11, 13, 36, 65], two main types of native T-type Ca^{2+} currents that display either “fast” or “slow” activation and inactivation kinetics were reported [45, 68]. Cloning of the three low-threshold Ca^{2+} channel genes (Cav3) further confirmed this crude categorization. Cav3.1 and Cav3.2 ($\alpha 1G$ and $\alpha 1H$, respectively) generate low-threshold Ca^{2+} currents displaying fast activation and inactivation mechanisms [18, 69], while Cav3.3 ($\alpha 1I$) shows much slower kinetics [52]. Regardless of this difference in gating kinetics, all native and recombinant channels share the same basic voltage-dependence with an activation threshold and a nearly complete steady-state inactivation around -60mV [68].

However, a closer look at the gating properties of the T-channels reveals that the steady-state activation and inactivation curves overlap (Fig. 1A). Therefore, in this voltage region that corresponds to neuronal resting membrane potentials, a few T-channels are not inactivated and their open probability is close but not equal to zero, hence a tonic T-current (i.e. I_{Twindow}) is generated. Since the activation and inactivation curves are obtained by fitting currents that in this voltage region are obviously very small, a precise estimation of I_{Twindow} , particularly for native channels, is difficult to achieve, and thus great caution should be used in interpreting these data. Nevertheless, investigations on recombinant channels have suggested that Cav3.3 channels may

generate a larger and more depolarized $I_{T\text{window}}$ (see Fig 7 in [68]; [15]) than that elicited by Cav3.1 and Cav3.2 channels. These differences in $I_{T\text{window}}$ should be carefully considered when assessing the precise role of this tonic current in the excitability of neurons that possess different complements (and different subcellular distributions) of the three isoforms of T-channels. In particular, the glutamatergic thalamocortical (TC) neurons only express Cav3.1 channels while the GABAergic neurons of the nucleus reticularis thalami (NRT) possess Cav3.3 channels in addition to a small component of Cav3.2 channels [45, 87].

Although $I_{T\text{window}}$ is an inherent biophysical property of all T-channels, its amplitude in some neurons may be too small, thus precluding a significant physiological role in cellular excitability. However, both TC and NRT neurons express especially large T-currents [6, 9, 31, 34] and a significant number of T-channels are still de-inactivated around -60mV. Thus, although the open probability of the channels is very low at these potentials, an $I_{T\text{window}}$ of about 30pA can be measured in TC neurons using voltage ramps that are slow enough to achieve steady-state equilibrium between activation and inactivation of the T-channels (Fig. 1B). Block of this tonic current with the specific T-channel antagonist, TTA-P2, induces a 3mV hyperpolarization of TC neurons held at -60mV but has no effect when the neuron is held at a membrane potential outside the voltage range of $I_{T\text{window}}$ activation (see Fig. 3 in [34]). As predicted from the biophysics of recombinant Cav3.3 channels, an even larger hyperpolarization (5mV) is observed upon application of TTA-P2 in NRT neurons [34], demonstrating that $I_{T\text{window}}$ play a crucial role in setting the resting membrane potential of both TC and NRT neurons. Moreover, when metabotropic glutamate or muscarinic receptors are activated the

interplay between the characteristic bell-shaped voltage-dependence of $I_{T\text{window}}$ (see inset in Fig. 1A) and the leak current creates a marked (up to 20mV) bistability of the resting membrane potential of TC and NRT neurons (Fig. 1C, middle trace) [20, 44]. The shift between these two stable membrane potentials can occur spontaneously as an intrinsic mechanism (leading to the appearance of repetitive Up and Down states of sleep slow waves, see next section) (Fig. 1C, top trace) [43] or can amplify small-amplitude subthreshold synaptic potentials leading to the generation of a rebound LTCP (see Fig. 6 in [93]).

Importantly, it is possible to block membrane bistability with the T-channel blocker TTA-P2 while leaving the LTCP and its associate firing almost intact (Fig. 1C, bottom trace). Indeed, Up and Down states quickly disappear upon a short period of TTA-P2 application that slightly reduces the functional T-channel population whereas the full block of LTCPs requires much longer antagonist application (see Fig. 7 in [34]). These two experiments clearly demonstrated that the high density of T-channels expressed in thalamic neurons far exceeds that required to generate a LTCP [9, 34]. Therefore, it is highly probable that such high density of T-channels provides the significant number of de-inactivated channels at depolarized potentials that are required for the full repertoire of the physiological responses of thalamic neurons. Finally, it is important to point out that these T-channels available at depolarized membrane potentials not only generate $I_{T\text{window}}$ in low-open probability conditions, but are also recruited by synaptic activities and intrinsic noise that, by mediating a drastic increase in open probability, generate additional transient T-currents that boost post-synaptic potentials (see Fig. 1 in [29]).

Thalamic and cortical T-channel contribution to sleep slow waves

Sleep slow waves are one of the fundamental EEG wave of non-REM sleep (Fig. 2A). They are present in almost all non-REM sleep stages [1-3, 37, 58, 75, 77], underlie sleep K-complexes [2, 3], and group together periods of sleep spindles [3, 62] and delta waves [2, 77]. The cellular counterpart of the sleep slow rhythm recorded in the EEG is the regular recurrence of a depolarized (Up) state and a hyperpolarized (Down) state of the membrane potential, that occurs synchronously in all cortical [77, 85, 86] and thalamic neurons [3, 16, 37, 58, 62, 75, 77, 79].

In cortical neurones, sleep slow waves result from intense excitatory and inhibitory synaptic barrages that generate the Up state and their absence that causes the Down state [73, 85, 86, 89]. Although T-currents are not considered to play a major role in this process, such powerful synaptic activity and the resulting changes in membrane potential do of course engage a variety of voltage-dependent channels, including T-channels. Indeed clear examples of LTCPs can be seen in recordings from cortical neurons during slow waves in anesthetised animals [16], as one would expect from the presence of i) all three T-channel isoforms in the neocortex [87] and ii) LTCPs in layers V-VI pyramidal neurons [28, 38, 41], as well as in somatostatin [48] and VIP [71] interneurons recorded in slices. It is surprising, therefore, that so far no study has directly investigated the role of T-channels in the activity of different cortical neurons during slow waves of non-REM sleep.

Because thalamic lesions do not suppress slow waves in anesthetized cats [85] and Up and Down states can be recorded in neocortical slices [17, 73] and in an isolated

cortical gyrus *in vivo* during anesthesia [88], these EEG slow waves were originally viewed as a cortically generated rhythm [12, 14, 88]. However, Up and Down states and slow waves similar to those observed *in vivo* can be recorded in thalamic slices [10, 21, 43], and a recent study has conclusively shown that selective block of thalamic firing by tetrodotoxin markedly reduces the frequency of EEG slow waves both in anesthetized and naturally sleeping rats [26]. Thus, a dynamic interplay between the synaptically driven neocortical oscillator and the thalamic oscillators of slow waves is necessary for the full expression of these waves of natural non-REM sleep.

As far as the thalamic oscillators are concerned many studies have clearly demonstrated that the T-channels of TC and NRT neurons contribute to the expression of sleep slow waves in three ways. Firstly, by evoking the LTCP (with high frequency burst of action potentials) that almost invariably marks the start of every Up state (Fig. 2C). Secondly (as mentioned in the previous section) by providing the membrane potential bistability that underlies the Up and Down state transitions, i.e. Up state = $I_{Twindow}$ “on” and Down state = $I_{Twindow}$ “off” (Fig. 2C) [19, 22]. Thirdly, by providing the selective Ca^{2+} entry that is required to activate i) the Ca^{2+} -activated, non-selective cation current (I_{CAN}), which tightly controls the durations of the Up states [43] (Fig. 2C), and ii) the Ca^{2+} -activated, K^+ channels (SK type) that contribute to the hyperpolarization that follows an LTCP [23, 24] (for a comprehensive biophysical description of these mechanisms and a list of the other currents contributing to sleep slow waves in thalamic neurons, see ref [22]).

On the basis of all these data, one would expect slow waves of natural sleep to be compromised in the absence of thalamic T-channels. Surprisingly, the original study in

mice with global knockout of the Cav3.1 isoform of the T-channels reported no change in EEG slow wave power [50]. Similar results were observed in mice with a “putative thalamic-selective” knockout of the same T-channel isoform, though recombination was also present in some cortical regions and hypothalamic nuclei [4]. However, the negative results of these two studies cannot be simply interpreted as indicating a lack of involvement of the Cav3.1 isoform in slow waves since i) compensation by other T-channel isoforms or other voltage- and transmitter-gated channels might have occurred in the thalamus of these two types of KO mice, and ii) Cav3.1 T-channels that are strongly expressed in brain areas other than the thalamus (see section below) were definitively knocked-out in these mice, with unpredictable consequences on slow waves and other sleep rhythms. Confirmation of an essential role for thalamic T-channels in sleep slow waves has finally been provided by experiments where optogenetics and neuronal ensemble recordings were combined with localized thalamic microdialysis injections of the selective T-channel antagonist TTA-P2 [26]. Thalamic dialysis concentration of TTA-P2 that fully blocks T-channel mediated burst firing produces a consistent reduction in the frequency of slow waves during anesthesia and natural non-REM sleep (Fig. 2A,B). In addition, block of thalamic T-channels suppresses the ability of selective optogenetic activation of TC neurons to entrain EEG slow waves [26]. These data, therefore, provide conclusive evidence on the essential role played by thalamic T-channels in slow wave of non-REM sleep.

Thalamic and cortical T-channel contribution to sleep spindles

A typical sleep spindle is a waxing and waning wave that lasts for a few seconds, has a frequency of 12-15 Hz in humans (but 8-12 Hz in rodents) and can occur in isolation from other sleep waves though it is mostly observed in close association with a K-complex (Fig. 3A1,A2) [3, 27, 62]. LTCPs are present at almost every cycle of the spindle wave in NRT neurons, and occasionally in TC neurones (Fig. 3A3,A4). The firing associated with the LTCPs generated by NRT neurons evokes GABA_A IPSPs in TC neurons, some of which provide enough time- and voltage-dependent removal of T-channel inactivation so that an LTCP (with or without the associated high-frequency burst) can then be generated. In turn, the LTCP-evoked firing of TC neurons elicits EPSPs in NRT neurons which help to trigger LTCPs at spindle frequency (Fig. 3A3,A4).

In addition to the LTCPs, another key contribution of T-channels to the electrical activity of TC neurons during sleep spindles is to provide the Ca²⁺ entry that regulates the cAMP-mediated up-regulation of I_h [56, 57]. It has been postulated that the potentiation of this inward current leads to a progressively larger depolarization of TC neurones during a spindle wave and ultimately to their inability to generate LTCPs, thus contributing to the spindle wave termination. A similar mechanism may occur in NRT neurons where the presence of HCN isoforms [63, 64] and I_h [10, 72] has now been demonstrated. Other roles of T-channels in NRT neurons during sleep spindles include the Ca²⁺ entry necessary to activate SK channels [24] and potentially I_{CAN} [10].

This well-accepted mechanism of spindle wave generation based on the recruitment of both TC and NRT T-channels has emerged from intracellular recordings in anesthetized animals (often after systemic injection of barbiturates to increase the occurrence of spindle waves) and in *in vitro* slice preparations. Therefore, caution should

be used when interpreting these results obtained in such different experimental conditions and extrapolating this mechanism to natural sleep spindles. Indeed, spindle waves tend not to occur in association with a K-complex under barbiturate anaesthesia, contrary to what is observed during ketamine/xylazine anaesthesia and natural sleep. In addition, the LTCPs of NRT neurons *in vivo* during spindle waves emerge from a depolarizing envelope, whereas spindle-like activity *in vitro* consists of LTCPs of increasing amplitude, each followed by a progressively larger afterhyperpolarization (Fig. 3A3).

Unfortunately, data obtained so far in naturally sleeping T-channel KO mice do not help to clarify the role of the different T-channel isoforms in the sleep spindle. Thus, genetic knockout of Cav3.1 channels (which abolishes T-current in all TC neurons) was originally reported to significantly decrease EEG power in the 8-10 Hz frequency band [50], although some remaining spindles showing a reduced amplitude were still present, whereas a later study by the same group reported no effect when spindle events were filtered at 6-15 Hz [51]. In Cav3.3 KO mice, no difference in the 10-12 Hz EEG power was observed when compared to wild type animals [6]. Only when the analysis was restricted to periods of transitions from non-NREM to REM sleep (i.e. when sleep spindles are more prominent), a reduction of the EEG power in the 10-12 Hz frequency band measured about 30s before REM sleep onset was observed [6]. Since in NRT neurons LTCPs are observed at every cycle of spindle waves both *in vivo* [30, 39] and *in vitro* [92] (see Fig. 3A3), the absence of a clear effect of genetically deleting Cav3.3 channels (the main isoform present in NRT neurons) on this sleep rhythm is highly surprising. These contradictory results may once again be in part explained by compensatory mechanisms that occur in these T-channel isoform KO mice but may also

result from difficulties in clearly identifying spindle episodes in the EEG of naturally sleeping mice. Indeed, although local field potential recordings in deep cortical areas reveal a comparable profile of spindles in humans and mice, spindles are not clearly apparent in EEG traces from mice and their identification require sophisticated analysis to assess EEG spectral changes at the level of individual sleep stage transitions [5, 94]. Notwithstanding these contrasting results in transgenic mice, recent experiments in rats, using microdialysis of TTA-P2 in the somatosensory thalamus and NRT, provide conclusive evidence of a drastic decrease of spindle waves both during anesthesia and natural sleep [26], confirming the key role of thalamic T-channels in the generation of this sleep rhythm.

Finally, since many cortical neurones possess a vast repertoire of T-channels (see previous section) and the waveform of cortical spindle waves spans the voltage region of T-channel activation/inactivation, a contribution of these cortical channels to the fine tuning of EEG sleep spindles would be expected, although to the best of our knowledge this has not so far been rigorously investigated.

Thalamic and cortical T-channel contribution to delta waves

In TC neurons, membrane potential oscillations at delta frequency (0.5-4 Hz) that consist of rhythmically occurring LTCPs were the first T-channel-dependent activity whose mechanism was fully elucidated *in vitro* [53, 54, 61, 76]. This work, together with *in vivo* studies in anesthetized animals [33, 66, 67, 83], strongly suggested that delta oscillations in TC neurones are fully determined in a pacemaker fashion by the time- and

voltage-dependencies of the h, T and K⁺ channels, in both TC and NRT neurons [10, 54, 61] stressing the key role of thalamic T-channels in the generation of these waves.

One important issue that has often been overlooked, however, is that the majority of EEG delta waves of natural sleep, and its thalamic counterpart the delta oscillation, do not occur in very long periods, as a somewhat inaccurate interpretation of the initial *in vitro* studies (Fig. 3B2, top trace) might lead to conclude. Indeed, thalamic delta oscillations appear to occur mostly in discrete groups during the Down state of slow waves in both TC and NRT neurons (Fig. 3B2, bottom trace) [43, 79, 90]. We are unaware of any evidence supporting the presence of long period of delta oscillations in TC neurons *in vivo* during natural sleep, raising the question of whether these long sequences of repetitive LTCPs are only observed in thalamic slices. An additional point of concern when interpreting EEG data is the ambiguity in the definition of delta waves of natural sleep and their potential overlap with slow waves. In other words, it is possible that EEG delta waves at the lower end of their frequency range (i.e. 0.5-2 Hz) may correspond to a thalamic (and cortical) cellular activity characterized by Up and Down states (i.e. slow waves) and not by repetitive LTCPs (compare Fig. 2A and Fig. 3B1).

Together with compensatory mechanisms (as highlighted in previous section), the above two issues might explain the contradictory results on delta waves obtained in mice with genetic ablation of T-channels. Thus, total KO of the Cav3.1 isoform was shown to induce a marked decrease in the power density of delta frequency band (selected as 2-6.5 Hz) [50] whereas in the “putative thalamic-selective” Cav3.1 KO mouse there is a moderate increase in EEG spectral power within the delta frequency range (selected as 1-4Hz) [4]. Moreover, no change in delta frequency power was observed in global Cav3.3

KO mice [6], while the impact of deleting the Cav3.2 isoform on sleep rhythms has not yet been analyzed. To complicate this picture further, systemic injection of selective antagonists of all T-channel isoforms has been shown to dose-dependently increase delta waves and reduce slow waves in wild-type mice and rats, respectively [26, 49].

As far as cortical T-channels are concerned, the firing input at delta frequency from TC to cortical neurones might clearly play an important contribution to the expression of delta waves of natural sleep in the EEG. However, membrane potential oscillations at delta frequency have been observed in cortical neurones [77] and the full mechanisms of the delta waves (i.e. the relative contribution of synaptic and intrinsic conductances, including a precise role for T-channels) in cortical neurons remains to be fully elucidated.

Contribution of T-channels in other brain regions to the expression of sleep waves

The difficulty in the interpretation of sleep studies using global, genetic or pharmacological T-channel block that we have outlined in the previous sections clearly stems from our lack of knowledge of the contribution of T-channels not only in cortical neurons but also in other brain networks that control and/or modulate natural sleep. In fact, neurons expressing LTCPs are present in the ventrolateral preoptic nucleus [40] and in the lateral hypothalamus (i.e. hypocretin-orexin expressing neurons) [35], two strongly interacting regions that belong, respectively, to the sleep-promoting and ascending arousal pathways [74] (Fig. 4). The potential presence of T-channels mediated electrical events (e.g. LTCPs, $I_{Twindow}$, T-channel dependent Ca^{2+} modulation or activation of various voltage-gated channels, etc.) in neurons of these pathways should therefore be

investigated to gain a full understanding of the role of T-channels in sleep. In this respect, it is clear that manipulating T-channels not only has an impact on various sleep rhythms but also on sleep architecture and the transitions between wake and sleep as well as between non-REM and REM sleep. Both global and “putative thalamic-selective” Cav3.1 KO mice (but not cortical KO mice) show an increased number of frequent brief awakenings that interrupt non-REM sleep and a delayed sleep onset [4, 50]. Moreover, systemic injection of TTA-A2/P2 in wild-type mice acutely reduces the mean time spent in active wake [49], and induces dose-dependent behavioral and EEG changes indicative of sedation/sleep in rats [59]

Conclusions

After 30 years since the discovery of T-channels in thalamic neurons, we have a very comprehensive view of their precise contribution to slow waves and spindles of natural sleep, including the role of LTCPs and $I_{Twindow}$. A similar understanding of delta waves is still missing, in part because of lack of appropriate studies and uncertainties on an appropriate classification of these waves. Moreover, we still know very little about the contribution of T-channels in neurons of the neocortex and other neuronal networks involved in sleep. Undoubtedly, the development of selective blockers for different T-channel isoforms and of conditional and area-selective KO animals will contribute to unravel the full role played by these widely expressed voltage-gated channels in sleep waves and architecture.

Acknowledgments

VC work in this field is supported by The Wellcome Trust (grant 91882), NL and RCL work by Agence Nationale de la Recherche (grant MNMP-2009) and CNRS (LEA 528).

References

1. Achermann P, Borbely AA (1997) Low-frequency (< 1 Hz) oscillations in the human sleep electroencephalogram. *Neuroscience* 81:213-22 DOI Electronic Resource Number
2. Amzica F, Steriade M (1997) The K-complex: its slow (<1-Hz) rhythmicity and relation to delta waves. *Neurology* 49:952-9 DOI Electronic Resource Number
3. Amzica F, Steriade M (2002) The functional significance of K-complexes. *Sleep medicine reviews* 6:139-49 DOI Electronic Resource Number
4. Anderson MP, Mochizuki T, Xie J, Fischler W, Manger JP, Talley EM, Scammell TE, Tonegawa S (2005) Thalamic Cav3.1 T-type Ca²⁺ channel plays a crucial role in stabilizing sleep. *Proceedings of the National Academy of Sciences of the United States of America* 102:1743-8 DOI Electronic Resource Number
5. Astori S, Wimmer RD, Luthi A (2013) Manipulating sleep spindles--expanding views on sleep, memory, and disease. *Trends in neurosciences* 36:738-48 DOI Electronic Resource Number
6. Astori S, Wimmer RD, Prosser HM, Corti C, Corsi M, Liaudet N, Volterra A, Franken P, Adelman JP, Luthi A (2011) The Ca(V)_{3.3} calcium channel is the major sleep spindle pacemaker in thalamus. *Proceedings of the National Academy of Sciences of the United States of America* 108:13823-8 DOI Electronic Resource Number
7. Bal T, von Krosigk M, McCormick DA (1995) Role of the ferret perigeniculate nucleus in the generation of synchronized oscillations in vitro. *J Physiol (Lond)* 483:665-85 DOI Electronic Resource Number
8. Bal T, von Krosigk M, McCormick DA (1995) Synaptic and membrane mechanisms underlying synchronized oscillations in the ferret lateral geniculate nucleus in vitro. *J Physiol (Lond)* 483:641-63 DOI Electronic Resource Number
9. Bessaih T, Leresche N, Lambert RC (2008) T current potentiation increases the occurrence and temporal fidelity of synaptically evoked burst firing in sensory thalamic neurons. *Proceedings of the National Academy of Sciences of the United States of America* 105:11376-81 DOI Electronic Resource Number
10. Blethyn KL, Hughes SW, Tóth TI, Cope DW, Crunelli V (2006) Neuronal basis of the slow (<1Hz) oscillation in neurons of the nucleus reticularis thalami in vitro. *J Neurosci* 26:2474-86 DOI Electronic Resource Number
11. Bossu JL, Feltz A (1986) Inactivation of the low-threshold transient calcium current in rat sensory neurones: evidence for a dual process. *J Physiol* 376:341-57 DOI Electronic Resource Number
12. Brown RE, Basheer R, McKenna JT, Strecker RE, McCarley RW (2012) Control of sleep and wakefulness. *Physiological reviews* 92:1087-187 DOI Electronic Resource Number
13. Carbone E, Lux HD (1984) A low voltage-activated, fully inactivating Ca channel in vertebrate sensory neurones. *Nature* 310:501-2 DOI Electronic Resource Number
14. Chauvette S, Crochet S, Volgushev M, Timofeev I (2011) Properties of slow oscillation during slow-wave sleep and anesthesia in cats. *The Journal of neuroscience* :

the official journal of the Society for Neuroscience 31:14998-5008 DOI Electronic Resource Number

15. Chemin J, Monteil A, Perez-Reyes E, Bourinet E, Nargeot J, Lory P (2002) Specific contribution of human T-type calcium channel isoforms ($\alpha 1G$, $\alpha 1H$ and $\alpha 1I$) to neuronal excitability. *J Physiol* 540:3-14. DOI Electronic Resource Number

16. Contreras D, Steriade M (1995) Cellular basis of EEG slow rhythms: a study of dynamic corticothalamic relationships. *J Neurosci* 15:604-22 DOI Electronic Resource Number

17. Cossart R, Aronov D, Yuste R (2003) Attractor dynamics of network UP states in the neocortex. *Nature* 423:283-8 DOI Electronic Resource Number

18. Cribbs LL, Lee JH, Yang J, Satin J, Zhang Y, Daud A, Barclay J, Williamson MP, Fox M, Rees M, Perez-Reyes E (1998) Cloning and characterization of $\alpha 1H$ from human heart, a member of the T-type Ca^{2+} channel gene family. *Circ Res* 83:103-9. DOI Electronic Resource Number

19. Crunelli V, Blethyn KL, Cope DW, Hughes SW, Parri HR, Turner JP, Toth TI, Williams SR (2002) Novel neuronal and astrocytic mechanisms in thalamocortical loop dynamics. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* 357:1675-93 DOI Electronic Resource Number

20. Crunelli V, Cope DW, Hughes SW (2006) Thalamic T-type Ca^{2+} channels and NREM sleep. *Cell Calcium* 40:175-90 DOI Electronic Resource Number

21. Crunelli V, Hughes SW (2010) The slow (<1 Hz) rhythm of non-REM sleep: a dialogue between three cardinal oscillators. *Nature neuroscience* 13:9-17 DOI Electronic Resource Number

22. Crunelli V, Toth TI, Cope DW, Blethyn K, Hughes SW (2005) The 'window' T-type calcium current in brain dynamics of different behavioural states. *The Journal of physiology* 562:121-9 DOI Electronic Resource Number

23. Cueni L, Canepari M, Adelman JP, Luthi A (2009) Ca^{2+} signaling by T-type Ca^{2+} channels in neurons. *Pflügers Archiv : European journal of physiology* 457:1161-72 DOI Electronic Resource Number

24. Cueni L, Canepari M, Lujan R, Emmenegger Y, Watanabe M, Bond CT, Franken P, Adelman JP, Luthi A (2008) T-type Ca^{2+} channels, SK2 channels and SERCAs gate sleep-related oscillations in thalamic dendrites. *Nature neuroscience* 11:683-92 DOI Electronic Resource Number

25. Cvetkovic-Lopes V, Eggermann E, Uschakov A, Grivel J, Bayer L, Jones BE, Serafin M, Muhlethaler M (2010) Rat hypocretin/orexin neurons are maintained in a depolarized state by TRPC channels. *PloS one* 5:e15673 DOI Electronic Resource Number

26. David F, Schmiedt JT, Taylor HL, Orban G, Di Giovanni G, Uebele VN, Renger JJ, Lambert RC, Leresche N, Crunelli V (2013) Essential thalamic contribution to slow waves of natural sleep. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 33:19599-610 DOI Electronic Resource Number

27. De Gennaro L, Ferrara M (2003) Sleep spindles: an overview. *Sleep medicine reviews* 7:423-40 DOI Electronic Resource Number

28. de la Pena E, Geijo-Barrientos E (1996) Laminar localization, morphology, and physiological properties of pyramidal neurons that have the low-threshold calcium current in the guinea-pig medial frontal cortex. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 16:5301-11 DOI Electronic Resource Number

29. Deleuze C, David F, Behuret S, Sadoc G, Shin HS, Uebele VN, Renger JJ, Lambert RC, Leresche N, Bal T (2012) T-type calcium channels consolidate tonic action potential output of thalamic neurons to neocortex. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 32:12228-36 DOI Electronic Resource Number
30. Deschenes M, Paradis M, Roy JP, Steriade M (1984) Electrophysiology of neurons of lateral thalamic nuclei in cat: resting properties and burst discharges. *J Neurophysiol* 51:1196-219 DOI Electronic Resource Number
31. Destexhe A, Neubig M, Ulrich D, Huguenard J (1998) Dendritic low-threshold calcium currents in thalamic relay cells. *J Neurosci* 18:3574-88 DOI Electronic Resource Number
32. Domich L, Oakson G, Steriade M (1986) Thalamic burst patterns in the naturally sleeping cat: a comparison between cortically projecting and reticularis neurones. *J Physiol (Lond)* 379:429-49 DOI Electronic Resource Number
33. Dossi RC, Nunez A, Steriade M (1992) Electrophysiology of a slow (0.5-4 Hz) intrinsic oscillation of cat thalamocortical neurones in vivo. *J Physiol (Lond)* 447:215-34 DOI Electronic Resource Number
34. Dreyfus FM, Tscherter A, Errington AC, Renger JJ, Shin HS, Uebele VN, Crunelli V, Lambert RC, Leresche N (2010) Selective T-type calcium channel block in thalamic neurons reveals channel redundancy and physiological impact of I(T)window. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 30:99-109 DOI Electronic Resource Number
35. Eggermann E, Bayer L, Serafin M, Saint-Mieux B, Bernheim L, Machard D, Jones BE, Muhlethaler M (2003) The wake-promoting hypocretin-orexin neurons are in an intrinsic state of membrane depolarization. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 23:1557-62 DOI Electronic Resource Number
36. Fedulova SA, Kostyuk PG, Veselovsky NS (1985) Two types of calcium channels in the somatic membrane of new-born rat dorsal root ganglion neurones. *The Journal of physiology* 359:431-46 DOI Electronic Resource Number
37. Ferri R, Rundo F, Bruni O, Terzano MG, Stam CJ (2005) Dynamics of the EEG slow-wave synchronization during sleep. *Clinical neurophysiology : official journal of the International Federation of Clinical Neurophysiology* 116:2783-95 DOI Electronic Resource Number
38. Friedman A, Gutnick MJ (1987) Low-threshold calcium electrogenesis in neocortical neurons. *Neuroscience letters* 81:117-22 DOI Electronic Resource Number
39. Fuentealba P, Timofeev I, Steriade M (2004) Prolonged hyperpolarizing potentials precede spindle oscillations in the thalamic reticular nucleus. *Proceedings of the National Academy of Sciences of the United States of America* 101:9816-21 DOI Electronic Resource Number
40. Gallopin T, Fort P, Eggermann E, Cauli B, Luppi PH, Rossier J, Audinat E, Muhlethaler M, Serafin M (2000) Identification of sleep-promoting neurons in vitro. *Nature* 404:992-5 DOI Electronic Resource Number
41. Hamill OP, Huguenard JR, Prince DA (1991) Patch-clamp studies of voltage-gated currents in identified neurons of the rat cerebral cortex. *Cereb Cortex* 1:48-61 DOI Electronic Resource Number

42. Hirsch JC, Fourment A, Marc ME (1983) Sleep-related variations of membrane potential in the lateral geniculate body relay neurons of the cat. *Brain research* 259:308-12 DOI Electronic Resource Number
43. Hughes SW, Cope DW, Blethyn KL, Crunelli V (2002) Cellular mechanisms of the slow (1 Hz) oscillation in thalamocortical neurons in vitro. *Neuron* 33:947-58. DOI Electronic Resource Number
44. Hughes SW, Cope DW, Toth TI, Williams SR, Crunelli V (1999) All thalamocortical neurones possess a T-type Ca^{2+} 'window' current that enables the expression of bistability-mediated activities. *J Physiol (Lond)* 517:805-15 DOI Electronic Resource Number
45. Huguenard JR, Prince DA (1992) A novel T-type current underlies prolonged Ca^{2+} -dependent burst firing in GABAergic neurons of rat thalamic reticular nucleus. *J Neurosci* 12:3804-17 DOI Electronic Resource Number
46. Jahnsen H, Llinas R (1984) Electrophysiological properties of guinea-pig thalamic neurones: an in vitro study. *J Physiol* 349:205-26 DOI Electronic Resource Number
47. Jahnsen H, Llinas R (1984) Ionic basis for the electro-responsiveness and oscillatory properties of guinea-pig thalamic neurones in vitro. *J Physiol* 349:227-47 DOI Electronic Resource Number
48. Kawaguchi Y, Kubota Y (1993) Correlation of physiological subgroupings of nonpyramidal cells with parvalbumin- and calbindinD28k-immunoreactive neurons in layer V of rat frontal cortex. *Journal of neurophysiology* 70:387-96 DOI Electronic Resource Number
49. Kraus RL, Li Y, Gregan Y, Gotter AL, Uebele VN, Fox SV, Doran SM, Barrow JC, Yang ZQ, Reger TS, Koblan KS, Renger JJ (2010) In vitro characterization of T-type calcium channel antagonist TTA-A2 and in vivo effects on arousal in mice. *The Journal of pharmacology and experimental therapeutics* 335:409-17 DOI Electronic Resource Number
50. Lee J, Kim D, Shin HS (2004) Lack of delta waves and sleep disturbances during non-rapid eye movement sleep in mice lacking $\alpha 1\text{G}$ -subunit of T-type calcium channels. *Proc Natl Acad Sci U S A* 101:18195-9 DOI Electronic Resource Number
51. Lee J, Song K, Lee K, Hong J, Lee H, Chae S, Cheong E, Shin HS (2013) Sleep spindles are generated in the absence of T-type calcium channel-mediated low-threshold burst firing of thalamocortical neurons. *Proceedings of the National Academy of Sciences of the United States of America* 110:20266-71 DOI Electronic Resource Number
52. Lee JH, Daud AN, Cribbs LL, Lacerda AE, Pereverzev A, Klockner U, Schneider T, Perez-Reyes E (1999) Cloning and expression of a novel member of the low voltage-activated T-type calcium channel family. *J Neurosci* 19:1912-21. DOI Electronic Resource Number
53. Leresche N, Jassik-Gerschenfeld D, Haby M, Soltesz I, Crunelli V (1990) Pacemaker-like and other types of spontaneous membrane potential oscillations of thalamocortical cells. *Neurosci Lett* 113:72-7 DOI Electronic Resource Number
54. Leresche N, Lightowler S, Soltesz I, Jassik-Gerschenfeld D, Crunelli V (1991) Low-frequency oscillatory activities intrinsic to rat and cat thalamocortical cells. *J Physiol (Lond)* 441:155-74 DOI Electronic Resource Number
55. Llinas R, Jahnsen H (1982) Electrophysiology of mammalian thalamic neurones in vitro. *Nature* 297:406-8 DOI Electronic Resource Number

56. Lüthi A, McCormick DA (1998) Periodicity of thalamic synchronized oscillations: the role of Ca²⁺- mediated upregulation of Ih. *Neuron* 20:553-63 DOI Electronic Resource Number
57. Lüthi A, McCormick DA (1999) Modulation of a pacemaker current through Ca(2+)-induced stimulation of cAMP production. *Nat Neurosci* 2:634-41 DOI Electronic Resource Number
58. Massimini M, Huber R, Ferrarelli F, Hill S, Tononi G (2004) The sleep slow oscillation as a traveling wave. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 24:6862-70 DOI Electronic Resource Number
59. McCafferty C, David F, Venzi M, Di Giovanni G, Orban G, Uebele VN, Renger JJ, Lambert RC, Leresche N, V C (2012) T-type calcium channels of cortical and thalamocortical neurons are necessary for absence seizures. *Society for Neuroscience*. New Orleans USA
60. McCormick DA (1992) Neurotransmitter actions in the thalamus and cerebral cortex and their role in neuromodulation of thalamocortical activity. *Prog Neurobiol* 39:337-88 DOI Electronic Resource Number
61. McCormick DA, Pape HC (1990) Properties of a hyperpolarization-activated cation current and its role in rhythmic oscillation in thalamic relay neurones. *J Physiol (Lond)* 431:291-318 DOI Electronic Resource Number
62. Molle M, Marshall L, Gais S, Born J (2002) Grouping of spindle activity during slow oscillations in human non-rapid eye movement sleep. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 22:10941-7 DOI Electronic Resource Number
63. Monteggia LM, Eisch AJ, Tang MD, Kaczmarek LK, Nestler EJ (2000) Cloning and localization of the hyperpolarization-activated cyclic nucleotide-gated channel family in rat brain. *Brain research. Molecular brain research* 81:129-39 DOI Electronic Resource Number
64. Notomi T, Shigemoto R (2004) Immunohistochemical localization of Ih channel subunits, HCN1-4, in the rat brain. *The Journal of comparative neurology* 471:241-76 DOI Electronic Resource Number
65. Nowycky MC, Fox AP, Tsien RW (1985) Three types of neuronal calcium channel with different calcium agonist sensitivity. *Nature* 316:440-3 DOI Electronic Resource Number
66. Nunez A, Amzica F, Steriade M (1992) Intrinsic and synaptically generated delta (1-4 Hz) rhythms in dorsal lateral geniculate neurons and their modulation by light-induced fast (30-70 Hz) events. *Neuroscience* 51:269-84 DOI Electronic Resource Number
67. Nunez A, Curro Dossi R, Contreras D, Steriade M (1992) Intracellular evidence for incompatibility between spindle and delta oscillations in thalamocortical neurons of cat. *Neuroscience* 48:75-85 DOI Electronic Resource Number
68. Perez-Reyes E (2003) Molecular physiology of low-voltage-activated t-type calcium channels. *Physiol Rev* 83:117-61 DOI Electronic Resource Number
69. Perez-Reyes E, Cribbs LL, Daud A, Lacerda AE, Barclay J, Williamson MP, Fox M, Rees M, Lee JH (1998) Molecular characterization of a neuronal low-voltage-activated T-type calcium channel. *Nature* 391:896-900. DOI Electronic Resource Number
70. Pirchio M, Turner JP, Williams SR, Asproдини E, Crunelli V (1997) Postnatal development of membrane properties and delta oscillations in thalamocortical neurons of

the cat dorsal lateral geniculate nucleus. *J Neurosci* 17:5428-44 DOI Electronic Resource Number

71. Porter JT, Cauli B, Tsuzuki K, Lambolez B, Rossier J, Audinat E (1999) Selective excitation of subtypes of neocortical interneurons by nicotinic receptors. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 19:5228-35 DOI Electronic Resource Number

72. Rateau Y, Ropert N (2006) Expression of a functional hyperpolarization-activated current (I_h) in the mouse nucleus reticularis thalami. *Journal of neurophysiology* 95:3073-85 DOI Electronic Resource Number

73. Sanchez-Vives MV, McCormick DA (2000) Cellular and network mechanisms of rhythmic recurrent activity in neocortex. *Nature neuroscience* 3:1027-34 DOI Electronic Resource Number

74. Saper CB, Fuller PM, Pedersen NP, Lu J, Scammell TE (2010) Sleep state switching. *Neuron* 68:1023-42 DOI Electronic Resource Number

75. Simon NR, Manshanden I, Lopes da Silva FH (2000) A MEG study of sleep. *Brain research* 860:64-76 DOI Electronic Resource Number

76. Soltesz I, Lightowler S, Leresche N, Jassik-Gerschenfeld D, Pollard CE, Crunelli V (1991) Two inward currents and the transformation of low-frequency oscillations of rat and cat thalamocortical cells. *J Physiol (Lond)* 441:175-97 DOI Electronic Resource Number

77. Steriade M (2003) The corticothalamic system in sleep. *Frontiers in bioscience : a journal and virtual library* 8:d878-99 DOI Electronic Resource Number

78. Steriade M, Amzica F, Nunez A (1993) Cholinergic and noradrenergic modulation of the slow (approximately 0.3 Hz) oscillation in neocortical cells. *J Neurophysiol* 70:1385-400 DOI Electronic Resource Number

79. Steriade M, Contreras D, Curro Dossi R, Nunez A (1993) The slow (< 1 Hz) oscillation in reticular thalamic and thalamocortical neurons: scenario of sleep rhythm generation in interacting thalamic and neocortical networks. *J Neurosci* 13:3284-99 DOI Electronic Resource Number

80. Steriade M, Deschênes M (1998) Cellular thalamic mechanismsElsevier, Amsterdam

81. Steriade M, Deschenes M, Domich L, Mulle C (1985) Abolition of spindle oscillations in thalamic neurons disconnected from nucleus reticularis thalami. *J Neurophysiol* 54:1473-97 DOI Electronic Resource Number

82. Steriade M, Domich L, Oakson G, Deschenes M (1987) The deafferented reticular thalamic nucleus generates spindle rhythmicity. *J Neurophysiol* 57:260-73 DOI Electronic Resource Number

83. Steriade M, Dossi RC, Nunez A (1991) Network modulation of a slow intrinsic oscillation of cat thalamocortical neurons implicated in sleep delta waves: cortically induced synchronization and brainstem cholinergic suppression. *J Neurosci* 11:3200-17 DOI Electronic Resource Number

84. Steriade M, McCormick DA, Sejnowski TJ (1993) Thalamocortical oscillations in the sleeping and aroused brain. *Science* 262:679-85 DOI Electronic Resource Number

85. Steriade M, Nunez A, Amzica F (1993) Intracellular analysis of relations between the slow (< 1 Hz) neocortical oscillation and other sleep rhythms of the electroencephalogram. *J Neurosci* 13:3266-83 DOI Electronic Resource Number

86. Steriade M, Nunez A, Amzica F (1993) A novel slow (< 1 Hz) oscillation of neocortical neurons in vivo: depolarizing and hyperpolarizing components. *J Neurosci* 13:3252-65 DOI Electronic Resource Number
87. Talley EM, Cribbs LL, Lee JH, Daud A, Perez-Reyes E, Bayliss DA (1999) Differential distribution of three members of a gene family encoding low voltage-activated (T-type) calcium channels. *J Neurosci* 19:1895-911 DOI Electronic Resource Number
88. Timofeev I, Grenier F, Bazhenov M, Sejnowski TJ, Steriade M (2000) Origin of slow cortical oscillations in deafferented cortical slabs. *Cerebral cortex* 10:1185-99 DOI Electronic Resource Number
89. Timofeev I, Grenier F, Steriade M (2001) Disfacilitation and active inhibition in the neocortex during the natural sleep-wake cycle: an intracellular study. *Proceedings of the National Academy of Sciences of the United States of America* 98:1924-9 DOI Electronic Resource Number
90. Timofeev I, Steriade M (1996) Low-frequency rhythms in the thalamus of intact-cortex and decorticated cats. *J Neurophysiol* 76:4152-68 DOI Electronic Resource Number
91. Toth TI, Hughes SW, Crunelli V (1998) Analysis and biophysical interpretation of bistable behaviour in thalamocortical neurons. *Neuroscience* 87:519-23 DOI Electronic Resource Number
92. von Krosigk M, Bal T, McCormick DA (1993) Cellular mechanisms of a synchronized oscillation in the thalamus. *Science* 261:361-4 DOI Electronic Resource Number
93. Williams SR, Toth TI, Turner JP, Hughes SW, Crunelli V (1997) The 'window' component of the low threshold Ca²⁺ current produces input signal amplification and bistability in cat and rat thalamocortical neurones. *J Physiol (Lond)* 505:689-705 DOI Electronic Resource Number
94. Wimmer RD, Astori S, Bond CT, Rovo Z, Chatton JY, Adelman JP, Franken P, Luthi A (2012) Sustaining sleep spindles through enhanced SK2-channel activity consolidates sleep and elevates arousal threshold. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 32:13917-28 DOI Electronic Resource Number

Figure 1. The window T-current tightly controls thalamic neuron excitability.

A. Normalized activation and steady-state inactivation curves of T-current recorded in a TC neuron from the rat ventrobasal thalamic nucleus. The activation curve was constructed by successive step depolarizations from -80mV to -45mV (2.5mV increments) preceded by a 1s hyperpolarizing pre-pulse to -100mV. Inactivation of the T-channels was induced using a 1s pre-pulse of increasing potential (from -100mV to -60mV with 2.5mV increments) and the resulting channel availability was estimated from the normalized current amplitude measured at -50mV. Data were fitted by Boltzman equations. Inset illustrates the voltage-dependence of the steady-state channel activation (window current) estimated from the product of the Boltzman fits of the normalized activation and inactivation curves.

B1. The window T-current evoked by a 10-s-long depolarizing voltage ramp from -100 to -40 mV preceded by a 1 s hyperpolarizing prepulse to -100 mV is fully blocked by the selective T-channel blocker TTA-P2 (1 μ M). **B2.** Voltage-dependence of the window T-current shown in B1.

C. In the continuous presence of trans-ACPD, recording from a cat TC neuron in an thalamic intralaminar nucleus reveals a slow oscillation (top trace) consisting of regularly recurring Up and Down states intermixed with much longer Up states with continuous tonic action potentials firing. Each Down state starts with a clear inflection point leading to a stereotypical large hyperpolarizing potential that, upon I_h activation, slowly repolarizes the neuron up to the LTCP-threshold (see also Fig. 2C). Following the block of I_h with ZD 7288 (middle trace), the neuron exhibits two stable resting membrane potentials. Transitions between stable equilibrium potentials are evoked by short steps of

positive or negative injected currents (I_{inj}) that trigger an LTCP and switch off $I_{Twindow}$, respectively. Upon application of TTA-P2 that progressively blocks the T-channel population, the bistable behaviour quickly disappears due to the decrease in $I_{Twindow}$ while enough T-channels remain to evoke LTCPs (lower trace). B1-B2: reproduced with permission from [34]

Figure 2. Contribution of thalamic T-channels to sleep slow waves.

A. EEG slow wave and corresponding wavelet scalograms recorded in a rat during natural sleep. Continuous thalamic microdialysis injection of a solution containing the T-channel antagonist TTA-P2 (3 mM) reduces the frequency of slow waves compared to the injection of artificial cerebrospinal fluid (Control).

B. EEG slow wave recorded in a rat during ketamine/kylazine anesthesia. Continuous thalamic microdialysis injection of a solution containing 3 mM of TTA-P2 decreases the frequency of slow waves compared to the injection of artificial cerebrospinal fluid (Control).

C. Schematic diagram of two cycles of slow waves in a TC neuron shows the various contributions of T-channels to this oscillation.

A-C: reproduced with permission from [26] and [20].

Figure 3. Contribution of thalamic T-channels to sleep spindles and comparison of delta waves recorded *in vivo* and *in vitro*.

A1. Filtered (5-15 Hz) (bottom trace) and unfiltered (upper trace) examples of a spindle recorded in a rat during natural sleep (graph shows the corresponding wavelet scalogram). Note the occurrence of the spindle immediately after a Down state.

A2. Spindles recorded in a cat during natural light sleep and anesthesia (top and bottom set of traces, respectively). Note the association with a K-complex in the boxed spindles.

A3. Top trace: intracellular recording from a cat NRT neuron under barbiturate anaesthesia shows rhythmic LTCPs superimposed on a depolarizing envelope during a spindle wave. Bottom trace: in contrast, *in vitro* intracellular activity from a ferret neuron during spindle-like oscillations recorded in a slice preparation revealed LTCPs of increasing amplitude followed by a progressively larger afterhyperpolarization.

A4. *In vivo* intracellular recording from a cat TC neurone under barbiturate anaesthesia shows that the NRT-evoked IPSPs occasionally give rise to LTCPs. The resulting increase in intracellular Ca^{2+} concentration can progressively activate I_h , producing an increasing depolarizing drive that contributes to spindle wave termination.

B1. EEG trace showing delta waves at 2.5-3.5 Hz in a naturally sleeping rat and corresponding wavelet scalogram.

B2. *In vitro* recording of a TC neuron in an adult cat slice displays continuous oscillations at delta frequency upon hyperpolarizing DC current injection (top trace). In contrast, *in vivo* recordings in a cat TC neuron under ketamine/xylazine anesthesia clearly showed that delta waves occur as transient episodes during the Down state of slow waves (bottom trace).

A1-B2: reproduced with permission from [92], [70], [79], [26] [80].

Figure 4. Interconnected brain areas involved in sleep control and EEG sleep waves generation that present T-type dependent activity.

Example of LTCPs evoked from hyperpolarized membrane potentials in response to depolarizing current injections in layer V/VI cortical neuron (1), TC (2) and NRT (3) neurons, ventrolateral preoptic neuron (VLPO, 4) and hypocretin-orexin expressing neurons (hcrt/orx, 5) of the lateral hypothalamus. Both tonic firing and LTCP are presented in 1, 4 and 5.

Traces reproduce with permission from [28], [34], [40], [25].