

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 1-(2-Bromo-4-chlorophenyl)-3,3-dimethylthiourea

Gamal A. El-Hiti,<sup>a\*</sup> Keith Smith,<sup>b</sup> Amany S. Hegazy,<sup>b</sup> Mohammad Hayal Alotaibi<sup>c</sup> and Benson M. Kariuki<sup>b\*</sup><sup>a</sup>Cornea Research Chair, Department of Optometry, College of Applied Medical Sciences, King Saud University, PO Box 10219, Riyadh 11433, Saudi Arabia,<sup>b</sup>School of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, Wales, and <sup>c</sup>Petrochemical Research Institute, King Abdulaziz City for Science and Technology, PO Box 6086, Riyadh 11442, Saudi Arabia

Correspondence e-mail: gelhiti@ksu.edu.sa, kariukib@cardiff.ac.uk


Received 9 May 2014; accepted 16 May 2014

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.028;  $wR$  factor = 0.077; data-to-parameter ratio = 17.3.

In the title compound,  $\text{C}_9\text{H}_{10}\text{BrClN}_2\text{S}$ , the dimethylthiourea group is twisted from the benzene ring plane by  $54.38$  ( $6$ )°. In the crystal, the amino groups are involved in the formation of  $\text{N}-\text{H}\cdots\text{S}$  hydrogen bonds, which link the molecules into chains along  $[010]$ . Weak  $\text{C}-\text{H}\cdots\text{Cl}$  interactions further link these chains into layers parallel to the  $ab$  plane.

## Related literature

For related compounds, see: Maddani & Prabhu (2010); Yahyazadeh & Ghasemi (2013); Zhao *et al.* (2013). For convenient routes for modifying urea derivatives *via* organolithium intermediates, see: Smith *et al.* (1996, 1999, 2009, 2010, 2012, 2014). For the structures of related compounds, see: Zhao *et al.* (2008); Ramnathan *et al.* (1996).



## Experimental

## Crystal data

$\text{C}_9\text{H}_{10}\text{BrClN}_2\text{S}$   
 $M_r = 293.61$   
 Monoclinic,  $P2_1/n$   
 $a = 12.1369$  (3) Å

$b = 7.9431$  (2) Å  
 $c = 13.2230$  (4) Å  
 $\beta = 115.386$  (3)°  
 $V = 1151.67$  (6) Å<sup>3</sup>

$Z = 4$   
 Cu  $K\alpha$  radiation  
 $\mu = 8.40$  mm<sup>-1</sup>

$T = 296$  K  
 $0.28 \times 0.20 \times 0.09$  mm

## Data collection

Agilent SuperNova (Dual, Cu at zero, Atlas) diffractometer  
 Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2014)  
 $T_{\min} = 0.580$ ,  $T_{\max} = 1.000$

4291 measured reflections  
 2245 independent reflections  
 2078 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.015$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$   
 $wR(F^2) = 0.077$   
 $S = 1.04$   
 2245 reflections

130 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.33$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.39$  e Å<sup>-3</sup>

**Table 1**  
 Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{S1}^i$	0.86	2.67	3.349 (2)	137
$\text{C9}-\text{H9B}\cdots\text{Cl1}^{\text{ii}}$	0.96	2.81	3.696 (2)	153

Symmetry codes: (i)  $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $x - 1, y, z$ .

Data collection: *CrysAlis PRO* (Agilent, 2014); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS2013* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *WinGX* (Farrugia, 2012).

The authors thank the College of Applied Medical Sciences Research Center and the Deanship of Scientific Research at King Saud University for funding this research.

Supporting information for this paper is available from the IUCr electronic archives (Reference: CV5457).

## References

- Agilent (2014). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.  
 Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.  
 Maddani, M. R. & Prabhu, K. R. (2010). *J. Org. Chem.* **75**, 2327–2332.  
 Ramnathan, A., Sivakumar, K., Janarthanan, N., Meerarani, D., Ramadas, K. & Fun, H.-K. (1996). *Acta Cryst.* **C52**, 411–414.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Smith, K., El-Hiti, G. A. & Alshammari, M. B. (2012). *Synthesis*, **44**, 2013–2022.  
 Smith, K., El-Hiti, G. A. & Alshammari, M. B. (2014). *Synthesis*, **46**, 394–402.  
 Smith, K., El-Hiti, G. A. & Hegazy, A. S. (2010). *Synthesis*, pp. 1371–1380.  
 Smith, K., El-Hiti, G. A., Hegazy, A. S., Fekri, A. & Kariuki, B. M. (2009). *Arkivoc* **xiv**, 266–300.  
 Smith, K., El-Hiti, G. A. & Shukla, A. P. (1999). *J. Chem. Soc. Perkin Trans. 1*, pp. 2305–2313.  
 Smith, K., Shukla, A. P. & Matthews, I. (1996). *Sulfur Lett.* **20**, 121–137.  
 Yahyazadeh, A. & Ghasemi, Z. (2013). *Eur. Chem. Bull.* **2**, 573–575.  
 Zhao, Q., Li, S., Huang, K., Wang, R. & Zhang, X. (2013). *Org. Lett.* **15**, 4014–4017.  
 Zhao, P. S., Qin, Y. Q., Zhang, J. & Jian, F. F. (2008). *Pol. J. Chem.* **82**, 2153–2165.

## supporting information

*Acta Cryst.* (2014). E70, o704 [doi:10.1107/S1600536814011350]

## 1-(2-Bromo-4-chlorophenyl)-3,3-dimethylthiourea

**Gamal A. El-Hiti, Keith Smith, Amany S. Hegazy, Mohammad Hayal Alotaibi and Benson M. Kariuki**

### S1. Structural commentary

Recently, various thiourea derivatives have been synthesised and showed broad interesting properties (Maddani & Prabhu, 2010; Yahyazadeh & Ghasemi, 2013; Zhao *et al.*, 2013). In a continuation of our research focused on new synthetic routes towards novel substituted urea derivatives (Smith *et al.*, 1996, 1999, 2009, 2010, 2012, 2014) we have synthesized 3-(2-bromo-4-chlorophenyl)-1,1-dimethylthiourea (I) in a high yield (Smith *et al.*, 1996). We have prepared the material again and crystallized it in high purity in order to obtain its crystal structure, which we present herein.


In (I) (Fig. 1), all bond lengths and angles are normal and correspond well to those observed in the related compounds (Zhao *et al.*, 2008; Ramnathan *et al.*, 1996). The non-hydrogen atoms in (I) fall into two planes with an interplanar angle of 54.38 (6)° between the bromo-chlorophenyl and dimethylthiourea groups. Each molecule is involved in N—H···S contacts (Table 1) with two neighbouring molecules, with one as an acceptor and the other as a donor, leading to the formation of zig-zag-chains in [010] (Fig 2). The bromo-chlorophenyl and dimethylthiourea groups of adjacent molecules are parallel in the stack forming chains of alternating S···Br···S groups with a separation of 4.07 Å and 4.11 Å between the atoms.

### S2. Synthesis and crystallization


To a stirred solution of 2-bromo-4-chloro-1-isothiocyanatobenzene (12.43 g, 50.0 mmol) in anhydrous dioxane (120 ml) dimethylamine (7.10 g of 33% solution in ethanol, 52.0 mmol) was slowly added in a drop-wise manner over 5 min. The reaction mixture was stirred at room temperature for an extra 1 h. The solid obtained was collected by filtration and washed with dioxane (2 x 20 ml) and dried. Recrystallization from ethyl acetate gave 3-(2-bromo-4-chlorophenyl)-1,1-dimethylthiourea (13.80 g, 47.0 mmol; 94%) as yellow crystals, m.p. 193–194 °C [lit. 184–185 °C (ethyl acetate); Smith *et al.* (1996)]. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>, δ, p.p.m.) 7.97 (d, *J* = 8.8 Hz, 1 H, H-6), 7.59 (d, *J* = 2.3 Hz, 1 H, H-3), 7.32 (dd, *J* = 2.3, 8.8 Hz, 1 H, H-5), 7.17 (br, exch., 1 H, NH), 3.43 [s, 6 H, N(CH<sub>3</sub>)<sub>2</sub>]. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>, δ, p.p.m.) 181.2 (s, C=S), 136.6 (s, C-1), 131.8 (d, C-3), 131.0 (s, C-4), 127.8 (d, C-6), 127.7 (d, C-5), 118.1 (s, C-2), 41.3 [q, N(CH<sub>3</sub>)<sub>2</sub>]. AP<sup>+</sup>—MS (*m/z*, %): 297 ([*MH*<sup>81</sup>Br<sup>37</sup>Cl]<sup>+</sup>, 34), 295 ([*MH*<sup>81</sup>Br<sup>35</sup>Cl and *MH*<sup>79</sup>Br<sup>37</sup>Cl]<sup>+</sup>, 100), 293 ([*MH*<sup>79</sup>Br<sup>35</sup>Cl]<sup>+</sup>, 80), 263 (12), 215 (22), 213 (50). HRMS (AP<sup>+</sup>): Calculated for C<sub>9</sub>H<sub>11</sub><sup>79</sup>Br<sup>35</sup>ClN<sub>2</sub>S [*MH*] 292.9515; found, 292.9515.

### S3. Refinement

H atoms were positioned geometrically and refined using a riding model with *U*<sub>iso</sub>(H) = 1.2 times *U*<sub>eq</sub> for the atom they are bonded to except for the methyl groups where 1.5 times *U*<sub>eq</sub> was used with free rotation about the C—C bond.

**Figure 1**

View of (I) showing atom labels and 50% probability displacement ellipsoids for non-H atoms.

**Figure 2**

A portion of the crystal packing viewed along the *a* axis. N—H...S contacts are shown as dotted lines.

### 1-(2-Bromo-4-chlorophenyl)-3,3-dimethylthiourea

#### Crystal data

$C_9H_{10}BrClN_2S$

$M_r = 293.61$

Monoclinic,  $P2_1/n$

$a = 12.1369 (3) \text{ \AA}$

$b = 7.9431 (2) \text{ \AA}$

$c = 13.2230 (4) \text{ \AA}$

$\beta = 115.386 (3)^\circ$

$V = 1151.67 (6) \text{ \AA}^3$

$Z = 4$

$F(000) = 584$

$D_x = 1.693 \text{ Mg m}^{-3}$

Cu  $K\alpha$  radiation,  $\lambda = 1.5418 \text{ \AA}$

Cell parameters from 2078 reflections

$\theta = 4.1\text{--}75.5^\circ$

$\mu = 8.40 \text{ mm}^{-1}$

$T = 296 \text{ K}$

Plate, colourless

$0.28 \times 0.20 \times 0.09 \text{ mm}$

Data collection

Agilent SuperNova (Dual, Cu at zero, Atlas) diffractometer	4291 measured reflections
Radiation source: SuperNova (Cu) X-ray Source	2245 independent reflections
Mirror monochromator	2078 reflections with $I > 2\sigma(I)$
$\omega$ scans	$R_{\text{int}} = 0.015$
Absorption correction: multi-scan (CrysAlis PRO; Agilent, 2014)	$\theta_{\text{max}} = 73.5^\circ$ , $\theta_{\text{min}} = 4.1^\circ$
$T_{\text{min}} = 0.580$ , $T_{\text{max}} = 1.000$	$h = -13 \rightarrow 14$
	$k = -9 \rightarrow 6$
	$l = -16 \rightarrow 15$

Refinement

Refinement on $F^2$	H-atom parameters constrained
Least-squares matrix: full	$w = 1/[\sigma^2(F_o^2) + (0.0429P)^2 + 0.4682P]$
$R[F^2 > 2\sigma(F^2)] = 0.028$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.077$	$(\Delta/\sigma)_{\text{max}} = 0.002$
$S = 1.04$	$\Delta\rho_{\text{max}} = 0.33 \text{ e } \text{\AA}^{-3}$
2245 reflections	$\Delta\rho_{\text{min}} = -0.39 \text{ e } \text{\AA}^{-3}$
130 parameters	Extinction correction: SHELXL2013 (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001x Fc^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
0 restraints	Extinction coefficient: 0.0048 (3)
Hydrogen site location: inferred from neighbouring sites	

Special details

**Experimental.** Absorption correction: CrysAlisPro (Agilent, 2014). Empirical absorption correction using spherical harmonics, implemented in SCALE3 ABSPACK scaling algorithm.

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.85876 (18)	0.0661 (3)	0.22866 (16)	0.0377 (4)
C2	0.85637 (19)	0.1344 (3)	0.13058 (16)	0.0404 (4)
C3	0.9605 (2)	0.1405 (3)	0.11271 (19)	0.0495 (5)
H3	0.9581	0.1851	0.0468	0.059*
C4	1.0675 (2)	0.0794 (3)	0.1942 (2)	0.0523 (5)
C5	1.0726 (2)	0.0087 (3)	0.29241 (19)	0.0497 (5)
H5	1.1455	-0.0334	0.3465	0.060*
C6	0.96763 (19)	0.0022 (3)	0.30797 (17)	0.0440 (5)
H6	0.9699	-0.0460	0.3729	0.053*
C7	0.72784 (18)	0.1177 (3)	0.32603 (16)	0.0379 (4)
C8	0.5801 (3)	0.1471 (4)	0.4043 (2)	0.0638 (7)
H8A	0.5786	0.0517	0.4483	0.096*
H8B	0.5006	0.1978	0.3710	0.096*
H8C	0.6383	0.2278	0.4515	0.096*
C9	0.5176 (2)	0.0190 (4)	0.2169 (2)	0.0569 (6)
H9A	0.5032	0.0890	0.1533	0.085*
H9B	0.4443	0.0110	0.2274	0.085*

H9C	0.5416	-0.0913	0.2045	0.085*
Br1	0.70976 (2)	0.22178 (4)	0.01974 (2)	0.05768 (14)
Cl1	1.19895 (7)	0.08827 (15)	0.17270 (8)	0.0936 (3)
N1	0.74917 (15)	0.0536 (3)	0.24110 (14)	0.0442 (4)
H1	0.6901	-0.0001	0.1897	0.053*
N2	0.61441 (17)	0.0925 (3)	0.31655 (16)	0.0470 (4)
S1	0.83592 (5)	0.22395 (7)	0.43366 (4)	0.04637 (16)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0377 (9)	0.0420 (10)	0.0314 (9)	-0.0038 (8)	0.0129 (7)	-0.0039 (8)
C2	0.0436 (10)	0.0403 (10)	0.0335 (9)	-0.0009 (8)	0.0128 (8)	-0.0013 (8)
C3	0.0570 (13)	0.0531 (13)	0.0436 (11)	-0.0046 (10)	0.0264 (10)	0.0013 (10)
C4	0.0437 (11)	0.0644 (15)	0.0526 (12)	-0.0043 (10)	0.0242 (10)	-0.0078 (11)
C5	0.0395 (10)	0.0595 (14)	0.0420 (11)	0.0018 (9)	0.0098 (9)	-0.0050 (10)
C6	0.0451 (10)	0.0495 (12)	0.0327 (9)	0.0004 (9)	0.0123 (8)	0.0013 (8)
C7	0.0420 (10)	0.0375 (10)	0.0312 (9)	0.0047 (8)	0.0128 (8)	0.0048 (7)
C8	0.0684 (15)	0.0766 (18)	0.0611 (15)	0.0111 (14)	0.0416 (13)	0.0006 (13)
C9	0.0376 (10)	0.0679 (16)	0.0603 (14)	0.0018 (10)	0.0164 (10)	-0.0056 (12)
Br1	0.05856 (19)	0.0600 (2)	0.04035 (17)	0.01302 (11)	0.00771 (12)	0.00683 (10)
Cl1	0.0558 (4)	0.1432 (9)	0.0976 (6)	-0.0020 (4)	0.0480 (4)	-0.0011 (6)
N1	0.0376 (8)	0.0595 (11)	0.0339 (8)	-0.0088 (8)	0.0138 (7)	-0.0088 (8)
N2	0.0435 (9)	0.0547 (11)	0.0451 (9)	0.0054 (8)	0.0213 (8)	-0.0008 (8)
S1	0.0545 (3)	0.0447 (3)	0.0308 (3)	0.0000 (2)	0.0096 (2)	-0.00197 (19)

*Geometric parameters (Å, °)*

C1—C6	1.384 (3)	C7—N2	1.343 (3)
C1—C2	1.395 (3)	C7—N1	1.355 (3)
C1—N1	1.412 (3)	C7—S1	1.690 (2)
C2—C3	1.383 (3)	C8—N2	1.457 (3)
C2—Br1	1.887 (2)	C8—H8A	0.9600
C3—C4	1.372 (3)	C8—H8B	0.9600
C3—H3	0.9300	C8—H8C	0.9600
C4—C5	1.392 (3)	C9—N2	1.459 (3)
C4—Cl1	1.738 (2)	C9—H9A	0.9600
C5—C6	1.375 (3)	C9—H9B	0.9600
C5—H5	0.9300	C9—H9C	0.9600
C6—H6	0.9300	N1—H1	0.8600
C6—C1—C2	118.68 (18)	N1—C7—S1	122.03 (16)
C6—C1—N1	121.79 (18)	N2—C8—H8A	109.5
C2—C1—N1	119.39 (18)	N2—C8—H8B	109.5
C3—C2—C1	121.09 (19)	H8A—C8—H8B	109.5
C3—C2—Br1	118.76 (16)	N2—C8—H8C	109.5
C1—C2—Br1	120.14 (15)	H8A—C8—H8C	109.5
C4—C3—C2	118.7 (2)	H8B—C8—H8C	109.5

C4—C3—H3	120.7	N2—C9—H9A	109.5
C2—C3—H3	120.7	N2—C9—H9B	109.5
C3—C4—C5	121.6 (2)	H9A—C9—H9B	109.5
C3—C4—Cl1	118.89 (19)	N2—C9—H9C	109.5
C5—C4—Cl1	119.51 (19)	H9A—C9—H9C	109.5
C6—C5—C4	118.8 (2)	H9B—C9—H9C	109.5
C6—C5—H5	120.6	C7—N1—C1	126.45 (17)
C4—C5—H5	120.6	C7—N1—H1	116.8
C5—C6—C1	121.2 (2)	C1—N1—H1	116.8
C5—C6—H6	119.4	C7—N2—C8	120.9 (2)
C1—C6—H6	119.4	C7—N2—C9	122.72 (18)
N2—C7—N1	114.77 (18)	C8—N2—C9	116.3 (2)
N2—C7—S1	123.19 (16)		

*Hydrogen-bond geometry (Å, °)*

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N1—H1...S1 <sup>i</sup>	0.86	2.67	3.3488 (19)	137
C9—H9B...Cl1 <sup>ii</sup>	0.96	2.81	3.696 (2)	153

Symmetry codes: (i)  $-x+3/2, y-1/2, -z+1/2$ ; (ii)  $x-1, y, z$ .