

The T cell antigen receptor: the Swiss army knife of the immune system

M. Attaf, M. Legut, D. K. Cole
and A. K. Sewell

*Division of Infection and Immunity, Cardiff
University School of Medicine, Cardiff, UK*

Summary

The mammalian T cell receptor (TCR) orchestrates immunity by responding to many billions of different ligands that it has never encountered before and cannot adapt to at the protein sequence level. This remarkable receptor exists in two main heterodimeric isoforms: $\alpha\beta$ TCR and $\gamma\delta$ TCR. The $\alpha\beta$ TCR is expressed on the majority of peripheral T cells. Most $\alpha\beta$ T cells recognize peptides, derived from degraded proteins, presented at the cell surface in molecular cradles called major histocompatibility complex (MHC) molecules. Recent reports have described other $\alpha\beta$ T cell subsets. These 'unconventional' T cells bear TCRs that are capable of recognizing lipid ligands presented in the context of the MHC-like CD1 protein family or bacterial metabolites bound to the MHC-related protein 1 (MR1). $\gamma\delta$ T cells constitute a minority of the T cell pool in human blood, but can represent up to half of total T cells in tissues such as the gut and skin. The identity of the preferred ligands for $\gamma\delta$ T cells remains obscure, but it is now known that this receptor can also functionally engage CD1-lipid, or immunoglobulin (Ig) superfamily proteins called butyrophilins in the presence of pyrophosphate intermediates of bacterial lipid biosynthesis. Interactions between TCRs and these ligands allow the host to discriminate between self and non-self and co-ordinate an attack on the latter. Here, we describe how cells of the T lymphocyte lineage and their antigen receptors are generated and discuss the various modes of antigen recognition by these extraordinarily versatile receptors.

Keywords: CD1, MHC-I, MHC-II, MHC-Ib, MR1, T cell, T cell receptor, $\alpha\beta$ TCR, $\gamma\delta$ TCR

Accepted for publication 4 March 2015

Correspondence: A. K. Sewell, Division of
Infection and Immunity, Henry Wellcome
Building, Cardiff University School of
Medicine, University Hospital, Cardiff CF14
4XN, UK.

E-mail: sewellak@cardiff.ac.uk

Introduction

Immune surveillance by T lymphocytes is critical for the immune integrity of all jawed vertebrates and is imposed through an intricate armoury of functions that eliminate pathogen-infected and neoplastic cells. The T cell pool consists of several functionally and phenotypically heterogeneous subpopulations. T cells are broadly classified as $\alpha\beta$ or $\gamma\delta$ according to the somatically rearranged T cell receptor (TCR) they express at their surface.

$\alpha\beta$ T cells are by far the most abundant and the best-characterized circulating T cells. Most $\alpha\beta$ T cells recognize peptides from degraded proteins bound to major histocompatibility complex (MHC) molecules at the cell surface. These peptide-MHC (pMHC)-recognizing T cells were the first to be described [1]. T cells that respond to pMHC are

said to be 'conventional'. However, a significant fraction of the $\alpha\beta$ T cell pool consists of rarer T lymphocytes that do not recognize pMHC. These 'unconventional' $\alpha\beta$ T cells include: (i) mucosal-associated invariant T (MAIT) cells that display limited diversity and are involved in anti-bacterial immunity [2,3]; (ii) invariant natural killer T (iNKT) cells; and (iii) germline-encoded mycolyl-reactive (GEM) T cells. iNKT and GEM T cells are dedicated to recognition of glycolipids in the context of CD1d and CD1b, respectively [4,5]. Other T cells are believed to recognize lipid antigens in the context of CD1a and CD1c, but these subsets have not been well characterized, or named, at the time of writing.

$\gamma\delta$ T cells are also grouped as being 'unconventional', as they are not MHC-restricted and do not appear to recognize

peptide antigens. One to 10% of circulating T cells express a $\gamma\delta$ TCR, but this fraction is considerably higher in epithelial tissues. These tissues form the main portal of entry for pathogens, suggesting an important role of $\gamma\delta$ T cells as early immune sentinels. Murine knock-out studies have demonstrated that $\gamma\delta$ T cells have a clear role in pathogen clearance and tumour surveillance [6]. Some T cells express a $\delta/\alpha\beta$ TCR hybrid. This is because the recombination mechanism used to generate TCR chains operates on the same locus for TCR- δ and TCR- α . Thus, some gene segments within the locus can, in some instances, be involved in the generation of TCR chains that are partially α and partially δ and are able to pair with canonical TCR- β chains [7]. These so-called $\delta/\alpha\beta$ T cells constitute a significant proportion of T cells that recognize the CD1d- α -galactosylceramide (α GalCer) complex [7]. Other hybrid TCRs further blur traditional segregation into $\alpha\beta$ and $\gamma\delta$ TCRs, as $V\gamma J\beta$, $V\beta J\gamma$, $V\delta J\beta$, $V\delta J\gamma$ and $V\gamma J\delta$ TCR chains have been described, with or without the inclusion of D β and D δ segments. Many of these combinations follow the 12/23 rule of V-(D)-J recombination, are transcribed into full transcripts and translated into hybrid proteins [8]. The antigens recognized by T cells bearing these non-canonical TCR chains remain unknown, although a subset of $V\gamma$ (D β) J β TCRs are reported to be MHC-restricted [9].

Despite the functional and phenotypical differences between subsets, all T cells arise from the same precursors and share their early differentiation history. Thymic progenitor cells seed the thymus from the fetal liver or adult bone marrow. In the thymus, these progenitors enter a complex differentiation programme which leads to irreversible acquisition of T cell identity and expression of clonally distributed, variegated TCRs. In this review, we describe the mechanisms through which MHC-restricted and MHC-independent TCRs are generated and discuss recognition of antigen by distinct T cell subsets. We also re-examine recent findings on the more 'unconventional' subsets.

Generation of diversity and thymic selection

TCR diversity is generated somatically by gene rearrangement, a process that allows a vast array of different receptors to be produced from a limited set of genes. At the TCR- α locus (*tra*), discrete variable (V) and junctional (J) gene segments are recombined and juxtaposed to a constant (C) segment (Fig. 1a). Recombination at the TCR- β locus (*trb*) is similar, but includes an additional diversity (D) segment and one of two C segments can be appended to the rearranged TCR- β chain (Fig. 1b). The variability of TCRs is confined predominantly to three short hairpin loops on each chain, called complementarity determining regions (CDR). Collectively, the six CDR loops sit at the membrane-distal end of the TCR extracellular domain to form the antigen-binding site (Fig. 2a).

The variable CDR1 and 2 loops are encoded in the germline by the T cell receptor alpha variable (TRAV) and T cell receptor beta variable (TRBV) gene segments. By contrast, the hypervariable CDR3 loops are generated by random deletion and addition of template and non-template nucleotides at the junction between recombining V, (D) and J gene segments (Fig. 2b). Polymorphism in the *tr* loci could add yet further diversity to the potential TCR repertoire at the population level [12]. Theoretically, gene rearrangement by V-(D)-J recombination alone can produce $\sim 10^{18}$ TCRs in humans [13] and $\sim 10^{15}$ TCRs in the mouse [14]. $\gamma\delta$ TCRs are also generated by V-(D)-J recombination (Fig. 1c,d). The TCR- δ chain is thought to be by far the most diverse TCR chain due to the inclusion of multiple D segments, which can be translated in any reading frame (Fig. 2b). Thus, the theoretical number of different $\gamma\delta$ TCRs that could be produced is potentially much greater than for the $\alpha\beta$ TCR.

The quasi-random process of generating $\alpha\beta$ TCRs described above has the capacity to generate receptors that are inept at recognizing self-MHC molecules and receptors that could be autoreactive. Thymic selection ensures that only T cells bearing a TCR that recognizes self-peptides in the context of self-MHC receive a survival signal. The majority of thymocytes do not receive this signal. Cells that express TCRs that cannot recognize self-pMHC are unlikely to be useful for recognizing foreign peptides. These cells do not receive a survival signal through their TCR and are said to 'die by neglect'. At the other extreme, thymocytes that bear TCRs that react strongly to self-pMHC have the capacity to be autoreactive and are culled through a process of negative selection. Together, positive and negative selection ensure that only those $\alpha\beta$ T cells that are restricted to recognizing self-pMHC within a low affinity range can populate the periphery. Thus, the thymic environment allows the generation of a pool of $\alpha\beta$ T cells that are self-restricted, but not self-reactive [15].

Much less is known about selection of other, MHC-independent, T cell subsets. Invariant (type I) NK T cells are selected on CD1d-expressing CD4⁺CD8⁺ double-positive thymocytes and acquire effector function before exiting the thymus [16–18]. Selection of MAIT cells has been shown recently to require MR1 expression on double-positive thymocytes [19]. Commitment to the $\gamma\delta$ T cell fate is thought to be a TCR-dependent process whereby strong $\gamma\delta$ TCR signals induce $\gamma\delta$ commitment and weak pre-TCR signals, in the absence of $\gamma\delta$ TCR signalling, instruct thymocytes to initiate *tra* rearrangement [20]. This model mirrors classic positive selection via the $\alpha\beta$ TCR and suggests that $\gamma\delta$ T cells may also need to encounter a cognate ligand in the thymus. Indeed, CD73 is up-regulated as a result of $\gamma\delta$ TCR activation in the thymus. As CD73 is expressed by the majority of peripheral $\gamma\delta$ T cells, ligand recognition in the thymus appears




Fig. 1. Generation of $\alpha\beta$ and $\gamma\delta$ T cell receptors (TCRs) by V-(D)-J recombination. (a) The *tra/trd* locus consists of a cluster of 46 functional T cell receptor alpha variable (TRAV) segments and eight T cell receptor delta variable (TRDV) segments, followed by three segments in the T cell receptor delta diversity (TRDD) cluster and four segments in the T cell receptor delta joining (TRDJ) cluster. A total of 51 functional TRAV segments lie between the TRDC and the T cell receptor alpha chain constant region (TRAC) segments. At the *tra/trd* locus, V-J recombination brings together one of many TRAV segments and one of many TRAJ segments. The intervening sequences are spliced out, producing a TCR- α transcript in which V, J and C segments are directly adjacent. (b) The *trb* locus comprises 48 functional T cell receptor beta variable (TRBV) segments followed by two D segments, 12 functional TRBJ segments and two TRBC segments. For TCR- β chain rearrangements, V-(D)-J recombination is a two-step, ordered process. D-J recombination occurs first, juxtaposing TRBD1 to one of many TRBJ1 segments or TRBD2 to one of many TRBJ2 segments. V-DJ recombination subsequently brings the rearranged DJ join to one of many TRBV segments. The intervening sequences are then spliced out, generating a TCR- β transcript in which V, D, J and C segments are directly adjacent. (c) The *trg* locus is composed of six functional TRGV segments and five TRGJ segments followed by two TRGC segments. At the *trg* locus, V-J recombination joins one of many TRGV segments to a TRGJ segment. Similar to TCR- α chains, the intervening regions are spliced, producing a TCR- γ transcript in which V, J and C segments are directly adjacent. (d) The generation of TCR- δ chains also occurs at the *tra/trd* locus. The 5' end of this locus consists of a cluster of V genes. All V genes in this cluster can recombine with TRAJ, but only a subset can recombine with TRDD. Thus, many V genes in the *tra/trd* locus are used exclusively for TCR- δ rearrangement in early thymic precursors, while others are used exclusively for TCR- α rearrangement in double-positive thymocytes. Some V genes can be used interchangeably. Similar to TCR- β chains, TCR- δ chains are produced by V-(D)-J recombination and splicing, producing a final transcript in which V, D, J and C segments are directly juxtaposed. Unlike the TCR- β chain, however, TCR- δ can incorporate multiple D segments.

to be a common occurrence during $\gamma\delta$ T cell development [21]. However, potential ligands for $\gamma\delta$ T cells are largely unknown. Skint-1 is the only known ligand required for maturation of V γ 5⁺ dendritic epidermal $\gamma\delta$ T cells in the mouse, although its role in selection remains controversial [22]. Interestingly, a study of murine T10/T22-reactive $\gamma\delta$ T cells has indicated that, in contrast to $\alpha\beta$ T cells, ligand recognition by the $\gamma\delta$ TCR imprints effector function on the $\gamma\delta$ T cell pool, but not antigen specificity [23]. Overall, $\gamma\delta$ T cell selection is poorly understood and the identity of potential positive selection ligands for $\gamma\delta$ TCRs continues to be a matter of debate.

Conventional, pMHC-restricted T cells

The cardinal feature of $\alpha\beta$ T cells is the recognition of peptides derived from self and foreign proteins in the context of self-MHC molecules. The *mhc* locus was first described more than half a century ago as the set of genes which determine the outcome of tissue transplantation in congenic mice. It is now clear that the products of the *mhc* have evolved to allow T cells to perform highly specific functions which are crucial to adaptive immunity and host defence as a whole. The *mhc* locus is the most gene-dense and the most polymorphic region known to date, with more than 12 000 different alleles already




Fig. 2. Structure of T cell receptor (TCR) proteins and mRNA. (a) $\alpha\beta$ [Protein Data Bank (PDB): 3HG1] [10] and $\gamma\delta$ (PDB: 1HXM) [11]. TCRs adopt similar tertiary structures that position the complementarity-determining regions (CDR) loops at the membrane distal end of the molecules. Together the six CDR loops form the antigen binding site. (b) The mRNA structures show that for each chain CDR1 and CDR2 are encoded in the germline. CDR3 is the product of junctional diversity at V–J joins of T cell receptor (TCR)- α and TCR- γ chains and V–D–J joins in TCR- β and TCR- δ chains. CDR3 is consequently hypervariable. The colour code adopted for the CDR loops is maintained throughout this paper. The areas coloured in grey represent the constant and variable domains of the TCRs (not including the hypervariable CDR loops).

described in humans [24]. Polymorphism at the *mhc* ensures diversity in peptide presentation at the population level.

There are two types of classical MHC molecules: MHC class I (MHC-I) and MHC class II (MHC-II). There are three classical human MHC-I genes [human leucocyte antigen (HLA)-A, -B and -C]. These genes encode a membrane-spanning α -chain associated with the non-polymorphic β_2 microglobulin (β_2m) protein (Fig. 3a). Polymorphism is confined mainly to the membrane-distal $\alpha 1$ and $\alpha 2$ domains, while the $\alpha 3$ domain is largely invariant. The human MHC-II genes (DP, DQ and DR) encode two distinct polymorphic α and β -chains. Each chain folds into a membrane-distal polymorphic domain followed by a membrane-proximal immunoglobulin (Ig)-like domain (Fig. 3d) [37]. The peptide-binding cleft of both MHC-I and MHC-II consists of two anti-parallel α -helices forming a channel in which the peptide can bind in an extended conformation on a platform of eight anti-parallel β -pleated sheets. Specific peptide-binding pockets in the base of this cleft vary between different MHC alleles [38]. The binding affinity for MHC differs between

peptides according to their primary sequence. A given peptide will, at best, bind a limited set of MHCs. Reciprocally, any MHC allele can only accommodate a small fraction of the peptide collection derived from a given protein.

Conventional $\alpha\beta$ TCR ligands

MHC-I molecules are expressed on nearly all nucleated cells and present peptides derived from endogenous proteins, allowing T cells to interrogate the internal proteome by scanning the surface of the target cell. MHC-II molecules differ from MHC-I in that they predominantly present peptides derived from exogenous proteins and are expressed primarily on professional antigen-presenting cells. Despite the similarities in overall conformation (Fig. 3), MHC-I and MHC-II present peptides in a distinct manner governed by the configuration of the peptide-binding cleft. Polymorphic residues define the size and chemical properties of the binding pockets within the cleft and therefore the peptide collection that can be accommodated by a given MHC-I molecule. The closed




Fig. 3. The structures of peptide-major histocompatibility complex class I (pMHC-I) and class II (pMHC-II). The two classes of classical MHC adopt similar overall structures despite being differently comprised. MHC-I [Protein Data Bank (PDB): 1ZHL] [25] (a) consists of a variable heavy chain (grey) folded with the invariant β -2-microglobulin molecule (cyan). (b) The ends of the MHC-I peptide-binding groove are closed. MHC-I presents peptides of 8–14 amino acids in length to T cells. (c) An 8-mer peptide can lie flat in the MHC-I groove. As additional amino acid residues are added, peptides have to bulge upwards and outwards in order to be accommodated within the groove. It has recently been established that MHC-I restricted T cells can recognize the length of a peptide in addition to its amino acid sequence (PDB: 1ZHL, 1XH3, 2FZ3, 3BW9, 1A1N, 1JF1, 1HHI) [25,27–33], whereas MHC-II (PDB: 1KG0) [26] (d) consists of an α -chain (grey) and a β -chain (cyan). Both MHC-I and MHC-II fold to present peptide (red) to T cells within a peptide-binding groove. (e) The open ends of the MHC-II peptide-binding groove allow presented peptides to extend at both the N- and C-terminus. Thus, MHC-II generally presents longer peptides than MHC-I. This mode of binding also means that the MHC-II can sometimes present the same peptide in different registers by using different amino acid side chains for anchoring into the MHC-binding pockets. (f) The open-ended MHC-II binding groove enables longer peptides to form an elongated conformation with peptide N- and C-terminal peptide flanking regions extending outside of the groove (PDB: 1KG0, 1UVQ, 2SEB, 2IAN) [26,34–36].

conformation of the MHC-I $\alpha_1\alpha_2$ binding cleft (Fig. 3b) restricts the length of peptides that can be accommodated. MHC-I molecules typically bind peptides of eight to 10 amino acids in length, but longer peptides have been observed in some instances [39]. Because the MHC-I peptide-binding groove is closed at both ends, long peptides bulge out (Fig. 3c), exposing peptide side chains for direct interaction with the TCR [39]. Curiously, our own recent studies have shown that MHC-I restricted TCRs appear to be predisposed to bind peptides of a defined

length [27]. In contrast, the MHC-II peptide-binding groove is an open-ended conformation (Fig. 3e) which allows the binding of N- and C-terminally extended peptides of up to 30 amino acids in length (Fig. 3f).

TCR recognition of pMHC

MHC restriction is a defining characteristic of ‘conventional’ $\alpha\beta$ T cells. Although MHC restriction was described more than 30 years ago, the molecular forces




Fig. 4. T cell receptor (TCR)–peptide-major histocompatibility complex (pMHC) structures. MHC molecules (in grey) and peptide (in red sticks) are overlaid with the docking footprints of the individual complementarity-determining regions (CDR) loops of the cognate $\alpha\beta$ TCR. The coloured footprints correspond to the colours of the CDR loops shown in Fig. 2. (a) Structure of MHC-I molecule HLA-A*0201 presenting the immunodominant GLCTLVAML peptide from Epstein–Barr virus (EBV) [Protein Data Bank (PDB): 3O4L] [40]. The coloured footprint shows how the CDR loops of the AS01 TCR sit on the pMHC complex. This complex adopts a canonical conformation where the germline-encoded CDR1 and 2 loops contact mainly the MHC and the hypervariable CDR3 loops sit over the peptide. (b) Structure of the MS2-3C8TCR docked on the MHC class II molecule human leucocyte antigen (HLA)-DR4. Here, HLA-DR4 presents a peptide from myelin basic protein (PDB: 3O6F) [41]. (c) Overlay of all MHC-I (grey cartoon and surface)-restricted TCRs (multi-coloured) in which co-complex structures have been solved. All complexes were aligned on the MHC-I molecule to demonstrate the flexible nature of TCR–pMHC binding.

driving this interaction are still fiercely debated. Structural analyses of TCRs in their free and MHC-bound states have established that the TCR–pMHC interaction conforms to some ‘rules of engagement’. Indeed, most TCRs bind pMHC in a diagonal or nearly orthogonal orientation relative to the long axis of the MHC peptide-binding groove (Fig. 4a and b) [37]. This conserved docking strategy, together with the observation that 20% of the pre-selection TCR repertoire is MHC-specific [42], has led to the idea that TCRs are genetically ‘hard-wired’ for MHC recognition [23–45]. These rules fit conveniently with the observation that the germline-encoded CDR1 α , CDR1 β , CDR2 α and CDR2 β often contact the MHC α -helices, whereas the somatically rearranged CDR3 α and CDR3 β loops contact the peptide (Fig. 4c). This dogma was long supported by the identification of the so-called ‘interaction codons’ and the amino acids defining the MHC-I ‘restriction triad’ [44,46]. More

recently, however, Stadinski *et al.* [47] challenged the germline theory by demonstrating that the putative ‘interaction codons’ in a murine TCR- β chain were not strictly conserved, but were largely dependent on the identity of the partner TCR- α chain. Mutational studies on MHC alleles also suggest that TCR–pMHC interactions allow for substantial plasticity within the confines of a common binding and orientation system [48]. A murine study in which CDR1 and CDR2 were randomized *in vivo* in the context of a fixed CDR3 failed to show preferential selection for the wild-type CDR1 and 2 sequences, contradicting the idea of TCRs being genetically ‘biased’ towards MHC ligands [49]. It remains possible that the semi-conserved angle and polarity of TCR–pMHC interaction is required for the correct orientation of intracellular signalling domains, the positioning of the co-receptor binding site on MHC relative to the TCR or imposed by extracellular sites of glycosylation, as indicated by the




Fig. 5. CD8 and CD4 co-receptors bind to invariant parts of peptide-major histocompatibility complex class I (pMHC-I) and pMHC-II, respectively. (a) Structural model of a T cell receptor (TCR)–pMHC-I–CD8 tripartite interaction [Protein Data Bank (PDB): 3O4L and 1AKJ] [40,52]. MHC-I (grey cartoon) forms the peptide binding cleft using its $\alpha 1$ and $\alpha 2$ domains. The membrane distal face of the molecule comprises the TCR (blue and slate cartoon) docking platform. CD8 (green and yellow cartoon) binds at a distinct site on the $\alpha 3$ domain of the molecule. The structure shown is human leucocyte antigen (HLA) A*0201 and a CD8 $\alpha\alpha$ homodimer [52]. (b) Tripartite complex structure of the TCR–pMHC-II–CD4 interaction (CD4 shown in orange cartoon) (PDB: 3T0E) [50]. Similar to the TCR–pMHC-I–CD8 model, the CD4 co-receptor binds to an invariant site distal from the TCR binding platform.

crystal structure of a ternary TCR–pMHC–CD4 complex [50]. Our recent studies show that TCR–peptide specificity overrides affinity-enhancing TCR–pMHC interactions to suggest that TCR CDR3 loops might need to ‘accept’ peptide prior to TCR engagement the MHC component of the ligand [51]. Further work is required in order to dissect the sequence of molecular events involved in TCR–pMHC binding.

An alternative to the germline theory of MHC restriction suggests that recognition of MHC-restricted ligands is the result of a selection process through which the peripheral repertoire is enriched with MHC-specific TCRs and purged of MHC-independent TCRs. In a series of studies involving MHC-I, MHC-II and co-receptor deficient-mice (referred to as ‘quad-deficient’ mice), it was shown that the selection of MHC-specific TCRs was strictly governed by the T cell co-receptors CD4 and CD8. CD4 and CD8 act as antigen co-receptors by binding to invariant regions of the MHC-II and MHC-I, respectively, at sites distinct from the TCR docking platform (Fig. 5). The cytoplasmic tail of the co-receptor binds to the protein tyrosine kinase Lck, which mediates key membrane-proximal phosphorylation events during T cell activation and selection. Singer and colleagues [53] first showed that the deletion of both co-receptors allowed selection of CD4⁺ and CD8⁺ T cells to take place on an MHC-deficient background. The authors worked on the premise that co-receptor-independent signalling can occur and is initiated by non-MHC ligand binding to the TCR, akin to cross-linking antibodies. By contrast, in the presence of co-receptors, which sequester all the available Lck needed for signalling, TCR signalling can be triggered only by MHC ligands. Thus, in the ‘quad-deficient’

mouse, non-MHC ligands can induce thymic selection of MHC-independent T cells – T cells which would otherwise die by neglect in a normal thymus [54–56]. Reports of MHC-independent ligands for $\alpha\beta$ T cells are scarce in the literature, although a number of examples have been observed in the mouse [57] and in humans [58,59]. The existence of MHC-independent $\alpha\beta$ TCRs in the periphery further supports the notion that MHC restriction is a TCR-extrinsic feature imposed on developing thymocytes through thymic selection.

$\alpha\beta$ TCR cross-reactivity

As described above, the *mhc* locus is an example of extreme polymorphism. The vast majority of polymorphic residues in MHC proteins cluster around the peptide-binding cleft to suggest that this diversity is upheld to expand the variety of peptides that can be displayed to the immune system [38]. The TCR must recognize peptides presented by all these variants. Beyond this, effective T cell immunity requires the TCR repertoire to recognize virtually any foreign peptide that can bind to host MHC molecules as a failure to recognize all possible foreign peptides would leave ‘blind spots’ that could be exploited by pathogens [60]. Unlike the B cell receptor, the protein sequence of the TCR is fixed and never undergoes affinity maturation, so the TCRs expressed on existing naive T cells must be capable of responding to all alien antigens, despite never having encountered them before and being unable to adapt to them. The size of this task becomes apparent once it is realized just how many potential foreign peptides there are. This major evolutionary challenge has been met by enabling each




Fig. 6. The plasticity of $\alpha\beta$ T cell receptor (TCR) binding to peptide-major histocompatibility complex (pMHC). Individual TCRs use multiple mechanisms to bind to pMHC. These effects can increase the number of individual peptides that can be recognized. (a) Macro-level changes enable the TCR to bind pMHC with an altered angle or altered register. The cartoon shows the ‘footprints’ of TCR complementarity-determining region (CDR) loops projected down onto the pMHC. (b) Relatively micro-level flexibility in the CDR loops allows them to accommodate a variety of different shapes. The cartoon shows a side view of a TCR engaging pMHC. (c) The existing database of TCR–pMHC structures indicates that TCRs tend to focus interaction on two to four upward-facing amino acid residues in the antigenic peptide (so-called ‘hotspots’ [63]). In this example a TCR might focus on two amino acids in the peptide (shown in red). Such residue-focused interaction then allows the TCR to accommodate multiple amino acid substitutions at other positions in the peptide (indicated by the use of different colours on the right). The three mechanisms described above are not mutually exclusive and represent just some of the possibilities. Many residues can also bind in individual MHC binding pockets. It is now understood that altering a primary MHC anchor can substantially change the way that a peptide might be viewed by incoming T cells [64,65].

TCR to interact with many – sometimes millions – of individual peptides bound in the groove of a single MHC [13,61]. TCRs are further capable of engaging peptides presented by different foreign MHC alleles to promote alloreactivity. This enormous receptor plasticity is bestowed via a number of different mechanisms, as TCR binding can vary from being rigid and very focused [62] to being flexible in terms of both binding register and individual CDR loops (Fig. 6). It is also possible that the ability of T cells to respond to different TCR ligands could be varied during T cell development. TCR engagement in the presence of inhibitory signals is likely to dampen the number of ligands, whereas co-stimulatory signals would be expected to generate a greater level of T cell sensitivity and might increase the number of ligands that could be recognized (Fig. 7a,b). The co-receptors, CD8 and CD4, fall into a class of their own by co-engaging the pMHC ligand, thereby generating capacity

for the molecules to alter which TCR ligands a T cell can respond to [66,67] (Fig. 7c).

Implications of T cell cross-reactivity

It is becoming apparent that effective immune cover requires that each TCR must allow the T cell that bears it to respond to a large array of different peptide ligands [60]. This extensive receptor cross-reactivity has a number of important consequences. First, it allows a relatively small number of TCRs (~25 million [68]) to provide effective immune cover for a vastly greater number of theoretical foreign peptides that could be encountered. Extensive T cell cross-reactivity also ensures that far fewer T cells are required to scan a cell displaying a foreign peptide before one reacts to this peptide, thus ensuring a more rapid response time. The corollary of far-reaching TCR cross-reactivity is that each peptide will be




Fig. 7. The peptide cross-reactivity of conventional T cells can be varied. In order to be positively selected in the thymus, a T cell must bear a T cell receptor (TCR) that allows it to recognize – and respond to – self-peptide. It should not respond to this peptide thereafter. T cell cross-reactivity could be regulated throughout the lifetime of a T cell. (a) Co-inhibitory (in red) or (b) co-stimulatory signals (in green) are likely to decrease and increase the number of ligands that can be recognized by tuning the sensitivity of TCR engagement that a T cell responds to. (c) The co-receptors, CD4 and CD8 (in purple), represent a special class of co-stimulation as these receptors bind to the same peptide–major histocompatibility complex (pMHC) ligand at the TCR. This could allow the co-receptors to discriminate between different TCR–pMHC dwell times and thus tune a T cell to recognize only certain ligands [66]. Regulation of the cell surface expression and/or glycosylation of key receptors might also be used to vary sensitivity to cross-reactive TCR ligands.

recognized by several different receptors (i.e. T cell responses are polyclonal). Escape from a polyclonal T cell response represents a far greater challenge for pathogens that can vary their antigens, as a mutation that escapes from one TCR may still be recognized by a different TCR. Heterologous immunity, where a single T cell can respond to several infections, is a further important consequence of individual TCRs being capable of responding to multiple peptides [69]. Once it is realized that T cells can respond to many different peptides it should come as little surprise that there are pre-existing populations of HIV-specific memory T cells in people who are uninfected with this virus [70]. The extent and implications of T cell heterologous immunity are described in detail elsewhere [13,69]. The advantages of having a broadly cross-reactive T cell repertoire must outweigh the negatives, or the system would have been expected to evolve differently. Nevertheless, it is believed that widely cross-reactive T cells can have detrimental consequences. Most notably, the concept of T cells becoming activated by pathogens and then cross-recognizing self-ligands – a phenomenon known as molecular mimicry [71] – is

believed to be the root cause of autoimmune disease (Fig. 8). A further important consequence of TCR binding plasticity is alloreactive recognition of non-self-HLA presenting self-peptide because such recognition causes acute rejection of HLA-mismatched grafted organs [72]. Alloreactive recognition of pMHC by the TCR represents the major barrier to organ transplantation. Most transplants are HLA mismatched and require that the recipient take lifelong immunosuppressive medication, with its associated expense and side effects.

Therapeutic use of the $\alpha\beta$ TCR

The transfer of TCR genes into recipient host T cells followed by the adoptive transfer of these cells to patients allows the passive transfer of immunity [73]. This strategy can provide a convenient means for breaking tolerance to tumour antigens and has already shown promise in patients with malignant melanoma [74]. As described above, T cell immunity is a compromise where a limited number of receptors are required to provide immune cover for a vastly greater number of potential foreign




Fig. 8. T cell cross-reactivity causes autoimmunity. T cells bearing autoreactive T cell receptors (TCRs) sometimes escape from thymic culling and populate the peripheral tissues. Such cells usually bear TCRs that bind very weakly to self-peptide and generally remain harmless. However, if such a T cell becomes activated in response to a pathogen-derived peptide it will be stimulated to become an effector T cell. Antigen-experienced cells are known to be more sensitive to TCR triggering. Activation of such cells by a cross-recognized self-peptide could then result in autoimmune attack.

peptides bound to self-MHC. This compromise ensures that individual TCR–pMHC interactions are rarely optimal. Recent developments mean that TCRs can now be affinity-enhanced by phage display [75], yeast display [76] and computational design [77,78]. Such enhancement has been used to build immune foresight by selecting TCRs that can recognize all known escape variants of HIV [79]. Enhanced TCRs are particularly useful for cancer immunotherapy, as thymic selection culls T cells whose TCRs strongly recognize self-antigens. This process is presumably responsible for the finding that natural anti-tumour TCRs bind with substantially weaker affinity than anti-pathogen TCRs [80,81]. This leaves cancer-specific T cells at a distinct disadvantage compared to their counterparts that respond to non-self-peptides. The enhancement processes described above can now be used to close the TCR binding affinity gap between optimal anti-pathogen TCRs and weaker anti-tumour TCRs. Enhanced optimal TCRs can then be used in TCR gene therapy for cancer [82]. Such approaches are currently showing great promise. However, as enhanced TCRs have not undergone the rigours of thymic selection, where cells with TCRs that react strongly to self are deleted, they carry an inherent, but small, risk of being autoreactive. A recent trial of a TCR specific for the cancer-specific protein melanoma-associated antigen (MAGE) highlighted the potential problem. When cells with an enhanced MAGE-specific TCR were transfused back into two cancer patients, these patients developed rapid and fatal heart disease. Subsequent studies determined that the modified MAGE-specific TCR was also capable of cross-reacting with an MHC-I-presented peptide from the heart protein titin [83,84]. Despite these teething problems, other trials with other receptors have been successful, and we anticipate that the use of such therapies will become commonplace in the next 20 years. Many groups are currently examining the use of various hybrid antigen receptors in

T cell therapy. Such strategies are beyond the scope of this review, but have been documented recently elsewhere [85]. Enhanced TCRs have also recently been used as soluble molecules to induce cancer regression [86], thus opening up a further exciting mechanism by which TCRs can be used for therapeutic benefit (Fig. 9). The use of TCRs as soluble ‘drugs’ does not incorporate the same dangers as cell-based TCR therapies, as dosage can be scaled or medication withdrawn if difficulties arise. The potential for commercial use of TCRs has been assessed elsewhere [87].

Unconventional T cells

The list of unconventional T cells that do not recognize pMHC ligands is growing steadily, further demonstrating the versatility of the TCR. Recent discoveries have highlighted the existence of $\alpha\beta$ T cells that recognize non-peptide ligands. Approximately 10% of all $\alpha\beta$ T cells are now thought to recognize lipid antigens. A further 10% of human $\alpha\beta$ T cells appear to recognize bacterial metabolites.

$\alpha\beta$ TCR recognition of lipids

Many human $\alpha\beta$ T cells have been identified recently that respond to non-peptide antigens presented by MHC class I-related molecules from the CD1 protein family [88]. These glycoproteins are ideally suited for presentation of lipid-based antigens to T cells due to the hydrophobic nature of their deep antigen-binding pockets. There are five isoforms of CD1 molecules in humans, CD1a–e, although CD1e is not involved in antigen presentation. The antigen-binding clefts of CD1a–d differ in shape and size permitting the presentation of different lipids to T cells (Fig.10). T cells restricted by the group 1 CD1 molecules (CD1a–c) are considerably more numeric than CD1d-restricted T cells, but far less is known about these T cells and the TCRs that they




Fig. 9. Soluble T cell receptor (TCR) therapy. (a) The MHC-I antigen presentation pathway is predicted to present at least one peptide from any protein present in a cell at 500 copies or more. This clever system allows the TCRs on the surface of MHC-I-restricted T cells to inspect the cellular proteome from the cell surface and detect internal anomalies. This 'X-ray vision' allows TCRs to access a far greater number of cellular targets than are available to monoclonal antibodies. Phage-display and directed evolution of TCRs [75] can generate very high-affinity molecules ($K_D < 10$ pM) that bind to the cognate peptide-MHC (pMHC) with very long half-lives (several hours). These molecules can then be used to deliver therapeutic payloads to specific cells *in vivo*. (b) High-affinity tumour-specific TCRs can be 'fused' to a CD3-specific Fab fragment to produce a bi-specific molecule. Such molecules have recently been used to induce cancer regression [86].

express. At the time of writing, structures of group 1 CD1-restricted TCRs in complex with their antigen are only just starting to emerge. It is expected that further structures will appear soon in the literature.

CD1d-lipid complexes bind to TCRs that are expressed by the innate-like NK T cells [93]. CD1d-restricted T cells are divided into two types based on their TCR gene usage and the antigens to which they respond. Human type I NK T cells respond strongly to the lipid α -GalCer. These T cells typically utilize a semi-invariant TCR, and have been termed iNK T cells to reflect this. The type I NK T TCR uses an invariant TCR- α chain (TRAV10/TRAJ18) paired with a TRBV25.1-encoded β chain. Type II NK T cells express a broader TCR repertoire and are not activated by α -GalCer. NK T TCR recognition appears to be far more rigid than that observed for conventional recognition of pMHC. Type I NK T TCRs adopt a common footprint on CD1d (Fig. 11a). This recognition has been compared to an innate pattern recognition receptor. Recent type II NK T TCR-CD1d-sulphatide and lysosulphatide complexes have revealed that the mode of type II NK T TCR recognition is different to type I NK T TCR recognition [97]. Type II NK T cells dock orthogonally and over the A'-pocket of CD1d and make a larger binding interface than type I NK T TCRs. The binding mode of type II NK T TCRs is closer to that used by conventional TCR docking on pMHC.

GEM T cells recognize mycobacteria-derived (glyco)-lipids in the context of CD1b, a molecule which has the potential to accommodate the largest and most diverse lipid ligands among the CD1 protein family [90]. Although no ternary structure of a TCR bound to the CD1b-lipid complex has been resolved so far, a recent study suggested that binding of the lipid antigen to CD1b could induce a conformational change of the latter. Thus, the TCR recognition would be mediated not only by the solvent-exposed polar headgroup of the lipid, but also by the rearranged residues from the antigen-presenting molecule [90]. CD1c is also capable of presenting mycobacterial lipids and lipopeptides [98]. Similarly to CD1b, TCR recognition of CD1c seems to be mediated by both the lipid antigen and CD1c residues; however, without showing a common pattern recognised by all TCRs responding to the same antigen [91]. Interestingly, CD1c can also present immunogenic self-derived lipids accumulated in leukaemia cells, contributing to $\alpha\beta$ T cell-mediated tumour surveillance [99].

Contrary to CD1b-d, CD1a can bind lipids lacking a polar headgroup. These hydrophobic lipid antigens are then not recognized directly by the TCR, but rather allow CD1a to adapt to a TCR-activating conformation [89] (Fig. 11b). In contrast, lipids containing a polar headgroup can also be bound by CD1a but disrupt the activating conformation of the antigen-presenting molecule, thus abolishing TCR-mediated recognition.

MR1-restricted $\alpha\beta$ TCRs

It has been shown recently that MAIT cells recognize intermediates in the riboflavin biosynthesis pathway




Fig. 10. CD1a–d presentation of lipids. (a) CD1a can bind lipids not endowed with polar headgroups. Such lipids do not disrupt the T cell receptor (TCR)-activating conformation of the CD1a molecule. The presented ligand is lysophosphatidylcholine (LPC) [Protein Data Bank (PDB): 4X6E] [89]. (b) CD1b can potentially present the largest and most diverse lipids of all CD1-family proteins. The lipid ligands can possess one, two or three alkyl chains buried in the hydrophobic pockets of CD1b while their polar headgroup can be recognized by germline-encoded mycolyl-reactive (GEM) T cells. The presented ligand is ganglioside GM2 (PDB: 1GZP) [90]. (c) CD1c can present mycobacterial lipids such as phosphomycoketide (shown) to $\alpha\beta$ T cells (PDB: 4ONO) [91]. (d) CD1d can present self-derived lipids such as sulphatide (shown) or α -galactosylceramide (α -GalCer) to both $\alpha\beta$ [type I and II natural killer (NK) T cells] and $\gamma\delta$ T cells (PDB: 4MQ7) [92].

bound to the MHC-like molecule MR1 [100–102]. MAIT cells constitute about 10% of all T cells in humans, so recognition is likely to represent the most dominant of all T cell antigen-specificities. Vitamin B is synthesized by some bacteria and yeast but not by mammals. Thus, such microbial-specific biosynthetic pathways provide a further mechanism that T cells can use to distinguish self from non-self. MAIT TCRs bind to riboflavin precursors and the presenting MR1 molecule (Fig. 11c). Interestingly, MR1 can also present non-activating metabolites belonging to the vitamin B group, namely folic acid derivatives. The invariant TCR- α chain binds MR1 in a conserved, innate-like manner, thus allowing the TCR to interact directly with the activating riboflavin precursor, while no such contact is made if the MR1-bound ligand is a folic acid derivative [95,103]. MAIT cells have set a new paradigm in T cell recognition. As yet it is unclear how many other bacterial metabolites can be recognized by T cells, as a recent study by Corbett *et al.* [104] showed that MR1 is capable of capturing and presenting unstable intermediates of riboflavin synthesis, which would otherwise be converted into

non-activating metabolites. However, it is established that the dominant subset of $\gamma\delta$ T cells in human peripheral blood recognize intermediates in the non-mevalonate microbial pathway of isoprenoid biosynthesis, so it remains possible that a sizable fraction of human T cells also target other microbial-specific synthetic pathways.

$\gamma\delta$ TCRs

Surprisingly little is known about human $\gamma\delta$ T cells or the ligands they recognize. Unlike $\alpha\beta$ T cells, $\gamma\delta$ T cells exhibit strict tissue-specific localization, which is determined by the identity of the rearranged TCR. This is because gene rearrangement at the *trg* and *trd* loci is programmed developmentally and different $\gamma\delta$ T cells carrying distinct TCR- γ and TCR- δ rearrangements arise in successive waves that colonize different tissues during embryonic development [105]. The tissue specificity of $\gamma\delta$ T cells may also reflect a difference in the expression of $\gamma\delta$ TCR ligands. Should this prove true, then $\gamma\delta$ TCRs and their ligands might become very useful for tissue




Fig. 11. Recognition of antigen by 'unconventional' T cell receptors (TCRs). (a) Type I natural killer (NK) TCR recognition. The TCR binds CD1d- α -galactosylceramide (α GalCer) in a parallel docking mode. The recognition is markedly different from any known $\alpha\beta$ TCR-peptide-MHC (pMHC) interactions [Protein Data Bank (PDB): 3VWK] [94]. (b) Recognition of CD1a-lipid complex. $\alpha\beta$ TCR recognizes a permissive conformation of CD1a surface that depends on the nature of the lipid ligand bound by the latter. Contrary to TCR-CD1d structures, no direct interactions between TCR and a lipid ligand bound to CD1a have been observed (PDB: 4X6C) [89]. (c) Mucosal-associated invariant T (MAIT) TCR recognition of MR1. MR1 can present both activating (riboflavin precursors) and non-activating (folic acid precursors) intermediates from vitamin B biosynthetic pathways. The recognition of MR1 is mediated by conserved, invariant residues in an innate-like manner. The activating metabolites directly contact the TCR while no such interactions are observed for the non-activating metabolites (PDB: 4L4V) [95]. (d) A hybrid $\delta/\alpha\beta$ TCR binds the CD1d- α GalCer complex. The TCR is composed of a variable V δ 1 domain fused with joining and constant α domains, and paired to a TCR- β chain. TCR-CD1d interactions are driven mainly by the germline-encoded V δ 1 residues while TCR- β mediates specific lipid recognition (PDB: 4W04) [7]. (e) The $\gamma\delta$ TCR recognizes CD1d-lipid complexes. Germline-encoded V δ 1 residues are responsible for CD1d binding while the lipid ligand recognition is fine-tuned by hypervariable complementarity-determining region (CDR) 3 loops (PDB: 4MNG) [92]. (f) A mouse $\gamma\delta$ TCR binds the MHC-like molecule T22. The binding occurs predominantly via a conserved motif within the hypervariable CDR3 δ loop, whereas the non-conserved CDR3 δ residues fine-tune the affinity of the receptor towards T22 (PDB: 1YPZ) [96].

targeting. It remains to be seen whether $\gamma\delta$ TCRs can ever bind to pMHC ligands. Nevertheless, structural studies of the complex between a murine $\gamma\delta$ TCR and an MHC class I-like molecule T22 (Fig. 11f) indicate that at least some $\gamma\delta$ TCRs can bind their ligand in a manner reminiscent of $\alpha\beta$ TCR-MHC interactions [96,106].

Most progress with $\gamma\delta$ TCRs has been made with the predominant subset of $\gamma\delta$ T cells found in human peripheral blood that express a TCR made from the V γ 9 and V δ 2 genes. V γ 9⁺V δ 2⁺ T cells have been shown to recognize small pyrophosphate intermediates of the lipid biosynthetic pathway used by some bacteria [107,108]. Notably, pyrophosphate antigens can accumulate in neoplastic cells as well, as a result of metabolic deregulation [109]. While the structure of a V γ 9V δ 2 TCR has been solved [110], there is not yet a structure of this TCR bound to its ligand. Recog-

nition of pyrophosphate ligands requires cellular presentation by a primate cell [111]. This species specificity suggested that pyrophosphate antigens might be presented to V γ 9⁺V δ 2⁺ T cells by an MHC-like molecule. Recently the candidate for a pyrophosphate antigen-presenting molecule has been narrowed down to butyrophilin 3A1 (BTN3A1), an immunoglobulin-like molecule encoded in the MHC I locus. Crystallographic studies revealed that pyrophosphate antigens can be bound either in a shallow extracellular pocket of BTN3A1 and presented directly to T cells [112] or in an intracellular domain, enforcing a TCR-activating conformational change of the presenting molecule [113]. These mechanisms, however, are not necessarily mutually exclusive, as both intra- and extracellular binding of phosphoantigens may be required for efficient TCR recognition.




Fig. 12. The versatility of the human T cell receptor (TCR). The $\alpha\beta$ TCR can engage intermediates in riboflavin biosynthesis in the context of MR1 and a variety of lipid molecules bound to CD1a, CD1b, CD1c and CD1d. Classically, this receptor is also capable of binding to the almost infinite array of different peptides bound to more than 12 000 human leucocyte antigen (HLA) alleles of the HLA-A, -B, -C, -DR, -DQ and -DP loci. $\gamma\delta$ TCRs can also bind to CD1d-lipid complexes [103,108] and a variety of ectopically expressed cell stress markers including endothelial protein C receptor (EPCR) [107] (shown), MIC A/B [114] and hMSH2 [115]. Murine $\gamma\delta$ TCRs bind to the MHC-I like molecules T10 and T22 that are not found in humans [96]. The best-studied set of human $\gamma\delta$ T cells express V γ 9V δ 2 TCRs and recognize pyrophosphate antigens in the context of an immunoglobulin-like molecule, butyrophillin 3A1 [112]. The exact mechanism by which these TCRs recognize phosphoantigens awaits a TCR-ligand structure.

Similarly to ‘unconventional’ $\alpha\beta$ TCRs, $\gamma\delta$ TCRs can mediate recognition of lipids in the context of CD1d [92]. However, the docking manner differed substantially from type I and II NK T cells (Fig. 11e). Importantly, the contacts with CD1d were mediated mainly via germline-encoded residues in CDR1 and 2 loops, while hypervariable CDR3 loops bound the presented lipid. Therefore, it is possible that $\gamma\delta$ TCR can distinguish between subtly different lipid cargoes bound to CD1d in an adaptive-like manner. Additionally, Willcox *et al.* [107] have demonstrated recently that $\gamma\delta$ TCRs can bind a ligand shared by cytomegalovirus-infected and malignantly transformed cells, namely endothelial protein C receptor (EPCR). EPCR shows homology with CD1 protein family at the level of amino acid sequence and structure, and is also capable of presenting lipids – however, the binding of the TCR to EPCR was lipid-independent (Fig. 12). The recog-

nition was mediated via the hypervariable CDR3 γ loop of the TCR, thus suggesting that EPCR binding was a part of an adaptive response rather than innate-like pattern recognition. Notably, EPCR was essential, but not sufficient, for the recognition of the target cells, highlighting the need for additional accessory molecules that would create a generalized cellular stress context. The identity of these accessory molecules on the T cell or target cell remains largely unknown.

EPCR is not the only example of a $\gamma\delta$ TCR ligand recognized in a cellular stress context. It has been known for some time that a portion of human $\gamma\delta$ T cells can bind MHC class I polypeptide-related sequence (MIC) A [114], a stress-related ligand of natural killer cells, and human homologue of the bacterial MutS protein (hMSH2) [115], an element of the DNA repair system. Both molecules can be up-regulated and expressed

ectopically upon DNA damage, oxidative stress, malignant transformation and Epstein–Barr virus infection [116–118]. Importantly, both MIC A [119] and hMSH2 [116] were dually recognized by the TCR and a natural killer receptor NKG2D. Biophysical studies indicated that the ligand (MIC A) was bound initially by NKG2D forming a transient complex, giving way to a more stable TCR–ligand complex [119]. It is therefore possible that this sequential recognition forms a critical part of immune surveillance, allowing the T cells to detect signs of cellular stress rapidly which could, in turn, indicate virally infected or transformed cells. The requirement for a multi-component stress signature could also serve as a preventive measure against triggering autoimmune reactions – a matter of particular importance when one considers that most of the known $\gamma\delta$ TCR ligands are essentially self-derived. In line with this fact, $\gamma\delta$ T cells have been shown to contribute to autoimmune diseases such as myositis and type I diabetes, as they can recognize aminoacyl-tRNA synthetases [120] and an insulin-derived peptide [121], respectively. The role of $\gamma\delta$ T cells in pathogenesis of autoimmune diseases has been reviewed recently elsewhere [122].

Summary

It has been known for several decades that the vast majority of TCRs recognize MHC-bound peptides. Recent developments have highlighted the astonishing versatility of this receptor (Fig. 12). Indeed, it is now widely accepted that a substantial fraction of T cell pool is dedicated to the recognition of non-MHC ligands. This includes recognition of various foreign lipids and metabolic intermediates and stress-related proteins. However, in the case of $\gamma\delta$ T cells, the identity of putative ligands remains largely unknown. Although monoclonal antibodies now account for almost half of the new drugs that come onto the market, several TCR-based therapeutic strategies are in development. Such strategies are likely to have a bright future, and further exciting advances are anticipated shortly.

Acknowledgements

A. K. S. and D. K. C. are Wellcome Trust Senior Investigators and Career Development Fellows, respectively. M. L. is a Cancer Research UK PhD student. Our work on TCR degeneracy is funded by the Biotechnology and Biological Sciences Research Council UK.

Disclosure

The authors declare no conflicts of interest.

References

- Zinkernagel RM, Doherty PC. Restriction of *in vitro* T cell-mediated cytotoxicity in lymphocytic choriomeningitis within a syngeneic or semiallogeneic system. *Nature* 1974; **248**:701–2.

- Treiner E, Duban L, Bahram S et al. Selection of evolutionarily conserved mucosal-associated invariant T cells by MR1. *Nature* 2003; **422**:164–9.
- Le Bourhis L, Martin E, Peguillet I et al. Antimicrobial activity of mucosal-associated invariant T cells. *Nat Immunol* 2010; **11**: 701–8.
- Beckman EM, Porcelli SA, Morita CT, Behar SM, Furlong ST, Brenner MB. Recognition of a lipid antigen by CD1-restricted $\alpha\beta$ T cells. *Nature* 1994; **372**:691–4.
- Van Rhijn I, Kasmar A, de Jong A et al. A conserved human T cell population targets mycobacterial antigens presented by CD1b. *Nat Immunol* 2014; **14**:706–13.
- Carding SR, Egan PJ. $\gamma\delta$ T cells: functional plasticity and heterogeneity. *Nat Rev Immunol* 2002; **2**:336–45.
- Pellicci DG, Uldrich AP, Le Nours J et al. The molecular bases of $\delta/\alpha\beta$ T cell-mediated antigen recognition. *J Exp Med* 2014; **211**:2599–615.
- Allam A, Kabelitz D. TCR trans-rearrangements: biological significance in antigen recognition vs the role as lymphoma biomarker. *J Immunol* 2006; **176**:5707–12.
- Bowen S, Sun P, Livak F, Sharrow S, Hodes RJ. A novel T cell subset with trans-rearranged V γ -C β TCRs shows V β expression is dispensable for lineage choice and MHC restriction. *J Immunol* 2014; **192**:169–77.
- Cole DK, Yuan F, Rizkallah PJ et al. Germ line-governed recognition of a cancer epitope by an immunodominant human T-cell receptor. *J Biol Chem* 2009; **284**:27281–9.
- Allison TJ, Winter CC, Fournié JJ, Bonneville M, Garboczi DN. Structure of a human gammadelta T-cell antigen receptor. *Nature* 2001; **411**:820–4.
- Gras S, Chen Z, Miles JJ et al. Allelic polymorphism in the T cell receptor and its impact on immune responses. *J Exp Med* 2010; **207**:1555–67.
- Sewell AK. Why must T cells be cross-reactive? *Nat Rev Immunol* 2012; **2**:669–77.
- Davis MM, Bjorkmann PJ. T cell antigen receptor genes and T cell recognition. *Nature* 1988; **334**:395–402.
- Van Boehmer H, Teh HS, Kisielow P. The thymus selects the useful, neglects the useless and destroys the harmful. *Immunol Today* 1989; **10**:57–61.
- Bendelac A. Positive selection of mouse NK1+ T cells by CD1-expressing cortical thymocytes. *J Exp Med* 1982; **156**:2091–6.
- Coles MC, Raulet DH. NK1.1+ T cells in the liver arise in the thymus and are selected by interactions with class I molecules on CD4+CD8+ cells. *J Immunol* 2000; **164**:2412–8.
- Wei DG, Lee H, Park HS et al. Expansion and long-range differentiation of the NKT cell lineage in mice expressing CD1d exclusively on cortical thymocytes. *J Exp Med* 2005; **202**:239–48.
- Seach N, Guerri L, Le Bourhis L et al. Double-positive thymocytes select mucosal-associated invariant T cells. *J Immunol* 2013; **191**:6002–9.
- Wilson A, de Villartay JP, MacDonal HR. T cell receptor δ gene rearrangement and T early α (TEA) expression in immature $\alpha\beta$ lineage thymocytes: implications for $\alpha\beta/\gamma\delta$ lineage commitment. *Immunity* 1996; **4**:37–45.
- Coffey F, Lee SY, Buus TB et al. The TCR ligand-inducible expression of CD73 marks $\gamma\delta$ lineage commitment and a metastable intermediate in effector specification. *J Exp Med* 2014; **211**:329–43.

- 22 Turchinovich G, Hayday AC. Skint-1 identifies a common molecular mechanism for the development of interferon- γ -secreting versus interleukin-17-secreting $\gamma\delta$ T cells. *Immunity* 2011; **35**:59–68.
- 23 Jensen KD, Su X, Shin S *et al.* Thymic selection determines gammadelta T cell effector fate: antigen-naive cells make interleukin-17 and antigen-experienced cells make interferon gamma. *Immunity* 2008; **29**:90–100.
- 24 Robinson J, Halliwell JA, Hayhurst JD, Flicek P, Parham P, Marsh SG. The IPD and IMGT/HLA database: allele variant databases. *Nucleic Acids Res* 2015; **43**:D423–31.
- 25 Tynan FE, Borg NA, Miles JJ *et al.* High resolution structures of highly bulged viral epitopes bound to major histocompatibility complex class I. Implications for T-cell receptor engagement and T-cell immunodominance. *J Biol Chem* 2005; **280**: 23900–9.
- 26 Mullen MM, Haan KM, Longnecker R, Jardetzky TS. Structure of the Epstein–Barr virus gp42 protein bound to the MHC class II receptor HLA-DR1. *Mol Cell* 2002; **9**:375–85.
- 27 Ekeruche-Makinde J, Miles JJ, van den Berg HA *et al.* Peptide length determines the outcome of TCR/peptide–MHC engagement. *Blood* 2013; **121**:1112–23.
- 28 Probst-Kepper M, Hecht HJ, Herrmann H *et al.* Conformational restraints and flexibility of 14-meric peptides in complex with HLA-B*3501. *J Immunol* 2004; **173**:5610–6.
- 29 Miles JJ, Borg NA, Brennan RM *et al.* TCR alpha genes direct MHC restriction in the potent human T cell response to a class I-bound viral epitope. *J Immunol* 2006; **177**:6804–14.
- 30 Wynn KK, Fulton Z, Cooper L *et al.* Impact of clonal competition for peptide–MHC complexes on the CD8+ T-cell repertoire selection in a persistent viral infection. *Blood* 2008; **111**: 4283–92.
- 31 Smith KJ, Reid SW, Stuart DI, McMichael AJ, Jones EY, Bell JI. An altered position of the alpha 2 helix of MHC class I is revealed by the crystal structure of HLA-B*3501. *Immunity* 1996; **4**:203–13.
- 32 Sliz P, Michielin O, Cerottini JC *et al.* Crystal structures of two closely related but antigenically distinct HLA-A2/melanocyte–melanoma tumor–antigen peptide complexes. *J Immunol* 2001; **167**:3276–84.
- 33 Madden DR, Garboczi DN, Wiley DC. The antigenic identity of peptide–MHC complexes: a comparison of the conformations of five viral peptides presented by HLA-A2. *Cell* 1993; **75**:693–708.
- 34 Siebold C, Hansen BE, Wyer JR *et al.* Crystal structure of HLA-DQ0602 that protects against type 1 diabetes and confers strong susceptibility to narcolepsy. *Proc Natl Acad Sci USA* 2004; **101**:1999–2004.
- 35 Dessen A, Lawrence CM, Cupo S, Zaller DM, Wiley DC. X-ray crystal structure of HLA-DR4 (DRA*0101, DRB1*0401) complexed with a peptide from human collagen II. *Immunity* 1997; **7**:473–81.
- 36 Deng L, Langley RJ, Brown PH *et al.* Structural basis for the recognition of mutant self by a tumor-specific, MHC class II-restricted T cell receptor. *Nat Immunol* 2007; **8**:398–408.
- 37 Rudolph MG, Stanfield RL, Wilson IA. How TCRs bind MHCs, peptides, and coreceptors. *Annu Rev Immunol* 2006; **24**:419–66.
- 38 Hughes AL, Ota T, Nei M. Positive Darwinian selection promotes charge profile diversity in the antigen-binding cleft of class I major-histocompatibility-complex molecules. *Mol Biol Evol* 1990; **7**:515–24.
- 39 Rossjohn J, Gras S, Miles JJ, Turner SJ, Godfrey DI, McCluskey J. T cell antigen receptor recognition of antigen-presenting molecules. *Annu Rev Immunol* 2014; **33**:169–200.
- 40 Miles JJ, Bulek AM, Cole DK *et al.* Genetic and structural basis for selection of a ubiquitous T cell receptor deployed in Epstein–Barr virus infection. *PLOS Pathog* 2010; **6**:e1001198.
- 41 Yin Y, Li Y, Kerzic MC, Martin R, Mariuzza RA. Structure of a TCR with high affinity for self-antigen reveals basis for escape from negative selection. *EMBO J* 2011; **30**:1137–48.
- 42 Merckenschlager M, Graf D, Lovatt M, Bommhardt U, Zamoyska R, Fisher AG. How many thymocytes audition for selection? *J Exp Med* 1997; **186**:1149–58.
- 43 Scott-Browne JP, Crawford E, Young MH, Kappler JW, Marrack P, Gapin L. Evolutionarily conserved features contribute to $\alpha\beta$ T cell receptor specificity. *Immunity* 2011; **35**:526–35.
- 44 Garcia KC, Adams JJ, Feng D, Ely LK. The molecular basis of TCR germline bias for MHC is surprisingly simple. *Nat Immunol* 2009; **10**:143.
- 45 Garcia KC. Reconciling views on T cell receptor germline bias for MHC. *Trends Immunol* 2012; **33**:429–36.
- 46 Feng D, Bond CJ, Ely LK, Maynard J, Garcia KC. Structural evidence for a germline-encoded T cell receptor–major histocompatibility complex interaction ‘codon’. *Nat Immunol* 2007; **8**:975–83.
- 47 Stadinski BD, Trenh P, Smith RL *et al.* A role for differential variable gene pairing in creating T cell receptors specific for unique major histocompatibility ligands. *Immunity* 2011; **35**:694–704.
- 48 Burrows SR, Chen Z, Archbold JK *et al.* Hard wiring of T cell receptor specificity for the major histocompatibility complex is underpinned by TCR adaptability. *Proc Natl Acad Sci USA* 2010; **107**:10608–13.
- 49 Holland SJ, Bartok I, Attaf M *et al.* The T-cell receptor is not hardwired to engage MHC ligands. *Proc Natl Acad Sci USA* 2012; **109**:E3111–8.
- 50 Yin Y, Wang XX, Mariuzza RA. Crystal structure of a complete ternary complex of T-cell receptor, peptide–MHC. *Proc Natl Acad Sci USA* 2012; **109**:5405–10.
- 51 Cole DK, Miles KM, Madura Fet *et al.* T-cell receptor (TCR)-peptide specificity overrides affinity-enhancing TCR–major histocompatibility complex interactions. *J Biol Chem* 2013; **289**: 628–38.
- 52 Gao GF, Tormo J, Gerth UC *et al.* Crystal structure of the complex between human CD8 α (α) and HLA-A2. *Nature* 1997; **387**:630–4.
- 53 Van Laethem F, Sarafova SD, Park JH *et al.* Deletion of CD4 and CD8 coreceptors permits generation of $\alpha\beta$ T cells that recognize antigens independently of the MHC. *Immunity* 2007; **27**:735–50.
- 54 Tikhonova AN, Van Laethem F, Hanada K *et al.* $\alpha\beta$ T cell receptors that do not undergo major histocompatibility complex-specific thymic selection possess antibody-like recognition specificities. *Immunity* 2012; **36**:79–91.
- 55 Van Laethem F, Tikhonova AN, Singer A. MHC restriction is imposed on a diverse T cell receptor repertoire by CD4 and CD8 co-receptors during thymic selection. *Trends Immunol* 2012; **33**:437–41.

- 56 Van Laethem F, Tikhonova AN, Pobeziński LA et al. Lck availability during thymic selection determines the recognition specificity of the T cell repertoire. *Cell* 2013; **154**:1326–41.
- 57 Rao A, Ko WW, Faas SJ, Cantor H. Binding of antigen in the absence of histocompatibility proteins by arsonate-reactive T-cell clones. *Cell* 1984; **36**:879–88.
- 58 Barnd DL, Lan MS, Metzgar RS, Finn OJ. Specific, major histocompatibility complex-unrestricted recognition of tumor-associated mucins by human cytotoxic T cells. *Proc Natl Acad Sci USA* 1989; **86**:7159–63.
- 59 Hanada K, Wang QJ, Inozume T, Yang JC. Molecular identification of an MHC-independent ligand recognized by a human α/β T-cell receptor. *Blood* 2011; **117**:4816–25.
- 60 Mason D. A very high level of crossreactivity is an essential feature of the T-cell receptor. *Immunol Today* 1998; **19**:395–404.
- 61 Wooldridge L, Ekeruche-Makinde J, van den Berg HA et al. A single autoimmune T cell receptor recognizes more than a million different peptides. *J Biol Chem* 2012; **287**:1168–77.
- 62 Bulek AM, Cole DK, Skowera A et al. Structural basis for the killing of human beta cells by CD8+ T cells in type 1 diabetes. *Nat Immunol* 2012; **13**:283–9.
- 63 Birnbaum ME, Mendoza JL, Sethi DK et al. Deconstructing the peptide-MHC specificity of T cell recognition. *Cell* 2014; **157**:1073–87.
- 64 Madura F, Rizkallah PJ, Holland CJ et al. Structural basis for ineffective T-cell responses to MHC anchor residue-improved 'heteroclitic' peptides. *Eur J Immunol* 2015; **45**:584–91.
- 65 Cole DK, Edwards ES, Wynn KK et al. Modification of MHC anchor residues generates heteroclitic peptides that alter TCR binding and T cell recognition. *J Immunol* 2010; **185**:2600–10.
- 66 van den Berg HA, Wooldridge L, Laugel B, Sewell AK. Coreceptor CD8-driven modulation of T cell antigen receptor specificity. *J Theor Biol* **249**:395–408.
- 67 Wooldridge L, Laugel B, Ekeruche J et al. CD8 controls T cell cross-reactivity. *J Immunol* 2010; **185**:4625–32.
- 68 Arstila T, Casrouge A, Baron V, Even J, Kanellopoulos J, Kourilsky P. A direct estimate of the human $\alpha\beta$ T cell receptor diversity. *Science* 1999; **286**:958–61.
- 69 Welsh RM, Selin LK. No one is naive: the significance of heterologous T-cell immunity. *Nat Rev Immunol* 2002; **2**:417–26.
- 70 Su LF, Kidd BA, Han A, Kotzin JJ, Davis MM. Virus-specific CD4+ memory-phenotype T cells are abundant in unexposed adults. *Immunity* 2013; **38**:373–83.
- 71 Wucherpfennig KW, Sethi D. T cell receptor recognition of self and foreign antigens in the induction of autoimmunity. *Semin Immunol* 2011; **23**:84–91.
- 72 Ely LK, Burrows SR, Purcell AW, Rossjohn J, McCluskey J. T-cells behaving badly: structural insights into alloreactivity and autoimmunity. *Curr Opin Immunol* 2008; **20**:575–80.
- 73 Restifo NP, Dudley ME, Rosenberg SA. Adoptive immunotherapy for cancer: harnessing the T cell response. *Nat Rev Immunol* 2012; **12**:269–81.
- 74 Morgan RA, Dudley ME, Wunderlich JR et al. Cancer regression in patients after transfer of genetically engineered lymphocytes. *Science* 2006; **314**:126–9.
- 75 Li Y, Moysey R, Molloy PE et al. Directed evolution of human T-cell receptors with picomolar affinities by phage display. *Nat Biotechnol* 2005; **23**:349–54.
- 76 Kieke MC, Shusta EV, Boder ET, Teyton L, Witttrup KD, Kranz DM. Selection of functional T cell receptor mutants from a yeast surface-display library. *Proc Natl Acad Sci USA* **96**:5651–6.
- 77 Hawse WF, Champion MM, Joyce MV et al. Cutting edge: evidence for a dynamically driven T cell signaling mechanism. *J Immunol* 2012; **188**:5819–23.
- 78 Irving M, Zoete V, Hebeisen M et al. Interplay between T cell receptor binding kinetics and the level of cognate peptide presented by major histocompatibility complexes governs CD8+ T cell responsiveness. *J Biol Chem* **287**:23068–78.
- 79 Varela-Rohena A, Molloy PE, Dunn SM et al. Control of HIV-1 immune escape by CD8 T cells expressing enhanced T-cell receptor. *Nat Med* 2008; **14**:1390–5.
- 80 Aleksic M, Liddy N, Molloy PE et al. Different affinity windows for virus and cancer-specific T-cell receptors: implications for therapeutic strategies. *Eur J Immunol* **42**:3174–9.
- 81 Cole DK, Pumphrey NJ, Boulter JM et al. Human TCR-binding affinity is governed by MHC class restriction. *J Immunol* 2007; **178**:5727–34.
- 82 Tan MP, Gerry AB, Brewer JE et al. TCR binding affinity governs the functional profile of cancer-specific CD8+ T cells. *Clin Exp Immunol* in press 2014.
- 83 Cameron BJ, Gerry AB, Dukes J et al. Identification of a Titin-derived HLA-A1-presented peptide as a cross-reactive target for engineered MAGE A3-directed T cells. *Sci Transl Med* 2013; **5**:197ra103.
- 84 Linette GP, Stadtmauer EA, Maus MV et al. Cardiovascular toxicity and titin cross-reactivity of affinity-enhanced T cells in myeloma and melanoma. *Blood* 2013; **22**:863–71.
- 85 Barrett DM, Singh N, Porter DL, Grupp SA, June CH. Chimeric antigen receptor therapy for cancer. *Annu Rev Med* 2014; **65**:333–47.
- 86 Liddy N, Bossi G, Adams KJ et al. Monoclonal TCR-redirectioned tumor cell killing. *Nat Med* 2012; **18**:980–7.
- 87 Sheridan C. Monoclonal T-cell receptor drugs pique pharma's interest. *Nat Biotechnol* 2013; **31**:950–1.
- 88 Mori L, De Libero G. T cells specific for lipid antigens. *Immunol Res* 2012; **53**:191–9.
- 89 Birkinshaw RW, Pellicci DG, Cheng TY et al. $\alpha\beta$ T cell antigen receptor recognition of CD1a presenting self lipid ligands. *Nat Immunol* 2015; **16**:258–66.
- 90 Gadola SD, Zaccari NR, Harlos K et al. Structure of human CD1b with bound ligands at 2.3 Å, a maze for alkyl chains. *Nat Immunol* 2002; **3**:721–6.
- 91 Roy S, Ly D, Li NS et al. Molecular basis of mycobacterial lipid antigen presentation by CD1c and its recognition by $\alpha\beta$ T cells. *Proc Natl Acad Sci USA* 2014; **111**:E4648–57.
- 92 Luoma AM, Castro CD, Mayassi T et al. Crystal structure of V δ 1 T cell receptor in complex with CD1d-sulfatide shows MHC-like recognition of a self-lipid by human $\gamma\delta$ T cells. *Immunity* **39**:1032–42.
- 93 Borg NA, Wun KS, Kjer-Nielsen L et al. CD1d-lipid-antigen recognition by the semi-invariant NKT T-cell receptor. *Nature* 2007; **448**:44–9.
- 94 Wun KS, Ross F, Patel O et al. Human and mouse type I natural killer T cell antigen receptors exhibit different fine specificities for CD1d-antigen complex. *J Biol Chem* 2012; **287**:39139–48.

- 95 Patel O, Kjer-Nielsen L, Le Nours J *et al.* Recognition of vitamin B metabolites by mucosal-associated invariant T cells. *Nat Commun* 2013; **4**:2142.
- 96 Adams EJ, Chien YH, Garcia KC. Structure of a gammadelta T cell receptor in complex with the nonclassical MHC T22. *Science* 2005; **308**:227–31.
- 97 Patel O, Pellicci DG, Gras S *et al.* Recognition of CD1d-sulfatide mediated by a type II natural killer T cell antigen receptor. *Nat Immunol* 2007; **13**:857–63.
- 98 Scharf L, Li NS, Hawk AJ *et al.* The 2.5 Å structure of CD1c in complex with a mycobacterial lipid reveals an open groove ideally suited for diverse antigen presentation. *Immunity* 2010; **33**:853–62.
- 99 Lepore M, de Lalla C, Gundimeda SR *et al.* A novel self-lipid antigen targets human T cells against CD1c(+) leukemias. *J Exp Med* 2014; **211**:1363–77.
- 100 Kjer-Nielsen L, Patel O, Corbett AJ *et al.* MR1 presents microbial vitamin B metabolites to MAIT cells. *Nature* 2012; **491**:717–23.
- 101 Reantragoon R, Kjer-Nielsen L, Patel O *et al.* Structural insight into MR1-mediated recognition of the mucosal associated invariant T cell receptor. *J Exp Med* 2012; **209**:761–74.
- 102 McWilliam HE, Birkinshaw RW, Villadangos JA, McCluskey J, Rossjohn J. MR1 presentation of vitamin B-based metabolite ligands. *Curr Opin Immunol* 2015; **34C**:28–34.
- 103 Eckle SB, Birkinshaw RW, Kostenko L *et al.* A molecular basis underpinning the T cell receptor heterogeneity of mucosal-associated invariant T cells. *J Exp Med* 2014; **211**:1585–600.
- 104 Corbett AJ, Eckle SB, Birkinshaw RW *et al.* T-cell activation by transitory neo-antigens derived from distinct microbial pathways. *Nature* 2014; **509**:361–5.
- 105 Cowan JE, Jenkinson WE, Anderson. The thymus medulla fosters generation of natural Treg cells, invariant $\gamma\delta$ T Cells and invariant NKT cells: what we learn from intrathymic migration. *Eur J Immunol* 2015; **45**:652–60.
- 106 Sandstrom A, Scharf L, McRae G, Hawk AJ, Meredith SC, Adams EJ. $\gamma\delta$ T cell receptors recognize the non-classical major histocompatibility complex (MHC) molecule T22 via conserved anchor residues in a MHC peptide-like fashion. *J Biol Chem* 2012; **287**:6035–43.
- 107 Willcox CR, Pitard V, Netzer S *et al.* Cytomegalovirus and tumor stress surveillance by binding of a human $\gamma\delta$ T cell antigen receptor to endothelial protein C receptor. *Nat Immunol* 2012; **13**:872–9.
- 108 Uldrich AP, Le Nours J, Pellicci DG *et al.* CD1d-lipid antigen recognition by the $\gamma\delta$ TCR. *Nat Immunol* 2013; **14**:1137–45.
- 109 Gober HJ, Kistowska M, Angman L, Jenö P, Mori L, De Libero G. Human T cell receptor gammadelta cells recognize endogenous mevalonate metabolites in tumor cells. *J Exp Med* 2003; **197**:163–8.
- 110 Morita CT, Beckman EM, Bukowski JF *et al.* Direct presentation of nonpeptide prenyl pyrophosphate antigens to human gamma delta T cells. *Immunity* 1995; **3**:495–507.
- 111 Tanaka Y, Morita CT, Tanaka Y, Nieves E, Brenner MB, Bloom BR. Natural and synthetic non-peptide antigens recognized by human gamma delta T cells. *Nature* 1995; **375**:155–8.
- 112 Vavassori S, Kumar A, Wan GS *et al.* Butyrophilin 3A1 binds phosphorylated antigens and stimulates human $\gamma\delta$ T cells. *Nat Immunol* 2013; **14**:908–16.
- 113 Sandstrom A, Peigné CM, Léger A *et al.* The intracellular B30.2 domain of butyrophilin 3A1 binds phosphoantigens to mediate activation of human V γ 9V δ 2 T cells. *Immunity* 2014; **40**:490–500.
- 114 Groh V, Steinle A, Bauer S, Spies T. Recognition of stress-induced MHC molecules by intestinal epithelial gammadelta T cells. *Science* 1998; **279**:1737–40.
- 115 Chen H, He X, Wang Z *et al.* Identification of human T cell receptor gammadelta-recognized epitopes/proteins via CDR3delta peptide-based immunobiochemical strategy. *J Biol Chem* 2008; **283**:12528–37.
- 116 Dai Y, Chen H, Mo C, Cui L, He W. Ectopically expressed human tumor biomarker MutS homologue 2 is a novel endogenous ligand that is recognized by human $\gamma\delta$ T cells to induce innate anti-tumor/virus immunity. *J Biol Chem* 2012; **287**:16812–9.
- 117 Mo C, Dai Y, Kang N, Cui L, He W. Ectopic expression of human MutS homologue 2 on renal carcinoma cells is induced by oxidative stress with interleukin-18 promotion via p38 mitogen-activated protein kinase (MAPK) and c-Jun N-terminal kinase (JNK) signaling pathways. *J Biol Chem* 2012; **287**:19242–54.
- 118 Lu J, Das M, Kanji S *et al.* Induction of ATM/ATR pathway combined with V γ 2V δ 2 T cells enhances cytotoxicity of ovarian cancer cells. *Biochim Biophys Acta* 2014; **1842**:1071–9.
- 119 Xu B, Pizarro JC, Holmes MA *et al.* Crystal structure of a gammadelta T-cell receptor specific for the human MHC class I homolog MICA. *Proc Natl Acad Sci USA* 2011; **108**:2414–9.
- 120 Bruder J, Siewert K, Obermeier B *et al.* Target specificity of an autoreactive pathogenic human $\gamma\delta$ -T cell receptor in myositis. *J Biol Chem* 2012; **287**:20986–95.
- 121 Aydinoglu MK, Zhang L, Wang C *et al.* $\gamma\delta$ T cells recognize the insulin B:9–23 peptide antigen when it is dimerized through thiol oxidation. *Mol Immunol* 2014; **60**:116–28.
- 122 Paul S, Shilpi, Lal G. Role of gamma-delta ($\gamma\delta$) T cells in autoimmunity. *J Leukoc Biol* 2014; **97**:259–71.