



Risk Factors for, and Effects of, Stress, Anxiety, and Depression in Adolescents

Gareth Richards^{1*} and Andrew P. Smith¹

¹Centre for Occupational & Health Psychology, School of Psychology, Cardiff University, United Kingdom.

Authors' contributions

This work was carried out in collaboration between both authors. Author GR carried out the statistical analysis and data interpretation, and drafted and revised the manuscript. Author APS designed and supervised the running of the project, and revised the manuscript for important intellectual content. Both authors read and approved the final manuscript prior to submission.

Article Information

DOI: 10.9734/BJESBS/2015/19501

Editor(s):

(1) Shao-I Chiu, Taipei College of Maritime Technology of Center for General Education, Taiwan.

Reviewers:

(1) Anonymous, Universite de Rouen, France.

(2) Anonymous, Vytautas Magnus University, Lithuania.

Complete Peer review History: <http://sciencedomain.org/review-history/10156>

Original Research Article

Received 12th June 2015
Accepted 3rd July 2015
Published 14th July 2015

ABSTRACT

Aims: Stress, anxiety, and depression are alarmingly common in the general population, can be extremely debilitating, and are a huge cost to public health services. Understanding risk factors associated with such conditions may therefore be useful in helping devise intervention strategies. Though much research has been conducted in the adult population, relatively few studies have investigated these problems in adolescents from the UK. This is a gap in the literature that the current paper aims to address.

Study Design: The current research utilised a cross-sectional design.

Place and Duration of Study: Data were collected from three secondary schools in Cornwall, UK, in June 2013.

Methodology: Data from the Cornish Academies Project were used to investigate links between demographic and lifestyle variables and single item measures of stress, anxiety, and depression. The sample included 2307 secondary school children aged 11-17 (48.5% male, 51.5% female) from the South West of England.

*Corresponding author: E-mail: RichardsG6@Cardiff.ac.uk;

Results: A number of risk factors were identified, including female gender, low sleep hours, and belonging to a higher school year. In addition to this, the effects of stress anxiety, and depression on school attendance, Key Stage 3/Key Stage 4 English and maths attainment, and the occurrence of behavioural sanctions were investigated. After demographic and lifestyle covariates had been controlled for, high stress was found to predict the occurrence of behavioural sanctions, and high depression was associated with below average English and maths attainment.

Conclusion: Though the findings presented are informative in themselves, the identification of correlates of mental health problems in this demographic group may also be of benefit to future studies that utilise multivariate approaches to data analysis.

Keywords: Adolescent behavior; anxiety; depression; stress.

1. INTRODUCTION

It has been estimated that approximately 450 million people worldwide have a mental health problem [1]. In the context of the UK, The Office for National Statistics [2] reported that 19% of those aged 16 and over attained a General Health Questionnaire score indicative of anxiety or depression. In addition to this, the UK Health and Safety Executive [3] stated that work-related stress, anxiety, and depression cost the UK economy in excess of £530 million per year.

Due to the complexity of relationships between stress, anxiety, and depression, it is considered important for research to take a multidimensional approach [4]. Not only does this allow for interactions between variables to be assessed, and for the influence of confounding factors to be controlled for statistically, it is also more likely to reflect real-world processes. In order to utilise such methods, it is therefore necessary to identify risk factors associated with poor mental health. A number of such variables have been identified in adult populations, such as female gender, low social class, low income and education, unemployment, not being married, and having a poor somatic health status [5-7]. Certain aspects of lifestyle have also been associated with mental health. For instance, aerobic exercise has been found to have antidepressant and anxiolytic effects in adults, and can protect against the harmful consequences of stress [8].

Considering the seriousness of the effects of stress, anxiety, and depression on society as a whole, it is surprising to find that few data have been published, which relate to such phenomena in British adolescents. A population-based sample of British children and adolescents aged 5-15 [9] did, however, identify the prevalence rate of DSM-IV disorders to be 9.5%. Furthermore, a longitudinal assessment of

prevalence rates from age 9 to 16 [10] determined that 36.7% of participants had at least one psychiatric disorder at some point during the study period.

As rates for psychiatric disorders have been shown to vary considerably across different cultures [6], the current paper aims to provide more recent estimates of the prevalence of stress, anxiety, and depression in British adolescents. Cross-sectional data are presented from the Cornish Academies Project, a large-scale longitudinal programme of research. Demographic and lifestyle risk factors of poor mental health were identified, and the effects of mental health on school performance outcomes were examined.

2. MATERIALS AND METHODS

2.1 Participants

The Cornish Academies Project was a longitudinal research programme investigating dietary effects on school performance and mental health in secondary school children from the South West of England. Participants in the current study came from the second cross-section of data, which has been described in detail elsewhere [11]. At this time-point, the cohort consisted of 3323 pupils (2610 of whom had consented to take part), and 2307 completed the questionnaires. An age range of 11-17 ($M = 13.6$, $SD = 1.49$) was observed, and the sex ratio was relatively balanced (48.5% male, 51.5% female).

2.2 Apparatus/Materials

Single item measures from the Wellbeing Process Questionnaire [12] were administered, for which participants were asked to state how frequently they had experienced stress, anxiety, and depression over the previous six months, on

a five-point scale (1 = not at all, 2 = rarely, 3 = sometimes, 4 = frequently, 5 = very frequently). Single items were used because they have been shown to be valid and reliable, can identify overall risk, and save time in comparison to full-length measures [13]. Three items were employed to measure the frequency of taking part in exercise; participants were asked to report how frequently they performed mildly energetic, moderately energetic, and vigorous exercise using a four-point scale (1 = three times a week or more, 2 = once or twice a week, 3 = about once to three times a month, 4 = never/hardly ever). These three items were factor analysed into a single-item solution, the details of which have been reported elsewhere [11]. In addition to this, participants were asked to state how many hours they typically spent asleep each night.

2.3 Design and Procedure

Pupils filled out the questionnaires after being administered them by teachers at their schools, and demographic and school performance data were acquired through the School Information Management System. This included age, sex, school year, school attendance, number of detentions/behavioural sanctions incurred, Key Stage 3/Key Stage 4 English and maths attainment, ethnicity, special educational needs status, eligibility/ineligibility to receive free school meals (used as an indication of socioeconomic status [14]), whether the child was cared for by a non-parental guardian, and whether English was spoken as an additional language. Statistical procedures were carried out using IBM SPSS version 20.

3. RESULTS

The percentage of pupils who reported to have 'frequently' or 'very frequently' experienced stress, anxiety, and depression over the previous six months was alarmingly high at 26%, 14.6%, and 13.2%, respectively (for frequency data, see Table 1). Though it should be acknowledged that one study using single item measurements of these concepts might not be an accurate representation of their true prevalence rates, these findings suggest that a large proportion of mental health problems begin relatively early in life. This is therefore an important area to address, as early intervention may potentially be able to reduce negative effects in both adolescents and adults.

3.1 Demographic and Lifestyle Risk Factors of Poor Mental Health

Data for the single item measures of mental health were recoded into dichotomous variables, with those who answered with 1 or 2 ('not at all' or 'rarely' experienced stress, anxiety, or depression) being placed into the above average group, and those who answered with 3, 4, or 5 ('sometimes', 'frequently', or 'very frequently' experienced stress, anxiety or depression) comprising the below average group. Initial Chi-square tests were performed to investigate associations between these variables and certain aspects of demography and lifestyle. Ethnicity, whether or not English was spoken as an additional language, and whether or not participants were looked after by a non-parental guardian were not investigated because the numbers of participants in the minority groups were particularly small: non-White $N = 84$ (2.8%), English spoken as an additional language $N = 51$ (1.7%), looked after by a non-parental guardian, $N = 17$ (.6%).

It was found that being female was associated with high stress, anxiety, and depression. Though no differences were observed in relation to the schools that the children came from, significant effects were observed regarding their year group. In each case, this reflected significant linear-by-linear trends in which mental health decreased in higher school years: stress, $\chi^2(1, N = 2193) = 28.289, P < .001$; anxiety, $\chi^2(1, N = 2183) = 42.181, P < .001$; depression, $\chi^2(1, N = 2181) = 5.593, P = .02$. Having a SEN status was also associated with high depression, though no effects were observed regarding stress or anxiety. Interestingly, high anxiety was associated with not being eligible to receive free school meals, though no such effects were observed for stress or depression. Sleeping for fewer than average hours per night and exercising infrequently were also associated with high stress, anxiety, and depression. For all Chi-square values and cross-tabulations between mental health and demographic variables, see Table 2; for mental health and lifestyle variables, see Table 3.

To further examine the nature of the effects observed in the univariate analysis, forwards logistic regression was used. The reason for doing this was so that associations between demographic and lifestyle variables and mental health could be examined independently.

Table 1. Frequency information for single item measures of stress, anxiety, and depression

	Not at all	Rarely	Sometimes	Frequently	Very frequently
Stress	216 (9.6%)	606 (26.9%)	842 (37.4%)	393 (17.5%)	192 (8.5%)
Anxiety	430 (19.2%)	856 (38.2%)	625 (27.9%)	225 (10%)	103 (4.6%)
Depression	808 (36.1%)	653 (29.2%)	481 (21.5%)	187 (8.4%)	108 (4.8%)

Note: Modal values are displayed in bold

The variables entered into the models were sex, school attended, school year, presence/absence of a special educational needs status, eligibility/ineligibility to receive free school meals, sleep, and exercise frequency.

High stress remained associated with being female, OR = 1.944, 95% CI [1.601, 2.362], Wald = 44.878, $P < .001$, sleeping for fewer hours than average per night, OR = 1.33, 95% CI [1.082, 1.635], Wald = 7.352, $P = .007$, and being in a higher school year, Wald = 20.863, $P < .001$. More specifically, in regards to this latter finding, compared to Year 7, higher stress levels were reported in Year 9, OR = 1.472, 95% CI [1.085, 1.999], Wald = 6.158, $P = .01$, Year 10, OR = 1.676, 95% CI [1.211, 2.321], Wald = 9.684, $P = .002$, and Year 11, OR = 1.533, 95% CI [1.11, 2.118], Wald = 6.712, $P = .01$, but not in Year 8, OR = .955, 95% CI [.71, 1.285], Wald = .092, $P = .76$.

In a similar manner to the risk factors of stress, high anxiety remained associated with being female, OR = 2.344, 95% CI [1.932, 2.845], Wald = 74.559, $P < .001$, sleeping for fewer hours than average, OR = 1.368, 95% CI [1.117, 1.677], Wald = 9.168, $P = .002$, and being in a higher school year, Wald = 23.474, $P < .001$. As with the effect of school year on stress, compared to Year 7, higher anxiety was reported in Year 9, OR = 1.397, 95% CI [1.024, 1.906], Wald = 4.463, $P = .04$, Year 10, OR = 1.709, 95% CI [1.238, 2.359], Wald = 10.63, $P = .001$, and Year 11, OR = 1.963, 95% CI [1.422, 2.711], Wald = 16.798, $P < .001$, but not in Year 8, OR = 1.122, 95% CI [.821, 1.535], Wald = .522, $P = .47$. Unlike stress, high anxiety also remained significantly associated with not being eligible to receive free school meals, OR = .663, 95% CI [.483, .911], Wald = 6.443, $P = .01$.

In a similar way to stress and anxiety, high depression remained associated with being female, OR = 1.934, 95% CI [1.58, 2.369], Wald = 40.744, $P < .001$, and sleeping for fewer hours than average per night, OR = 1.465, 95% CI [1.202, 1.785], Wald = 14.33, $P < .001$. In addition to this, having a special educational

needs status, OR = 1.302, 95% CI [1.037, 1.635], Wald = 5.156, $P = .02$, and exercising infrequently, OR = 1.343, 95% CI [1.1, 1.639], Wald = 8.411, $P = .004$, also both remained associated with high instances of depression.

3.2 Effects of Poor Mental Health on School Performance

Associations between school performance and mental health were initially examined at the univariate level using Chi-square tests. Low school attendance was found to be associated with high stress, anxiety and depression. Interestingly, high anxiety was also associated with high English attainment, though this effect is likely to be a Type 1 error considering that high maths attainment was related to low stress, depression, and anxiety (though the latter effect was only marginally significant). High stress and high depression were also associated with increased instances of behavioural sanctions (the latter effect again being marginally significant). For all Chi-square values and cross-tabulations between mental health variables and school performance, see Table 4.

In order to investigate the effects of mental health on school performance at the multivariate level, stress, anxiety, and depression (along with each of the demographic and lifestyle variables discussed in the previous section) were entered into forwards logistic regression models upon the dependent variables of school attendance, English attainment, maths attainment, and behavioural sanctions. Though no effects of mental health on school attendance remained significant, high stress was predictive of the occurrence of a high number of behavioural sanctions, OR = 1.388, 95% CI [1.07, 1.799], Wald = 6.102, $P = .01$. High depression was also found to predict both low English attainment, OR = 1.313, 95% CI [1.059, 1.629], Wald = 6.149, $P = .01$, and low maths attainment, OR = 1.287, 95% CI [1.042, 1.59], Wald = 5.481, $P = .02$. The effect on English attainment was particularly interesting considering that no such finding had been made at the univariate level.

Table 2. Chi-square values and cross-tabulations between demographic variables and mental health

		Sex		Academy			School year					SEN status		FSM		
		Male	Female	Academy 1	Academy 2	Academy 3	Year 7	Year 8	Year 9	Year 10	Year 11	Yes	No	Yes	No	
Stress	Low	Count	446	301	226	390	206	192	201	152	126	131	205	604	86	716
		Expected count	359.6	387.4	225.9	406.1	190.1	161.6	167.9	165.3	151.4	155.8	213.1	595.9	88.8	713.2
		Row %	59.7%	40.3%	27.5%	47.4%	25.1%	23.9%	25.1%	19%	15.7%	16.3%	25.3%	74.7%	10.7%	89.3%
		Adjusted residual	7.9	-7.9	.0	-1.4	1.7	3.4	3.6	-1.5	-2.9	-2.8	-.8	.8	-.4	.4
	High	Count	538	759	392	721	314	250	258	300	288	295	379	1029	157	1235
		Expected count	624.4	672.6	392.1	704.9	329.9	280.4	291.1	286.7	262.6	270.2	370.9	1037.1	154.2	1237.8
		Row %	41.5%	58.5%	27.5%	50.5%	22%	18%	18.5%	21.6%	20.7%	21.2%	26.9%	73.1%	11.3%	88.7%
		Adjusted residual	-7.9	7.9	.0	1.4	-1.7	-3.4	-3.6	1.5	2.9	2.8	.8	-.8	.4	-.4
	χ^2	63.064, $P < .001$			3.109, $P = .21$			33.93, $P < .001$.659, $P = .42$.159, $P = .69$	
Anxiety	Low	Count	674	502	348	624	314	291	289	256	213	203	330	933	157	1095
		Expected count	565.2	610.8	352.7	634.7	298.7	252.3	260.4	258.7	236.9	243.7	330.8	932.2	138.7	1113.3
		Row %	57.3%	42.7%	27.1%	48.5%	24.4%	23.2%	23.1%	20.4%	17%	16.2%	26.1%	73.9%	12.5%	87.5%
		Adjusted residual	9.8	-9.8	-.4	-.9	1.6	4.2	3.1	-.3	-2.6	-4.5	-.1	.1	2.5	-2.5
	High	Count	305	556	266	481	206	149	165	195	200	222	248	696	85	847
		Expected count	413.8	447.2	261.3	470.3	221.3	187.7	193.6	192.3	176.1	181.3	247.2	696.8	103.3	828.7
		Row %	35.4%	64.6%	27.9%	50.5%	21.6%	16%	17.7%	20.9%	21.5%	23.8%	26.3%	73.7%	9.1%	90.9%
		Adjusted residual	-9.8	9.8	.4	.9	-1.6	-4.2	-3.1	.3	2.6	4.5	.1	-.1	-2.5	2.5
	χ^2	95.408, $P < .001$			2.415, $P = .3$			42.931, $P < .001$.006, $P = .94$		6.342, $P = .01$	
Depression	Low	Count	723	615	392	715	354	304	318	280	252	271	357	1081	152	1273
		Expected count	644.7	693.3	401	721.7	338.3	287.5	299.9	294.7	269.2	273.8	379.6	1058.4	157.4	1267.6
		Row %	54%	46%	26.8%	48.9%	24.2%	21.3%	22.3%	19.6%	17.7%	19%	24.8%	75.2%	10.7%	89.3%
		Adjusted residual	7.3	-7.3	-.9	-.6	1.7	1.9	2	-1.6	-2	-.3	-2.3	2.3	-.8	.8
	High	Count	257	439	222	390	164	136	141	171	160	148	225	542	89	668
		Expected count	335.3	360.7	213	383.3	179.7	152.5	159.1	156.3	142.8	145.2	202.4	564.6	83.6	673.4
		Row %	36.9%	63.1%	28.6%	50.3%	21.1%	18%	18.7%	22.6%	21.2%	19.6%	29.3%	70.7%	11.8%	88.2%
		Adjusted residual	-7.3	7.3	.9	.6	-1.7	-1.9	-2	1.6	2	.3	2.3	-2.3	.8	-.8
	χ^2	53.688, $P < .001$			2.86, $P = .24$			11.244, $P = .02$					5.234, $P = .02$.598, $P = .44$	

Note: 'SEN' refers to the presence/absence of a special educational needs status; 'FSM' refers to eligibility/ineligibility to receive free school meals

Table 3. Chi-square values and cross-tabulations between lifestyle variables and mental health

		Sleep hours		Exercise frequency		
		High	Low	High	Low	
Stress	Low	Count	446	348	384	390
		Expected count	390.2	403.8	350.4	423.6
		Row %	56.2%	43.8%	49.6%	50.4%
		Adjusted residual	5	-5	3	-3
	High	Count	617	752	582	778
		Expected count	672.8	696.2	615.6	744.4
		Row %	45.1%	54.9%	42.8%	57.2%
		Adjusted residual	-5	5	-3	3
		χ^2	24.783, $p < .001$	9.256, $p = .002$		
Anxiety	Low	Count	673	571	581	631
		Expected count	611.6	632.4	549	663
		Row %	54.1%	45.9%	47.9%	52.1%
		Adjusted residual	5.4	-5.4	2.8	-2.8
	High	Count	387	525	382	532
		Expected count	448.4	463.6	414	500
		Row %	42.4%	57.6%	41.8%	58.2%
		Adjusted residual	-5.4	5.4	-2.8	2.8
		χ^2	28.652, $p < .001$	7.935, $p = .005$		
Depression	Low	Count	753	656	674	707
		Expected count	695	714	627.1	753.9
		Row %	53.4%	46.6%	48.8%	51.2%
		Adjusted residual	5.3	-5.3	4.3	-4.3
	High	Count	309	435	290	452
		Expected count	367	377	336.9	405.1
		Row %	41.5%	58.5%	39.1%	60.9%
		Adjusted residual	-5.3	5.3	-4.3	4.3
		χ^2	27.631, $p < .001$	18.402, $p < .001$		

4. DISCUSSION

The current study aimed to provide an indication of the levels of stress, anxiety, and depression present in secondary school children from the UK. The levels reported were worryingly high, though considering that the sample was relatively homogenous, may not be generalisable to other adolescents from the UK or other countries. In addition to this, a number of covariates of poor stress and mental health were identified, and associations between mental health and school performance were examined.

4.1 Covariates of Mental Health

Female gender was predictive of high stress, anxiety, and depression, even after controlling for other demographic and lifestyle covariates, a finding that broadly reflects those made in older populations [15,16]. Of particular concern was the observation that mental health decreased in higher school years.

Although the effect regarding depression did not retain statistical significance after controlling for covariates, those relating to stress and anxiety did. These results are comparable to those of Ford et al. [9], who found the overall prevalence of DSM-IV disorders to increase with age throughout adolescence, though may be different from those observed in older populations. For instance, a study of Turkish university students [17], found that first and second years reported higher levels of stress, anxiety, and depression compared to third, fourth, and fifth years. Differences in findings such as these therefore highlight the importance of taking demographic variance into account when investigating mental health.

High stress, anxiety, and depression were all associated with low sleep hours. This potentially mirrors observations that insomniacs are more likely to report anxiety and depression [18], and to perceive their lives as being more stressful compared to good sleepers [19].

Table 4. Chi-square values and cross-tabulations between school performance and mental health

			School attendance		KS3/KS4 English		KS3/KS4 Maths		Behavioral sanctions	
			High	Low	High	Low	High	Low	High	Low
Stress	Low	Count	456	346	382	414	409	386	152	654
		Expected count	433.9	368.1	394	402	382	413	170.8	635.2
		Row %	56.9%	43.1%	48%	52%	51.4%	48.6%	18.9%	81.1%
		Adjusted residual	2	-2	-1.1	1.1	2.4	-2.4	-2	2
	High	Count	732	662	696	686	636	744	314	1079
		Expected count	754.1	639.9	684	698	663	717	295.2	1097.8
		Row %	52.5%	47.5%	50.4%	49.6%	46.1%	53.9%	22.5%	77.5%
		Adjusted residual	-2	2	1.1	-1.1	-2.4	2.4	2	-2
χ^2			3.874, $p = .049$		1.137, $p = .29$		5.805, $p = .02$		4.146, $p = .04$	
Anxiety	Low	Count	706	546	595	648	618	624	251	1005
		Expected count	677.5	574.5	618.1	624.9	596.6	645.4	265.1	990.9
		Row %	56.4%	43.6%	47.9%	52.1%	49.8%	50.2%	20%	80%
		Adjusted residual	2.5	-2.5	-2	2	1.9	-1.9	-1.5	1.5
	High	Count	477	457	483	442	422	501	211	722
		Expected count	505.5	428.5	459.9	465.1	443.4	479.6	196.9	736.1
		Row %	51.1%	48.9%	52.2%	47.8%	45.7%	54.3%	22.6%	77.4%
		Adjusted residual	-2.5	2.5	2	-2	-1.9	1.9	1.5	-1.5
χ^2			6.095, $p = .01$		4.011, $p = .045$		3.459, $p = .063$		2.226, $p = .14$	
Depression	Low	Count	800	625	720	697	718	694	282	1146
		Expected count	769.9	655.1	702	715	678.3	733.7	299.7	1128.3
		Row %	56.1%	43.9%	50.8%	49.2%	50.8%	49.2%	19.7%	80.3%
		Adjusted residual	2.7	-2.7	1.6	-1.6	3.6	-3.6	-2	2
	High	Count	380	379	353	396	321	430	177	582
		Expected count	410.1	348.9	371	378	360.7	390.3	159.3	599.7
		Row %	50.1%	49.9%	47.1%	52.9%	42.7%	57.3%	23.3%	76.7%
		Adjusted residual	-2.7	2.7	-1.6	1.6	-3.6	3.6	2	-2
χ^2			7.357, $p = .007$		2.658, $p = .1$		12.908, $p < .001$		3.814, $p = .051$	

Note: 'KS3/KS4' refers to 'Key Stage 3/Key Stage 4'

Though it is difficult to infer the direction of the relationships observed (and they could indeed be bi-directional), these findings suggest that adolescents who sleep for relatively few hours per night may represent a group at increased risk of experiencing mental health problems.

High anxiety was associated with not being eligible to receive free school meals, and the effect remained significant after controlling for additional covariates. This finding is somewhat counterintuitive in that eligibility to receive free school meals is a proxy indication of low socioeconomic status [14], and financial difficulties have been shown to increase depression in British university students [20].

Having a special educational needs status remained associated with high depression, even after additional covariates had been controlled for. Infrequent exercise was also associated with high instances of depression, a finding that may mirror the observation that exercise can have antidepressant effects in adults [8].

4.2 Mental Health and School Performance

Though high stress, anxiety, and depression were all initially associated with low school attendance, none of these effects remained significant once demographic and lifestyle covariates had been controlled for. However, high depression was associated with low English and maths attainment, the former effect having not been detected at the univariate level. Differences between univariate and multivariate findings such as these may highlight the need to control for demographic and lifestyle covariates when investigating the effects of mental health on school performance outcomes. These results are also similar to the finding that depression can predict decreases in exam performance in university students [20]. Furthermore, high stress was found to remain predictive of accruing a high number of behavioural sanctions, an effect similar to the observation that a high number of stressful life events is associated with increased delinquency in adolescents [21].

4.3 Limitations

Though the current study benefitted from a large sample size, the population studied was somewhat homogenous (consisting mainly of White children from an area of relatively low

socioeconomic background). In addition to this, the study is limited in that the data used were only cross-sectional in nature. Though demographic and lifestyle factors were conceptualised as being risk factors of mental health, whereas school performance measures were conceptualised as being outcomes, the potential for reverse-causation and bi-directionality cannot be overlooked. For instance, though being depressed might negatively impact school attainment, achieving poor grades may also be a cause of depression.

5. CONCLUSION

The current study aimed to provide a profile of the levels of stress, anxiety, and depression present in a cohort of British secondary school children. In addition to this, a number of demographic and lifestyle risk factors of poor mental health were identified, which broadly reflect similar findings reported in older populations. Of particular concern are the findings that high stress, anxiety, and depression were relatively common, that the occurrence of poor mental health increased throughout secondary school education, and that it also appeared to negatively affect certain aspects of school performance. These findings therefore suggest strongly that mental health in adolescence should be an area of particular importance for future research.

CONSENT

Written informed consent was acquired from each participant prior to data collection.

ETHICAL APPROVAL

Ethical clearance was granted by Cardiff University's School of Psychology Ethics Committee (ethical clearance number: EC.12.09.11.3187).

ACKNOWLEDGEMENTS

The current research was supported by a grant from The Waterloo Foundation (grant number: 503692). Every person who contributed significantly to the current work has been acknowledged, and neither of the authors have any conflicts of interest to declare. The funders had no involvement in designing of the study, data collection, analysis and interpretation of data, or writing of the manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organization. Mental disorders affect one in four people. 2001; 10:4. Accessed 12 June 2015. Available:http://www.who.int/whr/2001/media_centre/press_release/en/
2. Office for National Statistics. Measuring national well-being - Health. 2013;6:19. Accessed 12 June 2015. Available:http://www.ons.gov.uk/ons/dcp171766_310300.pdf
3. Health and Safety Executive. Workplace stress costs Great Britain in excess of £530 million. 2007;11:7. Accessed 12 June 2015. Available:<http://www.hse.gov.uk/press/2007/c07021.htm>
4. Galvin J, Smith AP. Stress in U.K. mental health training: A multi-dimensional comparison study. *Br J Educ Soc Behav Sci.* 2015;9(3):161-175. DOI:10.9734/BJESBS/2015/18519#sthash.nkcCAdeT.dpuf
5. Jacobi F, Wittchen H-U, Hölling C, Höfler M, Pfister H, Müller N, et al. Prevalence, co-morbidity and correlates of mental disorders in the general population: Results from the German health interview and examination survey (GHS). *Psychol Med.* 1999;34(4):597-611. DOI: 10.1017/s0033291703001399
6. Andrade L, Caraveo-Anduaga JJ, Berglund P, Bijl R, Kessler RC, Demler O, et al. Cross-national comparisons of the prevalence and correlates of mental disorders. *Bull World Health Organ.* 2000; 78(4):413-426.
7. Eisenberg D, Gollust SE, Golberstein E, Hefner JL. Prevalence and correlates of depression, anxiety, and suicidality among university students. *Am J Orthopsychiatry.* 2007;77(4):534-542. DOI: 10.1037/0002-9432.77.4.534
8. Salmon P. Effects of physical exercise on anxiety, depression, and sensitivity to stress: A unifying theory. *Clin Psychol Rev.* 2001;21(1):33-61. DOI: 10.1016/s0272-7358(99)00032-x
9. Ford T, Goodman R, Meltzer H. The British Child and Adolescent Mental Health Survey 1999: The prevalence of DSM-IV disorders. *J Am Acad Child Adolesc Psychiatry.* 2003;42(10):1203-1211. DOI:10.1097/00004583-200310000-00011
10. Costello EJ, Mustillo S, Erkanli A, Keeler G, Angold A. Prevalence and development of psychiatric disorders in childhood and adolescence. *JAMA Psychiatry.* 2003; 60(8):837-844. DOI: 10.1001/archpsyc.60.8.837
11. Richards G, Malthouse A, Smith AP. The diet and behaviour scale (DABS): Testing a new measure of food and drink consumption in a cohort of secondary school children from the south west of England. *J Food Res.* 2015;4(3):148-161. DOI: 10.5539/jfr.v4n3p148
12. Williams G. Researching and developing mental health and well-being assessment tools for supporting employers and employees in Wales [dissertation]. Cardiff University; 2014.
13. Williams G, Smith AP. A holistic approach to stress and well-being. Part 6: The Wellbeing Process Questionnaire (WPQ Short Form). *Occup Health Work.* 2012; 9(1):29-31.
14. Shuttleworth I. The relationship between social deprivation, as measured by individual free school meal eligibility, and educational attainment at GCSE in Northern Ireland: A preliminary investigation. *Br Educ Res J.* 1995;21(4): 487-504. DOI: 10.1080/0141192950210404
15. Mahmoud JSR, Staten R, Hall LA, Lennie TA. The relationship among young adult college students' depression, anxiety, stress, demographics, life satisfaction, and coping styles. *Issues Ment Health Nurs.* 2012;33(3):149-156. DOI: 10.3109/01612840.2011.632708
16. Newbury-Birch D, Kamali F. Psychological stress, anxiety, depression, job satisfaction, and personality characteristics in preregistration house officers. *Postgrad Med J.* 2001;77(904):109-111. DOI: 10.1136/pmj.77.904.109
17. Bayram N, Bilgel N. The prevalence and socio-demographic correlations of depression, anxiety and stress among a group of university students. *Soc Psychiatry Psychiatr Epidemiol.* 2008; 43(8):667-672. DOI: 10.1007/s00127-008-0345-x

18. Taylor DJ, Lichstein KL, Durrence HH, Reidel BW, Bush AJ. Epidemiology of insomnia, depression, and anxiety. *Sleep*. 2005;28(11):1457-1464.
19. Morin CM, Rodrigue S, Ivers H. Role of stress, arousal, and coping skills in primary insomnia. *Psychosom Med*. 2003;65(2): 259-267.
DOI:10.1097/01.psy.0000030391.09558.a3
20. Andrews B, Wilding JM. The relation of depression and anxiety to life-stress and achievement in students. *Br J Psychol*. 2004;95(4):509-521.
DOI: 10.1348/0007126042369802.
21. Hoffmann JP, Cerbone FG. Stressful life events and delinquency escalation in early adolescence. *Criminology*. 1999;37(2): 343-374.
DOI: 10.1111/j.1745-9125.1999.tb00489.x

© 2015 Richards and Smith; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/10156>