

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/92522/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Hickey, Benjamin A., Cleves, Andrew, Alikhan, Raza, Pugh, Neil D. , Nokes, Leonard Derek Martin and Perera, Anthony 2017. The effect of active toe movement (AToM) on calf pump function and deep vein thrombosis in patients with acute foot and ankle trauma treated with cast - a prospective randomized study. *Foot and Ankle Surgery* 23 (3) , pp. 183-188. 10.1016/j.fas.2016.04.007

Publishers page: <http://dx.doi.org/10.1016/j.fas.2016.04.007>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Title Page

The effect of active toe movement (AToM) on calf pump function and deep vein thrombosis in patients with acute foot and ankle trauma treated with cast
– A Prospective Randomised Study

Authors:

Mr Ben A Hickey¹ – corresponding author email: drhickey@hotmail.co.uk

Mr Andrew Cleves¹

Dr Raza Alikhan¹

Professor Neil Pugh¹

Professor Len Nokes²

Mr Anthony Perera¹

Affiliation addresses:

1. University Hospital of Wales, Heath Park, Cardiff, CF14 4XW, Wales, UK
2. Cardiff University, Cardiff, Wales, UK

Corresponding author permanent address:

10 Trafalgar Road, Penylan, Cardiff, CF23 5BQ, Wales, UK

1. Abstract

Background

Patients with foot and ankle trauma treated with cast are advised to perform toe movements to prevent venous thromboembolism (VTE). Our aim was to determine the effect of active toe movement on asymptomatic deep vein thrombosis (DVT) and venous calf pump function.

Methods

Patients aged 18 to 60 years with acute foot and ankle trauma requiring below knee non weight bearing cast were randomized to intervention (regular active toe movement) or control groups (n=100). Patients had bilateral lower limb venous ultrasound to assess for DVT on discharge from clinic. Patients requiring chemical thromboprophylaxis were excluded.

Results

78 completed the study. 27% sustained asymptomatic DVT, with no statistically significant difference in calf pump function or DVT incidence between groups. All DVT's occurred in the injured lower limb.

Conclusion

Active toe movement is not a viable strategy for thromboprophylaxis in patients with acute foot and ankle trauma treated with cast.

Keywords: Thromboprophylaxis, Venous thrombosis, Deep vein thrombosis

2. Introduction

1
2
3
4
5 Thromboprophylaxis is clinically and financially beneficial compared with
6
7 treatment of thromboembolic events once they have occurred [1]. In patients
8
9 with foot and ankle trauma treated with cast, thromboprophylaxis significantly
10
11 reduces venous thromboembolism rate, with thirteen patients requiring
12
13 prophylaxis with low molecular weight heparin to prevent one asymptomatic
14
15 DVT [2-4]. Considering that chemical thromboprophylaxis for patients with
16
17 lower limb cast immobilisation represents 5.3% of the outpatient tariff for
18
19 trauma, this represent a significant cost [5]. This also represents a potential
20
21 risk, in view that chemical thromboprophylaxis can cause major bleeding [6].
22
23 Patients with foot and ankle trauma treated with casts are also advised to
24
25 participate in activities of early mobilisation and calf pumping exercises to
26
27 reduce their risk of developing DVT [7]. Although we have previously shown
28
29 that active toe movement increases venous velocities in the popliteal vein in
30
31 healthy volunteers, even with a below knee cast applied, the clinical
32
33 importance of these findings requires further investigation [8,9]. Most studies
34
35 rule out mechanical thromboprophylaxis in patients with below knee casts for
36
37 various reasons [10, 11]. However, considering that venous stasis is thought
38
39 to contribute to thrombogenesis, it is possible that regular active toe
40
41 movement (AToM) could reduce the development of deep vein thrombosis in
42
43 patients with injury and cast treatment [12]. Our primary aim was to determine
44
45 the effect of regular active toe movement (AToM) on rates of asymptomatic
46
47 DVT in patients with acute foot and ankle trauma treated with cast and define
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 which venous segments were involved. Our secondary aim was to determine
2 the effect of regular active toe movement on calf pump function.
3
4
5

6 7 3. Methods 8 9

10
11 In this prospective randomized controlled study, adult patients with acute foot
12 or ankle trauma treated with below knee cast [REDACTED]
13 [REDACTED] between 18th February 2014 and 21st January 2015 were recruited.
14
15
16

17
18 The local Ethics committee and Research and Development department
19 approved this study. The study was registered on [REDACTED]
20
21

22 [REDACTED] Database [REDACTED]
23
24

25 Funding for consumables [REDACTED]. In
26
27

28 accordance with NICE guidance and local hospital protocols, all patients were
29 assessed for risk of venous thromboembolism. If patients had any permanent
30 risk factors for VTE they were not eligible for the study and were provided with
31 Low molecular weight heparin (Enoxaparin) thromboprophylaxis 40mg
32 subcutaneous injection once daily and were not eligible for study participation.
33
34
35
36
37
38
39
40

41 Patients currently taking Warfarin or Heparin for any other reason were also
42 excluded.
43
44
45

46 47 48 3.1 Exclusion criteria 49 50 51

52
53 Risk factors for VTE (exclusion criteria) were: Age >60 years, Current
54 hormone therapy (Oral Contraceptive Pill, Hormone Replacement Therapy,
55 Tamoxifen), Personal or first degree relative VTE history, Any recent hospital
56
57
58
59
60
61
62
63
64
65

1 admission or major surgery (within 3 months), Pregnancy or immediate
2 postpartum (6 weeks), Any serious medical co-morbidity (including congestive
3 cardiac failure, chronic obstructive pulmonary disease, chronic kidney
4 disease, inflammatory bowel disease), Extensive varicosities, Active cancer,
5 Obesity (BMI over 30 kg/m²), Known thrombophilia, Achilles tendon rupture,
6 Equinus cast. All patients assessed as 'low risk' based on this risk
7 assessment were potentially eligible for the study and were referred to the
8 Chief Investigator for review in the next available fracture clinic. On review in
9 clinic, VTE risk assessment was repeated to confirm patients were 'low risk'
10 and eligible for the study. Patients who were anticipated to be treated in a
11 backslab or cast for less than 1 week were not recruited. Patients with injuries
12 requiring surgery were also not recruited because these patients are provided
13 with Low Molecular Weight Heparin. Patients presenting more than 72 hours
14 after initial injury were excluded.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33 34 35 36 3.2 Study design and intervention 37 38 39 40

41 The Chief Investigator recruited all patients. After consenting for participation
42 in this study, patients were allocated to either the intervention (AToM) or
43 control group according to a pre-defined computerized randomization
44 sequence with variable block sizes. Patients were not informed of their
45 randomization allocation group and all assessments of calf pump function and
46 ultrasound for deep vein thrombosis were performed by medical physicists
47 who were blinded to the randomization allocation. Participants who were
48 allocated to the intervention group were provided with an active toe movement
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 protocol to follow, with a compliance diary to complete. Patients allocated to
2 the intervention group were instructed to perform full active toe flexion and
3 extension movement exercises at the metatarsophalangeal and
4 interphalangeal joints as often as possible, with prescriptive advice to perform
5 these movements at a minimum of 60 movements every 6 hours. Patients
6 were managed in fracture clinic depending on their injury until discharge.
7 Patients were shown how to perform these simple exercises and provided
8 with written advice.
9
10
11
12
13
14
15
16
17
18
19
20
21

22 3.3 Outcome measures 23 24 25

26 When the patient no longer required their below knee cast or boot
27 immobilization as part of their treatment they attended the Medical Physics
28 department for assessment of calf pump function and bilateral above and
29 below knee venous Duplex scanning to screen for asymptomatic deep vein
30 thrombosis. Historically, venography was used to screen for DVT's, however
31 current clinical practice is to use Duplex ultrasonography. This has previously
32 been used in other similar studies [13]. Any patient who was discharged from
33 clinic prior to having Duplex scanning was sent an appointment letter to attend
34 for Duplex scanning. Senior Medical Physicists who work within the National
35 Health Service and perform these scan on patients routinely performed all
36 scans. If a deep vein thrombosis was present on Ultrasound scanning, the
37 anatomical location and laterality in relation to the injured limb was recorded.
38 The patient exited the study at this stage. Patients with positive Doppler scans
39 for DVT were provided with treatment if the DVT was located in the above
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 knee veins (Popliteal or more proximal). Patients with DVT in the below knee
2 veins had a repeat Duplex scan a week later to assess for resolution or
3 propagation of the DVT.
4
5
6
7
8

9 In patients who did not sustain deep vein thrombosis, calf pump function was
10 assessed using Doppler ultrasonography according to a standardized
11 protocol. With the patient seated with hips and knees at 90 degrees flexion,
12 baseline popliteal velocity was measured in the popliteal vein after cast
13 removal, using Doppler ultrasound at rest. Each patient then performed 3
14 consecutive active toe dorsiflexion exercises starting with the toe in neutral
15 position, with a rest of 30 seconds between to allow for venous refilling. Peak
16 popliteal velocity was recorded during active toe movement and the mean
17 was calculated. Peak velocity was then recorded during active toe
18 plantarflexion. This method has previously been used to assess calf pump
19 function in healthy volunteers with and without lower limb cast applied [8].
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 3.4 Statistical analysis 40 41 42

43 A power calculation was based on the prediction that asymptomatic DVT
44 would occur in 21.9% of patients with an anticipated intervention effect size of
45 63% reduction [14]. To allow for 5% withdrawal rate after randomization it was
46 planned to randomize 150 patients to each group to achieve 80% power. To
47 ensure randomization worked, baseline characteristics were analysed using t
48 tests for continuous variables and chi square tests for proportions. Shapiro
49 Wilk test was used to test for normality of calf pump function data and
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 analysis of non parametric results was performed with Mann Whitney U test.
2 A generalization of the Mann Whitney U statistic was also performed to
3 determine the effect size of AToM on calf pump function [15]. All statistical
4 analysis was performed using SPSS for Windows (SPSS, Chicago, Illinois).
5
6
7
8
9
10 Analysis of DVT outcomes between groups was performed using Chi square
11 test.
12
13
14
15

16 4. Results

17
18
19
20
21 During the recruitment, 342 patients between the ages of 18 and 60 years had
22 injuries treated with below knee casts, 242 were excluded prior to
23 randomization. 100 patients entered the study and were randomized to
24 intervention (AToM) or control groups. 78 patients completed the study
25 (Figure 1). There were no significant differences between the groups (Table
26 1). 22 (56%) of the patients randomized to the AToM group returned a fully
27 completed toe movement activity diary with number of active toe movements
28 recorded for each day during cast or boot treatment. 13 (59%) of these
29 patients recorded that they achieved the minimum number of active toe
30 movements (240) per day (range 16-600). 4 (31%) of these patients sustained
31 asymptomatic DVT. There were no reported adverse events.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

4.1 Calf pump function

After 100 patients had been recruited, analysis of the calf pump function between groups was performed to determine whether the intervention had any statistically significant effect on baseline or peak popliteal velocities. Calf pump function data was not normally distributed, and was analysed using Mann-Whitney test. (15,16) The U/nm statistic of 0.572 (95% CI 0.420 – 0.710) for both active toe dorsiflexion and active toe plantarflexion indicated that patients in the intervention group generally had greater peak velocities during these movements, however the effects were small (Table 2). The mean baseline popliteal vein diameter was 7.9mm in both groups. Although baseline popliteal velocity, peak velocity during active toe dorsiflexion and peak velocity during active toe plantarflexion were greater in the AToM group patients, none of these differences were statistically significant (Table 3).

4.2 Asymptomatic DVT

In view that AToM did not statistically significantly influence calf pump function in patients, it was decided by the Chief Investigator with agreement of the steering committee to request an interim analysis of the effect of the intervention on asymptomatic deep vein thrombosis. At this stage in the study, the overall event rate was 33.3% (n=13) in intervention and 20.5% (n=8) in control. Interim analysis found the confidence interval of the proportions ranged from 6.9% asymptomatic DVT reduction in the intervention to 31% increased incidence of asymptomatic DVT in the intervention group [16]. In

1 view that the anticipated effect size was a 63% asymptomatic DVT reduction
2 in the AToM intervention group, it appeared that the intervention was less
3 effective than expected and the steering committee decided to stop
4 recruitment [17].
5
6
7
8
9

10 11 4.3 Incidence and anatomical location of asymptomatic Deep Vein

12 13 Thrombosis 14

15
16
17
18
19 Overall, 21 (27%) patients were found to have asymptomatic deep vein
20 thrombosis on ultrasound examination (Table 4). All of these occurred in the
21 lower limb that had been injured and treated in cast. The DVT rate was 13/39
22 (33.3%) in intervention group, 8/39 (20.5%) in control group. These
23 differences were not statistically significant ($p=0.202$). 3 DVT's were above
24 knee and were treated for oral anticoagulants for 3 months. In all patients
25 except 2, the peroneal vein was thrombosed. In 12 patients the DVT was an
26 isolated peroneal DVT. 4 patients had combined peroneal and posterior tibial
27 vein DVT, 1 patient had isolated posterior tibial vein DVT, 2 had popliteal vein
28 DVT which appeared to arise from the peroneal vein, 1 patient had isolated
29 popliteal vein DVT. The mean age of patients who sustained DVT was 36
30 years (range 21 to 53) with a mean BMI of 25kgm^2 (range 19.7 to 30.4). 13
31 were male (65%). There was no statistically significant difference between
32 age or BMI between patients who sustained DVT and those who did not.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

5. Discussion

1
2
3
4
5 This is the first study to examine the effects of active toe movement on DVT in
6
7 patients with foot and ankle trauma treated with cast. Although we have
8
9 previously shown that active toe movement (AToM) can influence calf pump
10
11 function in healthy volunteers with below knee casts applied, we were unable
12
13 to find any evidence that this reduces the incidence of DVT [8]. We
14
15 anticipated that patients who performed regular active toe movements would
16
17 have had improved venous calf pump function parameters, however this
18
19 intervention had no significant effect.
20
21

22
23
24 The concept that lower limb muscular contraction can influence thrombosis is
25
26 not new. In a prospective randomized study performed almost 20 years ago,
27
28 38 patients who underwent total hip replacement were randomized to a
29
30 standardized foot and ankle movement protocol and venous plethysmography
31
32 was performed 4 days postoperatively [18]. Venous outflow was increased by
33
34 22% in the intervention group and it was recommended that regular foot and
35
36 ankle exercises should be part of a prophylactic regimen against venous
37
38 thrombosis [18]. Although these exercises may have influenced calf pump
39
40 function, there was no further study conducted to investigate the effects on
41
42 deep vein thrombosis. Similarly, Kaplan et al (2002) found that electrical
43
44 stimulation of the foot or calf muscles increased popliteal and femoral vein
45
46 velocities, but failed to investigate the effects on DVT [19]. More recently,
47
48 Fuchs et al (2005) conducted a prospective randomized controlled trial of 227
49
50 trauma patients with bony or ligamentous trauma to the spine, pelvis including
51
52 acetabulum, femur, tibia or ankle [20]. All patients received unfractionated
53
54
55
56
57
58
59
60
61
62
63
64
65

1 heparin, however the intervention group also received mechanical
2 thromboprophylaxis with an ankle continuous passive motion machine
3
4 (Arthroflow) for 30-minute sessions, three times per day (30 movements per
5
6 minute, range of motion 20 degree dorsiflexion to 40 degrees plantarflexion).
7
8 Asymptomatic DVT diagnosed on ultrasound or venogram was significantly
9
10 reduced in the Arthroflow group, from 25% to 3.6% ($p=0.001$). Whitelaw et al
11
12 (2001) compared the haemodynamic effects of 6 different mechanical
13
14 thromboprophylactic devices on calf pump function and found that there was
15
16 no statistically significant difference in increases in peak velocity between the
17
18 most effective device (AirCast VenaFlow) and active or passive dorsiflexion
19
20 [21]. An early prospective randomized controlled trial of 290 patients who
21
22 underwent total hip replacement, confirmed that not only did AV foot pumps
23
24 influence popliteal venous velocities, but also reduced asymptomatic DVT to
25
26 rates comparable to levels achieved with low molecular weight heparin [22].
27
28 Since then, intermittent pneumatic compression has been found to reduce
29
30 symptomatic DVT (from 16.7% in control group, to 7.3% in IPC) and
31
32 pulmonary embolism (from 2.8% to 1.2%) at meta-analysis level of 70 studies
33
34 including over 16000 medial and surgical patients [23]. It is also effective in
35
36 reducing VTE in patients with hip fracture, patients undergoing total knee
37
38 replacement and following stroke, all of which are considered high risk VTE
39
40 groups [24, 25, 26].
41
42
43
44
45
46
47
48
49
50
51
52

53 Considering that active toe movement influences venous haemodynamics in
54
55 healthy volunteers to a similar extent to intermittent pneumatic compression
56
57 foot pump devices, we anticipated that this may also reduce asymptomatic
58
59
60
61
62
63
64
65

1 DVT in patients with casts [8, 21]. One of the main reasons why AToM failed
2 to reduce DVT rates may relate to the relative infrequent performance of
3
4 exercises by patients. In our AToM study, participants in intervention group
5 performed 60 sets of movements 4 times per day, which equates to 240 sets
6
7 in a 24 hour period. In comparison to patients using A-V foot pumps in the
8
9 study by Warwick et al (1998) used them for a median of 15 hours in a 24
10
11 hour period at a 20 second cycle, equating to approximately 2700
12
13 compressions [22]. This is more than 10 times the number of compressions
14
15 participants in the AToM study would have performed. To achieve this with
16
17 AToM, assuming patients sleep for approximately 8 hours and would be
18
19 awake for 16 hours per day, participants would need to perform almost 170
20
21 sets of AToM per hour to achieve a similar amount of compression.
22
23

24
25
26
27
28 Considering that many patients in our study were unable to achieve even the
29
30 minimum suggested number of active toe movements (240 in 24 hours) and
31
32 that none of them reported performing more than 600 active toe movement
33
34 exercises in a 24 hour period it is unlikely that a patient would be able activate
35
36 their calf pump as frequently as an AV foot pump. Therefore, although the
37
38 effect of active toe movement on popliteal velocities is similar to the effect of
39
40 AV foot compression devices, there is a great difference in the number of calf
41
42 pump activations performed [21].
43
44
45
46
47

48 Considering that all patients were risk assessed for VTE and all were
49 considered low risk, we were surprised to find that 27% of all patients were
50 found to have an asymptomatic DVT on lower limb ultrasound examination. It
51 appears therefore that current risk assessment tools fail to identify a
52 significant proportion of patients who will develop asymptomatic venous
53
54
55
56
57
58
59
60
61
62
63
64
65

1 thrombosis. An important finding is that all of these venous thrombosis
2 occurred in the lower limb that had been injured and treated in cast.
3
4 Furthermore, the majority of these events occurred in the peroneal veins
5 (90%). In view of this, interventions to reduce venous stasis in these patients
6 may be more effective if they aim to reduce stasis in the peroneal veins, by
7 stimulation of the musculature in the peroneal compartment of the injured,
8 casted leg. Our intervention focused on active flexion and extension exercises
9 of the great toe, thereby using flexor and extensor hallucis longus to
10 influence calf pump mechanics. These muscles are located in the deep
11 posterior and anterior compartments of the leg, which may be a reason why
12 these exercises did not influence the incidence of peroneal vein thrombosis.
13
14 Other potential thromboprophylactic exercises could have included knee
15 flexion and extension to utilize contraction of gastrocnemius, which attaches
16 proximal to the knee joint. However, these muscles are located in the
17 superficial posterior compartment of the leg and may also fail to influence
18 stasis in the peroneal veins. This requires further study. To minimize the
19 potential for venous stasis, minimizing the time duration of rigid cast treatment
20 and using the least immobilizing cast/boot the injury will allow, may also
21 influence venous thrombosis rates.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 In view that the patient demographics in each study group were similar, it
49 appears that the randomization sequence used in our study was effective.
50
51 This is important, because unknown confounding factors are likely to have
52
53 also been similar between groups [27]. We achieved almost 80% follow up,
54
55
56
57
58 which is higher than many previous randomized controlled studies of
59
60
61
62
63
64
65

1 thromboprophylaxis in a similar patient group [28, 29, 30]. It has previously
2 been acknowledged that co-ordination of studies to reduce the risk of VTE is
3
4 challenging due the fact that patients with lower limb casts present and are
5
6 treated in several areas within the hospital e.g. Emergency department,
7
8 fracture clinic [31]. Although our intervention was relatively simple, this means
9
10 our findings have good external validity. It has previously been suggested that
11
12 prospective randomized controlled trials are at risk of lacking external validity
13
14 due to highly intensive or regimented intervention protocols [32]. Although
15
16 only 33% of patients in the intervention group reported achieving the minimum
17
18 recommended number of active toe movements, this is an effectiveness study
19
20 which assesses how the intervention works in the real world [27]. Although
21
22 multiple practitioners performed assessment for DVT, all had a minimum of 5
23
24 years experience in the NHS performing these scan as part of their clinical
25
26 practice. None of the ultrasound scans for DVT were likely to be false
27
28 positives in view that all were still positive on repeat scanning after a week
29
30 interval. All assessors of DVT were blinded to the patient randomization
31
32 group.
33
34
35
36
37
38
39
40
41
42

43 We acknowledge several limitations of our study. Firstly, we were only able to
44
45 recruit 100 (26%) patients after screening 380 for eligibility, due to the current
46
47 NICE guidelines that recommend chemical thromboprophylaxis for patients
48
49 with lower limb trauma treated with cast with additional risk factors. This
50
51 meant that none of the patients who underwent surgery could be included (n=
52
53 105), with an additional 72 patients being prescribed LMWH for other reasons.
54
55
56 We were also unable to recruit patients over the age of 60 years of age
57
58
59
60
61
62
63
64
65

(n=97) during the study period because this is also considered to increase VTE risk in patients with lower limb trauma treated with cast [6].

In conclusion, regular active toe movement by patients treated with lower limb cast for acute foot and ankle trauma did not significantly influence calf pump function or reduce the incidence of asymptomatic deep vein thrombosis. All cases of deep vein thrombosis occurred in the lower limb, which had been injured and treated in cast. The majority (90%) of venous thrombosis involved the peroneal veins. Advice to patients to perform regular active toe movement alone appears insufficient to reduce the risk of development of asymptomatic deep vein thrombosis.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References:

- [1]. Alikhan R, Cohen AT, Combe S, Samama MM, Desjardins L, Eldor A, et al. Risk factors for venous thromboembolism in hospitalized patients with acute medical illness: analysis of the MEDENOX Study. *Arch Intern Med. American Medical Association*; 2004 May 10;164(9):963–8.
- [2]. Roberts C. Towards evidence-based emergency medicine: best BETs from the Manchester Royal Infirmary. BET 4: thromboprophylaxis reduces venous thromboembolism rate in ambulatory patients immobilised in above-knee plaster cast. *Emergency Medicine Journal*. 2012 May;29(5):425–6.
- [3]. Testroote M, Stigter WAH, Janssen L, Janzing HMJ. Low molecular weight heparin for prevention of venous thromboembolism in patients with lower-leg immobilization. Testroote M, editor. *Cochrane Database Syst Rev*. Chichester, UK: John Wiley & Sons, Ltd; 2014;4:CD006681.
- [4]. Ettema HB, KOLLEN BJ, VERHEYEN CCPM, BÜLLER HR. Prevention of venous thromboembolism in patients with immobilization of the lower extremities: a meta-analysis of randomized controlled trials. *Journal of Thrombosis and Haemostasis*. 2008 Jul;6(7):1093–8.
- [5]. Menakaya CU, Pennington N, Muthukumar N, Joel J, Ramirez Jimenez AJ, Shaw CJ, et al. The cost of outpatient venous thromboembolism prophylaxis following lower limb injuries. *Bone Joint J. British Editorial Society of Bone and Joint Surgery*; 2013 May;95-B(5):673–7.
- [6]. National Clinical Guideline Centre – Acute and Chronic Conditions (UK). *Venous Thromboembolism: Reducing the Risk of Venous Thromboembolism (Deep Vein Thrombosis and Pulmonary Embolism) in Patients Admitted to Hospital*. London: Royal College of Physicians (UK); 2010.
- [7]. Le Sage S, Marianne McGee BSc MAC, Jessica D Emed RN BNMA. Knowledge of Venous Thromboembolism (VTE) prevention among hospitalized patients. *Journal of Vascular Nursing. Society for Vascular Nursing, Inc*; 2014 May 1;:1–9.
- [8]. Hickey BA, Morgan A, Pugh N, Perera A. The effect of lower limb cast immobilization on calf muscle pump function: a simple strategy of exercises can maintain flow. *Foot Ankle Int. SAGE Publications*; 2014 May;35(5):429–33.
- [9]. Parsons NR, Hiskens R, Price CL, Achten J, Costa ML. A systematic survey of the quality of research reporting in general orthopaedic journals. *J Bone Joint Surg Br. British Editorial Society of Bone and Joint Surgery*; 2011 Sep;93(9):1154–9.
- [10]. Parsonage I. Venous thromboembolism in patients with below-knee plaster casts. *Emerg Nurse*. 2009 Mar;16(10):32–5.
- [11]. Decramer A, Lowyck H, Demuyck M. Parameters influencing thromboprophylaxis management of a lower leg trauma treated with a cast/splint. *Acta Orthop Belg*. 2008 Oct;74(5):672–7.
- [12]. Lowe GDO. Virchow's triad revisited: abnormal flow. *Pathophysiol Haemost Thromb*. 2003 Dec;33(5-6):455–7.
- [13]. Selby R, Geerts WH, Kreder HJ, Crowther MA, Kaus L, Sealey F. A double-blind, randomized controlled trial of the prevention of clinically important venous thromboembolism after isolated lower leg fractures. *J Orthop Trauma*. 2015 May;29(5):224–30.
- [14]. Roderick P, Ferris G, Wilson K, Halls H, Jackson D, Collins R, et al. Towards evidence-based guidelines for the prevention of venous thromboembolism: systematic reviews of mechanical methods, oral anticoagulation, dextran and regional anaesthesia as thromboprophylaxis. *Health Technol Assess*. 2005 Dec;9(49):iii–iv–ix–x–1–78.
- [15]. Newcombe RG. Confidence intervals for an effect size measure based on the Mann-Whitney statistic. Part 1: general issues and tail-area-based methods. *Stat Med. John Wiley & Sons, Ltd*; 2006 Feb 28;25(4):543–57.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- [16]. Newcombe RG. Interval estimation for the difference between independent proportions: comparison of eleven methods. *Stat Med.* 1998 Apr 30;17(8):873–90.
 - [17]. Briel M, Bassler D, Wang AT, Guyatt GH, Montori VM. The dangers of stopping a trial too early. *J Bone Joint Surg Am.* 2012 Jul 18;94 Suppl 1:56–60.
 - [18]. McNally MA, Cooke EA, Mollan RA. The effect of active movement of the foot on venous blood flow after total hip replacement. *J Bone Joint Surg Am.* 1997 Aug;79(8):1198–201.
 - [19]. Kaplan RE, Czorny JJ, Fung TS, Unsworth JD, Hirsh J. Electrical foot stimulation and implications for the prevention of venous thromboembolic disease. *Thromb Haemost.* 2002 Aug;88(2):200–4.
 - [20]. Fuchs S, Heyse T, Rudofsky G, Gosheger G, Chylarecki C. Continuous passive motion in the prevention of deep-vein thrombosis: a randomised comparison in trauma patients. *J Bone Joint Surg Br. British Editorial Society of Bone and Joint Surgery;* 2005 Aug;87(8):1117–22.
 - [21]. Whitelaw GP, Oladipo OJ, Shah BP, DeMuth KA, Coffman J, Segal D. Evaluation of intermittent pneumatic compression devices. *Orthopedics.* 2001 Mar;24(3):257–61.
 - [22]. Warwick D, Harrison J, Glew D, Mitchelmore A, Peters TJ, Donovan J. Comparison of the use of a foot pump with the use of low-molecular-weight heparin for the prevention of deep-vein thrombosis after total hip replacement. A prospective, randomized trial. *J Bone Joint Surg Am.* 1998 Aug;80(8):1158–66.
 - [23]. Ho KM, Tan JA. Stratified meta-analysis of intermittent pneumatic compression of the lower limbs to prevent venous thromboembolism in hospitalized patients. *Circulation.* 2013 Aug 27;128(9):1003–20.
 - [24]. Handoll HH, Farrar MJ, McBirnie J, Tytherleigh-Strong G, Awal KA, Milne AA, et al. Heparin, low molecular weight heparin and physical methods for preventing deep vein thrombosis and pulmonary embolism following surgery for hip fractures. *Cochrane Database Syst Rev.* 2000;(2):CD000305.
 - [25]. Kakkos SK, Warwick D, Nicolaides AN, Stansby GP, Tsolakis IA. Combined (mechanical and pharmacological) modalities for the prevention of venous thromboembolism in joint replacement surgery. *J Bone Joint Surg Br.* 2012 Jun;94(6):729–34.
 - [26]. Dennis M, Sandercock P, Murray G, Forbes J. Intermittent pneumatic compression in patients with stroke--authors' reply. *Lancet.* 2013 Nov 2;382(9903):1481–2.
 - [27]. Bederman SS, Chundamala J, Wright JG. Randomized clinical trials in orthopaedic surgery: strategies to improve quantity and quality. *J Am Acad Orthop Surg.* 2010 Aug;18(8):454–63.
 - [28]. Wille-Jørgensen P, Jørgensen LN, Crawford M. Asymptomatic postoperative deep vein thrombosis and the development of postthrombotic syndrome. A systematic review and meta-analysis. *Thromb Haemost.* 2005 Feb;93(2):236–41.
 - [29]. Spannagel U, Kujath P. Low molecular weight heparin for the prevention of thromboembolism in outpatients immobilized by plaster cast. *Semin Thromb Hemost.* 1993;19 Suppl 1:131–41.
 - [30]. Lapidus LJ, Ponzer S, Elvin A, Levander C, Lårfars G, Rosfors S, et al. Prolonged thromboprophylaxis with Dalteparin during immobilization after ankle fracture surgery: a randomized placebo-controlled, double-blind study. *Acta Orthop.* 2007 Aug;78(4):528–35.
 - [31]. Nokes TJC, Keenan J. Thromboprophylaxis in patients with lower limb immobilisation - review of current status. *British Journal of Haematology.* 2009 Aug;146(4):361–8.
 - [32]. Castillo RC, Scharfstein DO, MacKenzie EJ. Observational studies in the era of randomized trials: finding the balance. *J Bone Joint Surg Am.* 2012 Jul 18;94 Suppl 1:112–7.

Table 1: Characteristics of participants

Characteristics	AToM (N = 39)	Control (N = 39)
Age (mean)	36.6 (18-60)	36.8 (18-60)
Male sex, n (%)	23 (59)	28 (72)
Bony injury, n (%)	36 (92)	36 (92)
Injuries, n		
Ankle fracture –Weber A	7	8
Ankle fracture – Weber B	15	16
Ankle sprain	3	3
Anterior process of calcaneus fracture	1	0
Cuboid fracture	1	1
Distal fibula avulsion fracture	1	0
Dorsal talonavicular ligament avulsion	3	1
Fifth metatarsal fracture	1	2
Lateral process of talus fracture	1	1
Lisfranc injury	0	1
Medial malleolus fracture	4	5
Navicular fracture	1	1
Posterior malleolus fracture	1	0
Duration of cast or boot, mean +/- SD (days)	42.7 (13.8)	39.6 (17.0)

Table 2: Mann Whitney U statistic analysis of calf pump function parameters

(U/mn)

	U/mn	95% confidence limits	
		Lower	Upper
Baseline popliteal vein diameter - injured/cast leg	0.495	0.349	0.642
Baseline popliteal vein velocity - injured/cast leg	0.533	0.384	0.676
Peak velocity during active toe dorsiflexion (average) - injured/cast leg	0.572	0.420	0.710
Peak velocity during active toe plantarflexion (average) - injured/casted	0.572	0.420	0.710

Table 3: Calf pump function results


Calf pump function parameter	AToM	Control	p-value
Baseline popliteal vein diameter (mm)	7.9 (1.736)	7.95 (SD 2.084)	0.933
Baseline popliteal vein velocity (cm/s)	7.33 (3.663)	6.8 (2.193)	0.579
Peak popliteal velocity during active toe dorsiflexion (cm/s)	44.04 (30.172)	34.20 (18.878)	0.361
Peak popliteal velocity during active toe plantarflexion (cm/s)	39.04 (23.699)	31.70 (16.398)	0.348

Table 4: Asymptomatic Deep Vein Thrombosis events

	AToM (N=39)	Control (N=39)
Asymptomatic DVT in Popliteal vein or more proximal	1	2
Asymptomatic below knee DVT details:		
Peroneal vein	8	5
Posterior tibial vein	1	0
Peroneal and posterior tibial veins	3	1
Total n (%)	13 (33.3)	8 (20.5)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1: Flow diagram of study participants



The effect of active toe movement (AToM) on calf pump function and deep vein thrombosis in patients with acute foot and ankle trauma treated with cast
– A Prospective Randomised Study

Highlights:

There is no current evidence that active toe movement reduces asymptomatic venous thromboembolism in patients with foot and ankle trauma treated with below knee cast

27% (n=21) of patients with acute foot and ankle trauma treated with below knee cast will develop asymptomatic deep vein thrombosis despite being assessed as low risk for thrombosis (above knee DVT incidence of 4% (n=3))

90% (n=19) of asymptomatic deep vein thrombosis events in this patient group involve the peroneal vein

All cases of asymptomatic deep vein thrombosis occur in the lower limb that is treated with below knee cast

Conflict of Interest Statement

The effect of active toe movement (AToM) on calf pump function and deep vein thrombosis in patients with acute foot and ankle trauma treated with cast
– A Prospective Randomised Study

Conflicts of interest statement:

None