

1 **Changes in temperature and precipitation extremes over the Greater Horn of Africa**  
2 **region from 1961 to 2010**

3

41. Philip Aming'o Omondi<sup>1</sup>,  
52. Joseph L. Awange<sup>2</sup>,  
63. Ehsan Forootan<sup>3\*</sup>,  
74. Laban Ayieko Ogallo<sup>1</sup>,  
85. Ruben Barakiza<sup>4</sup>,  
96. Gezahegn Bogale Girmaw<sup>5</sup>,  
107. Isaac Fesseha<sup>6</sup>,  
118. Venerabilis Kululetera<sup>7</sup>,  
129. Caroline Kilembe<sup>8</sup>,  
13 Mathieu Mugunga Mbat<sup>9</sup>,  
14 Mary Kilavi<sup>10</sup>,  
15 Stephen Mutua King'uyu<sup>10</sup>,  
16 Peter Adek Omeny<sup>10</sup>,  
17 Andrew Njogu<sup>10</sup>,  
18 Eldin Mamoun Badr<sup>11</sup>,  
19 Tibin Adam Musa<sup>12</sup>,  
20 Peris Muchiri<sup>13</sup>,  
21 Deus Bamanya<sup>14</sup>and  
22 Everline Komutunga<sup>15</sup>

23

24 **How to Cite:**

25 Omondi, P. A., Awange, J. L., Forootan, E., Ogallo, L. A., Barakiza, R., Girmaw, G. B., Fesseha,  
26 I., Kululetera, V., Kilembe, C., Mbat, M. M., Kilavi, M., King'uyu, S. M., Omeny, P. A., Njogu, A.,  
27 Badr, E. M., Musa, T. A., Muchiri, P., Bamanya, D. and Komutunga, E. (2014), Changes in  
28 temperature and precipitation extremes over the Greater Horn of Africa region from 1961 to  
29 2010. *Int. J. Climatol.*, 34: 1262–1277. doi: 10.1002/joc.3763

30 **URL:** <http://onlinelibrary.wiley.com/doi/10.1002/joc.3763/abstract>

31 **This is a report of the above cited paper. Please read the original paper from the URL link!**

32

# Changes in temperature and precipitation extremes over the Greater Horn of Africa region from 1961 to 2010

---

**ABSTRACT:** Recent special reports on climate extremes have shown evidences of changes in the patterns of climate extremes at global, regional and local scales. Understanding the characteristics of climate extremes at regional and local levels is critical not only for the development of preparedness and early warning systems, but is also fundamental in the development of any adaptation strategies. There is still very limited knowledge regarding the past, present and future patterns of climate extremes in the Greater Horn of Africa (GHA). This study, which was supported by the World Bank Global Facility for Disaster Reduction and Recovery (WB-GFDRR) and implemented by the World Meteorological Organization, was organized in terms of three workshops with three main objectives; (i) analysis of daily rainfall and temperature extremes for ten countries in the GHA region using observed in-situ data running from 1971 to 2006, (ii) assessing whether the United Kingdom Met-office and Hadley centre Providing REgional Climates for Impact Studies (UK-PRECIS) modeling system can provide realistic representation of the past and present climate extremes as observed by available in-situ data, and (iii) studying the future regional climate extremes under different scenarios to further assess the expected changes in climate extremes. This paper, therefore, uses the outputs of these workshops and also includes post-workshop analyses to assess the changes of climate extremes within the GHA. The results showed a significant decrease in total precipitation in wet days greater than 1 mm and increasing warm extremes, particularly at night, while cold extremes are decreasing. Considering a combination of geophysical models and satellite gravimetry observations from the Gravity Recovery And Climate Experiment (GRACE) mission in the frame of GRACE daily Kalman-smoothing models, for the years 2002 to 2010, we explored a decline in total water storage variations over the GHA.

---

**KEY WORDS:** climate extremes; Greater Horn of Africa; climate indices; water storage changes; PRECIS model

## 1 **1. Introduction**

2 Extreme events not only cause property damage, injury, hunger, loss of life and threaten the  
3 existence of some species (see, e.g., Downing 1991), but also drive changes in natural and  
4 human systems much more than average climate (Parmesan et al., 2000; Peterson et al., 2008,  
5 Tierney et al., 2013). Impacts of extreme climate change and variability lead to human suffering,  
6 particularly for the poor as witnessed, e.g., during the prevalent droughts and floods in the GHA  
7 region (Lyon and DeWitt, 2012). Analysis of changes in extreme climate events, therefore, is  
8 important due to the potentially high social, economic, and ecological impact of such events  
9 (e.g., Bohle et al., 1994; Arnell, 2004). Limited availability of long records of daily climate data  
10 in some parts of the world, including the GHA, hampers efforts to analyze the impacts of climate  
11 change and variability on the frequency and severity of climate extremes (Folland et al., 2001).  
12 Long records of daily climate data for the GHA region are discussed in Camberlin and Philippon  
13 (2002); WMO (2003); Brant et al. (2012) and Omondi (2011, 2012). Availability of long term  
14 high quality data in the region is hampered by inadequate monitoring networks; gaps in the  
15 records; a general decline of number of stations; chronic under-funding; differences in  
16 processing and quality control; and differences in data policies (WMO, 2003). It is evident that  
17 the GHA countries share pronounced climatic trends (Tierney et al., 2013) and variability and are  
18 vulnerable to extreme climatic conditions (e.g., Downing, 1991; Schreck and Semazzi, 2004;  
19 Anyah and Qiu, 2012 and Omondi et al., 2012).

20 It is increasingly becoming apparent that behind the ongoing research and debate on climate  
21 change, many parts of Africa are already witnessing dire consequences of erratic climatic  
22 conditions (Anyah and Qui, 2011; Shongwe et al., 2011) that are likely associated with regional  
23 climatic changes (Funk et al., 2008, 2012). This is expected to pose unprecedented challenges to  
24 most African economies that are significantly hinged on a predominantly rain-fed agriculture.  
25 Further challenges lie in understanding low frequency multi-decadal and centennial climate  
26 variability in the vastness and uniqueness of the complex African terrain and climate systems  
27 over eastern Africa (Omondi et al., 2012; Tierney et al., 2013).

28 In recent years over the GHA region, particularly in Kenya, Ethiopia and Somali, climate related  
29 extremes have been the dominant trigger of natural disasters. The region has recently witnessed  
30 frequent episodes of both excessive (e.g. Anyah and Semazzi, 2006; Hastenrath et al., 2010;  
31 Kijazi and Reason, 2009a; Viste et al., 2012) and deficient rainfall (e.g. Hastenrath et al., 2007,  
32 2010; Kijazi and Reason, 2009b; Omondi et al., 2012). Consequently, there has been an upsurge  
33 in interest both from the scientific communities (e.g., van Oldenborgh et al., 2005; Omumbo,  
34 2011) and policy makers to the risk of increased extreme climatic events. A recent climate  
35 analysis for East Africa was conducted by Christy and co-authors examining air temperature  
36 trends at 60 stations across Kenya (Christy et al., 2009). After spatially interpolating the station-  
37 based data, the study reports finding a statistically significant upward trend in minimum  
38 temperature ( $T_n$ ) in the Kenyan Highlands region. Omumbo et al. (2011) found evidence of a  
39 warming trend in observed maximum, minimum and mean temperatures at Kericho during the  
40 period 1979 to 2009 using gold standard meteorological observations. An upward trend of  
41  $\approx 0.2^\circ\text{C}/\text{decade}$  was observed in all three temperature variables ( $P < 0.01$ ). Mean temperature  
42 variations in Kericho were associated with large-scale climate variations including tropical SST  
43 ( $r = 0.50$ ;  $p < 0.01$ ). Other authors have also studied malaria resurgence using extreme climatic  
44 events over the region (e.g., Hay et al., 2002; Pascal et al., 2006 among others). Analysis of  
45 rainfall, minimum and maximum temperatures records showed increasing trends in annual  
46 minimum and maximum temperatures from 1951 to 2002 ( $0.4^\circ\text{C}/\text{decade}$  and  $0.2^\circ\text{C}/\text{decade}$ ,

1 respectively) but little trend in rainfall from 1901–50, 1951–2002 and 1901–2002 (Conway et al.,  
2 2004). In analysis of rainfall seasonality index (SI), precipitation concentration index (PCI) and  
3 modified Fournier index (MFI), Elagib (2010) found no statistically significant trends in  
4 observed rainfall for the hyper-arid region of Sudan during the common data period of 1945–  
5 2007.

6 Some recent studies using Global Climate Models (GCMs) have shown that changes in climate  
7 over the region are expected in a global warming scenario (IPCC, 2007; Shongwe, 2010; Anyah  
8 and Qiu, 2012). These are likely to include changes in the intensity, duration, and frequency of  
9 droughts and floods, heat waves, etc, and will have serious implications on agriculture, human  
10 health, as well as human activities. It is well documented that some parts of the GHA region is  
11 perennially prone to droughts and floods (Hastenrath et al., 2007; Kijazi and Reason, 2009a, b;  
12 Omondi, 2011; Bradfield and DeWitt, 2012).

13 Although there is a lack of long records of daily data for extreme climate change detection,  
14 international collaboration is significantly improving the situation, culminating in an analysis  
15 (see, e.g., Alexander et al., 2006) that provided a near-global perspective on changing climate  
16 extremes for the Intergovernmental Panel on Climate Change (IPCC) Fourth Assessment Report  
17 (IPCC, 2007). Yet quantifiable information, describing how weather and climate extremes are  
18 changing over the GHA region has been unavailable. In preparation for the Intergovernmental  
19 Panel on Climate Change (IPCC) Fifth Assessment Report, a major effort by the Expert Team on  
20 Climate Change Detection and Indices (ETCCDI) has been undertaken to analyze how extremes  
21 are changing over as much of the world as possible (Alexander et al., 2006). This included  
22 intensive international collaboration on data exchange and analysis, and, where data were not  
23 available, holding regional climate change workshops to generate information on extremes  
24 (Alexander et al., 2006). This global assessment initiative has greatly benefited from the  
25 contributions from a series of workshops (Peterson and Manton, 2008) coordinated by ETCCDI  
26 which is jointly sponsored by the WMO Commission for Climatology (CCI), the Joint  
27 Commission for Oceanography and Marine Meteorology (JCOMM), as well as the Research  
28 Programme on Climate Variability and Predictability (CLIVAR). The ETCCDI workshops seek  
29 to bring participants together from countries within a data sparse region to fill in data gaps and to  
30 provide capacity building.

31 The availability of daily observation data is steadily improving and has led to the development of  
32 gridded regional (Haylock et al., 2008) and global datasets (Caesar et al., 2006). For some areas  
33 such as the GHA region where daily data availability is still relatively poor, assembling and  
34 research using this data would be useful in decision making and policy formulation on the  
35 climate sensitive sectors of the region.

36 To address the issue of data shortage in GHA, therefore, and in line with global standard practice  
37 (e.g., Alexander et al., 2006), this study is intended to help fill the data gap in the region by  
38 assembling the necessary climate observations. Further, the understanding and use of regional  
39 climate downscaling tool known as PRECIS would help GHA countries design adaptation  
40 policies and reduce climate associated risks.

41 This study is divided into four parts namely (i) the assessment of the adequacy of regional  
42 climate observations and trends for adaptation purposes, (ii) the assessment of the adequacy and  
43 reliability of available model based climate projections for adaptation needs, (iii) the assessment  
44 of the expected changes in climate extremes needed to assist in developing effective adaptation  
45 and climate risk management strategies, and (iv) the analysis of changes in total water storage for  
46 the period 2002 to 2010 in order to assess recent vulnerability of GHA region to climatic

1 variations. The study is inherently regional in nature since, to be useful, regional climate models  
2 need observational support from as wide a region as possible. Climate does not recognize  
3 national boundaries, so in order to analyze and validate models, it is necessary to take a regional  
4 approach.

5 The presentation is organized as follows; in section 2, the study area, data used, and the analysis  
6 methods are presented. Section 3 presents the results, which are discussed in section 4. The study  
7 is then concluded in Section 5.

## 8 **2. Data and methods**

### 9 **2.1. The Greater Horn of Africa**

10 At the time of this study, the GHA region comprised of Burundi, Djibouti, Ethiopia, Eritrea,  
11 Kenya, Rwanda, Somalia, Sudan, Tanzania and Uganda (Figure 1). But currently, South Sudan  
12 that recently ceased and became sovereign state on 9<sup>th</sup> of July 2011 forms the 11<sup>th</sup> country in the  
13 GHA region. Majority of these countries are classified as least developed where most of the  
14 societies survive on less than one dollar per day (UNDP 2004). Its climate may be classified as  
15 arid and semi-arid with frequent recurrences of floods and droughts. The recurrences of floods  
16 and droughts have been associated with many socio-economic miseries. Furthermore, the region  
17 is often faced with serious food insecurity and resource-based conflicts. For example, 2010-2011  
18 has been shown to be the driest period in 60 years with more than 12 million people in need of  
19 emergency relief (CRS Report, 2012). Recent assessments (IPCC, 2007) showed that climate  
20 change is real and the poor are the most vulnerable due to the already high level of vulnerability  
21 and low coping capacity. The vulnerability is amplified by the fact that many of East Africans  
22 livelihoods are dependent on farming and livestock; two sectors that are especially sensitive to  
23 perturbations in the climate system. Climate change is, therefore, likely to set back development  
24 and food production in many of the predominantly agro-based economies of most communities.  
25 The GHA countries, like other African countries have short and/or fragmented climate records,  
26 often as a result of armed conflict at various times in the last 50 years. In a few cases, the  
27 available records are too short to be used for the adequate calculation of climate trends, but they  
28 are still valuable in providing a baseline for future analyses, as well as monitoring inter-annual  
29 climate variability.

### 30 **2.2. Station data and quality control**

31 It is important to note that three workshops were organized to aid with the collection of data  
32 from the member countries that would hitherto not be possible when organized by an individual  
33 country. The workshop participants brought along with them selection of their best quality  
34 digitized daily temperature and precipitation series (Figure 2).

35 Daily observed station data for maximum and minimum temperatures together with precipitation  
36 for a total of 73 stations from the 10 countries are employed in the analysis (Figure 1 and Table  
37 1). These data sets were subjected to quality control and homogeneity of their series using  
38 RCLimDex software (Peterson et al., 2002; Zhang et al., 2009). RHtest software was also used to  
39 perform homogeneity tests (Aguilar et al., 2005, 2009). These packages were downloaded from  
40 the ETCCDI website<sup>1</sup>.

---

<sup>1</sup><http://cccma.seos.uvic.ca/ETCCDI/>

1 Inhomogeneities in all the time series were identified and corrected. Linear trends were identified  
2 via a least-squares regression analysis with statistical significance assessed using a two-tailed t-  
3 test. These ‘standard’ meteorological observations were compared with spatially interpolated  
4 temperature datasets that have been developed for regional or global applications (Omumbo et  
5 al., 2011). The statistical and visual procedures contained in the RClimDex package were  
6 complemented by the required tests, following the guidelines given in Brunet et al. (2008). In  
7 fact, the tests are focused on the detection of nonsystematic errors usually caused by data  
8 processing, which happens most frequently during digitization procedure (Aguilar *et al.*, 2005,  
9 2009). Ambiguous values (e.g. negative precipitation or maximum temperature lower than  
10 minimum temperature) were identified. Also, the distribution of the precipitation data was  
11 visually inspected, as were plots of the temperature and precipitation time series in order to  
12 detect outlying values.

13 In the case of temperature, statistical outliers, identified as daily values outside a threshold of the  
14 mean value for that particular day plus/minus four standard deviations were also flagged (see  
15 Section. 2.4). The suspicious data were validated, set to missing value or corrected on the basis  
16 of subjective inspection of partial time series for the adjacent days at the same period with other  
17 years and by spatial comparison with those available close neighboring stations.

18 Once the data passed the quality control checks, they were evaluated for homogeneity. The  
19 station data underwent homogeneity testing using the RHtest software package (Aguilar et al.,  
20 2005, 2009), which helps in identifying step changes in a time series by comparing the goodness  
21 of fit of a two-phase regression model with that of a linear trend for the entire series (Wang,  
22 2003, 2008a, 2008b). RHtest is used to help identify series break points for further  
23 investigations. Selection of data for analysis was based on series length and completeness,  
24 quality control and homogeneity (Aguilar et al., 2005, 2009). In this study, we used a base period  
25 of 1961–1990 and 1971–2000 where most of the station data are available. In some cases such as  
26 in Somalia, Sudan and Rwanda; we used a shorter base period; for example, the data supplied for  
27 Gikongoro in Rwanda began in 1967, so we used a base period of 1967–1996. To be included in  
28 the analysis, time series need to be sampled for at least 30 years and contain fewer than 10% of  
29 missing/rejected values. The reference period of 1971–2000 was chosen to maximize the number  
30 of stations with available data for calculation of the percentile-based indices. Even with this  
31 maximization, only 8 countries with a total 58 of the 73 original stations had long enough  
32 homogeneous periods to be included in the analysis, and not all the indices were calculated for  
33 all the stations. Figure 1 shows the location of stations and Figure 2 the data availability for these  
34 stations.

35 Of the ten countries that had participants in the workshop, Somalia and Sudan had data time  
36 series falling outside the 1971–2000 period (Figure 2). Since many of the stations had data  
37 problems prior to 1970, the analysis was limited to the period 1971 to 2009. For percentile-based  
38 indices (e.g., the number of days exceeding the 90th percentile of minimum temperature), the  
39 methodology uses bootstrapping for calculating the baseline period values in order to avoid  
40 discontinuities in the indices time series at the beginning or end of the base period following the  
41 approach by Zhang et al. (2005).

42 The PRECIS modeling system was applied over the region to develop high-resolution climate  
43 scenarios. It was driven with initial and lateral boundary conditions using the UK Met Office  
44 Hadley Center Regional Climate Model (HadRM3P), which is a high-resolution atmospheric  
45 component of the Hadley Centre coupled ocean-atmosphere GCM-HadCM3, with a resolution of  
46  $1.875^{\circ}$  in longitude and  $1.25^{\circ}$  in latitude. Details of the Global Climate Models (GCMs) used in

1 verification of the PRECIS Regional Climate Model (RCM) can be obtained in Simon et al.  
2 (2004).

### 3 **2.3 Gravity Recovery And Climate Experiment (GRACE)**

4 Global and regional water cycles are related to different phenomena within the Earth system,  
5 including variations in atmosphere, hydrosphere, ice cover and land surface in various ways,  
6 while ranging from sub-seasonal and inter-annual to decadal and secular interactions. These all  
7 make it difficult to develop realistic models to simulate or predict water variations.

8 The gravity field of the Earth and its temporal variations, measured globally by GRACE (a joint  
9 US-German satellite mission launched in March 2002) however, are interrelated to total water  
10 storage (TWS) variations of the Earth (Wahr et al., 1998). This offers the unique opportunity to  
11 detect spatio-temporal variations of water within the Earth system from space (Tapley et al.,  
12 2004a, b).

13 The GRACE mission consists of two identical spacecrafts flying approximately 220 km apart in  
14 the same near-polar orbit of about 450 km (Tapley et al., 2004a, b). The main observable is the  
15 distance between the two satellites, measured using a microwave ranging system. Additional  
16 tracking information is provided by Global Positioning System (GPS) receivers on board each of  
17 the spacecraft and Satellite Laser Ranging (SLR) reflector. On-board accelerometers sense non-  
18 conservative forces such as atmospheric drag and solar radiation pressure. Time-variable gravity  
19 field solutions are obtained by the exploitation of GRACE observation data over certain time  
20 intervals, namely daily to monthly gravity field solutions. The solutions are computed in terms of  
21 spherical harmonics (SHs). GRACE time-variable products have been frequently used to study  
22 water variations and their relations to climate change, as documented, e.g., in Ramillien et al.  
23 (2004), Awange et al. (2008, 2009, 2011), Becker, et al. (2010) and Forootan et al. (2012).

24 This study made use of the ITG-GRACE2010 daily solutions (Kurtenbach et al., 2009), which  
25 are computed up to the SHs of degree and order 40, covering October 2002 to September 2009.  
26 For computation of such daily fields, the WaterGAP global hydrology model (WGHM), the  
27 atmospheric model European Centre for Medium-Range Weather Forecasts (ECMWF), and the  
28 ocean circulation model (OMCT) have been used, within the framework of a Kalman-smoother  
29 procedure, to derive the temporal correlations while using GRACE observations as the main  
30 information for computation of the gravity models. Details of computations can be found in  
31 (Kurtenbach, 2011).

32 The priority of using daily solutions for studying water variations when it is compared to those  
33 of monthly solutions is that it allows to recover fast gravity field (its equivalent water) variations  
34 as detailed as possible with reasonable temporal resolution. For studying TWS variation over  
35 the GHA, first 2588 daily gravity solutions covering October 2002 to September 2009 were  
36 downloaded from the official website of the Astronomical Physical und Mathematical Geodesy  
37 (APMG) group at Bonn University<sup>2</sup>. These fields were then used to generate the global TWS  
38 values according to the approach of Wahr et al. (1998) and the boundary of the GHA as it is  
39 shown in Figure 1 was extracted from the fields. Finally, spatially averaged TWS along with  
40 their corresponding cumulated TWS variations were computed (see Figure 7b). Daily GRACE  
41 products, in this study, have been used to examine the impact of climate variability on total  
42 water storage variations for recent years (2002 to 2009) when the station data sets were not

---

<sup>2</sup> <http://www.igg.uni-bonn.de/apmg/index.php?id=itg-grace2010>

1 available. In fact, TWS variability reflected in GRACE data indicates the prolonged impact of  
2 decadal climate change over the region. Discussions on the use of GRACE-TWS for drought  
3 monitoring, for instance, can be found in Houborg et al. (2012).

## 4 **2.4. Analysis methods**

### 5 **2.4.1 Trend calculation and Indices**

6 All stations from different countries were analyzed. Trends for individual stations were  
7 calculated by adapting Sen's (1968) slope estimator. This method has been applied in other  
8 similar works describing extreme indices (Aguilar et al., 2005, 2009; Zhang et al., 2005; Caesar  
9 et al., 2010) and also adapted to climatological data by Zhang et al. (2000) in a study of annual  
10 temperatures over Canada and by Wang and Swail (2001) in their analysis of extreme wave  
11 heights over the Northern Hemisphere. Trends are significant at the 5% level when results are  
12  $\pm 1.96$  standard deviations from the median trend. We require at least 70% of annual data to be  
13 non-missing to calculate a trend and refer to trends as being significant if they are determined to  
14 be statistically significant at the 5% level. To avoid biased estimates, station level trends were  
15 not calculated for series with excessive missing values.

16 A total of 27 indices, based upon recommendations of the ETCCDI, were calculated using  
17 RClimDex (Caesar et al., 2006, 2010). Many of the indices use locally defined thresholds,  
18 making it easier to compare results over a wide region. The indices are primarily based on station  
19 level thresholds calculated over a base period, such as the 90th percentile of minimum  
20 temperature. These thresholds are determined for each day of the year using data from that day  
21 and two days on either side of it over the course of the base period. Table 2 lists the indices  
22 presented in this paper while detailed descriptions of the indices and the exact formulae for  
23 calculating them are available on the ETCCDI web page<sup>3</sup>.

24 All the indices are essentially anomalies from the same base period. However, some precipitation  
25 indices could potentially be dominated by those stations with the greatest precipitation, as those  
26 stations may see precipitation vary from year to year by more than the total annual precipitation  
27 at stations with the least total precipitation (Aguilar et al., 2009). To determine whether this was  
28 the case for the stations analyzed, precipitation indices were also calculated by first standardizing  
29 the indices (dividing by the index's standard deviation). As a comparison of both approaches  
30 revealed similar shape and trends, the standardized indices are not used and the results are  
31 provided through the analysis of the simple anomaly series.

### 32 **2.4.2. Modeling of extreme rainfall and temperature**

33 Using the PRECIS regional climate modeling system, this study analyses the distribution of  
34 extremes of temperature and precipitation in GHA in the recent past (1961–1990) and in a future  
35 (2071–2100) climate under the IPCC SRES A2 and B2 emissions scenarios (IPCC 2007, chapter  
36 19 page 18). Rainfall and temperature were simulated by PRECIS RCM and compared with  
37 observations (in areas in which data are available) for the period 1961–1990. The main task here  
38 is to evaluate the simulations of current climate (particularly precipitation) of regional climate  
39 model by comparing them with currently available data, and thereby assessing the uncertainty  
40 associated with future climate predictions. The UK-Met Office high resolution PRECIS model  
41 runs are compared with available data to assess its performance. The model is used to generate  
42 future climate projections to demonstrate how they can be used and interpreted at the national

---

<sup>3</sup> <http://cccma.seos.uvic.ca/ETCCDI/>



1 level. Emphasis is placed on determining the extremes, trend and the variance explained by each  
2 model. The purpose of the simulation of regional climate is to examine and compare statistics  
3 relevant to the region in observations extremes and regional circulation model. This further  
4 demonstrates value of climate observations and regional models for decision making, to provide  
5 advice on model performance and limitations, and to improve capabilities across the region for  
6 using climate data records and model projections.

### 7 **2.4.3. Temporal Independent Component Analysis of GRACE spatio-temporal total water** 8 **storage products**

9 Independent component analysis (ICA) is a higher-order statistical technique, which can be  
10 viewed as an extension to the commonly used Principle Component Analysis (PCA) (Forootan  
11 and Kusche, 2012, 2013). Using an ICA algorithm, the input data (spatio-temporal observations)  
12 are assumed to consist of a linear mixture of unknown source signals, which cannot be directly  
13 measured. By incorporating higher order statistical information contained in the data in the  
14 decomposition procedure, ICA extracts statistically independent components that reflect spatial  
15 and temporal manifestations of physical processes hidden in the data (Lotsch et al., 2003;  
16 Hannachi et al., 2009).

17 Generally, there are two alternative ways to implement ICA on a temporal sequence of gridded  
18 datasets in which either temporally independent components or spatially independent time series  
19 can be estimated. The methods are respectively called temporal ICA and spatial ICA (for details  
20 see e.g., Forootan and Kusche, 2012 and Forootan et al., 2012). This study made use of temporal  
21 ICA method, since the scope of the study is to extract the temporal behavior of TWS changes for  
22 the period 2002 to 2010, to further investigate the impacts of rainfall after the long-term study of  
23 1970 to 2000.

## 24 **3. Results**

### 25 **3.1 Trends in temperature indices**

26 Trends for the temperature indices for some selected countries are shown in Table 3 in  
27 comparison to global and other regional indices. The two countries i.e., Ethiopia and Kenya are  
28 used for comparison purposes since their data coverage is relatively good and trends calculated  
29 include similar window period of 1971 to 2003. The warm extremes are increasing while cold  
30 extremes decreasing, these series clearly indicate significant warming. Individual stations show  
31 most spatial coherence in the TN90p index, that is, frequency of nights warmer than the 90th  
32 percentile. Nearly half of the available stations indicate a significant increase in this index over  
33 the period 1971–2004. Sample time series for the percentile-based temperature indices are shown  
34 in Fig. 3 for Asmara in Eritrea. The frequencies of warm days and nights, relative to the base  
35 period 1961–1990, increased strongly between 1961 and 1990, with a large increase in the  
36 number of nights per year exceeding the 90th percentile threshold. There were also large  
37 reductions in the frequency of cold nights and cold days over the 49 years. The warmest day and  
38 night of the year is warming at a rate approximately comparable to the global average.

39 In general, over the entire region, the frequency of warm days and warm nights has increased,  
40 and the frequency of cold days and cold nights has decreased. This agrees with the results from  
41 other studies that have analyzed these trends across different parts of the world (Griffiths et al.,  
42 2005; Klein Tank et al., 2006; Choi et al., 2009; Caesar et al., 2010). However, the results for the  
43 absolute temperature indices (TXx, TNx, etc.) defined for the entire region are sensitive to the  
44 large variability in these indices across the region. The percentile indices (e.g. TN90p) are more  
45 robust across large regions because they account for the influence of local climate effects. There

1 has been a significant increase in the absolute annual maximum of both daily maximum and  
2 minimum temperatures, again in common with the global picture (Griffiths et al., 2005; Klein  
3 Tank et al., 2006; Choi et al., 2009; Caesar et al., 2010). The coldest day and night of the year is  
4 warming slower than the global average, although planetary trend for the coldest day is not  
5 significant.

### 6 **3.2 Trends in precipitation indices**

7 The map in Figure 4b depicts total precipitation in wet days ( $> 1$  mm) (PRCPTOT) and it is an  
8 example indicating relative lack of spatial coherence for precipitation in the region. The number  
9 of stations with significant negative or positive trends is low. Sample graphs of trends, calculated  
10 for various precipitation indices are shown in Figures 4a, 5a and 5b, while Table 4 compares  
11 similar trends at regional and global scales (Alexander et al. 2006, Caesar et al., 2010) at similar  
12 window period. Western Lake Victoria, southern Sudan and western Ethiopia generally show  
13 significant decreases in total precipitation (e.g., see Figure 4b). For Asmara (Figure 4a) and  
14 Djibouti (Figure not shown), there are sharp drops in the total annual precipitation time series  
15 around 2000 to 2010. Likely associated with the decrease in total precipitation, the length of the  
16 maximum number of consecutive dry days is increasing in Asmara and Djibouti, while the length  
17 of the maximum number of consecutive wet days shows a significant decrease (figure not  
18 shown). The Simple Daily Intensity Index (SDII), which takes into account the number of days  
19 with rainfall greater than or equal to 1mm shows no significant changes. In general, decreasing  
20 trend in total precipitation in wet days ( $> 1$  mm) is observed in the north western sector (western  
21 Ethiopia and southern Sudan), and equatorial sector around Lake Victoria, while much of  
22 Ethiopia had significant positive increase (Figure 4b). The precipitation due to very wet days  
23 greater than 95<sup>th</sup> percentile (R95p) index (Figure 5(a)) indicate that the annual amount of  
24 precipitation contributed on days exceeding the long-term 95th percentile has decreased from  
25 about 50 mm to around 30 mm in Khartoum, but this change is non-significant (Figure 5a). The  
26 highest precipitation amount in five-day period or maximum 5-day precipitation (RX5day) index  
27 (Figure 5b) depicts significant reduction over the 40-year period.

28 Similar time series of R95p for the southern sector is represented by Dodoma (Figure 6a). There  
29 are increases in R95p over the southern sector, a marginal decrease over the equatorial sector,  
30 and a decrease over the northern sub-region. Between them, only the change within the southern  
31 sector is statistically significant.

32 Table 4 lists the regional trends for the precipitation indices and also the global trends. The same  
33 problems exist with defining some of the precipitation indices across the whole region that  
34 applied to the absolute temperature indices, and indices defined relative to a local climatology  
35 (e.g. percentile based) are preferable for comparing across such a large region.

36 Compared to the temperature indices, there are fewer significant trends in the precipitation  
37 indices. In contrast to the other sub-regions, the northern sector has decreasing trends in all  
38 precipitation indices, apart from the consecutive dry day index, suggesting a consistent change  
39 towards drier conditions. However, it must be emphasized that these trends are non-significant.  
40 Over the region as a whole, the precipitation trends are mixed (Funk et al., 2008, 2012). This  
41 does not parallel the global results of Caesar et al. (2010) indicating consistent trends towards  
42 wetter conditions across nearly all of the indices, although it should be noted that analysis was  
43 for a different time period (1971–2005) and had only limited coverage of the tropics.

### 44 **3.3 Relationship between precipitation and TWS changes**

1 Regarding the precipitation results, it was clear that the overall precipitation over the GHA is  
2 declining. To support the precipitation results of the last 7 years of the study (2002-2009), we  
3 used daily TWS products as described in Section 2.3. The goal was to see whether the total water  
4 availability of the region is affected by climate variations or not. As a matter of fact, TWS tells  
5 quite more sophisticated story of water variations over the study region by providing information  
6 on daily precipitation minus evaporation minus run-off over the region. Our results of spatially-  
7 averaged TWS over the GHA (Fig. 7) show that TWS declined between 2002 and 2007. An  
8 increase in TWS in 2007 is associated with the El Niño southern oscillation phenomenon, which  
9 usually brings rainfall to most parts of the region (Ogallo et al., 1988; Janowiak, 1988; Indeje,  
10 2000; Mutemi, 2003). This has been followed again by a decline in TWS variations over the  
11 GHA up to the end of the study period. The cumulative TWS over the study period supports the  
12 results of precipitation (e.g., Figure 4a), showing that the total water availability has decreased  
13 over the GHA during the last 7 years of the study (see Fig. 7b, bottom).

14 Applying the ICA method on GRACE-TWS data over the GHA shows that the first ICA mode  
15 (IC1) extracts 75% of variability in TWS changes. The spatial pattern of IC1 shows a dipole  
16 spatial structure with respect to the Equator. The temporal pattern of IC1 shows a dominant  
17 annual water cycle over the study area. The second ICA mode (IC2) corresponds to 15% of the  
18 variance, while the temporal pattern shows a summation of a long-term trend and an inter-annual  
19 variability. Considering the spatial pattern, therefore, a declining rate for the regions of the Lake  
20 Victoria Basin and the surrounding lakes were found. The declining pattern of TWS is also  
21 extended up to the south of Sudan (c.f., figure 4b). In contrast, over the tropical regions as well  
22 as Ethiopia a slight increase of TWS during 2002 to 2010 is seen (c.f., figure 4b). The spatial and  
23 temporal patterns of IC2, therefore, confirm the results of precipitation, which was derived for  
24 the long-term period during 1970 to 2000.

### 25 **3.4 Modeling precipitation extremes**

26 PRECIS Regional Climate Model (RCM) analyzed correctly and reproduced the mean seasonal  
27 and annual cycles of precipitation for the period 1961-1990 over the southern (Figure 8a),  
28 northern (Figure 8b) and equatorial (Figure 8c) sectors. The mean surface temperature  
29 climatology for all the four seasons of the region i.e. March-May (MAM), June-August (JJA),  
30 October-December (OND) and December-February (DJF) are spatially and temporally (Figure  
31 8d) simulated for the baseline period of 1961-1990 compared with the observed Climate  
32 Research Unit (CRU) data (gridded data based on the set aggregated to the RCM grid). The  
33 results show that, compared to CRU, the model underestimates rainfall over most parts of eastern  
34 highlands during OND while over the central sector (around Lake Victoria area) and southern  
35 sector of the region, the model overestimates rainfall (Figure 8a). On the contrary, over northern  
36 sector, the model produces the observed rainfall reasonably well (figure not shown). It is thus  
37 evident that the simulated rainfall by the PRECIS model is fairly consistent with the observed  
38 values over most parts of the study area. Results from the inter-annual variability of PRECIS  
39 simulated surface air temperature showed that during JJA season, some sections of the region  
40 recorded the lowest temperature and the warmest temperatures during the DJF season (Figure  
41 8b). The model results agree reasonably well with the observed patterns in terms of the spatial  
42 location of the extreme maximum temperatures.

43 The trends calculated for projected precipitation indices for the period 2010-2040 are shown in  
44 Figure 9. Only central Uganda represented by Mbarara, Masindi and Jinja show significant  
45 increases in total precipitation. Sample time series for Gulu is shown by Figure 9a and regional  
46 projected trends in Figure 9b. Total annual rainfall (PRCPTOT) showed a decreasing trend for

1 many of the stations over Sudan except for the Khartoum that had significant positive trend.  
2 Generally, consecutive wet days (CWD) showed a decreasing trend while the consecutive dry  
3 days (CDD) showed an increasing trend. While temperature indices vary from station to station,  
4 the dominant features seem to be an increasing trend in the number of cold nights (TN10p) with  
5 a decreasing trend in the number of warm nights (TN90p). Time series for projected Total  
6 Rainfall (PRCPTOT) over Rwanda, where three stations (Kigali, Kamembe, and Gikongoro)  
7 were analyzed showed evidence of decreasing trends in rainfall. Results show general increasing  
8 trend in Consecutive Dry Days (CDD) and decreasing trend for Consecutive Wet Days (CWD)  
9 for Gikongoro.

#### 10 **4. Discussion**

11 A set of daily station observations from countries in the GHA region were for the first time  
12 compiled and analyzed to enable assessment of changes in climate extremes over the region.  
13 Most stations showed decrease of total precipitation in wet days greater than 1 mm (PRCPTOT)  
14 as well as heavy rainy days (R10mm), maximum one-day precipitation (Rx1day), maximum  
15 five-day precipitation (Rx5day), heavy precipitation days (R10mm) and warm spell duration  
16 (WSDI). Index for warm days (Tx90P) showed increasing trend, while the index for cool days  
17 (TX10P) showed decreasing trend. The TXn index (monthly minimum value of daily maximum  
18 temperature) showed an increasing trend in most parts of the region. The TNx index (monthly  
19 maximum value of daily minimum temperature) showed an increasing trend over most parts of  
20 the region, and more significantly in the north-eastern and southern sectors of the region;  
21 indicating that there is a general increasing trend of warm nights for most of the stations in the  
22 region. Increasing trend in warm nights is indicative of significant night time warming.

23 Thus, increasing trends for warm nights were the most spatially coherent index consistent with  
24 the results of other regional workshops (Klein Tank et al., 2006, Choi et al., 2009) and the global  
25 analysis (Alexander et al., 2006, Caesar et al., 2010). Less spatial coherence trends in  
26 precipitation indices across the region and fewer trends that are locally significant when  
27 compared with the temperature indices are observed. In the few cases where statistically  
28 significant trends in precipitation indices are identified for regions and sub-regions, there is  
29 generally a trend towards wetter conditions consistent with the global results of Alexander et al.  
30 (2006).

#### 31 **5. Conclusion**

32 The study aimed at:

- 33 (i) Assessing the adequacy of regional climate observations and trends for adaptation  
34 purposes. In this regard, the study found that there is inadequate in-situ data for an  
35 individual country analysis but provided sufficient ground for regional level climate  
36 analysis. The results further showed increasing/decreasing trend in warm/cold extremes.  
37 Furthermore, frequencies of warm days and nights increased strongly, with a large  
38 increase in the number of nights per year exceeding the 90th percentile threshold  
39 between 1961 and 1990. On the contrary, precipitation patterns are mixed with fewer  
40 significant trends except significant decrease in total precipitation in wet days greater  
41 than 1 mm across the whole region.
- 42 (ii) Assessing the adequacy and reliability of available model based climate projections for  
43 adaptation needs. To this end, the study found that the simulated climate is fairly  
44 consistent with the observed values over most parts of the study area.

- 1 (iii) Assessing the expected changes in climate extremes needed to assist in developing  
2 effective adaptation and climate risk management strategies. Here, the study established  
3 that, generally, the model projected decreasing/increasing trend in consecutive wet/dry  
4 days with variations in temperature indices from one station to another. The dominant  
5 features in the region seem to be an increasing trend in the number of cold nights with a  
6 decreasing trend in the number of warm nights. Increasing trends for warm nights were  
7 the most spatially coherent index consistent with the results of other regions of the globe.
- 8 (iv) Assessing changes in the total water storage for the period 2002-2010. Here, the study  
9 established a decline in total water availability over the GHA region during the last 7  
10 years.

11 Our findings therefore showed that increasing trends in both night and day temperatures had the  
12 most spatially coherent indices. The model simulates well the spatial distribution of extreme  
13 temperature and rainfall events when compared with present climate observations, with  
14 temperature simulation being more realistic across the region. Overall, the future occurrence of  
15 warm days and nights are projected to be more frequent in the entire GHA, while the occurrence  
16 of cold night events is likely to decrease. The overall precipitation in the region decreased  
17 between 2002 and 2007 from the TWS products.

### 18 **Acknowledgements**

19 The authors would like to thank R. Huth (editor) and two anonymous reviewers whose  
20 comments considerably improved the manuscript. The Nairobi workshop was funded by the  
21 World Bank's Global Facility for Disaster Reduction and Recovery (GFDRR) Track II support  
22 and implemented through World Meteorological Organization (WMO); World Climate Research  
23 Programme (WCRP); Global Climate Observing System (GCOS). The workshop participants  
24 would like to thank all the National Meteorological and Hydrological Services (NMHSs) of the  
25 respective countries for providing daily rainfall and temperature data used in the analysis.  
26 Special thanks to United Kingdom (UK) Met Office for offering expertise to the workshops.

27

## 1 References

- 2 Alexander, L.V., Zhang X., Peterson, T.C., Caesar, J., Gleason, B., Tank, A.M.G.K., Haylock,  
3 M., Collins, D., Trewin, B., Rahimzadeh, F., Tagipour, A., Kumar, K.R., Revadekar, J.,  
4 Griffiths, G., Vincent, L., Stephenson, D.B., Burn, J., Aguilar, E., Brunet, M., Taylor, M.,  
5 New, M., Zhai, P., Rusticucci, M., Vazquez-Aguirre, J.L., 2006. Global observed changes  
6 in daily climate extremes of temperature and precipitation. *Journal of Geophysical*  
7 *Research* 111: D05109, DOI: 10.1029/2005JD006290.
- 8 Aguilar, E., Peterson, T. C., Ramírez Obando, P., Frutos, R., Retana, J. A., Solera, M., Soley,  
9 J., González Garcí a, I., Araujo, R. M., Rosa Santos, A., Valle, V. E., Brunet, M., Aguilar,  
10 L., ´lvarez, L. A., Bautista, M., Castañó n, C., Herrera, L., Ruano, E., Sinay, J. J.,  
11 Sa´nchez, E., Hern´andez Oviedo, G. I., Obed, F., Salgado, J. E., Va´zquez, J. L., Baca, M.,  
12 Guti´rrez, M., Centella, C., Espinosa, J., Mart´ınez, D., Olmedo, B., Ojeda Espinoza, C.  
13 E., Nu´ñez, R., Haylock, M., Benavides, H., Mayorga, R., 2005. Changes in precipitation  
14 and temperature extremes in Central America and northern South America, 1961 – 2003, *J.*  
15 *Geophys. Res.*, 110, D23107, doi: 10.1029/2005JD006119.
- 16 Aguilar, E., Aziz Barry, A., Brunet, M., Eakang, L., Fernandes, A., Massoukina, M., Mbah, J.,  
17 Mhanda, A., do Nascimento, D.J., Peterson, T.C., Thamba Umba, O., Tomou, M., Zhang,  
18 X., 2009. Changes in temperature and precipitation extremes in western central Africa,  
19 Guinea Conakry, and Zimbabwe, 1955–2006. *Journal of Geophysical Research* 114:  
20 D02115, DOI: 10.1029/2008JD011010.
- 21 Anyah, R.O., Semazzi, F.H., 2006. Climate variability over the Greater Horn of Africa based  
22 on NCAR AGCM ensemble. *Theoretical and Applied Climatology* 86: 39–62.
- 23 Anyah, R.O., Qiu, W., 2012. Characteristic 20th and 21st century rainfall and temperature  
24 patterns and changes over the Greater Horn of Africa. *International Journal of*  
25 *Climatology*, 32, 347–363; DOI: 10.1002/joc.2270.
- 26 Arnell, N.W., 2004. Climate change and global water resources: SRES emissions and socio-  
27 economic scenarios. *Global Environmental Change* 14(1): 31-52, doi:  
28 10.1016/j.gloenvcha.2003.10.006.
- 29 Awange, J., Sharifi, M., Ogonda, G., Wickert, J., Grafarend, E., Omulo, M., 2008. The falling  
30 Lake Victoria water level: GRACE, TRIMM and CHAMP satellite analysis of the lake  
31 basin. *Water Resource Management*, 22(7), 775–796. [http://dx.doi.org/10.1007/s11269-](http://dx.doi.org/10.1007/s11269-007-9191)  
32 [007-9191](http://dx.doi.org/10.1007/s11269-007-9191).
- 33 Awange, J., Sharifi, M., Baur, O., Keller, W., Featherstone, W., Kuhn, M., 2009. GRACE  
34 hydrological monitoring of Australia: Current limitations and future prospects. *Journal of*  
35 *Spatial Science*, 54(1), 23–36. <http://dx.doi.org/10.1080/14498596.2009.9635164>.
- 36 Awange, J., Fleming, K. M., Kuhn, M., Featherstone, W. E., Heck, B., Anjasmara, I., 2011. On  
37 the suitability of the 4°×4° GRACE mascon solutions for remote sensing Australian  
38 hydrology. *Remote Sensing of Environment*, 115(3), 864–875.  
39 <http://dx.doi.org/10.1016/j.rse.2010.11.014>.
- 40 Becker, M., Llovel, W., Cazenave, A., Günter, A., & Crétau, J. -F. 2010. Recent hydrological  
41 behavior of the East African great lakes region inferred from GRACE, satellite altimetry  
42 and rainfall observations. *Comptes Rendus Geoscience*, 342(3), 223–233.  
43 <http://dx.doi.org/10.1016/j.crte.2009.12.010>.

- 1 Bohle, H.G., Downing, T.E., Watts, M.J., 1994. Climate change and social vulnerability:  
2 Toward a sociology and geography of food insecurity. *Global Environmental Change* 4(1):  
3 37-48, doi: 10.1016/0959-3780(94)90020-5.
- 4 Bradfield, L., David, G., DeWitt, 2012. A recent and abrupt decline in the East African long  
5 rains, *Geophysical Research Letters*, vol. 39, 102702, doi: 10.1029/2011GL050337, 2012
- 6 Brant L., Ileana, B., George, N., Leila, K., Carvalho, M.V., Gabriel, B., Senay, D.A., Leroux,  
7 S., Funk, C., 2012. Seasonality of African Precipitation from 1996 to 2009, *Journal of*  
8 *Climate*, June 2012, Vol. 25, No. 12: pp. 4304-4322, (doi: 10.1175/JCLI-D-11-00157.1)
- 9 Brunet, M., Saladie´, O., Jones, P., Aguilar, E., Moberg, A., Lister, D., Walther, A., Almarza,  
10 C., 2008. A case-study/guidance on the development of long-term daily adjusted  
11 temperature datasets, WCDMP-66, 46 pp., World Meteorol. Org., Geneva.
- 12 Caesar, J., Alexander, L., Vose, R., 2006. Large-scale changes in observed daily maximum and  
13 minimum temperatures – creation and analysis of a new gridded dataset; *Journal of*  
14 *Geophysical Research* 111: D05101, DOI: 10.1029/2005JD006280.
- 15 Caesar, J.A., Alexander, L.V., Trewin, B., Tse-ring, K., Sorany, L., Vuniyayawa, V.,  
16 Keosavang, N., Shimana, A., Htay, M.M., Karmacharya, J., Jayasinghearachchi, D.A.,  
17 Sakkamart, J., Soares, E., Hung, L.T., Thuong, L.T., Hue, C.T., Dung, N.T.T., Hung, P.V.,  
18 Cuong, H.D., Cuong, N.M., Sirabaha, S., 2010. Changes in temperature and precipitation  
19 extremes over the Indo-Pacific region from 1971 to 2005, *Int. J. Climatol.*, Published  
20 online in Wiley InterScience ([www.interscience.wiley.com](http://www.interscience.wiley.com)) DOI: 10.1002/joc.2118
- 21 Camberlin, P., Philippon, N., 2002. The East African March-May rainy season, its  
22 teleconnections and predictability over the 1968-1997 period. *Journal of Climate*: 15,  
23 1002-1019.
- 24 Choi, G., Collins, D., Ren, G., Trewin, B., Baldi, M., Fukuda, Y., Afzaal, M., Pianmana, T.,  
25 Gomboluudev, P., Huong, P.T., Lias, N., Kwon, W.T., Boo, K.O., Cha, Y.M., Zhou, Y.,  
26 2009. Changes in means and extreme events of temperature and precipitation in the Asia-  
27 Pacific Network region, 1955–2007. *International Journal of Climatology* 29: 1906–1925.
- 28 Christy J. R, Norris W. B, McNider R. T., 2009: Surface Temperature Variations in East Africa  
29 and Possible Causes. *Journal of Climate* 2009, **22**:3342-3356.
- 30 Conway, D., C. Mould and W. Bewket, 2004: Over one century of rainfall and temperature  
31 observations in Addis Ababa, Ethiopia. *Int. J. Climatol.* **24**: 77–91 (2004)
- 32 CSR Report, 2012: Congressional Research Service report on Horn of Africa: The  
33 Humanitarian Crisis and International Response (<http://www.fas.org/sgp/crs/row/R42046.pdf>)
- 34 Downing, T.E., 1991. Vulnerability to hunger in Africa: A climate change perspective. *Global*  
35 *Environmental Change* 1(5): 365-380, doi: 10.1016/0959-3780(91)90003-C
- 36 Elagib N. A., 2010: Changing rainfall, seasonality and erosivity in the hyper-arid zone of  
37 Sudan. Published online 22 July 2010 in Wiley Online Library ([wileyonlinelibrary.com](http://wileyonlinelibrary.com))  
38 DOI: 10.1002/ldr.1023

- 1 Forootan, E., Awange, J., Kusche, J., Heck, B., A. Eicker, 2012. Independent patterns of water  
2 mass anomalies over Australia from satellite data and models. *Journal of Remote Sensing*  
3 of Environment, 124, 427-443, doi:0.1016/j.rse.2012.05.023.
- 4 Forootan, E., Kusche, J., 2012. Separation of global time-variable gravity signals into  
5 maximally independent components. *Journal of Geodesy*, 86 (7), 477-497, DOI:  
6 10.1007/s00190-011-0532-5.
- 7 Forootan, E., Kusche, J., 2013. Separation of deterministic signals, using independent  
8 component analysis (ICA). *Stud. Geophys. Geodetic.* 57 (1), 17-26, DOI: 10.1007/s11200-  
9 012-0718-1.
- 10 Folland, C.K., Karl, T.R., Christy, J.R., Clarke, R.A., Gruza, G.V., Jouzel, J., Mann, M.E.,  
11 Oerlemans, J., Salinger, M.J, Wang, S.W., 2001. Observed climate variability and change.  
12 In *Climate Change 2001. The Scientific Basis – Contribution of Working Group I to the*  
13 *Third Assessment Report of the Intergovernmental Panel on Climate Change.* Cambridge  
14 University Press: New York; pp 99–181.
- 15 Funk, C., Dettinger, M. D., Michaelsen, J. C., Verdin, J. P., Brown, M. E., Barlow, M., and  
16 Hoell, A. (2008). Warming of the Indian Ocean threatens eastern and southern African  
17 food security but could be mitigated by agricultural development. *Proceedings of the*  
18 *National Academy of Sciences*, 105(32), 11081-11086.
- 19 Funk, C., Michaelsen, J., and Marshall, M. T. (2012). 14 Mapping Recent Decadal Climate  
20 Variations in Precipitation and Temperature across Eastern Africa. *Remote Sensing of*  
21 *Drought: Innovative Monitoring Approaches*, 331
- 22 Griffiths, G.M., Chambers, L.E., Haylock, M.R., Manton, M.J., Nicholls, N., Baek, H.J., Choi,  
23 Y., Della Marta, P.M., Gosai, A., Iga, N., Lata, R., Laurent, V., Maitrepierre, L.,  
24 Nakamigawa, H., Ouprasitwong, N., Solofa, D., Tahani, L., Thuy, D.T., Tibig, L., Trewin,  
25 B., Vediapan, K., Zhai, P., 2005. Change in mean temperature as a predictor of extreme  
26 temperature change in the Asia-Pacific region. *International Journal of Climatology* 25:  
27 1301–1330.
- 28 Hannachi, A., Unkel, S., Trendafilov, N.T., Jolliffe, I.T., 2009. Independent component  
29 analysis of climate data: a new look at EOF rotation. *J Clim* 22:2797–2812.  
30 doi:10.1175/2008JCLI2571.1.
- 31 Hastenrath, S., Polzin, D., Mutai, C., 2007. Diagnosing the 2005 drought in equatorial East  
32 Africa. *Journal of Climate* 29: 4628-4637.
- 33 Hastenrath, S., Polzin, D., Mutai, C., 2010. Diagnosing the Droughts and Floods in Equatorial  
34 East Africa during Boreal Autumn 2005–08. *J Climate* 23: 813–817
- 35 Hay SI, Rogers DJ, Randolph SE, Stern DI, Cox J, Shanks GD, Snow RW: Hot topic or hot air?  
36 Climate change and malaria resurgence in East African highlands. *Trends Parasitol* 2002,  
37 18:530-534.
- 38 Haylock, M.R., Hofstra, N., Klein Tank, A.M.G., Klok, E.J., Jones, P.D., New, M., 2008. A  
39 European daily high-resolution gridded data set of surface temperature and precipitation  
40 for 1950–2006. *Journal of Geophysical Research* 113: D20119, DOI:  
41 10.1029/2008JD010201.




- 1 Houborg, R., Rodell, M., Li, B., Reichle, R., Zaitchik, B. F., 2012. Drought indicators based on  
2 model-assimilated Gravity Recovery and Climate Experiment (GRACE) terrestrial water  
3 storage observations. *Water Resources Research*, 48 (7), DOI: 10.1029/2011WR011291
- 4 Indeje, M., Semazzi, F.H.M., Ogallo, L.J., 2000. ENSO signals in East African rainfall and  
5 their prediction potentials. *Int. J. Climatol.* 20, 19-46.
- 6 IPCC, 2007. Climate change 2007: the physical science basis. In Contribution of Working  
7 Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate  
8 Change, Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M,  
9 Miller HL (eds). Cambridge University Press: Cambridge, New York.
- 10 Janowiak, J., 1988. An investigation of Interannual Rainfall Variability in Africa, *J. Clim.* 1,  
11 240-255.
- 12 Kijazi, A.L., Reason, C.J.C., 2009a. Analysis of the 2006 floods over northern Tanzania.  
13 *International Journal of climatology* 29: 955-970.
- 14 Kijazi, A.L., Reason, C.J.C., 2009b. Analysis of the 1998 to 2005 drought over the northeastern  
15 highlands of Tanzania. *Climate Research* 38: 209-223
- 16 Klein Tank, A.M.G., Peterson, T.C., Quadir, D.A., Dorji, S., Zou, X., Tang, H., Santhosh, K.,  
17 Joshi, U.R., Jaswal, A.K, Kolli, R.K, Sikder, A., Deshpande, N.R., Revadekar, J.V.,  
18 Yeleuova, K., Vandasheva, S., Faleyeva, M., Gomboluudev, P., Budhathoki, K.P, Hussain,  
19 A., Afzaal, M., Chandrapala, L., Anvar, H., Amanmurad, D., Asanova, V.S., Jones, P.D.,  
20 New, M.G., Spektorman, T., 2006. Changes in daily temperature and precipitation  
21 extremes in central and south Asia, *Journal of Geophysical Research- Atmospheres*  
22 111(D16105): DOI: 10.1029/2005JD006316.
- 23 Kurtenbach, E., Mayer-Gürr, T., Eicker, A., 2009. Deriving daily snapshots of the Earth's  
24 gravity field from GRACE LIB data using Kalman filtering, *Geophys. Res. Lett.*, 36,  
25 L17102, doi:10.1029/2009GL039564.
- 26 Kurtenbach, E., 2011. Entwicklung eines Kalman-Filters zur Bestimmung kurzzeitiger  
27 Variationen des Erdschwerefeldes aus Daten der Satellitenmission GRACE. PhD  
28 dissertation, Bonn University, Germany.
- 29 Lotsch, A., Friedl, M.A., Pinzón, J., 2003. Spatio-Temporal Deconvolution of NDVI Image  
30 Sequences Using Independent Component Analysis. *IEEE Transactions on Geoscience and*  
31 *Remote Sensing*, 41 (12)
- 32 Lyon, B., and DeWitt, D. G. (2012). A recent and abrupt decline in the East African long rains.  
33 *Geophysical Research Letters*, 39(2), L02702.
- 34 Mutemi, J.N., 2003. Climate anomalies over eastern Africa associated with various ENSO  
35 evolution phases: Ph.D. Thesis, University of Nairobi, Kenya.
- 36 Ogallo, L.J., Janowiak, J.E., Halpert, M.S., 1988. Teleconnection between seasonal rainfall  
37 over Eastern Africa and Global Sea surface temperature anomalies. *J. Meteor. Soc., Japan*,  
38 66, 807 – 822
- 39 Omondi, P., Awange, J.L., Ogallo, L.A., Okoola, R.A., Forootan, E., 2012. Decadal rainfall  
40 variability modes in observed rainfall records over East Africa and their relations to  
41 historical sea surface temperature changes. *J. Hydrol.*,  
42 <http://dx.doi.org/10.1016/j.jhydrol.2012.07.003>

- 1 Omondi, P.A., 2011. Agricultural Drought Indices, Proceedings of an Expert Meeting 2-4 June,  
2 2010, Murcia, Spain, WMO, AGM-11WMO/TD No. 1572, WAOB-2011, pp 106-112
- 3 Omondi, P., L. A. Ogallo, R. Anyah, J. M. Muthama and J. Ininda, 2012: Linkages between  
4 global sea surface temperatures and decadal rainfall variability over Eastern Africa region.  
5 Int. J. Climatol. DOI: 10.1002/joc.3578
- 6 Omumbo J. A., Bradfield L., Samuel M. W., Connor S. J., Madeleine C. T., 2011: Raised  
7 temperatures over the Kericho tea estates: revisiting the climate in the East African  
8 highlands malaria debate Malaria Journal 2011, 10:12  
9 <http://www.malariajournal.com/content/10/1/12>
- 10 Parmesan, C., Root, T.L. Willig, M.R., 2000. Impacts of extreme weather and climate on  
11 terrestrial biota, Bull. Am. Meteorol. Soc., 81, 443– 450, doi: 10.1175/1520-  
12 0477(2000)081<0443:IOEWAC>2.3.CO;2.
- 13 Pascal M., J. A. Ahumada, L. F. Chaves, X. Rodo, and M. Bouma, 2006: Malaria resurgence in  
14 the East African highlands: Temperature trends revisited. PNAS, April 11, 2006 vol. 103  
15 15, 5829–5834. [www.pnas.org/cgi/doi/10.1073/pnas.0508929103](http://www.pnas.org/cgi/doi/10.1073/pnas.0508929103)
- 16 Peterson, T.C., Manton, M.J., 2008. Monitoring changes in climate extremes: A tale of  
17 international collaboration, Bull. Am. Meteorol. Soc., 89, 1266– 1271.
- 18 Peterson, T.C., Easterling, D.R, Karl, T.R., Groisman, P., Nicholls, N., Plummer, N., Torok, S.,  
19 Auer, I., Bohm, R., Gullett, D., Vincent, L., Heino, R., Tuomenvirta, H., Mestre, O.,  
20 Szentimrey, T., Salinger, J., Førland, E.J., Hanssen-Bauer, I., Alexandersson, H., Jones, P.,  
21 Parker, D., 1998. Homogeneity adjustments of in situ atmospheric climate data: a review.  
22 International Journal of Climatology 18: 1493–1517.
- 23 Ramillien, G., Cazenave, A., Brunau, O., 2004. Global time variations of hydrological signals  
24 from GRACE satellite gravimetry. Geophysical Journal International, 158(3), 813–826.  
25 <http://dx.doi.org/10.1111/j.1365-246X.2004.02328.x>.
- 26 Tapley, B., Bettadpur, S., Ries, J., Thompson, P., Watkins, M., 2004a. GRACE measurements  
27 of mass variability in the Earth system. Science, 305, 503–505.  
28 <http://dx.doi.org/10.1126/science.1099192>.
- 29 Tapley, B., Bettadpur, S., Watkins, M., Reigber, C., 2004b. The gravity recovery and climate  
30 experiment: Mission overview and early results. Geophysical Research Letters,  
31 31(L09607). <http://dx.doi.org/10.1029/2004GL019920>.
- 32 Tierney, J.E.; Smerdon, J.E.; Anchukaitis, K.J.; Seager, R. (2013). Multidecadal variability in  
33 East African hydroclimate controlled by the Indian Ocean Nature (Lond.) 493(7432): 389-  
34 392. [hdl.handle.net/10.1038/nature11785](http://hdl.handle.net/10.1038/nature11785)
- 35 Sen, P.K., 1968. Estimates of regression coefficient based on Kendalls Tau, J. Am. Stat.  
36 Assoc., 63, 1379– 1389, doi:10.2307/2285891.
- 37 Schreck, C. J., Semazzi, F.H.M., 2004. Variability of the recent climate of eastern Africa. Int. J.  
38 Climatol., 24, 681-701.
- 39 Shongwe, M.E., van Oldenborgh, G.J., van den Hurk, B.J.J.M., van Aalst, M.K., 2010.  
40 Projected changes in mean and extreme precipitation in Africa under global warming. Part  
41 II: East Africa. J. Climate, 22, 3819–3837.

- 1 Shongwe, M.E., van Oldenborgh, G.J., van den Hurk, B., van Aalst, M., 2011. Projected  
2 changes in mean and extreme precipitation in Africa under global warming. Part II: East  
3 Africa. *J Clim* 24:3718–3733
- 4 Simon, W., Hassell, D., Hein, D., Jones, R., Taylor, R., 2004. Installing and Using the Hadley  
5 Centre Regional Climate Modelling System, PRECIS, Version 1.1. Met Office Hadley  
6 Centre: Exeter.
- 7 UNDP, 2004: Disasters for Least Developed Countries.
- 8 Viste, E., Korecha, D., and Sorteberg, A. (2012). Recent drought and precipitation tendencies  
9 in Ethiopia. *Theoretical and Applied Climatology*, 1-17.
- 10 van Oldenborgh, G.J., Philip, S.Y., Collins, M., 2005. El Niño in a changing climate: a multi-  
11 model study. *Ocean Science* 1: 81-95.
- 12 Wang, X.L., Swail, V.R., 2001. Changes of extreme wave heights in Northern Hemisphere  
13 oceans and related atmospheric circulation regimes, *J. Clim.*, 14, 2204 – 2221,  
14 doi:10.1175/1520-0442(2001)014<2204:COEWHI>2.0.CO;2.
- 15 Wang, X.L., 2003. Comments on “Detection of undocumented change points: a revision of the  
16 two-phase regression model”. *Journal of Climate* 16: 3383–3385.
- 17 Wang, X.L.L., 2008a. Accounting for autocorrelation in detecting mean shifts in climate data  
18 series using the penalized maximal t or F test. *Journal of Applied Meteorology and*  
19 *Climatology* 47: 2423–2444, DOI: 10.1175/2008JAMC1741.1.
- 20 Wang, X.L.L., 2008b. Penalized maximal F test for detecting undocumented mean shift without  
21 trend change. *Journal of Atmospheric and Oceanic Technology* 25: 368–384, DOI:  
22 10.1175/2007JTECHA982.1.
- 23 Wahr, J., Molenaar, M., Bryan, F., 1998. Time variability of the Earth's gravity field:  
24 Hydrological and oceanic effects and their possible detection using GRACE. *Journal of*  
25 *Geophysical Research*, 103(B12), 30205–30229. <http://dx.doi.org/10.1029/98JB02844>.
- 26 WMO., 2003. Report of the GCOS/GTOS/HWRP Expert Meeting on Hydrological Data for  
27 Global Studies, WMO/TD – No. 1156
- 28 Zhang, X. B., Vincent, L.A., Hogg, W.D., Niitso, A. 2000. Temperature and precipitation  
29 trends in Canada during the 20th century, *Atmos. Ocean*, 38, 395–429.
- 30 Zhang, X., Aguilar, E., Sensoy, S., Melkonyan, H., Tagiyeva, U., Ahmed, N., Kutaladze, N.,  
31 Rahimzadeh, F., Taghipour, A., Hantosh, T. H., Albert, P., Semawi, M., Ali, M.K., Al-  
32 Shabibi, S.H.M., Al-Oulan, Z., Taha Zatari, T., Al Dean Khelet, I.K., Hamoud, S.,  
33 Ramazan Sagir, R., Demircan, M., Eken, M., Adiguzel, M., Lisa Alexander, L., Thomas C.  
34 Peterson, C.T., Wallis, T., 2005. Trends in Middle East climate extreme indices from 1950  
35 to 2003, *J. Geophys. Res.*, 110, D22104, doi: 10.1029/2005JD006181.
- 36 Zhang, X., Zwiers, F.W., Hegerl, G., 2009. The influences of data precision on the calculation  
37 of temperature percentile indices; *International Journal of Climatology* 29: 321–327, DOI:  
38 10.1002/joc.1738.
- 39


1

## List of Figures




2


3 **Figure 1: Location of the Greater Horn of Africa region in Africa together with stations used**  
4 **in the study (see Table 1 for full names, the values of latitude and longitude for each station).**




**Figure 2: Available data for the time series used in the analysis. White indicates no data; upper box indicate rainfall while the lower one indicates temperature. The arrows show the common years available for all the countries.**




(a) Asmara TX10P



(b) Asmara TX90P




(c) Asmara TN10P




(d) Asmara TN90P

**Figure 3: 1971–2010 time series for (a) cold days, (b) warm days, (c) cold nights and (d) warm nights (units: %). Bold line indicates ordinary least squares fit for Asmara, Eritrea.**




(a) Asmara




(b) Regional trend

Figure 4: Precipitation Total (PRCPTOT). Individual station's time series and regional trend (a) Individual time series 1980–2010 for Asmara in Eritrea, (b) Regionally averaged station trends. Positive (negative) trends are shown in red (blue) circles. Large (small) circles indicate significant (insignificant) trends.




(a) Khartoum R95p




(b) Khartoum RX5day

Figure 5: 1961–2000 time series for (a) contribution from very wet days (R95p, units: mm) and (b) annual maximum 5-day precipitation amounts (RX5day, units: mm) for Khartoum. Bold lines indicate ordinary least squares fit.




(a) Dodoma R95p




(b) Dodoma RX5day

Figure 6: 1970 –2010 time series for (a) contribution from very wet days (R95p, units: mm) and (b) annual maximum 5-day precipitation amounts (RX5day, units: mm) for Dodoma. Bold lines indicate ordinary least squares fit.







**Figure 7a: (top) Spatially-averaged total water storage (EWT) variations over the GHA, derived from daily Kalman-smoother GRACE products, (bottom) Accumulated EWT changes over the GHA**




**Figure 7b: Implementing the temporal ICA method over 2588 days of TWS maps over the GHA, computed from Kalman-smoother daily GRACE products provided by the APMG group at Bonn University. Independent patterns are ordered according to the variance they represent. One can reconstruct each mode of TWS variability by multiplying the spatial patterns with their corresponding temporal components.**




(a) Southern sector



(b) Northern Sector




**(c) Equatorial sector**




**(d) Mean surface temperature**

**Figure 8: Simulated and observed mean climate cycles for some stations located in various sectors of the region.**



(a) Gulu



(b) Projected regional trend

**Figure 9: Projected Precipitation Total (PRCPTOT). Individual station time series and regional trend (a) Individual time series 2010–2040 for Gulu in Uganda, (b) Regionally averaged station trends. Positive (negative) trends are shown in red (blue) circles. Large (small) circles indicate significant (nonsignificant) trends.**

**Table 1. List of stations (see Fig. 1 for their exact location).**

NO	STATION	STN	CODE	LON	LAT	PERIOD
1	ABU HAMAD	ABH	62640	33.3	19.5	1960-2002
2	ARBA MINCH	ABM	63500	37.7	6.1	1986-2007
3	ASMARA	ASM	63021	38.9	15.3	1960-2009
4	ATBARA	ATB	62680	34.0	17.7	1960-2000
5	AWASSA	AWA	63460	38.5	7.1	1971-2007
6	BUJUMBURA	BUJ	64390	29.3	-3.3	1970-2010
7	BUKOBA	BUK	63729	31.8	-1.3	1960-1991
8	GIKONGORO	GIK		30.1	-1.6	1967-2010
9	GISENYI	GNY		29.3	-1.7	1971-2010
10	KAMEMBE	KMBE		28.9	-2.5	1971-2010
11	METE	METE		39.9	8.87	1983-2007
12	NAAMA	NAA		34.08	12.44	1963-2000
13	NAHOUD	NHD		28.26	12.42	1960-2000
14	OBIED	OBE		30.14	13.1	1960-2000
15	ORE	ORE		35.53	8.15	1951-2007
16	DAGORETTI	DAG	63741	36.8	-1.3	1960-1991
17	DAR.I.AIRP.	DIA	63894	39.2	-6.9	1971-2009
18	DEBRE MARCOS	DMKOS	63334	37.7	10.4	1970-2003
19	DIRE DAWA	DDAWA	63471	41.9	9.6	1970-2003
20	DJIBOUTI	DJB	63125	43.2	11.6	1979-2010
21	DODOMA	DOD	63862	35.8	-6.2	1971-2009
22	GENEINA	GEN	62770	22.5	13.5	1960-2001
23	GONDAR	GON	63331	37.4	12.5	1970-2003
24	GORE	GOR	63403	35.6	8.2	1970-2003
25	JINJA	JIN	63682	33.2	0.5	1970-2003
26	JUBA	JUB	62941	31.4	4.5	1960-2000
27	KHARTOUM	KHA	62721	32.6	15.6	1960-2000
28	KIGALI	KIG	64387	30.1	-2.0	1971-2010
29	KOSTI	KOS	62772	32.4	13.1	1960-2000
30	LODWAR	LOD	63612	35.6	3.1	1960-1991
31	MAKINDU	MAK	63766	37.8	-2.3	1960-1991
32	MALAKAL	MAL	62840	31.7	9.6	1960-2000
33	MASINDI	MAS	63654	31.7	1.7	1960-1991
34	MBEYA	MBE	63932	33.5	-8.9	1970-2009
35	MWANZA	MWA	63756	32.9	-2.5	1960-1991
36	NDONGOLA	NDO	62650	30.5	19.2	1960-2000
37	NEKEMTE	NEK	63340	36.6	9.1	1952-2007
38	NYALA	NYA	62790	24.9	12.1	1960-2000
39	ROBE	ROB	63474	40.0	7.1	1983-2007
40	*SOMALIA	SOM				

\* Missing Data

**Table 2. List of the ETCCDI indices used in this study**

<b>ID</b>	<b>Indicator Name</b>	<b>Description</b>	<b>Units</b>
<b>FD *</b>	frost days	count of days where TN (daily minimum temperature) < 0°C	Days
<b>SU</b>	summer days	count of days where TX (daily maximum temperature) > 25°C	Days
<b>ID</b>	icing days	count of days where TX < 0°C	Days
<b>TR</b>	tropical nights	count of days where TN > 20°C	Days
<b>GSL *</b>	growing season length	annual count of days between first span of at least six days where TG (daily mean temperature) > 5°C and first span in second half of the year of at least six days where TG < 5°C.	Days
<b>TXx</b>		monthly maximum value of daily maximum temperature	°C
<b>TNx</b>		monthly maximum value of daily minimum temperature	°C
<b>TXn</b>		monthly minimum value of daily maximum temperature	°C
<b>TNn</b>		monthly minimum value of daily minimum temperature	°C
<b>TN10p</b>	cold nights	count of days where TN < 10th percentile	%
<b>TX10p</b>	cold day-times	count of days where TX < 10th percentile	%
<b>TN90p*</b>	warm nights	count of days where TN > 90th percentile	%
<b>TX90p</b>	warm day-times	count of days where TX > 90th percentile	%
<b>WSDI *</b>	warm spell duration index	count of days in a span of at least six days where TX > 90th percentile	%
<b>CSDI</b>	cold spell duration index	count of days in a span of at least six days where TN > 10th percentile	Days
<b>DTR</b>	diurnal temperature range	mean difference between TX and TN (°C)	
<b>RX1day</b>	maximum one-day precipitation	highest precipitation amount in one-day period	mm
<b>RX5day *</b>	maximum 5-day precipitation	highest precipitation amount in five-day period	mm
<b>SDII *</b>	simple daily intensity index	mean precipitation amount on a wet day	mm
<b>R10mm *</b>	heavy precipitation days	count of days where RR (daily precipitation amount) ≥ 10 mm	Days
<b>R20mm</b>	very heavy precipitation days	count of days where RR ≥ 20 mm	Days
<b>R<sub>mm</sub>mm</b>		count of days where RR ≥ user-defined threshold in mm	Days
<b>CDD *</b>	consecutive dry days	maximum length of dry spell (RR < 1 mm)	Days
<b>CWD</b>	consecutive wet days	maximum length of wet spell (RR ≥ 1 mm)	Days
<b>R95p TOT*</b>		precipitation due to very wet days (> 95th percentile)	mm
<b>R99pTOT</b>		precipitation due to extremely wet days (> 99th percentile)	mm
<b>PRCPTOT</b>		total precipitation in wet days (> 1 mm)	mm

Full definitions are available from the ETCCDI website <http://cccma.seos.uvic.ca/ETCCDI/>

**Table 3: Regional Trends in Temperature Indices<sup>a</sup>**

Index	Guinea	Central Africa	Zimbabwe	Global	Kenya	Ethiopia	Units
Warmest day	<b>0.14</b>	<b>0.25</b>	0.15	<b>0.21</b>	<b>0.35</b>	0.11	° C /decade
Warmest night	<b>0.17</b>	<b>0.21</b>	0.10	<b>0.30</b>	0.17	<b>0.33</b>	° C /decade
Coldest day	<b>0.23</b>	<b>0.13</b>	0.00	0.37	0.02	0.10	° C /decade
Coldest night	0.04	<b>0.23</b>	0.02	<b>0.71</b>	<b>0.21</b>	<b>0.32</b>	° C /decade
DTR	0.12	0.00	0.11	-0.08	0.22	0.61	° C /decade
Cold Night frequency	-0.21	<b>-1.71</b>	<b>-1.24</b>	<b>-1.26</b>	<b>-1.10</b>	<b>-1.23</b>	% of days in a year/Decade
Cold Day frequency	<b>-2.15</b>	<b>-1.22</b>	<b>-1.05</b>	<b>-0.62</b>	<b>-1.6</b>	<b>-1.0</b>	% of days in a year/Decade
Warm night frequency	<b>1.19</b>	<b>3.24</b>	0.71	<b>1.58</b>	<b>1.44</b>	<b>2.14</b>	% of days in a year/Decade
Warm day frequency	<b>1.56</b>	<b>2.87</b>	<b>1.86</b>	<b>0.89</b>	<b>1.07</b>	0.65	% of days in a year/Decade

<sup>a</sup>The trends for the globe area from Alexander et al. (2006) and Caesar et al. (2010) based on the time period 1955-2003. A trend significant at the 5% level is marked with bold font.

**Table 4: Regional and global trends in precipitation Indices for the period 1971-2005**

Index	Indian Ocean	Himalayas	Indo-Pacific	Global	Northern sector	Equatorial sector	Southern sector	Units
PRCPTOT	<b>81.84</b>	<b>41.77</b>	-2.86	5.91	-2.92	-0.85	10.3	mm /decade
SDII	<b>1.05</b>	<b>1.55</b>	<b>0.25</b>	0.05	-0.81	-0.89	-0.13	mm/day/decade
CDD	0.66	2.61	-1.01	<b>-1.19</b>	0.37	0.32	0.45	Days /decade
CWD	0.10	-0.24	-0.13	<b>-0.07</b>	-0.05	-0.50	-0.07	Days /decade
RX1day	1.12	1.70	-1.12	0.26	0.48	-0.33	-0.72	mm /decade
RX5day	5.96	16.39	0.90	<b>0.73</b>	0.67	-0.67	0.12	mm /decade
R10mm	2.09	0.00	-0.14	<b>0.03</b>	-0.29	-0.28	0.35	Days /decade
R20mm	1.26	0.53	-0.04	0.06	-0.01	-0.04	0.14	Days /decade
R95p	22.66	<b>82.30</b>	12.24	<b>4.68</b>	12.9	-0.50	1.79	mm/Decade
R99p	-12.61	32.39	4.98	<b>3.38</b>	51.1	-2.30	8.05	mm/Decade

Note that global trends were calculated from Alexander et al. (2006) together with Caesar et al. (2010) data and referred to the period 1971 to 2003. Trends significant at the 5% level are shown in bold font.