

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/100905/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Pronzato, Luc, Wynn, Henry P. and Zhigljavsky, Anatoly 2017. Extended generalised variances, with applications. *Bernoulli* 23 (4A) , pp. 2617-2642. 10.3150/16-BEJ821

Publishers page: <http://dx.doi.org/10.3150/16-BEJ821>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Extended generalised variances, with applications

LUC PRONZATO¹, HENRY P. WYNN² and ANATOLY A. ZHIGLJAVSKY³

¹Laboratoire I3S, UMR 7172, UNS, CNRS; 2000, route des Lucioles, Les Algorithmes, bât. Euclide B, 06900 Sophia Antipolis, France. E-mail: pronzato@i3s.unice.fr

²London School of Economics, Houghton Street, London, WC2A 2AE, UK. E-mail: H.Wynn@lse.ac.uk

³School of Mathematics, Cardiff University, Senghennydd Road, Cardiff, CF24 4YH, UK.
E-mail: ZhigljavskyAA@cf.ac.uk

We consider a measure ψ_k of dispersion which extends the notion of Wilk's generalised variance for a d -dimensional distribution, and is based on the mean squared volume of simplices of dimension $k \leq d$ formed by $k + 1$ independent copies. We show how ψ_k can be expressed in terms of the eigenvalues of the covariance matrix of the distribution, also when a n -point sample is used for its estimation, and prove its concavity when raised at a suitable power. Some properties of dispersion-maximising distributions are derived, including a necessary and sufficient condition for optimality. Finally, we show how this measure of dispersion can be used for the design of optimal experiments, with equivalence to A and D -optimal design for $k = 1$ and $k = d$, respectively. Simple illustrative examples are presented.

Keywords: design of experiments; dispersion; generalised variance; maximum-dispersion measure; optimal design; quadratic entropy

1. Introduction

The idea of dispersion is fundamental to statistics and with different terminology, such as potential, diversity, entropy, information and capacity, stretches over a wide area. The variance and standard deviation are the most prevalent for a univariate distribution, and Wilks generalised variance is the term usually reserved for the determinant of the covariance matrix, V , of a multivariate distribution. Many other measures of dispersion have been introduced and a rich area comprises those that are order-preserving with respect to a dispersion ordering; see [5,13,24]. These are sometimes referred to as *measures of peakness* and *peakness ordering*, and are related to the large literature on dispersion measures which grew out of the Gini coefficient, used to measure income inequality [4] and diversity in biology, see [17], which we will discuss briefly below.

In the definitions, there are typically two kinds of dispersion, those measuring some kind of mean distance, or squared distance, from a central value, such as in the usual definition of variance, and those based on the expected distance, or squared distance, between two independent copies from the same distribution, such as the Gini coefficient. It is this second type that will concern us here and we will generalise the idea in several ways by replacing distance by volumes of simplices formed by k independent copies and by transforming the distance, both inside the expectation and outside. This use of volumes makes our measures of dispersion sensitive to the dimension of the subspace where the bulk of the data lives in.

The area of optimal experimental design is another which has provided a range of dispersion measures. Good designs, it is suggested, are those whose parameter estimates have low dispersion. Typically, this means that the design measure, the spread of the observation sites, *maximises* a measure of dispersion and we shall study this problem.

We think of a dispersion measure as a functional directly on the distribution. The basic functional is an integral, such as a moment. The property we shall stress for such functionals most is concavity: that a functional does not decrease under mixing of the distributions. A fundamental theorem in Bayesian learning is that we expect concave functionals to decrease through taking of observations, see Section 2.2 below.

Our central result (Section 3) is an identity for the mean squared volume of simplices of dimension k , formed by $k + 1$ independent copies, in terms of the eigenvalues of the covariance matrices or equivalently in terms of sums of the determinants of k -marginal covariance matrices. Second, we note that after an appropriate (exterior) power transformation the functional becomes concave. We can thus (i) derive properties of measures that maximise this functional (Section 4.1), (ii) use this functional to measure the dispersion of parameter estimates in regression problems, and hence design optimal experiments which minimise this measure of dispersion (Section 4.2).

2. Dispersion measures

2.1. Concave and homogeneous functionals

Let \mathcal{X} be a compact subset of \mathbb{R}^d , \mathcal{M} be the set of all probability measures on the Borel subsets of \mathcal{X} and $\phi : \mathcal{M} \rightarrow \mathbb{R}^+$ be a functional defined on \mathcal{M} . We will be interested in the functionals $\phi(\cdot)$ that are (see [Appendix](#) for precise definitions):

- (a) shift-invariant,
- (b) positively homogeneous of a given degree q , and
- (c) concave: $\phi[(1 - \alpha)\mu_1 + \alpha\mu_2] \geq (1 - \alpha)\phi(\mu_1) + \alpha\phi(\mu_2)$ for any $\alpha \in (0, 1)$ and any two measures μ_1, μ_2 in \mathcal{M} .

For $d = 1$, a common example of a functional satisfying the above properties, with $q = 2$ in (b), is the variance

$$\sigma^2(\mu) = E_\mu^{(2)} - E_\mu^2 = \frac{1}{2} \iint (x_1 - x_2)^2 \mu(dx_1)\mu(dx_2),$$

where $E_\mu = \mathbf{E}(x) = \int x \mu(dx)$ and $E_\mu^{(2)} = \int x^2 \mu(dx)$. Concavity follows from linearity of $E_\mu^{(2)}$, that is, $E_{(1-\alpha)\mu_1 + \alpha\mu_2}^{(2)} = (1 - \alpha)E_{\mu_1}^{(2)} + \alpha E_{\mu_2}^{(2)}$, and Jensen's inequality which implies $E_{(1-\alpha)\mu_1 + \alpha\mu_2}^2 \leq (1 - \alpha)E_{\mu_1}^2 + \alpha E_{\mu_2}^2$.

Any moment of $\mu \in \mathcal{M}$ is a homogeneous functional of a suitable degree. However, the variance is the only moment which satisfies (a) and (c). Indeed, the shift-invariance implies that the moment should be central, but the variance is the only concave functional among the central

moments, see [Appendix](#). In this sense, one of the aims of this paper is a generalisation of the concept of variance.

In the general case $d \geq 1$, the double variance $2\sigma^2(\mu)$ generalises to

$$\phi(\mu) = \iint \|x_1 - x_2\|^2 \mu(dx_1)\mu(dx_2) = 2 \int \|x - E_\mu\|^2 \mu(dx) = 2 \text{trace}(V_\mu), \tag{2.1}$$

where $\|\cdot\|$ is the L_2 -norm in \mathbb{R}^d and V_μ is the covariance matrix of μ . This functional, like the variance, satisfies conditions (a)–(c) with $q = 2$.

The functional (2.1) is the double integral of the squared distance between two random points distributed according to the measure μ . Our main interest will be concentrated around the general class of functionals defined by

$$\phi(\mu) = \phi_{[k],\delta,\tau}(\mu) = \left(\int \cdots \int \mathcal{V}_k^\delta(x_1, \dots, x_{k+1}) \mu(dx_1) \cdots \mu(dx_{k+1}) \right)^\tau, \quad k \geq 2 \tag{2.2}$$

for some δ and τ in \mathbb{R}^+ , where $\mathcal{V}_k(x_1, \dots, x_{k+1})$ is the volume of the k -dimensional simplex (its area when $k = 2$) formed by the $k + 1$ vertices x_1, \dots, x_{k+1} in \mathbb{R}^d , with $k = d$ as a special case. Property (a) for the functionals (2.2) is then a straightforward consequence of the shift-invariance of \mathcal{V}_k , and positive homogeneity of degree $q = k\delta\tau$ directly follows from the positive homogeneity of \mathcal{V}_k with degree k . Concavity will be proved to hold for $\delta = 2$ and $\tau \leq 1/k$ in Section 3. There, we show that this case can be considered as a natural extension of (2.1) (which corresponds to $k = 1$), with $\phi_{[k],2,\tau}(\mu)$ being expressed as a function of V_μ , the covariance matrix of μ . The concavity for $k = \tau = 1$ and all $0 < \delta \leq 2$, follows from the fact that $B(\lambda) = \lambda^\alpha$, $0 < \alpha \leq 1$, is a Bernstein function, which will be discussed briefly below. The functionals (2.2) with $\delta = 2$ and $\tau > 0$, $1 \leq k \leq d$, can be used to define a family of criteria for optimal experimental design, concave for $\tau \leq 1/k$, for which an equivalence theorem can be formulated.

2.2. Quadratic entropy and learning

In a series of papers [17–20], C.R. Rao and coworkers have introduced a quadratic entropy which is a generalised version of the $k = 2$ functional of this section but with a general kernel $K(x_1, x_2)$ in \mathbb{R}^d :

$$Q_R = \iint K(x_1, x_2) \mu(dx_1)\mu(dx_2). \tag{2.3}$$

For the discrete version

$$Q_R = \sum_{i=1}^N \sum_{j=1}^N K(x_i, x_j) p_i p_j,$$

Rao and coworkers developed a version of the Analysis of Variance (ANOVA), which they called Analysis of Quadratic Entropy (ANOQE), or Analysis of Diversity (ANODIV). The Gini coefficient, also used in the continuous and discrete form is a special case with $d = 1$ and $K(x_1, x_2) = |x_1 - x_2|$.

As pointed in [19], Chapter 3, a necessary and sufficient condition for the functional Q_R to be concave is

$$\iint K(x_1, x_2) \nu(dx_1) \nu(dx_2) \leq 0 \tag{2.4}$$

for all measures ν with $\int \nu(dx) = 0$. The discrete version of this is

$$\sum_{i=1}^N \sum_{j=1}^N K(x_i, x_j) q_i q_j \leq 0$$

for any choice of real numbers q_1, \dots, q_N such that $\sum_{i=1}^N q_i = 0$. Schilling, Song and Vondraček [22] discuss the general problem of finding for what class of continuous functions $B(\cdot)$ of $\|x_1 - x_2\|^2$ does the kernel $K(x_1, x_2) = B(\|x_1 - x_2\|^2)$ satisfy (2.4): the solution is that $B(\cdot)$ must be a so-called Bernstein function. We do not develop these ideas here, but note that $B(\lambda) = \lambda^\alpha$ is a Bernstein function for all $0 < \alpha \leq 1$. This is the reason that, above, we can claim concavity for $k = 1$ and all $0 < \delta \leq 2$ in (2.2).

Hainy, Müller and Wynn [6] discuss the link to embedding and review some basic results related to Bayesian learning. One asks what is the class of functionals ψ on a distribution $\mu(\theta)$ of a parameter in the Bayesian statistical learning such that for all $\mu(\theta)$ and all sampling distributions $\pi(x|\theta)$ one expects to learn, in the preposterior sense: $\psi(\mu(\theta)) \leq E_\nu \psi(\pi(\theta|X))$, with $X \sim \nu$. The condition is that ψ is convex, a result which has a history but is usually attributed to De-Groot [2]. This learning is enough to justify calling such a functional a generalised information functional, or a general learning functional. Shannon information falls in this class, and earlier versions of the result were for Shannon information. It follows that wherever, in this paper, we have a concave functional then its negative is a learning functional.

3. Functionals based on squared volume

In the rest of the paper, we focus our attention on the functional

$$\mu \in \mathcal{M} \quad \longrightarrow \quad \psi_k(\mu) = \phi_{[k],2,1}(\mu) = E\{\mathcal{V}_k^2(x_1, \dots, x_{k+1})\},$$

which corresponds to the mean squared volume of simplices of dimension k formed by $k + 1$ independent samples from μ . For instance,

$$\psi_2(\mu) = \iiint \mathcal{V}_2^2(x_1, x_2, x_3) \mu(dx_1) \mu(dx_2) \mu(dx_3), \tag{3.1}$$

with $\mathcal{V}_2(x_1, x_2, x_3)$ the area of the triangle formed by the three points with coordinates x_1, x_2 and x_3 in \mathbb{R}^d , $d \geq 2$. Functionals $\phi_{[k],\delta,\tau}(\mu)$ for $\delta \neq 2$ will be considered in another paper, including the case of negative δ and τ in connection with space-filling design for computer experiments.

Theorem 3.1 of Section 3.1 indicates how $\psi_k(\mu)$ can be expressed as a function of V_μ , the covariance matrix of μ , and shows that $\phi_{[k],2,1/k}(\cdot)$ satisfies properties (a), (b) and (c) of Section 2.1. The special case of $k = d$ was known to Wilks [28,29] in his introduction of generalised

variance, see also [27]. The connection with U-statistics is exploited in Section 3.3, where an unbiased minimum-variance estimator of $\psi_k(\mu)$ based on a sample x_1, \dots, x_n is expressed in terms of the empirical covariance matrix of the sample.

3.1. Expected squared k -simplex volume

Theorem 3.1. *Let the x_i be i.i.d. with the probability measure $\mu \in \mathcal{M}$. Then, for any $k \in \{1, \dots, d\}$, we have*

$$\psi_k(\mu) = \frac{k+1}{k!} \sum_{i_1 < i_2 < \dots < i_k} \det[\{V_\mu\}_{(i_1, \dots, i_k) \times (i_1, \dots, i_k)}] \tag{3.2}$$

$$= \frac{k+1}{k!} \sum_{i_1 < i_2 < \dots < i_k} \lambda_{i_1}[V_\mu] \times \dots \times \lambda_{i_k}[V_\mu], \tag{3.3}$$

where the $\lambda_i[V_\mu]$ denote the eigenvalues of the covariance matrix V_μ and all i_j belong to $\{1, \dots, d\}$. Moreover, the functional $\psi_k^{1/k}(\cdot)$ is shift-invariant, homogeneous of degree 2 and concave on \mathcal{M} .

The proof uses the following two lemmas, see [Appendix](#).

Lemma 3.1. *Let the $k+1$ vectors x_1, \dots, x_{k+1} of \mathbb{R}^k be i.i.d. with the probability measure μ , $k \geq 2$. For $i = 1, \dots, k+1$, denote $z_i = (x_i^\top 1)^\top$. Then*

$$\mathbb{E} \left\{ \det \left[\sum_{i=1}^{k+1} z_i z_i^\top \right] \right\} = (k+1)! \det[V_\mu].$$

Lemma 3.2. *The matrix functional $\mu \mapsto V_\mu$ is Loewner-concave on \mathcal{M} , in the sense that, for any μ_1, μ_2 in \mathcal{M} and any $\alpha \in (0, 1)$,*

$$V_{(1-\alpha)\mu_1 + \alpha\mu_2} \succeq (1-\alpha)V_{\mu_1} + \alpha V_{\mu_2}, \tag{3.4}$$

where $A \succeq B$ means that $A - B$ is nonnegative definite.

Proof of Theorem 3.1. When $k = 1$, the results follow from $\psi_1(\mu) = 2 \text{trace}(V_\mu)$, see (2.1). Using Binet–Cauchy formula, see, for example, [3], vol. 1, page 9, we obtain

$$\begin{aligned} & \mathcal{V}_k^2(x_1, \dots, x_{k+1}) \\ &= \frac{1}{(k!)^2} \det \left(\begin{array}{c} \left[\begin{array}{c} (x_2 - x_1)^\top \\ (x_3 - x_1)^\top \\ \vdots \\ (x_{k+1} - x_1)^\top \end{array} \right] \\ \left[(x_2 - x_1)(x_3 - x_1) \cdots (x_{k+1} - x_1) \right] \end{array} \right) \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{(k!)^2} \sum_{i_1 < i_2 < \dots < i_k} \det^2 \begin{bmatrix} \{x_2 - x_1\}_{i_1} & \dots & \{x_{k+1} - x_1\}_{i_1} \\ \vdots & \vdots & \vdots \\ \{x_2 - x_1\}_{i_k} & \dots & \{x_{k+1} - x_1\}_{i_k} \end{bmatrix} \\
 &= \frac{1}{(k!)^2} \sum_{i_1 < i_2 < \dots < i_k} \det^2 \begin{bmatrix} \{x_1\}_{i_1} & \dots & \{x_{k+1}\}_{i_1} \\ \vdots & \vdots & \vdots \\ \{x_1\}_{i_k} & \dots & \{x_{k+1}\}_{i_k} \\ 1 & \dots & 1 \end{bmatrix},
 \end{aligned}$$

where $\{x\}_i$ denotes the i th component of vector x . Also, for all $i_1 < i_2 < \dots < i_k$,

$$\det^2 \begin{bmatrix} \{x_1\}_{i_1} & \dots & \{x_{k+1}\}_{i_1} \\ \vdots & \vdots & \vdots \\ \{x_1\}_{i_k} & \dots & \{x_{k+1}\}_{i_k} \\ 1 & \dots & 1 \end{bmatrix} = \det \left(\sum_{j=1}^{k+1} z_j z_j^\top \right),$$

where we have denoted by z_j the $k + 1$ -dimensional vector with components $\{x_j\}_{i_\ell}$, $\ell = 1, \dots, k$, and 1. When the x_i are i.i.d. with the probability measure μ , using Lemma 3.1 we obtain (3.2), (3.3). Therefore

$$\psi_k(\mu) = \Psi_k[V_\mu] = \frac{k + 1}{k!} \mathcal{E}_k \{ \lambda_1[V_\mu], \dots, \lambda_d[V_\mu] \},$$

with $\mathcal{E}_k \{ \lambda_1[V_\mu], \dots, \lambda_d[V_\mu] \}$ the elementary symmetric function of degree k of the d eigenvalues of V_μ , see, for example, [12], page 10. Note that

$$\mathcal{E}_k[V_\mu] = \mathcal{E}_k \{ \lambda_1[V_\mu], \dots, \lambda_d[V_\mu] \} = (-1)^k a_{d-k},$$

with a_{d-k} the coefficient of the monomial of degree $d - k$ of the characteristic polynomial of V_μ ; see, for example, [12], page 21. We have in particular $\mathcal{E}_1[V_\mu] = \text{trace}[V_\mu]$ and $\mathcal{E}_d[V_\mu] = \det[V_\mu]$. The shift-invariance and homogeneity of degree 2 of $\psi_k^{1/k}(\cdot)$ follow from the shift-invariance and positive homogeneity of \mathcal{Y}_k with degree k . Concavity of $\Psi_k^{1/k}(\cdot)$ follows from [12], page 116 (take $p = k$ in equation (10), with $\mathcal{E}_0 = 1$). From [10], the $\Psi_k^{1/k}(\cdot)$ are also Loewner-increasing, so that from Lemma 3.2, for any μ_1, μ_2 in \mathcal{M} and any $\alpha \in (0, 1)$,

$$\begin{aligned}
 \psi_k^{1/k}[(1 - \alpha)\mu_1 + \alpha\mu_2] &= \Psi_k^{1/k}\{V_{(1-\alpha)\mu_1 + \alpha\mu_2}\} \\
 &\geq \Psi_k^{1/k}[(1 - \alpha)V_{\mu_1} + \alpha V_{\mu_2}] \\
 &\geq (1 - \alpha)\Psi_k^{1/k}[V_{\mu_1}] + \alpha\Psi_k^{1/k}[V_{\mu_2}] \\
 &= (1 - \alpha)\psi_k^{1/k}(\mu_1) + \alpha\psi_k^{1/k}(\mu_2). \quad \square
 \end{aligned}$$

The functionals $\mu \rightarrow \phi_{[k],2,\tau}(\mu) = \psi_k^\tau(\mu)$ are thus concave for $0 < \tau \leq 1/k$, with $\tau = 1/k$ yielding positive homogeneity of degree 2. The functional $\psi_1(\cdot)$ is a quadratic entropy \mathcal{Q}_R , see (2.3), or diversity measure [20]; $\psi_d(\mu)$ is proportional to Wilks generalised variance. Functionals $\psi_2^{1/2}(\cdot)$, see (3.1), and more generally $\psi_k^{1/k}(\cdot)$ for $k \geq 2$, can also be considered as diversity measures.

From the well-known expression of the coefficients of the characteristic polynomial of a matrix V , we have

$$\begin{aligned} \Psi_k(V) &= \frac{k+1}{k!} \mathcal{E}_k(V) \\ &= \frac{k+1}{(k!)^2} \det \begin{bmatrix} \text{trace}(V) & k-1 & 0 & \dots \\ \text{trace}(V^2) & \text{trace}(V) & k-2 & \dots \\ \dots & \dots & \dots & \dots \\ \text{trace}(V^{k-1}) & \text{trace}(V^{k-2}) & \dots & 1 \\ \text{trace}(V^k) & \text{trace}(V^{k-1}) & \dots & \text{trace}(V) \end{bmatrix}, \end{aligned} \tag{3.5}$$

see, for example, [11], page 28, and the $\mathcal{E}_k(V)$ satisfy the recurrence relations (Newton identities):

$$\mathcal{E}_k(V) = \frac{1}{k} \sum_{i=1}^k (-1)^{i-1} \mathcal{E}_{k-i}(V) \mathcal{E}_1(V^i), \tag{3.6}$$

see, for example, [3], vol. 1, page 88 and [10]. Particular forms of $\psi_k(\cdot)$ are

$$\begin{aligned} k = 1 : \quad & \psi_1(\mu) = 2 \text{trace}(V_\mu), \\ k = 2 : \quad & \psi_2(\mu) = \frac{3}{4} [\text{trace}^2(V_\mu) - \text{trace}(V_\mu^2)], \\ k = 3 : \quad & \psi_3(\mu) = \frac{1}{9} [\text{trace}^3(V_\mu) - 3 \text{trace}(V_\mu^2) \text{trace}(V_\mu) + 2 \text{trace}(V_\mu^3)], \\ k = d : \quad & \psi_d(\mu) = \frac{d+1}{d!} \det(V_\mu). \end{aligned}$$

3.2. Other concave homogeneous functionals

From the proof of Theorem 3.1, any Loewner-increasing, concave and homogeneous functional of the covariance matrix V_μ satisfies all properties (a)–(c) of Section 2.1. In particular, consider Kiefer’s Φ_p -class [8], defined by

$$\varphi_p(\mu) = \Phi_p(V_\mu) = \begin{cases} \lambda_{\max}(V_\mu), & \text{for } p = \infty, \\ \left\{ \frac{1}{d} \text{trace}(V_\mu^p) \right\}^{1/p}, & \text{for } p \neq 0, \pm\infty, \\ \det^{1/d}(V_\mu), & \text{for } p = 0, \\ \lambda_{\min}(V_\mu), & \text{for } p = -\infty, \end{cases} \tag{3.7}$$

with the continuous extension $\varphi_p(\mu) = 0$ for $p < 0$ when V_μ is singular. Notice that $\varphi_1(\cdot)$ and $\varphi_0(\cdot)$ respectively coincide with $\psi_1(\cdot)$ and $\psi_d^{1/d}(\cdot)$ (up to a multiplicative scalar).

The functionals $\varphi_p(\cdot)$ are homogeneous of degree 2, and concave for $p \in [-\infty, 1]$, see, for example, [16], Chapter 6. However, by construction, for any $p \leq 0$, $\varphi_p(\mu) = 0$ when μ is concentrated in a q -dimensional subspace of \mathbb{R}^d , for any $q < d$, whereas $\varphi_p(\mu) > 0$ for $p > 0$ and any $q > 0$. The family of functionals (3.7) is therefore unable to detect the true dimensionality of the data. On the other hand, $\psi_k(\mu) = 0$ for all $k > q$ when $\text{rank } V_\mu = q$.

3.3. Empirical version and unbiased estimates

Let x_1, \dots, x_n be a sample of n vectors of \mathbb{R}^d , i.i.d. with the measure μ . This sample can be used to obtain an empirical estimate $(\hat{\psi}_1)_n$ of $\psi_k(\mu)$, through the consideration of the $\binom{n}{k+1}$ k -dimensional simplices that can be constructed with the x_i . Below we show how a much simpler (and still unbiased) estimation of $\psi_k(\mu)$ can be obtained through the empirical variance-covariance matrix of the sample. See also [29,30].

Denote

$$\hat{x}_n = \frac{1}{n} \sum_{i=1}^n x_i,$$

$$\hat{V}_n = \frac{1}{n-1} \sum_{i=1}^n (x_i - \hat{x}_n)(x_i - \hat{x}_n)^\top = \frac{1}{n(n-1)} \sum_{i < j} (x_i - x_j)(x_i - x_j)^\top,$$

respectively the empirical mean and variance-covariance matrix of x_1 . Note that both are unbiased. We thus have

$$(\hat{\psi}_1)_n = \frac{2}{n(n-1)} \sum_{i < j} \|x_i - x_j\|^2 = 2 \text{trace}[\hat{V}_n] = \Psi_1(\hat{V}_n),$$

and the estimator $(\hat{\psi}_1)_n$ is an unbiased estimator of $\psi_1(\mu)$. For $k \geq 1$, consider the empirical estimate

$$(\hat{\psi}_k)_n = \binom{n}{k+1}^{-1} \sum_{j_1 < j_2 < \dots < j_{k+1}} \psi_k^2(x_{j_1}, \dots, x_{j_{k+1}}). \tag{3.8}$$

It satisfies the following.

Theorem 3.2. For x_1, \dots, x_n a sample of n vectors of \mathbb{R}^d , i.i.d. with the measure μ , and for any $k \in \{1, \dots, d\}$, we have

$$(\hat{\psi}_k)_n = \frac{(n-k-1)!(n-1)^k}{(n-1)!} \Psi_k(\hat{V}_n), \tag{3.9}$$

and $(\hat{\psi}_k)_n$ forms an unbiased estimator of $\psi_k(\mu)$ with minimum variance among all unbiased estimators.

This result generalises the main result of [27] to $k \leq d$, see Corollary 2.1 in that paper. The proof is given in Appendix.

Using the notation of Theorem 3.1, since $\mathcal{E}_k(V) = (-1)^k a_{d-k}(V)$, with $a_{d-k}(V)$ the coefficient of the monomial of degree $d - k$ of the characteristic polynomial of V , for a nonsingular V we obtain

$$\mathcal{E}_k(V) = \det(V)\mathcal{E}_{d-k}(V^{-1}), \tag{3.10}$$

see also [10], equation (4.2). Therefore, we also have

$$(\widehat{\psi}_{d-k})_n = \frac{(n-d+k-1)!(n-1)^{d-k}}{(n-1)!} \frac{(d-k+1)k!}{(k+1)(d-k)!} \det(\widehat{V}_n)\Psi_k(\widehat{V}_n^{-1}), \tag{3.11}$$

which forms an unbiased and minimum-variance estimator of $\psi_{d-k}(\mu)$. Note that the estimation of $\psi_k(\mu)$ is much simpler through (3.9) or (3.11) than using the direct construction (3.8).

One may notice that $\Psi_k(\widehat{V}_1)$ is clearly unbiased due to the linearity of $\Psi_1(\cdot)$, but it is remarkable that $\Psi_k(\widehat{V}_n)$ becomes unbiased after a suitable scaling, see (3.9). Since $\Psi_k(\cdot)$ is highly non-linear for $k > 1$, this property would not hold if \widehat{V}_n were replaced by another unbiased estimator of V_μ .

The value of $(\widehat{\psi}_k)_n$ only depends on \widehat{V}_n , with $E\{(\widehat{\psi}_k)_n\} = \psi_k(V_\mu)$, but its variance depends on the distribution itself. Assume $E\{\mathcal{Y}_k^4(x_1, \dots, x_{k+1})\} < \infty$. From [23], Lemma A, page 183, the variance of $(\widehat{\psi}_k)_n$ satisfies

$$\text{var}[(\widehat{\psi}_k)_n] = \frac{(k+1)^2}{n}\omega + O(n^{-2}),$$

where $\omega = \text{var}[h(x)]$, with $h(x) = E\{\mathcal{Y}_k^2(x_1, x_2, \dots, x_{k+1})|x_1 = x\}$. Obviously, $E[h(x)] = \psi_k(\mu)$ and calculations similar to those in the proof of Theorem 3.1 give

$$\begin{aligned} \omega &= \frac{1}{(k!)^2} \sum_{I,J} \det[\{V_\mu\}_{I \times I}] \det[\{V_\mu\}_{J \times J}] \\ &\times [E\{(E_\mu - x)_I^\top \{V_\mu\}_{I \times I}^{-1} (E_\mu - x)_I (E_\mu - x)_J^\top \{V_\mu\}_{J \times J}^{-1} (E_\mu - x)_J\} - k^2], \end{aligned} \tag{3.12}$$

where I and J respectively denote two sets of indices $i_1 < i_2 < \dots < i_k$ and $j_1 < j_2 < \dots < j_k$ in $\{1, \dots, d\}$, the summation being over all possible such sets. Simplifications occur in some particular cases. For instance, when μ is a normal measure, then

$$\begin{aligned} \omega &= \frac{2}{(k!)^2} \sum_{I,J} \det[\{V_\mu\}_{I \times I}] \det[\{V_\mu\}_{J \times J}] \\ &\times \text{trace}[\{V_\mu\}_{J \times J}^{-1} \{V_\mu\}_{J \times I} \{V_\mu\}_{I \times I}^{-1} \{V_\mu\}_{I \times J}]. \end{aligned}$$

If, moreover, V_μ is the diagonal matrix $\text{diag}\{\lambda_1, \dots, \lambda_d\}$, then

$$\omega = \frac{2}{(k!)^2} \sum_{I,J} \beta(I, J) \prod_I \lambda_i \prod_J \lambda_j,$$

with $\beta(I, J)$ denoting the number of coincident indices between I and J (i.e., the size of $I \cap J$). When μ is such that the components of x are i.i.d. with variance σ^2 , then $V_\mu = \sigma^2 I_d$, with I_d the d -dimensional identity matrix, and

$$\begin{aligned} & \mathbb{E}\{(E_\mu - x)_I^\top \{V_\mu\}_{I \times I}^{-1} (E_\mu - x)_I (E_\mu - x)_J^\top \{V_\mu\}_{J \times J}^{-1} (E_\mu - x)_J\} \\ &= \mathbb{E}\left\{\left(\sum_{i \in I} z_i^2\right)\left(\sum_{j \in J} z_j^2\right)\right\}, \end{aligned}$$

where the $z_i = \{x - E_\mu\}_i / \sigma$ are i.i.d. with mean 0 and variance 1. We then obtain

$$\omega = \frac{\sigma^{4k}}{(k!)^2} (\mathbb{E}\{z_i^4\} - 1) \beta_{d,k},$$

where

$$\begin{aligned} \beta_{d,k} &= \sum_{I,J} \beta(I, J) = \sum_{i=1}^k i \binom{d}{i} \binom{d-i}{k-i} \binom{d-i-(k-i)}{k-i} \\ &= \frac{(d-k+1)^2}{d} \binom{d}{k-1}. \end{aligned}$$

Example 1. We generate 1000 independent samples of n points for different measures μ . Figure 1 presents a box-plot of the ratios $(\hat{\psi}_k)_n / \psi_k(\mu)$ for various values of k and $n = 100$ (left), $n = 1000$ (right), when $\mu = \mu_1$ uniform in $[0, 1]^{10}$. Figure 2 presents the same information when $\mu = \mu_2$ which corresponds to the normal distribution $\mathcal{N}(0, I_{10}/12)$ in \mathbb{R}^{10} . Note that $V_{\mu_1} = V_{\mu_2}$ but the dispersions are different in the two figures. The fact that the variance of the




Figure 1. Box-plot of $(\hat{\psi}_k)_n / \psi_k(\mu)$ for different values of k : μ is uniform in $[0, 1]^{10}$, $n = 100$ (left) and $n = 1000$ (right)—1000 repetitions; minimum, median and maximum values are indicated, together with 25% and 75% quantiles.




Figure 2. Same as in Figure 1 but for μ normal $\mathcal{N}(0, I_{10}/12)$.

ratio $(\hat{\psi}_k)_n/\psi_k(\mu)$ increases with k is due to the decrease of $\psi_k(\mu)$, see Figure 3-left. Note that the values of $\psi_k(\mu)$ and empirical mean of $(\hat{\psi}_k)_n$ are extremely close. Figure 3-right presents the asymptotic and empirical variances of $(\hat{\psi}_k)_n/\psi_k(\mu)$ as functions of k .

Other properties of U-statistics apply to the estimator $(\hat{\psi}_k)_n$, including almost-sure consistency and the classical law of the iterated logarithm, see [23], Section 5.4. In particular, $(\hat{\psi}_k)_n$ is asymptotically normal, $\sqrt{n}[(\hat{\psi}_k)_n - \psi_k(\mu)] \xrightarrow{d} \mathcal{N}(0, (k + 1)^2\omega)$ with ω given by (3.12). This is illustrated in Figure 4-left below for μ uniform in $[0, 1]^{10}$, with $n = 1000$ and $k = 3$. The distribution is already reasonably close to normality for small values of n , see Figure 4-right for which $n = 20$.




Figure 3. Left: $\psi_k(\mu)$ (dots and solid line) and empirical mean of $(\hat{\psi}_k)_n$ (triangles and dashed line); right: asymptotic (dots and solid line) and empirical (triangles and dashed line) variances of $(\hat{\psi}_k)_n/\psi_k(\mu)$; μ is normal $\mathcal{N}(0, I_{10}/12)$, $n = 100, 1000$ repetitions.




Figure 4. Dots: empirical distribution of $(\widehat{\psi}_k)_n$ (histogram for 10,000 independent repetitions); solid line: asymptotic normal distribution $\mathcal{N}(\psi_k(\mu), (k+1)^2\omega/n)$; μ is uniform in $[0, 1]^{10}$ and $k = 3$; left: $n = 1000$; right: $n = 20$.

4. Maximum-diversity measures and optimal designs

In this section, we consider two types of optimisation problems on \mathcal{M} related to the functionals $\psi_k(\cdot)$ introduced in Theorem 3.1. First, in Section 4.1, we are interested in the characterisation and construction of maximum-diversity measures; that is, measures $\mu_k^* \in \mathcal{M}$ which maximise $\psi_k(\mu) = \Psi_k(V_\mu)$. The existence of an optimal measure follows from the compactness of \mathcal{X} and continuity of $\mathcal{Y}_k(x_1, \dots, x_{k+1})$ in each x_i , see [1], Theorem 1; the concavity and differentiability of the functional $\psi_k^{1/k}(\cdot)$ allow us to derive a necessary and sufficient condition for optimality.

In Section 4.2, we consider the problem of optimal design of experiments, where the covariance matrix V is the inverse of the information matrix $M(\xi)$ for some regression model.

4.1. Maximum-diversity measures

4.1.1. Necessary and sufficient condition

Since the functionals $\psi_k^{1/k}(\cdot)$ are concave and differentiable, for all $k = 1, \dots, d$, we can easily derive a necessary and sufficient condition for a probability measure μ_k^* on \mathcal{X} to maximise $\psi_k(\mu)$, in the spirit of the celebrated Equivalence Theorem of [9].

Denote by $\nabla_{\Psi_k}[V]$ the gradient of $\Psi_k(\cdot)$ at matrix V (a matrix of the same size as V) and by $F_{\psi_k}(\mu; \nu)$ the directional derivative of $\psi_k(\cdot)$ at μ in the direction ν ;

$$F_{\psi_k}(\mu; \nu) = \lim_{\alpha \rightarrow 0^+} \frac{\psi_k[(1 - \alpha)\mu + \alpha\nu] - \psi_k(\mu)}{\alpha}.$$

From the expression (3.5) of $\Psi_k(V)$, we have

$$\nabla_{\Psi_k}[V] = \frac{k+1}{k!} \nabla_{\mathcal{E}_k}[V],$$

where $\nabla_{\mathcal{E}_k}[V]$ denotes the gradient of $\mathcal{E}_k(\cdot)$ at V , which, using (3.6), can be shown by induction to satisfy

$$\nabla_{\mathcal{E}_k}[V] = \sum_{i=0}^{k-1} (-1)^i \mathcal{E}_{k-i-1}(V) V^i, \tag{4.1}$$

see [10]. We thus obtain in particular

$$\begin{aligned} k = 1 : \quad & \nabla_{\Psi_1}[V] = 2I_d, \\ k = 2 : \quad & \nabla_{\Psi_2}[V] = \frac{3}{2}[\text{trace}(V)I_d - V], \\ k = 3 : \quad & \nabla_{\Psi_3}[V] = \frac{1}{3}[\text{trace}^2(V) - \text{trace}(V^2)]I_d - \frac{2}{3}\text{trace}(V)V + \frac{2}{3}V^2, \\ k = d : \quad & \nabla_{\Psi_d}[V] = \frac{d+1}{d!} \det(V)V^{-1}. \end{aligned}$$

Using the differentiability of $\Psi_k(\cdot)$, direct calculation gives

$$F_{\psi_k}(\mu; \nu) = \text{trace} \left\{ \nabla_{\Psi_k}[V_\mu] \frac{dV_{(1-\alpha)\mu+\alpha\nu}}{d\alpha} \Big|_{\alpha=0} \right\},$$

with

$$\frac{dV_{(1-\alpha)\mu+\alpha\nu}}{d\alpha} \Big|_{\alpha=0} = \int [xx^\top - (E_\mu x^\top + x E_\mu^\top)] \nu(dx) - \int xx^\top \mu(dx) + 2E_\mu E_\mu^\top. \tag{4.2}$$

Notice that $dV_{(1-\alpha)\mu+\alpha\nu}/d\alpha|_{\alpha=0}$ is linear in ν .

Then, from the concavity of $\psi_k^{1/k}(\cdot)$, μ_k^* maximises $\psi_k(\mu)$ with respect to $\mu \in \mathcal{M}$ if and only if $\psi_k(\mu_k^*) > 0$ and $F_{\psi_k}(\mu_k^*; \nu) \leq 0$ for all $\nu \in \mathcal{M}$, that is

$$\text{trace} \left\{ \nabla_{\Psi_k}[V_{\mu_k^*}] \frac{dV_{(1-\alpha)\mu_k^*+\alpha\nu}}{d\alpha} \Big|_{\alpha=0} \right\} \leq 0, \quad \forall \nu \in \mathcal{M}. \tag{4.3}$$

We obtain the following.

Theorem 4.1. *The probability measure μ_k^* such that $\psi_k(\mu_k^*) > 0$ is ψ_k -optimal, that is, maximises $\psi_k(\mu)$ with respect to $\mu \in \mathcal{M}$, $k \in \{1, \dots, d\}$, if and only if*

$$\max_{x \in \mathcal{X}} (x - E_{\mu_k^*})^\top \frac{\nabla_{\Psi_k}[V_{\mu_k^*}]}{\Psi_k(V_{\mu_k^*})} (x - E_{\mu_k^*}) \leq k. \tag{4.4}$$

Moreover,

$$(x - E_{\mu_k^*})^\top \frac{\nabla_{\Psi_k}[V_{\mu_k^*}]}{\Psi_k(V_{\mu_k^*})} (x - E_{\mu_k^*}) = k \tag{4.5}$$

for all x in the support of μ_k^* .

Proof. First, note that the Newton equations (3.6) and the recurrence (4.1) for $\nabla_{\mathcal{E}_k}[\cdot]$ imply that $\text{trace}(V \nabla_{\Psi_k}[V]) = k \Psi_k(V)$ for all $k = 1, \dots, d$.

The condition (4.4) is sufficient. Indeed, suppose that μ_k^* such that $\psi_k(\mu_k^*) > 0$ satisfies (4.4). We obtain

$$\int (x - E_{\mu_k^*})^\top \nabla_{\Psi_k}[V_{\mu_k^*}](x - E_{\mu_k^*}) \nu(dx) \leq \text{trace}\{V_{\mu_k^*} \nabla_{\Psi_k}[V_{\mu_k^*}]\}$$

for any $\nu \in \mathcal{M}$, which gives (4.3) when we use (4.2). The condition is also necessary since (4.3) must be true in particular for δ_x , the delta measure at any $x \in \mathcal{X}$, which gives (4.4). The property (4.5) on the support of μ_k^* follows from the observation that $\int (x - E_{\mu_k^*})^\top \nabla_{\Psi_k}[V_{\mu_k^*}](x - E_{\mu_k^*}) \mu_k^*(dx) = \text{trace}\{V_{\mu_k^*} \nabla_{\Psi_k}[V_{\mu_k^*}]\}$. \square

Note that for $k < d$, the covariance matrix $V_{\mu_k^*}$ of a ψ_k -optimal measure μ_k^* is not necessarily unique and may be singular; see, for example, Examples 2 and 3 in Section 4.1.3. Also, $\psi_k(\mu) > 0$ implies that $\psi_{k-1}(\mu) > 0$, $k = 2, \dots, d$.

Remark 4.1. As a natural extension of the concept of potential in case of order-two interactions ($k = 1$), we call $P_{k,\mu}(x) = \psi_k(\mu, \dots, \mu, \delta_x)$ the potential of μ at x , where

$$\psi_k(\mu_1, \dots, \mu_{k+1}) = \int \dots \int \gamma_k^2(x_1, \dots, x_{k+1}) \mu_1(dx_1) \dots \mu_{k+1}(dx_{k+1}).$$

This yields $F_{\psi_k}(\mu; \nu) = (k + 1)[\psi_k(\mu, \dots, \mu, \nu) - \psi_k(\mu)]$, where μ appears k times in $\psi_k(\mu, \dots, \mu, \nu)$. Therefore, Theorem 4.1 states that μ_k^* with $\psi_k(\mu_k^*) > 0$ is ψ_k -optimal if and only if $\psi_k(\mu_k^*, \dots, \mu_k^*, \nu) \leq \psi_k(\mu_k^*)$ for any $\nu \in \mathcal{M}$, or equivalently $P_{k,\mu_k^*}(x) \leq \psi_k(\mu_k^*)$ for all $x \in \mathcal{X}$.

It can be shown that for any measure $\mu \in \mathcal{M}$, $\min_{x \in \mathcal{X}} P_{k,\mu}(x)$ is reached for $x = E_\mu$, which extends the result of [29] about the minimum property of the internal scatter.

Remark 4.2. Consider Kiefer’s Φ_p -class of orthogonally invariant criteria and their associated functional $\varphi_p(\cdot)$, see (3.7). From a result in [7], if a measure μ_p optimal for some $\varphi_p(\cdot)$ with $p \in (-\infty, 1]$ is such that V_{μ_p} is proportional to the identity matrix I_d , then μ_p is simultaneously optimal for all orthogonally invariant criteria. A measure μ_p having this property is therefore ψ_k -optimal for all $k = 1, \dots, d$.

Remark 4.3. Using (3.10), when V is nonsingular we obtain the property

$$\Psi_k(V) = \frac{(k + 1)(d - k)!}{(d - k + 1)k!} \det(V) \Psi_{d-k}(V^{-1})$$

which implies that maximising $\Psi_k(V)$ is equivalent to maximising $\log \det(V) + \log \Psi_{d-k}(V^{-1})$. Therefore, Theorem 4.1 implies that μ_k^* with nonsingular covariance matrix $V_{\mu_k^*}$ maximises $\psi_k(\mu)$ if and only if

$$\max_{x \in \mathcal{X}} (x - E_{\mu_k^*})^\top \left[V_{\mu_k^*}^{-1} - V_{\mu_k^*}^{-1} \frac{\nabla_{\Psi_{d-k}}[V_{\mu_k^*}^{-1}]}{\Psi_{d-k}(V_{\mu_k^*}^{-1})} V_{\mu_k^*}^{-1} \right] (x - E_{\mu_k^*}) \leq d$$

with equality for x in the support of μ_k^* . When k is large (and $d - k$ is small), one may thus check the optimality of μ_k^* without using the complicated expressions of $\Psi_k(V)$ and $\nabla_{\Psi_k}[V]$.

4.1.2. A duality property

The characterisation of maximum-diversity measures can also be approached from the point of view of duality theory.

When $k = 1$, the determination of a ψ_1 -optimal measure μ_1^* is equivalent to the dual problem of constructing the minimum-volume ball \mathcal{B}_d^* containing \mathcal{X} . If this ball has radius ρ , then $\psi_1(\mu_1^*) = 2\rho^2$, and the support points of μ_1^* are the points of contact between \mathcal{X} and \mathcal{B}_d^* ; see [1], Theorem 6. Moreover, there exists an optimal measure with no more than $d + 1$ points.

The determination of an optimal measure μ_d^* is also dual to a simple geometrical problem: it corresponds to the determination of the minimum-volume ellipsoid \mathcal{E}_d^* containing \mathcal{X} . This is equivalent to a D -optimal design problem in \mathbb{R}^{d+1} for the estimation of $\beta = (\beta_0, \beta_1^\top)^\top$, $\beta_1 \in \mathbb{R}^d$, in the linear regression model with intercept $\beta_0 + \beta_1^\top x$, $x \in \mathcal{X}$, see [26]. Indeed, denote

$$W_\mu = \int_{\mathcal{X}} \begin{pmatrix} 1 & x^\top \end{pmatrix}^\top \begin{pmatrix} 1 & x^\top \end{pmatrix} \mu(dx).$$

Then $\mathcal{E}_{d+1}^* = \{z \in \mathbb{R}^{d+1} : z^\top W_{\mu_d^*}^{-1} z \leq d + 1\}$, with μ_d^* maximising $\det(W_\mu)$, is the minimum-volume ellipsoid centered at the origin and containing the set $\{z \in \mathbb{R}^{d+1} : z = \begin{pmatrix} 1 & x^\top \end{pmatrix}^\top, x \in \mathcal{X}\}$. Moreover, \mathcal{E}_d^* corresponds to the intersection between \mathcal{E}_{d+1}^* and the hyperplane $\{z\}_1 = 1$; see, for example, [25]. This gives $\psi_d(\mu_d^*) = (d + 1)/d! \det(W_{\mu_d^*})$. The support points of μ_d^* are the points of contact between \mathcal{X} and \mathcal{E}_d^* , there exists an optimal measure with no more than $d(d + 3)/2 + 1$ points, see [26].

The property below generalises this duality property to any $k \in \{1, \dots, d\}$.

Theorem 4.2.

$$\max_{\mu \in \mathcal{M}} \Psi_k^{1/k}(V_\mu) = \min_{M, c: \mathcal{X} \subset \mathcal{E}(M, c)} \frac{1}{\phi_k^\infty(M)},$$

where $\mathcal{E}(M, c)$ denotes the ellipsoid $\mathcal{E}(M, c) = \{x \in \mathbb{R}^d : (x - c)^\top M(x - c) \leq 1\}$ and $\phi_k^\infty(M)$ is the polar function

$$\phi_k^\infty(M) = \inf_{V \geq 0: \text{trace}(MV) = 1} \frac{1}{\Psi_k^{1/k}(V)}. \tag{4.6}$$

The proof is given in [Appendix](#). The polar function $\phi_k^\infty(\cdot)$ possesses the properties of what is called an information function in [16], Chapter 5; in particular, it is concave on the set of symmetric non-negative definite matrices. This duality property has the following consequence.

Corollary 4.1. *The determination of a covariance matrix V_k^* that maximises $\Psi_k(V_\mu)$ with respect to $\mu \in \mathcal{M}$ is equivalent to the determination of an ellipsoid $\mathcal{E}(M_k^*, c_k^*)$ containing \mathcal{X} , minimum in the sense that M_k^* maximises $\phi_k^\infty(M)$. The points of contact between $\mathcal{E}(M_k^*, c_k^*)$ and \mathcal{X} form the support of μ_k^* .*

For any $V \geq 0$, denote by $M_*(V)$ the matrix

$$M_*(V) = \frac{\nabla \Psi_k[V]}{k \Psi_k(V)} = \frac{1}{k} \nabla_{\log \Psi_k}[V]. \tag{4.7}$$

Note that $M_*(V) \geq 0$, see [16], Lemma 7.5, and that

$$\text{trace}[VM_*(V)] = 1,$$

see the proof of Theorem 4.1. The matrix $V \geq 0$ maximises $\Psi_k(V)$ under the constraint $\text{trace}(MV) = 1$ for some $M \geq 0$ if and only if $V[M_*(V) - M] = 0$. Therefore, if M is such that there exists $V_* = V_*(M) \geq 0$ such that $M = M_*[V_*(M)]$, then $\phi_k^\infty(M) = \Psi_k^{-1/k}[V_*(M)]$. When $k < d$, the existence of such a V_* is not ensured for all $M \geq 0$, but happens when $M = M_k^*$ which maximises $\phi_k^\infty(M)$ under the constraint $\mathcal{X} \in \mathcal{E}(M, c)$. Moreover, in that case there exists a $\mu_k^* \in \mathcal{M}$ such that $M_k^* = M_*(V_{\mu_k^*})$, and this μ_k^* maximises $\psi_k(\mu)$ with respect to $\mu \in \mathcal{M}$.

Consider in particular the case $k = 1$. Then, $M_*(V) = I_d / \text{trace}(V)$ and $\phi_1^\infty(M) = \lambda_{\min}(M) / 2$. The matrix M_k^* of the optimal ellipsoid $\mathcal{E}(M_k^*, c_k^*)$ is proportional to the identity matrix and $\mathcal{E}(M_k^*, c_k^*)$ is the ball of minimum-volume that encloses \mathcal{X} .

When $k = 2$ and $I_d \geq (d - 1)M / \text{trace}(M)$, direct calculations show that $\phi_2^\infty(M) = \Psi_2^{-1/2}[V_*(M)]$, with

$$V_*(M) = [I_d \text{trace}(M) / (d - 1) - M][\text{trace}^2(M) / (d - 1) - \text{trace}(M^2)]^{-1};$$

the optimal ellipsoid is then such that $\text{trace}^2(M) / (d - 1) - \text{trace}(M^2)$ is maximised.

4.1.3. Examples

Example 2. Take $\mathcal{X} = [0, 1]^d$, $d \geq 1$ and denote by v_i , $i = 1, \dots, 2^d$ the 2^d vertices of \mathcal{X} . Consider $\mu^* = (1/2^d) \sum_{i=1}^{2^d} \delta_{v_i}$, with δ_v the Dirac delta measure at v . Then, $V_{\mu^*} = I_d / 4$ and one can easily check that μ^* is ψ_1 -optimal. Indeed, $E_{\mu^*} = \mathbf{1}_d / 2$, with $\mathbf{1}_d$ the d -dimensional vector of ones, and $\max_{x \in \mathcal{X}} (x - \mathbf{1}_d / 2)^\top (2I_d) (x - \mathbf{1}_d / 2) = d / 2 = \text{trace}\{V_{\mu^*} \nabla_{\Psi_1}[V_{\mu^*}]\}$. From Remark 4.2, the measure μ^* is ψ_k -optimal for all $k = 1, \dots, d$.

Note that the two-point measure $\mu_1^* = (1/2)[\delta_{\mathbf{0}} + \delta_{\mathbf{1}_d}]$ is such that $V_{\mu_1^*} = (\mathbf{1}_d \mathbf{1}_d^\top) / 4$ and $\psi_1(\mu_1^*) = d / 2 = \psi_1(\mu^*)$, and is therefore ψ_1 -optimal too. It is not ψ_k -optimal for $k > 1$, since $\psi_k(\mu_1^*) = 0$, $k > 1$.

Example 3. Take $\mathcal{X} = \mathcal{B}_d(\mathbf{0}, \rho)$, the closed ball of \mathbb{R}^d centered at the origin $\mathbf{0}$ with radius ρ . Let μ_0 be the uniform measure on the sphere $\mathcal{S}_d(\mathbf{0}, \rho)$ (the boundary of $\mathcal{B}_d(\mathbf{0}, \rho)$). Then, V_{μ_0} is proportional to the identity matrix I_d , and $\text{trace}[V_{\mu_0}] = \rho^2$ implies that $V_{\mu_0} = \rho^2 I_d / d$. Take $k = d$. We have $E_{\mu_0} = 0$ and

$$\max_{x \in \mathcal{X}} (x - E_{\mu_0})^\top \nabla_{\Psi_d}[V_{\mu_0}](x - E_{\mu_0}) = \frac{(d + 1)\rho^{2d}}{d^{d-1}d!} = \text{trace}\{V_{\mu_0} \nabla_{\Psi_d}[V_{\mu_0}]\},$$

so that μ_0 is ψ_d -optimal from (4.4).

Let μ_d be the measure that allocates mass $1/(d + 1)$ at each vertex of a d regular simplex having its $d + 1$ vertices on $\mathcal{S}_d(\mathbf{0}, \rho)$, with squared volume $\rho^{2d}(d + 1)^{d+1}/[d^d(d!)^2]$. We also have $V_{\mu_d} = \rho^2 I_d/d$, so that μ_d is ψ_d -optimal too. In view of Remark 4.2, μ_0 and μ_d are ψ_k -optimal for all k in $\{1, \dots, d\}$.

Let now μ_k be the measure that allocates mass $1/(k + 1)$ at each vertex of a k regular simplex \mathcal{P}_k , centered at the origin, with its vertices on $\mathcal{S}_d(\mathbf{0}, \rho)$. The squared volume of \mathcal{P}_k equals $\rho^{2k}(k + 1)^{k+1}/[k^k(k!)^2]$. Without any loss of generality, we can choose the orientation of the space so that V_{μ_k} is diagonal, with its first k diagonal elements equal to ρ^2/k and the other elements equal to zero. Note that $\psi_{k'}(\mu_k) = 0$ for $k' > k$. Direct calculations based on (3.5) give

$$\psi_k(\mu_k) = \frac{k + 1}{k!} \frac{\rho^{2k}}{k^k} \leq \psi_k(\mu_0) = \frac{k + 1}{k!} \binom{d}{k} \frac{\rho^{2k}}{d^k},$$

with equality for $k = 1$ and $k = d$, the inequality being strict otherwise.

4.2. Optimal design in regression models

In this section, we consider the case when $V = M^{-1}(\xi)$, where $M(\xi)$ is the information matrix

$$M(\xi) = \int_{\mathbb{T}} f(t) f^\top(t) \xi(dt)$$

in a regression model $Y_j = \theta^\top f(t_j) + \varepsilon_j$ with parameters $\theta \in \mathbb{R}^d$, for a design measure $\xi \in \Xi$. Here Ξ denotes the set of probability measures on a set \mathbb{T} such that $\{f(t) : t \in \mathbb{T}\}$ is compact, and $M^{-1}(\xi)$ is the (asymptotic) covariance matrix of an estimator $\hat{\theta}$ of θ when the design variables t are distributed according to ξ . The value $\psi_k(\mu)$ of Theorem 3.1 defines a measure of dispersion for $\hat{\theta}$, that depends on ξ through $V_\mu = M^{-1}(\xi)$. The design problem we consider consists in choosing ξ that minimises this dispersion, as measured by $\Psi_k[M^{-1}(\xi)]$, or equivalently that maximises $\Psi_k^{-1}[M^{-1}(\xi)]$.

4.2.1. Properties

It is customary in optimal design theory to maximise a concave and Loewner-increasing function of $M(\xi)$, see [16], Chapter 5, for desirable properties of optimal design criteria. Here we have the following.

Theorem 4.3. *The functions $M \rightarrow \Psi_k^{-1/k}(M^{-1})$, $k = 1, \dots, d$, are Loewner-increasing, concave and differentiable on the set \mathbb{M}^+ of $d \times d$ symmetric positive-definite matrices. The functions $\Psi_k(\cdot)$ are also orthogonally invariant.*

Proof. The property (3.10) yields

$$\Psi_k^{-1/k}(M^{-1}) = \left(\frac{k + 1}{k!}\right)^{-1/k} \frac{\det^{1/k}(M)}{\mathcal{E}_{d-k}^{1/k}(M)} \tag{4.8}$$

which is a concave function of M , see equation (10) of [12], page 116. Since $\Psi_k(\cdot)$ is Loewner-increasing, see [10], the function $M \rightarrow \Psi_k^{-1/k}(M^{-1})$ is Loewner-increasing too. Its orthogonal invariance follows from the fact that it is defined in terms of the eigenvalues of M . \square

Note that Theorems 3.1 and 4.3 imply that the functions $M \rightarrow -\log \Psi_k(M)$ and $M \rightarrow \log \Psi_k(M^{-1})$ are convex for all $k = 1, \dots, d$, a question which was left open in [10], and that $M \rightarrow \Psi_k(M^{-1})$ is convex, see [21].

As a consequence of Theorem 4.3, we can derive a necessary and sufficient condition for a design measure ξ_k^* to maximise $\Psi_k^{-1/k}[M^{-1}(\xi)]$ with respect to $\xi \in \Xi$, for $k = 1, \dots, d$.

Theorem 4.4. *The design measure ξ_k^* such that $M(\xi_k^*) \in \mathbb{M}^+$ maximises $\tilde{\psi}_k(\xi) = \Psi_k^{-1/k}[M^{-1}(\xi)]$ with respect to $\xi \in \Xi$ if and only if*

$$\max_{t \in \mathbb{T}} f^\top(t) M^{-1}(\xi_k^*) \frac{\nabla_{\Psi_k}[M^{-1}(\xi_k^*)]}{\Psi_k[M^{-1}(\xi_k^*)]} M^{-1}(\xi_k^*) f(t) \leq k \tag{4.9}$$

or, equivalently,

$$\max_{t \in \mathbb{T}} \left\{ f^\top(t) M^{-1}(\xi_k^*) f(t) - f^\top(t) \frac{\nabla_{\Psi_{d-k}}[M(\xi_k^*)]}{\Psi_{d-k}[M(\xi_k^*)]} f(t) \right\} \leq k. \tag{4.10}$$

Moreover, there is equality in (4.9) and (4.10) for all t in the support of ξ_k^* .

Proof. From (4.8), the maximisation of $\tilde{\psi}_k(\xi)$ is equivalent to the maximisation of $\mathcal{L}_k(\xi) = \log \tilde{\psi}_k(\xi)$ and $\tilde{\phi}_k(\xi) = \log \det[M(\xi)] - \log \Psi_{d-k}[M(\xi)]$. The proof is similar to that of Theorem 4.1 and is based on the following expressions for the directional derivatives of these two functionals at ξ in the direction $v \in \Xi$,

$$F_{\mathcal{L}_k}(\xi; v) = \text{trace} \left(\frac{1}{k} M^{-1}(\xi) \frac{\nabla_{\Psi_k}[M^{-1}(\xi)]}{\Psi_k[M^{-1}(\xi)]} M^{-1}(\xi) [M(v) - M(\xi)] \right)$$

and

$$F_{\tilde{\phi}_k}(\xi; v) = \text{trace} \left(\left\{ M^{-1}(\xi) - \frac{\nabla_{\Psi_{d-k}}[M(\xi)]}{\Psi_{d-k}[M(\xi)]} \right\} [M(v) - M(\xi)] \right),$$

and on the property $\text{trace}\{M \nabla_{\Psi_j}[M]\} = j \Psi_j(M)$. \square

In particular, consider the following special cases for k (note that $\Psi_0(M) = \mathcal{E}_0(M) = 1$ for any M).

- $k = d$: $\tilde{\phi}_d(\xi) = \log \det[M(\xi)],$
- $k = d - 1$: $\tilde{\phi}_{d-1}(\xi) = \log \det[M(\xi)] - \log \text{trace}[M(\xi)] - \log 2,$
- $k = d - 2$: $\tilde{\phi}_{d-2}(\xi) = \log \det[M(\xi)] - \log \{ \text{trace}^2[M(\xi)] - \text{trace}[M^2(\xi)] \} - \log(3/4).$

The necessary and sufficient condition (4.10) then takes the following form:

$$\begin{aligned}
 k = d : \quad & \max_{t \in \mathbb{T}} f^\top(t) M^{-1}(\xi_k^*) f(t) \leq d, \\
 k = d - 1 : \quad & \max_{t \in \mathbb{T}} \left\{ f^\top(t) M^{-1}(\xi_k^*) f(t) - \frac{f^\top(t) f(t)}{\text{trace}[M(\xi_k^*)]} \right\} \leq d - 1, \\
 k = d - 2 : \quad & \max_{t \in \mathbb{T}} \left\{ f^\top(t) M^{-1}(\xi_k^*) f(t) - 2 \frac{\text{trace}[M(\xi_k^*)] f^\top(t) f(t) - f^\top(t) M(\xi_k^*) f(t)}{\text{trace}^2[M(\xi_k^*)] - \text{trace}[M^2(\xi_k^*)]} \right\} \\
 & \leq d - 2.
 \end{aligned}$$

Also, for $k = 1$ condition (4.9) gives

$$\max_{t \in \mathbb{T}} f^\top(t) \frac{M^{-2}(\xi_1^*)}{\text{trace}[M^{-1}(\xi_1^*)]} f(t) \leq 1$$

(which corresponds to A -optimal design), and for $k = 2$

$$\max_{t \in \mathbb{T}} \frac{\text{trace}[M^{-1}(\xi_2^*)] f^\top(t) M^{-2}(\xi_2^*) f(t) - f^\top(t) M^{-3}(\xi_2^*) f(t)}{\text{trace}^2[M^{-1}(\xi_2^*)] - \text{trace}[M^{-2}(\xi_2^*)]} \leq 1.$$

It is well known that a D -optimal design measure maximising $\tilde{\psi}_d(\xi)$ minimises the (squared) volume of confidence ellipsoids \mathcal{E} , and that an A -optimal measure maximizing $\tilde{\psi}_1(\xi)$ minimises the sum of squared lengths of the principal axes of \mathcal{E} , see, e.g., [15], Lemma 5.1. More generally, as discussed in [21], the criteria $\tilde{\psi}_k(\xi)$ have interpretations in terms of confidence ellipsoids \mathcal{E} : a design measure ξ_k^* that maximises $\tilde{\psi}_k(\xi)$ minimises the sum of the squared volumes of the projections of \mathcal{E} on its principal k -dimensional linear subspaces.

Finally, note that a duality theorem, in the spirit of Theorem 4.2, can be formulated for the maximisation of $\Psi_k^{-1/k}[M^{-1}(\xi)]$; see [16], Theorem 7.12, for the general form of such duality properties in optimal experimental design.

4.2.2. Examples

Example 4. For the linear regression model on $\theta_0 + \theta_1 x$ on $[-1, 1]$, the optimal design for $\tilde{\psi}_k(\cdot)$ with $k = d = 2$ or $k = 1$ is

$$\xi_k^* = \left\{ \begin{array}{cc} -1 & 1 \\ 1/2 & 1/2 \end{array} \right\},$$

where the first line corresponds to support points and the second indicates their respective weights.

Example 5. For linear regression with the quadratic polynomial model $\theta_0 + \theta_1 t + \theta_2 t^2$ on $[-1, 1]$, the optimal designs for $\tilde{\psi}_k(\cdot)$ have the form

$$\xi_k^* = \left\{ \begin{array}{ccc} -1 & 0 & 1 \\ w_k & 1 - 2w_k & w_k \end{array} \right\},$$

Table 1. Efficiencies $\text{Eff}_k(\xi_j^*)$ for $j, k = 1, \dots, d$ in Example 5

	Eff_1	Eff_2	Eff_3
ξ_1^*	1	0.9770	0.9449
ξ_2^*	0.9654	1	0.9886
ξ_3^*	0.8889	0.9848	1

with $w_3 = 1/3$, $w_2 = (\sqrt{33} - 1)/16 \simeq 0.2965352$ and $w_1 = 1/4$. Define the efficiency $\text{Eff}_k(\xi)$ of a design ξ as

$$\text{Eff}_k(\xi) = \frac{\tilde{\psi}_k(\xi)}{\tilde{\psi}_k(\xi_k^*)}.$$

Table 1 gives the efficiencies $\text{Eff}_k(\xi_j^*)$ for $j, k = 1, \dots, d = 3$. The design ξ_2^* , optimal for $\tilde{\psi}_2(\cdot)$, appears to make a good compromise between A -optimality (which corresponds to $\tilde{\psi}_1(\cdot)$) and D -optimality (which corresponds to $\tilde{\psi}_3(\cdot)$).

Example 6. For linear regression with the cubic polynomial model $\theta_0 + \theta_1 t + \theta_2 t^2 + \theta_3 t^3$ on $[-1, 1]$, the optimal designs for $\tilde{\psi}_k(\cdot)$ have the form

$$\xi_k^* = \left\{ \begin{array}{cccc} -1 & -z_k & z_k & 1 \\ w_k & 1/2 - w_k & 1/2 - w_k & w_k \end{array} \right\},$$

where

$$\begin{aligned} z_4 &= 1/\sqrt{5} \simeq 0.4472136, & w_4 &= 0.25, \\ z_3 &\simeq 0.4350486, & w_3 &\simeq 0.2149859, \\ z_2 &\simeq 0.4240013, & w_2 &\simeq 0.1730987, \\ z_1 &= \sqrt{3\sqrt{7} - 6}/3 \simeq 0.4639509, & w_1 &= (4 - \sqrt{7})/9 \simeq 0.1504721, \end{aligned}$$

with z_3 satisfying the equation $2z^6 - 3z^5 - 45z^4 + 6z^3 - 4z^2 - 15z + 3 = 0$ and

$$w_3 = \frac{5z^6 + 5z^4 + 5z^2 + 1 - \sqrt{z^{12} + 2z^{10} + 3z^8 + 60z^6 + 59z^4 + 58z^2 + 73}}{12(z^6 + z^4 + z^2 - 3)},$$

with $z = z_3$. For $k = d - 2 = 2$, the numbers z_2 and w_2 are too difficult to express analytically. Table 2 gives the efficiencies $\text{Eff}_k(\xi_j^*)$ for $j, k = 1, \dots, d$. Here again the design ξ_2^* appears to make a good compromise: it maximises the minimum efficiency $\min_k \text{Eff}_f(\cdot)$ among the designs considered. One may refer to [21] for more examples, including polynomials of degree up to 6.

Table 2. Efficiencies $\text{Eff}_k(\xi_j^*)$ for $j, k = 1, \dots, d$ in Example 6

	Eff ₁	Eff ₂	Eff ₃	Eff ₄
ξ_1^*	1	0.9785	0.9478	0.9166
ξ_2^*	0.9694	1	0.9804	0.9499
ξ_3^*	0.9180	0.9753	1	0.9897
ξ_4^*	0.8527	0.9213	0.9872	1

Appendix

Shift-invariance and positive homogeneity

Denote by \mathcal{M} the set of probability measures defined on the Borel subsets of \mathcal{X} , a compact subset of \mathbb{R}^d . For any $\mu \in \mathcal{M}$, any $\theta \in \mathbb{R}^d$ and any $\lambda \in \mathbb{R}^+$, respectively denote by $T_{-\theta}[\mu]$ and $H_{\lambda^{-1}}[\mu]$ the measures defined by:

$$\text{for any } \mu\text{-measurable } \mathcal{A} \subseteq \mathcal{X}, \quad T_{-\theta}[\mu](\mathcal{A} + \theta) = \mu(\mathcal{A}), \quad H_{\lambda^{-1}}[\mu](\lambda\mathcal{A}) = \mu(\mathcal{A}),$$

where $\mathcal{A} + \theta = \{x + \theta : x \in \mathcal{A}\}$ and $\lambda\mathcal{A} = \{\lambda x : x \in \mathcal{A}\}$. The shift-invariance of $\phi(\cdot)$ then means that $\phi(T_{-\theta}[\mu]) = \phi(\mu)$ for any $\mu \in \mathcal{M}$ and any $\theta \in \mathbb{R}^d$, positive homogeneity of degree q means that $\phi(H_{\lambda^{-1}}[\mu]) = \lambda^q \phi(\mu)$ for any $\mu \in \mathcal{M}$ and any $\lambda \in \mathbb{R}^+$.

The variance is the only concave central moment

For $q \neq 2$, the q th central moment $\Delta_q(\mu) = \int |x - E_\mu|^q \mu(dx)$ is shift-invariant and homogeneous of degree q , but it is not concave on \mathcal{M} . Indeed, consider for instance the two-point probability measures

$$\mu_1 = \left\{ \begin{array}{cc} 0 & 1 \\ 1/2 & 1/2 \end{array} \right\} \quad \text{and} \quad \mu_2 = \left\{ \begin{array}{cc} 0 & 101 \\ w & 1 - w \end{array} \right\},$$

where the first line denotes the support points and the second one their respective weights. Then, for

$$w = 1 - \frac{1}{404} \frac{201^{q-1} - 202q + 405}{201^{q-1} - 101q + 102}$$

one has $\partial^2 \Delta_q[(1 - \alpha)\mu_1 + \alpha\mu_2] / \partial \alpha^2|_{\alpha=0} \geq 0$ for all $q \geq 1.84$, the equality being obtained at $q = 2$ only. Counterexamples are easily constructed for values of q smaller than 1.84.

Proof of Lemma 3.1. We have

$$\mathbb{E} \left\{ \det \left[\sum_{i=1}^{k+1} z_i z_i^\top \right] \right\} = (k+1)! \det \begin{bmatrix} \mathbb{E}(x_1 x_1^\top) & E_\mu \\ E_\mu^\top & 1 \end{bmatrix} = (k+1)! \det[V_\mu],$$

see for instance [14], Theorem 1. □

Proof of Lemma 3.2. Take any vector z of the same dimension as x . Then $z^\top V_\mu z = \text{var}_\mu(z^\top x)$, which is a concave functional of μ , see Section 2.1. This implies that $z^\top V_{(1-\alpha)\mu_1 + \alpha\mu_2} z = \text{var}_{(1-\alpha)\mu_1 + \alpha\mu_2}(z^\top x) \geq (1-\alpha) \text{var}_{\mu_1}(z^\top x) + \alpha \text{var}_{\mu_2}(z^\top x) = (1-\alpha)z^\top V_{\mu_1} z + \alpha z^\top V_{\mu_2} z$, for any μ_1, μ_2 in \mathcal{M} and any $\alpha \in (0, 1)$. Since z is arbitrary, this implies (3.4). □

Proof of Theorem 3.2. The estimate (3.8) forms a U-statistics for the estimation of $\psi_k(\mu)$ and is thus unbiased and has minimum variance, see, for example, [23], Chapter 5. We only need to show that it can be written as (3.9).

We can write

$$\begin{aligned} (\widehat{\psi}_k)_n &= \binom{n}{k+1}^{-1} \\ &\times \sum_{j_1 < j_2 < \dots < j_{k+1}} \frac{1}{(k!)^2} \sum_{i_1 < i_2 < \dots < i_k} \det^2 \begin{bmatrix} \{x_{j_1}\}_{i_1} & \dots & \{x_{j_{k+1}}\}_{i_1} \\ \vdots & & \vdots \\ \{x_{j_1}\}_{i_k} & \dots & \{x_{j_{k+1}}\}_{i_k} \\ 1 & \dots & 1 \end{bmatrix} \\ &= \binom{n}{k+1}^{-1} \frac{1}{(k!)^2} \sum_{i_1 < i_2 < \dots < i_k} \det \left(\sum_{j=1}^n \{z_j\}_{i_1, \dots, i_k} \{z_j\}_{i_1, \dots, i_k}^\top \right), \end{aligned}$$

where we have used Binet–Cauchy formula and where $\{z_j\}_{i_1, \dots, i_k}$ denotes the $k+1$ dimensional vector with components $\{x_j\}_{i_\ell}$, $\ell = 1, \dots, k$, and 1. This gives

$$\begin{aligned} (\widehat{\psi}_k)_n &= \binom{n}{k+1}^{-1} \frac{n^{k+1}}{(k!)^2} \sum_{i_1 < i_2 < \dots < i_k} \det \left(\frac{1}{n} \sum_{j=1}^n \{z_j\}_{i_1, \dots, i_k} \{z_j\}_{i_1, \dots, i_k}^\top \right) \\ &= \binom{n}{k+1}^{-1} \frac{n^{k+1}}{(k!)^2} \\ &\times \sum_{i_1 < i_2 < \dots < i_k} \det \begin{bmatrix} (1/n) \left\{ \sum_{j=1}^n x_j x_j^\top \right\}_{(i_1, \dots, i_k) \times (i_1, \dots, i_k)} & \{\widehat{x}_n\}_{i_1, \dots, i_k} \\ \{\widehat{x}_n\}_{i_1, \dots, i_k}^\top & 1 \end{bmatrix} \\ &= \binom{n}{k+1}^{-1} \frac{n^{k+1}}{(k!)^2} \sum_{i_1 < i_2 < \dots < i_k} \det \left[\frac{n-1}{n} \{\widehat{V}_n\}_{(i_1, \dots, i_k) \times (i_1, \dots, i_k)} \right], \end{aligned}$$

and thus (3.9). □

Proof of Theorem 4.2. (i) The fact that $\max_{\mu \in \mathcal{M}} \Psi_k^{1/k}(V_\mu) \geq \min_{M,c: \mathcal{X} \subset \mathcal{E}(M,c)} 1/\phi_k^\infty(M)$ is a consequence of Theorem 4.1. Indeed, the measure μ_k^* maximises $\Psi_k^{1/k}(V_\mu)$ if and only if

$$(x - E_{\mu_k^*})^\top M_*(V_{\mu_k^*})(x - E_{\mu_k^*}) \leq 1 \quad \text{for all } x \text{ in } \mathcal{X}. \tag{A.1}$$

Denote $M_k^* = M_*(V_{\mu_k^*})$, $c_k^* = E_{\mu_k^*}$, and consider the Lagrangian $L(V, \alpha; M)$ for the maximisation of $(1/k) \log \Psi_k(V)$ with respect to $V \geq 0$ under the constraint $\text{trace}(MV) = 1$: $L(V, \alpha; M) = (1/k) \log \Psi_k(V) - \alpha[\text{trace}(MV) - 1]$. We have

$$\left. \frac{\partial L(V, 1; M_k^*)}{\partial V} \right|_{V=V_{\mu_k^*}} = M_k^* - M_k^* = 0$$

and $\text{trace}(M_k^* V_{\mu_k^*}) = 1$, with $V_{\mu_k^*} \geq 0$. Therefore, $V_{\mu_k^*}$ maximises $\Psi_k(V)$ under the constraint $\text{trace}(M_k^* V) = 1$, and, moreover, $\mathcal{X} \subset \mathcal{E}(M_k^*, c_k^*)$ from (A.1). This implies

$$\begin{aligned} \Psi_k^{1/k}(V_{\mu_k^*}) &= \max_{V \geq 0: \text{trace}(M_k^* V) = 1} \Psi_k^{1/k}(V) \\ &\geq \min_{M,c: \mathcal{X} \subset \mathcal{E}(M,c)} \max_{V \geq 0: \text{trace}(MV) = 1} \Psi_k^{1/k}(V) = \min_{M,c: \mathcal{X} \subset \mathcal{E}(M,c)} \frac{1}{\phi_k^\infty(M)}. \end{aligned}$$

(ii) We prove now that $\min_{M,c: \mathcal{X} \subset \mathcal{E}(M,c)} 1/\phi_k^\infty(M) \geq \max_{\mu \in \mathcal{M}} \Psi_k^{1/k}(V_\mu)$. Note that we do not have an explicit form for $\phi_k^\infty(M)$ and that the infimum in (4.6) can be attained at a singular V , not necessarily unique, so that we cannot differentiate $\phi_k^\infty(M)$. Also note that compared to the developments in [16], Chapter 7, here we consider covariance matrices instead of moment matrices.

Consider the maximisation of $\log \phi_k^\infty(M)$ with respect to M and c such that $\mathcal{X} \subset \mathcal{E}(M, c)$, with Lagrangian

$$L(M, c, \beta) = \log \phi_k^\infty(M) + \sum_{x \in \mathcal{X}} \beta_x [1 - (x - c)^\top M(x - c)], \quad \beta_x \geq 0 \text{ for all } x \text{ in } \mathcal{X}.$$

For the sake of simplicity, we consider here \mathcal{X} to be finite, but β may denote any positive measure on \mathcal{X} otherwise. Denote the optimum by

$$T^* = \max_{M,c: \mathcal{X} \subset \mathcal{E}(M,c)} \log \phi_k^\infty(M).$$

It satisfies

$$T^* = \max_{M,c} \min_{\beta \geq 0} L(M, c, \beta) \leq \min_{\beta \geq 0} \max_{M,c} L(M, c, \beta)$$

and $\max_{M,c} L(M, c, \beta)$ is attained for any c such that

$$Mc = M \sum_{x \in \mathcal{X}} \beta_x x / \left(\sum_{x \in \mathcal{X}} \beta_x \right),$$

that is, in particular for

$$c^* = \frac{\sum_{x \in \mathcal{X}} \beta_x x}{\sum_{x \in \mathcal{X}} \beta_x},$$

and for M^* such that $0 \in \partial_M L(M, c^*, \beta)|_{M=M^*}$, the subdifferential of $L(M, c^*, \beta)$ with respect to M at M^* . This condition can be written as

$$\sum_{x \in \mathcal{X}} \beta_x (x - c^*)(x - c^*)^\top = \tilde{V} \in \partial \log \phi_k^\infty(M)|_{M=M^*},$$

with $\partial \log \phi_k^\infty(M)$ the subdifferential of $\log \phi_k^\infty(M)$,

$$\partial \log \phi_k^\infty(M) = \{V \geq 0 : \Psi_k^{1/k}(V)\phi_k^\infty(M) = \text{trace}(MV) = 1\},$$

see [16], Theorem 7.9. Since $\text{trace}(MV) = 1$ for all $V \in \partial \log \phi_k^\infty(M)$, $\text{trace}(M^*\tilde{V}) = 1$ and thus $\sum_{x \in \mathcal{X}} \beta_x (x - c^*)^\top M^*(x - c^*) = 1$. Also, $\Psi_k^{1/k}(\tilde{V}) = 1/\phi_k^\infty(M^*)$, which gives

$$L(M^*, c^*, \beta) = -\log \Psi_k^{1/k} \left[\sum_{x \in \mathcal{X}} \beta_x (x - c^*)(x - c^*)^\top \right] + \sum_{x \in \mathcal{X}} \beta_x - 1.$$

We obtain finally

$$\begin{aligned} & \min_{\beta \geq 0} L(M^*, c^*, \beta) \\ &= \min_{\gamma > 0, \alpha \geq 0} \left\{ -\log \Psi_k^{1/k} \left[\sum_{x \in \mathcal{X}} \alpha_x (x - c^*)(x - c^*)^\top \right] + \gamma - \log(\gamma) - 1 \right\} \\ &= \min_{\alpha \geq 0} -\log \Psi_k^{1/k} \left[\sum_{x \in \mathcal{X}} \alpha_x (x - c^*)(x - c^*)^\top \right] = -\log \Psi_k^{1/k}(V_k^*), \end{aligned}$$

where we have denoted $\gamma = \sum_{x \in \mathcal{X}} \beta_x$ and $\alpha_x = \beta_x/\gamma$ for all x . Therefore, $T^* \leq -\log \Psi_k^{1/k}(V_k^*)$, that is, $\log[\min_{M, c: \mathcal{X} \subset \mathcal{E}(M, c)} 1/\phi_k^\infty(M)] \geq \log \Psi_k^{1/k}(V_k^*)$. \square

Acknowledgment

The work of the first author was partly supported by the ANR project 2011-IS01-001-01 DESIRE (DESIGns for spatial Random fiElds).

References

[1] Björck, G. (1956). Distributions of positive mass, which maximize a certain generalized energy integral. *Ark. Mat.* **3** 255–269. MR0078470

- [2] DeGroot, M.H. (1962). Uncertainty, information, and sequential experiments. *Ann. Math. Statist.* **33** 404–419. [MR0139242](#)
- [3] Gantmacher, F. (1966). *Théorie des Matrices*. Paris: Dunod.
- [4] Gini, C. (1921). Measurement of inequality of incomes. *Econ. J.* **31** 124–126.
- [5] Giovagnoli, A. and Wynn, H.P. (1995). Multivariate dispersion orderings. *Statist. Probab. Lett.* **22** 325–332. [MR1333191](#)
- [6] Hainy, M., Müller, W.G. and Wynn, H.P. (2014). Learning functions and approximate Bayesian computation design: ABCD. *Entropy* **16** 4353–4374. [MR3255991](#)
- [7] Harman, R. (2004). Lower bounds on efficiency ratios based on Φ_p -optimal designs. In *MODa 7—Advances in Model-Oriented Design and Analysis. Contrib. Statist.* 89–96. Heidelberg: Physica. [MR2089329](#)
- [8] Kiefer, J. (1974). General equivalence theory for optimum designs (approximate theory). *Ann. Statist.* **2** 849–879. [MR0356386](#)
- [9] Kiefer, J. and Wolfowitz, J. (1960). The equivalence of two extremum problems. *Canad. J. Math.* **12** 363–366. [MR0117842](#)
- [10] López-Fidalgo, J. and Rodríguez-Díaz, J.M. (1998). Characteristic polynomial criteria in optimal experimental design. In *MODa 5—Advances in Model-Oriented Data Analysis and Experimental Design (Marseilles, 1998). Contrib. Statist.* 31–38. Heidelberg: Physica. [MR1652210](#)
- [11] Macdonald, I.G. (1995). *Symmetric Functions and Hall Polynomials*, 2nd ed. *Oxford Mathematical Monographs*. New York: Oxford Univ. Press. [MR1354144](#)
- [12] Marcus, M. and Minc, H. (1992). *A Survey of Matrix Theory and Matrix Inequalities*. New York: Dover Publications. [MR1215484](#)
- [13] Oja, H. (1983). Descriptive statistics for multivariate distributions. *Statist. Probab. Lett.* **1** 327–332. [MR0721446](#)
- [14] Pronzato, L. (1998). On a property of the expected value of a determinant. *Statist. Probab. Lett.* **39** 161–165. [MR1652548](#)
- [15] Pronzato, L. and Pázman, A. (2013). *Design of Experiments in Nonlinear Models: Asymptotic Normality, Optimality Criteria and Small-Sample Properties. Lecture Notes in Statistics* **212**. New York: Springer. [MR3058804](#)
- [16] Pukelsheim, F. (1993). *Optimal Design of Experiments. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics*. New York: Wiley. [MR1211416](#)
- [17] Rao, C.R. (1982). Diversity and dissimilarity coefficients: A unified approach. *Theoret. Population Biol.* **21** 24–43. [MR0662520](#)
- [18] Rao, C.R. (1982). Diversity: Its measurement, decomposition, apportionment and analysis. *Sankhyā Ser. A* **44** 1–22. [MR0753075](#)
- [19] Rao, C.R. (1984). Convexity properties of entropy functions and analysis of diversity. In *Inequalities in Statistics and Probability (Lincoln, Neb., 1982). Institute of Mathematical Statistics Lecture Notes—Monograph Series* **5** 68–77. Hayward, CA: IMS. [MR0789236](#)
- [20] Rao, C.R. (2010). Quadratic entropy and analysis of diversity. *Sankhya A* **72** 70–80. [MR2658164](#)
- [21] Rodríguez-Díaz, J.M. and López-Fidalgo, J. (2003). A bidimensional class of optimality criteria involving ϕ_p and characteristic criteria. *Statistics* **37** 325–334. [MR1997183](#)
- [22] Schilling, R.L., Song, R. and Vondraček, Z. (2012). *Bernstein Functions: Theory and Applications*, 2nd ed. *de Gruyter Studies in Mathematics* **37**. Berlin: de Gruyter. [MR2978140](#)
- [23] Serfling, R.J. (1980). *Approximation Theorems of Mathematical Statistics*. New York: Wiley. [MR0595165](#)
- [24] Shaked, M. (1982). Dispersive ordering of distributions. *J. Appl. Probab.* **19** 310–320. [MR0649969](#)
- [25] Shor, N. and Berezovski, O. (1992). New algorithms for constructing optimal circumscribed and inscribed ellipsoids. *Optim. Methods Softw.* **1** 283–299.

- [26] Titterton, D.M. (1975). Optimal design: Some geometrical aspects of D -optimality. *Biometrika* **62** 313–320. [MR0418355](#)
- [27] van der Vaart, H.R. (1965). A note on Wilks' internal scatter. *Ann. Math. Statist.* **36** 1308–1312. [MR0178533](#)
- [28] Wilks, S. (1932). Certain generalizations in the analysis of variance. *Biometrika* **24** 471–494.
- [29] Wilks, S.S. (1960). Multidimensional statistical scatter. In *Contributions to Probability and Statistics* 486–503. Stanford, CA: Stanford Univ. Press. [MR0120721](#)
- [30] Wilks, S.S. (1962). *Mathematical Statistics. A Wiley Publication in Mathematical Statistics*. New York: Wiley. [MR0144404](#)

Received June 2015 and revised January 2016