

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/103192/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Panesar, Sukhmeet Singh, deSilva, Debra, Carson-Stevens, Andrew , Cresswell, Kathrin M, Salvilla, Sarah Angostora, Slight, Sarah Patricia, Javad, Sundas, Netuveli, Gopalakrishnan, Larizgoitia, Itziar, Donaldson, Liam J, Bates, David W and Sheikh, Aziz 2016. How safe is primary care? A systematic review. *BMJ Quality & Safety* 25 (7) , pp. 544-553. 10.1136/bmjqs-2015-004178

Publishers page: <http://dx.doi.org/10.1136/bmjqs-2015-004178>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



# BMJ Quality & Safety

## How Safe is Primary Care? A Systematic Review

Journal:	<i>BMJ Quality &amp; Safety</i>
Manuscript ID:	bmjqs-2015-004178.R1
Article Type:	Systematic Review
Keywords:	Patient safety, Primary care, Risk management

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1     **How Safe is Primary Care? A Systematic Review**

4     **Key words:** primary care; patient safety; incidents; harm; errors

6     **Word count:** 3,836

Confidential: For Review Only

## ABSTRACT

---

**Importance:** Improving patient safety is at the forefront of policy and practice. While considerable progress has been made in understanding the frequency, causes, and consequences of error in hospitals, less is known about the safety of primary care.

**Objective:** We investigated how often patient safety incidents occur in primary care and how often these were associated with patient harm.

**Evidence Review:** We searched 18 databases and contacted international experts to identify published and unpublished studies available between January 1, 1980 and July 31, 2014. Patient safety incidents of any type were eligible. Eligible studies were critically appraised using validated instruments and data were descriptively and narratively synthesized.

**Findings:** Nine systematic reviews and 100 primary studies were included. Studies reported between <1 and 24 patient safety incidents per 100 consultations. The median from population-based record review studies was 2 to 3 incidents for every 100 consultations / records reviewed. It was estimated that around 4% of these incidents may be associated with severe harm, defined as significantly impacting on a patient's wellbeing, including long-term physical or psychological issues or death (range <1% to 44% of incidents). Incidents relating to diagnosis and prescribing were most likely to result in severe harm.

**Conclusions and Relevance:** Millions of people throughout the world use primary care services on any given day. This review suggests that safety incidents are relatively common, but most do not result in serious harm that reaches the patient. Diagnostic and prescribing incidents are the most likely to result in avoidable harm.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Systematic Review Registration:** This systematic review is registered with the International  
Prospective Register of Systematic Reviews (PROSPERO CRD42012002304).

Confidential: For Review Only

## INTRODUCTION

---

Health services strive to provide good quality care, but sometimes people are inadvertently harmed.[1] Between 3% and 17% of people admitted to hospital may experience a safety incident,[1-6] and it is commonly reported that about 10% of hospitalized patients may experience harm.[7-9] Over the last two decades, a substantial body of work has been undertaken to understand the reasons that patient safety incidents occur in hospitals and the effectiveness of interventions to avoid and reduce the impact of such incidents.[10-15] Far less is known about the nature, causes or consequences of incidents in primary care.[16] This may be due to many factors, including the assumption that primary care is safer than hospital care, because primary care is in the early stages of development in some parts of the world, and because primary care medical records may not always be easily accessible, thus making it difficult to study patient safety incidents.

An important first step in preventing harm in primary care is to understand how often patient safety incidents occur, what type of incidents occur, and what impact they have. This is particularly important given the drive for universal access, which is predicated on enhanced provision of primary care.[17, 18] The global drive towards primary care-based models of care has been supported by the World Health Organization (WHO) in low- and middle-income countries and economic pressures in industrialized nations. This is particularly true in the US, which is expanding primary care through the creation of Accountable Care Organizations (ACOs) and Patient-Centered Medical Homes (PCMH). It is important to understand how this expansion can proceed in a safe, sustainable manner. We were commissioned by WHO to investigate the frequency of patient safety incidents in primary care and the resulting harm in order to set the scene for deliberations on how to prevent incidents and minimize their impact.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

59   **METHODS**

---

61   This systematic review of published and unpublished literature was conducted according to  
62   PRISMA guidelines.[19] Our review is registered with the PROSPERO database (PROSPERO  
63   CRD42012002304). We provide a summary of our methods below. Readers are referred to full  
64   details about the methodology which are freely available online, including as part of the online  
65   supplement.[20]

67   **Inclusion and Exclusion Criteria**

68   Studies were eligible for inclusion if they were systematic reviews or primary research  
69   conducted in humans and focused on patient safety incidents in primary care. Box 1 outlines  
70   how we defined primary care, safety incidents, severity of harm, and other key terms. We were  
71   interested in studies that included data about one or more of the following:

- 72       1. Number of safety incidents
- 73       2. Type of safety incidents
- 74       3. Severity of harm associated with safety incidents

76   Primary care varies widely between and within countries so our search strategy covered a broad  
77   range of care delivered outside hospital.[21-27] However, for the purposes of this article, we  
78   focused on studies describing models of care that were comparable with US notions of primary  
79   care and incidents of commission rather than omission (see Box 1 for definitions). A broader  
80   range of studies were identified on topics such as community pharmacy, but these have not  
81   been summarized here. Similarly, studies with a broad ‘ambulatory care’ focus were not  
82   included if these combined settings such as hospital, outpatients and primary care. Only studies  
83   with a primary care focus were included where that focus could be ascertained from the title and  
84   from the abstract.



85

86 Studies that aimed to test an intervention and collected safety data incidentally were not  
87 included, because the aim was to investigate the frequency of safety incidents and harm in  
88 routine practice, not when an intervention was undertaken.

89

90 Published and unpublished research available between January 1, 1980 and July 31, 2014 was  
91 eligible. Eligible study types were 1) systematic reviews; 2) primary studies not included in the  
92 reviews; and 3) primary studies included in the reviews only if they contained empirical data to  
93 feed into the calculation of specific estimates of harm that were not available in the reviews  
94 themselves (33 studies). Non-systematic reviews, case series, or case reports were not eligible  
95 for inclusion and nor were studies included in other systematic reviews that did not contain  
96 exact rates of harms for use in our calculations

97

98

99



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Box 1: Definitions of terms used in the review**

**Harm – no harm, low harm, moderate harm and severe harm**

Harm was defined as impairing the structure or function of the body or mind. This may include pain, nausea, psychological distress, disability, or death. The criteria used in individual studies included in the review were extracted but we standardized the descriptors of harm based on methods suggested by the United Kingdom’s (UK) National Patient Safety Agency:[35]

- No harm: any patient safety incident that had the potential to cause harm but was prevented, resulting in no harm, or that ran to completion but no harm occurred
- Low harm: required extra observation or minor treatment and caused minimal harm
- Moderate harm: resulted in a moderate increase in treatment and caused significant but not permanent harm (an example would be hospitalization)
- Severe harm: resulted in permanent harm such as disability, death, or long-lasting physical or mental consequences

**Incidents of commission**

Incidents of commission were defined as those occurring when something was actively done incorrectly or inappropriately, such as prescribing the wrong dose of medication.

**Incidents of omission**

Incidents of omission were defined as those occurring when there was a lapse in the quality of care. Such incidents were outside the scope of the review.

**Patient safety incidents**

Patient safety incidents were defined as any unintended or unexpected incident(s) that could have or were judged to have led to patient harm. ‘Patient safety incident’ is an umbrella term which is used to describe a single incident or a series of incidents that occur over time. The LINNAEUS taxonomy is a primary care patient safety classification system that categorizes incidents into ‘Process incidents’ or ‘Knowledge or skill incidents’ and then further divides these into sub-categories.[34] This taxonomy was used as an initial coding framework to group studies based on their primary focus of investigation. Studies including incidents that could not be classified in this way were categorized as ‘Other.’

**Primary care**

Primary care was defined as ‘first port of call’ generalist care, delivered outside hospital inpatient settings. Although in some countries primary care includes, for example, community nursing and community pharmacy in addition to family practice clinics, for the purposes of this review we used the following definition of the US primary care workforce: “...includes the specialties of family practice, general practice, general internal medicine, and general paediatrics and, for women patients, obstetricians and gynaecologists.”[35]

## Search Strategy and Study Selection

Search terms were developed based on an international taxonomy for patient safety and previous work.[28-34, 36] Our search terms are available via the online supplement.[20]

Eighteen databases containing published and unpublished literature were searched, including: African Index Medicus, African Journals Online, Bioline International, CINAHL, Embase, IndMED, HINARI, Iran MEDEX, Korean MED, Latin American and Caribbean Health Sciences, Medline, NepJOL, PsycINFO, Thai Index Medicus, WHOLIS, Google Scholar, SIGLE. The final three databases in this list include gray literature. We also searched 'The Grey Literature Report' (<http://www.greylit.org/>). We also searched AHRQ Patient Safety Network (<http://www.psnet.arhq.gov>) which is a patient safety literature clearing house. WHO invited an international panel of primary care clinicians and policy-makers to identify additional published and unpublished studies. Further material was sought using the bibliographies of identified papers and by contacting experts through WHO's six regional offices.

Where primary studies appeared in the systematic reviews that met our inclusion criteria, these were not analysed separately.

Studies identified as potentially suitable were assessed for inclusion by two independent reviewers (SSP and AC-S), with arbitration by a third reviewer (AS), if necessary. The full text of all papers was rescreened by a third reviewer (DdS) when revising the manuscript.

## Quality Assessment

Studies were quality appraised to assess internal and external validity[37] using the Critical Appraisal Skills Programme (CASP) for systematic reviews[38] and the Evidence Based Library and Information Practice (EBLIP) Critical Appraisal Checklist for epidemiologic studies.[39] An

overall grading of the individual components was given for each study. Quality appraisal was independently carried out by two reviewers (SSP and AC-S, or KC and SAS). Disagreements were resolved through discussion, with arbitration by an additional reviewer, if necessary (AS).

**Data Extraction and Synthesis**

Preliminary data were abstracted onto a customized data extraction sheet by two independent reviewers (SSP and AC-S, or KC and SAS), with arbitration by an additional reviewer if necessary (AS). Data were then re-extracted by a third reviewer (DdS) about country of origin; study design; measurement methods; frequencies of patient safety incidents, and burden of harm.

There was significant heterogeneity in the countries of origin, research methods used, type of safety incidents analyzed and metrics so it was not appropriate to combine the data using meta-analysis. A descriptive and narrative synthesis of the data was undertaken. The frequency of incidents and harm were tabulated and graphed. Median incident rates were calculated based on population-based record review studies.

This review does not aim to provide a definitive summary statistic for the frequency of incidents, but rather to show the range in estimates. The rate of per 100 consultations / people / prescriptions was either drawn directly from articles or calculated from data provided in the articles. For example, if articles provided a percentage, we reframed this as a rate out of 100. Equally, if articles provided a numerator and denominator, we converted this to a denominator of 100, if appropriate. This does not allow exact comparability because the unit of measurement differed between studies (people / prescriptions / consultations), but provides a summary of broad trends.

## FINDINGS

---

### Number, Type, and Setting of Studies Included

We screened 61,521 articles and 109 studies met our inclusion criteria: nine systematic reviews and 100 primary studies (see Figure 1). Eighty-eight percent of the systematic reviews (8/9) and 12% of the primary studies (12/100) were judged to be of high quality.

Thirty-six percent of studies were from the US or Canada (39/109), 39% were from Europe (43/109), 13% were from other Organization for Economic Co-operation and Development (OECD) countries (14/109), and 12% were from non-OECD countries (13/109). All studies focused on primary care / family practice / general practice clinics (or aggregated data drawn from these services). Most studies used record review or prescription review to measure safety incidents so they were using an epidemiologic measurement approach. A minority used incident reporting systems and very few used interviews or surveys. The online supplement provides a summary of the characteristics, methods and findings of each study.[20]

### Frequency of Patient Safety Incidents in Primary Care

Fifty-nine studies (some of which were reported in more than one paper) provided an estimate of the frequency of patient safety incidents (see Table 1 in the online supplement). Twelve studies collating 'any type of patient safety incident' reported between <1 and 24 incidents per 100 consultations (see Figure 2). Most of the studies reporting incident rates were not of high quality, though most involved record review or review of prescriptions. Coupled with the large variation in estimates, this means that we cannot confidently state the rate of patient safety incidents in primary care. However, the median of studies based on record review was 2 to 3 incidents per 100 consultations / patient records reviewed.

**Types of Safety Incidents**

Studies documenting the type of safety incidents identified that the three most common categories were: administrative and communication incidents; diagnostic incidents; and prescribing and medication management incidents. Although some studies defined 'communication issues' as safety incidents, the majority of these incidents did not result in harm. The severity of harm associated with incidents is described below.

It is outside the scope of this review to comment about all the types of incidents and their relative frequency, but we provide some broad ranges to show the variation in research. For instance, the proportion of incidents relating to administrative and communication issues ranged between 6% and 67% of all incidents in individual studies.[40-49] Some studies estimated that administration incidents occurred in at least 6% of patient contacts.[50] Most of these incidents related to issues such as incomplete, unavailable, unclear or incorrect documentation;[7, 34, 51, 52] inappropriate monitoring of laboratory tests;[53] or insufficient communication between providers or between professionals and patients.[54]

Studies of incident reporting systems suggest diagnostic incidents were responsible for 4% to 45% of all reported patient safety related incidents.[51, 54, 55] Common diagnostic incidents related to misdiagnosis or missed diagnoses.

Thirty-five studies focused explicitly on prescribing incidents, where the rate was between 1 and 90 out of 100 prescriptions issued (see Table 2 in the online supplement). The figures were higher in studies that focused on particular subgroups, such as the elderly or those taking multiple medications.[56-59] It was difficult to compare these studies because they used different measurement approaches and focused on specific patient populations.

Estimates of the rate of dispensing incidents in primary care also varied widely, from less than 2% of prescriptions[51, 60-64] to up to 65%.[52] These variations are likely to reflect the different definitions used (such as whether or not 'not specifying the route of administration' was listed as an incident),[65] study designs,[51, 54, 67, 68] and focusing on certain sub-sets of patients such as those receiving psychotropic medications,[69] those with poly-pharmacy,[57] or those in care homes.[59]

Results varied depending on whether the studies were high or lower quality. For example, a systematic review found that retrospective studies yielded a lower estimate of adverse drug events (3%),[70] compared with prospective evaluations (10%).[71] Therefore, as with the overall rate of safety incidents, it is not possible to draw firm conclusions about the rate of consultations or people who experience diagnosis, communication, or medication incidents, but we can say that these three broad categories made up the bulk of incidents recorded.

### **Harm Associated with Patient Safety Related Incidents**

Although patient safety incidents may be relatively common in primary care, many incidents did not result in actual harm. For instance, 'safety incidents' may include illegible handwriting on prescriptions, even if such incidents do not ultimately impact on wellbeing. The definition of 'safety incidents' often included processes rather than direct patient impacts.

As with estimates of the quantity and type of incidents in primary care, estimates of harm also varied widely. Table 3 (in the online supplement) lists the severity of harm recorded in 33 individual studies. Many of these studies reviewed incident reports. Serious incidents may be more likely to have been reported, so these studies probably over-estimated the severity of harm. Figure 3 illustrates the range of estimates from record review studies only. These types of studies are more likely to give a representative picture because they do not rely on incident

reports or significant event analysis. Studies based on record review had a median estimate of 4% of incidents being associated with severe harm, defined as significantly impacting on a patient's wellbeing, including long-term physical or psychological issues or death (range <1% to 44%).

**Incidents Associated with Harm**

Diagnostic and medication-related incidents were most commonly associated with harm to patients. For example, one study found that 58% of reported misdiagnoses were associated with harm (severity not described).[54] Between 8%[72] and 11%[73] of medication incidents were reported to result in harm (of any severity). These proportions varied depending on the population studied, research design, and outcome of interest. Results also varied depending on whether the studies were high or lower quality; however, the exact proportions are perhaps less important than the fact that it was diagnostic and prescribing errors that were associated with most severe harm.



## DISCUSSION

---

### Statement of Principal Findings

This extensive review suggests that patient safety incidents are a relatively frequent occurrence in primary care, but that most do not result in significant harm to patients. The heterogeneity of studies means that it is not possible to provide a point estimate of the frequency of incidents, but record review studies suggested a median of around 2 to 3 incidents per 100 consultations / patient records reviewed. About 4% of these incidents were associated with severe harm (median of record review studies). Diagnostic and medication incidents were most likely to result in harm and most likely to result in severe harm.

### Strengths and Limitations

This is the most comprehensive synthesis of the evidence available about patient safety incidents in primary care. Our search strategy was broad and looked for published and unpublished studies, with particular effort made to identify research from low- and middle-income country settings (though few studies were found).

The review provides, for the first time, a comprehensive assessment of how common patient safety incidents are in core primary care contexts and how frequently these are associated with harm. When these estimates are considered in light of data about the high rates of use of primary care services,[74] the absolute burden of iatrogenic harm may be large and may increase if primary care expansion continues in a similar fashion to the current models of care.

Understanding the epidemiology of errors in primary care contexts is crucial to baselining, understanding risk factors, and ultimately developing and evaluating strategies to reduce the risk of iatrogenic harm. We have provided a baseline from which to work.

1  
2  
3 317  
4  
5 318 We have also identified some key methodological insights that need to be considered when  
6  
7 319 planning future epidemiological studies. Key amongst these are the need for multi-stakeholder  
8  
9 320 perspectives, validated tools, and longitudinal study designs in representative populations, with  
10  
11 321 boosted samples in high-risk patients (e.g. those with multi-morbidity and/or poly-pharmacy).  
12  
13 322  
14  
15  
16 323 An important issue is whether the findings of the review represent 'typical' primary care practice.  
17  
18 324 To assist this we limited the review to studies focusing on primary care clinics similar to those  
19  
20 325 run in a US context (rather than a wider definition of primary care as may be common in some  
21  
22 326 parts of the world), but this means that other primary care contexts were excluded. It was  
23  
24 327 sometimes differentiate studies for inclusion because they contained a mix of primary care and  
25  
26 328 other care or because the definition and scope of ambulatory care was not included in papers.  
27  
28 329  
29  
30  
31 330 A major limitation is that there is no widely used standardized taxonomy for classifying incidents  
32  
33 331 in primary care settings.[8,75,76] This means that studies defined and measured incidents  
34  
35 332 differently, resulting in variations in the estimated proportion of incidents and harms. Higher  
36  
37 333 quality studies, those with a broad population focus, and those based on record review were  
38  
39 334 more likely to have lower estimates of the frequency and severity of harm. Relationships  
40  
41 335 between country, data quality, setting, and severe harm deserve further attention.  
42  
43 336  
44  
45  
46 337 Most studies used a single method to assess the frequency of incidents, rather than a  
47  
48 338 triangulation of approaches. This may under-estimate the frequency of incidents. The primary  
49  
50 339 care record systems used to record incidents may also be open to coding errors.  
51  
52 340  
53  
54  
55  
56  
57  
58  
59  
60

As with all systematic reviews, publication bias may be present, whereby certain types of studies may be more likely to be published. We sought to address this by searching extensively for unpublished literature and by sense-checking findings with international experts. The large variability of findings suggests that our results were not unduly tarnished by only the highest levels of incidents or harms being reported. The variability of estimates remained regardless of whether studies were rated as high or low quality.

Our graphical representation of the severity of harm is based on studies that reviewed records, rather than relying on incident reports. This is because incident reports may be less likely to capture incidents that have low severity and people may be more likely to report some types of incidents than others. Studies of malpractice claims may be particularly open to bias so we excluded these when reporting median rates.

### **Recommendations for Policy, Practice, and Future Research**

Patient safety is high on the policy radar when developing and rolling-out new models of primary care (e.g. PCMHs). But these 'solutions' can introduce new risks which need to be proactively identified. For example, health information technology (HIT) has been seen as an important solution to enhancing safety, but it is now appreciated that such technology can also introduce some new risks.[77] Having standardized methods to identify and quantify these risks is essential.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Key implications of this review include the need to develop a standardized set of definitions of core terminology, the need to promote mixed-methods evaluations that triangulate different sources of evidence, and a particular focus on diagnostic and medication errors, which appear to be most frequently associated with severe patient harm. There is also a need to better identify those at greatest risk of experiencing patient safety incidents and the nature of the incidents that occur, because such insights will be crucial to developing interventions to decrease the burden of iatrogenic harm.

A standardized taxonomy for classifying incidents and harm would allow comparisons across settings, countries, and over time. Longitudinal, multi-methods investigations would provide more insight into the extent of harm associated with different types of error. More in-depth analysis into particular areas of high risk is warranted, in particular people with multiple long-term conditions and associated poly-pharmacy. The paucity of evidence about low- and middle-income countries highlights the need for more work to understand the nature of incidents and the opportunities for prevention in these resource-scarce contexts.

The estimated proportion of patient safety incidents in primary care is generally lower than the estimated 10% of people who experience events in hospital,[9,78] but primary care and hospital encounters are not easily comparable because hospitalized patients experience multiple clinical encounters during a single admission. Furthermore, the overall volume of people using primary care is substantially higher than those using hospital services in many parts of the world, so even if incidents occur in a lower proportion of visits, this translates into a considerable burden of potential harm, though most is not severe.[79] However, incidents occur in all care settings so this finding is only useful if it prompts policy-makers and clinicians to do something about it.[80]

Better prediction tools and more experimental studies are needed to understand which incidents we can avoid, and how best to do so.

388

389 It is important to consider whether adverse drug events and similar are 'incidents' if the  
390 medication was correctly prescribed, dispensed, and monitored. In this review, all incidents  
391 were counted, as it was not possible to distinguish 'correct' prescribing from individual studies.  
392 However, it may be more useful to consider whether incidents are preventable versus a result of  
393 proper care. Research has attempted to determine what proportion of safety incidents may be  
394 preventable. Most studies of this type relate to prescribing and medication management  
395 incidents,[53, 69, 78, 81, 82] and use observational cross-sectional designs, so it can be  
396 challenging to interpret the estimates. A small number of more robust before-and-after studies  
397 and randomized controlled trials have found that up to half of all incidents may be preventable  
398 using interventions such as pharmacist-led medication review, computerized physician order  
399 entry (CPOE) and computerized decision support (CDS) systems, error alert systems, and  
400 education of professionals.[83-91]

401

402 **Conclusions**

403 Primary care services are expanding globally, providing a first port of call to millions of people  
404 every day.[18] Universal access to healthcare remains firmly on the agenda of policy-makers  
405 however, these services are not without potential harms. This review has suggested major gaps  
406 in the evidence base which now need to be filled. WHO's forthcoming roadmap on *Safer*  
407 *Primary Care for All* will explore the most effective ways to prevent incidents, particularly those  
408 most likely to cause serious harm.

409

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

410 To further support this journey, there is a need for researchers to use existing well-developed  
411 definitions, taxonomies, and tools, such as the NPSA definition and the Linnaeus  
412 taxonomy,[35,92] to allow greater comparability between studies and research contexts. There  
413 is also a need for better quality epidemiological studies, but the review shows that focusing on  
414 diagnostic and prescribing errors – which are the most frequent and hence important sources of  
415 significant iatrogenic harm – needs to be a priority for research and policy.[93,94]

Confidential:- For Review Only

**FIGURE LEGENDS**

Figure 1: PRISMA diagram of studies included in the review

Figure 2: Safety incidents per 100 primary care consultations / records – results from 12 studies

Figure 3: Proportion of safety incidents in primary care resulting in severe harm – results from 7 studies based on record review



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

422    **Acknowledgements:**  
423    **Competing interests:**  
424  
425  
426  
427  
428  
429    **Funding:**  
430  
431  
432  
433  
434  
435    **Transparency declaration**  
436  
437  
438  
439  
440    **Contributorship:**  
441  
442  
443  
444  
445    **Copyright:** The Corresponding Author has the right to grant on behalf of all authors and does  
446 grant, on behalf of all authors, a worldwide license to the Publishers and its licensees in  
447 perpetuity, in all forms, formats, and media (whether known now or created in the future), to i)

1  
2  
3 448 publish, reproduce, distribute, display, and store the Contribution, ii) translate the Contribution  
4  
5 449 into other languages, create adaptations, reprints, include within collections and create  
6  
7 450 summaries, extracts and/or, abstracts of the Contribution, iii) create any other derivative work(s)  
8  
9 451 based on the Contribution, iv) to exploit all subsidiary rights in the Contribution, v) the inclusion  
10  
11 452 of electronic links from the Contribution to third party material wherever it may be located; and,  
12  
13 453 vi) license any third party to do any or all of the above.  
14  
15  
16 454  
17  
18 455  
19  
20 456  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

REFERENCES

1. Landrigan CP, Parry GJ, Bones CB, et al. Temporal trends in rates of patient harm resulting from medical care. *N Engl J Med*. 2010;363(22):2124-2134.

2. Baker GR, Norton PG, Flintoft V, et al. The Canadian adverse events study: the incidence of adverse events among hospital patients in Canada. *CMAJ*. 2004;170:1678-1686.

3. Brennan TA, Leape LL, Laird NM, et al. Incidence of adverse events and negligence in hospitalised patients: results of the Harvard Medical Practice Study I. 1991. *Qual Saf Health Care*. 2004;13:145-151.

4. Davis P, Lay-Yee R, Briant R, et al. Adverse events in New Zealand public hospitals. I. Occurrence and impact. *N Z Med J*. 2002;115:U271.

5. Schioler T, Lipczak H, Pedersen BL, et al. Incidence of adverse events in hospitals. A retrospective study of medical records. *Ugeskr Laeger*. 2001;163:5370-5378.

6. Thomas EJ, Studdert DM, Burstin HR, et al. Incidence and types of adverse events and negligent care in Utah and Colorado. *Med Care*. 2000;38:261-271.

7. Vincent C, Neale G, Woloshynowych M. Adverse events in British hospitals: preliminary retrospective record review. *BMJ*. 2001;322:517-519.

8. Makeham M, Dovey S, Runciman W, Larizgiotia I. *Methods and measures used in primary care patient safety research*, 2008. [www.who.int/patientsafety/research/methods\\_measures/makeham\\_dovey\\_full.pdf](http://www.who.int/patientsafety/research/methods_measures/makeham_dovey_full.pdf). Accessed 1 January 2015.

9. Kohn LT, Corrigan JM, Donaldson MS (Institute of Medicine). *To Err is Human: Building a Safer Health System*. Washington, DC: National Academy Press, 2000.

10. Clinton HR, Obama B. Making patient safety the centerpiece of medical liability reform. *N Engl J Med*. 2006;354(21):2205-2208.

11. Leape LL, Brennan TA, Laird NM, et al. The nature of adverse events in hospitalised patients: Results from the Harvard Medical Practice Study II. *N Engl J Med*. 1991; 324:377-384.

12. Bates DW, Cullen DJ, Laird N, et al. Incidence of adverse drug events and potential adverse drug events. Implications for prevention. ADE Prevention Study Group. *JAMA*. 1995;274(1):29-34.

13. World Health Organization. *Clean Care is Safer Care*. [www.who.int/gpsc/en/index.html](http://www.who.int/gpsc/en/index.html). Accessed 1 January 2015.

14. World Health Organization. *Safe Surgery Saves Lives*. [www.who.int/patientsafety/safesurgery/en/](http://www.who.int/patientsafety/safesurgery/en/). Accessed 1 January 2015.

15. Haynes AB, Weiser TG, Berry WR, et al. Safe Surgery Saves Lives Study Group. A surgical safety checklist to reduce morbidity and mortality in a global population. *N Engl J Med*. 2009;360(5):491-499.

16. World Health Organization. *Safer Primary care – A Global Challenge*. [www.who.int/patientsafety/summary\\_report\\_of\\_primary\\_care\\_consultation.pdf](http://www.who.int/patientsafety/summary_report_of_primary_care_consultation.pdf). Accessed 1 January 2015.

17. World Health Organization. *The World Health Report 2008 – Primary Health Care (Now More than Ever)*. [www.who.int/whr/2008/en/index.html](http://www.who.int/whr/2008/en/index.html). Accessed 22 September 2014.

18. Sheikh A, Panesar SS, Larizgiotia I, Bates DW, Donaldson LJ. Safer Primary care for All – A Global Imperative. *Lancet Global Health* (accepted July 2013).

19. Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* 6(6): e1000097. doi:10.1371/journal.pmed1000097.
20. Panesar SS, Carson-Stevens A, Salvilla SA, et al. Estimating the frequency of errors and the global burden from iatrogenic harm in primary care: protocol for a systematic review and meta-analysis (Accepted April 2012)  
[www.crd.york.ac.uk/PROSPEROFILES/2304\\_PROTOCOL\\_20120323.pdf](http://www.crd.york.ac.uk/PROSPEROFILES/2304_PROTOCOL_20120323.pdf). Accessed 1 January 2015.
21. Declaration of Alma-Ata. International Conference on Primary Health Care. 1978. Alma-Ata, USSR: World Health Organization.  
[http://www.who.int/publications/almaata\\_declaration\\_en.pdf](http://www.who.int/publications/almaata_declaration_en.pdf) Accessed 1 January 2015.
22. Institute of Medicine. *Primary care: America's Health in a New Era*.  
[www.nap.edu/openbook.php?record\\_id=5152&page=27](http://www.nap.edu/openbook.php?record_id=5152&page=27) Accessed 1 January 2015.
23. Starfield B. Basic concepts in population health and health care. *Journal of Epidemiology and Community Health*. 2001;55:452-454.
24. WHO. Health Systems Strengthening Glossary.  
[www.who.int/healthsystems/hss\\_glossary/en/index8.html#8](http://www.who.int/healthsystems/hss_glossary/en/index8.html#8). Accessed 1 January 2015.
25. Hospital Report Research Collaborative. New measures of ambulatory care performance in Ontario. 2006.  
[www.hospitalreport.ca/downloads/otherreports/SNAPSHOT\\_REPORT\\_Amb\\_FINAL.pdf](http://www.hospitalreport.ca/downloads/otherreports/SNAPSHOT_REPORT_Amb_FINAL.pdf). Accessed 1 January 2015.
26. World Health Organization. More than words. Conceptual Framework for the International Classification for Patient Safety. 2009.  
[www.who.int/patientsafety/implementation/taxonomy/icps\\_technical\\_report\\_en.pdf](http://www.who.int/patientsafety/implementation/taxonomy/icps_technical_report_en.pdf). Accessed 1 January 2015.
27. World Bank. How we classify countries. 2012. <http://data.worldbank.org/about/country-classifications>. Accessed 1 January 2015.
28. Kingston-Reichers J, Ospina M, Jonsson E, et al. *Patient Safety in Primary care*. Edmonton, AB: Canadian Patient Safety Institute and BC Patient Safety and Quality Council. 2010.
29. Makeham M, Dovey S, Runciman W, Larizgoitia I. Methods and measures used in primary care patient safety research.  
[www.who.int/patientsafety/research/methods\\_measures/primary\\_care\\_ps\\_research/en/index.html](http://www.who.int/patientsafety/research/methods_measures/primary_care_ps_research/en/index.html). Accessed 1 January 2015.
30. Tanon AA, Champagne F, Contandriopoulos AP, et al. Patient safety and systematic reviews: finding papers indexed in MEDLINE, EMBASE and CINAHL. *Qual Saf Health Care*. 2010;19(5):452-461.
31. Pearson A, Aromataris A. Patient Safety Primary care: a review of the literature. Australian Commission on Safety and Quality in healthcare. <http://www.safetyandquality.gov.au/wp-content/uploads/2009/01/Patient-Safety-in-Primary-Health-Care-A-Review-of-the-Literature-2009.pdf> Accessed 1 January 2015.
32. Anesthesia Patient Safety Foundation. [www.apsf.org/about\\_safety.php](http://www.apsf.org/about_safety.php). Accessed 1 January 2015.
33. Dovey SM, Meyers DS, Phillips Jr RL, et al. A preliminary taxonomy of medical errors in family practice. *Qual Saf Health Care*. 2002;11(3):233-238.
34. Singh, R. Singh, A. Singh, S. Singh, G. Creating a Common Vision for all Stakeholders to Make Healthcare Safer with Interactive Visual Modeling. Special Issue on Soft Computing, Simulation, and Web centric Computing on Selected Papers from NASTEC 2009, *International Journal of Advanced Intelligence Paradigms* (IJAIIP). 2011;3(1).

1  
2  
3 550 35. National Patient Safety Agency. Seven Steps to Patient Safety. 2004.  
4 551 [www.nrls.npsa.nhs.uk/resources/collections/seven-steps-to-patient-safety/?entryid45=59787](http://www.nrls.npsa.nhs.uk/resources/collections/seven-steps-to-patient-safety/?entryid45=59787).  
5 552 Accessed 1 January 2015.

6  
7 553 36. Bindman Andrew B, Majeed Azeem. Organisation of primary care in the United States *BMJ*.  
8 554 2003;326:631.

9  
10 555 37. Cochrane Group. Assessing risk of bias in included studies. 2011.  
11 556 [http://handbook.cochrane.org/chapter\\_8/8\\_assessing\\_risk\\_of\\_bias\\_in\\_included\\_studies.htm](http://handbook.cochrane.org/chapter_8/8_assessing_risk_of_bias_in_included_studies.htm).  
12 557 Accessed 1 January 2015.

13 558 38. Solutions for Public Health. Critical Appraisal Skills Programme. 2007. [http://www.casp-](http://www.casp-uk.net/#!/casp-tools-checklists/c18f8)  
14 559 [uk.net/#!/casp-tools-checklists/c18f8](http://www.casp-uk.net/#!/casp-tools-checklists/c18f8). Accessed 1 January 2015.

15 560 39. Glynn L. EBLIP Critical Appraisal Checklist. 2006.  
16 561 <http://ebltoolkit.pbworks.com/f/EBLCriticalAppraisalChecklist.pdf>. Accessed 1 January 2015.

17  
18 562 40. Cox SJ, Holden JD. A retrospective review of significant events reported in one district in  
19 563 2004-2005. *British Journal of General Practice*. 2007;57(542):732-736.

20 564 41. Hansen LB, Fernald D, Araya-Guerra R, et al. Pharmacy clarification of prescriptions ordered  
21 565 in primary care: A report from the applied strategies for improving patient safety (ASIPS)  
22 566 collaborative. *Journal of the American Board of Family Medicine*. 2006;19(1):24-30.

23 567 42. McKay J, Bradley N, Lough M, Bowie P. A review of significant events analysed in general  
24 568 practice: implications for the quality and safety of patient care. *BMC Family Practice*.  
25 569 2009;10:61.

26 570 43. Diamond MR, Kamien M, Sim MG, Davis J. A critical incident study of general practice  
27 571 trainees in their basic general practice term. *Med J Aust*. 1995;162(6):321-324.

28 572 44. Elder NC, Vonder Meulen M, Cassidy A. The identification of medical errors by family  
29 573 physicians during outpatient visits. *Annals of Family Medicine*. 2004;2(2):125-129.

30 574 45. Rosser W, Dovey S, Bordman R, White D, Crighton E, Drummond N. Medical errors in primary  
31 575 care: results of an international study of family practice. Canadian Family Physician *Medecin*  
32 576 *de Famille Canadien*. 2005;51:386-387.

33 577 46. Fit KE, Burkiewicz JS, Sweeney BL. PDA-based documentation of medication interventions in  
34 578 an ambulatory care setting. *Journal of Pharmacy Technology*. 2007;23(1):9-16.

35 579 47. Bhasale AL, Miller GC, Reid SE, Britt HC. Analysing potential harm in Australian general  
36 580 practice: an incident-monitoring study. *Med J Aust*. 1998;169(2):73-76.

37 581 48. Kuzel AJ, Woolf SH, Gilchrist VJ, et al. Patient reports of preventable problems and harms in  
38 582 primary health care. *Ann Fam Med*. 2004;2(4):333-340.

39 583 49. Hickner J, Graham DG, Elder NC, et al. Testing process errors and their harms and  
40 584 consequences reported from family medicine practices: a study of the American Academy of  
41 585 Family Physicians National Research Network. *Qual Saf Health Care*. 2008;17(3):194-200.

42 586 50. Sayers YM, Armstrong P, Hanley K. Prescribing errors in general practice: A prospective  
43 587 study. *European Journal of General Practice*. 2009;15(2):81-83.

44 588 51. Teagarden JR. Dispensing error rate in a highly-automated mail-service pharmacy practice.  
45 589 *Pharmacotherapy*. 2005;25(11):1629-1635.

46 590 52. Teinila T, Gronroos V, Airaksinen M. Survey of dispensing error practices in community  
47 591 pharmacies in Finland: a nationwide study. *Journal of the American Pharmacists Association*.  
48 592 2009;49(5):604-610.

49 593 53. Mitchell ED, Rubin G, Macleod U. Understanding diagnosis of lung cancer in primary care:  
50 594 qualitative synthesis of significant event audit reports. *Br J Gen Pract*. 2013;63(606):e37-46.

51  
52  
53  
54  
55  
56  
57  
58  
59  
60



54. Hoffmann B, Beyer M, Rohe J, Gensichen J, Gerlach FM. "Every error counts": a web-based incident reporting and learning system for general practice. *Qual Saf Health Care*. 2008;17(4):307-312.
55. Ilboudo TP, Chou YJ, Huang N. Assessment of providers' referral decisions in rural Burkina Faso: a retrospective analysis of medical records. *BMC Health Serv Res* 2012;12:54.
56. Stewart AL, Lynch KJ. Identifying discrepancies in electronic medical records through pharmacist medication reconciliation. *J Am Pharm Assoc*. 2012;52:59-68.
57. Bradley MC, Fahey T, Cahir C, et al. Potentially inappropriate prescribing and cost outcomes for older people: a cross-sectional study using the Northern Ireland Enhanced Prescribing Database. *Eur J Clin Pharmacol*. 2012;68(10):1425-1433.
58. Taylor LK, Kawasumi Y, Bartlett G, Tamblyn R. Inappropriate prescribing practices: the challenge and opportunity for patient safety. *Health Q*. 2005;8 Spec No:81-85.
59. Van Der Hooft CS, Jong GW, Dieleman JP, et al. Inappropriate drug prescribing in older adults: The updated 2002 Beers criteria - A population-based cohort study. *British Journal of Clinical Pharmacology*. 2005;60(2):137-144.
60. Chua SS, Wong ICK, Edmondson H, et al. A feasibility study for recording of dispensing errors and 'near misses' in four UK primary care pharmacies. *Drug Safety*. 2003;26(11):803-813.
61. Flynn EA. Medication dispensing errors in community pharmacies: a nationwide study. *Proc Hum Fact Ergonom Soc*. 2002;48-51.
62. Flynn EA. National observational study of prescription dispensing accuracy and safety in 50 pharmacies. *J Am Pharm Assoc*. 2003;43:191-200.
63. Kayne S. Negligence and the pharmacist. Part 3. Dispensing and prescribing errors. *Pharm J*. 1996;257:32-35.
64. Hoxsie DM. Analysis of community pharmacy workflow processes in preventing dispensing errors. *J Pharm Pract*. 2006;19:124-130.
65. Al Khaja KAJ, Damanhori AHH, Al-Ansari TM, Sequeira RP. Topical corticosteroids in infants: prescribing pattern and prescribing errors in Bahrain. *Pharmacy World & Science*. 2007;29(4):395-399.
66. Honigman B, Lee J, Rothschild J, et al. Using computerized data to identify adverse drug events in outpatients. *J Am Med Inform Assoc*. 2001;8(3):254-66.
67. James KL, Barlow D, McArtney R, et al. Incidence, type and causes of dispensing errors: A review of the literature. *International Journal of Pharmacy Practice*. 2009;17(1):9-30.
68. Ashcroft DM, Quinlan P, Blenkinsopp A. Prospective study of the incidence, nature and causes of dispensing errors in community pharmacies. *Pharmacoepidemiology and Drug Safety*. 2005;14(5):327-332.
69. Guthrie B, McCowan C, Davey P, et al. High risk prescribing in primary care patients particularly vulnerable to adverse drug events: cross sectional population database analysis in Scottish general practice. *BMJ (Overseas & Retired Doctors Edition)*. 2011;342(7812):1406-1407.
70. Makeham MA, Kidd MR, Saltman DC, et al. The Threats to Australian Patient Safety (TAPS) study: incidence of reported errors in general practice. *Med J Aust*. 2006;185(2):95-98.
71. Martinez Sanchez A, Campos RM. Detection of prescribing related problems at the community pharmacy. *International Journal of Clinical Pharmacy*. 2011;33(1):66-69.
72. van Dulmen SA, Tacken MA, Staal JB, Gaal S, Wensing M, Nijhuis-van der Sanden MW. Patient safety in primary allied health care: what can we learn from incidents in a Dutch exploratory cohort study? *Med Care*. 2011;49(12):1089-1096.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

73. Hickner J, Zafar A, Kuo GM, et al. Field test results of a new ambulatory care Medication Error and Adverse Drug Event Reporting System-MEADERS. *Ann Fam Med*. 2010;8(6):517-525.

74. NHS England. Improving general practice – a call to action. <http://www.england.nhs.uk/ourwork/qual-clin-lead/calltoaction/igp-cta/>. Accessed 1 January 2015.

75. Elder NC, Dovey SM. Classification of medical errors and preventable adverse events in primary care: A synthesis of the literature. *Journal of Family Practice*. 2002;51(11):927-932.

76. Sandars J, Esmail A. The frequency and nature of medical error in primary care: understanding the diversity across studies. *Fam Pract*. 2003;20(3):231-236.

77. Institute of Medicine. *Health IT and Patient Safety: Building Safer Systems for Better Care*. 2011. <http://www.iom.edu/Reports/2011/Health-IT-and-Patient-Safety-Building-Safer-Systems-for-Better-Care.aspx>. Accessed 1 January 2015.

78. Vincent C, Neale G, Woloshynowych M. Adverse events in British hospitals: preliminary retrospective record review. *BMJ*. 2001;322(7285):517-519. Erratum in: *BMJ*. 2001;322(7299):1395.

79. Sheikh A, Panesar SS, Larizgoitia A, Bates DW, Donaldson LJ. Safer primary care for all: a global imperative. *Lancet Global Health*. 2013;1:e182-e183.

80. Cresswell K, Panesar SS, Salvilla SA, et al. Global research priorities to better understand the burden of iatrogenic harm in primary care: an international Delphi exercise. *PLoS Med*. 2013;10(11):e1001554.

81. Al Khaja KA, Sequeira RP, Damanhori AH. Medication prescribing errors pertaining to cardiovascular/antidiabetic medications: a prescription audit in primary care. *Fundam Clin Pharmacol*. 2011 Jan 25.

82. Olaniyan JO, Ghaleb M, Dhillon S, Robinson P. Safety of medication use in primary care. *Int J Pharm Pract* (published online June 2014).

83. Al Khaja KAJ, Damanhori AHH, Al-Ansari TM, Sequeira RP. Topical corticosteroids in infants: prescribing pattern and prescribing errors in Bahrain. *Pharmacy World & Science*. 2007;29(4):395-399.

84. Kane-Gill SL, Van JB, Handler SM. Adverse drug reactions in hospital and ambulatory care settings identified using a large administrative database. *Annals of Pharmacotherapy*. 2010;44(6):983-994.

85. Zavaleta-Bustos M, Castro-Pastrana LI, Reyes-Hernandez I, Lopez-Luna MA, Bermudez-Camps IB. Prescription errors in a primary care university unit: Urgency of pharmaceutical care in Mexico.

86. Hider P, Lay-Yee R, Davis P, Briant R, Scott A. Monitoring the quality of primary care: Use of hospital-based audit studies. *In J Risk and Safety in Medicine*. 2005;17(1-2):81-89.

87. Krska J, Cromarty JA, Arris F, et al. Pharmacist-led medication review in patients over 65: a randomized, controlled trial in primary care. *Age Ageing*. 2003;30(3):205-211.

88. Granas AG, Berg C, Hjellvik V, et al. Evaluating categorisation and clinical relevance of drug-related problems in medication reviews. *Pharmacy World and Science*. 2010;32(3):394-403.

89. Tamblyn R, Huang A, Taylor L, et al. A randomized trial of the effectiveness of on-demand versus computer-triggered drug decision support in primary care. *J Am Med Informatics Assoc*. 2008;15(4):430-439.

90. Palen TE, Raebel M, Lyons E, Magid DM. Evaluation of laboratory monitoring alerts within a computerized physician order entry system for medication orders. *Am J Managed Care*. 2006;12(7):389-396.



91. Batuwitage BT, Kingham JGC, Morgan NE, Bartlett RL. Inappropriate prescribing of proton pump inhibitors in primary care. *Postgraduate Medical Journal*. 2007;83(975):66-68.
92. Makeham MA, Dovey SM, County M, et al. An international taxonomy for errors in general practice: A pilot study. *Med J Aust*. 2002;177(2):68-72.
93. Cresswell KM, Panesar SS, Salvilla SA, et al. Global Research Priorities to Better Understand the Burden of Iatrogenic Harm in Primary Care: An International Delphi Exercise. *PLoS Med*. 2013;10(11):e1001554. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3833831/> Accessed 1 January 2015.
94. Avery AJ, Rodgers S, Cantrill JA, et al. A pharmacist-led information technology intervention for medication errors (PINCER): a multicentre, cluster randomised, controlled trial and cost-effectiveness analysis. *Lancet*. 2012;379(9823):1310-1319. Abramson EL, Bates DW, Jenter C, et al. Ambulatory prescribing errors among community-based providers in two states. *J Am Med Inform Assoc*. 2012;19(4):644-648.
95. Abramson EL, Malhotra S, Osorio SN, et al. A long-term follow-up evaluation of electronic health record prescribing safety. *J Am Med Inform Assoc*. 2013;20(e1):e52-58.
96. A-Elgayoum SME, El-Feki AE-KA, Mahgoub BA, El-Rayah E-A, Giha HA: Malaria overdiagnosis and burden of malaria misdiagnosis in the suburbs of central Sudan: special emphasis on artemisinin-based combination therapy era. *Diagnostic Microbiology & Infectious Disease*. 2009;64(1):20-26.
97. Al Khaja KAJ, Al Ansari TM, Damanhori AHH, Sequeira R. Evaluation of drug utilization and prescribing errors in infants: a primary care prescription-based study. *Health Policy*. 2007;81(2-3):350-357.
98. Al Khaja KAJ, Sequeira RP, Al-Ansari T, Damanhori AHH, James H, Handu SS. Pediatric iron preparations for infants in Bahrain: some therapeutic concerns. *International Journal of Clinical Pharmacology & Therapeutics*. 2010;48(3):200-205.
99. Al-Agilly S, Neville RG, Robb H, Riddell S. Involving patients in checking the validity of the NHS shared record: A single practice pilot. *Informatics in Primary Care*. 2007;15(4):217-220.
100. APEAS Study. Patient Safety in primary Health Care. Madrid: Ministry of Health and Consumer Affairs, 2008.
101. Aranaz-Andrés JM, Aibar C, Limón R, et al. A study of the prevalence of adverse events in primary care in Spain. *Eur J Public Health*. 2012;22(6):921-925.
102. Avery AJ, Ghaleb M, Barber N, et al. The prevalence and nature of prescribing and monitoring errors in English general practice: a retrospective case note review. *Br J Gen Pract*. 2013;63(613):e543-553.
103. Bradbury F. How important is the role of the physician in the correct use of a drug? An observational cohort study in general practice. *International Journal of Clinical Practice* (Supplement). 2004;58:27-33.
104. Bradley MC, Fahey T, Cahir C, et al. Potentially inappropriate prescribing and cost outcomes for older people: a cross-sectional study using the Northern Ireland Enhanced Prescribing Database. *Eur J Clin Pharmacol*. 2012;68(10):1425-1433.
105. Bregnhøj L, Thirstrup S, Kristensen MB, Bjerrum L, Sonne J. Prevalence of inappropriate prescribing in primary care. *Pharmacy World and Science*. 2007;29(3):109-115.
106. Brekke M, Rognstad S, Straand J, et al. Pharmacologically inappropriate prescriptions for elderly patients in general practice: How common? Baseline data from The Prescription Peer Academic Detailing (Rx-PAD) study. *Scand J Prim Health Care*. 2008;26(2):80-85.
107. Casalino LP, Dunham D, Chin MH, et al. Frequency of failure to inform patients of clinically significant outpatient test results. *Archives of Internal Medicine*. 2009;169(12):1123-1129.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

108. Clark RC, Maxwell SRJ, Kerr S, et al. The influence of primary care prescribing rates for new drugs on spontaneous reporting of adverse drug reactions. *Drug Safety*. 2007;30(4):357-366.

109. De Wet C, Bowie P. The preliminary development and testing of a global trigger tool to detect error and patient harm in primary-care records. *Postgraduate Medical Journal*. 2009;85(1002):176-180.

110. De Wilde S, Carey IM, Harris T, et al. Trends in potentially inappropriate prescribing amongst older UK primary care patients. *Pharmacoepidemiol Drug Saf*. 2007;16(6):658-667.

111. Eggleton KS, Dovey SM. Using triggers in primary care patient records to flag increased adverse event risk and measure patient safety at clinic level. *NZ Med J*. 2014;127(1390):45-52.

112. Elder NC, Vonder Meulen M, Cassedy A. The identification of medical errors by family physicians during outpatient visits. *Annals of Family Medicine*. 2004;2(2):125-129.

113. Farrow SC, Zeuner D, Hall C. Improving infection control in general practice. *The Journal of the Royal Society for the Promotion of Health*. 1999;119(1):17-22.

114. Field TS, Gurwitz JH, Harrold LR, et al. Strategies for detecting adverse drug events among older persons in the ambulatory setting. *Journal of the American Medical Informatics Association*. 2004;11(6):492-498.

115. Gaal S, Verstappen W, Wolters R, Lankveld H, van Weel C, Wensing M. Prevalence and consequences of patient safety incidents in general practice in the Netherlands: a retrospective medical record review study. *Implementation Science*. 2011;6(1):37.

116. Gurwitz JH, Field TS, Harrold LR, et al. Incidence and preventability of adverse drug events among older persons in the ambulatory setting. *JAMA*. 2003; 289(9):1107-1117.

117. Howard M, Dolovich L, Kaczorowski J, Sellors C, Sellors J. Prescribing of potentially inappropriate medications to elderly people. *Family Practice*. 2004;21(3):244-247.

118. Ilboudo TP, Chou YJ, Huang N. Assessment of providers' referral decisions in rural Burkina Faso: a retrospective analysis of medical records. *BMC Health Serv Res* 2012;12:54.

119. Khoja T, Neyaz Y, Qureshi NA, Magzoub MA, Haycox A, Walley T. Medication errors in primary care in Riyadh city, Saudi Arabia. *Eastern Mediterranean Health Journal*. 2011;17(2):156-160.

120. Khoja TA, Al-Shammari SAI, Farag MK, Al-Mazrou Y. Quality of prescribing at primary care centers in Saudi Arabia. *Journal of Pharmacy Technology*. 1996;12(6):284-288.

121. Khoo EM, Lee WK, Sararaks S, et al. Medical errors in primary care clinics – a cross sectional study. *BMC Fam Pract*. 2012;13:127.

122. Koper D, Kamenski G, Flamm M, Böhmendorfer B, Sönnichsen A. Frequency of medication errors in primary care patients with polypharmacy. *Fam Pract*. 2013;30(3):313-319.

123. Leon AC, Portera L, Olfson M, et al. Diagnostic errors of primary care screens for depression and panic disorder. *International Journal of Psychiatry in Medicine*. 1999;29(1):1-12.

124. Lund BC, Carnahan RM, Egge JA, Chrischilles EA, Kaboli PJ. Inappropriate prescribing predicts adverse drug events in older adults. *Annals of Pharmacotherapy*. 2010;44(6):957-963.

125. Makeham M, Dovey S, Runciman W, Larizgoitia I. Methods and measures used in primary care patient safety research.  
[www.who.int/patientsafety/research/methods\\_measures/primary\\_care\\_ps\\_research/en/index.html](http://www.who.int/patientsafety/research/methods_measures/primary_care_ps_research/en/index.html) Accessed 01 January 2015.

126. McKay J, de Wet C, Kelly M, Bowie P. Applying the Trigger Review Method after a brief educational intervention: potential for teaching and improving safety in GP specialty training? *BMC Med Educ.* 2013;13:117.
127. Neville RG, Robertson F, Livingstone S, Crombie IK. A classification of prescription errors. *The Journal of the Royal College of General Practitioners.* 1989;39(320):110-112.
128. Nicholson D, Hersh W, Gandhi TK, Weingart SN, Bates DW. Medication errors: not just a "few bad apples". *Journal of Clinical Outcomes Management.* 2006;13(2):114-116.
129. Paille F, Pissiochet P. Drug interactions in primary health care: Prospective study of 896 patients treated for hypertension. [French]. *Therapie.* 1995;50(3):253-258.
130. Pandit NB, Choudhary SK. Unsafe injection practices in Gujarat, India. *Singapore Medical Journal.* 2008;49(11):936-939.
131. Ryan C, O'Mahony D, Kennedy J, Weedle P, Byrne S. Potentially inappropriate prescribing in an Irish elderly population in primary care. *British Journal of Clinical Pharmacology.* 2009;68(6):936-947.
132. Sandars J, Esmail A. The frequency and nature of medical error in primary care: understanding the diversity across studies. *Fam Pract.* 2003;20(3):231-236.
133. Sayers YM, Armstrong P, Hanley K. Prescribing errors in general practice: A prospective study. *European Journal of General Practice.* 2009;15(2):81-83.
134. Schneider JK, Mion LC, Frengley JD. Adverse drug reactions in an elderly outpatient population. *Am J Hosp Pharm.* 1992;49(1):90-96.
135. Smith PC, Araya-Guerra R, Bublitz C, et al. Missing clinical information during primary care visits. *Journal of the American Medical Association.* 2005;293(5):565-571.
136. Smits M, Huibers L, Kerssemeijer B, de Feijter E, Wensing M, Giesen P. Patient safety in out-of-hours primary care: a review of patient records. *BMC Health Services Research.* 2010;10:335.
137. Straand J, Rokstad KS. Elderly patients in general practice: diagnoses, drugs and inappropriate prescriptions. A report from the More & Romsdal Prescription Study. *Family Practice.* 1999;16(4):380-388.
138. Tomlin A, Reith D, Dovey S, Tilyard M. Methods for retrospective detection of drug safety signals and adverse events in electronic general practice records. *Drug Saf.* 2012;35(9):733-743.
139. Tsang C, Majeed A, Aylin P. Routinely recorded patient safety events in primary care: a literature review. *Fam Pract.* 2012;29(1):8-15.
140. Tsang C, Bottle A, Majeed A, Aylin P. Adverse events recorded in English primary care: observational study using the General Practice Research Database. *Br J Gen Pract.* 2013;63(613):e534-542.
141. Van Der Hooft CS, Jong GW, Dieleman JP, et al. Inappropriate drug prescribing in older adults: The updated 2002 Beers criteria - A population-based cohort study. *British Journal of Clinical Pharmacology.* 2005;60(2):137-144.
142. Wahls TL, Cram PM. The frequency of missed test results and associated treatment delays in a highly computerized health system. *BMC Fam Pract.* 2007;8:32.
143. Weingart SN, Gandhi TK, Seger AC, et al. Patient-reported medication symptoms in primary care. *Archives of Internal Medicine.* 2005;165(2):234-240.
144. APEAS Study. Patient Safety in Primary Health Care. Madrid: Ministry of Health and Consumer Affairs, 2008.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

145. Aranaz-Andrés JM, Aibar C, Limón R, et al. A study of the prevalence of adverse events in primary care in Spain. *Eur J Public Health*. 2012;22(6):921-925.

146. Avery AJ, Ghaleb M, Barber N, et al. The prevalence and nature of prescribing and monitoring errors in English general practice: a retrospective case note review. *Br J Gen Pract*. 2013;63(613):e543-553.

147. Beyer M, Rohe J, Rusitska M, Blauth E, Gerlach FM. The German incident reporting system for general practice: Structure, first results. [German]. *Zeitschrift fur Allgemeinmedizin*. 2005;81(4):147-153.

148. Bhasale A. The wrong diagnosis: identifying causes of potentially adverse events in general practice using incident monitoring. *Fam Pract*. 1998;15(4):308-318.

149. De Wet C, Bowie P. The preliminary development and testing of a global trigger tool to detect incident and patient harm in primary-care records. *Postgraduate Medical Journal*. 2009;85(1002):176-180.

150. Elder NC, Vonder Meulen M, Cassedy A. The identification of medical incidents by family physicians during outpatient visits. *Annals of Family Medicine*. 2004;2(2):125-129.

151. Fernald DH, Pace WD, Harris DM, West DR, Main DS, Westfall JM. Event reporting to a primary care patient safety reporting system: a report from the ASIPS collaborative. *Annals of Family Medicine*. 2004;2(4):327-332.

152. Gaal S, Verstappen W, Wolters R, Lankveld H, van Weel C, Wensing M. Prevalence and consequences of patient safety incidents in general practice in the Netherlands: a retrospective medical record review study. *Implementation Science*. 2011;6(1):37.

153. Gurwitz JH, Field TS, Harrold LR, et al. Incidence and preventability of adverse drug events among older persons in the ambulatory setting. *JAMA*. 2003;289(9):1107-1117.

154. Hickner J, Zafar A, Kuo GM, et al. Field test results of a new ambulatory care Medication Incident and Adverse Drug Event Reporting System-MEADERS. *Ann Fam Med*. 2010;8(6):517-525.

155. Hoffmann B, Beyer M, Rohe J, Gensichen J, Gerlach FM. "Every incident counts": a web-based incident reporting and learning system for general practice. *Qual Saf Health Care*. 2008;17(4):307-312.

156. Kennedy AG, Littenberg B, Senders JW. Using nurses and office staff to report prescribing incidents in primary care. *Int J Qual Health Care*. 2008;20(4):238-245.

157. Khoja T, Neyaz Y, Qureshi NA, Magzoub MA, Haycox A, Walley T. Medication incidents in primary care in Riyadh city, Saudi Arabia. *Eastern Mediterranean Health Journal*. 2011;17(2):156-160.

158. Khoo EM, Lee WK, Sararaks S, et al. Medical incidents in primary care clinics--a cross sectional study. *BMC Fam Pract*. 2012;13:127.

159. Kingston-Reichers J, Ospina M, Jonsson E, Childs P, Mcleod L, Mazted J. Patient Safety in Primary care. Edmonton, AB: Canadian Patient Safety Institute and BC Patient Safety and Quality Council. 2010.

160. Kostopoulou O, Delaney B. Confidential reporting of patient safety events in primary care: results from a multilevel classification of cognitive and system factors. *Qual Saf Health Care*. 2007;16(2):95-100.


161. Kuo GM, Phillips RL, Graham D, Hickner JM. Medication incidents reported by USA family physicians and their office staff. *Qual Saf Health Care*. 2008;17(4):286-290.

162. Kuzel AJ, Woolf SH, Gilchrist VJ, et al. Patient reports of preventable problems and harms in primary health care. *Ann Fam Med*. 2004;2(4):333-340.




- 1  
2  
3 869 163. Makeham M, Dovey S, Runciman W, Larizgoitia I. Methods and measures used in primary  
4 870 care patient safety research.  
5 871 [www.who.int/patientsafety/research/methods\\_measures/primary\\_care\\_ps\\_research/en/index.ht](http://www.who.int/patientsafety/research/methods_measures/primary_care_ps_research/en/index.html)  
6 872 [ml](http://www.who.int/patientsafety/research/methods_measures/primary_care_ps_research/en/index.html) Accessed 1 January 2015.
- 7  
8 873 164. McKay J, Bradley N, Lough M, Bowie P. A review of significant events analysed in general  
9 874 practice: implications for the quality and safety of patient care. *BMC Family Practice*.  
10 875 2009;10:61.
- 11 876 165. McKay J, de Wet C, Kelly M, Bowie P. Applying the Trigger Review Method after a brief  
12 877 educational intervention: potential for teaching and improving safety in GP specialty training?  
13 878 *BMC Med Educ*. 2013;13:117.
- 14  
15 879 166. Murie J, McGhee C. Clinical governance in action. Assessing risk by analysing significant  
16 880 events in primary care. *Quality in Primary Care*. 2003;11(3):205-211.
- 17 881 167. O'Beirne M, Sterling PD, Zwicker K, Hebert P, Norton PG. Safety incidents in family medicine.  
18 882 *BMJ Qual Saf*. 2011;20(12):1005-1010.
- 19  
20 883 168. Pearson A, Aromataris A. *Patient Safety Primary care: a review of the literature*. Australian  
21 884 Commission on Safety and Quality in healthcare. 2010.  
22 885 [www.health.gov.au/internet/safety/publishing.nsf/Content/DBDB4EAE1386D1ABCA25775300](http://www.health.gov.au/internet/safety/publishing.nsf/Content/DBDB4EAE1386D1ABCA257753001ECA09/$File/26889-Literature-Review.PDF)  
23 886 [1ECA09/\\$File/26889-Literature-Review.PDF](http://www.health.gov.au/internet/safety/publishing.nsf/Content/DBDB4EAE1386D1ABCA257753001ECA09/$File/26889-Literature-Review.PDF) Accessed 1 January 2015.
- 24 887 169. Phillips Jr RL, Bartholomew LA, Dovey SM, et al. Learning from malpractice claims about  
25 888 negligent, adverse events in primary care in the United States. *Quality and Safety in Health*  
26 889 *Care*. 2004;13(2):121-126.
- 27  
28 890 170. Phillips RL, Dovey SM, Graham D, Elder NC, Hickner JM. Learning from different lenses:  
29 891 reports of medical incidents in primary care by clinicians, staff and patients. *J Patient Saf*.  
30 892 2006;2(3):140 -146.
- 31 893 171. Schneider JK, Mion LC, Frengley JD. Adverse drug reactions in an elderly outpatient  
32 894 population. *Am J Hosp Pharm*. 1992;49(1):90-96.
- 33  
34 895 172. Singh H, Giardina TD, Meyer AN, Forjuoh SN, Reis MD, Thomas EJ. Types and origins of  
35 896 diagnostic incidents in primary care settings. *JAMA Intern Med*. 2013;173(6):418-425.
- 36 897 173. Smits M, Huibers L, Kerssemeijer B, de Feijter E, Wensing M, Giesen P. Patient safety in out-  
37 898 of-hours primary care: a review of patient records. *BMC Health Services Research*.  
38 899 2010;10:335.
- 39 900 174. Statham MO, Sharma A, Pane AR. Misdiagnosis of acute eye diseases by primary health care  
40 901 providers: incidence and implications. *Medical Journal of Australia*. 2008;189(7):402-404.
- 41  
42 902  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 1: PRISMA diagram of studies included in the review



254x190mm (96 x 96 DPI)


Review Only



Note: the rate per 100 consultations, records, or people is provided as specified in the text next to the study (consultations / records / people). Details of the first author are provided to allow cross-checking to the study details in the online supplement. All studies were observational unless listed as a review. The quality rating is based on validated scales.

197x144mm (300 x 300 DPI)





Note: Details of the first author are provided to allow cross-checking to the individual study details in the online supplement. All studies were observational unless listed as a review. The quality rating is based on validated scales.

192x142mm (300 x 300 DPI)

Review Only

## How Safe is Primary Care? A Systematic Review

### ONLINE SUPPLEMENT: TABLES

**Table 1: Summary of studies about frequency of safety incidents in primary care**

Author	Study type	Number analyzed	Region	Type of incidents	Rate	Study quality
Bregnhøj L 2007 <sup>106</sup>	Observational	1621	OECD	Prescribing / medication	39.5 per 100 medicines	High
Gurwitz JH 2003 <sup>117</sup>	Observational	30397	OECD	Prescribing / medication	5 per 100 people	High
Honigman B 2001 <sup>66</sup>	Observational	25056	OECD	Prescribing / medication	5.5 per 100 people	High
Lund BC 2010 <sup>125</sup>	Observational	236	OECD	Prescribing / medication	14.4 per 100 people	High
Olaniyan JO 2014 <sup>82</sup>	Systematic review	33	OECD	Prescribing / medication	1 to 90 out of 100 prescriptions	High
Sandars J 2003 <sup>133</sup>	Systematic review	280	OECD	Prescribing / medication	11 per 100 prescriptions	High
Tsang C 2012 <sup>140</sup>	Systematic review	15	OECD	Prescribing / medication	0.49 per 100 people aged 65+	High
Abramson 2012 <sup>95</sup>	Observational	9385	OECD	Prescribing / medication	36.7 per 100 prescriptions	Moderate
Al Khaja KA 2011 <sup>81</sup>	Observational	2773	Other	Prescribing / medication	26.4 per 100 prescriptions	Moderate
Al Khaja KAJ 2007 <sup>65,98</sup>	Observational	2282	Other	Prescribing / medication	90.5 per 100 prescriptions	Moderate
Al Khaja KAJ 2010 <sup>99</sup>	Observational	86	Other	Prescribing / medication	87.2 per 100 prescriptions	Moderate
Avery AJ 2013 <sup>103</sup>	Observational	6048	OECD	Prescribing / medication	4.9 per 100 prescribed items	Moderate
Bradbury F 2004 <sup>104</sup>	Observational	8830	OECD	Prescribing / medication	16.6 per 100 people	Moderate
Bradley MC 2012 <sup>105</sup>	Observational	166108	OECD	Prescribing / medication	34 per 100 people	Moderate
Brekke M 2008 <sup>107</sup>	Observational	86000	OECD	Prescribing / medication	18.4 per 100 people	Moderate
Brenner S 2012	Observational	516	OECD	Prescribing / medication	15 per 100 records	Moderate
Clark RC 2007 <sup>109</sup>	Observational	127582000	OECD	Prescribing / medication	3.1 per 100 people	Moderate
De Wilde S 2007 <sup>111</sup>	Observational	171690	OECD	Prescribing / medication	32.2 per 100 people	Moderate
Field TS 2004 <sup>115</sup>	Observational	31757	OECD	Prescribing / medication	4.8 per 100 people	Moderate
Howard M 2004 <sup>118</sup>	Observational	777	OECD	Prescribing / medication	16.3 per 100 people	Moderate
Khoja T 2011 <sup>120</sup>	Observational	5299	Other	Prescribing / medication	18.7 per 100 prescriptions	Moderate
Khoo	Observational	1753	Other	Prescribing /	41.1 per 100	Moderate

Author	Study type	Number analyzed	Region	Type of incidents	Rate	Study quality
2012 <sup>122</sup>				medication	records	
Martinez Sanchez A 2011 <sup>71</sup>	Observational	23995	OECD	Prescribing / medication	1.5 per 100 prescriptions	Moderate
Ryan C 2009 <sup>132</sup>	Observational	500	OECD	Prescribing / medication	13 per 100 people	Moderate
Schneider JK 1992 <sup>135</sup>	Observational	463	OECD	Prescribing / medication	21 per 100 people	Moderate
Straand J 1999 <sup>138</sup>	Observational	16774	OECD	Prescribing / medication	13.5 per 100 prescriptions	Moderate
Tomlin A 2012 <sup>139</sup>	Observational	173478	OECD	Prescribing / medication	14.4 per 100 people	Moderate
Van Der Hooff CS 2005 <sup>142</sup>	Observational	25258	OECD	Prescribing / medication	20 per 100 older people	Moderate
Weingart SN 2005 <sup>144</sup>	Observational	661	OECD	Prescribing / medication	4.7 per 100 people	Moderate
Abramson EL 2013 <sup>96</sup>	Observational	1905	OECD	Prescribing / medication	3.8 per 100 prescriptions	Low
Khoja TA 1996 <sup>121</sup>	Observational	6350	Other	Prescribing / medication	11.6 per 100 prescriptions	Low
Koper 2013 <sup>123</sup>	Observational	169	OECD	Prescribing / medication	56.2 per 100 people	Low
Neville RG 1989 <sup>128</sup>	Observational	15916	OECD	Prescribing / medication	3.2 per 100 prescribed items	Low
Nicholson D 2006 <sup>129</sup>	Observational	24	OECD	Prescribing / medication	91.7 per 100 clinicians prescribing	Low
Paille F 1995 <sup>130</sup>	Observational	4080	OECD	Prescribing / medication	32.5 per 100 prescriptions	Low
Sayers YM 2009 <sup>134</sup>	Observational	3948	OECD	Prescribing / medication	12.4 per 100 prescriptions	Low
Apeas 2008 <sup>101,102</sup>	Observational	96047	OECD	Any safety incident	0.8 to 17.93 per 100 consultations	High
Makeham M 2008 <sup>126</sup>	Systematic review	49	OECD	Any safety incident	0.0004 to 24 per 100 consultations	High
Makeham MA 2006 <sup>70</sup>	Observational	166569	OECD	Any safety incident	0.2 per 100 people	High
De Wet C 2009 <sup>110</sup>	Observational	500	OECD	Any safety incident	9.4 per 100 records	Moderate
Elder NC 2004 <sup>113</sup>	Observational	351	OECD	Any safety incident	24 per 100 consultations	Moderate
Gaal S 2011 <sup>116</sup>	Observational	8401	OECD	Any safety incident	2.2 per 100 consultations	Moderate
Smits M 2010 <sup>137</sup>	Observational	145	OECD	Any safety incident	18.6 per 100 records	Moderate
Tsang C 2013 <sup>141</sup>	Observational	74763	OECD	Any safety incident	0.6 per 100 people	Moderate
Van Dulmen SA 2011 <sup>72</sup>	Observational	1000	OECD	Any safety incident	1.8 per 100 records	Moderate
Eggleton KS 2014 <sup>112</sup>	Observational	170	OECD	Any safety incident	7 per 100 consultations	Low
McKay J 2013 <sup>127</sup>	Observational	520	OECD	Any safety incident	15 out of 100 records	Low

Author	Study type	Number analyzed	Region	Type of incidents	Rate	Study quality
A-Elgayoum SME 2009 <sup>97</sup>	Observational	3203	Other	Diagnosis	70 per 100 blood smears	Moderate
Illboudo TP 2012 <sup>119</sup>	Observational	1331	Other	Diagnosis	94.1 per 100 people	Moderate
Leon AC 1999 <sup>124</sup>	Observational	1000	OECD	Diagnosis	16 out of 100 people	Moderate
Wahls TL 2007 <sup>143</sup>	Observational	9116	OECD	Diagnosis	0.7 per 100 people	Moderate
Casalino LP 2009 <sup>108</sup>	Observational	1889	OECD	Failure to notify of abnormal result	7.1 per 100 abnormal results	Moderate
Al-Agilly S 2007 <sup>100</sup>	Observational	258	OECD	Inaccurate records	27.5 per 100 records	Moderate
Farrow SC 1999 <sup>114</sup>	Observational	82	OECD	Infection control	24.4 per 100 practices	Low
Pandit NB 2008 <sup>131</sup>	Observational	182	Other	Injection practices	77 out of 100 clinics	Low
Smith PC 2005 <sup>136</sup>	Observational	1614	OECD	Missing information	13.6 per 100 consultations	Moderate

Note: Studies are arranged in order of type of incident, followed by study quality. Harm rates were devised by extracting figures directly from articles. No recalculations were performed. If studies used slightly different terminology to the definitions of severity of harm listed in Box 1, then the data were categorized into the Box 1 definitions based on descriptions in the articles themselves. For example, if an article stated that 10% of incidents resulted in hospitalization, this would be listed as a moderate harm. Death rates were reported separately where available.

**Table 2: Prescribing incidents in primary care per 100 prescriptions / people – results from 33 studies arranged in order of study quality**

Study	Study type	Prescribing incidents	Study quality
Bregnhøj <sup>106</sup>	Observational	39.5 / 100 medicines	High
Gurwitz <sup>117</sup>	Observational	5 / 100 people	High
Honigman <sup>66</sup>	Observational	5.5 / 100 people	High
Lund <sup>125</sup>	Observational	14.4 / 100 people	High
Olaniyan <sup>82</sup>	Systematic review	90 / 100 prescriptions	High
Sandars <sup>133</sup>	Systematic review	11 / 100 prescriptions	High
Tsang <sup>140</sup>	Systematic review	0.49 / 100 people	High
Abramson <sup>95</sup>	Observational	36.7 / 100 prescriptions	Moderate
Al Khaja <sup>65</sup>	Observational	90.5 / 100 prescriptions	Moderate
Al Khaja <sup>81</sup>	Observational	26.4 / 100 prescriptions	Moderate
Al Khaja <sup>98</sup>	Observational	87.2 / 100 prescriptions	Moderate
Avery <sup>94</sup>	Observational	4.9 / 100 prescribed items	Moderate
Bradbury <sup>104</sup>	Observational	16.6 / 100 people	Moderate
Bradley <sup>105</sup>	Observational	34 / 100 people	Moderate
Brekke <sup>107</sup>	Observational	18.4 / 100 people	Moderate
Clark <sup>108</sup>	Observational	3.1 / 100 people	Moderate
De Wilde <sup>111</sup>	Observational	32.2 / 100 people	Moderate
Field <sup>115</sup>	Observational	4.8 / 100 people	Moderate
Howard <sup>118</sup>	Observational	16.3 / 100 people	Moderate
Khoja <sup>120</sup>	Observational	18.7 / 100 prescriptions	Moderate
Khoo <sup>122</sup>	Observational	41.1 / 100 records	Moderate
Martinez Sanchez <sup>71</sup>	Observational	1.5 / 100 prescriptions	Moderate
Ryan <sup>132</sup>	Observational	13 / 100 people	Moderate
Schneider <sup>135</sup>	Observational	21 / 100 people	Moderate
Straand <sup>138</sup>	Observational	13.5 / 100 prescriptions	Moderate
Tomlin <sup>139</sup>	Observational	14.4 / 100 people	Moderate
Van Der Hooft <sup>142</sup>	Observational	20 / 100 people	Moderate
Weingart <sup>144</sup>	Observational	4.7 / 100 people	Moderate
Abramson <sup>95</sup>	Observational	3.8 / 100 prescriptions	Low
Khoja <sup>120</sup>	Observational	11.6 / 100 prescriptions	Low
Koper <sup>123</sup>	Observational	56.2 / 100 people	Low
Neville <sup>128</sup>	Observational	3.2 / 100 prescribed items	Low
Nicholson <sup>129</sup>	Observational	91.7 / 100 clinicians	Low

Note: the number per 100 prescriptions, medications prescribed, or people is provided as specified. Details of the first author are provided to allow cross-checking to the individual study. Details are in the online supplement.<sup>20</sup> The quality rating is based on validated scales.

**Table 3: Summary of studies about severity of harm from safety incidents in primary care**

Author	Study type	Number of events analyzed	Region	Severity of harm found in studies based on chart review	Severity of harm found in studies based on prescription reviews	Severity of harm found in studies based on incident reports / claims
Apeas 2008 <sup>145</sup>	Observational	1108	OECD			23.6% no harm 38.6% low harm 32% moderate harm 5.8% severe harm 70.2% preventable
Aranaz-Andrés JM 2012 <sup>146</sup>	Observational	773	OECD	5.9% severe harm 64.3% preventable		
Avery AJ 2013 <sup>147</sup>	Observational	6048	OECD		0.2% severe harm	
Beyer M 2005 <sup>148</sup>	Observational	85	OECD			34% temporary or permanent harm
Bhasale A 1998 <sup>149</sup>	Observational	142	OECD			42.3% no harm 25.4% low harm 10.6% moderate harm 8.5% severe harm 13.4% death
De Wet C 2009 <sup>150</sup>	Observational	500	OECD	82.9% low to moderate harm		
Dovey SM 2002 <sup>33</sup>	Observational	330	OECD			55.8% no harm 12.1% low harm 7.0% moderate harm 5.5% severe harm 0.3% death
Elder NC 2004 <sup>151</sup>	Observational	351	OECD			24% actual harm (severity not specified)
Fernald DH 2004 <sup>152</sup>	Observational	209	OECD			64.1% no harm 15.3% low harm 10.1% moderate harm 10.5% severe harm

Author	Study type	Number of events analyzed	Region	Severity of harm found in studies based on chart review	Severity of harm found in studies based on prescription reviews	Severity of harm found in studies based on incident reports / claims
Gaal S 2011 <sup>153</sup>	Observational	1000	OECD	38.6% no harm 50.5% low harm 6.9% moderate harm 4.0% 'unknown harm'		
Gurwitz JH 2003 <sup>154</sup>	Observational	30397	OECD			38% severe harm (serious, life threatening or fatal)
Hickner J 2010 <sup>155</sup>	Observational	507	OECD			1.6% severe harm
Hoffmann B 2008 <sup>156</sup>	Observational	78	OECD			3.9% no harm 25.6% actual harm (severity unknown)
Kennedy AG 2008 <sup>157</sup>	Observational	216	OECD			90% no harm 8.7% low harm 1.8% moderate harm 0% severe harm
Khoja T 2011 <sup>158</sup>	Observational	5299	Other		53.2% no harm 8.7% low harm 37.3% moderate harm 0.8% severe harm	
Khoo 2012 <sup>159</sup>	Observational	1753	Other	39.9% severe harm 93.5% preventable		
Kingston-Reichers J 2010 <sup>160</sup>	Systematic review	Review	OECD	9% to 52% harm (severity not specified) 42% to 83% preventable		
Kostopoulou O 2007 <sup>161</sup>	Observational	78	OECD			3.9% no harm 16.7% severe harm or death
Kuo GM 2008 <sup>162</sup>	Observational	194	OECD			41% no harm 35% low harm 21% moderate harm 3% severe harm



Author	Study type	Number of events analyzed	Region	Severity of harm found in studies based on chart review	Severity of harm found in studies based on prescription reviews	Severity of harm found in studies based on incident reports / claims
Kuzel AJ 2004 <sup>163</sup>	Observational	170	OECD			76.9% harm (severity unknown)
Makeham M 2008 <sup>164</sup>	Systematic review	Review	OECD	17% to 39% harm (severity not specified) 0% to 4% severe harm 45% to 76% preventable		
Makeham MA 2006 <sup>70</sup>	Observational	166569	OECD			0.25% harm (severity not specified)
McKay J 2009 <sup>165</sup>	Observational	191	OECD			57.1% no harm 7.3% low harm 11.5% moderate harm 4.7% severe harm 1.6% death 17.8% not classified
McKay J 2013 <sup>166</sup>	Observational	520	OECD	44% moderate to severe harm 45% preventable		
Murie J 2003 <sup>167</sup>	Observational	55	OECD			47.3% no harm 9.1% low harm 3.6% moderate harm 18.2% severe harm 21.8% death
O'Beirne M, 2013 <sup>168</sup>	Observational	264	OECD			50% harm (any type) 1% severe harm 93% preventable
Pearson A 2009 <sup>169</sup>	Systematic review	Review	OECD	24% to 42% harm (severity not specified)		
Phillips Jr RL 2004 <sup>170</sup>	Observational	26126	OECD			18.7% low harm 26.0% moderate harm 19.0% severe harm 36.3% death

Author	Study type	Number of events analyzed	Region	Severity of harm found in studies based on chart review	Severity of harm found in studies based on prescription reviews	Severity of harm found in studies based on incident reports / claims
Phillips RL 2006 <sup>171</sup>	Observational	701	OECD			20.7% no harm 30.0% low harm 30.0% moderate harm 14.3% severe harm 8.1% death
Schneider JK 1992 <sup>172</sup>	Observational	332	OECD	3.6% severe harm		
Singh H 2013 <sup>173</sup>	Observational	190	OECD	86% moderate or severe harm		
Smits M 2010 <sup>174</sup>	Observational	145	OECD	29.6% moderate harm 0% death		
Statham MO 2008 <sup>175</sup>	Observational	123	OECD			51.2% low harm 39.8% moderate harm 8.9% severe harm 11.6% preventable

## Online Supplement

### **Summary of studies included in the review**

This supplement provides a summary of each of the studies identified for the review and their quality (rated using validated appraisal tools).

For systematic reviews, quality was assessed using the Critical Appraisal Skills Programme (CASP) tool. For observational (epidemiological and other) studies, quality was assessed using the Evidence Based Library and Information Practice (EBLIP) Critical Appraisal Checklist.

The table is ordered alphabetically.

Findings are listed in terms of:

- the number of safety incidents
- the type of safety incidents
- the level of harm associated with incidents

Not all studies included findings in each of these areas, so parts of the table are left blank if there were no relevant key findings.

### **Studies included in systematic reviews**

Studies summarized in systematic reviews identified as part of this review are listed in alphabetical order. This is to illustrate the range of studies drawn on, even if the original primary studies were not incorporated into the narrative synthesis for our review.

### **Online content**

The final page of this document lists where the full protocol can be found online (including the full search terms and search strategy) as well as where freely available online content can be sourced.

Online table 1: Summary of key findings of studies included in the review

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Abramson 2012 <sup>1</sup>	Observational	Primary care clinic	USA	Review of prescriptions	Reviewed 9385 prescriptions from 78 providers and found 36.7 prescribing incidents per 100 prescriptions (95% CI 30.7 to 44.0), excluding incidents associated with legibility. Non-legibility incidents were found in 175.0 per 100 prescriptions (95% CI 169.1 to 181.3), inappropriate abbreviation incidents in 13.4 and direction incidents in 4.2 per 100 prescriptions. The majority of incidents were judged to be preventable.			Moderate
Abramson EL 2013 <sup>2</sup>	Observational	Primary care	USA	Record review	Chart review of patient notes from 16 clinicians over a three month period analysed 1905 prescriptions. The prescribing incident rate was 3.8 per 100 prescriptions (95% CI 2.8 to 5.1).			Low
A-Elgayoum SME 2009 <sup>3</sup>	Observational	Primary care clinic	Sudan	Test review	3203 blood smears from patients clinically suspected to have malaria were examined. Of these 2253/3203 (70.3%) were found to be a misdiagnosis.			Moderate
Al Khaja KA 2011 <sup>4</sup>	Observational	Primary care clinics and pharmacy	Bahrain	Review of prescriptions	Audited prescriptions issued by 20 primary care centers and found 733/2773 (26.4%) medical prescribing incidents.			Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Al Khaja KAJ 2007 <sup>5</sup>	Observational	Primary care clinics	Bahrain	Review of prescriptions		Incidents in prescriptions included 4972/7139 (69.7%) incidents of omission such as strength/dose: 573/4972 (11.5%); dosage form: 2815/4972 (56.6%); dosage frequency: 261/4972 (5.3%); and length of treatment: 1323/4972 (26.6%). Incidents of commission accounted for 1759/7139 (24.6%) of incidents in prescriptions. These included dosage form: 159/1759 (9.0%); dosage frequency: 105/1759 (6.0%); and length of treatment: 1419/1759 (80.7%).		Moderate
Al Khaja KAJ 2007 <sup>6</sup>	Observational	Primary care clinic	Bahrain	Review of prescriptions	2282 prescriptions dispensed for infants for corticosteroids were analyzed. The frequency of dosing and length of therapy were not stated in 21.6% and 43.6% of prescriptions, respectively.			Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Al Khaja KAJ 2007 <sup>7</sup>	Observational	Primary care clinic	Bahrain	Review of prescriptions	Reviewed prescriptions for infants. Drug-related incidents were present in 2066/2282 (90.5%) of the prescriptions reviewed. 5745 prescribed drug items were reviewed. There were 4282/5745 (74.5%) drug-related incidents.	Incidents of omission accounted for 4146/5745 (72.2%) of drug-related incidents. The breakdown of this category was: strength/dose: 424/4146 (10.2%); dosage form: 979/4146 (23.6%); dosage frequency: 621/4146 (15.0%); and length of treatment: 2122/4146 (51.2%). Incidents of commission accounted for 3338/5745 (58.1%). The breakdown of this category was: dosage form: 1354/3338 (40.6%); dosage frequency: 4/3338 (0.1%); length of treatment: 1594/3338 (47.8%); and incidents of commission: 386/3338 (11.6%). Incidents of integration accounted for 183/5745 (3.2%).		Moderate
Al Khaja KAJ 2010 <sup>8</sup>	Observational	Primary care clinics and pharmacy	Bahrain	Review of prescriptions	Audited prescriptions issued by 20 primary care centers for pediatric iron preparations. Found 75/86 (87.2%) of prescriptions contained omission incidents.	Omission incidents were subcategorized as: unclear names of iron preps: 15/75 (20.0%); prescriptions without specifying the daily dosage of iron: 4/75 (5.3%); prescriptions without stating the daily dosage of iron: 42/75 (56.0%); and prescriptions without specifying the duration of therapy: 14/75 (18.7%).		Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Al-Agilly S 2007 <sup>9</sup>	Observational	General practice	UK	Record review	Patients were invited to check their records. 187 out of 258 patients (72.5%) said their records were accurate. There were 89 inaccuracies reported by patients. 42 (47%) were of obvious clinical importance including wrongly entered diagnoses, or missing major morbidity such as an operation, or incidents in repeat medication. There were 47 (53%) inaccuracies in lifestyle data (smoking, alcohol history or weight), or dates of illnesses.			Moderate
Apeas 2008 <sup>10</sup>	Observational	Primary care clinic	Spain	Incident reporting system	Of 96,047 visits, the health care professional detected some possible adverse effect in 1,932 visits, generating a total of 2,059 reports. A total of 1,074 adverse effects were identified corresponding to 971 different patients. The prevalence of adverse effects per visit was 11.18% (1,074/96,047, 95% CI 10.52% to 11.85%). The prevalence of visits which experience some adverse event was 17.93% (1,722/96,047, 95% CI 17.09% to 18.77%).	The causes of these incidents included medication (534/1180, 45.3%), provision of care (285/1080, 26.4%), communication (273/1080, 25.3%), diagnosis (159/1080, 14.7%) and management (99/1080, 9.2%).	In 23.6% of the cases, the consequences of the adverse effect did not affect the care provided, in 33.1% a higher level of observation and monitoring were required, in the remaining 7.5%, the adverse effect required an additional test and in 17.1%, an additional medical or surgical treatment was performed. In 14.9%, the consequence of the adverse effect required another visit or referral to specialized care (without hospitalization), and in 5.8% hospitalization of the patients for some life support treatment was required. 778 / 1108 (70.2%) adverse events were deemed to be preventable and 330/1108 (29.8%) could not have been prevented.	High



First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Aranaz-Andrés JM 2012 <sup>11</sup>	Observational	General practice	Spain	Record review and survey	Data were collected for 52 healthcare professionals who attended 96,047 consultations. 773 adverse events were identified, giving a point prevalence of 0.8% (95% CI 0.76 to 0.85).	55.5% (429) of the AE stemmed from problems with the medication prescribed; 17.1% (132) involved a worsening of the clinical course of the underlying disease; 7.8% (60) involved complications from a medical procedure; 7.4% (57) involved health care-related infection; and 6.1% (50) stemmed from problems with the care dispensed (wound cures, catheter care, etc).	Most adverse events (64.3%) were considered preventable and only 5.9% were severe, usually related to medication (odds ratio 4.6; 95% CI 2.1 to 10.3).	Low
Avery AJ 2013 <sup>12</sup>	Observational	General practice	UK	Record review	6048 unique items prescribed over a 12 month period for 1777 patients were examined. 4.9% of all prescription items had a prescribing or monitoring incident (95% CI 4.4% to 5.5%).		Most incidents were of mild to moderate severity. 0.2% were classed as severe. Children under 15 years, adults over 64 years and people with higher numbers of unique medication items prescribed were at greater risk of incidents.	Moderate
Beyer M 2003 <sup>13</sup>	Observational	General Practice	Germany	Incident reporting system		Examined 51 medication incidents. These were subcategorized as follows: failure to give medicine: 3/51 (5.9%); wrong medication: 8/51 (15.7%); wrong dose: 9/51 (17.7%); drug omitted: 1/51 (2.0%); adverse events: 6/51 (11.8%); drug interactions: 6/51 (11.8%); other: 10/51 (19.6%); no indication: 8/51 (15.7%).		Low

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Beyer M 2005 <sup>14</sup>	Observational	General practice	Germany	Incident reporting system		Analyzed 85 incident reports. These included wrong diagnosis: 17/85 (20.0%); wrong clinical decision: 9/85 (11.0%); not applicable: 1/85 (1.0%); administration incidents: 9/85 (11.0%); investigation incidents: 2/85 (2.0%); treatment incidents: 23/85 (27.0%); communication incidents : 18/85 (21.0%); wrong payment: 4/85 (5.0%); wrong treatment: 2/85 (2.0%).	34% of incidents caused temporary or permanent harm.	Moderate
Bhasale A 1998 <sup>15</sup>	Observational	General practice	Australia	Incident reporting system		Types of diagnostic incident included: missed diagnosis: 59/142 (41.6%); misdiagnosis: 38/142 (26.8%); delayed diagnosis: 31/142 (21.8%); and diagnostic procedural complication: 18/142 (12.7%). Contributing factors were listed as: incident in judgment: 63/142 (44.4%); failure in recognizing signs and symptoms: 57/142 (40.1%); poor communication between patient and health team: 33/142 (23.2%); poor communication between health professionals: 30/142 (21.1%); inappropriate action of others: 30/142 (21.1%); inappropriate patient assessment: 27/142 (19.0%); inappropriate follow up of patient: 19/142 (13.4%); and inappropriate second opinion : 33/142 (23.2%).	60/142 (42.3%) patients suffered no harm; 36/142 (25.4%) suffered low harm; 15/142 (10.6%) suffered moderate harm; 12/142 (8.5%) suffered severe harm; and that 19/142 (13.4%) patients died.	Moderate
Bradbury F 2004 <sup>16</sup>	Observational	General practice	Ireland	Record review	Of the 8830 patient records reviewed for people using NSAIDs, 1462/8830 (16.6%) showed evidence of an adverse drug reaction.			Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Bradley MC 2012 <sup>17</sup>	Observational	Primary care	UK	Database	The overall prevalence of potentially inappropriate prescribing in over 70 year olds (n =166,108) was 34%.			Moderate
Bregnhøj L 2007 <sup>18</sup>	Observational	General Practice	Denmark	Records review	212 older people with polypharmacy were prescribed 1621 medications by their GPs. 640/1621 (39.5%) of the medications were likely to be inappropriate. Most of the patients (200/212, 94.3%) had one or more inappropriate ratings among their medications.			High
Brekke M 2008 <sup>19</sup>	Observational	General practice	Norway	Record review	15,790/86,000 (18.4%) elderly people received one or more potentially harmful medications / inappropriate prescriptions from their GP.			Moderate
Brenner S 2012 <sup>20</sup>	Observational	Primary and urgent care	US	Record review	A trigger tool identified that 15% of patients had adverse drug events.	54% of adverse drug events occurred during medication monitoring, and 45% during patient self-administration.		Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Britt H 1997 <sup>21</sup>	Observational	General practice	Australia	Incident reporting system		Examination of 500 incident reports found pharmacological treatments: 259/500 (51.8%); non-pharmacological treatments: 183/500 (36.5%); diagnostic incidents: 142/500 (28.3%); equipment incidents: 26/500 (5.2%); poor communication: 130/500 (26.0%); incident in judgment: 120/500 (24.0%); action of others: 120/500 (24.0%); poor communication between health professionals: 100/500 (20.0%); patient consulted other medical officer: 80/500 (16.0%); failure to recognize symptoms: 70/500 (14.0%); patients history not adequately reviewed: 55/500 (11.0); omission of checking procedure: 55/500 (11.0%); and GP tired/rushed/running late: 50/500 (10.0%).		Moderate
Casalingo LP 2009 <sup>22</sup>	Observational	Primary care	USA	Record review	Retrospective medical record review of 5434 randomly selected patients aged 50 to 69 years in 19 community-based and 4 academic medical center primary care practices found the rate of apparent failures to inform people of an abnormal test result or to document informing the patient was 7.1% (135 failures / 1889 abnormal results), with a range of 0% to 26.2%.			Moderate
Clark RC 2007 <sup>23</sup>	Observational	General practice	UK	Record review	127,582,000 patient records were reviewed and adverse drug reactions were found in 3,968,000/127,582,000 (3.1%) cases.			Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Cox SJ 2004 <sup>24</sup>	Observational	General practice	UK	Significant Event Audit		This study analyzed significant events. Administrative incidents made up the highest proportion of events: 95/337 (28.2%). Other events were categorized as follows: other medical: 70/337 (20.8%); prescribing-related events: 46/337 (13.7%); missed new cancer diagnosis: 32/337 (9.5%); other: 28/337 (8.31%); nursing-related events: 23/337 (6.8%); patient complaint: 17/337 (5.0%); avoidable death where terminal care took place at home: 15/337 (4.5%); section under Mental Health Act: 8/337 (2.4%); and suicide: 3/337 (0.89%).		Low
De Wet C 2009 <sup>25</sup>	Observational	General practice	UK	Record review	Review of 500 records found an adverse event in 47 records (9.4%), indicating that harm occurred at a rate of one event per 48 consultations. A further 17 records (3.4%) contained evidence of a potential adverse event.	Incident and harm rates were higher in those aged >60 years, and most were medication-related (59%).	Harm severity was low to moderate for most patients (82.9%).	Moderate
De Wilde S 2007 <sup>26</sup>	Observational	General practice	UK	Record review	Looked at 171,690 records of elderly people and found 55,325 / 171,690 (32.2%) patients received potentially inappropriate medications.			Moderate
Diamond MR 1995 <sup>27</sup>	Observational	General practice	Australia	Interviews		Interviews with 39 trainees in general practice found that of 180 incident reports 70/180 (38.9%) were related to communication; 17/180 (9.4%) were related to management; 56/180 31.11% to diagnosis; 17/180 9.44% to administration; and 20/180 11.11% were classified as 'other'.		Low

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Dovey SM 2002 <sup>28</sup>	Observational	Family practice	USA	Incident reporting system		Knowledge and skills incidents made up 46/330 (13.9%) of medical incidents. Process incidents accounted for 284/330 (86.1%) of medical incidents. The process incidents were further broken down as: office administration: 102/284 (35.9%); investigations: 82/284 (28.9%); treatment 76/284 (26.8%); communication: 19/284 (6.7%); and insurance-related incidents: 1/284 (0.4%). When looking at incident reports, process incidents made up 284/330 (86.1%) of incident reports. The process incidents were further broken down as: office administration: 102/284 (35.9%); investigations: 82/284 (28.9%); other investigations: 6/284 (2.11%); treatments: 76/284 (26.8%); communication: 19/284 (6.7%); payment: 4/284 (1.4%). Knowledge and skill incidents made up 46/330 (13.9%) of incident reports.	184/330 (55.8 %) incident reports recorded that the patient suffered no harm; 40/330 (12.1%) reports showed that the patient suffered low harm; 23/330 (7.0%) reports showed that the patient suffered moderate harm; 18/330 (5.5%) reports showed that the patient suffered severe harm; and 1/330 (0.3%) report recorded the death of a patient.	Moderate
Eggleton KS 2014 <sup>29</sup>	Observational	General practice	New Zealand	Record review	Harm was identified in 46 out of 170 patient records (27%). This equates to 7 occurrences of harm per 100 consultations.	All harms related to medication use.		Low

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Elder NC 2002 <sup>30</sup>	Systematic review	Primary care	USA	Various		Four studies described medical incidents and adverse events in primary care, and three other studies peripherally addressed primary care medical incidents. There were three main types of preventable adverse events: diagnosis, treatment, and preventive services. Process incidents were classified into four categories: clinician, communication, administration and blunt end.		High
Elder NC 2004 <sup>31</sup>	Observational	Family practice	USA	Survey	Fifteen physicians in 7 practices completed forms for 351 visits. Incidents and preventable adverse events were identified in 24% of these visits. There was wide variation in how often individual physicians identified incidents (3% to 60% of visits).	57/351 (16.2%) of the reports related to office administration incidents. Of these, 37/57 (64.9%) related to charting; and general office administration accounted for 21/57 (36.8%) incidents. Physician-related incidents accounted for 28/351 (8.0%); patient communication incidents accounted for 16/351; and 15/351 (4.3%) related to preventable adverse events.	Harm was believed to have occurred as a result of 24% of the incidents, and was a potential in another 70%.	Moderate
Ely JW 2012 <sup>32</sup>	Observational	Primary care	USA	Survey		200 family physicians, general internists and general pediatricians were surveyed about diagnostic incidents. Common presenting complaints included abdominal pain (n = 27 of 202 patients, 13%); fever (n =19; 9%); and fatigue (n = 15, 7%).		Low



First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Farrow SC 1999 <sup>33</sup>	Observational	General practice	UK	Survey	This study looked at infection control in general practice. In the 82 practices, it found the following issues. Failure to access an autoclave: 56/82(68.3%); failure to have an autoclave with maintenance contract: 34/82 (41.5%); lack of access to Central Sterile Supply Department: 16/82 (19.5%); ineffective decontamination: 20/82 (24.4%); and inappropriate use of chemical disinfectants: 33/82 (40.24%). It also found a lack of hepatitis B vaccination in exposed staff in 31/82 (37.8%) practices; no infection control guidelines/sharps injury protocols in 60/82 (73.2%) practices; and a lack of training in 54/82 (65.85%) practices.			Low

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Fernald DH 2004 <sup>34</sup>	Observational	Primary care	USA	Incident reporting system		Examined reports to primary care incident report system. Diagnostic testing incidents accounted for 325/708 (45.0%) of all incident reports. This was further broken down into: blood test: 129/325 (39.7%); other specimen: 67/325 (20.6%); imaging: 58/325 (17.9%); and 'other or unspecified test': 31/325 (9.5%). Medication incidents accounted for 165/708 (23.3%) of all incident reports. This category was further categorized as follows: wrong drug: 32/165 (19.4%); right drug: wrong dose or timing: 99/ 165 (60.0%); right drug: wrong administration or dispensing: 39/165 (23.6%); drugs not prescribed: 6/165 (3.64%). Communication incidents accounted for 437/708 (61.7%) of all incident reports.	134/209 patients (64.1%) suffered no harm; 32/209 (15.3%) suffered low harm; 21/209 patients (10.1%) suffered moderate harm; and 22/209 patients (10.5%) suffered severe harm.	High
Field TS 2004 <sup>35</sup>	Observational	Primary care clinic	USA	Record review and incident reports	Among 31,757 older people, 1,523 adverse drug events were identified, a rate of 48.0 per 1,000 person-years. The rate of preventable adverse drug events was 13.3 per 1,000 person-years.			Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Gaal S 2011 <sup>36</sup>	Observational	General practice	Netherlands	Record review	A random sample of 1,000 patients from 20 general practices was obtained. The 1,000 patient records included a total of 8,401 patient contacts with the practice. A total of 211 patient safety incidents were identified (95% CI 185 to 241). These incidents concerned 186 patients. In other words, a total of 1 to 4 patient safety incidents per patient were detected per year for a prevalence of 2.2% for all patient contacts (186/8401).	116/211 (55.0%) adverse events were related to poor organization. 31/211 (14.7%) were related to treatments; 26/211 (12.3%) were related to communication issues; 21/211 (10.0%) were related to diagnosis; 14/211 (6.6%) were categorized as 'prevention'; and triage accounted for 3/211 (1.4%) adverse events. In terms of cause, 4/358 (1.1%) were categorized as technical; 176/358 (49.2%) as human; 97/358 (27.1%) as organizational; and 81/358 (22.6%) as patient-related.	58 out of 211 patient safety incidents affected patients; seven were associated with hospital admission; none resulted in permanent disability or death. 51/101 (50.5%) of patients suffered low harm; 39/101 (38.6%) suffered no harm; 7/101 (6.9%) suffered moderate harm; and 4/101 (4.0%) were categorized as 'unknown harm'.	Moderate
Garfield S 2009 <sup>37</sup>	Systematic review	Primary care	UK	Various		Review of 27 papers to map the medicines management system in primary care in the UK. The proportion of incidents in managing medicines in primary care was high. Several stages of the process had incident rates of 50% or more: repeat prescribing reviews, interface prescribing and communication and patient adherence.		High

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Gehring 2012 <sup>38</sup>	Observational	General practice	Switzerland	Survey		Cross-sectional survey of 630 nurses and doctors in 472 practices asking about the occurrence of patient safety incidents and analyzed data from the Swiss primary care critical incident reporting systems. The incidents with highest frequency were incomplete, unavailable, unclear, or incorrect patient documentation (88.4%); missing, incomplete, or erroneous information from external providers (81.7%); and required medication was not prescribed, administered, or dispensed (80.6%).		Low
Gurwitz JH 2003 <sup>39</sup>	Observational	Ambulatory care clinic (including primary care)	USA	Record review and incident reports	In 30,397 older person-years there were 1523 adverse drug events. The overall rate of adverse drug events among older people was 50.1 per 1000 person-years, with a rate of 13.8 preventable adverse drug events per 1000 person-years.	Incidents associated with preventable drug events were most often related to prescribing and monitoring.	38% (578) of drug events were categorized as serious, life threatening or fatal.	High
Hansen LB 2006 <sup>40</sup>	Observational	Primary care	USA	Survey		22 practices recorded 567 clarification calls over a two week period from pharmacies, most frequently for prior authorization issues (n = 209; 37%), formulary issues (n = 148; 26%), and unclear/missing prescription dosages (n = 117; 21%). Drug classes most frequently requiring clarifications were gastrointestinal (n = 122; 21.7%), cardiovascular (n = 278; 13.9%), and analgesic / anesthetic (n = 74; 13.2%) agents.		Low

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Hickner J 2008 <sup>41</sup>	Observational	Family practice	USA	Incident reporting system		Examined 590 event reports of 966 testing process incidents. Incidents occurred in ordering tests (12.9%), implementing tests (17.9%), reporting results to clinicians (24.6%), clinicians responding to results (6.6%), notifying patient of results (6.8%), general administration (17.6%), communication (5.7%) and other categories (7.8%). Charting or filing incidents accounted for 14.5% of incidents.		Moderate
Hickner J 2010 <sup>42</sup>	Observational	Primary care clinic	USA	Incident reporting system		507 anonymous event reports were submitted by 24 practices. Of these reports, 357 (70%) included medication incidents only, 138 (27%) involved adverse drug events only, and 12 (2.4%) included both.	Eight (1.6%) of the reported events led to hospitalization.	High
Hildebrandt DE 2006 <sup>43</sup>	Observational	Family practice	USA	Record review			Examined level of harm suffered when people were not appropriately triaged when they rang primary care for help out of hours. 31/119 (26.1%) patients suffered low harm (discomfort due to pain); and 4/119 (3.4%) patients suffered moderate harm.	Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Hoffmann B 2008 <sup>44</sup>	Observational	General practice	Germany	Incident reporting system		Most of the incident reports (52/78, 66.7%) – related to ‘appropriate care obstructed or delayed/inappropriate care provided’. Within this category, ‘prescribing or medication review’ accounted for 11/52 (21.2%) reports; 10/52 (19.2%) reports related to ‘delays or inappropriate care in hospital. ‘Lack of information, communication failures’ were responsible for 7/52 (13.5%) reports. The rest of the breakdown is as follows: dealing with test results or hospital correspondence: 5/52 (9.6%) reports; referrals (delayed/forgotten): 5/52 (9.6%) reports; vaccination/ drug administration: 4/52 (7.7%) reports; judging urgency of patient’s condition: 2/52 (3.9%) reports; external factors/equipment failures: 3/52 (5.8%) reports; failing to home visit: 2/52 (3.85%); dispensing incidents: 2/52 (3.9%).	‘Actual or potential consequence’ related to 20/78 (25.6%) incident reports. ‘No apparent potential for harm to patients’ related to 3/78 (3.9%) reports; and 3/78 (3.9%) were labelled as ‘other’.	Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Holden J 1998 <sup>45</sup>	Observational	General practice	UK	Record review		Audit of 1263 deaths found that GP factors occurred in 65/1263 (5.2%) cases. Within this category, the highest proportion was 25/65 (38.5%) reports related to delayed referral, diagnosis, treatment. This was followed by non-prescription of aspirin: 22/65 (3.9%) reports; failure to check/control blood pressure: 12/65 (18.5%) reports; side-effects from aspirin: 3/65 (4.6%) reports; poor diabetic control: 1/65 (1.5%) report; failure to challenge hospital drug treatment: 1/65 (1.5%) report; and failure to treat osteoporosis: 1/65 (1.5%)		Moderate
Honigman B 2001 <sup>46</sup>	Observational	Primary and ambulatory care	USA	Record review	There were adverse drug events in 864/25056 (3.5%) of cases. The adverse drug event rate was rate was 5.5 (95% CI 5.2 to 5.9) per 100 patients coming for care.			High
Howard M 2004 <sup>47</sup>	Observational	Family practice	Canada	Record review	127/777 (16.3%) older people were prescribed one or more potentially inappropriate medications.			Moderate



First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Illboudo TP, 2012 <sup>48</sup>	Observational	Primary care clinic	Burkina Faso	Record review	This study assessed the correctness of diagnoses from 12 health centers among patients with severe malaria and pneumonia. Among the 457 malaria cases affecting children under five, 66 cases (14.4%) were correctly diagnosed and of those 66 correctly diagnosed cases, 40 cases (60.6%) received an appropriate referral decision from their providers. Among the adult pneumonia cases, 5.9% (79/1331) of the diagnoses were correctly diagnosed; however, the appropriateness rate of the provider's referral decision was 98.7% (78/79).			Moderate
Kennedy AG 2008 <sup>49</sup>	Observational	Primary care	USA	Incident reporting system			216 incident reports were submitted. Nearly 90% (142/165) were incidents that did not reach the patient (low severity). Nineteen incidents reached the patient without causing harm (8.7%) and 4 incidents caused temporary harm requiring intervention (1.8%)..	Moderate
Khoja T 2011 <sup>50</sup>	Observational	Primary care	Saudi Arabia	Review of prescriptions	This study looked at 5299 prescriptions and found prescription incidents in 990/5299 (18.7%) of cases.		8/990 (0.8%) were classified as serious, and were potentially life-threatening incidents; 369/990 (37.3%) were classified as 'major nuisance'; 86/990 (8.7%) were classified as 'minor nuisance'; and 527/990 were classified as trivial (53.2%).	Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Khoja TA 1996 <sup>51</sup>	Observational	Primary care	Saudi Arabia	Review of prescriptions	Out of 6350 prescriptions, 11.6% were found to have at least one incident.	Of 20320 individual drug entries, incidents included strength not stated: 772/20320 (3.8%); wrong dose: 329/20320 (1.6%); tablet instead of capsule: 197/20320 (1.0%); quantity and/or frequency missing: 652/20320 (3.2%); drug interactions: 24/20320 (0.1%); contraindication: 59/20320 (0.3%); wrong drug: 193/20320 (1.0%); incomplete drug: 136/20320 (0.7%); frequency of the daily dose missing: 18186/20320 (89.5%); generic name of the drug missing: 17475/20320 (86.0%); strength missing: 15504/20320 (76.3%); illegible/incomplete personal and diagnostic data: 12801/20320 (63.0%); duration of treatment missing: 7681/20320 (37.8%); poor/fair handwriting and abbreviations: 3861/20320 (19.0%); very poor handwriting and abbreviations: 3658/20320 (18.0%); and form of the drug missing: 2723/20320 (13.4%).		Low

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Khoo 2012 <sup>52</sup>	Observational	Primary care clinic	Malaysia	Record review	A sample of 1753 medical records from 12 primary care clinics were scanned for diagnostic, management and documentation incidents. They found that 3.6% of medical records contained diagnostic incidents (95% CI 2.2 to 5.0), 53.2% contained management incidents (95% CI 46.3 to 60.2), 41.1% contained medication incidents (95% CI 35.8 to 46.4), 21.7% contained investigation incidents (95% CI 16.5 to 26.8), and 14.5% contained decision-making incidents (95% CI 10.8 to 18.2).		39.9% (95% CI 33.1 to 46.7) of incidents were potentially seriously harmful and nearly all (93.5%) were considered preventable.	Moderate
Kingston-Reichers J 2010 <sup>53</sup>	Systematic review	Primary care	Canada	Various		Two major forms of harm were missed or delayed diagnosis and medication harm.	Included 46 studies. Proportion of incidents associated with harm was between 9% and 52% in individual studies. Between 42% and 83% were thought to be preventable.	High
Koper 2013 <sup>54</sup>	Observational	General practice	Austria	Review of prescriptions	Examined medications of 169 patients with polypharmacy in 22 GP clinics. 74 patients (56.2%) had at least one dosing incident, four patients (2.4%) had at least one interaction in the most severe category. 158 patients were elderly (≥65 years), and of these 37.3% (n = 59) had at least one potentially inappropriate medication for the elderly.			Low

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Koper D 2013 <sup>55</sup>	Observational	General practice	Austria	Record review	The medications of 169 patients with polypharmacy treated in 22 GP surgeries were analysed. Patients took an average of nine medicines each day. At least one dosing incident was found in 56% of patients.			Low
Kostopoulou O 2007 <sup>56</sup>	Observational	General practice	UK	Incident reporting system		'Appropriate care obstructed or delayed/inappropriate care provided' accounted for 52/78 (66.7%) reports.	78 incident reports were relevant to patient safety and analysable. They included 21 (27%) adverse events and 50 (64%) near misses. 16.7% (13/71) had serious patient consequences, including one death. 75.7% (59/78) had the potential for serious patient harm. 'No apparent potential for harm to patients' accounted for 3/78 (3.9%) incident reports.	Moderate
Kuo GM 2008 <sup>57</sup>	Observational	Family practice	USA	Incident reporting system		126 out of 194 (70%) medication incidents were prescribing incidents, 17 (10%) were medication administration incidents, 17 (10%) documentation incidents, 13 (7%) dispensing incidents and 5 (3%) were monitoring incidents. Adverse drug events resulted from 16% of reported medication incidents.	Harm from reported incidents was categorised as: prevented and did not reach patients, (72, 41%), reached patients but did not require monitoring (63, 35%), reached patients and required monitoring (15, 8%), reached patients and required intervention (23, 13%) and reached patients and resulted in hospitalisation (5, 3%). No deaths were reported.	Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Kuzel AJ 2004 <sup>58</sup>	Observational	Primary care	USA	Interviews		People identified 221 events. The highest proportion of adverse events reported were related to breakdowns in the clinician-patient relationship: 82/221 (37.1%). This was followed by access breakdown: 63/221 (28.5%); and technical incident: 54/221 (24.4%). Communication breakdown related to 17/221 (7.7%) of adverse events; with inefficiency of care accounting for 5/221 (2.3%) of adverse events.	107/221 events had harm (76.9%). 119/170 (70.0%) of the harms were psychological. 39/170 of the harms were physical (22.9%).	Moderate
Leon AC 1999 <sup>59</sup>	Observational	Primary care	USA	Interviews	This study examined the diagnosis of mental health issues in primary care. It found: misdiagnosis of major depression in 160/1000 (16.0%) patients; and misdiagnosis of panic disorder in 96/1001 (9.6%) patients.			Moderate
Lund BC 2010 <sup>60</sup>	Observational (part of trial)	Primary care	USA	Survey	Of 236 patients, 34 (14.4%) experienced an adverse drug event.			High
Makeham M 2008 <sup>61</sup>	Systematic review	Primary care	Australia	Various	Review of 49 studies found estimates of patient safety incidents in primary care were 0.004 to 240 per 1000 primary care consultations.	26% to 57% of incidents involved diagnostic "incidents"; 7% to 52% involved treatment; 13% to 47% involved investigations; 9% to 56% involved office administration; 5% to 72% were communication incidents.	45% to 76% of all 'incidents' were preventable. Harm from safety incidents ranged from 1.3 significant minor incidents per 1000 treatments to 4% of incidents resulting in death, 17% to 39% resulting in harm, and 70% to 76% had potential for harm.	High

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Makeham MA 2002 <sup>62</sup>	Observational	General practice	Australia, Canada, Netherlands, New Zealand, UK, USA	Incident reporting system		Of 132 incident reports, 104/132 (78.8%) were related to process incidents. Of these, 26/104 (25.0%) related to incidents in office administration; 17/104 (16.4%) related to investigation incidents; 38/104 (36.6%) related to treatment incidents; 20/104 (19.2%) related to communication incidents; 1/104 (1.0%) related to payment incidents; and incidents in health care workforce management accounted for 2/104 (1.9%) of reports. 28 of the 132 incident reports were related to knowledge and skill incidents (21.2%). These were further broken down into: incidents in execution of a clinical task 7/28 (25.0%); incidents in diagnosis: 18/28 (64.3%); wrong treatment decision with right diagnosis: 3/28 (10.7%).		Moderate
Makeham MA 2006 <sup>63</sup>	Observational	General Practice	Australia	Incident reporting system	84 GPs submitted 418 incident reports, claimed for 490864 consultations and saw 166569 individuals in one year. The incidence of reported incident per consultation per year was 0.078 (95% CI 0.076% to 0.080%). The incidence of reported incidents per patient seen per year was 0.24% (95% CI 0.235% to 0.245%). Two incidents are reported for every 1000 individual patients seen by a GP.		418/166,569 (0.25%) incidents led to patient harm.	High

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Makeham MA 2008 <sup>64</sup>	Observational	General Practice	Australia	Incident reporting system		Incidents relating to the processes of healthcare (n=365, 69.5%) were more common than those relating to deficiencies in the knowledge and skills of health professionals (n=160, 30.5%).		High
Martinez Sanchez A 2011 <sup>65</sup>	Observational	Community pharmacy (review of primary care prescriptions)	Spain	Review of prescriptions	Review of community pharmacy records for primary care indicated prescription incidents in 355/23995 cases (1.5%).	Most incidents were due to incomplete/incorrect information or prescribed items being unavailable: 247/355 (69.6%). Inappropriate doses accounted for 27/355 incidents (7.6%); and inappropriate direction or instruction accounted for 25/355 incidents (7.0%).		Moderate
McKay J 2009 <sup>66</sup>	Observational	General practice	UK	Significant Event Audit		259 significant event analyses were reviewed with the following breakdown of incidents: disease diagnosis and disease management: 46/259 (17.8%); prescribing, dispensing and other drugs 46/259 (17.8%); patient and relatives: 43/259 (16.6%); investigations and results: 37/259 (14.29%); communication: 23/259 (8.88%); administration: 16/259 (6.18%); medical records and confidentiality: 15/259 (5.79%); appointments and surgeries: 12/259 (4.63%); home visits and external care: 10/259 (3.86%); equipment: 7/259 (2.70%); miscellaneous: 2/259 (0.77%); health and safety: 2/259 (0.77%).	191 reports were reviewed regarding harm. Most patients did not suffer any harm: 109/191 (57.1%); low harm occurred in 14/191 cases (7.33%); moderate harm in 22/191 cases 11.52%; and severe harm in 9/191 cases (4.71%). Three deaths occurred overall: 3/191 (1.57%). 34/191 (17.80%) cases were not classified. The 109 incidents (57.1%) which did not lead to any harm were made up of incidents which had the potential to cause patient harm but were prevented, or incidents that ran to completion without harm occurring – ‘near misses’.	Moderate



First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
McKay J 2013 <sup>67</sup>	Observational	General practice	UK	Record review	Twenty-one GP trainees attended a two hour workshop about how to use a trigger tool to identify patient safety incidents and then reviewed 25 clinical records each. 80 out of 520 records (15%) identified previously undetected patient safety incidents.		30 out of these 80 incidents were judged potentially preventable (45%). 35 were judged to cause moderate to severe harm (44%).	Low
Miller GC 2006 <sup>68</sup>	Observational	General practice	Australia	Incident reporting system	852/8215 patients (10.4%) suffered adverse events.		Harm to patients was recorded as: mild: 297/551 (53.9%); moderate: 197/551 (35.8%); severe: 55/551 (10.0%); and 'don't know': 2/551 (0.4%).	Low
Montastruc P 1993 <sup>69</sup>	Observational	Primary care clinic	France	Record review		49 adverse events identified in a rural area over a one year period were reviewed. Cutaneous adverse drug reactions accounted for 15/49 (30.6%) events. This was followed by digestive adverse drug reactions: 14/49 (28.6%); neurological adverse drug reactions: 11/49 (22.5%); pulmonary adverse drug reactions: 3/49 (6.1%); cardiovascular adverse drug reactions: 2/49 (4.1%); and others: 4/49 (8.2%).		Low
Murie J 2003 <sup>70</sup>	Observational	General practice	UK	Significant Event Audit		Looked at 55 significant events: operational accounted for 24/55 (43.6%), followed by clinical 20/55 (36.4%); strategic failures: 9/55 (16.4%); human resources: 2/55 (3.6%); environmental: 3/55 (5.5%); political: 1/55 (1.8%) and legislative: 1/55 (1.8%).	Of the 55 cases reviewed, 12/55 (21.8%) resulted in death; 10/55 (18.2%) resulted in severe harm; 2/55 (3.6%) resulted in moderate harm; 5/55 (9.1%) resulted in low harm; and 26/55 (47.3%) resulted in no harm.	Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Neville RG 1989 <sup>71</sup>	Observational	General practice	UK	Review of prescriptions	There were a total of 504 incidents from 15,916 prescription items (3.17%) during a three month observation period at one clinic.	212/504 incidents (42.1%) related to dose. This was further broken down as: strength of preparation not stated: 162/212 (76.4%); dose wrong by multiple of 10: 4/212 (1.9%); and other incorrect dose: 46/212 (21.7%). 187/504 (37.1%) related to quantity (wrong pack size). 40/504 (7.9%) related to naming of drugs, further subcategorized as follow: incomplete description: 33/40 (82.5%); confusion of similar names: 3/40 (7.5%); wrong drug: 3/40 (7.5%); and controlled drug regulations not followed: 1/40 (2.5%). 57/504 (11.3%) of the prescription incidents related to formulation. 8/504 (1.6%) related to 'limited list (prep not available on NHS)'.		Low
Nicholson D 2006 <sup>72</sup>	Observational	Primary care	USA	Record review	Of the 24 participating physicians, 22/24 (91.7%) made at least one prescribing incident over the seven month period that led to an adverse event. All of the incidents leading to an adverse event were described as preventable or ameliorable.			Low
O'beirne M, 2013 <sup>73</sup>	Observational	Family practice	Canada	Incident reporting system		264 incident reports were submitted by 191 practices. The top four types of incidents reported were documentation (41.4%), medication (29.7%), clinical administration (18.7%) and clinical process (17.5%).	Most reported incidents were judged to have 'virtually certain evidence of preventability' (93%). Harm was associated with 50% of incidents. Only 1% of the incidents had a severe impact.	Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Olaniyan JO 2014 <sup>74</sup>	Systematic review	Primary care	UK	Various	Thirty-three studies estimating the incidence of medication incidents in primary care were identified and thirty-six studies evaluated the impact of incident-prevention interventions. Medication incidents were found to be common, with incident rates ranging from less than 1% to more than 90%, depending on the definitions and methods used.	The prescribing stage was most susceptible to incidents. Those aged over 65 years and children were more likely to experience significant incidents.		High
Paille F 1995 <sup>75</sup>	Observational	General practice	France	Review of prescriptions	Focused on incidents for people with hypertension. Found that 1324/4080 (32.5%) prescriptions contained potentially inappropriate medications.			Low
Pandit NB 2008 <sup>76</sup>	Observational	Primary care	India	Survey	Reviewed 182 organizations. 77% of had unsafe injection practices, including the use of a boiling pan for sterilisation, recapping of needles and exposure to body fluids. The prevalence of needle stick injuries among service providers was 52.2% and the annual incidence of needle stick injuries was 19%.			Low
Pariser RJ 1987 <sup>77</sup>	Observational	Primary care	USA	Record review		Reviewed 319 medical incidents in 260 patients with skin conditions. 281/319 (88.0%) were diagnosis incidents.		Low
Pearson A 2009 <sup>78</sup>	Systematic review	Primary care	Australia	Various		Incidents related to administration, knowledge and skills, prescribing, processes and communication.	Review of 33 studies found that harm rates ranged from 24% to 42%.	High

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Phillips Jr RL 2004 <sup>79</sup>	Observational	Primary care clinic	USA	Malpractice claims			26,126 peer-reviewed malpractice claims were reviewed. 5921/26126 (22.7%) were assessed as negligent claims. The malpractice claims data identified the following levels of harm to patients. 2148/5921 (36.3%) resulted in a death; 1124/5921 (19.0%) resulted in severe harm; 1542/5921 (26.0%) resulted in moderate harm; and 1107/5921 (18.7%) resulted in low harm.	Moderate
Phillips RL 2006 <sup>80</sup>	Observational	Family medicine clinics	USA	Incident reporting system		Examined reports to primary care incident report system. 898/935 (96.0%) were process incidents and 37/935 (4.0%) were knowledge and skill incidents.	145 / 701 (20.7%) incidents resulted in no harm to the patient, 196/701 (30.0%) in low harm, 203/701 (30.0%) in moderate harm, 100/701 (14.3%) in severe harm and 57 / 701 (8.1%) in death.	High
Rosser W 2005 <sup>81</sup>	Observational	Family medicine clinics	Canada, England, Netherlands, New Zealand, USA, Australia	Incident reporting system		Examined 508 incident reports. The most common cause of incident was classified as office processes: 160/508 (31.5%). This was followed by: treatment incidents: 109/508 (21.5%); clinical knowledge: 89/508 (17.5%); external investigations: 73/508 (14.4%); communication: 62/508 (12.2%); work force management: 10/508 (2.0%); and financial accounting: 5/508 (1.0%).		High

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Rubin G 2003 <sup>82</sup>	Observational	General practice	UK	Incident reporting system		Examined 940 incident reports. These were subcategorized as follows: prescription incidents: 397/940 (42.2%); communication incidents: 282/940 (30.0%); equipment incidents: 153/940 (16.3%); appointments incidents: 63/940 (6.7%); clinical incidents: 24/940 (2.6%); other incidents: 21/940 (2.2%).		Low
Ryan C 2009 <sup>83</sup>	Observational	General practice	Ireland	Record review	Records of 500 elderly people were screened. One tool found 69 medicines were prescribed inappropriately in 65 patients (13%). Another tool identified 63 potentially inappropriate medicines in 52 patients (10.4%).			Moderate
Sandars J 2003 <sup>84</sup>	Systematic review	Primary care	UK	Various	Review of 280 studies found wide differences in rates of incidents in primary care, varying from five to 80 per 100,000 consultations. Prescribing and prescription incidents have been found to occur in up to 11% of all prescriptions, mainly related to incidents in dose.	Incidents related to diagnosis were the most common across all studies, varying from 26% to 78% of identified incidents. Incidents associated with diagnosis, either delayed or missed, were most likely to result in major harm.		High

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Sayers YM 2009 <sup>85</sup>	Observational	General practice	Ireland	Review of prescriptions	From a total of 3,948 prescriptions, 491 (12.4%) contained one or more incidents. From a total of 8,686 drug items, 546 (6.2%) contained one or more incidents.	The most common incident was 'no direction': 226/491 (46.0%), followed by 'other prescription-related cause': 93/491 (18.9%). The rest of the breakdown was as follows: not dated: 71/491 (14.46%); inadequate information: 60/491 (12.2%); mix up of prescriptions: 51/491 (10.4%); wrong dose: 43/491 (8.8%); no age given: 37/491 (7.5%); CD incident: 27/491 (5.5%); illegible: 27/491 (5.5%); off market: 20/491 (4.1%); not signed: 17/491 (3.5%).		Low
Schiff GD 2013 <sup>86</sup>	Observational	Primary care	USA	Malpractice claims		Examined 551 malpractice claims from primary care. Allegations were related to diagnosis (72.1%), medications (12.3%), other medical treatment (7.4%), communication (2.7%), patient rights (2.0%), and patient safety or security (1.5%).		Low
Schneider JK 1992 <sup>87</sup>	Observational	General medicine	USA	Record review	Records for 332 elderly people who attended a general medical clinic and 131 who attended a geriatric clinic showed potential drug interactions in 143 people (31%). There were 107 documented adverse drug reactions in 97 patients (21%). Of these patients, 86 were noted by doctors as having had a reaction.		Twelve reactions led to hospitalization.	Moderate
Schweppach DL 2012 <sup>88</sup>	Observational	Primary care	Switzerland	Survey		630 doctors and nurses were surveyed and 391 (31%) described 936 threats to patient safety. Safety of medication (8.8%), triage by nurses (7.2%) and drug interactions (6.8%) were the threats cited most frequently.		Low

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Shaughnessy AF 1989 <sup>89</sup>	Observational	Family practice	USA	Review of prescriptions		1814 prescriptions by family medicine residents were reviewed and incidents were categorized as: omissions: 707/1814 (38.97%); incorrect dosage: 254/1814 (14.00%); legal issues: 109/1814 (6.01%); non-prescription based: 417/1814 (22.99%); inaccurate/vague quantity: 218/1814 (12.02%); and incomplete direction: 109/1814 (6.01%).		Moderate
Singh H 2013 <sup>90</sup>	Observational	Primary care clinic	USA	Record review		Reviewed 190 diagnostic incidents identified at two primary care facilities. Found missed diagnoses in 68 cases, relating to pneumonia (6.7%), decompensated congestive heart failure (5.7%), acute renal failure (5.3%), cancer (5.3%), and urinary tract infection or pyelonephritis (4.8%). These were due to issues in one or more of the following: the clinical encounter (78.9%), referrals (19.5%), patient-related factors (16.3%), follow-up (14.7%), and diagnostic tests (13.7%).	86% of incidents were classed as potentially moderately or severely harmful.	Moderate
Smith PC 2005 <sup>91</sup>	Observational	Primary care	USA	Survey	253 clinicians were surveyed about 1614 patient visits. Clinicians reported missing clinical information in 13.6% of visits.	Missing information included laboratory results (6.1% of all visits), letters/dictation (5.4%), radiology results (3.8%), history and physical examination (3.7%), and medications (3.2%).		Moderate



First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Smits M 2010 <sup>92</sup>	Observational	Family practice	Netherlands	Record review	145 patient records were reviewed and 27/145 (18.6%) patient safety incidents were identified.	15/27 incidents (55.6%) related to inadequate or suboptimal treatment; 9/27 (33.3%) related to triage incidents; and 6 /27 (22.2%) related to wrong or misguided diagnosis.	Of the 27 identified patient safety incidents, 8/27 (29.6%) had consequences for patients: an extra intervention was needed in 6/27 (22.2%) cases, and 2/27 (7.4%) patients had to be admitted to a hospital. No incidents resulted in permanent harm or death.	Moderate
Statham MO 2008 <sup>93</sup>	Observational	General practice	Australia	Record review			1062 people with acute eye disease were assessed. Incorrect diagnoses accounted for 642/1062 (60.5%) patients. conditions that were misdiagnosed and subsequently associated with severe adverse patient outcome occurred in 91/123 (74.0%) patients. The remainder had a correct initial diagnosis, but subsequent incorrect treatment: 32/123 (26.0%). 63/123 (51.2%) suffered low harm; 49/123 (39.8%) suffered moderate harm; and 11/123 (8.9%) suffered severe harm. There were judged to be 123/1062 (11.6%) patients who suffered preventable adverse outcomes.	Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Stewart L, 2012 <sup>94</sup>	Observational	Primary care clinic	USA	Interviews		Of 219 patients interviewed to compare their medication use with their medical record, 162 (74%) had at least one discrepancy. The most common type of discrepancy was an incorrect medication documented on the chart. The most common reasons included over-the-counter (OTC) use of medications and patients not reporting use of medications.		Low
Straand J 1999 <sup>95</sup>	Observational	General practice	Norway	Review of prescriptions	Reviewed 16774 prescription incidents among elderly people. 13.5% of prescriptions met criteria for inappropriate prescribing.			Moderate
Tam KWT 2008 <sup>96</sup>	Observational	Primary care	Hong Kong	Record review and incident reports		Of the 132 adverse drug events in four clinics, 108/132 (81.82%) were categorized as actual adverse drug events (as opposed to potential adverse drug events). Of these, 5/108 (4.6%) were preventable; and 103/108 (95.4%) were non-preventable. Of the 108 adverse drug events, 5/108 (4.6%) were detected as preventable.		Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Taylor LK 2005 <sup>107</sup>	Observational	Primary care clinic	USA	Incident reporting system		Looked at alerts generated by prescriptions over a nine-month period, and found alerts for prescription incidents in 6428/22419 (28.7%) cases. These were subcategorized as: drug-disease contraindication : 2644/6428 (41.1%); drug-drug interactions: 1522/6428 (23.7%); potential toxicity: 1022/6428 (16.0%); drug duplication: 731/6428 (11.4%); contraindicated for patient age: 249/6428 (3.9%); potential dosing incident: 221/6428 (3.4%); and other: 39/6428 (0.6%).		Low
Tilyard M 2005 <sup>97</sup>	Observational	General practice	Australia, Canada, Netherlands, New Zealand, UK, USA	Incident reporting system		431 incident reports which were categorized as: treatment process incident: 110/431 (25.5%); office administration incident: 82/431 (19.0%); investigation process incident: 73/431 (16.9%); communication incidents: 62/431 (14.4%); wrong diagnosis: 54/431 (12.5%); and other: 50/431(11.6%).		Moderate
Tomlin A, 2012 <sup>98</sup>	Observational	General practice	New Zealand	Record review	During a 6-year period, 173,478 patients from 30 practices received 4,811,561 prescriptions. There were 37,397 allergies, adverse events and other warnings recorded for 24994 patients (14%).			Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Tsang C 2012 <sup>99</sup>	Systematic review	Primary care	UK	Various	Review of 15 studies of the types of adverse events that are routinely recorded in primary care found about 6.5% of adult emergency admissions were due to drug-related events. Between 0.7% and 2.3% of deaths following adverse events were attributed to treatment in primary care. In patients aged >65 years, the occurrence of adverse drug events was estimated at 4.9 per 1000 population			High
Tsang C 2013 <sup>100</sup>	Observational	General practice	UK	Record review	Examined 74,763 people's records. Incidence was 6 adverse events per 1000 person-years (95% CI 5.74 to 6.27), which is equivalent to 8 adverse events per 10,000 consultations. Those at greatest risk were patients aged 65-84 years, those with the most consultations, five or more emergency admissions, and those with the greatest number of conditions. People registered at their practice for the longest periods of time had a lower risk of an adverse event.			Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Tse J 2011 <sup>101</sup>	Observational	General practice	Australia	Record review		33 patient records were reviewed. High levels of accuracy were found in the area of demographic details (94%). Moderately high levels of accuracy were reported for allergies (61%) but also a considerable percentage of non-recorded information was present (36%). Inaccuracies in medication lists were reported in 51% of records reviewed with 32.1% of all medications being inaccurately recorded. While over 91% of participants had a history summary with eight or less items present, omissions were reported for one in every five participants.		Low
Van Der Hooft CS 2005 <sup>102</sup>	Observational	Primary care	Netherlands	Record review	Looked at the risk of inappropriate drug prescriptions. In the calendar year 2001, found the risk of receiving at least one inappropriate drug prescription in older adults was 5052/25258 (20%). Between 1997 and 2001, the 1-year risk of receiving at least one inappropriate drug prescription for older adults ranged between 16.8% (95% CI: 16.3–17.3%) and 18.5% (18.3–18.7%).			Moderate
Van Dulmen SA 2011 <sup>103</sup>	Observational	Primary care	Netherlands	Record review and incident reports	1000 records were reviewed from 20 practices. In 18 out of 1000 (1.8%; 95% CI 1.0 to 2.6) records an incident was detected.	The main causes of incidents were incidents in clinical decisions (89%), communication with other healthcare providers (67%), and monitoring (56%).		Moderate

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Wahls TL 2007 <sup>104</sup>	Observational	Primary care	USA	Survey	Diagnostic incidents associated with the failure to follow up on abnormal diagnostic studies ("missed results") are a potential cause of treatment delay. 106 clinicians were surveyed who saw an average of 86 patients per two week period. Providers encountered 64 patients with missed results during the two week period leading up to the study and 52 patients with treatment delays.	The most common missed results included imaging studies (29%), clinical laboratory (22%), anatomic pathology (9%), and other (40%). The most common diagnostic delays were cancer (34%), endocrine problems (26%), cardiac problems (16%), and others (24%).		Moderate
Wallace F 2013 <sup>105</sup>	Systematic review	Primary care	Ireland	Various		Examined the epidemiology of malpractice claims in primary care (written demands for compensation for medical injury). Studies reporting on original data with ten or more cases were eligible. Thirty-four studies were included. Twenty-eight studies included data from medical indemnity malpractice claims databases and six studies reported survey data. Fifteen studies were from the USA, nine from the UK, seven from Australia, two from France and one from Canada. Diagnosis incidents were the most common, accounting for 26% to 63% of all claims. Medication incident was the second most common, accounting for 5.6% to 20% of all claims.		Low

First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Weingart SN 2005 <sup>106</sup>	Observational	Primary care	USA	Record review and interviews	661 patients who received prescriptions from 4 primary care practices were interviewed and their records reviewed. Patients' failure to discuss 90 medication symptoms resulted in 19 (21%) ameliorable and 2 (2%) preventable adverse drug events. Physicians' failure to change therapy in 48 cases resulted in 31 (65%) ameliorable adverse drug events.			Moderate
Wetzels R 2009 <sup>107</sup>	Observational	General practice	Netherlands	Record review and incident reporting system		31 incidents were identified and classified as follows: Practice administration incidents accounted for 10/31 (32.3%). This was subcategorized as medical records: 7/10 (70.0%); appointments: 2/10 (20.0%); other – administration: 1/10 (10.0%). Diagnostic incidents accounted for 6/31 (19.4%). Broken down as: wrong diagnosis: 1/6 (16.7%); delayed diagnosis: 4/6 (66.7%); other diagnosis: 1/6 (16.7%). Therapeutic incidents accounted for 7/31 (22.6%). Subcategorized as: wrong, incomplete treatment: 5/7 (71.4%); other treatment: 2/7 (28.6%). Communication incidents accounted for 8/31 (25.8%). Subcategorized as: communication with patients: 3/8 (37.5%); and communication with caregivers: 5/8 (62.5%).		Moderate



First author	Study type	Setting	Country	Measurement method	Number of safety incidents	Type of safety incidents	Harm associated with incidents	Quality rating
Woolf SH 2004 <sup>108</sup>	Observational	Family medicine clinics	USA	Incident reporting system		Examined incident reports. Process incidents accounted for 135/184 (73.4%) of reports. Process incidents were further subcategorized as follows: treatment: 50/135 (37.0%); office administration: 33/135 (24.4%); investigations: 27/135 (20.0%); communication: 23/135 (17.0%); and insurance-related incidents: 2/135 (1.5%). Knowledge and skill incidents accounted for 49/184 (26.6%) of incident reports. These were further broken down into the below categories: execution of a clinical task: 6/49 (12.2%); wrong diagnosis: 26/49 (53.0%); and wrong treatment decision: 17/49(34.7%).		High
Zavaleta-Bustos M 2008 <sup>109</sup>	Observational	Primary care clinic	Mexico	Review of prescriptions		Reviewed 370 prescriptions. The most common incident was found to be inappropriate prescriptions: 268/370 (72.4%). This was followed by inappropriate dosage regime: 102/268 (38.1%); inappropriate indication: 92/268 (34.3%); unnecessary medications: 24/268 (9.0%); medication duplicity: 14/268 (5.2%); potential drug-drug interactions: 23/268 (8.6%); missing medication: 8/268 (3.0%); and inappropriate administration route: 5/268 (1.9%).		Moderate

**Additional material included in systematic reviews so not duplicated in the review**

Al Khaja KA et al. An evaluation of prescribing errors in primary care in Bahrain. *Int J Clin Pharmacol Ther* 2005; 43: 294–301.

Australian Institute of Health and Welfare. Public and Private sector medical indemnity claims in Australia. <http://www.aihw.gov.au/search/?q=medical+indemnity+claims>

Avery AJ et al. Prevalence and Causes of Prescribing Errors in General Practice: The PRACtiCE Study. London: Division of Primary Care, School of Community Health Sciences, University of Nottingham: General Medical Council, 2012: 1–259.

Avery T, Barber N, Ghaleb M, et al. Investigating the prevalence and causes of prescribing errors in general practice: Report for the General Medical Council (GMC); May 2012.

Baker R, Sullivan E, Camosso-Stepinovic R, Rashid A, et al. Making use of mortality data to improve quality and safety in general practice: a review of current approaches. *Qual Saf Health Care* 2007; 16(2): 84-9.

Bhasale AL, Miller GC, Reid S, Britt HC. Analysing potential harm in Australian general practice; an incident-monitoring study. *Med J Aust* 1998; 169:73–6.

Britten N, Stevenson FA, Barry CA, Barber N, Bradley CP. Misunderstandings in prescribing decisions in general practice: qualitative study. *BMJ* 2000; 320:484–8.

Buck MD, Atreja A, Brunker CP, Jain A, Suh TT, Palmer RM, et al. Potentially inappropriate medication prescribing in outpatient practices: prevalence and patient characteristics based on electronic health records. *The American Journal of Geriatric Pharmacotherapy* 2009;7(2):84-92.

Budnitz DS, Pollock DA, Weidenbach KN et al. National surveillance of emergency department visits for outpatient adverse drug events. *JAMA* 2006;

Canadian Medical Protective Association (CMPA). Critical incident groupings, family, practitioner settlement cases, CMPA 2005–2009. Courtesy of the Research Department, Canadian Medical Protective Association.

Chen YF et al. Incidence and possible causes of prescribing potentially hazardous/contraindicated drug combinations in general practice. *Drug Safety* 2005; 28: 67–80.

Conradi MH, de Mol BAJM. Research on errors and safety in Dutch general and hospital practice. In Rosenthal AR, Mulcahy L, Lloyd-Bostock S (eds). *Medical Mishaps: Pieces of the Puzzle*. Buckingham (UK): Open University Press, 1999: 74–84.

Cox SaH, JD. A retrospective review of significant events reported in one district in 2004–2005. *British Journal of General Practice*. 2007;

Cox S, Holden J. A retrospective review of significant events reported in one district in 2004–2005. *Br J Gen Pract* 2007; 57(542):

Devine EB et al. The impact of computerized provider order entry on medication errors in a multispecialty group practice. *J Am Med Inform Assoc* 2010; 17: 78–84.

Dhabali AAH et al. Pharmaco-epidemiologic study of the prescription of contraindicated drugs in a primary care setting of a university: a retrospective review of drug prescription. *Int J Clin Pharmacol Ther* 2011; 49: 500–509.

Diamond M, Kamien M, Sim M, Davis J. A critical incident study of general practice trainees in their basic general practice term. *Med J Aust* 1995; 162(6): 321-4.

Dovey S, Green L, Fryer GF. Identifying Threats to Patient Safety in Family Practice . [www.aafppolicy.org/posters/errors/](http://www.aafppolicy.org/posters/errors/),

Dovey S, Phillips R, Green L, Fryer G. Consequences of medical errors observed by family physicians. *Am Fam Physician* 2003; 67(5): 915.

Dovey S, Phillips R, Green L, Fryer G. Types of medical errors commonly reported by family physicians. *Am Fam Physician* 2003; 67(4): 697.

Dovey S. Advancing Understanding of medical errors in general practice: A discussion of recent research from the American Academy of Family Physicians. *New Zealand Journal of Family Practice*. 2003;30:242

Duggan C, Bates I, Hough J: Discrepancies in prescribing. Where do they occur? *Pharmaceutical J* 1996, 256: 65-67.

Duggan C, Feldman R, Hough J, Bates I: Reducing adverse prescribing discrepancies following hospital discharge. *Int J Pharmacy Practice* 1998, 6: 77-82.

Elder N, Vonder Meulen M, Cassidy A. The identification of Medical errors by family physicians during outpatient visits. *Annals of Family Medicine* . 2004

Elder NC, McEwen TR, Flach JM, Gallimore JJ. Management of test results in family medicine offices. *Annals of Family Medicine* 2009;7(4):343-51.

Elder NC. Quality and Safety in Outpatient Laboratory Testing. *Clinics in Laboratory Medicine* 2008;28(2):295-303.

Ely JW, Levinson W, Elder NC, Mainous AG III, Vinson DC. Perceived causes of family physicians' errors. *J Fam Pract* 1995; 40:337–44.

Fenn P, Gray A, Rivero-Arias O, et al. The epidemiology of error: an analysis of databases of clinical negligence litigation. University of Manchester, Manchester, 2004.

Field TS et al. Adverse drug events resulting from patient errors in older adults. *J Am Geriatr Soc* 2007; 55: 271–276.

*How Safe is Primary Care? A Systematic Review*

- 1 Fischer G, Feters MD, Munro AP, Goldman EB. Adverse events in primary care identified from a risk-management  
2 database. *J Fam Pract* 1997; 45:40–6.
- 3 Flannery FT, Parikh PD, Oetgen WJ. Characteristics of medical professional liability claims in patients treated by family  
4 medicine physicians. *J Am Board Fam Med* 2010;23:753–61.
- 5 Franklin BD, O'Grady K, Donyai P, Jacklin A, Barber N: The impact of a closed-loop prescribing and administration  
6 system on prescribing errors, administration errors and staff time. *Qual Safe Health Care* 2007, 16: 279-284.
- 7 Gandhi TK et al. Adverse drug events in ambulatory care. *NEJM* 2003; 348: 1556–1564.
- 8 Gandhi TK et al. Impact of basic computerized prescribing on outpatient medication errors and adverse drug events. *J Am*  
9 *Med Inform Assoc* 2002; 9: S48–S49.
- 10 Gandhi TK et al. Outpatient prescribing errors and the impact of computerized prescribing. *J Gen Intern Med* 2005; 20:  
11 837–841.
- 12 Gandhi TK, Kachalia A, Thomas EJ, et al. Missed and delayed diagnoses in the ambulatory setting: a study of closed  
13 malpractice claims. *Ann Intern Med* 2006;145:488–96.
- 14 Gandhi TK, Kachalia A, Thomas EJ, Puopolo AL, Yoon C, Brennan TA, Studdert DM. Missed and delayed diagnoses in  
15 the ambulatory setting: A study of closed malpractice claims. *Annals of Internal Medicine* 2006;145(7):488-96.
- 16 Gandhi TK, Sittig DF, Franklin M, Sussman AJ, Fairchild DG, Bates DW. Communication breakdown in the outpatient  
17 referral process. *J Gen Intern Med* 2000; 15:626–31.
- 18 Gandhi TK, Weingart SN, Borus J, Seger AC, Peterson J, Burdick E, et al. Patient safety: adverse drug events in  
19 ambulatory care. *New England Journal of Medicine* 2003;348(16):1556-64.
- 20 Gandhi TK, Weingart SN, Seger AC, Borus J, Burdick E, Poon EG, et al. Outpatient prescribing errors and the impact of  
21 computerized prescribing. *Journal of General Internal Medicine* 2005;20(9):837-41.
- 22 Guerra R, Bublitz C, Parnes B, Dickinson LM, Van Vorst R, et al. Missing clinical information during primary  
23 care visits. *JAMA* . 2005 Feb 2;
- 24 Inman W. Postmarketing surveillance of adverse drug reactions in general practice. I: search for new methods. *Br Med J*  
25 *(Clin Res Ed)* 1981; 282(6270): 1131-2.
- 26 Jacobs S, O'Beirne M, Derflinger L-P, Vlach L, et al. Errors and adverse events in family medicine: Developing and  
27 validating a Canadian taxonomy of errors. *Can Fam Physician* 2007; 53: 270-6.
- 28 Jani YH, Ghaleb MA, Marks SD, Cope J, Barber N, Wong ICK: Electronic prescribing reduced prescribing errors in a  
29 pediatric renal outpatient clinic. *J Pediat* 2008, 152:214-218.
- 30 Jena AB, Seabury S, Lakdawalla D, et al . Malpractice risk according to physician specialty. *N Engl J Med* 2011;365:629–  
31 36.
- 32 Kaushal R et al. Adverse drug events in pediatric outpatients. *Ambul Pediatr* 2007; 7: 383–389.
- 33 Kaushal R et al. Medication errors in paediatric outpatients. *Qual Saf Health Care* 2010; 19: 10.
- 34 Kostopoulou O, Oudhoff J, Nath R, Delaney BC, Munro CW, Harries C, et al. Predictors of diagnostic  
35 accuracy and safe management in difficult diagnostic problems in family medicine. *Medical Decision Making*.  
36 2008;
- 37 Kostopoulou O, Delaney BC, Munro CW. Diagnostic difficulty and error in primary care—a systematic review. *Fam Pract*  
38 Kravitz RL, Rolph JE, Petersen L. Omission-related malpractice claims and the limits of defensive medicine. *Med Care*  
39 *Res Rev* 1997;54:456–71.
- 40 Kriisa I. Swedish malpractice reports and convictions. *Qual Assur Health Care* 1990;
- 41 Leape LL. Reporting of adverse events. *New England Journal of Medicine* 2002;
- 42 Mackenzie P . Learning from clinical claims in primary care. *Medical Protection Society (MPS) Casebook*. 2011  
43 September 2011:7–9.
- 44 Makeham MA. The Measurement of Threats to Patient Safety in Australian General Practice: University of Sydney;  
45 2008.
- 46 Marwaha M et al. A retrospective analysis on a survey of handwritten prescription errors in general practice. *Int J Pharm*  
47 *Pharm Sci* 2010; 2(Suppl. 3): 80–82.
- 48 Meredith V, Cook CB, Penman A. Use of the Physician Insurers Association of America database as a surveillance tool  
49 for diabetes-related malpractice claims in the U.S. *Diabetes Care* 1998;21:1096–100.
- 50 Michel P, Wilson R. Adverse events in developing countries. In preparation.
- 51 Morcos S, Francis S-A, Duggan C: Where are the weakest links? A descriptive study of discrepancies in prescribing  
52 between primary and secondary sectors of mental health service provision. *Psychiatric Bull* 2002, 26: 371-374.
- 53 Morris C, Rodgers S, Hammersley V, Avery A, Cantrill J. Indicators for preventable drug related morbidity: application in  
54 primary care. *Qual Saf Health Care* 2004;
- 55 Murie J, McGhee C. Assessing risk by analysing significant events in primary care. *Qual Prim Care* 2003; 11(3): 205-10.
- 56 Najaf-Zadeh A, Dubos F, Aurel M, et al. Epidemiology of malpractice lawsuits in paediatrics. *Acta Paediatr* 2008;97:1486–  
57 91.
- 58 Najaf-Zadeh A, Dubos F, Pruvost I, et al. Epidemiology and aetiology of paediatric malpractice claims in France. *Arch Dis*  
59 *Child* 2011;96:127–30.

1 Nanji KC et al. Errors associated with outpatient computerized prescribing systems. *J Am Med Inform Assoc* 2011; 18:  
2 767–773.

3 Perrio M, Wilton L, Shakir S. A modified prescription-event monitoring study to assess the introduction of Seretide  
4 Evohaler in England: an example of studying risk monitoring in pharmacovigilance. *Drug Saf* 2007; 30(8): 681-95.

5 Persell SD, Heiman HL, Weingart SN, Burdick E, Borus J, Murff HJ, et al. Understanding of drug  
6 indications by ambulatory care patients. *American Journal of Health System Pharmacy* 2004;

7 Phillips RL Jr., Bartholomew LA, Dovey SM, et al. Learning from malpractice claims about negligent, adverse events in  
8 primary care in the United States. *Qual Saf Health Care* 2004;13:121–6.

9 Pirohamed M, James S, Meakin S, Green C, Scott AK, Walley TJ, Farrar K, Park BK, Breckenridge AM: Adverse drug  
10 reactions as cause of admission to hospital: prospective analysis of 18 820 patients. *Br Med J* 2004, 329: 15-19.

11 primary care patients. *Pharmacoepidemiology and Drug Safety* 2007;16(6):658-67.

12 Research Programme Medication Errors 2: Pilot Study Volume 16 . London: The Department of Health; 2007:40-44.

13 Rothschild JM, Federico FA, Gandhi TK, et al. Analysis of medication-related malpractice claims: causes, preventability,  
14 and costs. *Arch Intern Med* 2002;162:2414–20.

15 Runciman WB, Roughhead EE, Semple SJ, Adams RJ. Adverse drug events and medication errors in Australia. *Int J Qual*  
16 *Health Care* 2003; 15(suppl\_1): i49-59.

17 Schadlich P. [Quality of event data in detection of unwanted drug side-effects in general practice of established  
18 physicians]. *Gesundheitswesen* 1993; 55(Suppl 1): 8-12.

19 Schiff GD, Hasan O, Kim S, Abrams R, Cosby K, Lambert BL, et al. Diagnostic Error in Medicine: Analysis of 583  
20 Physician-Reported Errors. *Archives of Internal Medicine* 2009;169(20):1881-87.

21 Shah SNH, Aslam M, Avery AJ: A survey of prescription errors in general practice. *Pharmaceutical J* 2001, 267: 860-862.

22 Shaw R, Drever F, Hughes H, Osborn S, Williams S. Adverse events and near miss reporting in the NHS. *Qual Saf Health*  
23 *Care* 2005; 14(4): 279-83.

24 Silk N. An analysis of 1000 consecutive general practice negligence claims. Medical Protection Society (MPS), 2000.

25 Singh H, Thomas EJ, Khan MM, Petersen LA. Identifying diagnostic errors in primary care using an electronic screening  
26 Singh R, Lean-Plunckett EA, Kee R, Wisniewski A, Cadzow R, Okazaki S, et al. Experience with a trigger tool for  
27 identifying adverse drug events among older adults in ambulatory primary care. *Quality & Safety in Health Care*  
28 2009;18(3):199-204.

29 Soendergaard B, Kirkeby B, Dinsen C, Herborg H, Kjellberg J, Staehr P. Drug-related problems in general practice:  
30 results from a development project in Denmark. *Pharm World Sci* 2006;28(2):61-4.(46)

31 Sweeney G, Westcott R, Stead J. The benefits of significant event audit in primary care: a case study. *J Clin Governance*

32 Thomsen LA et al. Systematic review of the incidence and characteristics of preventable adverse drug events in  
33 ambulatory care. *Ann Pharmacother* 2007; 41: 1411–1426.

34 TorelloLserte J, Ferrando JC, Lainez M, Morillas MG, Gonzalez AA. [Adverse reactions to drugs reported by the primary  
35 care physicians of Andalusia. Analysis of underreporting] *Atencion Primaria* 1994; 13(6): 307-11.

36 van der Hooft CS, Jong GW, Dieleman JP, Verhamme KMC, van der Cammen TJM, Stricker BHCH, Sturkenboom MCJM.  
37 Inappropriate drug prescribing in older adults: The updated 2002 Beers criteria - A population-based cohort study.  
38 *British Journal of Clinical Pharmacology* 2005; 60(2):137-44.

39 Velo GP, Minuz P. Medication errors: Prescribing faults and prescription errors. *British Journal of Clinical Pharmacology*  
40 2009;67(6):624-28.

41 Wetzels R, Wolters R, van Weel C, Wensing M. Mix of methods is needed to identify adverse events in general practice: a  
42 prospective observational study. *BMC Family Practice* 2008;9:35.

43 Wilson R, Runciman W, Gibberd R, Harrison B, et al. The Quality in Australian Health Care Study. *Med J Aust* 1995;  
44 163(9): 458-71.

45 Woods D, Thomas E, Holl J, Weiss K, Brennan T. Ambulatory care adverse events and preventable adverse events  
46 leading to a hospital admission. *Qual Saf Health Care* 2007; 16(2): 127-31.

47 Zuccotti G, Sato L . Malpractice risk in ambulatory settings: an increasing and underrecognized problem. *JAMA*  
48 2011;305:2464–5.

## Online content

Online material is available at:

[http://www.crd.york.ac.uk/PROSPERO/display\\_record.asp?ID=CRD42012002304](http://www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42012002304)

and the PDFs are available as follows:

- Search strategy:  
[http://www.crd.york.ac.uk/PROSPEROFILES/2304\\_STRATEGY\\_20120323.pdf](http://www.crd.york.ac.uk/PROSPEROFILES/2304_STRATEGY_20120323.pdf)
- Full protocol:  
[http://www.crd.york.ac.uk/PROSPEROFILES/2304\\_PROTOCOL\\_20120323.pdf](http://www.crd.york.ac.uk/PROSPEROFILES/2304_PROTOCOL_20120323.pdf)



References

1 Abramson EL, Bates DW, Jenter C, Volk LA, Barron Y, Quaresimo J, et al. Ambulatory prescribing incidents among community-based providers in two states. *J Am Med Inform Assoc* 2012;19(4):644-8

2 Abramson EL1, Malhotra S, Osorio SN, Edwards A, Cheriff A, Cole C, Kaushal R. A long-term follow-up evaluation of electronic health record prescribing safety. *J Am Med Inform Assoc* 2013;20(e1):e52-58

3 A-Elgayoum SME, El-Feki AE-KA, Mahgoub BA, El-Rayah E-A, Giha HA: Malaria overdiagnosis and burden of malaria misdiagnosis in the suburbs of central Sudan: special emphasis on artemisinin-based combination therapy era. *Diagnostic Microbiology & Infectious Disease*, 2009; 64(1):20-26.

4 Al Khaja KA, Sequeira RP, Damanhori AH. Medication prescribing incidents pertaining to cardiovascular/antidiabetic medications: a prescription audit in primary care. *Fundam Clin Pharmacol* 2011 Jan 25.

5 Al Khaja KAJ, Sequeira RP, Al-Ansari TM, Damanhori AHH: Prescription writing skills of residents in a family practice residency programme in Bahrain. *Postgraduate Medical Journal*, 2007; 84(990):198-204.

6 Al Khaja KAJ, Damanhori AHH, Al-Ansari TM, Sequeira RP. Topical corticosteroids in infants: prescribing pattern and prescribing incidents in Bahrain. *Pharmacy World & Science* 2007;29(4):395-399.

7 Al Khaja KAJ, Al Ansari TM, Damanhori AHH, Sequeira R. Evaluation of drug utilization and prescribing incidents in infants: a primary care prescription-based study. *Health Policy* 2007;81(2-3):350-357.

8 Al Khaja KAJ, Sequeira RP, Al-Ansari T, Damanhori AHH, James H, Handu SS. Pediatric iron preparations for infants in Bahrain: some therapeutic concerns. *International Journal of Clinical Pharmacology & Therapeutics* 2010;48(3):200-205.

9 Al-Agilly S, Neville RG, Robb H, Riddell S: Involving patients in checking the validity of the NHS shared record: A single practice pilot. *Informatics in primary care* 2007, 15 (4):217-220.

10 APEAS Study. Patient Safety in primary Health Care. Madrid: Ministry of Health and Consumer Affairs, 2008

11 Aranaz-Andrés JM, Aibar C, Limón R, Mira JJ, Vitaller J, Agra Y, Terol E. A study of the prevalence of adverse events in primary healthcare in Spain. *Eur J Public Health* 2012, 22(6):921-925.

12 Avery AJ1, Ghaleb M, Barber N, Dean Franklin B, Armstrong SJ, Serumaga B, Dhillon S, Freyer A, Howard R, Talabi O, Mehta RL. The prevalence and nature of prescribing and monitoring incidents in English general practice: a retrospective case note review. *Br J Gen Pract* 2013;63(613):e543-553.

13 Beyer M, Gerlach FM: Medication incidents in German general practice: A research report. [German]. *Zeitschrift fur Allgemeinmedizin* 2003, 79 (7):332-335.

14 Beyer M, Rohe J, Rusitska M, Blauth E, Gerlach FM: The German incident reporting system for general practice: Structure, first results. [German]. *Zeitschrift fur Allgemeinmedizin* 2005, 81 (4):147-153.

15 Bhasale A: The wrong diagnosis: identifying causes of potentially adverse events in general practice using incident monitoring. *Fam Pract* 1998, 15(4):308-318

16 Bradbury F: How important is the role of the physician in the correct use of a drug? An observational cohort study in general practice. *International Journal of Clinical Practice* (Supplement) 2004;58:27-33.

17 Bradley MC, Fahey T, Cahir C, Bennett K, O'Reilly D, Parsons C, Hughes CM. Potentially inappropriate prescribing and cost outcomes for older people: a cross-sectional study using the Northern Ireland Enhanced Prescribing Database. *Eur J Clin Pharmacol*. 2012 Oct;68(10):1425-33.

18 Bregnhøj L, Thirstrup S, Kristensen MB, Bjerrum L, Sonne J: Prevalence of inappropriate prescribing in primary care. *Pharmacy World and Science* 2007;29(3):109-115.

19 Brekke M, Rognstad S, Straand J, Furu K, Gjelstad S, Bjørner T, Dalen I: Pharmacologically inappropriate prescriptions for elderly patients in general practice: How common? Baseline data from The Prescription Peer Academic Detailing (Rx-PAD) study. *Scand J Prim Health Care* 2008, 26(2):80-85.

20 Brenner S, Detz A, López A, Horton C, Sarkar U.: Signal and noise: applying a laboratory trigger tool to identify adverse drug events among primary care patients. *BMJ Qual Saf* 2012;21(8):670-675.

21 Britt H, Miller GC, Steven ID, Howarth GC, Nicholson PA, Bhasale AL, Norton KJ: Collecting data on potentially harmful events: a method for monitoring incidents in general practice. *Fam Pract* 1997, 14(2):101-106.

22 Casalino LP, Dunham D, Chin MH, Bielang R, Kistner EO, Karrison TG, Ong MK, Sarkar U, McLaughlin MA, Meltzer DO: Frequency of failure to inform patients of clinically significant outpatient test results. *Archives of Internal Medicine*

## How Safe is Primary Care? A Systematic Review

2009;169(12):1123-1129.

- 23 Clark RC, Maxwell SRJ, Kerr S, Cuthbert M, Buchanan D, Steinke D, Webb DJ, Bateman ND: The influence of primary care prescribing rates for new drugs on spontaneous reporting of adverse drug reactions. *Drug Safety* 2007;30(4):357-366.
- 24 Britten N, Brant S, Cairns A, Hall WW, Jones I, Salisbury C, Virji A, Herxheimer A. Continued prescribing of inappropriate drugs in general practice. *Journal of Clinical Pharmacy & Therapeutics*, 1995;20(4):199-205.
- 25 De Wet C, Bowie P: The preliminary development and testing of a global trigger tool to detect incident and patient harm in primary-care records. *Postgraduate Medical Journal* 2009;85(1002):176-180.
- 26 De Wilde S, Carey IM, Harris T, Richards N, Victor C, Hilton SR, Cook DG. Trends in potentially inappropriate prescribing amongst older UK primary care patients. *Pharmacoepidemiol Drug Saf* 2007;16(6):658-667.
- 27 Diamond MR, Kamien M, Sim MG, Davis J: A critical incident study of general practice trainees in their basic general practice term. *Med J Aust* 1995, 162(6):321-324.
- 28 Dovey SM, Meyers DS, Phillips Jr RL, Green LA, Fryer GE, Galliher JM, Kappus J, Grob P: A preliminary taxonomy of medical incidents in family practice. *Quality and Safety in Health Care* 2002; 11(3):233-238.
- 29 Eggleton KS1, Dovey SM. Using triggers in primary care patient records to flag increased adverse event risk and measure patient safety at clinic level. *N Z Med J*. 2014 Mar 7;127(1390):45-52.
- 30 Elder NC, Dovey SM: Classification of medical incidents and preventable adverse events in primary care: A synthesis of the literature. *Journal of Family Practice* 2002, 51 (11):927-932.
- 31 Elder NC, Vonder Meulen M, Cassedy A: The identification of medical incidents by family physicians during outpatient visits. *Annals of Family Medicine*, 2004; 2(2):125-129.
- 32 Ely JW, Kaldjian LC, D'Alessandro DM. Diagnostic incidents in primary care: lessons learned. *J Am Board Fam Med* 2012;25(1):87-97.
- 33 Farrow SC, Zeuner D, Hall C: Improving infection control in general practice. *The Journal of the Royal Society for the Promotion of Health* 1999, 119(1):17-22.
- 34 Fernald DH, Pace WD, Harris DM, West DR, Main DS, Westfall JM: Event reporting to a primary care patient safety reporting system: a report from the ASIPS collaborative. *Annals of Family Medicine*, 2004; 2(4):327-332.
- 35 Field TS, Gurwitz JH, Harrold LR, Rothschild JM, Debellis K, Seger AC, Fish LS, Garber L, Kelleher M, Bates DW: Strategies for detecting adverse drug events among older persons in the ambulatory setting. *Journal of the American Medical Association* 2004, 11 (6):492-498.
- 36 Gaal S, Verstappen W, Wolters R, Lankveld H, van Weel C, Wensing M: Prevalence and consequences of patient safety incidents in general practice in the Netherlands: a retrospective medical record review study. *Implementation Science*;2011;6(1):37.
- 37 Garfield S, Barber N, Walley P, Willson A, Eliasson L. Quality of medication use in primary care--mapping the problem, working to a solution: a systematic review of the literature. *BMC Med*. 2009 Sep 21;7:50
- 38 Gehring K, Schwappach DLB, Battaglia M, Buff R, Huber F, Sauter P, Wieser M. Frequency of and Harm Associated With Primary Care Safety Incidents. *Am J Manag Care*. 2012;18(9):e323-e337.
- 39 Gurwitz JH, Field TS, Harrold LR, Rothschild J, Debellis K, Seger AC, Cadoret C, Fish LS, Garber L, Kelleher M et al: Incidence and preventability of adverse drug events among older persons in the ambulatory setting. *JAMA: Journal of the American Medical Association* 2003, 289(9):1107-1117.
- 40 Hansen LB, Fernald D, Araya-Guerra R, Westfall JM, West D, Pace W. Pharmacy clarification of prescriptions ordered in primary care: A report from the applied strategies for improving patient safety (ASIPS) collaborative. *Journal of the American Board of Family Medicine* 2006;19(1):24-30.
- 41 Hickner J, Graham DG, Elder NC, Brandt E, Emsermann CB, Dovey S, Phillips R: Testing process incidents and their harms and consequences reported from family medicine practices: a study of the American Academy of Family Physicians National Research Network. *Qual Saf Health Care* 2008, 17(3):194-200.
- 42 Hickner J, Zafar A, Kuo GM, Fagnan LJ, Forjuoh SN, Knox LM, Lynch JT, Stevens BK, Pace WD, Hamlin BN, Scherer H, Hudson BL, Oppenheimer CC, Tierney WM. Field test results of a new ambulatory care Medication Incident and Adverse Drug Event Reporting System-MEADERS. *Ann Fam Med*. 2010 Nov-Dec;8(6):517-25.
- 43 Hildebrandt DE, Westfall JM, Fernald DH, Pace WD: Harm resulting from inappropriate telephone triage in primary care. *Journal of the American Board of Family Medicine* 2006, 19 (5):437-442.
- 44 Hoffmann B, Beyer M, Rohe J, Gensichen J, Gerlach FM: "Every incident counts": a web-based incident reporting and learning system for general practice. *Qual Saf Health Care* 2008, 17(4):307-312.
- 45 Holden J, O'Donnell S, Brindley J, Miles L: Analysis of 1263 deaths in four general practices. *Br J Gen Pract* 1998, 48(432):1409-1412.



1  
2 46 Honigman B, Lee J, Rothschild J, Light P, Pulling RM, Yu T, Bates DW. Using computerized data to identify adverse drug events  
3 in outpatients. *J Am Med Inform Assoc.* 2001 May-Jun;8(3):254-66  
4  
5 47 Howard M, Dolovich L, Kaczorowski J, Sellors C, Sellors J. Prescribing of potentially inappropriate medications to elderly people.  
6 *Family Practice* 2004;21(3):244-247.  
7  
8 48 Ilboudo TP, Chou YJ, Huang N. Assessment of providers' referral decisions in rural Burkina Faso: a retrospective analysis of  
9 medical records. *BMC Health Serv Res.* 2012 Mar 8;12:54  
10  
11 49 Kennedy AG, Littenberg B, Senders JW: Using nurses and office staff to report prescribing incidents in primary care. *Int J Qual*  
12 *Health Care* 2008, 20(4):238-245.  
13  
14 50 Khoja T, Neyaz Y, Qureshi NA, Magzoub MA, Haycox A, Walley T. Medication incidents in primary care in Riyadh city, Saudi  
15 Arabia. *Eastern Mediterranean Health Journal* 2011;17(2):156-160.  
16  
17 51 Khoja TA, Al-Shammari SAI, Farag MK, Al-Mazrou Y: Quality of prescribing at primary care centers in Saudi Arabia. *Journal of*  
18 *Pharmacy Technology* 1996, 12 (6):284-288.  
19  
20 52 Khoo EM, Lee WK, Sararaks S, Abdul Samad A, Liew SM, Cheong AT, Ibrahim MY, Su SH, Mohd Hanafiah AN, Maskon K, Ismail  
21 R, Hamid MA. Medical incidents in primary care clinics--a cross sectional study. *BMC Fam Pract.* 2012 Dec 26;13:127.  
22  
23 53 Kingston-Reichers J, Ospina M, Jonsson E, Childs P, Mcleod L and Mazted J. (2010). Patient Safety in Primary care. Edmonton,  
24 AB: Canadian Patient Safety Institute and BC Patient Safety and Quality Council  
25  
26 54 Koper D, Kamenski G, Flamm M, Böhmendorfer B, Sönnichsen A. Frequency of medication incidents in primary care patients with  
27 polypharmacy. *Fam Pract.* 2013;30(3):313-9.  
28  
29 55 Koper D1, Kamenski G, Flamm M, Böhmendorfer B, Sönnichsen A. Frequency of medication incidents in primary care patients with  
30 polypharmacy. *Fam Pract* 2013;30(3):313-319.  
31  
32 56 Kostopoulou O, Delaney B: Confidential reporting of patient safety events in primary care: results from a multilevel classification of  
33 cognitive and system factors. *Qual Saf Health Care* 2007, 16(2):95-100.  
34  
35 57 Kuo GM, Phillips RL, Graham D, Hickner JM: Medication incidents reported by USA family physicians and their office staff. *Qual*  
36 *Saf Health Care* 2008, 17(4):286-290.  
37  
38 58 Kuzel AJ, Woolf SH, Gilchrist VJ, Engel JD, LaVeist TA, Vincent C, Frankel RM: Patient reports of preventable problems and  
39 harms in primary health care. *Ann Fam Med* 2004, 2(4):333-340.  
40  
41 59 Leon AC, Portera L, Olsson M, Kathol R, Farber L, Lowell KN, Sheehan DV: Diagnostic incidents of primary care screens for  
42 depression and panic disorder. *International Journal of Psychiatry in Medicine* 1999, 29(1):1-12.  
43  
44 60 Lund BC, Carnahan RM, Egge JA, Chrischilles EA, Kaboli PJ. Inappropriate prescribing predicts adverse drug events in older  
45 adults. *Annals of Pharmacotherapy* 2010;44(6):957-963.  
46  
47 61 Makeham M, Dovey S, Runciman W and Larizgoitia I. Methods and measures used in primary care patient safety research.  
48 Available online at [www.who.int/patientsafety/research/methods\\_measures/primary\\_care\\_ps\\_research/en/index.html](http://www.who.int/patientsafety/research/methods_measures/primary_care_ps_research/en/index.html) (Last  
49 accessed 31 July 2014)  
50  
51 62 Makeham MA, Dovey SM, County M, Kidd MR: An international taxonomy for incidents in general practice: a pilot study. *Med J*  
52 *Aust* 2002, 177(2):68-72.  
53  
54 63 Makeham MA, Kidd MR, Saltman DC, Mira M, Bridges-Webb C, Cooper C, Stromer S: The Threats to Australian Patient Safety  
55 (TAPS) study: incidence of reported incidents in general practice. *Med J Aust* 2006;185(2):95-98.  
56  
57 64 Makeham MA, Stromer S, Bridges-Webb C, Mira M, Saltman DC, Cooper C, Kidd MR: Patient safety events reported in general  
58 practice: a taxonomy. *Qual Saf Health Care* 2008, 17(1):53-57.  
59  
60 65 Martinez Sanchez A, Campos RM: Detection of prescribing related problems at the community pharmacy. *International Journal of*  
*Clinical Pharmacy* 2011;33(1):66-69.  
61  
62 66 McKay J, Bradley N, Lough M, Bowie P. A review of significant events analysed in general practice: implications for the quality  
63 and safety of patient care. *BMC family practice* 2009;10:61.  
64  
65 67 McKay J, de Wet C, Kelly M, Bowie P. Applying the Trigger Review Method after a brief educational intervention: potential for  
66 teaching and improving safety in GP specialty training? *BMC Med Educ* 2013;13:117.  
67  
68 68 Miller GC, Britth HC, Valenti L: Adverse drug events in general practice patients in Australia. *Medical Journal of Australia*, 2006;  
69 184(7):321-324.  
70  
71 69 Montastruc P, Rey S, Rey C, M.-E LL, Montastruc JL: A prospective intensive study of adverse drug reactions in rural medicine.  
72 *Drug Investigation* 1993, 6 (6):337-339.  
73  
74 70 Murie J, McGhee C: Clinical governance in action. Assessing risk by analysing significant events in primary care. *Quality in*  
75 *Primary Care* 2003, 11(3):205-211.

## How Safe is Primary Care? A Systematic Review

- 71 Neville RG, Robertson F, Livingstone S, Crombie IK: A classification of prescription incidents. *The Journal of the Royal College of General Practitioners* 1989;39(320):110-112.
- 72 Nicholson D, Hersh W, Gandhi TK, Weingart SN, Bates DW: Medication incidents: not just a "few bad apples". *Journal of Clinical Outcomes Management* 2006, 13(2):114-116.
- 73 O'Beirne M, Sterling PD, Zwicker K, Hebert P, Norton PG. Safety incidents in family medicine. *BMJ Qual Saf* 2011;20(12):1005-10
- 74 Olaniyan JO1, Ghaleb M, Dhillon S, Robinson P. Safety of medication use in primary care. *Int J Pharm Pract* (published online June 2014).
- 75 Paille F, Pissochet P. Drug interactions in primary health care: Prospective study of 896 patients treated for hypertension. [French]. *Therapie* 1995;50(3):253-258.
- 76 Pandit NB, Choudhary SK: Unsafe injection practices in Gujarat, India. *Singapore Medical Journal* 2008, 49 (11):936-939.
- 77 Pariser RJ, Pariser DM: Primary care physicians' incidents in handling cutaneous disorders. A prospective survey. *Journal of the American Academy of Dermatology*, 1987; 17(2 Pt 1):239-245.
- 78 Pearson A and Aromataris A. Patient Safety Primary Healthcare: a review of the literature (2009 – Australian Commission on Safety and Quality in healthcare. Available online at [www.health.gov.au/internet/safety/publishing.nsf/Content/DBDB4EAE1386D1ABCA257753001ECA09/\\$File/26889-Literature-Review.PDF](http://www.health.gov.au/internet/safety/publishing.nsf/Content/DBDB4EAE1386D1ABCA257753001ECA09/$File/26889-Literature-Review.PDF) (Last accessed December 2014)
- 79 Phillips Jr RL, Bartholomew LA, Dovey SM, Fryer Jr GE, Miyoshi TJ, Green LA: Learning from malpractice claims about negligent, adverse events in primary care in the United States. *Quality and Safety in Health Care* 2004, 13 (2):121-126.
- 80 Phillips RL, Dovey SM, Graham D, Elder NC and Hickner JM. Learning from different lenses: reports of medical incidents in primary care by clinicians, staff and patients. *J Patient Saf* 2006; 2(3): 140 -146
- 81 Rosser W, Dovey S, Bordman R, White D, Crighton E, Drummond N: Medical incidents in primary care: results of an international study of family practice. *Canadian family physician Medecin de famille canadien* 2005, 51:386-387.
- 82 Rubin G, George A, Chinn DJ, Richardson C: Incidents in general practice: development of an incident classification and pilot study of a method for detecting incidents. *Qual Saf Health Care* 2003, 12(6):443-447.
- 83 Ryan C, O'Mahony D, Kennedy J, Weedle P, Byrne S. Potentially inappropriate prescribing in an Irish elderly population in primary care. *British Journal of Clinical Pharmacology* 2009;68(6):936-947.
- 84 Sandars J, Esmail A: The frequency and nature of medical incident in primary care: understanding the diversity across studies. *Fam Pract* 2003, 20(3):231-236.
- 85 Sayers YM, Armstrong P, Hanley K: Prescribing incidents in general practice: A prospective study. *European Journal of General Practice* 2009;15(2):81-83.
- 86 Schiff GD, Puopolo AL, Huben-Kearney A, Yu W, Keohane C, McDonough P, Ellis BR, Bates DW, Biondolillo M. Primary care closed claims experience of Massachusetts malpractice insurers. *JAMA Intern Med* 2013;173(22):2063-2068.
- 87 Schneider JK, Mion LC, Frengley JD. Adverse drug reactions in an elderly outpatient population. *Am J Hosp Pharm.* 1992;49(1):90-96.
- 88 Schwappach DL, Gehring K, Battaglia M, Buff R, Huber F, Sauter P, Wieser M. Threats to patient safety in the primary care office: concerns of physicians and nurses. *Swiss Med Wkly.* 2012 Jun 6;142:w13601
- 89 Shaughnessy AF, Nickel RO. Prescription-writing patterns and incidents in a family medicine residency program. *Journal of Family Practice* 1989;29(3):290-295.
- 90 Singh H, Giardina TD, Meyer AN, Forjuoh SN, Reis MD, Thomas EJ. Types and origins of diagnostic incidents in primary care settings. *JAMA Intern Med.* 2013 Mar 25;173(6):418-25.
- 91 Smith PC, Araya-Guerra R, Bublitz C, Parnes B, Dickinson LM, Van Vorst R, Westfall JM, Pace WD: Missing clinical information during primary care visits. *Journal of the American Medical Association* 2005, 293 (5):565-571.
- 92 Smits M, Huibers L, Kerssemeijer B, de Feijter E, Wensing M, Giesen P: Patient safety in out-of-hours primary care: a review of patient records. *BMC Health Services Research*, 2010; 10:335.
- 93 Statham MO, Sharma A, Pane AR: Misdiagnosis of acute eye diseases by primary health care providers: incidence and implications. *Medical Journal of Australia*, 2008; 189(7):402-404.
- 94 Stewart AL, Lynch KJ. Identifying discrepancies in electronic medical records through pharmacist medication reconciliation. *J Am Pharm Assoc.* 2012;52:59-68.
- 95 Straand J, Rokstad KS. Elderly patients in general practice: diagnoses, drugs and inappropriate prescriptions. A report from the More & Romsdal Prescription Study. *Family Practice* 1999;16(4):380-388.

1  
2 96 Tam KWT, Kwok HK, Fan YMC, Tsui KB, Ng KK, Ho KYA, Lau KT, Chan YC, Tse CWC, Lau CM: Detection and prevention of  
3 medication misadventures in general practice. *International Journal for Quality in Health Care* 2008, 20 (3):192-199.  
4  
5 97 Tilyard M, Dovey S, Hall K: Avoiding and fixing medical incidents in general practice: prevention strategies reported in the  
6 Linnaeus Collaboration's Primary Care International Study of Medical Incidents. *N Z Med J* 2005, 118(1208):U1264.  
7  
8 98 Tomlin A, Reith D, Dovey S, Tilyard M. Methods for retrospective detection of drug safety signals and adverse events in electronic  
9 general practice records. *Drug Saf.* 2012;35(9):733-43.  
10  
11 99 Tsang C, Majeed A, Aylin P. Routinely recorded patient safety events in primary care: a literature review. *Fam Pract* 2012;29(1):8-  
12 15  
13 100 Tsang C, Bottle A, Majeed A, Aylin P. Adverse events recorded in English primary care: observational study using the General  
14 Practice Research Database. *Br J Gen Pract* 2013;63(613):e534-42.  
15  
16 101 Tse J, You W. How accurate is the electronic health record? - a pilot study evaluating information accuracy in a primary care  
17 setting. *Stud Health Technol Inform* 2011;168:158-64  
18  
19 102 Van Der Hooft CS, G.W T.J, Dieleman JP, Verdamme KMC, Van Der Cammen TJM, Stricker BHC, Sturkenboom MCJM:  
20 Inappropriate drug prescribing in older adults: The updated 2002 Beers criteria - A population-based cohort study. *British Journal*  
21 *of Clinical Pharmacology* 2005, 60 (2):137-144.  
22  
23 103 van Dulmen SA, Tacke MA, Staal JB, Gaal S, Wensing M, Nijhuis-van der Sanden MW. Patient safety in primary allied health  
24 care: what can we learn from incidents in a Dutch exploratory cohort study? *Med Care.* 2011, 49(12):1089-1096.  
25  
26 104 Wahls TL, Cram PM: The frequency of missed test results and associated treatment delays in a highly computerized health  
27 system. *BMC Fam Pract* 2007, 8:32.  
28  
29 105 Wallace E1, Lowry J, Smith SM, Fahey T. The epidemiology of malpractice claims in primary care: a systematic review. *BMJ Open*  
30 2013;3(7). pii: e002929.  
31  
32 106 Weingart SN, Gandhi TK, Seger AC, Seger DL, Borus J, Burdick E, Leape LL, Bates DW: Patient-reported medication symptoms  
33 in primary care. *Archives of Internal Medicine* 2005, 165 (2):234-240.  
34  
35 107 Wetzels R, Wolters R, Van Weel C, Wensing M: Harm caused by adverse events in primary care: A clinical observational study.  
36 *Journal of Evaluation in Clinical Practice* 2009, 15 (2):323-327.  
37  
38 108 Woolf SH, Kuzel AJ, Dovey SM, Phillips RL, Jr.: A string of mistakes: the importance of cascade analysis in describing, counting,  
39 and preventing medical incidents. *Ann Fam Med* 2004, 2(4):317-326.  
40  
41 109 Zavaleta-Bustos M, Castro-Pastrana LI, Reyes-Hernandez I, Lopez-Luna MA, Bermudez-Camps IB: Prescription incidents in a  
42 primary care university unit: Urgency of pharmaceutical care in Mexico. *Revista Brasileira de Ciencias Farmaceuticas/Brazilian*  
43 *Journal of Pharmaceutical Sciences* 2008;44(1):115-125.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60