

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/105301/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Meltzer-Brody, S., Larsen, J. T., Petersen, L., Guintivano, J., Di Florio, Arianna , Miller, W. C., Sullivan, P. F. and Munk-Olsen, T. 2018. Adverse life events increase risk for postpartum psychiatric episodes: a population based epidemiologic study. *Depression and Anxiety* 35 (2) , pp. 160-167. 10.1002/da.22697

Publishers page: <http://dx.doi.org/10.1002/da.22697>


Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Figure 1. Cumulative incidence of any postpartum psychiatric diagnosis by number of different adversities experienced at age 0-15 years.



Cumulative incidence calculated using competing risk regression with death as competing event.