

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/108387/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

El-Hiti, Gamal A., Smith, Keith, Hegazy, Amany S., Alshammari, Mohammed B. and Kariuki, Benson M. 2018. 1-(2-Bromo-4-methylphenyl)-3,3-dimethylthiourea. IUCrData 3 (1) 10.1107/S2414314618000457 file

Publishers page: <http://dx.doi.org/10.1107/S2414314618000457>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1-(2-Bromo-4-methylphenyl)-3,3-dimethylthiourea

Gamal A. El-Hiti,^{a*} Keith Smith,^b Amany S. Hegazy,^b Mohammed B. Alshammari^c and Benson M. Kariuki^{b‡}

^aCornea Research Chair, Department of Optometry, College of Applied Medical Sciences, King Saud University, PO Box 10219, Riyadh 11433, Saudi Arabia, ^bSchool of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, Wales, and ^cChemistry Department, College of Sciences and Humanities, Prince Sattam bin Abdulaziz University, PO Box 83, Al-Kharj 11942, Saudi Arabia. *Correspondence e-mail: gelhiti@ksu.edu.sa

Received 3 January 2018

Accepted 8 January 2018

Edited by J. Simpson, University of Otago, New Zealand


‡ Additional corresponding author, e-mail: kariukib@cardiff.ac.uk.

Keywords: crystal structure; hydrogen bonding; thiourea; synthesis.

CCDC reference: 1815356

Structural data: full structural data are available from iucrdata.iucr.org

The bromomethylphenyl and dimethylthiourea groups of the molecule of the title compound, C₁₀H₁₃BrN₂S, are inclined to one another at an interplanar angle of 55.13 (6)°. In the crystal, molecules are stacked along the *b* axis and intermolecular N—H···S contacts form chains of molecules along [010].



Structure description

Thioureas show various biological activities (Yao *et al.*, 2012; Kocyigit-Kaymakcioglu *et al.*, 2013; Korkmaz *et al.*, 2015; Yang *et al.*, 2015; Tahir *et al.*, 2015; Shakeel *et al.*, 2016) and therefore their syntheses are always of interest (Kong *et al.*, 2015; Nguyen *et al.*, 2014; Maki *et al.*, 2014; Chau *et al.* 2014; Maddani & Prabhu, 2010). The X-ray crystal structures of some 1-(2-bromophenyl)-3,3-dimethylthiourea derivatives have been published recently (El-Hiti *et al.*, 2014, 2017): as part of our ongoing studies in this area, we now describe the synthesis and structure of the title compound.

The asymmetric unit comprises one molecule of the title compound (Fig. 1). The molecule is not planar, as indicated by an intramolecular interplanar angle of 55.13 (6) between the bromomethylphenyl and dimethylthiourea groups.

In the crystal, molecules are stacked along the *b*-axis and N—H···S intermolecular contacts, Table 1, form chains of molecules along [010], Fig. 2.

Synthesis and crystallization

The title compound was synthesized by the reaction of equimolar quantities of 2-bromo-4-methylphenyl isothiocyanate and dimethylamine in dry dichloromethane at 20°C for 1 h. Water was added and the organic layer was separated, dried over anhydrous




Figure 1
ORTEP representation (50% probability) of the asymmetric unit of $C_{10}H_{13}BrN_2S$.




Figure 2
Crystal packing showing N—H...S contacts as dashed lines. H atoms not involved in hydrogen bonding have been omitted for clarity.

magnesium sulfate and evaporated under vacuum. The crude product was recrystallized using a solvent mixture of diethyl ether and hexane (1:2 by volume) to give colourless crystals of the title compound, m.p. 174–175°C (lit. 173–175°C; Smith *et al.*, 1996).

Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2.

Funding information

The project was supported by King Saud University, Deanship of Scientific Research, Research Chairs and Cardiff University.

References

Agilent (2014). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
 Cambridge Soft (2001). *CHEM3D Ultra*. Cambridge Soft Corporation, Cambridge, Massachusetts, USA.
 Chau, C.-M., Chuan, T.-J. & Liu, K.-L. (2014). *RSC Adv.* **4**, 1276–1282.
 El-Hiti, G. A., Smith, K., Hegazy, A. S., Alotaibi, M. H. & Kariuki, B. M. (2014). *Acta Cryst.* **E70**, o704.

Table 1
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N1-H1\cdots S1^i$	0.86	2.60	3.309 (2)	140

Symmetry code: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$.

Table 2
Experimental details.

Crystal data	
Chemical formula	$C_{10}H_{13}BrN_2S$
M_r	273.19
Crystal system, space group	Monoclinic, $P2_1/n$
Temperature (K)	296
a, b, c (Å)	12.2617 (7), 8.0222 (4), 12.7397 (7)
β (°)	112.380 (6)
V (Å ³)	1158.76 (12)
Z	4
Radiation type	Cu $K\alpha$
μ (mm ⁻¹)	6.22
Crystal size (mm)	0.22 × 0.12 × 0.05
Data collection	
Diffractometer	Agilent SuperNova, Dual, Cu at zero, Atlas
Absorption correction	Gaussian (<i>CrysAlis PRO</i> ; Agilent, 2014)
T_{min}, T_{max}	0.926, 0.976
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	7605, 2318, 2035
R_{int}	0.030
$(\sin \theta/\lambda)_{max}$ (Å ⁻¹)	0.624
Refinement	
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.038, 0.115, 1.06
No. of reflections	2318
No. of parameters	130
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{max}, \Delta\rho_{min}$ (e Å ⁻³)	0.54, -0.64

Computer programs: *CrysAlis PRO* (Agilent, 2014), *SHELXS2013* (Sheldrick, 2008), *SHELXL2013* (Sheldrick, 2015), *ORTEP-3 for Windows* and *WinGX* (Farrugia, 2012) and *CHEM3D Ultra* (Cambridge Soft, 2001).

El-Hiti, G. A., Smith, K., Hegazy, A. S., Alotaibi, M. H. & Kariuki, B. M. (2017). *Z. Kristallogr. New Cryst. Struct.* **232**, 31–32.
 Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
 Kocyyigit-Kaymakcioglu, B., Celen, A. O., Tabanca, N., Ali, A., Khan, S. I., Khan, I. A. & Wedge, D. E. (2013). *Molecules*, **18**, 3562–3576.
 Kong, X., Yao, Z., He, Z., Xu, W. & Yao, J. (2015). *Med. Chem. Commun.* **6**, 867–870.
 Korkmaz, N., Obaidi, O. A., Senturk, M., Astley, D., Ekinci, D. & Supuran, C. T. (2015). *J. Enzyme Inhib. Med. Chem.* **30**, 75–80.
 Maddani, M. R. & Prabhu, K. R. (2010). *J. Org. Chem.* **75**, 2327–2332.
 Maki, T., Tsuritani, T. & Yasukata, T. (2014). *Org. Lett.* **16**, 1868–1871.
 Nguyen, T. B., Ermolenko, L. & Al-Mourabit, A. (2014). *Synthesis*, **46**, 3172–3179.
 Shakeel, A., Altaf, A. A., Qureshi, A. M. & Badshah, A. (2016). *J. Drug Des. Med. Chem.* **2**, 10–20.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
 Smith, K., Shukla, A. P. & Matthews, I. (1996). *Sulfur Lett.* **20**, 121–137.
 Tahir, S., Badshah, A., Hussain, R. A., Tahir, M. N., Tabassum, S., Patujo, J. A. & Rauf, M. K. (2015). *J. Mol. Struct.* **1099**, 215–225.
 Yang, M., Pickard, A. J., Qiao, X., Gueble, M. J., Day, C. S., Kucera, G. L. & Bierbach, U. (2015). *Inorg. Chem.* **54**, 3316–3324.
 Yao, J., Chen, J., He, Z., Sun, W. & Xu, W. (2012). *Bioorg. Med. Chem.* **20**, 2923–2929.

full crystallographic data

IUCrData (2018). 3, x180045 [https://doi.org/10.1107/S2414314618000457]

1-(2-Bromo-4-methylphenyl)-3,3-dimethylthiourea

Gamal A. El-Hiti, Keith Smith, Amany S. Hegazy, Mohammed B. Alshammari and Benson M. Kariuki

1-(2-Bromo-4-methylphenyl)-3,3-dimethylthiourea

Crystal data

$C_{10}H_{13}BrN_2S$

$M_r = 273.19$

Monoclinic, $P2_1/n$

$a = 12.2617$ (7) Å

$b = 8.0222$ (4) Å

$c = 12.7397$ (7) Å

$\beta = 112.380$ (6)°

$V = 1158.76$ (12) Å³

$Z = 4$

$F(000) = 552$

$D_x = 1.566$ Mg m⁻³

Cu $K\alpha$ radiation, $\lambda = 1.54184$ Å

Cell parameters from 3751 reflections

$\theta = 4.2$ – 73.6 °

$\mu = 6.22$ mm⁻¹

$T = 296$ K

Block, colourless

$0.22 \times 0.12 \times 0.05$ mm

Data collection

Agilent SuperNova, Dual, Cu at zero, Atlas diffractometer

ω scans

Absorption correction: gaussian
(CrysAlis PRO; Agilent, 2014)

$T_{\min} = 0.926$, $T_{\max} = 0.976$

7605 measured reflections

2318 independent reflections

2035 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.030$

$\theta_{\max} = 74.1$ °, $\theta_{\min} = 4.3$ °

$h = -14 \rightarrow 15$

$k = -9 \rightarrow 9$

$l = -12 \rightarrow 15$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.038$

$wR(F^2) = 0.115$

$S = 1.06$

2318 reflections

130 parameters

0 restraints

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.072P)^2 + 0.3056P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.54$ e Å⁻³

$\Delta\rho_{\min} = -0.64$ e Å⁻³

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.26065 (3)	0.22909 (5)	0.48048 (3)	0.06494 (17)
S1	0.18412 (6)	0.22540 (8)	0.06795 (5)	0.0495 (2)
C2	0.1250 (2)	0.1405 (3)	0.3645 (2)	0.0454 (5)
N1	0.24617 (18)	0.0533 (3)	0.26170 (17)	0.0477 (5)
H1	0.2981	-0.0043	0.3139	0.057*
C1	0.1340 (2)	0.0699 (3)	0.26865 (19)	0.0416 (5)
N2	0.3930 (2)	0.0961 (3)	0.19709 (19)	0.0511 (5)
C8	0.2794 (2)	0.1197 (3)	0.18062 (19)	0.0422 (5)
C6	0.0329 (2)	0.0058 (3)	0.1861 (2)	0.0473 (6)
H6	0.0372	-0.0438	0.1218	0.057*
C10	0.4382 (3)	0.1476 (5)	0.1115 (3)	0.0662 (8)
H10A	0.3850	0.1113	0.0380	0.099*
H10B	0.5144	0.0984	0.1281	0.099*
H10C	0.4450	0.2668	0.1120	0.099*
C5	-0.0743 (2)	0.0146 (4)	0.1981 (2)	0.0495 (6)
H5	-0.1414	-0.0287	0.1415	0.059*
C3	0.0176 (3)	0.1482 (4)	0.3771 (2)	0.0524 (6)
H3	0.0137	0.1948	0.4424	0.063*
C4	-0.0834 (2)	0.0873 (4)	0.2936 (2)	0.0509 (6)
C9	0.4791 (2)	0.0276 (4)	0.3016 (3)	0.0614 (7)
H9A	0.4670	0.0750	0.3655	0.092*
H9B	0.5573	0.0538	0.3063	0.092*
H9C	0.4699	-0.0912	0.3019	0.092*
C7	-0.2015 (3)	0.0970 (5)	0.3057 (4)	0.0756 (9)
H7A	-0.1911	0.0734	0.3828	0.113*
H7B	-0.2545	0.0169	0.2562	0.113*
H7C	-0.2337	0.2069	0.2857	0.113*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0668 (3)	0.0716 (3)	0.0472 (2)	-0.01503 (14)	0.01142 (17)	-0.00997 (12)
S1	0.0580 (4)	0.0512 (4)	0.0351 (3)	-0.0007 (3)	0.0132 (3)	0.0014 (2)
C2	0.0495 (13)	0.0465 (13)	0.0379 (11)	-0.0033 (10)	0.0139 (10)	-0.0002 (9)
N1	0.0414 (10)	0.0630 (13)	0.0398 (10)	0.0077 (9)	0.0166 (9)	0.0107 (9)
C1	0.0420 (11)	0.0471 (12)	0.0380 (11)	0.0036 (9)	0.0179 (9)	0.0049 (9)
N2	0.0460 (11)	0.0609 (13)	0.0501 (11)	-0.0049 (10)	0.0226 (9)	0.0027 (10)
C8	0.0458 (12)	0.0453 (12)	0.0360 (11)	-0.0041 (9)	0.0161 (9)	-0.0058 (9)
C6	0.0494 (13)	0.0545 (14)	0.0372 (11)	0.0016 (11)	0.0155 (10)	-0.0024 (10)
C10	0.0652 (17)	0.081 (2)	0.0662 (18)	-0.0097 (16)	0.0408 (15)	-0.0004 (16)
C5	0.0405 (12)	0.0562 (14)	0.0474 (13)	0.0003 (10)	0.0117 (10)	0.0027 (11)
C3	0.0627 (15)	0.0547 (15)	0.0486 (13)	0.0032 (12)	0.0311 (12)	-0.0037 (11)
C4	0.0475 (13)	0.0550 (15)	0.0549 (14)	0.0060 (11)	0.0248 (11)	0.0075 (11)
C9	0.0436 (13)	0.0743 (19)	0.0638 (17)	-0.0032 (13)	0.0178 (12)	0.0084 (15)
C7	0.0566 (17)	0.091 (3)	0.090 (2)	0.0030 (16)	0.0412 (17)	0.000 (2)

Geometric parameters (\AA , $^\circ$)

Br1—C2	1.893 (2)	C10—H10B	0.9600
S1—C8	1.693 (2)	C10—H10C	0.9600
C2—C3	1.388 (4)	C5—C4	1.392 (4)
C2—C1	1.388 (3)	C5—H5	0.9300
N1—C8	1.355 (3)	C3—C4	1.379 (4)
N1—C1	1.418 (3)	C3—H3	0.9300
N1—H1	0.8600	C4—C7	1.515 (4)
C1—C6	1.383 (4)	C9—H9A	0.9600
N2—C8	1.340 (3)	C9—H9B	0.9600
N2—C9	1.456 (4)	C9—H9C	0.9600
N2—C10	1.459 (3)	C7—H7A	0.9600
C6—C5	1.382 (4)	C7—H7B	0.9600
C6—H6	0.9300	C7—H7C	0.9600
C10—H10A	0.9600		
C3—C2—C1	121.1 (2)	H10B—C10—H10C	109.5
C3—C2—Br1	118.91 (19)	C6—C5—C4	121.0 (2)
C1—C2—Br1	119.99 (19)	C6—C5—H5	119.5
C8—N1—C1	126.2 (2)	C4—C5—H5	119.5
C8—N1—H1	116.9	C4—C3—C2	120.5 (2)
C1—N1—H1	116.9	C4—C3—H3	119.7
C6—C1—C2	118.2 (2)	C2—C3—H3	119.7
C6—C1—N1	121.8 (2)	C3—C4—C5	118.4 (2)
C2—C1—N1	119.8 (2)	C3—C4—C7	121.0 (3)
C8—N2—C9	123.0 (2)	C5—C4—C7	120.5 (3)
C8—N2—C10	120.7 (2)	N2—C9—H9A	109.5
C9—N2—C10	116.1 (2)	N2—C9—H9B	109.5
N2—C8—N1	114.9 (2)	H9A—C9—H9B	109.5
N2—C8—S1	122.94 (19)	N2—C9—H9C	109.5
N1—C8—S1	122.14 (19)	H9A—C9—H9C	109.5
C5—C6—C1	120.7 (2)	H9B—C9—H9C	109.5
C5—C6—H6	119.6	C4—C7—H7A	109.5
C1—C6—H6	119.6	C4—C7—H7B	109.5
N2—C10—H10A	109.5	H7A—C7—H7B	109.5
N2—C10—H10B	109.5	C4—C7—H7C	109.5
H10A—C10—H10B	109.5	H7A—C7—H7C	109.5
N2—C10—H10C	109.5	H7B—C7—H7C	109.5
H10A—C10—H10C	109.5		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 \cdots S1 ⁱ	0.86	2.60	3.309 (2)	140

Symmetry code: (i) $-x+1/2, y-1/2, -z+1/2$.