

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/109625/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Myers, Kenneth A., White, Susan M., Mohammed, Shehla, Metcalfe, Kay A., Fry, Andrew E. , Wraige, Elisabeth, Vasudevan, Pradeep C., Balasubramanian, Meena and Scheffer, Ingrid E. 2018. Childhood-onset generalized epilepsy in Bainbridge-Ropers syndrome. *Epilepsy Research* 140 , pp. 166-170. 10.1016/j.eplepsyres.2018.01.014

Publishers page: <http://dx.doi.org/10.1016/j.eplepsyres.2018.01.014>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



## **Supporting Information**

Supplementary Video - Atypical absence seizure with loss of tone. An atypical absence seizure in patient #1. The patient is awake and standing when his father notices his breathing change.

The patient progressively loses tone and becomes completely limp. Following the event, he returns to baseline awareness almost immediately. Event duration is approximately 15 seconds.

The typical EEG pattern is shown in Figure B.