

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/111258/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Sosdian, S. M. , Rosenthal, Y. and Toggweiler, J. R. 2018. Deep Atlantic carbonate ion and CaCO_3 compensation during the Ice Ages. *Paleoceanography and Paleoclimatology* 33 (6) , pp. 546-562.
10.1029/2017PA003312

Publishers page: <http://dx.doi.org/10.1029/2017PA003312>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Deep Atlantic carbonate ion and CaCO_3 Compensation during the Ice Ages

S. M. Sosdian^{1*}, Y. Rosenthal^{2,3}, & J.R. Toggweiler⁴

¹Cardiff University, School of Earth & Ocean Sciences, Cardiff, CF10 3AT, UK

²Rutgers University, Department of Marine and Coastal Sciences, New Brunswick, NJ 08901, USA

³Rutgers University, Department of Earth and Planetary Sciences

⁴Geophysical Fluid Dynamics Laboratory, NOAA, Princeton, New Jersey, USA

*Corresponding author

Higher alkalinity is compensation for reduced CaCO_3 burial in the deep ocean in response to increased carbon sequestration in the deep ocean. This process accounts for about half of the reduction in glacial atmospheric CO_2 . To date our understanding of this process comes from benthic carbon isotope and CaCO_3 burial records. Here we present a 1.5 My orbitally resolved deep ocean calcite saturation record (ΔCO_3^{2-}) derived from benthic foraminiferal B/Ca ratios in the North Atlantic. Glacial ΔCO_3^{2-} declines across the mid-Pleistocene transition (MPT) suggesting increased sequestration of carbon in the deep Atlantic. The magnitude, timing, and structure of deep Atlantic Ocean ΔCO_3^{2-} parallels changes in % CaCO_3 and contrasts the small amplitude, anti-phased swings in IndoPacific ΔCO_3^{2-} and % CaCO_3 during the mid-to-late Pleistocene questioning the classic view of CaCO_3 compensatory mechanism. We propose that the increasing corrosivity of the deep Atlantic causes the locus of CaCO_3 burial to shift into the equatorial Pacific where the flux of CaCO_3 to the seafloor was sufficiently high to overcome low saturation and establish a new burial “hot spot”. Based on this mechanism, we propose that the persistently low ΔCO_3^{2-} levels at Marine Isotope Stages (MIS) 12, set the stage for the high pCO_2 levels at MIS 11 and subsequent interglacials via large swings in ocean alkalinity caused by shifts in CaCO_3 burial. Similarly, the development of classic (‘anti-correlated’) CaCO_3 patterns was driven by enhanced ocean stratification and an increase in deep ocean corrosivity in response to MPT cooling.

Keywords: B/Ca, benthic foraminifera, calcite saturation, North Atlantic, carbon dioxide, Pleistocene, CaCO_3 compensation

Key Points:

- North Atlantic deep ocean calcite saturation is reconstructed using B/Ca ratios in benthic foraminifera across the mid-to-late Pleistocene
- Deep ocean ΔCO_3^{2-} levels declined across the mid-Pleistocene and G/IG ΔCO_3^{2-} amplitude increased as ΔCO_3^{2-} became persistently under-saturated 400 ky ago
- Shifts in the locus of CaCO_3 burial maintains the global CaCO_3 balance on G/IG timescales

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1029/2017PA003312

1. Introduction

Mechanisms proposed to explain Pleistocene changes in atmospheric $p\text{CO}_2$ call for a transfer of carbon dioxide from the atmosphere to the glacial deep ocean by way of increased efficiency of the biological pump (Broecker & Peng, 1982; Broecker & Peng, 1987; Boyle et al. 1988). In contrast, upwelling of respired CO_2 and its transfer back to the atmosphere would explain the $p\text{CO}_2$ increase during the interglacials. However, as previously pointed out biological mechanisms cannot account for the full G/IG amplitude and duration of atmospheric $p\text{CO}_2$ during the late Pleistocene (Broecker and Peng, 1982; Boyle, 1988; Broecker and Peng, 1989; Archer and Maier-Reimer 1994; Toggweiler, 1999; Sigman & Boyle, 2000). This raises the likelihood that calcium carbonate (CaCO_3) compensation in response to changes in the amount of carbon sequestered in the deep ocean might have had substantial contribution to the G/IG changes in $p\text{CO}_2$ (Broecker and Peng, 1982; Boyle, 1988; Broecker and Peng, 1987; Toggweiler, 1999).

In this scenario, the sequestration of excess glacial metabolic CO_2 reduces carbon ion (CO_3^{2-}) in the deep ocean. As the deep ocean CO_3^{2-} is controlled by the balance between the supply of ingredients for CaCO_3 (calcium (Ca^{2+}) and bicarbonate (HCO_3^-)) into the ocean and removal via burial of CaCO_3 in the sediments, this imbalance promotes CaCO_3 dissolution until a new steady state is reached via adjustments in the lysocline, the depth in the ocean where CaCO_3 preservation (above) gives way to dissolution (below). The shoaling of the lysocline raises ocean alkalinity and acts to further lower atmosphere CO_2 . Current CaCO_3 compensation theory predicts, the ocean's CO_3^{2-} response should be fairly fast, within the range of 5-10 ky and atmospheric $p\text{CO}_2$ would change on the timescale set by the CO_3^{2-} response time (Broecker & Peng, 1987; Keir, 1988; Archer et al. 1997; Kohler & Fischer, 2006). If this mechanism did operate on G/IG timescales across the Pleistocene, we would expect to find evidence for relevant oscillations in deep ocean carbon storage, CO_3^{2-} content, lysocline in tandem with CaCO_3 preservation and dissolution patterns.

A second mechanism proposed to explain G/IG changes in deep ocean CO_3^{2-} and CaCO_3 burial patterns invokes sea level driven shifts in carbonate deposition a.k.a coral reef hypothesis (Berger, 1982; Berger and Keir, 1984; Opdyke and Walker, 1992). In this scenario, during glaciation, sea level drop reduces the shelf area and CaCO_3 deposition on continental shelves. This would act to increase whole ocean alkalinity and deep ocean CO_3^{2-} and plays a critical role in reducing atmospheric CO_2 . This idea implies that on G/IG

timescales whole ocean alkalinity and $p\text{CO}_2$ is dictated by sea level driven shifts in the partitioning of CaCO_3 deposition between continental shelf and deep ocean, or more specifically enhanced glacial preservation is tied to alkalinity supplied from the shelf areas during this period of sea level fall. If this hypothesis did operate on G/IG timescales then there should be a correlation between timing and amplitude of long-term changes in deep ocean CO_3^{2-} and sea level in the past.

Most of our knowledge for changes in Pleistocene deep ocean G/IG carbon inventory and sedimentary carbonate processes derives from benthic carbon isotope records ($\delta^{13}\text{C}_b$) and CaCO_3 burial and preservation fluxes. $\delta^{13}\text{C}_b$ records have been used to track the transfer of carbon between the organic and inorganic pools in the ocean, and records spanning the Pleistocene have also been interpreted to reflect large G/IG shifts in water mass distributions with implications for carbon storage in the deep ocean (Raymo et al. 1997; Clark et al. 2006; Skinner et al. 2009). However, interpretations of $\delta^{13}\text{C}_b$ records in terms of carbon storage and circulation are confounded as these proxies are impacted by a range of biogeochemical and physical processes, such as changes in preformed $\delta^{13}\text{C}_b$ signals of the source water (Lynch-Stieglitz et al. 1995; Raymo et al. 1997; Yu et al. 2008 and references herein). Pleistocene % CaCO_3 records from the deep Atlantic show enhanced dissolution during glacial periods relative to interglacials, whereas Pacific % CaCO_3 records indicate enhanced preservation during glacials relative to interglacials (Arrhenius, 1952; Farrell & Prell, 1989; Howard & Prell, 1994; deMenocal et al. 1997; Hodell et al. 2001; Sexton & Barker, 2012 and references here in). This classic ‘seesaw’ CaCO_3 preservation pattern has persisted across the last 800 ky and has been suggested to vary in response to vertical repositioning of the Atlantic lysocline (Berger & Winterer, 1974; Boyle, 1983; Farrell & Prell, 1989; Howard & Prell, 1994; deMenocal et al. 1997). However, interpretations of % CaCO_3 records in terms of variations of deep ocean CO_3^{2-} are confounded by the fact that in addition to dissolution they also reflect carbonate production and dilution by terrigenous material and thus cannot directly be related to changes in atmospheric $p\text{CO}_2$.

A robust assessment of deep ocean CO_3^{2-} across Pleistocene G/IG cycles awaits the development of basin-wide records of ocean carbonate chemistry to fully resolve the nature of the ocean buffering capacity and its role in driving G/IG $p\text{CO}_2$ cycles. As the CO_3^{2-} of the deep ocean is regulated by changes in mean ocean alkalinity, records of deep ocean CO_3^{2-} and % CaCO_3 provide insights into the compensation mechanism, coral reef hypothesis, and the

ability of the ocean to buffer changes in atmospheric CO₂. Also, the CO₃²⁻ of the deep ocean's respired carbon pool should track atmospheric pCO₂ which makes it a good proxy to investigate mechanisms affecting the carbon cycle and atmospheric pCO₂ (Broecker & Peng, 1987; Yu et al. 2010). Here we use boron to calcium ratios in the benthic foraminifer *Cibicidoides wuellerstorfi* as a means to investigate Pleistocene G/IG changes in deep North Atlantic Ocean CO₃²⁻.

Boron to calcium (B/Ca) ratios in epifaunal benthic foraminifera have been used to reconstruct seawater calcite saturation state ($\Delta[\text{CO}_3^{2-}] = [\text{CO}_3^{2-} \text{ in situ}] - [\text{CO}_3^{2-} \text{ sat}]$) based on available empirical calibrations (Yu & Elderfield, 2007; Brown et al. 2011; Rae et al. 2011; Raitzsch et al. 2011; Yu et al. 2013). Overall, the available evidence from B/Ca-based ΔCO_3^{2-} records show a divergent response related to depth and ocean basin across the last glacial cycle (Yu et al. 2010; Yu et al. 2013; Doss & Marchitto, 2013; Yu et al. 2014; Allen et al. 2015; Yu et al. 2016). Currently available, long Pleistocene records are mostly of low resolution documenting long-term trends in ΔCO_3^{2-} but not resolving the G/IG cyclicity (Rickaby et al. 2010; Elmore et al. 2015; Kender et al. 2016) except to two recent studies. In the North Atlantic, Lear et al (2016) put forth a suite (n=80) of trace metal records (Cd/Ca, B/Ca, U/Ca) across the mid-Pleistocene (0.4 to 1.1 Ma) in Deep Sea Drilling Program (DSDP) site 607 and show a decrease in glacial B/Ca values across the MPT. As G/IG changes in $\delta^{13}\text{C}_b$ occur alongside shifts in nutrient content and deep ocean CO₃²⁻, the authors attributed the cause to a change from dominantly northern water mass to dominantly southern source waters across the MPT and an increase in respired carbon pool of the glacial Atlantic. Beyond this they suggest that the increased corrosivity of bottom waters during glacials might have increased the ocean alkalinity and lowered atmospheric CO₂ (Lear et al. 2016). Additionally a study by Kerr et al. (2017), using orbitally resolved IndoPacific B/Ca records show that deep Pacific CO₃²⁻ concentration slightly increased during the glacial intervals while falling during the interglacials across the last 500 ky with similar patterns to the sediment CaCO₃ content. They argue that a global increase in CO₃²⁻ during glacial intervals is caused by the reduction of carbonate deposition on the shelves during low sea level stands (a.k.a 'coral reef hypothesis') across the last 450 ky (Berger, 1982; Opdyke & Walker, 1992). Here, we explore alternative mechanism, whereby the post-MPT Pacific and Atlantic CO₃²⁻ records are driven by basin-to-basin CaCO₃ compensation

In this paper we extend the DSDP site 607 North Atlantic record of Lear et al. (2016; n=80) and increase the number of samples by 280 analyses, alongside including a nearby piston core record (Chain 84-24-23 PC) to provide a complete orbitally resolved benthic foraminifera B/Ca reconstruction for the last 1.5 My through the mid to late Pleistocene. Combined with existing IndoPacific B/Ca records (Kerr et al., 2017), carbon isotope, and %CaCO₃ records (Ruddiman et al. 1989; Raymo et al. 1997) we investigate mechanisms controlling the deep ocean carbon budget of the North Atlantic and examine the physical and biogeochemical processes that can explain the observed G/IG changes in ΔCO_3^{2-} . The focus on the deep Atlantic stems from geochemical and modeling evidence suggesting the deep North Atlantic has played a significant role in carbon sequestration on G/IG timescales during the late Pleistocene (Toggweiler, 1999; Hain et al. 2010; Yu et al. 2016; Howe et al. 2016). We discuss the importance of the CaCO₃ cycle for the G/IG pCO₂ change in light of this 1.5 My Atlantic ΔCO_3^{2-} stack and propose that the enhancement of atmospheric CO₂ by CaCO₃ compensation is due to a change in the locus of CaCO₃ burial between the Atlantic and IndoPacific in response to the transfer of alkalinity between these ocean basins. Our new record also provides new insights into the mechanisms associated with the mid-Pleistocene transition (MPT) and mid-Brunhes Event (MBE).

2. Core material and methods

2.1 Hydrographic conditions & Age Model

We present benthic foraminiferal B/Ca records for Deep Sea Drilling Program (DSDP) site 607 (41°00'N; 32°57'W, 3427 m) on the western flank of the Mid-Atlantic Ridge supplemented with measurements from a nearby piston core (Chain 82-24-23 Piston Core (referred to hereafter as CHN82-24-23PC; 43°N, 31°W; water depth 3406 m; Figure 1). Both sites are situated within the path of North Atlantic Deep Water (NADW; T=2.6 °C, S=34.1) close to its northern sources. In the modern ocean, bottom water at these sites reflects the influence mostly of NADW (low nutrient, high $\delta^{13}\text{C}_b$ and ΔCO_3^{2-}) and to a lesser extent Antarctic Bottom Water (AABW; enriched nutrient, low $\delta^{13}\text{C}_b$ and ΔCO_3^{2-}) (Yu et al. 2008) (Figure 2). Modern ΔCO_3^{2-} (33.2 $\mu\text{mol/kg}$) are estimated from nearby GEOSECS data. The age model for both sites are based on LRO4 stack Lisiecki & Raymo (2005) and previously used in Sosdian & Rosenthal (2009).

2.2 Analytical Method

Well-preserved benthic foraminifera of the species *Cibicidoides wuellerstorfi* were picked from >150 μm fraction at sampling intervals of 15 cm providing an average temporal resolution of ~ 3 ky. Picked samples restricted to the morphology of *C. wuellerstorfi sensu stricto* for B/Ca analyses throughout the core (Rae et al. 2011). Pleistocene records of B/Ca should mainly reflect deep ocean ΔCO_3^{2-} variability due to the relatively long oceanic residence time of B (~ 10 My) and Ca (~ 1 My) and their conservative nature (LeMarchand et al., 2002). *C. wuellerstorfi* B/Ca were converted to deep sea ΔCO_3^{2-} using a sensitivity of $1.14 \mu\text{mol/mol}$ per $\mu\text{mol/kg}$ specific to this species obtained from the global core-top calibration from Yu et al. (2007) where $\text{B/Ca} (\mu\text{mol/mol}) = 1.14(\pm 0.04) \times \Delta\text{CO}_3^{2-} (\mu\text{mol/kg}) + 177.(\pm 1.0)$.

Samples were gently crushed between two glass plates and metal oxides visible within chambers were removed from the sample to minimize contamination. The fragments were transferred into acid-cleaned, dry 0.5 ml micro-centrifuge tubes. Crushed foraminiferal samples contained 3-15 individuals and ideally weighed from 0.200 mg to 0.400 mg, but due to low species abundance in certain intervals samples occasionally weighed < 0.200 mg. Samples were cleaned using the reductive or 'long' cleaning protocol (Boyle and Keigwin, 1985/6; Rosenthal et al., 1997). The effect of different cleaning methods on B/Ca ratios is minimal (Yu et al., 2007). Samples were cleaned and dissolved using reagents prepared with boron-free water as prepared by passing double distilled water over an anion exchange resin (Aldrich, amberlite IRA-743) to remove boron. Following cleaning, samples were dissolved in trace metal clean 0.065 N Nitric Acid (SEASTAR) and 100 μl of this solution was diluted with 300 μl of trace metal clean 0.5 N Nitric Acid to obtain a Ca concentration of about $3 \pm 1 \text{ mmol mol}^{-1}$. Diluted samples were analyzed for a suite of trace metal ratios using the Rutgers Sector Field Inductively Coupled Plasma-Mass Spectrometer (ICP-MS), on a Finnigan MAT Element-1 and Element-2 XR located at the Department of Marine and Coastal Sciences using methods adapted from Rosenthal et al., (1999). The method was slightly modified to accommodate for measurement of B/Ca ratios, specifically, ammonia gas was introduced in the Teflon-spray chamber during sample analysis to remove the memory effect associated with analysis of boron (Babila et al., 2014). Long-term precision, based on analysis of consistency standards across a range of B/Ca ratios is $\pm 4\%$. Analysis runs including CHN82-24-23PC B/Ca records were offset from the long-term mean B/Ca, as determined by the long-term average of the consistency standards, and to account for this we applied an analytical

offset to CHN82-24-23PC B/Ca values of 5% prior to including them in Figure 3. Al, and Ti were used as contaminant indicators and no long-term correspondence in trend exists between B/Ca and each elemental ratio. However, across MIS 25, high B/Ca values correspond with elevated Al and Ti and were removed from the final record.

2.3 Cross-Spectral Analysis & Trends in glacial and interglacial maxima

Here we use the ARAND package (Howell, 2001) to evaluate the phase and coherency between elemental and isotopic records from DSDP 607 and compare variables relative to benthic $\delta^{18}\text{O}$ in 100-ky world (i.e. from 140 to 750 ka, 100 ky periodicity). We interpolated all records to even intervals of 3-ky resolution prior to cross-spectral analysis and used the inverse of benthic $\delta^{18}\text{O}$ in the analysis. Proxy phase relationships are compared and presented in the representative time window during the “100 ky world” of the late Pleistocene.

To explore the trends in glacial and interglacial extremes in B/Ca ratios, we define glacial and interglacial maxima across the last 1.5 My from the benthic oxygen isotope record. ΔCO_3^{2-} maxima and minima were picked within a ka window around the interglacial or glacial peak, to allow for uncertainty associated with identifying the interglacial or glacial maxima for each site, the tuning approach, and to account for changes in the length and structure of interglacial periods (Tzedakis et al., 2009). G-I amplitude is taken as the absolute value of the difference between successive average glacial/interglacial values (e.g. G1-IG1, G2-IG1, G2-IG2).

3. Results

3.1 Down core B/Ca record

During the last 1.5 My, *C. wuellerstorfi* B/Ca at site 607 exhibits G/IG variability with a decrease in average B/Ca minima of 21 $\mu\text{mol/mol}$ across the Pleistocene mostly driven by changes in the glacial values primarily decreasing at MIS 23 from 180 $\mu\text{mol/mol}$ to an average 161 across MIS 20-18. Glacial B/Ca then increase to 179 $\mu\text{mol/mol}$ at MIS 14. At MIS 12, glacial B/Ca values decrease by 25 $\mu\text{mol/mol}$ and was at this level also during MIS 10 and 6 (Figure 3C, 4). In contrast, interglacial B/Ca ratios show a small long-term decline throughout the record resulting in the G/IG amplitude varying across the Pleistocene; during the pre-MPT interval (MIS 49 to 25, 1500 to 950 ka), the average G/IG amplitude in B/Ca is $\sim 24 \mu\text{mol/mol}$ (IG: 227 ± 4 ; G: 203 ± 5 ; ± 1 standard errors of the mean, SEE). The G/IG

amplitude increases to 48 (IG: 224 ± 3 ; G: 170 ± 1 ; ± 1 SEE) $\mu\text{mol/mol}$ across MIS 23 to 13 and throughout the late Pleistocene interval. Interglacial B/Ca values decrease at MIS 23 from average steady values of 227 to 206 $\mu\text{mol/mol}$ at MIS 19 and increase at MIS 17 to average IG value of 227 $\mu\text{mol/mol}$ until MIS 11 where it declines to average IG value of 200 $\mu\text{mol/mol}$ where it remains steady for the rest of the Pleistocene (Figure 4). Notably, the B/Ca record shows a remarkable correlation with the benthic foraminiferal $\delta^{13}\text{C}$ record both on G/IG and longer time scales (Ruddiman et al. 1989; Figure 3B,C).

Our piston core record encompasses the 165 to 75 and 55 to 10 ka interval. The CHN82-24-23PC B/Ca variability generally agrees with available records across these overlapping time intervals. For comparison, we look toward the published *C. wuellerstorfi* B/Ca records from Yu et al. (2008, 2016) from the deep North Atlantic across the LGM to Holocene (BOFS 8K; water depth 4045 m) and from 60 to 90 ka (MD95-2039; MD01-2446; Figure 3C). CHN82-24-23PC B/Ca tracks the rise in MIS 5 and decline into MIS 4 as shown in records from MD01-2446 and MD95-2039. The BOFS 8K record shows a similar amplitude change in B/Ca ($\Delta\text{G/IG}=43$; IG: 213; G:170) $\mu\text{mol/mol}$ to the G/IG change exhibited across the last 400 ky from site 607 (Figure S3). LGM CHN82-24-23PC B/Ca values do not reach the low values of BOFS 8K as the record is of lower resolution and likely not capturing the minimum in B/Ca.

The record of deep Atlantic B/Ca at site 607 from this study agrees well with the low resolution MPT record presented by Lear et al. (2016; Figure 3C). Therefore, we compile their data alongside our record and use this to examine the spectral properties of the record alongside examining deep ocean ΔCO_3^{2-} changes. The B/Ca record exhibits orbital frequencies similar to the $\delta^{18}\text{O}$ record with a strong 100 ky peak present in the 140 to 700 ka time interval, following the MPT (Figure S2). Prior to the MPT, from 1000-1500 ka, the record shows precession-like periodicities (18, 21 ky) but does not resolve a 41 ky cyclicity, possibly due to small gaps in the record. Also, B/Ca changes lag $\delta^{18}\text{O}_b$ by 3 ky similar to the $\delta^{13}\text{C}_b$ and %NCW record in the 100 ky world (Figure S3; Table S1) whereas % CaCO_3 lags $\delta^{18}\text{O}_b$ by 8 ky.

Moving forward we combine all B/Ca records from this study (DSDP 607; CHN82-24-23PC) and previous studies (DSDP 607; BOFS 8K; MD95-2039; MD01-2446) from the deep North

Atlantic to produce a composite Atlantic B/Ca record. (Figure 4D). Applying the calibration of Yu et al. (2007) yields a long-term change in ΔCO_3^{2-} of 20 $\mu\text{mol/kg}$ with a G/IG amplitude shift of 23 $\mu\text{mol/kg}$ prior to the MPT (1000-1500 ka), and 38 $\mu\text{mol/kg}$ post MPT (0-750 ka; Figure 5D; calibration uncertainty $\pm 9 \mu\text{mol/kg}$). Interglacial ΔCO_3^{2-} values decrease by 10 $\mu\text{mol/kg}$ from the pre- to post- MPT with average interglacial levels of the last 4 cycles similar or slightly higher than modern ΔCO_3^{2-} of 33 $\mu\text{mol/kg}$ at site 607. Glacial ΔCO_3^{2-} levels decreased by 27 $\mu\text{mol/kg}$ from pre- to post- MPT conditions. Our ΔCO_3^{2-} record shows distinct variations on G/IG timescales indicating modifications in deep North Atlantic Ocean carbon storage across the Pleistocene. The range of deepwater ΔCO_3^{2-} variability ($>30 \mu\text{mol/kg}$), recorded at Site 607 exhibits changes beyond the modern ΔCO_3^{2-} range of the NADW and AABW end-members suggesting that in addition to shifts in water mass distribution, other processes (discussed section 4.1) are driving the observed G/IG cycles in ΔCO_3^{2-} .

Changes in CaCO_3 content in deep North Atlantic sediments have been variably related to changes in biological production at the surface, dissolution/preservation on the seafloor and dilution by terrigenous inputs (Ruddiman et al. 1989). LGM CaCO_3 content in various sites in the deep Atlantic has been variably linked to lower CaCO_3 production and higher dissolution alongside a higher terrigenous input (Crowley, 1983; Francois et al. 1990; Francois & Bacon, 1991; Barker et al., 2004; Sexton & Barker, 2012). However, at site 607 the ΔCO_3^{2-} records change in parallel with % CaCO_3 suggesting that ΔCO_3^{2-} ion variations are largely associated with the dissolution and preservation cycles in the deep North Atlantic (Figure 5). Deep Atlantic ΔCO_3^{2-} values observed across the Pleistocene are consistent with Atlantic carbonate preservation records showing high preservation during interglacials and enhanced dissolution during glacials (Crowley, 1983; Ruddiman et al. 1989; Sexton & Barker 2012). Below (see section 4.2), we further explore the relationship between CaCO_3 preservation dissolution cycles and deep ocean ΔCO_3^{2-} content.

4. Discussion

4.1 Atlantic Deep ocean ΔCO_3^{2-} evolution

Benthic $\delta^{13}\text{C}$ in the deep North Atlantic varies in response to changes in the amount of organic carbon/nutrient, the influence of NADW, air-sea gas exchange and shifts in mean ocean $\delta^{13}\text{C}$ (Raymo et al., 1997; Raymo, 2004). In present-day, site 607 is bathed by NADW, a well-ventilated water mass with high $\delta^{13}\text{C}_b$ ($>1.0\text{‰}$) and in saturated conditions

($\Delta\text{CO}_3^{2-}=33\text{ }\mu\text{mol/kg}$). In comparison, southern sourced water mass, Antarctic Bottom Water (AABW), is relatively undersaturated ($\Delta\text{CO}_3^{2-}=0\text{ }\mu\text{mol/kg}$) and has lower $\delta^{13}\text{C}_b$ ($\sim 0.4\text{‰}$) due to higher nutrient content. The ΔCO_3^{2-} and $\delta^{13}\text{C}_b$ records closely covary across the last 1 My and display similar shifts during the 1000 to 800 ka interval, thereby suggesting that waxing and waning of northern and southern component waters played a key role in deep Atlantic ΔCO_3^{2-} as previously proposed for the last glacial maximum and recently for the Pleistocene (Marchitto et al. 2005; Yu et al. 2008; Lear et al. 2016; Figure 5). Pena & Goldstein (2014), using ϵ_{Nd} proxy, show a decrease in the relative proportion of NADW versus southern sourced water from 1000 to 900 ka, supporting the notion of a greater contribution of Southern Component Water (SCW) at Site 607. However, the benthic $\delta^{13}\text{C}$ and ΔCO_3^{2-} composition of source waters is also impacted by air-sea exchange driven by temperature, and CO_2 outgassing and invasion in the Southern Ocean and North Atlantic (Lynch-Stieglitz et al. 1995). Considering the impact of air-sea gas exchange across the MPT, Lear et al (2016) estimated that the increase in $\delta^{13}\text{C}_b$ partially reflects reduction of outgassing due to an increase in sea ice extent in the Southern Ocean leading to a decline in SCW $\delta^{13}\text{C}$ and ΔCO_3^{2-} alongside the increase in SCW at expense of Northern Component Water (NCW) at site 607.

The decrease in $\delta^{13}\text{C}_b$ from 1000 to 800 ka is associated with a decrease in ΔCO_3^{2-} , which suggests the incursions of SCW were associated with lower ΔCO_3^{2-} and increased nutrient content, supported by higher Cd/Ca during glacials from a reconstruction at Site 607 (Lear et al. 2016). To examine this further and determine whether this reflects changes in corrosivity of southern sourced deep waters or changes in the volume of SSW, we compare our ΔCO_3^{2-} record to a %NCW record (Raymo et al. 1997; Lang et al. 2016; Figure 5A, D). The % NCW record documents the circulation component at site 607 scaled in terms of relative changes in preformed $\delta^{13}\text{C}_b$ composition of NADW (ODP Site 982) and $\delta^{13}\text{C}_b$ composition of deep Pacific (ODP Site 849).

From 1000 to 800 ka, the glacial decrease evident in the ΔCO_3^{2-} record, with a pronounced decline mirrored in both the decrease of $\delta^{13}\text{C}_b$ and %NCW record, suggest an increase in SCW penetration to this site, as supported by ϵ_{Nd} records from Pena & Goldstein (2014). Pleistocene vertical $\delta^{13}\text{C}_b$ gradients between intermediate and deep waters in the South Atlantic show that since 1100 ka a sharp chemocline developed existed between mid-depth

and deep waters during glacial periods. This sharp chemocline separated well-ventilated intermediate waters from poorly ventilated deep waters (Hodell et al. 2003) consistent with the decrease in ΔCO_3^{2-} recorded at Site 607. Using a stacked magnetic susceptibility record Schmieder et al. (2000) show that during the MPT interim state (from 900 to 600 ka), CaCO_3 accumulation rates in the subtropical South Atlantic are notably reduced suggesting greater influence of Antarctic bottom waters. Using planktonic foraminifera fragmentation index, Groger et al. (2003) show the enhancement of G/IG preservation and dissolution cycles at that time. These lines of evidence indicate that changes in our B/Ca- ΔCO_3^{2-} record across the MPT are related to an increase of SCW at the expense of NCW.

Following the MPT, from MIS 20 to 14, both records show similar shifts in glacial values with glacial ΔCO_3^{2-} increasing until MIS 12 paralleling the increase in glacial $\delta^{13}\text{C}_b$ values. Interglacial ΔCO_3^{2-} slightly increase from MIS 21 to 17 and then plateau in tandem with the interglacial $\delta^{13}\text{C}_b$ trend (Figures 4, 5). Both features suggest a tight coupling on G/IG timescales. In contrast, the %NCW record shows only a small (~10%) increase in glacial maxima across this time interval. Thus, the inferred small decrease in the volumetric contribution of SCW likely contributed negligibly to the increase ΔCO_3^{2-} during this time interval. Instead, the close correspondence between the glacial increase in ΔCO_3^{2-} (by about 20 $\mu\text{mol/kg}$) and the increase in glacial $\delta^{13}\text{C}_b$ suggests that both were driven by whole ocean changes in carbon. From MIS 15 onwards, Lear et al. (2016) propose that the interglacial increase in $\delta^{13}\text{C}_b$ is related to reduced interglacial CO_2 evasion into North Atlantic source waters, impacting both the $\delta^{13}\text{C}_b$ and ΔCO_3^{2-} records. Following the MPT, ΔCO_3^{2-} remains at relatively similar levels suggesting air-sea gas exchange minimally impacted the interglacial ΔCO_3^{2-} North Atlantic source waters (Figure 4).

Stacked planktic and benthic $\delta^{13}\text{C}$ foraminifera records, show a decrease by 0.35‰ around 900 ka, gradual return to pre-1 Ma values at 500 ka and a final rapid decrease at 420 ka (Wang et al. 2004; Hoogakker et al. 2006; Lisiecki, 2010). The global long-term increase of 0.6 per mil lasting until 500 ka and the subsequent decrease at MIS 12 likely reflect changes in the carbon burial to the deep ocean modified by changes in the rain ratio, bottom water ventilation, and surface ocean production (Wang et al. 2004; Hoogakker et al. 2006; Lisiecki, 2010). The similarity between the ΔCO_3^{2-} and $\delta^{13}\text{C}_b$ records both in the long-term trends and G/IG variability throughout this time window suggests it is recording changes in oceanic carbon-carbonate system and changes in deep ocean circulation.

4.2 Deep ocean CO_3^{2-} and CaCO_3 preservation/dissolution cycles

A recent study, using orbitally resolved IndoPacific B/Ca records show that over the past 500 ky, deep Pacific ΔCO_3^{2-} G/IG variability parallels changes in sediment % CaCO_3 content, apparently in contrast with the expectation of anticorrelation between these properties derived from the classic compensation theory (Kerr et al., 2017). To explain the apparent contradiction between the expectation from the compensation theory and their observed records, Kerr et al., (2017) argue that a global increase in CO_3^{2-} during glacial intervals is caused by the delivery of alkalinity from the shelves due to the reduction of carbonate deposition on the shelves during low sea level stands (a.k.a ‘coral reef hypothesis’) across the last 450 ky (Berger, 1982; Opdyke & Walker, 1992). Here, we propose an alternative mechanism, whereby the post-MPT Pacific and Atlantic CO_3^{2-} records are driven by basin-to-basin alkalinity transfer, rather than the classic vertical CaCO_3 compensation.

Over the past 400 ky, the Atlantic ΔCO_3^{2-} record shows G/IG swings of 30-40 $\mu\text{mol/kg}$ in the 100-ky band which decline into the glacial and recover across the terminations (Fig. 5D; 6D). In contrast, the IndoPacific records show smaller ($<10 \mu\text{mol/kg}$) changes in ΔCO_3^{2-} as reported by Kerr et al. (2017) (Figure 6C), which rise into the glacial and shoot down across the terminations. This behavior is also evident in the % CaCO_3 records from the Atlantic, IndoPacific, and Southern Ocean, in support of our deep ocean ΔCO_3^{2-} records (Figure 5C;6A,B). The nature of the deep ocean ΔCO_3^{2-} records contrasts the expected changes as estimated from box models and predicted by CaCO_3 compensation theory (Boyle, 1988; Broecker & Peng, 1989; Toggweiler, 1999; Toggweiler, 2008).

Using a seven-box model Toggweiler (2008), estimates that in response to deep ocean ventilation the response to expected change in whole ocean ΔCO_3^{2-} below 1000 m, is on the order of $\sim 9 \mu\text{mol/kg}$, significantly lower than ΔCO_3^{2-} swings observed at DSDP site 607 across the last 400 ky (Figure 7). Also, the structure of the observed ΔCO_3^{2-} changes lacks the expected spike up on the terminations and down on glacial initiations followed by rapid return to the steady state baseline as expected from the classic compensation theory.

Additionally, the large swings in Atlantic ΔCO_3^{2-} are not compensated quickly as expected and the deep Atlantic CO_3^{2-} remains above or below its long-term mean for periods substantially greater than the assumed response time of $\sim 5\text{-}10$ kyrs (Figure 5D;6D; Broecker & Peng, 1987; Keir, 1988; Archer et al. 1997; Kohler & Fischer, 2006). It is noteworthy that the ΔCO_3^{2-} reconstructions show that both the deep Atlantic and Pacific were undersaturated

with respect to calcite during glacial intervals suggesting that dissolution was not the only control on CaCO_3 preservation in the deep glacial ocean. The difference in the structure of G/IG change, magnitude, and relative timing of G/IG cycles between the Atlantic and Pacific deep ΔCO_3^{2-} records questions the classic view of CaCO_3 compensatory mechanism and calls for a different approach (Toggweiler, 1999; 2008).

We propose that over the past ~ 1 My, CaCO_3 compensation occurred in response to changes in the places where CaCO_3 is buried rather than vertical shifts in the lysocline. This explains the Atlantic-Pacific difference and the contrast to the box model predictions. We hypothesize that deep ocean ΔCO_3^{2-} and $\%\text{CaCO}_3$ were driven by changes in corrosivity of the deep North Atlantic and interbasin migration of calcite burial or 'hot spots' across G/IG cycles (Dunne et al. 2012). A "hot spot" is an area of the ocean where the enhanced production of CaCO_3 at the surface overlies a favorable burial environment on the sea floor. The net result is a local burial flux of CaCO_3 that is higher than average. The concept is important because the cumulative burial in a few large hot spots could, in principle, balance the delivery of Ca^{2+} and HCO_3^- ions to the ocean from rivers. At present, the main hotspot or locus of CaCO_3 burial is in the deep Atlantic and much of the global burial of CaCO_3 occurs there due to the formation of the youngest, least corrosive deep waters in this region (see Fig. 1c in Dunne et al. 2012). In contrast, the Pacific Ocean has a less propensity for CaCO_3 burial due to the oldest, undersaturated deep waters residing there. Also, in the present day the production and flux of CaCO_3 to the sea floor is not uniform across the ocean and burial of CaCO_3 is more of a kinetic process than a purely equilibrium process (Li et al. 1969; Broecker & Peng, 1982; Dunne et al. 2012). In this sense, favorable burial environments can also include areas where the water above the sea floor is mildly undersaturated with respect to calcite and enhanced fluxes of CaCO_3 to the sea floor overwhelm the dissolution in the sediments (due to the slow kinetics of the dissolution reactions). This non-uniformity, we claim, creates the potential for "hot spots" of CaCO_3 burial and preservation, which result when enhanced fluxes to the seafloor coincide with a relatively shallow seafloor and/or the presence of NADW. In this instance, all the CaCO_3 production in areas outside the hot spots tends to dissolve.

Here we build upon these observations and propose that incursion of dense corrosive waters from the southern hemisphere into the glacial deep Atlantic decreased Atlantic calcite burial thus raising the ocean's average CO_3^{2-} . We propose this is compensated by a shift in the locus of CaCO_3 burial into the equatorial Pacific to compensate for the declining burial in the other

areas. This is exhibited during glacials, when ΔCO_3^{2-} and % CaCO_3 records show an increase in glacial deep Atlantic corrosivity and undersaturation that inhibits burial across the last 800 ky (e.g. MIS 12) (Fig. 6B, D). Also, at the same time, empirically, it seems that the locus of burial shifts to the tropical Pacific evident from sedimentary carbonate records which shows that the area of CaCO_3 -rich sediments expands in the Pacific during the LGM and shrinks everywhere else (Farrell & Prell, 1989; Catubig et al. 1998). This suggests that the Pacific became a hot spot for CaCO_3 burial despite the undersaturated state during glacials. We propose this is likely caused by the following effects i) increase in CaCO_3 export that overcame the apparent corrosivity of the water and ii) the depth at which major CaCO_3 dissolution starts occurs is not $\Delta\text{CO}_3^{2-}=0$ but in fact lower. This is consistent with Subhas et al. (2017) suggesting that CaCO_3 dissolution starts only at a saturation state of 0.7 not 1. Therefore, it seems that the small increase in deep Pacific ΔCO_3^{2-} was sufficient to shift the balance from carbonates dissolution to preservation. In this way, the ΔCO_3^{2-} in the deep IndoPacific rises in response to the same mechanism that keeps the ΔCO_3^{2-} in the deep Atlantic low and the atmospheric CO_2 low.

When the locus of burial shifts to the Pacific, the ocean's average CO_3^{2-} and alkalinity of the ocean must increase to compensate, or rise to a point where the burial in the new hot spot is sufficiently high to overcome the general corrosivity of the deep Pacific. These excursions lead to a larger compensation response that is spread out over longer periods of time and would make the amount of alkalinity and dissolved inorganic carbon in the ocean swing widely over time. This is reflected in the slow rise of ΔCO_3^{2-} through stages 4, 3, and 2 in TT013-PC72 (Kerr et al. 2017) and in the late glacial increases in % CaCO_3 at site 1089, a stand in for burial in the Indo Pacific (Figure 6A). The gradual increase in alkalinity over this time helps keep the pCO_2 of the atmosphere low for tens of thousands of years and the ocean cold, which reinforces the stratification of the deep Atlantic and times when the ΔCO_3^{2-} falls over time at site 607 as seen particularly well during MIS 12 and 10. So, instead of the CO_3^{2-} simply ticking up and down within a G/IG transition, as in the classical response, a shift of the locus of CaCO_3 burial tends to hold the deep CO_3^{2-} up or down over entire glacials or interglacials.

Conversely, when the locus of burial shifts back to the Atlantic during interglacials, the ocean's average CO_3^{2-} and alkalinity of the ocean falls because it is suddenly easy to bury

CaCO₃ again in the less corrosive Atlantic. This seems to be seen especially well during Terminations II and I at VM 28-122 in Yu et al. (2013) and in the %CaCO₃ decreases at site 1089 (Figure 6A). The net loss of alkalinity from the ocean helps keep the atmospheric pCO₂ high and consequently the ocean warm during interglacials. Also, CO₃²⁻ in the deep Atlantic is “super-saturated” during IG intervals and as such needs to decrease a lot to reach undersaturation and start dissolution as expressed by the large swings in ΔCO₃²⁻ (30-40 μmol/kg).

The response of %CaCO₃ and deep ocean CO₃²⁻ content to variations in the Atlantic meridional overturning circulation (AMOC) strength has been previously explored in several modeling studies (Chikamoto et al. 2008; Yu et al. 2016). Chikamoto et al. (2008) show that during an ‘AMOC shutdown’, initiated by a freshwater perturbation, the average carbonate ion concentration in the North Atlantic is reduced by ~37 μmol/kg and the CaCO₃ burial flux is reduced in the North Atlantic and increased in the Pacific. In addition, Yu et al. (2016) investigate the influence of AMOC changes on deep Atlantic CO₃²⁻ and show that a cessation of NADW leads to a decrease in deep (>3 km) North Atlantic and Equatorial Atlantic CO₃²⁻ by 20-40 μmol/kg. Although these model results provide some justification for our argument at least, the Chikamota et al (2008) model results have a negligible effect on atmospheric pCO₂ in contrast to our conjecture that G-I compensation changes played a role in atmospheric pCO₂. We suggest here that their model does not fully incorporate regional production differences in CaCO₃, which when in combination with favorable burial environments produces hotspots, does not replicate fully the scenario where the burial of CaCO₃ shifts from one ocean to another and concomitant G-I changes in pCO₂.

In previous studies it has been argued that sea-level variations drive the cycles of CaCO₃ burial and dissolution in the ocean and can explain the low amplitude variations in the Pacific and glacial increase in ΔCO₃²⁻ and covariation in %CaCO₃ and ΔCO₃²⁻ (Yu et al., 2013; Kerr et al. 2017). Kerr et al. (2017) use the broad correlation of the amplitude of glacial increase in CO₃²⁻ with the amplitude of sea-level decline as support. In doing so, the authors suggest that shifts in Atlantic ΔCO₃²⁻ are less sensitive to whole ocean carbonate system changes, as past water mass distribution are strong in this region and impacting deep carbon storage. Furthermore, the ΔCO₃ records presented here resemble the δ¹³C_b (proxy of carbon storage and circulation) rather than the δ¹⁸O_b record (proxy of ice volume and sea level). This is very

pronounced between MIS 20 and 14 and then 12 and 6 when glacial ΔCO_3 follow closely the $\delta^{13}\text{C}_b$ (Figure 5). These lines of evidence suggest that changes in carbon storage in the deep Atlantic are tightly coupled to the very thing that is driving the variations in atmospheric CO_2 , namely changes in the ventilation of southern-sourced bottom water. Building on this idea, we argue that hypothesized enhanced preservation, due to alkalinity supplied from the shelf areas during sea level fall, could affect the Atlantic but not enough to overcome the dissolution changes caused by shifts in ventilation/carbon storage. Therefore, we argue that CaCO_3 dissolution in the Atlantic was the dominant source of alkalinity to the deep Pacific and hence played the key role in controlling atmospheric CO_2 across mid-to-late Pleistocene. Indeed, CaCO_3 compensation, as viewed from changes in deep water ΔCO_3^{2-} occurred throughout the past 1.5 My and apparently scaled with the changes in atmospheric pCO_2 (Figure 5).

Across the MPT around 1.1 Ma, the classic ('anti-correlated') CaCO_3 patterns (i.e. dissolution in Atlantic and preservation in Pacific during glacials) developed. Sexton & Barker (2012) suggest this was due to the strengthening of deepwater ventilation in the abyssal Pacific during glacials and weakening during IGs, which led to glacial Pacific CaCO_3 dissolution to diminish while driving IG Pacific CaCO_3 dissolution to intensify. The proposed idea by these authors would, however, let the respired CO_2 in the deep Pacific escape to the atmosphere during glacials. Here we propose an alternative scenario whereby the development of extremely poor ventilation and increased stratification in the deep Atlantic provides a better explanation for the Pleistocene CaCO_3 observations and atmospheric CO_2 . Deep ocean cooling starts at Site 607 in the North Atlantic at ~1.2 Ma (Sosdian and Rosenthal 2009, Ford et al., 2016), while based on the record from ODP Site 1123, the deep glacial Southern Ocean was close to freezing throughout this period (Elderfield et al., 2012). While it is possible that some of the oxygen isotopic change of seawater change that Elderfield et al (2012) document at the MPT reflects increase salinity, either way the isotope data reflect an increase in the density of the Southern Ocean deep waters with increased stratification and hence weaker ventilation as seen in the last glacial period in the South Atlantic (Adkins et al. 2001). The cooling of deep water at Site 607 (Sosdian and Rosenthal 2009, Ford et al., 2016) likely reflects increased stratification of the deep Atlantic. A colder ocean led to northward migration of the Antarctic polar fronts and likely resulted in thicker and more expanded southern sea ice (Kemp et al., 2010) that made the deep ocean more saline more stratified and less ventilated over time. We propose the

initial cause of the anti-phased behavior between the Atlantic and Pacific relates to the deep ocean cooling across the MPT that led to changes in ventilation and drove the deep Atlantic into under-saturation consistent with the appearance of anti-phased behavior in %CaCO₃, as pinpointed by Sexton & Barker (2012) at ~1.1 Ma.

Following the cooling at 1.2 Ma, both the $\delta^{13}\text{C}_b$ and %NCW decline from 1 to 0.8 Ma highlighting changes in ventilation. The coherency between the $\delta^{13}\text{C}_b$ and Atlantic ΔCO_3^{2-} record across the MPT indicates changes in deep ocean ventilation are driving deep ocean CO_3^{2-} changes (Figure 3; 5). Thus, across the MPT, we propose that the increase in deep ocean corrosivity led to the development of the classic CaCO₃ pattern whereby the locus of CaCO₃ burial shifts to the Pacific during glacials and switches back the Atlantic during interglacials.

4.3 The Mid-Bruhnes Event

Glacial ΔCO_3^{2-} levels drop by ~ 30 $\mu\text{mol/kg}$ after MIS 14, becoming for the first time under-saturated through the last 1.5 My in the deep North Atlantic at MIS 12, when it reaches -20 ± 9 $\mu\text{mol/kg}$ (Figure 5; 6). This period of corrosive deep waters corresponds with the mid-Brunhes dissolution event, beginning at ~600 ka and lasting for 400 ky, peaking around ~400 ka (MIS 11), exhibited by records showing enhanced dissolution in the deep ocean occurring in the North Atlantic (Crowley, 1985; Jansen et al. 1986; Groger et al. 2003; Barker et al. 2006), in the Indian Ocean (Bassinot et al. 1994), and in the Pacific (Farrell & Prell, 1989).

The decrease in the saturation in the deep North Atlantic occurs alongside the development of the largest ice sheets of the Pleistocene, cooling of the deep Southern Ocean and North Atlantic (as recorded in ODP Site 1123 and DSDP Site 607; Elderfield et al., 2012; Sosdian and Rosenthal, 2009) and intensification of the 100 ky cycles in sea level and pCO₂ (Lisiecki & Raymo, 2005; Hönlisch et al. 2009; Bereiter et al. 2015). Cooling of the deep Atlantic, which culminated at MIS 12, likely resulted in greater stratification and sluggish ventilation, which enhanced carbon sequestration in the deep Atlantic possible augmented by increased surface ocean productivity (Lawrence et al. 2013).

The % NCW record shows a prolonged period of near 0% levels, the longest duration of reduced NCWs across the last 800 ky (Figure 5). Records of ocean circulation show

exceptionally prolonged (5 ky) collapse of the Atlantic Meridional Overturning Circulation (Riveiros et al. 2013) and increased influence of SSW (Hall & Becker, 2007) at MIS 12 suggesting a longer residence time of seawater in the glacial deep Atlantic. The isolation of these deep waters allowed greater accumulation of respired organic carbon as surface ocean productivity was higher at Site 607/U1313 as indicated by alkenone mass accumulate rates (C_{37} total MAR) and total organic carbon records (Stein et al. 2009; Lawrence et al. 2013). Both records indicate higher surface ocean productivity possibly tied to the more southerly position of the polar front and/or increased dust supply (Stein et al. 2009; Lawrence et al. 2013).

Modeling studies show that a 50% NADW reduction leads to a decrease of $\sim 30 \mu\text{mol/kg}$ in the deep Atlantic ($>3 \text{ km}$) and NADW cessation leads to a greater decrease of $20\text{-}40 \mu\text{mol/kg}$ (Yu et al. 2016). We propose here that additional decrease in ΔCO_3^{2-} minima starting at MIS 12, set the stage for the high pCO_2 levels at the mid-Bruhnes event via an increase in whole ocean alkalinity followed by enhanced CaCO_3 preservation at MIS 11, 9 and MIS 5. Specifically, we propose that whole ocean alkalinity became higher than "normal" during MIS 12, in response to cooling and reduced ventilation of the deep Atlantic, which caused increase corrosivity. Then, when the deep ocean became better ventilated during termination V, more CaCO_3 was preserved everywhere for a short period of time. The enhanced removal of alkalinity then pushed the atmospheric pCO_2 higher. The larger response time than previously assumed contributes to this effect (i.e. the excess alkalinity is not completely removed during the deglaciation but throughout the interglacial maintaining the high pCO_2 levels). This suggests that IG pCO_2 levels are partly determined by the preceding G/IG changes and the response time of the compensation. A noteworthy corollary of the proposed mechanism, is that the lower limit of the glacial pCO_2 , which has been $\sim 180\text{ppmV}$ during the past 800 ky, is not strictly set by the ocean alkalinity and consequently the "extra" alkalinity generated at the corrosive glacials (MIS 11, 9, 5) was available for removal during the subsequent interglacials (e.g., MIS 11, 9, 5) leading to higher pCO_2 levels.

5. Conclusions

To understand the nature of the ocean's buffering capacity and its role in modulating pCO_2 , reconstructions of the deep ocean carbonate system parameters are necessary.

Pleistocene benthic foraminiferal B/Ca record from DSDP site 607 suggests 30-40 $\mu\text{mol/kg}$ G/IG variability in deep ocean ΔCO_3^{2-} , substantially larger than seen in the IndoPacific records. The observed timing, magnitude and structure of swings in G/IG deep Atlantic ocean ΔCO_3^{2-} are also larger than expected based on thermodynamic CaCO_3 compensation mechanism. To explain this difference, we suggest that during glacials, when ventilation weakens, the more corrosive and lower ΔCO_3^{2-} in the deep Atlantic, shifts the locus of CaCO_3 burial to the equatorial Pacific to maintain the long-term CaCO_3 balance via a new carbonate burial ‘hot spot’, where the carbonate flux to the seafloor and the small increase in ΔCO_3^{2-} are sufficient to overcome the under-saturated conditions characteristic of the deep Pacific during interglacial intervals. We argue that, the small increase in IndoPacific ΔCO_3^{2-} is driven by the same mechanism that keeps the deep Atlantic ΔCO_3^{2-} and atmospheric CO_2 low and is not due to the delivery of alkalinity from the shelves during low sea level stands.

Shifts in the global carbon cycle and deep ocean circulation play a role in driving the long-term trend in Atlantic ΔCO_3^{2-} across the mid-to-late Pleistocene. The ΔCO_3^{2-} record from DSDP site 607 shows shifts in G/IG variability associated with the MPT and mid-Brunhes event. The basin-basin fractionation likely initiated across the MPT as the deep Atlantic Ocean began to play a large role in enhancing pCO_2 due to changes in ventilation, likely due to cooling and enhanced stratification, which drove deep Atlantic into under-saturation during glacial intervals. Deep ocean ΔCO_3^{2-} levels reached the lowest levels of the last 1.5 My at Marine Isotope Stage (MIS) 12 and became persistently under-saturated. We propose the low ΔCO_3^{2-} of MIS 12 were possibly initiated by a combination of sluggish circulation and high surface ocean productivity allowing for increased storage of CO_2 in the deep Atlantic. The low ΔCO_3^{2-} of the deep Atlantic set the stage for the high pCO_2 levels at MIS 11 via an increase in whole ocean alkalinity followed by enhanced CaCO_3 preservation at MIS 11 and the subsequent interglacials of MIS 9 and 5.

Acknowledgements

Thanks to useful discussions with Phil Sexton, help with time series analysis from Henrieka Detlef and Kira Lawrence, and ICPMS technical support from Paul Field. We also thank anonymous reviewers for comments that greatly improved the manuscript. Sindia Sosdian acknowledges the financial support provided by the Welsh Government and Higher Education Funding Council for Wales through the Sêr Cymru National Research Network for Low Carbon, Energy and Environment. Yair Rosenthal acknowledges NSF support (OCE 1334691). All data presented in this study are given in the Supplementary Tables.

References

- Adkins, J. F., and D. P. Schrag (2001), Pore fluid constraints on deep ocean temperature and salinity during the last glacial maximum, *Geophysical Research Letters*, 28(5), 771-774.
- Allen, K. A., E. L. Sikes, B. Hönisch, A. C. Elmore, T. P. Guilderson, Y. Rosenthal, and R. F. Anderson (2015), Southwest Pacific deep water carbonate chemistry linked to high southern latitude climate and atmospheric CO₂ during the Last Glacial Termination, *Quaternary Science Reviews*, 122, 180-191.
- Archer, D., and E. Maierreimer (1994), Effect of deep-sea sedimentary calcite preservation on atmospheric CO₂ concentration, *Nature*, 367(6460), 260-263.
- Archer, D., H. Kheshgi, and E. MaierReimer (1997), Multiple timescales for neutralization of fossil fuel CO₂, *Geophysical Research Letters*, 24(4), 405-408.
- Arrhenius, G. (1952), Sediment cores from the East Pacific. Reports of the Swedish Deep Sea Expeditions, 1947-1948, 1-227.
- Babila, T. L., Y. Rosenthal, and M. H. Conte (2014), Evaluation of the biogeochemical controls on B/Ca of Globigerinoides ruber white from the Oceanic Flux Program, Bermuda, *Earth and Planetary Science Letters*, 404, 67-76
- Barker, S., Kiefer, T., Elderfield, H., 2004. Temporal changes in North Atlantic circulation constrained by planktonic foraminiferal shell weights. *Paleoceanography* 19. PA3008, doi:10.1029/2004PA001004.
- Barker, S., D. Archer, L. Booth, H. Elderfield, J. Henderiks, and R. E. M. Rickaby (2006), Globally increased pelagic carbonate production during the Mid-Brunhes dissolution interval and the CO₂ paradox of the MIS 11, *Quaternary Science Reviews*, 25(23-24), 3278-3293.
- Bassinot, F. C., L. Beaufort, E. Vincent, L. D. Labeyrie, F. Rostek, P. J. Muller, X. Quidelleur, and Y. Lancelot (1994), Coarse fraction fluctuations in pelagic carbonate sediments from the tropical Indian Ocean- a 1500-kyr record of carbonate dissolution, *Paleoceanography*, 9(4), 579-600.
- Bereiter, B. *et al.* Revision of the EPICA Dome C CO₂ record from 800 to 600 kyr before present. *Geophysical Research Letters* 42, 542-549, doi:10.1002/2014gl061957 (2015).

Berger, W. H., (1982) Increase of carbon dioxide in the atmosphere during deglaciation: the coral reef hypothesis. *Naturwissenschaften*, 69, 87-88. <http://dx.doi.org/10.1007/BF00441228>

Berger, W. H., and R. S. Keir (1984), Glacial - Holocene changes in atmospheric CO₂ and the deep - sea record, in *Climate Processes and Climate Sensitivity*, Geophys. Monogr. Ser., vol. 29, edited by J. E. Hansen and T. Takahashi, pp. 337-351, AGU, Washington, D. C.

Berger, W. H. and E.L. Winterer (1974) Plate stratigraphy and the fluctuating carbonate line, *Pelagic Sediments on Land and Under the Sea*, edited by K.J. Hsu and H.C. Jeakyns, Spec. Publ. Int. Assoc. Sedimentol., 1, 11-48, 1974.

Past Interglacials Working Group of PAGES (2016), Interglacials of the last 800,000 years, *Reviews of Geophysics*, 54(1), 162-219.

Boyle, E. A. (1983) Chemical accumulation variations under the Peru current during the past 130,000 years, *J. Geophys. Res.*, 88, 7667-7680, 1983.

Boyle, E. A. (1988), Vertical oceanic nutrient fractionation and glacial-interglacial CO₂ cycles, *Nature*, 331(6151), 55-56.

Boyle, E.A., & Keigwin, L. (1985/85) Comparison of Atlantic and Pacific paleochemical records for the last 215,000 years — changes in deep ocean circulation and chemical Inventories, *Earth Planet. Sci. Lett.*, 76 (1985/86), pp. 135-150

Broecker, W. S., and T. H. Peng (1987) The role of CaCO₃ compensation in the glacial to interglacial atmospheric CO₂ change. *Global Biogeochem. Cycles* 1, 15-29.

Broecker, W. S. and T.-H.-Peng, *Tracers in the Sea*, Lamont-Doherty Geol. Obs., Palisades, N.Y., 1982.

Brown, R. E., L. D. Anderson, E. Thomas, and J. C. Zachos (2011), A core-top calibration of B/Ca in the benthic foraminifers *Nuttallides umbonifera* and *Oridorsalis umbonatus*: A proxy for Cenozoic bottom water carbonate saturation, *Earth and Planetary Science Letters*, 310(3-4), 360-368.

Catubig, N. R., D. E. Archer, R. Francois, P. deMenocal, W. Howard, and E.-F. Yu (1998), Global deep-sea burial rate of calcium carbonate during the last glacial maximum, *Paleoceanography*, 13, 298-310.

- Chalk, T.B, Hain, M.P., Foster, G.L., Rohling, E.J., Sexton, P.F., Badger, M.P.S., Cherry, S.G., Hasenfratz, A.P., Haug, G.H., Jaccard, S.L., Martínez-García, A., Pälike, H., Pancost, R.D., Wilson, P.A. (2017) Causes of ice age intensification across the Mid-Pleistocene Transition PNAS Proceedings of National Academy Science, doi:10.1073/pnas.1702143114

Chikamota, M.O., Matsumoto, K. Ridgwell, A. (2008) Response of deep-sea CaCO₃ sedimentation to Atlantic meridional overturning circulation shutdown, *Journal of Geophysical Research*, doi:10.1029/2007JG000669

Clark, P. U., D. Archer, D. Pollard, J. D. Blum, J. A. Rial, V. Brovkin, A. C. Mix, N. G. Pisias, and M. Roy (2006), The middle Pleistocene transition: characteristics, mechanisms, and implications for long-term changes in atmospheric pCO₂, *Quaternary Science Reviews*, 25(23-24), 3150-3184.

Crowley, T. J. (1983a), Calcium-carbonate preservation patterns in the central North Atlantic during the last 150,000 years, *Marine Geology*, 51(1-2), 1-14.

Crowley, T.J., 1985. Late Quaternary carbonate changes in the North Atlantic and Atlantic/Pacific comparisons. In: Sundquist, E.T., Broecker, W.S. (Eds.), *The Carbon Cycle and Atmospheric CO₂: Natural Variations. Archean to Present*, AGU, Washington, D. C., pp. 271–284.

deMenocal, P., D. Archer, and P. Leth (1997) Pleistocene variations in deep Atlantic circulation and calcite burial between 1.2-0.6 Ma: a combined data-model approach, *Proceedings of the Ocean Drilling Program, Scientific Results*, Vol. 154, 285-297

Doss, W., and T. M. Marchitto (2013), Glacial deep ocean sequestration of CO₂ driven by the eastern equatorial Pacific biologic pump, *Earth and Planetary Science Letters*, 377, 43-54.

Dunne, J. P., B. Hales, and J. R. Toggweiler (2012), Global calcite cycling constrained by sediment preservation controls, *Global Biogeochemical Cycles*, 26.

Elderfield, H., P. Ferretti, M. Greaves, S. Crowhurst, I. N. McCave, D. Hodell, and A. M. Piotrowski (2012), Evolution of Ocean Temperature and Ice Volume Through the Mid-Pleistocene Climate Transition, *Science*, 337(6095), 704-709.

Elmore, A. C., E. L. McClymont, H. Elderfield, S. Kender, M. R. Cook, M. J. Leng, M. Greaves, and S. Misra (2015), Antarctic Intermediate Water properties since 400 ka recorded in infaunal (*Uvigerina peregrina*) and epifaunal (*Planulina wuellerstorfi*) benthic foraminifera, *Earth and Planetary Science Letters*, 428, 193-203.

Farrell, J. W., and W. L. Prell (1989), Climatic change and CaCO₃ preservation: an 800,000 year bathymetric reconstruction from central equatorial Pacific Ocean, *Paleoceanography*, 4(4), 447-466.

Ford, H., S.M. Sosdian, Y. Rosenthal, M.E. Raymo (2016) Gradual and abrupt changes during the Mid-Pleistocene Transition, *Quaternary Science Reviews*, 149, 222-233.

Francois, R., and M. P. Bacon (1991), Variations in terrigenous input into the deep equatorial Atlantic during the past 24,000 years, *Science*, 251(5000), 1473-1476.

Francois, R., M. P. Bacon, and D. O. Suman (1990), Thorium-230 profiling in deep sea sediments: high resolution records of flux and dissolution of carbonate in the equatorial Atlantic during the last 24,000 years, *Paleoceanography*, 5(5), 761-787.

Groger, M., R. Henrich, and T. Bickert (2003), Glacial-interglacial variability in lower North Atlantic deep water: inference from silt grain-size analysis and carbonate preservation in the western equatorial Atlantic, *Marine Geology*, 201(4), 321-332.

Hain, M. P., D. M. Sigman, and G. H. Haug (2010), Carbon dioxide effects of Antarctic stratification, North Atlantic Intermediate Water formation, and subantarctic nutrient drawdown during the last ice age: Diagnosis and synthesis in a geochemical box model, *Global Biogeochemical Cycles*, 24.

Hall, I. R., and J. Becker (2007), Deep Western Boundary Current variability in the subtropical northwest Atlantic Ocean during marine isotope stages 12-10, *Geochemistry Geophysics Geosystems*, 8)

1. Higgins JA, et al. (2015) Atmospheric composition 1 million years ago from blue ice in the Allan Hills, Antarctica. *Proc Natl Acad Sci USA* **112**:6887–6891

Hodell, D. A., K. A. Venz, C. D. Charles, and U. S. Ninnemann (2003), Pleistocene vertical carbon isotope and carbonate gradients in the South Atlantic sector of the Southern Ocean, *Geochemistry Geophysics Geosystems*, 4, 39.

Hodell, D. A., C. D. Charles, and F. J. Sierro (2001), Late Pleistocene evolution of the ocean's carbonate system, *Earth and Planetary Science Letters*, 192(2), 109-124.

Hönisch, B., N. G. Hemming, D. Archer, M. Siddall, and J. F. McManus (2009), Atmospheric Carbon Dioxide Concentration Across the Mid-Pleistocene Transition, *Science*, 324(5934), 1551-1554.

Hoogakker, B. A. A., E. J. Rohling, M. R. Palmer, T. Tyrrell, and R. G. Rothwell (2006), Underlying causes for long-term global ocean delta C-13 fluctuations over the last 1.20 Myr, *Earth and Planetary Science Letters*, 248(1-2), 15-29.

Howard, W. R., and W. L. Prell (1994), Late Quaternary CaCO_3 production and preservation in Southern Ocean- Implications of for oceanic and atmospheric carbon cycling, *Paleoceanography*, 9(3), 453-482.

Howe, J. N. W., A. M. Piotrowski, T. L. Noble, S. Mulitza, C. M. Chiessi, and G. Bayon (2016), North Atlantic Deep Water Production during the Last Glacial Maximum, *Nature Communications*, 7.

Howell, P., et al. (2006), Arand time series analysis software, Brown Univ., Providence, R. I.

Jansen, J. H. F., A. Kuijpers, and S. R. Troelstra (1986), A mid-Brunhes climatic event-long term changes in global atmosphere and ocean circulation, *Science*, 232(4750), 619-622.

Keir, R. S. (1988), On the late Pleistocene ocean geochemistry and circulation, *Paleoceanography*, 3(4), 413-445.

Kemp, A.E.S., Grigorov, I., Pearce, R.B., Naveira Garabato, A.C., 2010. Migration of the Antarctic Polar Front through the mid-Pleistocene transition: evidence and climatic implications. *Quaternary Science Reviews* 29, 1993-2009.

Kender, S., E. L. McClymont, A. C. Elmore, D. Emanuele, M. J. Leng, and H. Elderfield (2016), Mid Pleistocene foraminiferal mass extinction coupled with phytoplankton evolution, *Nature Communications*, 7.

Kerr, J., R. Rickaby, J. Yu, H. Elderfield, A. Y. Sadekov (2017), The effect of ocean alkalinity and carbon transfer on deep-sea carbonate ion concentration during the past five glacial cycles, *Earth and Planetary Science Letters*, 471, 42-53.

Key, R.M., Kozyr, A., Sabine, C.L., Lee, K., Wanninkhof, R., Bullister, J.L., Feely, R.A., Millero, F.J., Mordy, C., Peng, T.H., 2004. A global ocean carbon climatology: Results from Global Data Analysis Project (GLODAP). *Global Biogeochemical Cycles* 18.

Koehler, P., and R. Bintanja (2008), The carbon cycle during the Mid Pleistocene Transition: the Southern Ocean Decoupling Hypothesis, *Climate of the Past*, 4(4), 311-332.

Lang, D. C., I. Bailey, P. A. Wilson, T. B. Chalk, G. L. Foster, and M. Gutjahr (2016), Incursions of southern-sourced water into the deep North Atlantic during late Pliocene glacial intensification, *Nature Geoscience*, 9(5), 375-379.

Lawrence, K. T., D. M. Sigman, T. D. Herbert, C. A. Riihimaki, C. T. Bolton, A. Martinez-Garcia, A. Rosell-Mele, and G. H. Haug (2013), Time-transgressive North Atlantic productivity changes upon Northern Hemisphere glaciation, *Paleoceanography*, 28(4), 740-751.

Lear, C.H., K. Billups, R.E.M. Rickaby, L. Diester-Haass, E. Mawbey, and S.M. Sosdian (2016), Breathing more deeply: Deep ocean carbon storage during the mid-Pleistocene climate transition. *Geology* (10.1130/G38636.1)

Lemarchand, D., J. Gaillardet, E. Lewin, and C. J. Allegre (2002), Boron isotope systematics in large rivers: implications for the marine boron budget and paleo-pH reconstruction over the Cenozoic, *Chemical Geology*, 190(1-4), 123-140.

Li, Y.-H., T. Takahashi, and W. S. Broecker, Degree of saturation of CaCO_3 in the oceans, J. *Geophys. Res.*, 74, 5507-5525, 1969

Lisiecki, L. E. (2010), A benthic delta C-13-based proxy for atmospheric pCO_2 over the last 1.5 Myr, *Geophysical Research Letters*, 37.

Lisiecki, L. E., and M. E. Raymo (2005), A Pliocene-Pleistocene stack of 57 globally distributed benthic delta O-18 records, *Paleoceanography*, 20(1).

Luethi, D., et al. (2008), High-resolution carbon dioxide concentration record 650,000-800,000 years before present, *Nature*, 453(7193), 379-382.

Lynch-Stieglitz, J., Stocker, T.F., Broecker, W.S., and Fairbanks, R.G., 1995, The influence of air-sea exchange on the isotopic composition of oceanic carbon: Observations and modelling, *global Biogeochemical Cycles*, v. 9, p. 653-665.

Marchitto, T. M., J. Lynch-Stieglitz, and S. R. Hemming (2005), Deep pacific CaCO_3 compensation and glacial-interglacial atmospheric CO_2 , *Earth and Planetary Science Letters*, 231(3-4), 317-336.

Opdyke, B. N., and J. C. G. Walker (1992), Return of the coral-reef hypothesis- basin to shelf partitioning of CaCO_3 and its effect on atmospheric CO_2 , *Geology*, 20(8), 733-736.

Pena, L. D., and S. L. Goldstein (2014), Thermohaline circulation crisis and impacts during the mid-Pleistocene transition, *Science*, 345(6194), 318-322.

Petit, J. R., et al. (1999), Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica, *Nature*, 399(6735), 429-436.

Rae, J. W. B., G. L. Foster, D. N. Schmidt, and T. Elliott (2011), Boron isotopes and B/Ca in benthic foraminifera: Proxies for the deep ocean carbonate system, *Earth and Planetary Science Letters*, 302(3-4), 403-413.

Raitzsch, M., E. C. Hathorne, H. Kuhnert, J. Groeneveld, and T. Bickert (2011), Modern and late Pleistocene B/Ca ratios of the benthic foraminifer *Planulina wuellerstorfi* determined with laser ablation ICP-MS, *Geology*, 39(11), 1039-1042.

Raymo, M. E., D. W. Oppo, and W. Curry (1997), The mid-Pleistocene climate transition: A deep sea carbon isotopic perspective, *Paleoceanography*, 12(4), 546-559.

Raymo, M. E., D. W. Oppo, B. P. Flower, D. A. Hodell, J. F. McManus, K. A. Venz, K. F. Kleiven, and K. McIntyre (2004), Stability of North Atlantic water masses in face of pronounced climate variability during the Pleistocene, *Paleoceanography*, 19(2).

Rickaby, R. E. M., H. Elderfield, N. Roberts, C. D. Hillenbrand, and A. Mackensen (2010), Evidence for elevated alkalinity in the glacial Southern Ocean, *Paleoceanography*, 25.

Vazquez Riveiros, N., C. Waelbroeck, L. Skinner, J. C. Duplessy, J. F. McManus, E. S. Kandiano, and H. A. Bauch (2013), The MIS 11 paradox and ocean circulation: Role of millennial scale events, *Earth and Planetary Science Letters*, 371, 258-268.

Rosenthal, Y., E. A. Boyle, and L. Labeyrie (1997), Last glacial maximum paleochemistry and deepwater circulation in the Southern Ocean: Evidence from foraminiferal cadmium, *Paleoceanography*, 12(6), 787-796.

Rosenthal, Y., M. P. Field, and R. M. Sherrell (1999), Precise determination of element/calcium ratios in calcareous samples using sector field inductively coupled plasma mass spectrometry, *Analytical Chemistry*, 71(15), 3248-3253.

Ruddiman, W. F., M. E. Raymo, D. G. Martinson, B. M. Clement, and J. Backman (1989), Pleistocene evolution: Northern hemisphere ice sheets and North Atlantic Ocean, *Paleoceanography*, 4(4), 353-412.

Schlitzer, R. 2012. Ocean Data View <http://odv.awi.de>.

Schmieder, F., T. von Dobeneck, and U. Bleil (2000), The Mid-Pleistocene climate transition as documented in the deep South Atlantic Ocean: initiation, interim state and terminal event, *Earth and Planetary Science Letters*, 179(3-4), 539-549.

Sexton, P. F., and S. Barker (2012), Onset of 'Pacific-style' deep-sea sedimentary carbonate cycles at the mid-Pleistocene transition, *Earth and Planetary Science Letters*, 321, 81-94.

Siegenthaler, U., et al. (2005), Stable carbon cycle-climate relationship during the late Pleistocene, *Science*, 310(5752), 1313-1317.

Sigman, D. M., and E. A. Boyle (2000), Glacial/interglacial variations in atmospheric carbon dioxide, *Nature*, 407(6806), 859-869.

Skinner, L. C. (2009), Glacial-interglacial atmospheric CO₂ change: a possible standing volume effect on deep-ocean carbon sequestration, *Climate of the Past*, 5(3), 537-550.

Sosdian, S., and Y. Rosenthal (2009), Deep-Sea Temperature and Ice Volume Changes Across the Pliocene-Pleistocene Climate Transitions, *Science*, 325(5938), 306-310.

Stein, R., J. Hefter, J. Grutzner, A. Voelker, and B. D. A. Naafs (2009), Variability of surface water characteristics and Heinrich-like events in the Pleistocene midlatitude North Atlantic Ocean: Biomarker and XRD records from IODP Site U1313 (MIS 16-9), *Paleoceanography*, 24.

Subhas, A.V., Adkins, J.F., Rollins, N.E., Naviaux, J, Erez, J., and W.M. Berelson (2017) Catalysis and chemical mechanisms of calcite dissolution in seawater, *PNAS*, 114, 8175-8180.

Toggweiler, J. R. (1999), Variation of atmospheric CO₂ by ventilation of the ocean's deepest water, *Paleoceanography*, 14(5), 571-588.

Toggweiler, J. R. (2008) Origin of the 100,000-year timescale in Antarctica temperatures and atmospheric CO₂, *Paleoceanography*, Vol. 23, PA2211, doi:10.1029/2006PA001405,

Tzedakis, P. C., D. Raynaud, J. F. McManus, A. Berger, V. Brovkin, and T. Kiefer (2009), Interglacial diversity, *Nature Geoscience*, 2(11), 751-755.

Wang, P. X., J. Tian, X. R. Cheng, C. L. Liu, and J. Xu (2003), Carbon reservoir changes preceded major ice-sheet expansion at the mid-Brunhes event, *Geology*, 31(3), 239-242.

Yu, J., and H. Elderfield (2007), Benthic foraminiferal B/Ca ratios reflect deep water carbonate saturation state, *Earth and Planetary Science Letters*, 258(1-2), 73-86.

Yu, J., H. Elderfield, and A. M. Piotrowski (2008), Seawater carbonate ion-delta C-13 systematics and application to glacial-interglacial North Atlantic ocean circulation, *Earth and Planetary Science Letters*, 271(1-4), 209-220.

Accepted Article
Yu, J., W. S. Broecker, H. Elderfield, Z. Jin, J. McManus, and F. Zhang (2010), Loss of Carbon from the Deep Sea Since the Last Glacial Maximum, *Science*, 330(6007), 1084-1087.

Yu, J., R. F. Anderson, Z. Jin, J. W. B. Rae, B. N. Opdyke, and S. M. Eggins (2013), Responses of the deep ocean carbonate system to carbon reorganization during the Last Glacial-interglacial cycle, *Quaternary Science Reviews*, 76, 39-52.

Yu, J., R. F. Anderson, Z. Jin, L. Menviel, F. Zhang, F. J. Ryerson, and E. J. Rohling (2014), Deep South Atlantic carbonate chemistry and increased interocean deep water exchange during last deglaciation, *Quaternary Science Reviews*, 90, 80-89.

Yu, J., et al. (2016), Sequestration of carbon in the deep Atlantic during the last glaciation, *Nature Geoscience*, 9(4), 319-324.




Figure 1 Ocean Data Viewer bathymetric map showing modern locations of DSDP site 607 (depth=3.4 km; red symbol) used in this study and from previously published data (grey/black symbols; BOFS8K depth=4.0 km; MD95-2039 depth=3.4 km; MD01-2446 depth=3.6km; ODP 1089 depth=4.6 km; TT013-PC72 depth=4.3 km) are shown in the map (Schlitzer, 2012).




Figure 2 Deep CO_3^{2-} profiles from (A) the North Atlantic near DSDP site 607 and (B) equatorial IndoPacific, nearby site TT013-PC72, used in Kerr et al. (2017) B/Ca record. In the modern ocean, basin-specific differences in deep ocean CO_3^{2-} occur as a result of the formation of North Atlantic Deep Water (NADW), which fills the deep Atlantic today with preservation-friendly, high- CO_3^{2-} water whereas the deep Pacific is filled with corrosive low- CO_3^{2-} water in response. The saturation state ($\Delta[\text{CO}_3^{2-}]$) of the deep ocean modulates the depth in the ocean where CaCO_3 preservation (above) gives way to dissolution (below) termed the lysocline whereby its position is deeper in the Atlantic with more CaCO_3 burial per unit area relative to the Pacific (Dunne et al. 2012). Carbonate system data are from nearby GLODAP sites (Key et al. 2004)




Figure 3 DSDP site 607 records of benthic foraminiferal (A) $\delta^{18}\text{O}$ and (B) $\delta^{13}\text{C}$ (Ruddiman et al., 1989) and (C) B/Ca ratios from this study (DSDP 607 and CHN82-24-23PC) and previously published deep North Atlantic records from BOFS 8K (Yu et al. 2008), MD95-2039 (Yu et al. 2016), MD01-2446 (Yu et al. 2016) and DSDP site 607 (Lear et al. 2016). All records are on the LRO4 age model and MIS are highlighted based on LRO4 stack alignment (Lisiecki & Raymo, 2005). Error bars show analytical precision for B/Ca (r.s.d. = 4.0%).




Figure 4 Interglacial and glacial B/Ca trends across the mid-to-late Pleistocene determined from the composite DSDP 607 and CHN82-24-23PC records. Data from BOFS 8K are used to estimate G/IG change across the H-LGM. Linear trends in panel A showcase the overall long-term decrease in interglacial and glacial B/Ca values. Note the steeper decrease in glacial relative to interglacial B/Ca values.




Figure 5 DSDP site 607 (A) %NCW at site 607 and the re-occupation site U1313 plotted with 95% confidence intervals (Raymo et al. 1997; Lang et al. 2016) (B) $\delta^{13}C_b$ (Ruddiman et al. 1989) (C) %CaCO₃ from DSDP site 607 and VM30-97 (Ruddiman et al. 1989), (D) compiled deep North Atlantic ocean B/Ca and ΔCO_3^{2-} from DSDP 607 and CHN82-24-23PC and others (Yu et al. 2008; Yu et al. 2016; Lear et al. 2016) and (E) atmospheric CO₂ reconstructions from ice cores (Petit et al. 1999; Siegenthaler et al. 2005; Luethi et al. 2008; Bereiter et al. 2015), blue ice (Higgins et al. 2015) and boron isotopes (Hönisch et al. 2009; Chalk et al. 2017). Error bars show ΔCO_3^{2-} uncertainty window of $\pm 9 \mu\text{mol/kg}$ from Yu et al. (2007) calibration. Boxes around the 1 Ma data indicate an age uncertainty of $\pm 89\text{ky}$ as presented in Higgins et al. (2015). The Mid-Brunhes Event and Mid-Pleistocene transition are denoted as MBE and MPT.




Figure 6 %CaCO₃ records from (A) ODP 1089 (Hodell et al (2001)) in the deep South Atlantic representing the deep Pacific and (B) DSDP 607 and VM30-97 (Ruddiman et al. 1989) in the deep North Atlantic alongside B/Ca-based ΔCO₃²⁻ reconstruction from the (C) equatorial Indo-Pacific (core TT013-PC72; Kerr et al. 2017) and (D) compiled deep North Atlantic ΔCO₃²⁻ with (E) atmospheric CO₂ ice core (Petit et al. 1999; Siegenthaler et al. 2005; Luethi et al. 2008; Bereiter et al. 2015) and B-isotope derived reconstructions (Hönisch et al. 2009; Chalk et al. 2017). Dashed line on panel D represents 0 μmol/kg. Error bars show ΔCO₃²⁻ uncertainty window of ±9 μmol/kg from Yu et al. (2007) calibration. Note inverted axis in panel B.




Figure 7 Deep CO_3^{2-} cycles across the Pleistocene as estimated by a seven-box model presented in Toggweiler (2008; see Figure 4) showcasing the classic CaCO_3 compensation mechanism. Note the time axis is inverted.