

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:<https://orca.cardiff.ac.uk/id/eprint/114406/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Ayorinde, A.A., Bhattacharya, S. , Druce, K.L., Jones, G.T. and Macfarlane, G.J. 2017. Chronic pelvic pain in women of reproductive and post-reproductive age: a population-based study. *European Journal of Pain* 21 (3) , pp. 445-455. 10.1002/ejp.938

Publishers page: <http://dx.doi.org/10.1002/ejp.938>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Table S1: Odds ratios (ORs) for interaction terms to assess the differences in factors associated with CPP in women of reproductive age compared to post-reproductive age women

| Interaction term | OR (95%CI) |
|---|-------------------|
| Reproductive age X infertility | 1.96 (0.82–4.68) |
| Reproductive age X termination of pregnancy | 1.04 (0.49–2.18) |
| Reproductive age X miscarriage | 1.97 (0.99–3.91) |
| Reproductive age X nulliparity | 0.73 (0.37–1.42) |
| Reproductive age X caesarean delivery | 1.34 (0.58–3.11) |
| Reproductive age X mild fatigue | 1.05 (0.52–2.10) |
| Reproductive age X moderate/severe fatigue | 1.01 (0.52–1.96) |
| Reproductive age X somatic symptoms | 0.51 (0.28–0.92) |
| Reproductive age X depression | 0.74 (0.38–1.46) |