

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/116113/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Hewitt, Jonathan , Marke, Margaret, Honeyman, Calum, Huf, Simon, Lai, Aida, Dong, Anni, Wright, Tom, Blake, Sarah, Fallaize, Rebecca, Hughes, Jane L, Pearce, Lyndsay and McCarthy, Kathryn 2018. Cognitive impairment in older patients undergoing colorectal surgery. *Scottish Medical Journal* 63 (1) , pp. 11-15.
10.1177/0036933017750988

Publishers page: <http://dx.doi.org/10.1177/0036933017750988>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.






Page Proof Instructions and Queries

Journal Title: Scottish Medical Journal (SCM)

Article Number: 750988

Thank you for choosing to publish with us. This is your final opportunity to ensure your article will be accurate at publication. Please review your proof carefully and respond to the queries using the circled tools in the image below, which are available by clicking “Comment” from the right-side menu in Adobe Reader DC.*

Please use *only* the tools circled in the image, as edits via other tools/methods can be lost during file conversion. For comments, questions, or formatting requests, please use . Please do *not* use comment bubbles/sticky notes .



*If you do not see these tools, please ensure you have opened this file with Adobe Reader DC, available for free at get.adobe.com/reader or by going to Help > Check for Updates within other versions of Reader. For more detailed instructions, please see us.sagepub.com/ReaderXProofs.

No.	Query
	Please confirm that all author information, including names, affiliations, sequence, and contact details, is correct.
	Please review the entire document for typographical errors, mathematical errors, and any other necessary corrections; check headings, tables, and figures.
	Please confirm that the Funding and Conflict of Interest statements are accurate.
	Please ensure that you have obtained and enclosed all necessary permissions for the reproduction of artistic works, (e.g. illustrations, photographs, charts, maps, other visual material, etc.) not owned by yourself. Please refer to your publishing agreement for further information.
	Please note that this proof represents your final opportunity to review your article prior to publication, so please do send all of your changes now.
AQ: 1	Please provide the job titles of all the authors, per journal style.
AQ: 2	Please provide the postal code for correspondence.



Cognitive impairment in older patients undergoing colorectal surgery

Jonathan Hewitt, Margaret Marke, Calum Honeyman, Simon Huf, Aida Lai, Anni Dong, Tom Wright, Sarah Blake, Rebecca Fallaize, Jane L Hughes, Lyndsay Pearce and Kathryn McCarthy

Abstract

Background: With increasing numbers of older people being referred for elective colorectal surgery, cognitive impairment is likely to be present and affect many aspects of the surgical pathway. This study is aimed to determine the prevalence of cognitive impairment and assess it against surgical outcomes.

Methods: The Montreal Cognitive Assessment (MoCA) was carried out in patients aged more than 65 years. We recorded demographic information. Data were collected on length of hospital stay, complications and 30-day mortality.

Results: There were 101 patients assessed, median age was 74 years (interquartile range = 68–80), 54 (53.5%) were women. In total, 58 people (57.4%) ‘failed’ the Montreal Cognitive Assessment test (score ≤ 25). There were two deaths (3.4%) within 30 days of surgery in the abnormal Montreal Cognitive Assessment group and none in the normal group. Twenty-nine (28.7%) people experienced a complication. The percentage of patients with complications was higher in the group with normal Montreal Cognitive Assessment (41.9%) than abnormal Montreal Cognitive Assessment (19.9%) ($p=0.01$) and the severity of those complications were greater (chi-squared for trend $p=0.01$). The length of stay was longer in people with an abnormal Montreal Cognitive Assessment (mean 8.1 days vs. 5.8 days, $p=0.03$).

Conclusion: Cognitive impairment was common, which has implications for informed consent. Cognitive impairment was associated with less postoperative complications but a longer length of hospital stay.

Keywords

Older people, surgery, cognitive impairment, Montreal Cognitive Assessment

Introduction

In the developed world, the number of surgical procedures being performed on older patients is rising faster than the rate of population increase.¹ This is indicative of age no longer being a contraindication to surgical intervention. Older people are being referred to secondary care for elective colorectal surgery, including cancer surgery. This population includes those with cognitive impairment. The presence of cognitive impairment has several implications. Firstly, it has been shown that people with cognitive impairment have worse outcomes following surgery.^{2,3} Secondly, informed consent may be influenced by the presence of cognitive impairment.⁴ Finally, enhanced recovery programmes are now commonplace in elective colorectal surgery, with proven benefits in terms of length of hospital stay and reduced

complication rates.⁵ Whether these regimens are suitable and applicable to the older confused person is less clear.

In 2005, Nasreddine et al. characterised the Montreal Cognitive Assessment (MoCA) (www.mocat-est.org), a tool that is particularly good at detecting mild cognitive impairment. It has subsequently become one of the commonest and most validated tools for assessing cognitive function.⁶ Increasing use

Division of Population Medicine, Cardiff University, University Hospital Wales, UK [AQ1]

Corresponding author:

Jonathan Hewitt, Division of Population Medicine, Cardiff University, University Hospital Wales, Cardiff, UK. [AQ2]
Email: hewittj2@cardiff.ac.uk

in the UK was, in part, driven by copyright restrictions regarding the use of the Mini Mental State Examination (MMSE),⁷ previously the cognitive assessment tool of choice. In contrast to the MMSE, the MoCA is cost-free to use in clinical and educational based settings.

The MoCA has been assessed across a range of surgical settings^{4,8,9} but never in elective colorectal surgery. The aim of this study was to assess the prevalence of cognitive impairment using the MoCA and assess any relationships between these results and a range of surgical outcomes; complications, death and length of hospital stay.

Methods

The MoCA is a 30-point questionnaire (see Figure 1, the MoCA assessment) (www.mocatest.com). Any score of 26 and above is considered normal. The MoCA has been used across a broad spectrum of clinical conditions and is well tested and validated against other cognitive assessment methods.^{10,11} It is available in almost every major language.

We used the MoCA test in English (Original Version, 7.1). Staff gathering MoCA data underwent training in the implementation and use of the questionnaire, prior to commencing the study.

From June 2012 to June 2014, we prospectively assessed pre-operative cognition in all patients 65 years and older who were undergoing elective colorectal surgery for benign (predominantly diverticular disease) or malignant disease (colorectal cancer). Only patients unable to complete the test owing to inadequate visual perception were excluded from the study. During routine preoperative assessment, researchers based at North Bristol NHS Trust collected the MoCA data. Participant's age and sex were recorded. Patients were grouped by age (65–74, 75–84, and above 85 years). To characterise co-morbidity data were collected on the number of current medications¹² (grouped into < or \geq 5). Delirium was not assessed. Data were collected on length of hospital stay (rounded up as whole day integers), complications and 30-day mortality. Complications were characterised using the Clavien-Dindo scoring system.¹³

Data were collected from the hospital electronic patient records, clinical case notes and prescribing charts. All data were collated using a password-protected spreadsheet. Patients were anonymised, with identifiable data removed. Data were handled and stored according to local data management guidelines.

The study was registered according to local guidelines. As the study collated information collected as part of routine clinical care, mental state estimation being recommended as trust policy, the study was

deemed a service evaluation project and as such did not require ethical approval.

Data analysis was carried out using STATA version 13. Continuous data are summarised as mean and median (interquartile range (IQR)) values and categorical data as frequencies with percentages. Comparisons were performed using Wilcoxon rank sum and chi-squared tests.

Results

There were 101 patients included in the study. Median age was 74 years (IQR = 68–80), two patients had missing data for age. Fifty patients were aged between 65 and 74 years, 37 between 75 and 84 years and 10 older than 85 years. There were 54 (53.5%) women. There were 58 people (57.4%) who failed the MoCA test (score \leq 25). Overall, the mean score for the MoCA examination was 24.1 ('normal' \geq 26, range = 8–30). Participants were taking an average number of 4.5 medications each (range = 0–11). There were 46 people (45.5%) taking five or more medications.

An abnormal MoCA was associated with increasing age group ($p=0.03$) (test for trend) or taking five or more medications ($p=0.01$). There were two deaths (3.4%) within 30 days of surgery in the abnormal MoCA group, there were no deaths in the normal MoCA group. Twenty-nine (28.7%) patients experienced a complication (Clavien-Dindo Classification I–V). The percentage of patients with complications was higher in the group with normal MoCA (41.9%) than abnormal MoCA (19.9%) ($p=0.01$) and the severity of those complications was greater (chi-squared for trend $p=0.01$). People with an abnormal MoCA remained in hospital for longer (mean 8.1 days vs. 5.8 days, $p=0.03$). These results are summarised in Table 1.

Procedures included right hemicolectomy, small bowel resection, reversal of Hartmans procedure, ileostomy, anterior resection and ventral mesh rectopexy. They were performed by both open and laparoscopic technique.

Discussion

This study has demonstrated that cognitive impairment, measured using the MoCA, is common in this elective older colorectal population, with 57.4% of patients having an abnormal score. Cognitive impairment was associated with a longer length of hospital stay. Despite this, people with cognitive impairment (MoCA score \leq 25) experienced lower rates of complications. Furthermore, these complications were less severe.

This is the first study to document the prevalence of preoperative cognitive impairment in patients






MONTREAL COGNITIVE ASSESSMENT (MOCA) Version 7.1 Original Version					NAME : Education : Sex :		Date of birth : DATE :																				
VISUOSPATIAL / EXECUTIVE					Copy cube 		Draw CLOCK (Ten past eleven) (3 points)		POINTS ___/5																		
																											
NAMING											___/3																
MEMORY					Read list of words, subject must repeat them. Do 2 trials, even if 1st trial is successful. Do a recall after 5 minutes.		<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td></td> <td style="text-align: center;">FACE</td> <td style="text-align: center;">VELVET</td> <td style="text-align: center;">CHURCH</td> <td style="text-align: center;">DAISY</td> <td style="text-align: center;">RED</td> </tr> <tr> <td style="text-align: center;">1st trial</td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td style="text-align: center;">2nd trial</td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </table>			FACE	VELVET	CHURCH	DAISY	RED	1st trial						2nd trial						No points
	FACE	VELVET	CHURCH	DAISY	RED																						
1st trial																											
2nd trial																											
ATTENTION					Read list of digits (1 digit/ sec.).		Subject has to repeat them in the forward order [] 2 1 8 5 4 Subject has to repeat them in the backward order [] 7 4 2		___/2																		
ATTENTION					Read list of letters. The subject must tap with his hand at each letter A. No points if ≥ 2 errors		[] FBACMNAAJKLBAFAKDEAAAJAMOF AAB		___/1																		
ATTENTION					Serial 7 subtraction starting at 100 [] 93 [] 86 [] 79 [] 72 [] 65		4 or 5 correct subtractions: 3 pts , 2 or 3 correct: 2 pts , 1 correct: 1 pt , 0 correct: 0 pt		___/3																		
LANGUAGE					Repeat : I only know that John is the one to help today. [] The cat always hid under the couch when dogs were in the room. []				___/2																		
LANGUAGE					Fluency / Name maximum number of words in one minute that begin with the letter F [] ____ (N ≥ 11 words)				___/1																		
ABSTRACTION					Similarity between e.g. banana - orange = fruit [] train - bicycle [] watch - ruler				___/2																		
DELAYED RECALL					Has to recall words WITH NO CUE		<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="text-align: center;">FACE</td> <td style="text-align: center;">VELVET</td> <td style="text-align: center;">CHURCH</td> <td style="text-align: center;">DAISY</td> <td style="text-align: center;">RED</td> </tr> <tr> <td style="text-align: center;">[]</td> <td style="text-align: center;">[]</td> <td style="text-align: center;">[]</td> <td style="text-align: center;">[]</td> <td style="text-align: center;">[]</td> </tr> </table>		FACE	VELVET	CHURCH	DAISY	RED	[]	[]	[]	[]	[]	Points for UNCUEDE recall only								
FACE	VELVET	CHURCH	DAISY	RED																							
[]	[]	[]	[]	[]																							
Optional					Category cue		<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </table>																				
Optional					Multiple choice cue		<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </table>																				
ORIENTATION					[] Date [] Month [] Year [] Day [] Place [] City				___/6																		
© Z.Nasreddine MD					www.mocatest.org		Normal ≥ 26 / 30		TOTAL ___/30																		
Administered by: _____							Add 1 point if ≤ 12 yr edu																				

Figure 1. MoCA test.

Table 1. Complications and length of stay.

	Overall	Normal MoCA (n = 43) (42.6%) (≥26)	Abnormal MoCA (n = 58) (57.4%) (≤25)	p Value
Complications				
Yes	29	18 (41.9)	11 (19.0)	0.01 ^a
No	72	25 (58.1)	47 (81.0)	
Clavien Dindo				
0	72	25 (58.1)	47 (81.0)	0.02 ^b
Class				
I	8	4 (9.3)	4 (6.9)	
II	15	11 (25.6)	3 (5.2)	
III	0	0	1 (1.7)	
IV	4	3 (7.0)	1 (1.7)	
V	2	0	2 (3.5)	
Length of stay				
Median (IQR)	5 (3–8)	5 (3–6)	8 (4–9)	0.03 ^c

Note: Clavien_Dindo classification – I: no need for treatment; II: pharmacological treatment; III: requiring surgical or endoscopic or radiological treatment (IIIa – not under GA, IIIb – under GA); IV: life threatening complication (IVa – single organ dysfunction, IVb – multi organ dysfunction); V: death of a patient.

^aChi-square.

^bChi-square for trend.

^cWilcoxon Ranksum.

undergoing elective colorectal surgery. Therefore, direct comparisons with other studies are limited. One of the more comparable studies was conducted by our team in acute surgery. That paper assessed the prevalence of cognitive impairment in emergency general surgical patients in three UK centres. MoCA detected cognitive impairment in over 70% of the population,⁴ higher than the present study. One possible explanation is delirium. Delirium is common in all acute hospital presentations of the older person,¹⁴ estimated to be present in up to one-third of acute vascular¹⁵ and a quarter of acute orthopaedic patients.¹⁶ The MoCA score does not test for delirium, hence that study would not differentiate between the two. Thus, the difference between the two figures (57.4% vs. 70%) may be attributable to delirium. Delirium screening might be advocated for future research studies in this area.

There are clinical implications of managing the older elective surgical patient with cognitive impairment. People with cognitive impairment are more likely to develop incident delirium³ and long-term cognitive dysfunction following surgery. For example, Monk et al. reported cognitive dysfunction following surgery in 41.4% of older patients discharged after major non-cardiac surgery.¹⁷ Kline and colleagues also

demonstrated that postoperative cognitive dysfunction is higher in people with pre-existing disease,¹⁸ and a recent systematic review confirmed a higher rate of death in this patient group.¹⁹

Modification of anaesthetic techniques, have been shown to reduce the incidence of post-operative delirium.²⁰ Similarly, multi-task exercise programmes have shown improvements in function and cognition in elderly patients.²¹ Ultimately, pre-operative identification and optimisation of those most at risk of delirium and post-operative cognitive dysfunction may lead to improvements in post-operative recovery, quality of life and mortality in older patients who are most at risk.

Perhaps the most striking finding of these results is the reduced rate of complications demonstrated in our population with an abnormal MoCA, both in the absolute number of complications and the severity of those recorded. This may be attributable to under-reporting by cognitively impaired patients leading to an under-diagnosis of complications by medical staff. For example, cognitively impaired people are less likely to report pain²² and other physical symptoms. However, this might arguably result in increasing severity of complications when recognised. Prolonged length of stay in the abnormal MoCA group may be attributable to increase in required community care and social support on discharge. It may also reflect unrecognised (potentially minor) complications delaying the discharge from hospital in this group. Another potential complication may be that only physically fitter people, with cognitive impairment, were offered surgery, hence the lower rate of complications. However, we do not data to support that assumption.

The other major implication of these findings is whether the 57.4% of participants with abnormal MoCA scores are able to give informed consent for the operation they are about to undergo. The MoCA test is sensitive enough to detect mild cognitive impairment.^{6,23} Therefore, some (perhaps a majority) of the abnormal results obtained represent people with substantial residual cognitive ability. While consent is decision-specific, it seems highly likely, based on these data that many of the individuals in this study may not have had sufficient capacity to complete fully informed consent. Whilst not addressed in this study, the study raises concerns that consent may not always be valid, an area for consenting surgeons to consider. Similarly, our other concern is whether cognitively impaired patients are fully able to comply with the enhanced recovery protocol, which involves early mobilisation, enhanced nutrition and the need to retain information and follow instructions. Perhaps there is a need to tailor a recovery pathway and indeed consenting process specifically for patients with cognitive impairment. Both these aspects of surgical care also highlight the

importance of comprehensive assessment of older surgical patients and particularly engaging patients, their relatives and their carers in managing expectations and clinical decision-making.

These data illustrate the high level of cognitive impairment in a population undergoing elective colorectal surgery. They also suggest a reduced rate of complications, and a greater length of hospital stay. Further larger scale studies may fully elucidate the impact of cognitive impairment on elective colorectal surgery and whether enhanced recovery programs are fully tailored to the cognitively impaired.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

References

- Partridge JS, Harari D and Dhesi JK. Frailty in the older surgical patient: a review. *Age Ageing* 2012; 41: 142–147.
- Boos GL, Soares LF and Oliveira Filho GR. Postoperative cognitive dysfunction: prevalence and associated factors. *Revista Brasileira de Anestesiologia* 2005; 55: 517–524.
- Ni Chonchubhair A, Valacio R, Kelly J, et al. Use of the abbreviated mental test to detect postoperative delirium in elderly people. *Br J Anaesth* 1995; 75: 481–482.
- Hewitt J, Williams M, Pearce L, et al. The prevalence of cognitive impairment in emergency general surgery. *Int J Surg* 2014; 12: 1031–1035.
- Zhao JH, Sun JX, Gao P, et al. Fast-track surgery versus traditional perioperative care in laparoscopic colorectal cancer surgery: a meta-analysis. *BMC Cancer* 2014; 14: 607.
- Nasreddine ZS, Phillips NA, Bedirian V, et al. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc* 2005; 53: 695–699.
- Newman JC and Feldman R. Copyright and open access at the bedside. *N Engl J Med* 2011; 365: 2447–2449.
- de Guise E, Leblanc J, Champoux MC, et al. The minimal state examination and the Montreal Cognitive Assessment after traumatic brain injury: an early predictive study. *Brain Inj* 2013; 27: 1428–1434.
- Wong GK, Ngai K, Lam SW, et al. Validity of the Montreal Cognitive Assessment for traumatic brain injury patients with intracranial haemorrhage. *Brain Inj* 2013; 27: 394–398.
- McLennan SN, Mathias JL, Brennan LC, et al. Validity of the Montreal Cognitive Assessment (MoCA) as a screening test for mild cognitive impairment (MCI) in a cardiovascular population. *J Geriatr Psychiatry Neurol* 2011; 24: 33–38.
- Sweet L, Van Adel M, Metcalf V, et al. The Montreal Cognitive Assessment (MoCA) in geriatric rehabilitation: psychometric properties and association with rehabilitation outcomes. *Int Psychogeriatr* 2011; 23: 1582–1591.
- George J, Vuong T, Bailey MJ, et al. Development and validation of the medication-based disease burden index. *Ann Pharmacother* 2006; 40: 645–650.
- Dindo D, Demartines N and Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004; 240: 205–213.
- Fong TG, Davis D, Growdon ME, et al. The interface between delirium and dementia in elderly adults. *Lancet Neurol* 2015; 14: 823–832.
- Sasajima Y, Sasajima T, Azuma N, et al. Factors related to postoperative delirium in patients with lower limb ischaemia: a prospective cohort study. *Eur J Vasc Endovasc Surg* 2012; 44: 411–415.
- Evered LA, Silbert BS, Scott DA, et al. Preexisting cognitive impairment and mild cognitive impairment in subjects presenting for total hip joint replacement. *Anesthesiology* 2011; 114: 1297–1304.
- Monk TG, Weldon BC, Garvan CW, et al. Predictors of cognitive dysfunction after major noncardiac surgery. *Anesthesiology* 2008; 108: 18–30.
- Kline RP, et al. Surgery and brain atrophy in cognitively normal elderly subjects and subjects diagnosed with mild cognitive impairment. *Anesthesiology* 2012; 116: 603–612.
- Salluh JI, Wang H, Schneider EB, et al. Outcome of delirium in critically ill patients: systematic review and meta-analysis. *BMJ* 2015; 350: h2538.
- Chan MT, Cheng BC, Lee TM, et al. Bis-guided anesthesia decreases postoperative delirium and cognitive decline. *J Neurosurg Anesthesiol* 2013; 25: 33–42.
- Hars M, Herrmann FR, Gold G, et al. Effect of music-based multitask training on cognition and mood in older adults. *Age Ageing* 2014; 43: 196–200.
- Hadjistavropoulos T, Herr K, Turk DC, et al. An interdisciplinary expert consensus statement on assessment of pain in older persons. *Clin J Pain* 2007; 23: S1–S43.
- Luis CA, Keegan AP and Mullan M. Cross validation of the Montreal Cognitive Assessment in community dwelling older adults residing in the southeastern US. *Int J Geriatr Psychiatry* 2009; 24: 197–201.