

# Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <http://orca.cf.ac.uk/117716/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Silva, Ana Rita R., Santos, Catia and Ferreira, Nuno 2019. Multigenerational effects of carbendazim in daphnia magna: from a subcellular to a population level. *Environmental Toxicology and Chemistry* 38 (2), pp. 412-422. 10.1002/etc.4307 file

Publishers page: <https://doi.org/10.1002/etc.4307> <<https://doi.org/10.1002/etc.4307>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



ろ Multigenerational effects of carbendazim in *Daphnia magna*: from a subcellular to  
わ a population level

わ Ana Rita R. Silva\*<sup>1</sup>, Célia Santos<sup>1,2</sup>, Nuno G. C. Ferreira<sup>1</sup>, Rui Morgado<sup>1</sup>, Diogo N. Cardoso<sup>1</sup>,  
み Andreia Cruz<sup>1</sup>, Sílvia Mendo<sup>1</sup>, Amadeu M. V. M. Soares<sup>1</sup> and Susana Loureiro\*<sup>1</sup>

を <sup>1</sup>Department of Biology & CESAM, University of Aveiro, 3810-193 Aveiro, Portugal

ん <sup>2</sup>Department of Biology, Terrestrial Ecology Unit, Ghent University, K.L. Ledeganckstraat 35, 9000  
う Ghent, Belgium

、 \*E mail: [ritas@ua.pt](mailto:ritas@ua.pt); [sloureiro@ua.pt](mailto:sloureiro@ua.pt)

れ

ろ Abstract

ろ Anthropogenic activities such as the use of pesticides may affect in some way  
わ aquatic biota populations, due to potential agricultural runoffs or disposals.  
ろ Carbendazim is one example of a widely used fungicide with a high potential to end up  
ろ in aquatic ecosystems through runoff. Deleterious effects observed at the individual  
ろ level are possibly explained by changes in homeostasis at cellular and both can then be  
ろ used to predict effects at the population level. In the present study, an isoclinal  
ろ population of *Daphnia magna* (clone k6) was exposed to concentration that mimics  
ろ relevant levels of carbendazim in the environment during twelve generations. The  
れ effects of carbendazim on biochemical biomarkers (cholinesterase, catalase and  
ろ glutathione S-transferase), lipid peroxidation and energy-related parameters  
ろ (carbohydrates, lipids and proteins jointly with energy available and energy  
ろ consumption), parental longevity, and population growth ( $r$ ) were assessed in some  
ろ generations. The long-term exposure to carbendazim presented no effect on the intrinsic  
ろ rate of natural increase ( $r$ ) of adult *D. magna*. However, daphnids longevity decreased at  
ろ the F12 when compared to daphnids from control. Cholinesterases, glutathione S-  
ろ transferase and lipid peroxidation showed differences between the exposed and non-  
ろ exposed populations. However, for catalase and energy related-parameters no  
ろ differences were observed between these two populations. Natural variability was  
ろ observed throughout the test period, under control conditions, within the twelve  
ろ generations. Overall, carbendazim induced some effects at the subcellular level that  
ろ were translated to longevity, but latter vanishing in terms of population effects.

ろ

ろ Key words *Daphnia magna*, multigenerations, carbendazim, biochemical biomarkers,  
ろ energy reserves, DNA damage



これ Carbendazim (CBZ: methyl-2-benzimidazole carbamate) has been used for many years as a fungicide in several agricultural crops, including potatoes, strawberries, onions, wheat, oranges, among others (EU Pesticide Database, 2016) and consequently it is likely to be released from spring to autumn. CBZ is the active breakdown product of benomyl, which is also a systemic fungicide (Davidse, 1973). Emissions to aquatic systems include spray-drift or run-off from crops and soils (after rainfall events), that may occur in a cadence of continuity or pulses (WHO, 1993). CBZ was considered persistent in the water layer (Cuppen et al., 2000), and the maximum reported concentrations was 4.5 µg/L in surface waters of the basin of the Traiguera river in Chile (Palma et al., 2004). In addition, in a previous study, a multigenerational test showed that DNA damage (genotoxicity) increased throughout generations of *D. magna* exposed to CBZ (Silva et al., 2017). Therefore selecting several biomarkers that may provide additional information on other mechanisms induced by CBZ under long term exposures may be useful to understand effects at the individual and population level. Several biomarkers were selected: cholinesterase (ChE) activity, a well-known target site of carbamate pesticides, which inhibits its activity triggering neurotoxic effects in *D. magna* (Barata et al., 2004); catalase (CAT) as an antioxidant enzyme (Brown et al., 2004); glutathione S-transferases (GST) which is related with biotransformation and antioxidant defense (Hyne and Maher, 2003); lipid peroxidation (LPO) rate, which is associated with cell damage (Barata et al., 2005); and energy reserves related parameters.

## 2. Materials and methods

### 2.1 Test organism and test chemicals

The water flea *D. magna* Straus clone K6 (originally from Antwerp, Belgium) was obtained from continuous culture maintained in a laboratory at the University of Aveiro (Portugal) and cultured in American Society for Testing and Materials moderated-hard-water medium (ASTM, 1980) (temperature 20 ± 1 °C; photoperiod 16h:8h (light:dark)). Daphnids were fed with the microalga *Raphidocelis subcapitata* at a concentration of 3x10<sup>5</sup> cells/mL and supplemented with an organic extract (Marinure seaweed extract, supplied by Glenside Organics Ltd.).

A stock solution of carbendazim (CAS No. 10605-21-7, 99.4% purity, Bayer Crop Science) was prepared in ASTM medium and used to maintain the multigenerational test. Chemical analyses were performed to confirm concentrations of CBZ in the test

medium at Marchwood Scientific Services, Southampton, UK. CBZ was analysed by Liquid Chromatography-Mass Spectrometry (LCMS-MS) using the QuERCHERS (quick, easy, cheap, effective, rugged, safe) method (details for the chemical analyses can be found in supplementary material).

## 2.2 Multigenerational experimental setup

For the multigenerational approach the concentration of 5 µg/L of CBZ was chosen based on the results from a reproduction test, where an equivalent no-observed-effect-concentration (NOEC) was derived (Silva et al., 2015). The experimental design of the multigenerational experiment is shown in Figure 1 and will be further described in the following subtopics (where F represents generation). For additional details please check Figure 1 SD.




Figure 1. Multigenerational experimental approach with *Daphnia magna*. Each box represents a generation and the respective endpoints evaluated or bioassays carried out. F represents generation.

During the multigenerational experiment, an isoclinal population of *D. magna* was exposed to 5 µg/L of CBZ throughout 12 generations, with the intention to look forward and test the effects in a large temporal time scale. The choice for the 12 generations allows the comparison to previous results available (Silva et al., 2017) along with the fact that this fungicide may be present during several months in the field,

as its application can occur for different agricultural crops and at different times during the year (EU Pesticide Database, 2016). Simultaneously, a second isoclonal population of daphnids was maintained under clean medium and used as control. The population of daphnids that was maintained in a control/clean condition (ASTM, *R. subcapitata* and organic extract but no CBZ) will be designated throughout the study as Dph\_Clean and population of daphnids exposed to CBZ (*ASTM, R. subcapitata*, organic extract and CBZ) as Dph\_CBZ.

*D. magna* multigenerational bioassays were carried out in triplicate using glass vessels (1L volume capacity) with 20 daphnids each (<24h neonates), for both isoclonal populations. Each replicate consisted in ASTM medium with *R. subcapitata* (concentration of  $3 \times 10^5$  cells/mL) and organic extract (Marinure seaweed extract, supplied by Glenside Organics Ltd.), spiked with CBZ in the case of the Dph\_CBZ population. The medium was completely renewed three times a week. When neonates were not required for any parameter evaluation, they were discarded and removed on a daily basis. Each subsequent generation was always initiated by using third brood neonates (<24h) of the previous one and maintained in the same conditions (either Dph\_Clean or Dph\_CBZ).

In the F0, F3, F6 and F12 generations several parameters were evaluated (see sections below). Adults' reproduction was reported in time, along with the time for the first brood, and also survival and longevity. Their broods were used to evaluate offspring fitness through sensitivity tests and biochemical markers. In order to control differences in daphnids' responses from sensitivity variations/biological variation in organisms (Loureiro et al., 2010; Novais and Amorim, 2015) all endpoints were simultaneously assessed in neonates from Dph\_Clean and Dph\_CBZ throughout generations.

### 2.2.1. Individual and Population level endpoints

The total number of neonates till the fifth brood (21 days) and parental survival were recorded for both populations and the intrinsic rate of natural increase ( $r$ ) calculated, using the Euler Lotka equation (Lotka, 1913):

(Equation 1) 
$$\sum_{x=0}^{\infty} l_x m_x e^{-rx} = 1$$

where  $l_x$  is the proportion of individuals surviving to age  $x$ ,  $m_x$  is per-capita fecundity, and  $x$  represents days.

Parental longevity (lifespan in days) was also recorded. Adult daphnids were kept under the same conditions as in the test, until they die. The day of death was recorded and time-response relationship, using the 50% lethal time ( $LT_{50}$ ) values, was determined.

Additionally, a sensitivity test with potassium dichromate ( $K_2Cr_2O_7$ ) was performed with multigenerational offspring, according to the OECD procedure (OECD, 2004). In brief, neonates (<24h old) were exposed to a concentration range of  $K_2Cr_2O_7$  for 24h and their immobilization recorded.

### 2.2.2. Subcellular level endpoints

All biochemical analyses were carried out in 7 days old organisms. For that, and from each treatment, *D. magna* neonates (<24h) were randomly sampled from the three multigenerational replicates, in order to accomplish a 7 day exposure design with five replicates, of 16 organisms each, for the enzymatic activities and LPO measurements, and three replicates with 20 organisms each, for the energy reserves assays (Figure 1 SD). These exposures were maintained in similar conditions as their parental exposures.

For the enzymatic determinations and LPO, organisms were collected to 1.5 mL Eppendorfs, with maximum media removal, shock frozen in liquid nitrogen and then stored at -80°C until analyses. Prior to analysis, samples were prepared using an adapted protocol described by Ferreira et al. (2010). ChE activity was measured according to the Ellman method (Ellman et al., 1961) adapted to a 96 well microplate as described in Guilhermino et al. (1996). CAT activity was determined based on the methodology described by Claiborne (1985) adapted to microplate (Ferreira et al., 2015). GST activity was determined according to the method described by Habig et al. (1974) adapted to microplate (Frasco and Guilhermino, 2002). Details for all enzyme analyses are presented in supplementary data. LPO was determined as described by Ohkawa et al. (1979) and Bird and Draper (1984), adapted to microplate, by measuring the production of thiobarbituric acid-reactive substances (TBARS) at 535 nm. For details please check the supplementary data.

Energy reserves were measured using a protocol adapted from Ferreira et al. (2015) previously described by De Coen and Janssen (1997). Total proteins, carbohydrates and lipid contents; energy consumption ( $E_c$ ), as electron transport

activity - ETS and available energy (Ea) were determined (protocol details can be found as well in supplementary data), and calculated as:

(Equation 2)  $E_c = \text{ETS activity (mj/org/min)}$

(Equation 3)  $E_a = \text{carbohydrates + lipids + proteins (mj/org)}$

### 2.3 Statistical Analysis

The mean value of the intrinsic rate of natural increase (r) was determined using the Jackknife method (Pestana et al., 2010; Taberner et al., 1993). The 50% lethal time (LT<sub>50</sub>) values were calculated using a nonlinear regression with a three-parameter logistic function using SigmaPlot v11.0 software (Systat Software Inc., 2008). To compare the LT<sub>50</sub> values obtained for Dph\_Clean and Dph\_CBZ, a generalized likelihood ratio test was applied using statistical package SPSS (SPSS 20.0.0, 2011). Normality was assessed using the Shapiro-Wilk test and homoscedasticity using Levene's equal variance test (Systat Software Inc., 2008). GST and E<sub>c</sub> data were square-root transformed to correct for normality.

Significant differences between exposure (Dph\_Clean and Dph\_CBZ) and generations (time) were checked for all endpoints (except longevity) using a two-way ANOVA with Bonferroni post-test; and generations (time) and exposure were used as fixed factors. The Two-way ANOVA were performed in SigmaPlot v11.0 software as well (Systat Software Inc., 2008). The R-squared (R<sup>2</sup>) was calculated by dividing the sum of squares of each factor and of their interaction by the total sums of squares of the two-way ANOVAs (Hullett and Levine, 2003), to evaluate the percentage of variance accounted for each factor in the ANOVAs.

## 3. Results and Discussion

### 3.1. Chemical analysis

The results of the chemical analysis showed that in the ASTM, CBZ concentration decreased in time, with a decay rate (K<sub>0</sub>) of 0.03/hour (st. error= 0.005), showing that only 18% of the initial concentration (7.21 g/L) left after 48h (described already in Silva et al. (2015)).



## 3.2. Individual and Population level endpoints

### 3.2.1. Intrinsic rate of natural increase (r)

The intrinsic rate of natural increase (r) is a representative endpoint that can provide information at the population level by traducing offspring production and adults' survival, within a time frame, into an indication of population growth (Buhl et al., 1993). This endpoint (r) usually represents a more sensitive parameter than considering only the number of neonates, because it integrates the reproduction output, number of mothers, number of broods and time (days) to the brood release. This enables also to bridge the gap on extrapolations from individuals to populations. However, this pattern was not observed in the present study, with no significant differences for r between Dph\_Clean and Dph\_CBZ (two-way ANOVA,  $F_{1,23} = 2.78$ ,  $p > 0.05$ ) and neither an interaction between exposure and generations (two-way ANOVA,  $F_{3,23} = 3.12$ ,  $p > 0.05$ ) (Fig. 2 and Table 1 SD). Similar findings by Zalizniak and Nugegoda (2006) showed no clear effects of chlorpyrifos on three successive generations of *Daphnia carinata* for the intrinsic rate of natural increase (r). This might be related with a compensation between survival, fecundity and maturation time (Zalizniak and Nugegoda, 2006), and potential an differential allocation of energy.

Some variability on data from non-exposed daphnids (Dph\_Clean) was also observed (Fig. 2), which might be related with non-fully controlled exposure conditions such as food quality and/or small variations in room temperature. Although all conditions are intended to be constant a slight inherent variability is likely to occur. In the work of Clubbs and Brooks (2007), a slight variance in the mean number of neonates/intrinsic rate of population growth was also shown even in controls of F0 and F1. In the present study, the difference obtained in r from F0 to F3 may be related to the time for the first brood release. While F0 daphnids released their first brood at day 8 (average), the F3 daphnids only released their first brood at day 10 (average). Although it is acceptable that daphnids release their first brood between 8 and 10 days old, this may provide some slight differences when evaluating datasets. Howsoever, as previously described, endpoints were compared between Dph\_Clean and Dph\_CBZ within generations, to control differences in organisms sensitivity as well (Loureiro et al., 2010).





Figure 2. Intrinsic rate of natural increase ( $r$ ) of *Daphnia magna* populations exposed throughout 12 generations to control conditions (Dph\_Clean, white dots) and to carbendazim (Dph\_CBZ, black dots). Data are expressed as mean values and standard error ( $n=3$ ).

### 3.2.2. Longevity in parental organisms

In the F0 generation, the pattern for longevity was similar between Dph\_Clean and Dph\_CBZ (Fig. 3), which was reinforced by the similar  $LT_{50}$  values: 61.97 days and 60.52 days, respectively (Table 2 SD), with no significant differences in slopes of the probit regressions between both  $LT_{50}$  values ( $\text{value} = 0.91$ ,  $p > 0.05$ ). After twelve generations (F12), longevity was affected by CBZ, with a  $LT_{50}$  value of 57.87 days, significantly lower than the 76.18 days for Dph\_Clean ( $\text{value} = 676.2$ ,  $p < 0.05$ ) (Table 2 SD). Throughout the generations the  $LT_{50}$  values were always in the same order in the Dph\_CBZ population, however in the F12 the longevity in Dph\_CBZ was lower when compared to F12 Dph\_Clean and this should be taken into account. Survival has been evaluated in daphnids in multigenerational tests, however this assessment is usually carried out only until the 21 d (corresponding to a standard reproduction test with *Daphnia*) (Chen et al., 2013; Schez et al., 2004; Tanaka and Nakanishi, 2002). Chen et al. (2013) observed that the pesticide pentachlorophenol caused an earlier mortality in F2 comparing with F0 daphnids, representing an enhanced toxic effect in the F2 generation. Increase in sensitivity due to a continuous exposure probably results from chemical bioaccumulation or transgenerational reductions in fitness (Kimberly and Salice, 2014). In addition, in the present study the  $r$  did not show any significant

われ differences between and within treatments and generations, probably representing an  
 わる higher energy investment by daphnids on reproduction upon exposure to CBZ in  
 わわ detriment of an investment in long-term survival.



わわ

わぬ Figure 3. Longevity of *Daphnia magna* populations in control conditions (Dph\_Clean, white dots) and  
 わる exposed to carbendazim (Dph\_CBZ, black dots) for several generations: F0 generation, F3 generation, F6  
 わを generation and F12 generation (n=3). Longevity (in days) is expressed as mean values of live adults and  
 わん standard error, for every 10 days.

わづ  
 わ  
 われ

### 3.2.3. Neonates fitness- sensitivity test

わる  
 わわ  
 わわ  
 わわ  
 わわ  
 わわ  
 わわ

Throughout the multigenerational experiment, the two isoclonal populations (Dph\_Clean and Dph\_CBZ) were relatively synchronised in their reproduction (with a difference of only some hours). The physiological conditions of *D. magna* were measured by looking at their sensitivity towards exposure to the reference chemical potassium dichromate. The 24h-EC50 values obtained in control daphnids (Dph\_Clean) and in carbendazim (Dph\_CBZ) from F0, F3, F6 and F12 were always within the

recommended range of 0.6 mg/l to 2.1 mg/l (Table 3 SD) (EN ISO 6341, 1996), assuring that stock organisms are reliable.

机`

### 机机 3.3. Subcellular level endpoints

机机 Several endpoints have been evaluated in multigenerational tests with *Daphnia* sp. such as mortality, reproduction or body length (Jacobasch et al., 2014; Kim et al., 2012; S@nchez et al., 2004), however the study of physiological pathways, such as neurotransmission capabilities, detoxification potential or antioxidant capacity is less common. Carbamate pesticides are known to inhibit the ChE activity in *D. magna* (Barata et al., 2004). Though, in the present study ChE levels were significantly higher for the population maintained in CBZ (Dph\_CBZ) than for those in clean medium (Dph\_Clean) (two-way ANOVA,  $F_{1,39} = 11.737$ ,  $p < 0.05$ ) (Table 4 SD), at all generations (Fig. 4a). An increase in the ChE activity in *D. magna* exposed to low concentrations of cadmium and the carbamate propoxur has been already reported (Jemec et al., 2007a; Printes and Callaghan, 2004). Increases in ChE levels at low doses might be explained by compensatory mechanisms after the disruption of homeostasis (Calabrese and Baldwin, 2003). Andrade et al. (2016) observed an increase in ChE activity after exposure of zebrafish (*Danio rerio*) embryos to CBZ, which the authors hypothesized to be related with apoptosis mechanisms, and has also been associated to the induction of gene expression related with apoptosis (Jiang et al., 2014). Although the exact mechanisms are not yet understood, indirect evidences that ChE participates in the regulation of apoptosis and cell proliferation have been theorised (Jiang and Zhang, 2008). A similar mechanism related with mediation of cell apoptosis could be playing a role in the ChE activity increase observed in the present study, however additional studies should be performed to confirm this hypothesis. Both exposure and generation showed some significant interaction, indicating that populations responded differently throughout the generations (two-way ANOVA,  $F_{3,39} = 3.393$ ,  $p < 0.05$ ) (Table 4 SD). However, throughout the successive generations (F0 to F12), there was an overall trend for the attenuation of this effect with ChE activity in F12 becoming similar between both populations (Fig. 4a). In the present study, variation in ChE activity was observed throughout generations, including in Dph\_Clean. Variation in ChE activity amongst individuals of the same species has been observed for *D. magna* in previous studies, including under control conditions. In the literature, under control conditions *D. magna* ChE levels vary from 0.034 to approximately 1.5 nmol/mg prot/min (Jemec et al.,

2007a; Qi et al., 2013), which is in accordance with the ChE activity reported in the present study for daphnids kept in clean medium (Dph\_Clean). On the other hand, the values reported here for ChE in Dph\_CBZ ranged from 0.74 (F0) to 0.60 nmol/mg prot/min (F12).

Although the main toxicity mechanism of carbamates is usually through ChE inhibition, exposure to carbamate pesticides has shown to trigger also other toxicity effects such as oxidative stress, by inducing generation of reactive oxygen species (ROS) (Milatovic et al., 2006). CAT is an antioxidant enzyme, which is responsible for breaking down hydrogen peroxide into water and molecular oxygen (Claiborne, 1985). CAT activity values reported in the literature for *D. magna* control groups range from 62.4  $\mu$ mol/mg prot/min (for daphnids with 22 d) to 250  $\mu$ mol/mg prot/min (for daphnids with 6 d) (Barata et al., 2005; Jemec et al., 2007b). In the present work, CAT levels determined for Dph\_Clean (between 25.92 and 39.21  $\mu$ mol/mg prot/min) were slightly lower comparing with the values reported in literature and were maintained at similar levels from F0 to F12 generations (Fig. 4b). However, several factors might cause this variability, including for instance the type of food provided (e.g. algae species), daphnids age and experimental conditions (e.g. temperature or photoperiod) (Rose et al., 2004). Comparing both populations, the exposure did not induce statistically differences for CAT levels (two-way ANOVA,  $F_{1,37} = 0.348$ ,  $p > 0.05$ ), though both factors interacted, meaning that populations responded differently throughout the generations due to exposure (two-way ANOVA,  $F_{3,37} = 10.271$ ,  $p < 0.001$ ) (Table 4 SD). Two patterns were observed when comparing both populations within the same generation: in F0 and F6, CAT activity increased upon exposure to CBZ, while in F3 and F12 a decrease was depicted (Fig. 4b). The stimulation in the initial response of F0 daphnids is possibly a typical response to the low CBZ concentration, to reduce oxidative stress (Vega and Pizarro, 2000). This was followed by a decrease in activity, followed by another increase and decrease, possibly meaning that the physiology of the organisms was working towards releasing ROS, whenever needed. CAT activity decreased in the herb fenugreek *Trigonella foenum-graecum* exposed to CBZ (Sangeetha, 2010), which was justified by a decrease in ROS, which was due to the previous activity of other antioxidative stress enzymes. Reduction in CAT activity was also observed in fish tissues (Palanikumar et al., 2014), rats (A dedara et al., 2013) and goats (Prashantkumar et al., 2013) after exposure to CBZ, being surely a dose-related response. Besides this, CAT activity in F3 daphnids exposed to CBZ was highly decreased which can be a

result of a simultaneous activation of another antioxidant defense mechanism, considering that in F6, this enzymatic activity showed a recovery (Sies, 1993; Wu et al., 2011). This may be the case of reduced glutathione (GSH), which is also involved in the removal of hydroperoxides (e.g. H<sub>2</sub>O<sub>2</sub>) (Sies, 1993; Wu et al., 2011), and occurs by its oxidation (mediated by H<sub>2</sub>O<sub>2</sub>) into glutathione disulfide (GSSG), therefore reducing the amount of substrate available to induce CAT (Wu et al., 2011).

Regarding GST activity, the reported values in the literature for control groups of *D. magna* varied from 42 nmol/mg prot/min (*D. magna* with 7 d), to 70 nmol/mg prot/min (*D. magna* with 21 d) and reached 235.2 nmol/mg prot/min (age not reported) (Borgeraas and Hessen, 2002; Chen et al., 2005; Domingues et al., 2015). Levels for detoxification looking at GST activity were different between both populations (*Dph\_Clean* and *Dph\_CBZ*) (two-way ANOVA,  $F_{1,33} = 4.557$ ,  $p < 0.05$ ) (Fig. 4c and Table 4 SD). Both factors, generation and exposure, interacted, indicating that populations responded differently throughout generations (two-way ANOVA,  $F_{3,33} = 2.951$ ,  $p < 0.05$ ) (Table 4 SD). This enzyme plays an important role in cellular detoxification processes of several chemicals and defense against peroxidative products of DNA (Henson et al., 2001). Pesticides can promote the consumption of glutathione in exposed organisms through a GST-catalyzed reaction in detoxification processes, and therefore GST induction aims to protect the organism (Ezemonye and Tongo, 2010; Timur et al., 2002). A multigenerational experiment with *D. magna* exposed to microcystins showed that upon parental exposure for 7 d, a higher GST activity in their offspring was observed when compared to those from controls (Ortiz-Rodriguez et al., 2012).

In the case of LPO, both populations (*Dph\_Clean* and *Dph\_CBZ*) showed differences (two-way ANOVA,  $F_{1,39} = 12.957$ ,  $p < 0.001$ ) and there was an interaction between generations and exposure, meaning that, throughout the generations, both populations reacted differently for this biomarker (two-way ANOVA,  $F_{3,39} = 6.612$ ,  $p < 0.001$ ) (Fig. 4d and Table 4 SD). In the F0 generation, a decrease in LPO in offspring from *Dph\_CBZ* was observed, comparing with *Dph\_Clean*. Vernouillet et al. (2010) observed a similar decrease in lipid peroxidation when exposed the crustacean *Thamnocephalus platyurus* to the pharmaceutical carbamazepine and suggested that carbamazepine might have preventing fatty acid oxidation in the membranes, by acting as a radical scavenger or by directly downregulate the cytosolic phospholipase A<sub>2</sub> activity. However, in the F6 generation differences in LPO between *Dph\_Clean* and

Dph\_CBZ were attenuated (Fig. 4d). In the last generation tested (F12), there was a slightly increase in LPO for neonates of Dph\_CBZ comparing with Dph\_Clean. This seems to indicate an imbalance in organisms redox equilibrium towards a situation of oxidative stress as previously described in several organisms (including the european eel and collembola) when exposed to harbor water, carbamazepine, fluoxetine and nanoparticle fullerene C60 (Ahmad et al., 2004; Oliveira et al., 2015; Zhu et al., 2006).

In the study of Palanikumar et al. (2014), the milkfish *Chanos chanos* was exposed to CBZ and chlorpyrifos and a relationship between DNA damage and the fluctuation in antioxidant enzymes responses might exist. In addition, an increase in DNA damage in *D. magna* was already reported from generation F0 to F12 under a similar approach, where daphnids were exposed to CBZ in multigenerational experiment as well (Silva et al., 2017), yet such straight relationship between DNA damage with antioxidant enzymes could not be established.




Figure 4. Biomarkers activities in *Daphnia magna* exposed (Dph\_CBZ, black dots) and non-exposed (Dph\_Clean, white dots) to carbendazim throughout generations: a) Cholinesterase (ChE) activity b) Catalase (CAT) activity c) Glutathione S-transferase (GST) activity and d) Lipid peroxidation (LPO) rate (n=5). Data are expressed as mean values and standard error.

When under stress and in order to survive, organisms undergo numerous alterations at a low level of biological organization. These alterations include metabolic changes that may end up affecting their energy-reserve fraction and energy consumption (Jeon et al., 2013; Vandenbrouck et al., 2009). Considering the multigenerational effects observed in several life traits, some energy-related parameters were measured in different generations as an attempt to detect and track possible CBZ induced changes in resource allocation. Carbohydrates, which are considered the first energy fraction to be consumed, presented an almost similar pattern between both isoclinal populations Dph\_Clean and Dph\_CBZ (two-way ANOVA,  $F_{1,23} = 0.167$ ,  $p > 0.05$ ) (Fig. 5a and Table 5 SD), with no interaction between both factors, generations and exposure (two-way ANOVA,  $F_{3,23} = 2.202$ ,  $p > 0.05$ ) (Table 5 SD). Carbohydrates contents in Dph\_Clean neonates are within the range of those found in literature, which vary highly between 199 to 2054 mj/organism (for neonates <24h exposed for 48h/96h) (De Coen and Janssen, 1997). This variability may be justified by the different food sources and quantities provided to organisms, along with changes in temperature and dissolved oxygen, among others (Bergman Filho et al., 2011).

For the lipids reserves, a similar pattern was observed for both populations Dph\_Clean and Dph\_CBZ (two-way ANOVA,  $F_{1,22} = 0.113$ ,  $p > 0.05$ ) and no interaction between both factors (generations and exposure) was found (two-way ANOVA,  $F_{3,22} = 1.334$ ,  $p > 0.05$ ) (Fig. 5b and Table 5 SD). Lipidic levels obtained in this work for daphnids in clean medium (Dph\_Clean) are within the same range reported in literature for the Daphnia species (approx. 1000 mj/organism) (Bergman Filho et al., 2011).

A similar trend was obtained for proteins between Dph\_Clean and Dph\_CBZ populations (two-way ANOVA,  $F_{1,23} = 0.00115$ ,  $p > 0.05$ ) (Fig. 5c and Table 5 SD) with no interaction between both factors, generations and exposure (two-way ANOVA,  $F_{3,23} = 3.129$ ,  $p > 0.05$ ) (Table 5 SD). In literature, protein values for *D. magna* in control situation range from 1694 mj/organism to 5518 mj/organism (exposed for 48h) (De Coen and Janssen, 1997). The Dph\_Clean population presented protein values ranging from 2000 mj/org to 3000 mj/org, which were maintained throughout generations, as well as those for the population exposed to CBZ. This was somehow contrary to the expected and reported by several authors for several species, where an increase in protein content was observed for *D. magna*, *Danio rerio* and *E. albidus* exposed to lindane, effluents and also CBZ, respectively (De Coen and Janssen, 2003a; Novais and Amorim, 2013; Smolders et al., 2003). In addition, Kim et al. (2014) studied the effects



of tetracycline in four generations of *D. magna*; depletions in proteins, carbohydrates and lipid reserves were found in consequence of the stress caused by tetracycline. However, throughout the generations, these reductions were recovered (comparing with the control group), suggesting some adaptation (Kim et al., 2014).




Figure 5

Figure 5. Energy-related parameters in *Daphnia magna* exposed (Dph\_CBZ, black dots) and non-exposed (Dph\_Clean, white dots) to carbendazim throughout generations: a) Carbohydrates b) Lipids and c) Protein contents (n=3). Data are expressed as mean values and standard error.

This lack of differences in lipidic, protein and carbohydrates contents (Table 5 SD), derive similar Energy available (Ea) or Energy consumed (Ec). While for Ea no interaction between generations and exposure was obtained, the same was not true for Ec (Table 6 SD; Fig. 6). Although no differences in patterns were observed for energy reserves within generations and exposure to CBZ, a trade-off seemed to have occurred, while looking at  $r$  and organism's longevity. Several examples found in literature reported that *D. magna* showed an ability to switch its life history responses while exposed to stressors (Minguez et al., 2015), and in the present study reproduction was more favoured than survival. Different patterns may be justified by the chemical nature, but also the level of concentration used should be considered. In the present study, the concentration used is a NOEC for reproduction in a 21d exposure. In addition, in the present study, some differences were observed in terms of oxidative stress related biomarkers, showing that in some cases enzymatic activities were activated to achieve homeostasis, while decreasing their activity afterwards, or being compensated by other enzymatic processes within the same molecular pathway. These physiological processes might have helped the organism to maintain a healthy status, and few effects depicted under a population level.




Figure 6.

Figure 6. Energy-related parameters on individuals of *Daphnia magna* exposed (Dph\_CBZ, black dots) and non-exposed (Dph\_Clean, white dots) to carbendazim throughout generations. Data are expressed as mean values and standard error.

The variability observed in several parameters recorded for *D. magna* under control exposure throughout the generations is a pattern that should be further discussed

and not ignored. Besides the enlightenments referred previously (e.g. algae quality), Traudt et al. (2016) also demonstrated that exposing daphnids to cadmium using a narrow age window of less than 24h reflected an EC50 value 10 times different between them (when comparing daphnids with 0-4h and daphnids with 20-24h). In the present work, some of the observed effects can be attributed partly to inter-generational environmental and natural variation. In fact, inter-generational effects due to, for instance, changes in parental food environment have already been demonstrated (Plaistow et al., 2005). In the same study, evidences that life history traits are the result of interactions between past but also present environments and that these traits vary from one generation to the subsequent one were demonstrated. In addition, trade-offs between endpoints were found (as observed in the present work) and changed depending on the high (trade-off between fecundity and survival) or low food supply (trade-off between age and size) (Plaistow et al., 2005). Variability in responses has been discussed since the 90's for daphnids, where apart from the genotype and environmental variables and their interaction, an additional variable, the 'residual variability component' is present (Soares et al., 1992). This former variable may include for instance measurement/pure errors and others unexplained/natural (Falconer and Mackay, 1996).

#### 4. Conclusions

The multigenerational exposure of *D. magna* to a NOEC equivalent concentration of CBZ induced low effects but provided useful information to understand how populations react to long-term exposure to chemicals. One of the first highlights derived from the present study is that the continuous exposure to CBZ did not induce changes in the intrinsic rate of natural increase ( $r$ ) but deeply affected their longevity, with a notorious decrease in the lifespan found in daphnids exposed after 12 generations to CBZ. Considering that energy related-parameters showed no significant differences between both populations, a trade-off in energy allocation possibly occurred, with more energy being allocated in the reproduction of daphnids, diminishing energy available for survival in a long term. Energy was also partly allocated to detoxification, when looking at biomarkers' patterns. Although ChE, GST and LPO showed differences between clean and exposed isoclonal populations, the identification of a clear detoxification mechanism could not be depict.

Although *D. magna* is a parthenogenic organism generally expected to show low variability in responses, several life-trait parameters within generations in control exposures presented some inconsistency.

## Acknowledgments

This work was supported by the project RePulse- Responses of *Daphnia magna* exposed to chemical pulses and mixtures throughout generations (FCOMP-01-0124-FEDER-019321; Refł FCT PTDC/AAC-AMB/117178/2010), by funding FEDER through COMPETE- Programa Operacional Factores de Competitividade, and by the Portuguese Science Foundation (FCT) through CESAM: UID/AMB/50017/2013. Ana Rita R. Silva was funded by the project `MARPRO- Conservation of marine protected species in Mainland Portugal\_ through the doctoral fellowship (BD/UI88/5534/2011). The Special Research Fund (BOF) of Ghent University supported the doctoral fellowship of C@ia S. A. Santos (B/13833/01 - BOF13/DOC/034). Diogo Cardoso was funded by the doctoral grant (PD/BD/52569/2014) and Andreia Cruz was funded by an individual post-doctoral grant (BPD/UI88/2886/2013), within the project `Sustainable Use of Marine Resources\_ - MARES (CENTRO-07-ST24-FEDER-002033), financed by QREN, Mais Centro- Programa Operacional Regional do Centro e Uni<sup>2</sup>o Europeia/ Fundo Europeu de Desenvolvimento Regional. The authors would like to thank the laboratorial support given by Dr. Abel Ferreira.

Data availability` Data and calculation tools are available from the corresponding author ([ritas@ua.pt](mailto:ritas@ua.pt)).

## References

- A dedara, I. A., Vaithinathan, S., Jubendradass, R., Mathur, P. P., Farombi, E. O., 2013. Kolaviron prevents carbendazim-induced steroidogenic dysfunction and apoptosis in testes of rats. *Environ Toxicol Pharmacol.* 35, 444-53.
- Ahmad, I., Pacheco, M., Santos, M. A., 2004. Enzymatic and nonenzymatic antioxidants as an adaptation to phagocyte-induced damage in *Anguilla anguilla* L. following in situ harbor water exposure. *Ecotoxicology and Environmental Safety.* 57, 290-302.
- Andrade, T. S., Henriques, J. F., Almeida, A. R., Machado, A. L., Koba, O., Giang, P. T., Soares, A. M., Domingues, I., 2016. Carbendazim exposure induces developmental, biochemical and behavioural disturbance in zebrafish embryos. *Aquatic Toxicology.* 170, 390-9.
- ASTM, 1980. Standard practice for conducting acute toxicity tests with fishes, macroinvertebrates and amphibians. Report E-729-80. American Standards for Testing and Materials. Philadelphia, P.A.
- Barata, C., Solayan, A., Porte, C., 2004. Role of B-esterases in assessing toxicity of organophosphorus (chlorpyrifos, malathion) and carbamate (carbofuran) pesticides to *Daphnia magna*. *Aquat Toxicol.* 66, 125-39.
- Barata, C., Varo, I., Navarro, J. C., Arun, S., Porte, C., 2005. Antioxidant enzyme activities and lipid peroxidation in the freshwater cladoceran *Daphnia magna* exposed to redox cycling compounds. *Comp Biochem Physiol C Toxicol Pharmacol.* 140, 175-86.

- 魚れ Bergman Filho, T. U., Soares, A. M., Loureiro, S., 2011. Energy budget in *Daphnia magna* exposed to natural stressors. *Environ Sci Pollut Res Int.* 18, 655-62.  
 魚子 Bird, R., Draper, H., 1984. Comparative studies on different methods of malonaldehyde determination. *Methods in Enzymology* 105, 299-305.  
 魚わ Borgeraas, J., Hessen, D. O., 2002. Variations of antioxidant enzymes in *Daphnia* species and populations as related to ambient UV exposure. *Hydrobiologia.* 47, 15-30.  
 魚ゆ Brusch, J. M., Smith, P. N., 2009. Development of resistance to cyfluthrin and naphthalene among *Daphnia magna*. *Ecotoxicology.* 18, 600-609.  
 魚魚 Brown, P. J., Long, S. M., Spurgeon, D. J., Svendsen, C., Hankard, P. K., 2004. Toxicological and biochemical responses of the earthworm *Lumbricus rubellus* to pyrene, a non-carcinogenic polycyclic aromatic hydrocarbon. *Chemosphere.* 57, 1675-81.  
 魚在 Chen, Y., Huang, J., Xing, L., Liu, H., Giesy, J. P., Yu, H., Zhang, X., 2013. Effects of multigenerational exposures of *D. magna* to environmentally relevant concentrations of pentachlorophenol. *Environmental Science and Pollution Research.* 21, 234-243.  
 魚ん Buhl, K. J., Hamilton, S. J., Schmulbach, J. C., 1993. Chronic toxicity of the bromoxynil formulation Buctril to *Daphnia magna* exposed continuously and intermittently. *Arch. Environ. Contam. Toxicol.* 25, 152-159.  
 魚ん Calabrese, E. J., Baldwin, L. A., 2003. Inorganics and hormesis. *Critical Reviews in Toxicology.* 33, 215-304.  
 魚ん Chen, W., Song, L., Ou, D., Gan, N., 2005. Chronic toxicity and responses of several important enzymes in *Daphnia magna* on exposure to sublethal microcystin-LR. *Environ Toxicol.* 20, 323-30.  
 魚ん Chen, Y., Huang, J., Xing, L., Liu, H., Giesy, J. P., Yu, H., Zhang, X., 2013. Effects of multigenerational exposures of *D. magna* to environmentally relevant concentrations of pentachlorophenol. *Environmental Science and Pollution Research.* 21, 234-243.  
 魚れ Claiborne, A., 1985. Catalase activity. In: Greenwald, R.A. (Ed.), *Handbook of Methods for Oxygen Research.* CRC Press, Boca Raton, FL.  
 魚ろ Clubbs, R. L., Brooks, B. W., 2007. *Daphnia magna* responses to a vertebrate estrogen receptor agonist and an antagonist: a multigenerational study. *Ecotoxicol Environ Saf.* 67, 385-98.  
 魚わ Cuppen, J. G. M., Van Den Brink, P. J., Camps, E., Uil, K. F., Brock, T. C. M., 2000. Impact of the fungicide carbendazim in freshwater microcosms. I. Water quality, breakdown of particulate organic matter and responses of macroinvertebrates. *Aquatic Toxicology.* 48, 233-250.  
 魚を Davidse, L. C., 1973. Antimitotic activity of methyl benzimidazol-2-yl carbamate (MBC) in *Aspergillus nidulans*. *Pesticide Biochemistry and Physiology.* 3, 317-325.  
 魚ん De Coen, W., Janssen, C. R., 2003a. The missing biomarker link: Relationships between effects on the cellular energy allocation biomarker of toxicant-stressed *Daphnia magna* and corresponding population characteristics. *Environ Toxicol Chem.* 22, 1632-1641.  
 魚ん De Coen, W. M., Janssen, C., 2003b. A multivariate biomarker-based model predicting population-level responses of *Daphnia magna*. *Environmental Toxicology and Chemistry.* 22, 2195-2201.  
 魚ん De Coen, W. M., Janssen, C. R., 1997. The use of biomarkers in *Daphnia magna* toxicity testing. IV. Cellular Energy Allocation: a new methodology to assess the energy budget of toxicant-stressed *Daphnia* populations. *Journal of Aquatic Ecosystem Stress and Recovery.* 6, 43-55.  
 魚ん Domingues, I., Santos, C. S., Ferreira, N. G., Machado, L., Oliveira, R., Ferreira, A., Lopes, I., Loureiro, S., Soares, A. M., 2015. Suitability of enzymatic markers to assess the environmental condition of natural populations of *Gambusia affinis* and *Daphnia magna*-a case study. *Environ Monit Assess.* 187, 208.  
 魚ん Ecobichon, D. J., 2001. Pesticide use in developing countries. *Toxicology.* 160, 27-33.  
 魚ん Ellman, G. L., Courtney, K. D., Andres, V., Featherstone, R. M., 1961. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochemical Pharmacology* 7, 88-90.  
 魚ん EN ISO 6341, Water Quality-Determination of the Inhibition of the Mobility of *Daphnia magna* Straus (Cladocera, Crustacea)-Acute Toxicity Test in ISO 6341, Cor.1(98). International Organisation for Standardisation, Geneva, Switzerland., 1996.

EU Pesticide Database, 2016. EU Pesticide Database, European Commission. <http://ec.europa.eu/food/plant/pesticides/eu-pesticides-database/public/?event=activeSubstanceDetail&language=EN&selectedID=1080> accessed in September 2017.

Ezemonye, L., Tongo, I., 2010. Sublethal effects of endosulfan and diazinon pesticides on glutathione-S-transferase (GST) in various tissues of adult amphibians (*Bufo regularis*). *Chemosphere*. 81, 214-7.

Falconer, D. S., Mackay, T. F. C., 1996. Introduction to quantitative genetics. Fourth edition. Longman Science and Technology, Harlow, UK.

Ferreira, N. G., Morgado, R., Santos, M. J., Soares, A. M. V. M., Loureiro, S., 2015. Biomarkers and energy reserves in the isopod *Porcellionides pruinosus*: the effects of long-term exposure to dimethoate. *Sci Total Environ*. 502, 91-102.

Ferreira, N. G. C., Santos, M. J. G., Domingues, I., Calhã, C. F., Monteiro, M., Amorim, M. J. B., Soares, A. M. V. M., Loureiro, S., 2010. Basal levels of enzymatic biomarkers and energy reserves in *Porcellionides pruinosus*. *Soil Biology and Biochemistry*. 42, 2128-2136.

Frasco, M. F., Guilhermino, L., 2002. Effects of dimethoate and beta-naphthoflavone on selected biomarkers of *Poecilia reticulata*. *Fish Physiology and Biochemistry*. 26, 149-156.

Guilhermino, L., Lopes, M. C., Carvalho, A. P., Soares, A. M. V. M., 1996. Inhibition of acetylcholinesterase activity as effect criterion in acute tests with juvenile *Daphnia magna*. *Chemosphere*. 32, 727-38.

Habig, W. H., Pabst, M. J., Jakoby, W. B., 1974. Glutathione S-transferases: a first enzymatic step on mercapturic acid formation. *Journal of Biological Chemistry*. 249, 7130-7139.

Hebert, P. D. N., Ward, R. D., 1972. Inheritance during parthenogenesis in *Daphnia magna*. *Genetics*. 71, 639-642.

Henson, K. L., Stauffer, G., Gallagher, E. P., 2001. Induction of glutathione S-transferase activity and protein expression in brown bullhead (*Ameiurus nebulosus*) liver by ethoxyquin. *Toxicol Sci* 62, 54-60.

Hullett, C. R., Levine, T. R., 2003. The overestimation of effect sizes from F values in meta-analysis: the cause of a solution. *Communication Monographs*. 70, 52-67.

Hyne, R. V., Maher, W. A., 2003. Invertebrate biomarkers: links to toxicosis that predict population decline. *Ecotoxicology and Environmental Safety*. 54, 366-374.

Jacobasch, C., Volker, C., Giebner, S., Volker, J., Alsenz, H., Potouridis, T., Heidenreich, H., Kayser, G., Oehlmann, J., Oetken, M., 2014. Long-term effects of nanoscaled titanium dioxide on the cladoceran *Daphnia magna* over six generations. *Environ Pollut*. 186, 180-6.

Jemec, A., Drobne, D., Tisler, T., Trebse, P., Ros, M., Sepcic, K., 2007a. The applicability of acetylcholinesterase and glutathione S-transferase in *Daphnia magna* toxicity test. *Comp Biochem Physiol C Toxicol Pharmacol*. 144, 303-9.

Jemec, A., Tisler, T., Drobne, D., Sepcic, K., Fournier, D., Trebse, P., 2007b. Comparative toxicity of imidacloprid, of its commercial liquid formulation and of diazinon to a non-target arthropod, the microcrustacean *Daphnia magna*. *Chemosphere*. 68, 1408-18.

Jeon, J., Kretschmann, A., Escher, B. I., Hollender, J., 2013. Characterization of acetylcholinesterase inhibition and energy allocation in *Daphnia magna* exposed to carbaryl. *Ecotoxicol Environ Saf*. 98, 28-35.

Jiang, H., Zhang, X. J., 2008. Acetylcholinesterase and apoptosis. A novel perspective for an old enzyme. *FEBS J*. 275, 612-7.

Jiang, J., Wu, S., Wu, C., An, X., Cai, L., Zhao, X., 2014. Embryonic exposure to carbendazim induces the transcription of genes related to apoptosis, immunotoxicity and endocrine disruption in zebrafish (*Danio rerio*). *Fish Shellfish Immunol*. 41, 493-500.

Kim, H. Y., Lee, M. J., Yu, S. H., Kim, S. D., 2012. The individual and population effects of tetracycline on *Daphnia magna* in multigenerational exposure. *Ecotoxicology*. 21, 993-1002.

Kim, H. Y., Yu, S., Jeong, T. Y., Kim, S. D., 2014. Relationship between trans-generational effects of tetracycline on *Daphnia magna* at the physiological and whole organism level. *Environ Pollut.* 191, 111-8.

Kimberly, D. A., Salice, C. J., 2014. If you could turn back time: understanding transgenerational latent effects of developmental exposure to contaminants. *Environ Pollut.* 184, 419-25.

Liess, M., Foit, K., Becker, A., Hassold, E., Dolciotti, I., Kattwinkel, M., Duquesne, S., 2013. Culmination of low-dose pesticide effects. *Environ Sci Technol.* 47, 8862-8.

Lokta, A. J., 1913. Vital statistics - a natural population norm. *Journal - Washington Academy of Sciences.* 3, 241-248, 289-293.

Loureiro, S., Svendsen, C., Ferreira, A. L. G., Pinheiro, C., Ribeiro, F., Soares, A. M. V. M., 2010. Toxicity of three binary mixtures to *Daphnia magna*: Comparing chemical modes of action and deviations from conceptual models. *Environmental Toxicology and Chemistry.* 29, 1716-1726.

Milatovic, D., Gupta, R. C., Aschner, M., 2006. Anticholinesterase toxicity and oxidative stress. *Scientific World Journal.* 6, 295-310.

Minguez, L., Ballandonne, C., Rakotomalala, C., Dubreule, C., Kientz-Bouchart, V., Halm-Lemeille, M. P., 2015. Transgenerational effects of two antidepressants (sertraline and venlafaxine) on *Daphnia magna* life history traits. *Environ Sci Technol.* 49, 1148-55.

Novais, S. C., Amorim, M. J., 2013. Changes in cellular energy allocation in *Enchytraeus albidus* when exposed to dimethoate, atrazine, and carbendazim. *Environ Toxicol Chem.* 32, 2800-7.

Novais, S. C., Amorim, M. J. B., 2015. Normal operating range (NOR) in *Enchytraeus albidus* - Transcriptional responses to control conditions. *Applied Soil Ecology.* 85, 1-10.

OECD, 2004. OECD guidelines for testing of chemicals. Guideline 202: *Daphnia sp.*, Acute immobilisation test, adopted April 2004.

Ohkawa, H., Ohishi, N., Yagi, K., 1979. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Analytical Biochemistry.* 95, 351-358.

Oliveira, M., Cardoso, D. N., Soares, A. M., Loureiro, S., 2015. Effects of short-term exposure to fluoxetine and carbamazepine to the collembolan *Folsomia candida*. *Chemosphere.* 120, 86-91.

Ortiz-Rodriguez, R., Dao, T. S., Wiegand, C., 2012. Transgenerational effects of microcystin-LR on *Daphnia magna*. *J Exp Biol.* 215, 2795-805.

Palanikumar, L., Kumaraguru, A. K., Ramakritinan, C. M., Anand, M., 2014. Toxicity, biochemical and clastogenic response of chlorpyrifos and carbendazim in milkfish *Chanos chanos*. *International Journal of Environmental Science and Technology.* 11, 765-774.

Palma, G., Sanchez, A., Olave, Y., Encina, F., Palma, R., Barra, R., 2004. Pesticide levels in surface waters in an agricultural-forestry basin in Southern Chile. *Chemosphere.* 57, 763-770.

Pestana, J. L., Loureiro, S., Baird, D. J., Soares, A. M., 2010. Pesticide exposure and inducible antipredator responses in the zooplankton grazer, *Daphnia magna* Straus. *Chemosphere.* 78, 241-8.

Plaistow, S. J., Lapsley, C. T., Benton, T. G., 2005. Context-dependent intergenerational effects: the interaction between past and present environments and its effect on population dynamics. *The American Naturalist.* 167, 206-215.

Prashantkumar, W., Rampal, S., Saini, S. P. S., Prakash, N., Lokesh, L. V., Ahsan-Ul-Haq, S., 2013. Sub - chronic exposure of carbendazim (bavistin 50% wp) alters antioxidant status in male goats (*Capra hircus*). *Adv. Pharmacol. Toxicol.* 14, 59-66.

Printes, L. B., Callaghan, A., 2004. A comparative study on the relationship between acetylcholinesterase activity and acute toxicity in *Daphnia magna* exposed to anticholinesterase insecticides. *Environ. Toxicol. Chem.* 23, 1241-1247.

Qi, S., Wang, C., Chen, X., Qin, Z., Li, X., Wang, C., 2013. Toxicity assessments with *Daphnia magna* of Guadipyr, a new neonicotinoid insecticide and studies of its effect on



acetylcholinesterase (AChE), glutathione S-transferase (GST), catalase (CAT) and chitinase activities. *Ecotoxicol Environ Saf.* 98, 339-44.

Rose, R. M., Warne, M. S., Lim, R. P., 2004. Sensitivity of offspring to chronic 3,4-dichloroaniline exposure varies with maternal exposure. *Ecotoxicol Environ Saf.* 58, 405-12.

Sánchez, M., Andreu-Moliner, E., Ferrando, M. D., 2004. Laboratory investigation into the development of resistance of *Daphnia magna* to the herbicide molinate. *Ecotoxicology and Environmental Safety.* 59, 316-323.

Sangeetha, R., 2010. Activity of Superoxide Dismutase and Catalase in Fenugreek (*Trigonella foenum-graecum*) in Response to Carbendazim. *Indian Journal of Pharmaceutical Sciences.* 72, 112-114.

Sies, H., 1993. Strategies of antioxidant defense. *EJB Reviews.* 215, 213-219.

Silva, A. R. R., Cardoso, D. N., Cruz, A., Lourenco, J., Mendo, S., Soares, A. M. V. M., Loureiro, S., 2015. Ecotoxicity and genotoxicity of a binary combination of triclosan and carbendazim to *Daphnia magna*. *Ecotoxicol Environ Safety.* 115, 279-290.

Silva, A. R. R., Cardoso, D. N., Cruz, A., Pestana, J. L., Mendo, S., Soares, A. M. V. M., Loureiro, S., 2017. Multigenerational effects of carbendazim in *Daphnia magna*. *Environ Toxicol Chem.* 1-12.

Smolders, R., De Boeck, G., Blust, R., 2003. Changes in cellular energy budget as a measure of whole effluent toxicity in zebrafish (*Danio rerio*). *Environ Toxicol Chem* 22, 890-899.

Soares, A. M., Baird, D. J., Calow, P., 1992. Interclonal variation in the performance of *Daphnia magna* Straus in chronic bioassays. *Environmental Toxicology and Chemistry.* 11, 1477-148.

SPSS 20.0.0, SPSS (Statistical Package for the Social Sciences). SPSS for Windows. 2011.

Systat Software Inc., 2008. SigmaPlot for Windows (version 11.0). San Jose, California, USA.

Taberner, A., Castanera, P., Silvestre, E., Dopazo, J., 1993. Estimation of the intrinsic rate of natural increase and its error by both algebraic and resampling approaches. *CABIOS.* 9, 535-540.

Tanaka, Y., Nakanishi, J., 2002. Chronic effects of p-nonylphenol on survival and reproduction of *Daphnia galeata*: multigenerational life table experiment. *Environ Toxicol.* 17, 487-92.

Timur, S., Onal, S., Karabay, U. N., Sayim, F., Zihniogul, F., 2002. In vivo effects of malathion on glutathione-S-transferase and acetylcholinesterase activities in various tissues of neonatal rats. *Turkish Journal of Zoology.* 27, 247-252.

Vandenbrouck, T., Soetaert, A., van der Ven, K., Blust, R., De Coen, W., 2009. Nickel and binary metal mixture responses in *Daphnia magna*: molecular fingerprints and (sub)organismal effects. *Aquat Toxicol.* 92, 18-29.

Vega, M. P., Pizarro, R. A., 2000. Oxidative stress and defence mechanisms of the freshwater cladoceran *Daphnia longispina* exposed to UV radiation. *Journal of Photochemistry and Photobiology B: Biology.* 54, 121-125.

Vernouillet, G., Eullaffroy, P., Lajeunesse, A., Blaise, C., Gagne, F., Juneau, P., 2010. Toxic effects and bioaccumulation of carbamazepine evaluated by biomarkers measured in organisms of different trophic levels. *Chemosphere.* 80, 1062-8.

WHO, 1993. Environmental Health Criteria 149. International Programme on Chemical Safety, Geneva. Available from: [http://www.inchem.org/documents/hsg/hsg/hsg82\\_e.htm#SectionNumber:1.4](http://www.inchem.org/documents/hsg/hsg/hsg82_e.htm#SectionNumber:1.4), accessed in September 2017.

Wu, H. Y., Xu, H. X., Hong, Y. B., Zhang, J. F., Wu, J. C., 2011. The use of biomarkers in the antioxidant responses of *Daphnia magna* to the acute and chronic exposure to no. 20 diesel oil and 2,4-dichlorophenol. *Chemical Speciation and Bioavailability.* 23, 80-87.

Zalizniak, L., Nuggeoda, D., 2006. Effect of sublethal concentrations of chlorpyrifos on three successive generations of *Daphnia carinata*. *Ecotoxicol Environ Saf.* 64, 207-14.

Zhu, S., Oberdorster, E., Haasch, M. L., 2006. Toxicity of an engineered nanoparticle (fullerene, C60) in two aquatic species, *Daphnia* and fathead minnow. *Mar Environ Res.* 62 Suppl, S5-9.

h<sub>2</sub>

h<sub>2</sub>