

# ORCA - Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository:https://orca.cardiff.ac.uk/id/eprint/119703/

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Karatzias, Thanos, Murphy, Philip, Cloitre, Marylene, Bisson, Jonathan , Shevlin, Mark, Hyland, Philip, Maercker, Andreas, Ben-Ezra, Menachem, Coventry, Peter, Mason-Roberts, Susan, Bradley, Aoife and Hutton, Paul 2019. Psychological interventions for ICD-11 complex PTSD symptoms: systematic review and meta-analysis. Psychological Medicine 49 (11) , pp. 1761-1775. 10.1017/S0033291719000436

Publishers page: https://doi.org/10.1017/S0033291719000436

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See http://orca.cf.ac.uk/policies.html for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



# Title

Psychological interventions for ICD-11 Complex PTSD symptoms: Systematic review and meta-analysis

# <u>Running Head</u>

Psychological therapies for CPTSD

# <u>Authors</u>

Thanos Karatzias<sup>1,2</sup> Philip Murphy<sup>1</sup> Marylene Cloitre<sup>3,4</sup> Jonathan Bisson<sup>5</sup> Neil Roberts<sup>5,6</sup> Mark Shevlin<sup>7</sup> Philip Hyland<sup>8</sup> Andreas Maercker<sup>9</sup> Menachem Ben-Ezra<sup>10</sup> Peter Coventry<sup>11</sup> Susan Mason-Roberts<sup>1</sup> Aoife Bradley<sup>1</sup> Paul Hutton<sup>1</sup>

# **Affiliations**

<sup>1</sup>Edinburgh Napier University, School of Health & Social Care, Edinburgh, UK
 <sup>2</sup>NHS Lothian, Rivers Centre for Traumatic Stress, Edinburgh, UK
 <sup>3</sup>Stanford University, Department of Psychiatry and Behavioral Sciences, USA
 <sup>4</sup>National Center for PTSD, Veterans Affairs Palo Alto Health Care System, Palo Alto, CA, USA
 <sup>5</sup>Cardiff University, School of Medicine, Cardiff, UK
 <sup>6</sup>Psychology and Counselling Directorate, Cardiff and Vale University Health Board, Cardiff, UK
 <sup>6</sup>Psychology and Counselling Directorate, Cardiff and Vale University Health Board, Cardiff, UK
 <sup>6</sup>Psychology of Psychology, Derry, UK
 <sup>8</sup>National College of Ireland, School of Business, Dublin, Ireland
 <sup>9</sup>University of Zurich, Department of Psychology, Psychopathology and Clinical Interventions, Zurich, Switzerland
 <sup>10</sup>School of Social Work, Ariel University, Ariel, Israel
 <sup>11</sup>Department of Health Sciences and Centre for Reviews and Dissemination, University of York

# <sup>1</sup><u>Address for Correspondence</u>:

Thanos Karatzias, Edinburgh Napier University, Sighthill Campus, Sighthill Court, Edinburgh EH11 4BN, Scotland UK, Tel. (+44) (0) 131 455 5345, Email.<u>t.karatzias@napier.ac.uk</u>

#### Abstract

Background: The 11<sup>th</sup> revision to the WHO International Classification of Diseases (ICD-11) identified Complex Posttraumatic Stress Disorder (CPTSD) as a new condition. There is a pressing need to identify effective CPTSD interventions. Methods: We conducted a systematic review and meta-analysis of Randomised Controlled Trials (RCTs) of psychological interventions for Posttraumatic Stress Disorder (PTSD), where participants were likely to have clinically significant baseline levels of one or more CPTSD symptom clusters (affect dysregulation, negative self-concept and/or disturbed relationships). We searched MEDLINE, PsycINFO, EMBASE and PILOTS databases (January 2018), and examined study and outcome quality. Results: Fifty-one RCTs met inclusion criteria. Cognitive Behavioural Therapy (CBT), Exposure alone (EA), and Eye Movement Desensitization and Reprocessing (EMDR) were superior to usual care for PTSD symptoms, with effects ranging from g = -0.90 (CBT; k=27, 95% CI -1.11, -0.68; moderate quality) to g = -1.26 (EMDR; k=4, 95% CI -2.01, -0.51; low quality). CBT and EA each had moderatelarge or large effects on negative self-concept, but only 1 trial of EMDR reported this outcome. CBT, EA and EMDR each had moderate or moderate-large effects on disturbed relationships. Few RCTs reported affect dysregulation data. The benefits of all interventions were smaller when compared to non-specific interventions (e.g., befriending). Multivariate meta-regression suggested childhood-onset trauma was associated with a poorer outcome. **Conclusions**: The development of effective interventions for CPTSD can build upon the success of PTSD interventions. Further research should assess the benefits of flexibility in intervention selection, sequencing and delivery, based on clinical need and patient preferences.

**Keywords**: CPTSD, psychological therapies, childhood trauma, systematic review, metaanalysis, randomised controlled trials

2

#### Introduction

The 11<sup>th</sup> revision to the World Health Organization's International Classification of Diseases (ICD-11) (WHO, 2018) includes two distinct sibling conditions, Posttraumatic Stress Disorder (PTSD) (code 6B40) and Complex PTSD (CPTSD) (code 6B41), under a general parent category of 'Disorders specifically associated with stress'. PTSD is comprised of three symptom clusters including (1) re-experiencing of the trauma in the here and now, (2) avoidance of traumatic reminders, and (3) a persistent sense of current threat that is manifested by exaggerated startle and hypervigilance. ICD-11 CPTSD includes the three PTSD clusters and three additional clusters that reflect 'disturbances in self-organization' (DSO); (1) affect dysregulation, (2) negative self-concept, and (3) disturbances in relationships (Maercker et al., 2013). These disturbances are proposed to be typically associated with sustained, repeated, or multiple forms of traumatic exposure (e.g., genocide campaigns, childhood sexual abuse, child soldiering, severe domestic violence, torture, or slavery) (Karatzias et al., 2017), reflecting loss of emotional, psychological, and social resources under conditions of prolonged adversity (Cloitre et al., 2013).

The qualitative distinction between PTSD and CPTSD symptomatology has been supported in different trauma samples (see Brewin et al., 2017) including those experiencing interpersonal violence (Cloitre et al., 2013), rape, domestic violence, traumatic bereavement (Elklit, Hyland, & Shevlin, 2014), survivors of institutional abuse such as that occurring within foster care and religious organizations (Knefel et al., 2015) and refugees (Hyland et al., 2018). The distinction between PTSD and CPTSD has also been confirmed in samples of young adults (Perkonigg et al., 2014) and children (Sachser, Keller, & Goldbeck, 2016). The second-order factorial structure of CPTSD in which the disorder is comprised of both PTSD and DSO has also been supported in previous research (e.g. Karatzias et al., 2016; Hyland et al., 2017a; Hyland et al., 2017b; Shevlin et al., 2017).

To date a number of meta-analyses and systematic reviews have investigated the effectiveness of PTSD treatments in general (Barrera et al, 2013; Bisson & Andrews, 2005, 2007; Bisson et al., 2007; Bisson et al., 2013; Callahan et al 2004; DeJong & Gorey, 1996; Ehring et al, 2014; Pelekis & Dahl, 2005; Roberts et al, 2015; Sloan et al, 2013; Taylor & Harvey, 2009; Taylor & Harvey 2010; Watts et al, 2013). Overall, previous meta-analyses have supported the efficacy of trauma-focused psychological treatments, such as Cognitive Behavioural Therapy (CBT) and Eye Movement Desensitization and Reprocessing (EMDR), for the treatment of DSM-IV PTSD, a condition of three clusters of symptoms including reexperience, avoidance of the traumatic reminders and hyperarousal. CBT and EMDR target patients' memories of their traumatic events and the personal meanings of the trauma and typically include repeated in vivo and/or imaginal exposure to the trauma, reappraisal of the meaning of the trauma and its consequences, or some combination of these techniques (e.g. Bisson et al., 2013). These approaches have been identified as efficacious for a range of PTSD survivors, including rape victims, survivors of childhood abuse, refugees, combat veterans, and victims of motor vehicle accidents (Foa et al. 2009), although most existing evidence on these interventions concerns single adult traumas (e.g. Bisson et al., 2013). There is disagreement whether trauma focused treatments are optimal for more complex traumatic presentations such as CPTSD. For complex traumatic presentations, a phase-based model, originally proposed by Herman (1992), has been suggested as the preferred treatment option (Cloitre et al., 2012).

Phased interventions address disturbances in self-organization and related problems in day to day functioning (e.g., improving safety, emotion regulation and social skills) first, while explicit exploration of the trauma (e.g., exposure) is subsequently introduced (Cloitre et al., 2012b). The rationale for this sequencing is two-fold; firstly to increase emotional, psychological and social resources to improve functioning in daily life and secondly, to use these resources to enhance the effectiveness of trauma-focused work. Whilst there is some support for this approach (e.g., Cloitre et al., 2010), it is uncertain if a stabilisation phase is necessary and it might lead to unhelpful delays in using more trauma-focused interventions (De Jongh et al., 2016). Another approach to managing complex traumatisation focuses on treating symptoms that are co-morbid with PTSD. Empirical investigations have generally demonstrated the feasibility and effectiveness of these approaches. Examples include PTSD with Substance Use Disorder (SUD) (Mills et al., 2012) where SUD and PTSD interventions are integrated and implemented relatively simultaneously and PTSD with Borderline Personality Disorder (BPD) (Harned, Korslund and Linehan, 2014) where ideally the BPD and PTSD interventions occur concurrently (but only once the patient has developed the emotional and behavioural control to tolerate the PTSD and its co-morbidity but is rather a distinct disorder with a specific symptom profile.

Considering that ICD-11 CPTSD is a new condition, it will take a substantial amount of time before an evidence base accumulates regarding its treatment. However, there is evidence on interventions that addressed at least partially the symptoms of CPTSD, including those of DSO. The aim of this systematic review and meta-analysis was to synthesise the evidence on effectiveness of treatments for the symptoms of CPTSD and identify therapies that look most promising for treating the symptoms of CPTSD. To achieve this goal, we examined evidence from trials for PTSD where participants were also likely to have clinically significant levels of one or more CPTSD DSO symptom clusters at baseline, and where usable data on the effect of interventions on these symptoms were reported. We also aimed to explore the moderating effect of RCT quality, the developmental timing of traumatic exposure (childhood vs. adulthood), phased vs. non-phased interventions, and individual vs. group interventions on treatment outcome. Our ultimate goal was to create a list of research priorities to inspire future research in the treatment of ICD-11 CPTSD.

# Method

#### Protocol registration

A protocol for this systematic review and meta-analysis was pre-registered (CRD42017055305) on February 2017. Changes to the protocol are listed in the supplement.

#### Search strategy and study selection

The search process was conducted in three main phases. First, MEDLINE, PsycINFO, EMBASE and PILOTS databases were searched for studies published from database inception to October 2017 using the following search terms: ("PTSD" or "posttrauma\*" or "psychological stress\*" or "combat" or "post-trauma\*" or "gross stress reaction" or "stress disorder\*" or "trauma\*" or "psychological trauma") AND ("randomised" or "randomized" or "randomised controlled trial" or "randomized controlled trial" or "RCT") AND ("therapy" or "psychological therapy" or "psychological intervention" or "intervention" or "treatment"). The only limiter applied in this search was language (English only). Second, to update the search, the same databases were searched for studies published from database inception to January 2018 using similar search terms: ("PTSD" or "posttrauma\*" or "psychological stress\*" or "combat" or "post-trauma\*" or "gross stress reaction" or "stress disorder\*" or "trauma\*" or "psychological trauma") AND ("randomised" or "randomized" or "RCT") AND ("therapy" or "intervention" or "treatment"). Limiters applied in this search were language (English only), humans, age group (adolescence, defined as between 13 and 17 years old, and adulthood, defined as 18 years and older), treatment and prevention, and randomised controlled trials. Third, the reference lists of earlier systematic reviews and meta-analyses of clinical trials for PTSD were screened for additional studies (Bisson et al., 2013; Bradlev et al., 2005; Cusack et al., 2016; Ehring et al., 2014; Imel et al., 2013; Kline et al., 2018). Three independent investigators (AB, SR, PM) carried out the search. Any discrepancies between search results were discussed and resolved with members of the research team (PHU, TK). As a final step, unpublished data were identified through contacting investigators and searching clinical trial registries (ClinicalTrials.gov and the UK Clinical Trials Gateway).

Studies were eligible for inclusion if they were randomised controlled trials (RCTs) reporting the effects of an individual or group-based psychological intervention for adults (mean age  $\geq 16$  years) with PTSD (ICD-10 and/or DSM-III-IV criteria), if participants experienced at least one of the additional CPTSD criteria at baseline (affect dysregulation, negative self-concept and disturbances in relationships, as defined in ICD-11), and if participants were free from developmental or intellectual disability, neurodegenerative disorders and acquired and/or traumatic brain injury. Studies where participants had comorbid substance misuse difficulties or other mental health conditions were included, but studies where participants had a primary diagnosis of substance misuse disorder were excluded. Case studies, uncontrolled trials and crossover trials were not included.

To establish whether participants had clinically significant levels of one or more of the additional CPTSD symptom clusters at baseline, any published clinical cut-offs relating to the CPTSD syndrome or individual CPTSD DSO symptoms were referred to in the first instance. If these were not available, any original validation study of the CPTSD index was referred to in order to try to identify relevant healthy norms; if the mean of the participants was more than one standard deviation (SD) away from the mean of these norms (in the direction of impairment), participants were considered to have clinically significant levels of the relevant CPTSD index. If there was no original validation study or if studies did not contain relevant healthy norms, studies that contained such norms was then searched for; if there were multiple studies, those with the largest sample sizes were prioritised. If the above clinical cut-offs or relevant norms could not be obtained, a decision about clinical significance was made on a case-by-case basis (e.g., if the participants' mean on a CPTSD DSO symptom indicated that

they were closer to being intact than impaired, they were not considered to have clinically significant levels of the relevant CPTSD symptom).

We defined a 'psychological intervention' as a talk-based intervention delivered by a trained therapist who adapted the treatment to patients on the basis of a therapeutic relationship (i.e., no delivery of a non-modifiable standard protocol, e.g., progressive muscle relaxation) (Benish, Imel and Wampold, 2008), and met at least two of the following four criteria: (a) a citation to an established school or approach to psychotherapy; (b) a description of the therapy that contained a reference to a psychological process (e.g., operant conditioning); (c) a reference to a treatment manual that was used to guide the delivery of the treatment; (d) the identification of active ingredients of the treatment and citations for these ingredients. Some of the face-to-face interventions we included did not meet these criteria (e.g., mindfulness, yoga), however we decided to report their effects in the interests of completeness. Online or other non-face-to-face interventions, even though they may meet these criteria, were excluded because of their different method of delivery and in an effort to reduce heterogeneity.

We further categorized psychological interventions into four different groups; (a) CBT (see definition below); (b) exposure therapy alone (i.e., psychological interventions, which were not better defined as CBT, emphasizing exposure to the trauma memory as the principal active treatment component, such as PE and imaginal exposure); (c) EMDR (i.e., psychological interventions consistent with the manual by Shapiro, 1995); (d) other psychological interventions (e.g., mindfulness). As per NICE guidelines, CBT was defined as a discrete psychological intervention where service users: (i) establish links between thoughts, feelings or actions with respect to the current or past symptoms, and/or functioning; (ii) re-evaluate their perceptions, beliefs or reasoning in relation to the target symptoms (National Collaborating Centre for Mental Health, 2014). To be categorized as CBT, the intervention also had to focus on at least one of the following: (iii) service users monitoring their own

thoughts, feelings or behaviours with respect to the symptom or recurrence of symptoms; (iv) promotion of alternative ways of coping with the target symptom (National Collaborating Centre for Mental Health, 2014). Given this broad definition of CBT, psychological interventions which involved cognitive/imagery modification with or without exposure therapy were considered to be CBT in nature.

We compared psychological intervention(s) to each other or to a control condition, which could be treatment as usual (TAU; also included 'waiting list control'), or TAU plus a non-specific therapeutic intervention (i.e. befriending, counselling).

#### Outcomes and data extraction

Our primary outcome was the standardised difference between groups at end of treatment in severity of (a) PTSD symptoms (as per ICD-11, DSM III-IV criteria) and (b) affect dysregulation, negative self-concept and disturbances in relationships. These were also used to calculate the associated number needed to treat (NNT) for clinically significant response, based on different estimates of response rates in the control condition.

Two reviewers (PHU, AB) extracted data relating to study characteristics, including details on participants, interventions received and outcomes assessed. Three reviewers (PM, AB, SR) also completed independent assessments of whether participants' mean baseline scores on measures of CPTSD symptoms were within the clinical range, which were then discussed and approved by two other reviewers (TK, PHU). Study authors were contacted in every case where CPTSD-relevant outcomes appeared to have been assessed but not reported. To assess outcomes, we extracted means and standard deviations (SD) where possible. If SDs were not reported, then these were derived from standard errors, confidence intervals, p-values or t-values where possible, following Cochrane Handbook procedures (Higgins and Green, 2011).

# Analysis

We used Comprehensive Meta-Analysis software (version 3) for the meta-analyses. We first calculated the post-intervention standardised mean difference (Hedges' g) and standard error (SE) for each individual study on each outcome (PTSD, affect dysregulation, negative self-concept, disturbances in relationships). Hedges' g was selected as the effect size measure because it accounts for variation in sample size and sample variance (Deeks, Altman and Bradburn, 2001). A composite effect was also computed for each study by combining PTSD and any available CPTSD DSO outcome data. To do this, we computed the average Hedges' g and associated SE across the outcomes. The range of measures used to assess these meant it was not feasible to adjust the composite estimate for the between-outcome correlation, and had to instead assume this was zero. When the number of participants (N) contributing data to each domain differed, we used the smallest N for the composite estimate. When there was sufficient data (at least two studies), we calculated the differences between interventions and controls on PTSD, affect dysregulation, negative self-concept, and disturbances in relationships individually, using DerSimonian and Laird (1986) random-effects meta-analyses. We then pooled data from studies reporting PTSD plus (a) 1, 2 or 3 CPTSD DSO outcomes, (b) 2 or 3 CPTSD DSO outcomes, and (c) all 3 CPTSD DSO outcomes. The estimates were expressed in units of Hedges' g with associated 95% confidence intervals. Between group differences in clinically significant change were derived from the Hedges' g estimate and an assumed control event response rate (CER) using the Furukawa method (Furukawa, 1999; Furukawa and Leucht, 2011; http://rpsychologist.com/d3/cohend/) and presented as NNT for benefit or harm. Morina et al., (2014) report a CER of 44% for PTSD however because CPTSD is assumed to have a poorer prognosis we estimated what the NNT to benefit or harm would be if we halved this value to 22%. We also estimated what the NNT would be if the natural remission rate in the control conditions was either very high (50%) or very low (10%). Using the relative group

difference and a range of assumed CERs to compute NNT is the method recommended by the Cochrane Handbook, since this "helps users to understand the important impact that typical baseline risks have on the absolute benefit that they can expect" (Higgins and Green., 2011).

The potential impact of publication bias was assessed using funnel plots, Egger's test and Duval and Tweedie's Trim-and-Fill procedure (random-effects) (Duval and Tweedie, 2000; Egger et al., 1997), but only for analyses derived from at least 10 studies (Higgins and Green, 2011). Cohen's (1988) established conventions (small = 0.2, moderate = 0.5, large = 0.8) were used to interpret individual and meta-analytical estimates of Hedges' g. Statistical significance was inferred when p-values were below 0.05, although values between 0.01 and 0.09 were downgraded for imprecision. Heterogeneity was assessed using the I<sup>2</sup> statistic, and compared with thresholds specified in the Cochrane Handbook (<40% low; 30-60% moderate; 50-90% substantial; 75-100% considerable) (Higgins and Green, 2011).

#### Assessment of study and outcome quality

Individual study quality was assessed using the Cochrane Collaboration Risk of Bias tool (Higgins et al., 2011) and meta-analytical estimates were assessed using the GRADE approach (Guyatt et al., 2008) (see supplement). The GRADE approach considers the quality of studies contributing to each analysis, the consistency, directness and precision of the pooled estimate, and the risk of publication bias.

Cochrane risk of bias ratings were completed by two reviewers independently (PM, AB), and checked by a third (PHU). An overall individual study quality rating was also produced (see supplement for criteria). GRADE ratings were performed by one reviewer (PHU) and checked by two others (PM, TK). An overall GRADE assessment is provided alongside each outcome to inform the interpretation of these findings.

#### Moderator analyses

We combined all studies into a single dataset to conduct a series of pre-specified univariate moderator analyses, and one multivariate analysis, again using Comprehensive Meta-Analysis software (version 3). The outcome for each meta-regression analysis was the post-treatment group difference in CPTSD symptom severity. For this we used, in order of preference, the composite estimates of differences in (1) PTSD plus the three CPTSD DSO symptom clusters; (2) PTSD plus two CPTSD DSO symptom clusters (3) PTSD plus one CPTSD DSO symptom cluster or (4) PTSD alone.

Pre-specified univariate analyses included the relevant Cochrane Risk of Bias parameters (sequence generation, allocation concealment, detection bias, reporting bias, attrition bias), onset of trauma (childhood vs adulthood), degree to which sample met CPTSD criteria (i.e., whether data on PTSD plus three, two, one or no CPTSD DSO symptom clusters were used) and therapy format (individual vs group). There was insufficient data to support pre-specified analysis of phased vs non-phased interventions. We also examined the effect of therapy type (individual CBT, group CBT, EMDR, exposure alone, group IPT), and the effect of using a non-specific control condition (i.e., versus a usual care / waiting list control group). To ensure that all studies with 3 or more arms could be included without double-counting of participants, we split the sample size of any shared treatment or control arms in half for these comparisons, as recommended in the Cochrane Handbook (Higgins and Green, 2011), and revised the individual study effect sizes accordingly. To ensure power for the multivariate analyses, we limited this to 5 variables; study quality, therapy type, degree to which sample met CPTSD criteria, trauma onset, and use of a non-specific control condition.

#### Results

#### Study selection

The search returned 28,521 results, of which 28,310 were excluded on the basis of title or abstract (see Figure 1). Following title and abstract screening, the full texts of the remaining 211 articles were examined. One hundred and forty one full text articles were excluded. A further 19 full text articles were excluded because they described studies that did not include clinically significant levels of one or more CPTSD DSO symptom clusters at baseline. Fifty-one studies met full inclusion criteria and were included in the current study. Of these, 35 studies had a CBT arm, 11 had an exposure only arm, 9 had an EMDR arm, and 9 assessed the effect of other interventions, including interpersonal psychotherapy (IPT), mindfulness, trauma management training (TMT), dialogical exposure therapy (DET), dialectical behaviour therapy, CBT plus emotion regulation training, and stabilisation therapy. Figure 2 provides an overview of studies contributing to each analysis. A table of included study characteristics and a table of excluded studies, with reasons for exclusion, are provided in the supplement.

# Insert Figure 1

#### **Insert Figure 2**

#### Quality assessment

The results of the Cochrane risk of bias assessment are shown in the supplement and GRADE ratings for each meta-analytical outcome are shown below and in the far right column of Tables 1-4 and Table J.1 (supplement). Just over half of the included studies used appropriate methods to generate a random sequence to allocate participants to groups, but poor reporting limited our assessment of this domain. A slightly smaller proportion had a low risk of bias for

allocation sequence concealment, but again poor reporting prevented a clear assessment of this domain. The majority of studies had a low risk of detection bias because assessors were unaware of the group that participants had been allocated to. Most also had a low risk of attrition bias with acceptable rates of missing post-intervention data (<25%). However, most had a high risk of reporting bias primarily due to a lack of a preregistered protocol. The risk of performance bias was unavoidably high across all studies due to the nature of the interventions, which precluded blinding of participants. Overall, we rated the majority of studies as high in methodological quality.

# Meta-analytical outcomes

#### Cognitive behavioural therapy (CBT) (Table 1, and supplement)

As shown in Table 1, compared to usual care, CBT had a moderate-large effect on disturbances in relationships (k=16, g = -0.66; 95% CI = -0.84, -0.48) and large effects on affect dysregulation (k=3, g = -1.42; 95% CI = -2.20, -0.65), negative self-concept (k=9, g = -0.82; 95% CI = -1.19, -0.44) and PTSD symptoms (k=27, g = -0.90; 95% CI = -1.11, -0.68) (all moderate quality evidence), with the NNT varying from 2 (affect dysregulation assuming CER of 22%) to 6 (disturbances in relationships assuming CER of 10%). Moderate to large effects were also observed on the composite estimates of PTSD and CPTSD DSO symptoms (low to high quality evidence), with NNTs of between 3 (PTSD + 1, 2, or 3 CPTSD DSO outcomes assuming CER of 50%) and 8 (PTSD + 3 CPTSD DSO outcomes assuming CER of 10%). However few studies measured more than one type of CPTSD DSO symptom. Significant publication bias was detected whenever there were sufficient studies to assess this, however only the estimate for disturbances in relationships was reduced when trim-and-fill analysis was applied. Compared to non-specific control interventions, CBT had a small effect on disturbances in relationships (k=3, g = -0.32; 95% CI = -0.60, -0.03) and a small-moderate

effect on PTSD symptoms (k=9, g = -0.37; 95% CI = -0.66, -0.09) (moderate quality evidence), with NNTs varying between 7 (PTSD assuming 50% CER) and 15 (disturbances in relationships assuming 10% CER). Although there was no evidence it had significant effects on affect dysregulation and negative self-concept, few studies reported usable data. When we pooled effects from all 9 studies reporting data on PTSD and at least one CPTSD DSO domain, a small effect was observed (k=9, g = -0.34; 95% CI = -0.62, -0.06; low quality evidence), with NNTs of between 8 (50% CER) and 14 (10% CER), but no studies measured more than one domain.

# Exposure therapy alone (Table 2, and supplement)

As shown in Table 2, compared to usual care, exposure therapy alone had a moderate effect on disturbances in relationships (k=4, g = -0.59; 95% CI = -1.12, -0.07; moderate quality evidence), a moderate-large effect on negative self-concept (k=3, g = -0.73; 95% CI = -1.03, - 0.43; moderate quality evidence), and a large effect on PTSD symptoms (k=6, g = -1.05; 95% CI = -1.52, -0.58; low quality evidence), with NNTs of between 3 (PTSD - all assumed CERs) and 7 (disturbances in relationships, assuming 10% CER). No studies examined whether exposure was superior to usual care in relation to affect dysregulation. Moderate to large effects on the composite outcomes of PTSD and CPTSD DSO symptoms were observed (low to high quality evidence), with NNTs ranging from 3 (PTSD + 1, 2 or 3 CPTSD DSO outcomes, CERs of 22% and 50%) to 7 (PTSD + 2 or 3 CPTSD DSO outcomes, assuming 10% CER), however only one study provided usable data on more than one type of CPTSD DSO symptom. There was no evidence that exposure alone was superior to non-specific therapies in relation to disturbances in relationships, but only one study provided usable data. No studies reported whether exposure alone was superior to non-specific therapies in relation to either affect dysregulation or negative self-concept. Two studies found no effect of exposure alone on either

PTSD data, or the composite outcome of PTSD plus CPTSD DSO symptoms (low quality evidence). No studies provided data on more than one CPTSD DSO symptom.

*Eye-Movement and Desensitisation and Reprocessing therapy (EMDR) (Table 3, and supplement)* 

As shown in Table 3, compared to usual care, the few available studies suggested EMDR had a moderate effect on negative self-concept (k=1, g = -0.61; 95% CI = -1.04, -0.17; low quality evidence), a moderate-large effect on disturbances in relationships (k=4, g=-0.76; 95% CI = -1.35, -0.16; moderate quality evidence), and large effects on affect dysregulation (k=1, g=-1.64; 95% CI = -2.56, -0.72; very low quality evidence) and PTSD symptoms (k=4, 1)g = -1.26; 95% CI = -2.01, -0.51; low quality evidence), with NNTs ranging from 2 (affect dysregulation, all CERs) to 7 (disturbances in relationships, assuming CER of 10%). EMDR also had a large effect on the composite outcome of PTSD and at least one CPTSD DSO symptom (k=4, g = -1.15; 95% CI = -1.92, -0.37; low quality evidence), with NNTs of 2 (CER of 22%) or 3 (CER of 10% or 50%), but it did not have an effect on the composite outcome of PTSD and more than one CPTSD DSO symptom (very low quality evidence). There was no evidence that EMDR was superior to non-specific interventions in relation to disturbances in relationships or affect dysregulation (very low quality evidence). Although moderate-large effects on negative self-concept (k=2, g = -0.78; 95% CI = -1.56, -0.01) and PTSD symptoms (k=3, g=-0.69; 95% CI=-1.35, -0.03) (very low quality evidence) were observed, with NNTs of between 4 (negative self-concept, all CERs) and 6 (PTSD; CER of 10%), these analyses were based on only 2-3 studies. A moderate effect on the composite outcome of PTSD and at least one CPTSD DSO symptom was observed (k=3, g = -0.52; 95% CI = -0.97, -0.08; low quality evidence), with NNTs of between 5 (CER 50%) and 8 (CER 10%), but no effect was

found on the composite outcome of PTSD and more than one CPTSD DSO symptom (very low quality evidence).

# Comparison of CBT, Exposure and EMDR (Table 4, and supplement)

As shown in Table 4, there was very limited evidence that EMDR had a small-moderate advantage over CBT in relation to PTSD symptoms (k=2, g = 0.37; 95% CI = 0.03, 0.71; low quality evidence), with an NNT of 7-12, but no differences between CBT, exposure alone or EMDR were observed for any other outcomes

# Other comparisons (supplement)

As shown in Table J.1 (supplement), one small study (Krupnick 2008) found IPT had an advantage over usual care in reducing PTSD plus disturbances in relationships (k=1, g = -1.02; 95% CI = -1.65, -0.39; very low quality evidence), with an NNT of 3-4, and another small study (Azad marzabadi 2014) found mindfulness was more effective than usual care in relation to disturbances in relationships (k=1, g = -1.60; 95% CI = -2.43, -0.77; very low quality evidence), with an NNT of 2-3. Several other small studies compared various psychotherapeutic interventions to other interventions, or to CBT, exposure or EMDR. We found no evidence to favour any particular intervention in relation to the composite outcome of PTSD plus CPTSD DSO symptoms (very low to low quality evidence).

#### Moderator analyses (Figure 3, and supplement)

As shown in Table L.1 (supplement), use of a non-specific control condition rather than usual care or waiting list was associated with a smaller benefit of psychological therapy in univariate meta-regression, with a reduction in Hedges' g of 0.48, (95% CI = 0.18, 0.77). No other moderators were significant when examined individually. As shown in Table M.1

(supplement), the effect of using a non-specific control condition was larger in multivariate meta-regression, with a reduction in Hedges' g of 0.69 (95% CI = 0.39, 1.00) in this analysis. Study quality and age of trauma onset also emerged as significant moderators of therapy effects in this analysis. Low quality studies were associated with a significantly *lower* effect size, with a reduction in Hedges' g of 0.30 (95% CI = 0.00, 0.61). Studies where participants had predominantly childhood-onset trauma were associated with a reduction in Hedges' g of 0.35 (95% CI = 0.02, 0.69), when compared to trials where most participants had adult-onset trauma (Figure 3).

# Discussion

We examined evidence from RCTs of psychological treatments for PTSD where participants were also likely to have clinically significant levels of one or more CPTSD DSO symptoms at baseline, and where usable data on the effect of interventions on these symptoms were reported. A total of 51 studies met inclusion criteria. Overall, results indicate that when compared to usual care, CBT, Exposure alone and EMDR perform relatively equally for symptoms of PTSD and the DSO symptoms of negative self-concept and disturbances in relationships. While the quality of this evidence was moderate for CBT, it ranged from low to moderate for Exposure alone and EMDR. Few trials reported the effectiveness of psychological therapies for symptoms of affect dysregulation. Low quality evidence suggests that EMDR has a small-moderate advantage over CBT in relation to PTSD symptoms, but there was no evidence of any differences between CBT, Exposure alone or EMDR for the other outcomes including DSO symptoms. Univariate and multivariate meta-regression confirmed that the effectiveness of psychological therapies was considerably lower when compared to nonspecific therapies, which suggests that non-specific effects may account for a large proportion of therapeutic change in symptoms of CPTSD in these trials. The multivariate meta-regression also found that treatment outcome may be moderated by the developmental time of the onset of psychological trauma, with childhood trauma being associated with smaller effects of psychological therapies on CPTSD symptoms.

The data are encouraging in that the accumulation of evidence suggests that there are specific interventions that work for several of the CPTSD symptom clusters. The data also suggest that no particular type of intervention (exposure, cognitive re-appraisal, bilateral stimulation) is necessary to resolve any one symptom cluster. A critical question is whether current treatments devised for PTSD are equally effective for those who will be diagnosed with CPTSD. Our results replicate earlier findings that individual trauma-focused treatments show large effect sizes. Although the evidence is at a very early stage, we found that some nontrauma-focused therapies, such as mindfulness and IPT, may also reduce PTSD and interpersonal disturbance, suggesting alternative options. Importantly, childhood abuse was found to moderate all outcomes across all types of treatments, suggesting those with a history of childhood trauma may experience less improvement, and that current treatments for this patient population can be improved. These results have implications for the treatment of CPTSD as those with childhood abuse are at risk for CPTSD and in this meta-analysis may represent those more likely to have the full symptom profile.

Research is needed to determine how to optimize treatment outcomes for those with childhood abuse and other populations at risk for CPTSD. This includes identifying which treatment interventions are most effective for specific symptom clusters, which are most acceptable to patients, in what order to present interventions and the optimal duration of different types of interventions. Considering current debates in the literature, it would have been useful to explore the usefulness of phased vs. non-phased interventions and individual vs. group interventions for CPTSD. Unfortunately, we did not find adequate evidence to enable further analysis of these treatment outcome moderators. There is substantial evidence

indicating that CPTSD and PTSD represent distinct patient populations with different symptoms profiles (Brewin et al., 2017), suggesting the value of developing treatments that more precisely and effectively resolve the differing effects of trauma exposure by systematically testing type, order and duration of interventions specific to each disorder and taking into account patient preferences across both disorders (Cloitre, 2015).

Our meta-analysis has a number of strengths. We minimised the risk of bias by preregistering the review, and we minimised errors and omissions by having two or more reviewers conduct comprehensive searches, assess study quality and extract descriptive data. We considered a range of treatments from different countries and included participants with a range of backgrounds and types of psychological trauma including military, civilian and childhood trauma. Many studies have used qualified therapists and considered assessments of adherence to the protocol. However, most of the research was conducted in western countries, thus limiting the extent to which the findings may generalise to non-western countries. Furthermore, the evidence we have reviewed as part of this meta-analysis was predominantly on DSM-IV PTSD. Most studies did not present data on multiple traumatisation which typically results in CPTSD (Karatzias et al., 2016). Even when the index trauma that was targeted occurred in adulthood in included studies, it would be useful to assess lifetime traumatic history and consider the accumulative effect of multiple traumatisation. In relation to outcomes, we have only considered therapeutic gains at post-treatment. Future research should explore long-term outcomes of these interventions. Furthermore, for this meta-analysis we have used proxy measures for the CPTSD constructs. It might well be the case that a number of studies that included people with CPTSD have not been included in the study as they have not reported outcomes on relevant constructs or reported outcomes have not met clinical thresholds or our definition of 'clinical significance'. It might also be the case that the measures employed in included studies do not accurately reflect the corresponding DSO clusters, thus introducing some measurement bias. Moreover, while the quality of the meta-analytical evidence was high or moderate for some of the outcomes (e.g., when CBT was compared with usual care or non-specific control interventions), it was low or very low for most of the outcomes. Related to this, there was substantial heterogeneity for just over half of the outcomes. Thus, there is some uncertainty in the conclusions that can be drawn. It is also worth noting that we did not downgrade the meta-analytical outcomes for indirectness, as indirect evidence of psychological interventions for CPTSD was the focus of this review. If, on the other hand, we had been interested in direct evidence of psychological interventions for CPTSD, most if not all the outcomes would have been downgraded for indirectness.

There is clearly a need for further well-designed trials of psychological therapies that incorporate appropriate methods of randomisation, blinding of assessors, long-term follow up and appropriate training of therapists and monitoring of treatment adherence. We have identified a set of research priorities to benefit people with CPTSD in the future that might directly or indirectly result from the findings of this review:

- Effectiveness of phased vs. non-phased interventions for CPTSD: Very few included studies in this meta-analysis have incorporated a phased approach to treatment and it was not possible to address this question.
- Effectiveness of trauma focused treatments vs. non trauma focused treatments. Existing evidence is predominantly focused on trauma-focused treatments.
- Head-to-head comparisons between trauma focused treatments for CPTSD. Most studies explored the effectiveness of interventions against standard care or no treatment.
- Exploring safety of trauma focused therapies for CPTSD. It is essential that future research in this area provides information on adverse effects.
- Investigation of whether diagnosis of CPTSD moderates outcomes when compared against those who do not meet diagnosis in standard treatments. Clinical reality suggests

that many people do not meet full diagnostic criteria but still suffer from a number of debilitating symptoms that relate to that condition.

- Appropriateness and effectiveness of trauma focused treatments for CPTSD following childhood trauma. In this meta-analysis, childhood trauma was found to negatively moderate the effect of trauma focused interventions.
- Comparing pharmacotherapy vs. psychotherapy for CPTSD. In this meta-analysis we did not address the effectiveness of pharmacotherapies alone or in combination with psychotherapy.
- Considering the nature of the three DSO factors, it is worth exploring the effectiveness of attachment based interventions and relational therapies as limited evidence is currently present for these interventions.
- Exploring the effectiveness of individual vs. group interventions for CPTSD. We found no evidence addressing this question for people with CPTSD.
- Exploring the effectiveness of interventions that tackle all CPTSD symptom clusters in
  a single study using as a primary outcome of CPTSD based on a dedicated measure.
  The present review extracted proxy data from existing trials that measure the CPTSD
  constructs.

In conclusion, this meta-analysis is the first step in identifying effective treatments for CPTSD. Findings regarding the usefulness of trauma-focused interventions look promising but less so for CPTSD symptoms following childhood trauma. Further research is needed to explore and develop existing and new treatments for CPTSD.

References

- Azad Marzabadi E and Hashemi Zadeh SM (2014) The Effectiveness of Mindfulness Training in Improving the Quality of Life of the War Victims with Post Traumatic stress disorder (PTSD) *Iranian Journal of Psychiatry* **9**, 228–236.
- Barrera TL, Mott JM, Hofstein RF and Teng EJ (2013) A meta-analytic review of exposure in group cognitive behavioral therapy for posttraumatic stress disorder. *Clinical Psychology Review* 33, 24–32.
- Benish SG, Imel ZE and Wampold BE (2008) The relative efficacy of bona fide psychotherapies for treating post-traumatic stress disorder: A meta-analysis of direct comparisons. *Clinical Psychology Review* 28, 746–758.
- **Bisson J and Andrew M** (2005) Psychological treatment of post-traumatic stress disorder (PTSD) *Cochrane Database of Systematic Reviews* **18**, CD003388.
- Bisson J and Andrew M (2007) Psychological treatment of post-traumatic stress disorder (PTSD) *Cochrane Database of Systematic Reviews* **18**, CD003388.
- Bisson JI, Ehlers A, Matthews R, Pilling S, Richards D and Turner S (2007)
  Psychological treatments for chronic post-traumatic stress disorder. Systematic review and meta-analysis. *British Journal of Psychiatry* 190, 97-104.
- Bisson JI, Roberts NP, Andrew M, Cooper R and Lewis C (2013) Psychological therapies for chronic post-traumatic stress disorder (PTSD) in adults. *Cochrane Database of Systemic Reviews* 13, CD003388.
- Bradley R, Greene J, Russ E, Dutra L and Westen D (2005)A multidimensional metaanalysis of psychotherapy for PTSD. *American Journal of Psychiatry* **162**, 214-27.

Brewin CR, Cloitre M, Hyland P, Shevlin M, Maercker A, Bryant RA, Humayun A, Jones LM, Kagee A, Rousseau C, Somasundaram D, Suzuki Y, Wessely S, van Ommeren M and Reed GM (2017) A review of current evidence regarding the ICD-11 proposals for diagnosing PTSD and complex PTSD. *Clinical Psychology Review* 58, 1-15.

- **Callahan KL, Price JL and Hilsenroth MJ** (2004) A review of interpersonal psychodynamic group psychotherapy outcomes for adult survivors of childhood sexual abuse. *International Journal of Group Psychotherapy* **54**, 491-519.
- Cloitre M, Garvert DW, Brewin CR, Bryant RA and Maercker A (2013) Evidence for proposed ICD-11 PTSD and complex PTSD: a latent profile analysis. *European Journal of Psychotraumatology* **4**, doi: 10.3402/ejpt.v4i0.20706
- Cloitre M, Courtois CA, Ford JD, Green BL, Alexander P, Briere J, Herman JL, Lanius R, Stolbach BC, Spinazzola J, Van der Kolk BA and Van der Hart O (2012) The ISTSS Expert Consensus Treatment Guidelines for Complex PTSD in Adults. Retrieved from http:// www.istss.org/ISTSS\_Main/media/Documents/ISTSS-Expert-Concesnsus-Guidelines-for-Complex-PTSD-Updated-060315.pdf
- Cloitre M, Petkova E, Wang J and Lu Lassell F (2012) An examination of the influence of a sequential treatment on the course and impact of dissociation among women with PTSD related to childhood abuse. *Depression and Anxiety* 29, 709–717.
- Cloitre M, Stovall-McClough KC, Nooner K, Zorbas P, Cherry S, Jackson CL, Gan W and Petkova E (2010) Treatment for PTSD related to childhood abuse: A randomized controlled trial. *American Journal of Psychiatry* **167**, 915–924.
- Cloitre M (2015) The "one size fits all" approach to trauma treatment: Should we be satisfied? *European Journal of Psychotraumatology* 6, 27344. http://dx.doi.org/10.3402/ejpt.v6.27344

- **Cohen J** (1988) *Statistical Power Analysis in the Behavioral Sciences*. Hillsdale, NJ: Erlbaum.
- Cusack K, Jonas DE, Forneris CA, Wines C, Sonis J, Middleton JC, Feltner C,
  Brownley KA, Olmsted KR, Greenblatt A, Weil A and Gaynes BN (2016)
  Psychological treatments for adults with PTSD: A systematic review and metaanalysis. *Clinical Psychology Review* 43, 128-141.
- Deeks JJ, Altman, DG and Bradburn MJ (2001) Statistical methods for examining heterogeneity and combining results from several studies in meta-analysis. Systematic reviews in health care: meta-analysis in context, 285-312. https://doi.org/10.1002/9780470693926.ch15.
- De Jongh A, Resick PA, Zoellner LA, van Minnen A, Lee CW, Monson CM, Foa EB,
  Wheeler K, Broeke ET, Feeny N, Rauch SA, Chard KM, Mueser KT, Sloan DM,
  van der Gaag M, Rothbaum BO, Neuner F, de Roos C, Hehenkamp LM, Rosner
  R, Bicanic IA (2016) Critical analysis of the current treatment guidelines for
  complex PTSD in adults. *Depression and Anxiety* 33, 359–369.
- DerSimonian R and Laird N (1986) Meta-analysis in clinical trials. *Control Clinical Trials*7, 177–188.
- **Duval S and Tweedie R (2000)** Trim and fill: A Simple Funnel Plot Based Method of Testing and Adjusting for Publication Bias in Meta-analysis. *Biometrics* **56**, 455–463.
- Egger M, Smith GD, Schneider M and Minder C (1997) Bias in meta-analysis detected by a simple, graphical test. *British Medical Journal (Clinical Research Ed.)* 315, 629– 634.
- Ehring T, Welboren R, Morina N, Wicherts JM, Freitag J and Emmelkamp PMG (2014) Meta-analysis of psychological treatments for posttraumatic stress disorder in adult survivors of childhood abuse. *Clinical Psychology Review* 34, 645-657.

- Elklit A, Hyland P and Shevlin M (2014) Evidence of symptom profiles consistent with posttraumatic stress disorder and complex posttraumatic stress disorder in different trauma samples. *European Journal of Psychotraumatology* 5, 24221. http://dx.doi.org/10.3402/ ejpt.v5.24221
- Foa EB, Keane TM, Friedman MJ and Cohen JA (2009) Effective treatments for PTSD: Practice guidelines from the International Society for Traumatic Stress Studies. New York: Guilford Press.

Furukawa TA (1999) From effect size into number needed to treat. Lancet 353, 1680.

Furukawa TA and Leucht S (2011) How to obtain NNT from Cohen's d: comparison of two methods. *PLoS One* 6, e19070. doi: 10.1371/journal.pone.0019070.

Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P,

Schünemann HJ and GRADE Working Group (2008) GRADE: An emerging consensus on rating quality of evidence and strength of recommendations. *British Medical Journal* **336**, 924–926.

- Harned MS, Korslund KE and Linehan MM (2014) A pilot randomized controlled trial of Dialectical Behavior Therapy with and without the Dialectical Behavior Therapy
   Prolonged Exposure protocol for suicidal and self-injuring women with borderline personality disorder and PTSD. *Behaviour Research and Therapy* 55, 7–17.
- Herman J (1992) Complex PTSD: A Syndrome in Survivors of Prolonged and Repeated Trauma. *Journal of Traumatic Stress* 5, 377-391.
- Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, Savovic J, Schulz KF, Weeks L, Sterne JA, Cochrane Bias Methods Group and Cochrane Statistical Methods Group (2011) The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *British Medical Journal* 343, d5928. doi: 10.1136/bmj.d5928.

Higgins JPT and Green S (2011) Cochrane Handbook for Systematic Reviews of Interventions. The Cochrane Collaboration and John Wiley & Sons Ltd: England.

- Hyland P, Murphy J, Shevlin M, Vallières F, McElroy E, Elklit A, Christoffersen M and Cloitre M (2017a) Variation in post-traumatic response: The role of trauma type in predicting ICD-11 PTSD and CPTSD symptoms. *Social Psychiatry and Psychiatric Epidemiology* 52, 727-736.
- Hyland P, Shevlin M, Brewin CR, Cloitre M, Downes A, Jumbe S, Karatzias T, Bisson JI and Roberts NP (2017b) Validation of post-traumatic stress disorder (PTSD) and complex PTSD using the International Trauma Questionnaire. *Acta Psychiatrica Scandinavica* 136, 313-322.
- Hyland P, Ceannt R, Daccache F, Abou Daher R, Sleiman J, Gilmore B, Byrne S,
  Shevlin M, Murphy J and Vallières F (2018) Are posttraumatic stress disorder
  (PTSD) and complex-PTSD distinguishable within a treatment-seeking sample of
  Syrian refugees living in Lebanon? *Global Mental Health*, 5, p.e14. doi:
  10.1017/gmh.2018.2.
- Imel ZE, Laska K, Jakupcak M and Simpson TL (2013) Meta-analysis of dropout in treatments for PTSD. *Journal of Consulting and Clinical Psychology* 81, 394-404.
- Karatzias T, Shevlin M, Fyvie C, Hyland P, Efthymiadou E, Wilson D, Roberts N,
  Bisson JI, Brewin CR and Cloitre M (2016) Evidence of Distinct Profiles of
  Posttraumatic Stress Disorder (PTSD) and Complex Posttraumatic Stress Disorder
  (CPTSD) based on the New ICD-11 Trauma Questionnaire (ICD-TQ) *Journal of Affective Disorders* 207, 181 -187.

Karatzias T, Shevlin M, Fyvie C, Hyland P, Efthymiadou E, Wilson D, Roberts N,
Bisson JI, Brewin CR and Cloitre M (2017) Evidence of distinct profiles of
Posttraumatic Stress Disorder (PTSD) and Complex Posttraumatic Stress Disorder
(CPTSD) based on the new ICD-11 Trauma Questionnaire (ICD-TQ). Journal of
Anxiety Disorders 207, 181-187.

- Kline AC, Cooper AA, Rytwinksi NK and Feeny NC (2018) Long-term efficacy of psychotherapy for PTSD: A meta-analysis of randomized controlled trials. *Clinical Psychology Review* 59, 30-40.
- Knefel M, Garvert DW, Cloitre M and Lueger-Schuster B (2015) Update to an evaluation of ICD-11 PTSD and complex PTSD criteria in a sample of adult survivors of childhood institutional abuse by Knefel & Lueger-Schuster (2013): a latent profile analysis. *European Journal of Psychotraumatology* 6, 25290. http://dx.doi.org/10.3402/ ejpt.v6.25290
- Krupnick JL, Green BL, Stockton P, Miranda J, Krause E and Mete M (2008) Group interpersonal psychotherapy for low-income women with posttraumatic stress disorder. *Psychotherapy Research : Journal of the Society for Psychotherapy Research* 18, 497–507.
- Maercker A, Brewin CR, Bryant RA, Cloitre M, Reed GM, van Ommeren M,
  Humayun A, Jones LM, Kagee A, Llosa AE, Rousseau C, Somasundaram DJ,
  Souza R, Suzuki Y, Weissbecker I, Wessely SC, First MB and Saxena S (2013)
  Proposals for mental disorders specifically associated with stress in the ICD-11. *Lancet* 381, 1683-1685.
- Mills KL, Teesson M, Back SE, Brady KT, Baker AL, Hopwood S, Sannibale C, Barrett EL, Merz S, Rosenfeld J and Ewer PL (2012) Integrated exposure-based therapy for co-occurring posttraumatic stress disorder and substance dependence: A

randomized controlled trial. *Journal of the American Medical Association* **308**, 690–699.

- Morina N, Wicherts JM, Lobbrecht J and Priebe S (2014) Remission from posttraumatic stress disorder in adults: A systematic review and meta-analysis of long term outcome studies. *Clinical Psychology Review* **34**, 249–255.
- National Collaborating Centre for Mental Health (2014) Psychosis and Schizophrenia in adults: Treatment and management: Updated edition 2014. *National Institute for Health and Care Excellence (UK)* 14–36.
- Peleikis DE and Dahl AA (2005) A systematic review of empirical studies of psychotherapy with women who were sexually abused as children. *Psychotherapy Research* 15, 304–315.
- Perkonigg A, Höfler M, Cloitre M, Wittchen HU, Trautmann S and Maercker A (2015) Evidence for two different ICD-11 posttraumatic stress disorder in a community sample of adolescents and young adults. *European Archives of Psychiatry* and Clinical Neuroscience 266, 317-28.
- Roberts NP, Roberts PA, Jones N and Bisson JI (2015) Psychological interventions for post-traumatic stress disorder and comorbid substance use disorder: A systematic review and meta-analysis. *Clinical Psychology Review* 38, 25-38.
- Sachser C, Keller F and Goldbeck L (2016) Complex PTSD as proposed for ICD-11: validation of a new disorder in children and adolescents and their response to Trauma-Focused Cognitive Behavioral Therapy. *Journal of Child Psychology and Psychiatry* 58, 160–168.
- Shapiro F (1995) Eye movement desensitization and reprocessing (EMDR): Basic principles, protocols, and procedures. New York: Guilford Press.

- Shevlin M, Hyland P, Karatzias T, Roberts N and Bisson J (2017) Examining the disconnect between psychometric models and clinical reality. *Journal of Anxiety Disorders* 24, 54–59.
- Sloan DM, Feinstein BA, Gallagher MW, Beck JG and Keane TM (2013) Efficacy of group treatment for posttraumatic stress disorder symptoms: A meta-analysis. *Psychological Trauma: Theory, Research, Practice, and Policy* 5, 176–183.
- **Taylor JE and Harvey ST** (2010) A meta-analysis of the effects of psychotherapy with adults sexually abused in childhood. *Clinical Psychology Review* **30**, 749-767.
- **Taylor JE and Harvey ST** (2009) Effects of psychotherapy with people who have been sexually assaulted: A meta-analysis. *Aggression and Violent Behaviour* **14**, 273–285.
- Watts BV, Schnurr PP, Mayo L, Young-Xu Y, Weeks WB and Friedman MJ (2013) Meta-analysis of the efficacy of treatments for posttraumatic stress disorder. *The Journal of Clinical Psychiatry* 74, 541-550.
- **World Health Organization (2018)** The ICD-11 for mortality and morbidity statistics. Retrieved from https://icd.who.int/browse11/l-m/e

# **Supplementary Appendix to:**

Karatzias, T., Murphy, P., Cloitre, M., Bisson, J., Roberts, N., Shevlin, M., Hyland, P., Maercker, A., Ben-Ezra, M, Coventry, P., Mason-Roberts, S., Bradley, A., Hutton, P. (2019). Psychological interventions for ICD-11 Complex PTSD symptoms: Systematic review and meta-analysis. *Psychological Medicine*.

# **Content of Supplementary Appendix**

- A. Protocol
- B. Changes from Protocol
- C. Excluded Studies
- D. Summary of Characteristics of Included Studies
- E. Participants' baseline scores on the CPTSD symptom clusters and corresponding norms
- F. Risk of Bias Criteria
- G. Risk of Bias in Included Studies Summary
- H. Risk of Bias in Included Studies Detailed
- I. GRADE Assessment Criteria
- J. Table Other Comparisons
- K. Table Clinically Significant Response (number needed to treat per control group rate)
- L. Table Meta-regression Moderators (univariate)
- M. Table Meta-regression Moderators (multivariate)
- N. Forest Plots Cognitive/imagery modification with or without exposure vs TAU/WL or non-specific control
- O. Forest Plots Exposure only vs TAU/WL or non-specific control
- P. Forest Plots EMDR vs TAU/WL or non-specific control
- Q. Forest Plots Comparison of CBT, Exposure and EMDR
- R. Bubble Plots Meta-regression Moderators (univariate)
- S. Bubble Plots Meta-regression Moderators (multivariate)
- T. Funnel Plots for Meta-analyses (where publication bias is indicated)
- U. PRISMA Checklist
- V. References for the Included Studies
- W. References for the Excluded Studies
- X. Other References

# A. Protocol

Title: Psychological interventions for Complex PTSD: systematic review and meta-analysis.

*Reviewers:* Thanos Karatzias, Mick Fleming, Susan Roberts, Aoife Bradley, Claire Fyvie, Jonathan Bisson, Neil Roberts, Philip Hyland, Marylene Cloitre, Tobias Hecker, Andreas Maercker, Paul Hutton

# *Review question(s)*

What psychological interventions are effective for complex post traumatic stress disorder, and how effective are they? What is the safety and acceptability of psychological interventions for complex post traumatic stress disorder?

# Searches

Searches of MEDLINE, PsycINFO, EMBASE and PILOTS will be conducted using the search terms listed below. Unpublished trials will be identified through contacting investigators and through searching clinical trial registries such as Clinicaltrials.gov. Language will be restricted to English. There will be no time period restrictions. #1. PTSD or posttrauma\* or psychological stress\* or combat or post-trauma\* or gross stress reaction or stress disorder\* or trauma\* or psychological trauma. #2. randomized or randomized or randomized controlled trial or RCT or randomized controlled trial. #3.therapy or psychological therapy or psychological intervention or intervention or treatment).mp. [mp=ti, ab, ot, nm, hw, kf, px, rx, ui, an, eu, pm, sy, tn, dm, mf, dv, kw, fs].

# Types of study to be included

Randomised controlled trials with or without rater masking will be included. Uncontrolled, non-randomised and crossover trials, qualitative studies and case studies will be excluded.

# Condition or domain being studied

Complex post traumatic stress disorder (CPTSD) as described in ICD-11 proposals. According to ICD-11 individuals meet diagnostic criteria for complex PTSD if they meet existing criteria for post traumatic stress disorder (PTSD) (re-experiencing of the trauma, avoidance of reminders of the trauma, enhanced sense of threat indicated by hypervigilance and hyperarousal) and have clinically significant difficulties in affect dysregulation, a pervasive negative self-concept and experience interpersonal disturbances (Cloitre et al. 2013). Individuals meet diagnostic criteria for complex PTSD if they meet existing criteria for post traumatic stress disorder (PTSD) (re-experiencing of the trauma, avoidance of reminders of the trauma, enhanced sense of threat indicated by hypervigilance and hyperarousal) and have clinically significant difficulties in affect dysregulation, a pervasive traumatic stress disorder (PTSD) (re-experiencing of the trauma, avoidance of reminders of the trauma, enhanced sense of threat indicated by hypervigilance and hyperarousal) and have clinically significant difficulties in affect dysregulation, a negative self-concept and experience interpersonal disturbances (Cloitre et al. 2013).

# Participants/population

We are interested in the effect of psychological interventions on adults who meet criteria for CPTSD. However, since CPTSD is a new diagnostic category, we anticipate that few studies have explicitly included this group. For this reason we will only include trials of interventions where participants meet ICD and DSM – III and IV criteria for PTSD and present with clinically significant symptoms of re-experience, avoidance hyperarousal and score within the clinically significant range on at least one of the additional CPTSD criteria, namely emotion dysregulation, negative self-concept and interpersonal disturbance. We will only include trials where the mean or median age of participants is at least 16, and we will only include trials

where participants with developmental or intellectual disability, neurodegenerative disorders and acquired or traumatic brain injury are excluded. We will include studies where participants have comorbid substance misuse difficulties, but we will exclude trials where participants have a primary diagnosis of substance misuse disorder.

# *Intervention(s), exposure(s)*

We will include trials where participants in at least one arm receive 'bona fide' psychological interventions (defined according to criteria developed by Benish et al (2008),\* delivered in group or individual format, including but not limited to CBT, interpersonal therapy, psychodynamic therapy, EMDR or psychoeducation. \*The bona fide definition (Benish, Imel and Wampold, 2008) requires that treatments had to be delivered by a trained therapist who adapted the treatment to patients on the basis of a therapeutic relationship (i.e., no delivery of a non-modifiable standard protocol, e.g., progressive muscle relaxation); treatments also needed to be conducted personally and face-to-face (i.e., no online treatments or treatments conducted with, e.g., audio material). Moreover, at least two of the following four criteria had to be fulfilled with regards to their descriptions in the studies: (a) a citation to an established school or approach to psychotherapy; (b) a description of the therapy that contained a reference to a psychological process (e.g., operant conditioning); (c) a reference to a treatment manual that was used to guide the delivery of the treatment; (d) the identification of active ingredients of the treatment and citations for these ingredients.

\*Benish SG, Imel ZE, Wampold BE. The relative efficacy of bona fide psychotherapies for treating posttraumatic stress disorder: A meta-analysis of direct comparisons. Clin Psychol Rev. 2008;28:746–58. doi:10.1016/j.cpr.2007.10.005.

# *Comparator(s)/control*

Psychological interventions will be compared against one another and also against no additional treatment (i.e., 'treatment as usual' and 'waiting list control') and non-active interventions, such as befriending.

# Context

All settings to be included.

# *Primary outcome(s)*

The primary outcome will be twofold: 1. The between group difference, at end of treatment and 12-months post-randomisation, in severity of (a) PTSD symptoms as per ICD-11 and DSM III and IV and (b) emotion dysregulation, negative self-concept and/or interpersonal disturbance. To compute this composite outcome, we will calculate the average standardised mean difference across these outcomes taking into account the correlation between these variables where / if possible. 2. The between group difference at end of treatment and at 12-months post randomisation, in the relative and absolute risk of not achieving a clinically significant response in PTSD symptoms, defined using Jacobson criteria.

# Timing and effect measures

End of treatment and 12-months post randomisation (or nearest time-point within a 3-month range).

# Secondary outcome(s)

1. Safety, as measured by the between group difference, at end of treatment and at 12-months postrandomisation, in the relative risk of serious adverse events (death, suicide, attempted suicide, significant deterioration in symptoms, admission to hospital). 2. The acceptability of

the interventions, as measured by the between group difference in the relative risk of dropping out early from either the treatment or the trial (where the comparator is treatment as usual).

# *Timing and effect measures*

End of treatment and 12-months post randomisation (or nearest time-point within a 3-month range) for safety, and end of treatment for acceptability.

# Data extraction (selection and coding)

We will extract group means and associated standard deviations (and N contributing to those means) for continuous outcomes, and number of events (denominator = number randomised to arm) for dichotomous outcomes, using a spreadsheet. We will use the total number randomised if reported. For all outcomes except acceptability, we will assume those not including in the reported analyses are either missing completely at random (for continuous outcomes) or had no change from randomisation (dichotomous outcomes). Two researchers will double-extract data for all outcomes, and a third rater will be consulted in relation to any discrepancies and / or disagreements. If means and standard deviations are not reported, then we will estimate the between group difference from other statistical parameters, such as confidence intervals, standard errors, p-values, t-values or F-values, following procedures in the Cochrane Handbook, and using the Campbell Effect Size Calculator, if possible. If we need to combine data from 2 groups before entry into the meta-analysis, we will do so following the formulae specified in the Cochrane Handbook.

# Risk of bias (quality) assessment

At the study level the risk of bias will be assessed using the Cochrane collection Risk of Bias Tool (Higgins et al 2011). This involves categorising studies as having a low, high or unclear risk of bias in the areas of selection and allocation of participants, intervention concealment, attrition and reporting. The results of this assessment will be used to inform interpretation of reported effect sizes and overall conclusions. The quality of overall outcomes will be assessed using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) (Guyatt et al 2011).

# Strategy for data synthesis

For continuous outcomes, we will perform random-effects meta-analyses to compute an overall standardised mean difference and associated 95% confidence intervals, with Hedges's g adjustment. For dichotomous outcomes, we will also perform random-effects meta-analyses to compute an overall relative risk, an overall difference in absolute risk, and the number needed to treat for one to experience benefit / harm (computed as the inverse of the absolute risk).

# Analysis of subgroups or subsets

We will examine whether the results are moderated by the degree to which the sample population meet CPTSD criteria (whether the sample score within the clinical range on 1, 2 or 3 CPTD criteria at baseline). We will also examine the potential moderating role of quality parameters including rater blinding, attrition, random sequence generation description. We will also examine whether the effectiveness of psychological treatment for CPTSD is moderated by the following: - individual vs. group format - adult onset trauma vs.. childhood onset trauma vs. both - phased / staged interventions vs. non-phased / non-staged interventions.

Contact details for further information Aoife Bradley a.bradley@napier.ac.uk
Organisational affiliation of the review Edinburgh Napier University

Review team members and their organisational affiliations Professor Thanos Karatzias. Edinburgh Napier University Dr Mick Fleming. Edinburgh Napier University Ms Susan Roberts. Edinburgh Napier University Ms Aoife Bradley. Edinburgh Napier University Dr Claire Fyvie. Rivers Centre for Traumatic Stress Professor Jonathan Bisson. Cardiff University Dr Neil Roberts. Cardiff University Dr Neil Roberts. Cardiff University Dr Philip Hyland. National College of Ireland Professor Marylene Cloitre. New York University Dr Tobias Hecker. University of Bielefeld Professor Andreas Maercker. University of Zurich Dr Paul Hutton. Edinburgh Napier University

Anticipated or actual start date 16 January 2017

Anticipated completion date 16 January 2018

*Funding sources/sponsors* None

*Conflicts of interest* None known

*Language* English

*Country* Scotland

Stage of review Review\_Ongoing

Subject index terms status Subject indexing assigned by CRD

Subject index terms Humans; Psychotherapy; Stress Disorders, Post-Traumatic

*Date of registration in PROSPERO* 10 February 2017

Date of publication of this version 10 February 2017 Details of any existing review of the same topic by the same authors

## Stage of review at time of this submission

Stage	Started	Completed
Preliminary searches	Yes	No
Piloting of the study selection process	No	No
Formal screening of search results against eligibility criteria	No	No
Data extraction	No	No
Risk of bias (quality) assessment	No	No
Data analysis	No	No

Available from:

http://www.crd.york.ac.uk/PROSPERO/display\_record.php?ID=CRD42017055305

## **B.** Changes from Protocol

A subsequent change was our inclusion of some psychological interventions (e.g., mindfulness, yoga) which were not strictly 'bona fide' psychological interventions (Benish, Imel, & Wampold, 2008). We made this decision in the interests of completeness.

While we had planned to take into account the correlation between variables when computing the composite outcome, this was not possible due the range of measures used to assess the variables; instead we had to assume the correlation was zero.

Additional changes included abandoning our pre-specified moderator analysis of phased vs non-phased interventions due to insufficient data.

We were also unable to determine rates of clinically significant response according to our prespecified method (Jacobsen's criteria). We instead converted the SMDs to NNTs using the Furukawa approach, under 3 assumptions of the control event rate (10%, 50% and 22%, which is half the control event rate observed for PTSD). Secondary outcomes (safety, drop-out) and follow-up data will be reported separately.

## C. Excluded Studies

The following table (**Table C.1**) details studies or reports excluded after inspection of the fulltext report, or via correspondence with authors. Studies or reports excluded on basis of title or abstract alone are not detailed as these are too numerous and the vast majority were of different conditions or were otherwise unrelated to the review question.

Study reference	Reason for exclusion
Acarturk 2016	No useable CPTSD index
Acierno 2017	No relevant comparator
Akbarian 2015	No useable CPTSD index
Alghamdi 2015	Sample not suitable
Arntz 2007	Baseline CPTSD index/indices not clinically significant
Asukai 2010	No useable CPTSD index
Badura-Brack 2018	No relevant psychological intervention
Bass 2016	Sample not suitable
Beidel 2017	Not RCT
Belleau 2017	No useable CPTSD index
Betancourt 2014	Sample not suitable
Bichescu 2007	No useable CPTSD index
Blanchard 2003	Baseline CPTSD index/indices not clinically significant
Boden 2012	No useable CPTSD index
Bohus 2013	No useable CPTSD index
Bormann 2013	No useable CPTSD index
Bormann 2014	No useable CPTSD index
Bradley 2003	Sample not suitable
Bremner 2017	Baseline CPTSD index/indices not clinically significant
Brom 1989	Baseline CPTSD index/indices not clinically significant
Brunet 2013	Sample not suitable
Bryan 2016	No useable CPTSD index
Bryant 2003	No useable CPTSD index
Bryant 2008	No useable CPTSD index
Buhmann 2016	Baseline CPTSD index/indices not clinically significant
Carlson 1998	No useable CPTSD index
Castillo 2016	No useable CPTSD index
Catani 2009	Sample not suitable
Chard 2005	No useable CPTSD index
Classen 2001	Sample not suitable
Cloitre 2010	Baseline CPTSD index/indices not clinically significant
Cloitre 2012	Study has overlapping sample with another study
Coffey 2016	No useable CPTSD index
Cook 2010	No useable CPTSD index
Cooper 1989	No useable CPTSD index
Cooper 2017	No relevant comparator
Cottraux 2008	No useable CPTSD index
de Bont 2016	No useable CPTSD index
Devilly 1998	No useable CPTSD index

Study reference	Reason for exclusion
Devilly 1999	No useable CPTSD index
Echeburua 1997	No useable CPTSD index
Edmond 1999	Sample not suitable
Edmond 2004	Study has overlapping sample with another study
Ertl 2011	Baseline CPTSD index/indices not clinically significant
Fecteau 1999	No useable CPTSD index
Feeny 2002	Study has overlapping sample with another study
Feske 2008	Unable to obtain relevant norms
Foa 1991	No useable CPTSD index
Foa 2004	Study has overlapping sample with another study
Foa 2013	No useable CPTSD index
Foa 2017	No useable CPTSD index
Fredman 2016	No relevant psychological intervention
Gamito 2010	No useable CPTSD index
Gersons 2000	No useable CPTSD index
Gilboa-Schechtman	Sample not suitable
2010	
Glynn 1999	No useable CPTSD index
Goldstein 2018	No relevant psychological intervention
Gutner 2016	Study has overlapping sample with another study
Hien 2004	No useable CPTSD index
Hien 2009	No useable CPTSD index
Hien 2017	Baseline CPTSD index/indices not clinically significant
Hien 2017 (b)	Study has overlapping sample with another study
Hinton 2005	No useable CPTSD index
Holliday 2014	Study has overlapping sample with another study
Holliday 2015	Study has overlapping sample with another study
Ironson 2002	No useable CPTSD index
Jacob 2014	No useable CPTSD index
Jensen 1994	No useable CPTSD index
Jindani 2015	Baseline CPTSD index/indices not clinically significant
Johnson 2011	No useable CPTSD index
Johnson 2016	Not RCT
Kearney 2013	Baseline CPTSD index/indices not clinically significant
Kearney 2016	Baseline CPTSD index/indices not clinically significant
Kip 2014	Study has overlapping sample with another study
Konig 2016	Study has overlapping sample with another study
Kruse 2009	Not RCT
Lange 2003	No relevant psychological intervention
Lee 2002	No useable CPTSD index
Levi 2016	Not RCT
Liedl 2011	Retracted
Litz 2007	No relevant psychological intervention
Lovell 2011	Study has overlapping sample with another study
Macdonald 2016	No relevant psychological intervention

Study reference	Reason for exclusion
Marcus 1997	No useable CPTSD index
Markowitz 2015	Baseline CPTSD index/indices not clinically significant
Markowitz 2017	Study has overlapping sample with another study
Maxwell 2016	No useable CPTSD index
McGovern 2015	No useable CPTSD index
McLay 2017	No useable CPTSD index
McLean 2014	No useable CPTSD index
Meier 2015	No useable CPTSD index
Mills 2012	No useable CPTSD index
Monson 2012	No relevant psychological intervention
Moradi 2014	No useable CPTSD index
Morath 2014	No useable CPTSD index
Morland 2014	No relevant psychological intervention
Moser 2010	Study has overlapping sample with another study
Nacasch 2011	No useable CPTSD index
Nacasch 2015	No relevant comparator
Neuner 2004	No useable CPTSD index
Neuner 2008	No useable CPTSD index
Neuner 2010	No useable CPTSD index
Nijdam 2018	No useable CPTSD index
Nosen 2014	No useable CPTSD index
Oktedalen 2015	Baseline CPTSD index/indices not clinically significant
Pabst 2014	No useable CPTSD index
Paivio 2010	No relevant comparator
Paunovic 2001	No useable CPTSD index
Peniston 1991	No useable CPTSD index
Polusny 2015	Baseline CPTSD index/indices not clinically significant
Possemato 2016	Sample not suitable
Pruiksma 2016	Study has overlapping sample with another study
Reger 2016	No useable CPTSD index
Resick 2003	Study has overlapping sample with another study
Resick 2008	No relevant comparator
Resick 2015	No useable CPTSD index
Resick 2017	No relevant comparator
Roberts 2016	Not RCT
Rothbaum 1997	No useable CPTSD index
Rothbaum 2005	No useable CPTSD index
Rothbaum 2006	No useable CPTSD index
Rothbaum 2014	No relevant psychological intervention
Ruglass 2017	No useable CPTSD index
Sack 2016	No relevant comparator
Sannibale 2013	No useable CPTSD index
Sautter 2015	No relevant psychological intervention
Schaal 2009	Baseline CPTSD index/indices not clinically significant
Schneier 2012	No relevant psychological intervention

Study reference	Reason for exclusion
Schnurr 2003	No useable CPTSD index
Schnurr 2007	No useable CPTSD index
Schnyder 2011	No useable CPTSD index
Scott 2017	Study has overlapping sample with another study
Shea 2013	Sample not suitable
Shnaider 2017	No relevant psychological intervention
Sikkema 2007	No useable CPTSD index
Sin 2017	Not RCT
Slade 2017	Study has overlapping sample with another study
Sloan 2012	No useable CPTSD index
Spence 2011	No relevant psychological intervention
Steinert 2017	No useable CPTSD index
Study Ref	Reason for Exclusion
Tarrier 1999 (a)	No useable CPTSD index
Tarrier 1999 (b)	No useable CPTSD index
Taylor 2003	Baseline CPTSD index/indices not clinically significant
van den Berg 2016	Study has overlapping sample with another study
van der Kolk 2007	No relevant comparator
van der Kolk 2016	No relevant psychological intervention
van Emmerik 2008	No useable CPTSD index
Vaughan 1994	No useable CPTSD index
Wahbeh 2016	Baseline CPTSD index/indices not clinically significant
Wells 2015	No useable CPTSD index
Wilson 1995	No useable CPTSD index
Wilson 1997	Study has overlapping sample with another study
Wolf 2015	Not RCT
Zang 2013	No useable CPTSD index
Zang 2014	No useable CPTSD index
Zang 2017	No useable CPTSD index
Zlotnick 1997	No useable CPTSD index
Zlotnick 2009	No useable CPTSD index
Zoellner 1999	Study has overlapping sample with another study
Zoellner 2017	No relevant psychological intervention

D.	Table D.1. Summary of Characteristics of the 51 Included Studies
----	--

Study Ref	Groups included	N	Country	Participants	Age, mean (SD)	% female	Duration & N sessions available	Drops outs N (%)	Trauma exposure	Type of Trauma	Trauma onset	Treatment setting
Ahmadi 2015	EMDR	16	Iran	Military servicemen	29.9 (7.8)	0	Unclear	5 (31.3)	Single	Non- sexual	Adulthood	Community
	REM	16					Unclear	6 (37.5)				
	Control	16						4 (25.0)				
Azad marzabadi 2014	Mindfulness	14	Iran	War victims with PTSD	Not reported	0	90mins, 8	2 (14.3)	Single	Non- sexual	Adulthood	Community
	Control	14						2 (14.3)				
Basoglu 2007	SSBT	16	Turkey	Earthquake survivors	34.0 (11)	87	60 mins, 1	1 (6.3)	Single	Non- sexual	Adulthood	Community
	RA	15						0(0)				
Beidel 2011	TMT	18	USA	Combat Veterans	58.93(N R)	0	90 mins, 29	4 (22.2)	Single	Non- sexual	Adulthood	Community
	EXP	17			59.76(N R)		90 mins, 29	1 (5.9)				
Beidel 2019	TMT	49	USA	Military veterans	37.67 (8.51)	7	90 mins, 29	14 (28)	Single	Non- sexual	Adulthood	Community
	EXP	43			33.26 (11.31)		90 mins, 29	22 (50)				
1	1		1		1							

Study Ref	Groups included	N	Country	Participants	Age, mean (SD)	% female	Duration & N sessions available	Drops outs N (%)	Trauma exposure	Type of Trauma	Trauma onset	Treatment setting
Bryant 2013	Support/CB T	34	Australia	Adult civilian patients	41.15 (12.92)	54	90 mins, 12	13 (38.2)	Single	Non- sexual	Adulthood	Community
	Skills/CBT	36			37.86 (12.70)		90 mins, 12	3 (8.3)				
Butollo 2016	DET	74	German y	Trauma survivors	37.99 (12.1)	66	90 mins,max.24	9 (12.2)	Multiple	Sexual and non-sexual	Adulthood	Community
	СРТ	67			33.67 (10.3)		90 mins, max.24	11(14.9)				
Cloitre 2002	STAIR+MP E	31	USA	CSA survivors	34 (7.22)	100	60 -90 mins, 16	9 (29)	Single & Multiple	Sexual & Non-sexual	Childhood	Community
	MA WL	27					15 mins,12	3 (11)				
Difede 2007	CBT	15	USA	Disaster workers	45.77 (7.72)	NR	75mins ,12	8 (53.3)	Single	Non- sexual	Adulthood	Community
	TAU	16						2(12.5)				
Dorrepaal 2012	EXP	38	Netherla nds	CSA survivors	40.3 (10.7)	NR	120mins, 20	7(18.4)	Multiple	Sexual and non-sexual	Childhood & Adulthood	Community
	TAU	33			37.1 (10.3)			5 (15.1)				
Duffy 2007	СТ	29	Ireland	Trauma survivors	NR	NR	NR ,12	9 (31.0)	Multiple	Non- sexual	Adulthood	Community
	WL	29						3 (10.3)				
Dunn 2007	SMT	51	USA	Veterans	54.7 (6.9)	0	90 mins, 14	17(33.3)	Single	Non-sexual	Adulthood	Community

Study Ref	Groups included	N	Country	Participants	Age, mean (SD)	% female	Duration & N sessions available	Drops outs N (%)	Trauma exposure	Type of Trauma	Trauma onset	Treatment setting
	PGT	50			55.0 (7.6)		90 mins, 14	6(12.0)				
Dunne 2012	TF-CBT	13	Australia	MVA survivors	32.54 (7.09)	50	60mins,10	1 (7.7)	Single	Non-sexual	Adulthood	Community
	WL	13						2(15.4)				
Ehlers 2003	СТ	28	UK	MVA survivors	NR	NR	60-90mins, 12	0 (0)	Single	Non- sexual	Adulthood	Community
	SHB	28					40mins,1	3 (10.7)				
	RA	29					20mins, 1	2 (6.9)				
Ehlers 2005	СТ	14	UK	PTSD patients	35.4 (10.9)	53.6	60-90mins, 4-20	0 (0)	Single	Non-sexual	Adulthood	Community
	WL	14			37.8 (11.2)			0 (0)				
Ehlers 2014	Intensive CT	30	UK	Chronic PTSD	39.7 (12.4)	58.7	18hrs over 5-7 days	1(3.3)	Multiple	Sexual and /or non- sexual	Childhood & Adulthood	Community
	Weekly CT	31			41.5 (11.7)		100mins,12	1(3.2)				
	Weekly ST	30			37.8 (9.9)		100mins ,12	3 (10)				
	WL	30			36.8 (10.5)			0 (0)				
Foa 1999	PE	25	USA	Chronic PTSD	34.9 (10.6)	100	90-120mins , 9	2 (8)	Single	Sexual or non-sexual	Childhood or Adulthood	Community
	SIT	26					90-120mins, 9	7(27)				
	PE-SIT	30					90-120mins, 9	8 (27)				
	WL	15						0 (0)				

Study Ref	Groups included	N	Country	Participants	Age, mean (SD)	% female	Duration & N sessions available	Drops outs N (%)	Trauma exposure	Type of Trauma	Trauma onset	Treatment setting
Foa 2005	PE	79	USA	Assault survivors	31.3 (9.8)	100	90-120mins, 9-12	27 (34.1)	Single	Sexual or non-sexual	Childhood or Adulthood	Community or University - based
	PE-CR	74					90-120mins, 9-12	30(40.5)				
	WL	26						1 (3.8)				
Forbes 2012	СРТ	30	Australia	Military veteran	53.13 (13.97)	3.4	60-90 mins, 12	9(30.0)	Single	Non-sexual	Adulthood	Community
	TAU	29			53.62 (13.33)			9 (31.1)				
Ford 2011	TARGET	48	USA	Mothers with PTSD	30.7(6.9)	100	50mins, 12	12(25)	Unclear	Unclear	Adulthood	Community
	РСТ	53					NR, 12	14 (26)				
	WL	45										
Galovski 2012	МСРТ	53	USA	Trauma survivors	39.80 (11.74)	69	NR, 12	14 (26.4)	Single	Sexual or non-sexual	Childhood or Adulthood	Community
	SMDT	47					NR12,	7 (14.9)				
Ghafoori 2017	PE	24	USA	Trauma survivors	35.2 (12.0)	83.1	60-90 mins, 12	34 (72)	Multiple	Sexual and /or non- sexual	Adulthood	Community
	РСТ	47					60-90 mins,12	16 (66)				
Harned 2014	DBT	9	USA	Women with BPS & PTSD	32.6 (12.0)	100	1 year of treatment	4(44.4)	Single	Sexual or non-sexual	Childhood or Adulthood	Community
	DBT -PE	17					1 year of treatment	7(41.2)				

Study Ref	Groups included	N	Country	Participants	Age, mean (SD)	% female	Duration & N sessions available	Drops outs N (%)	Trauma exposure	Type of Trauma	Trauma onset	Treatment setting
Hinton 2009	IT-CBT	12	USA	Refugees	49.92 (9.23)	60	NR, 12	0 (0)	Single	Non sexual	Adulthood	Community
	DT- CBT	12			49.08 (7.56)		NR,12	0 (0)				
Hinton 2011	CA-CBT	12	USA	Female Latino patients	47.6 (8.2)	100	60 mins, 14	0	Unclear	Unclear	Unclear	Community
	AMR	12			51.4 (5.9)		60 mins, 14	0				
Hogberg 2007	EMDR	13	Sweden	Public transportation workers	43 (8)	20.8	90mins, 5	0(0)	Single	Non sexual	Adulthood	Community
	WL	11			43 (11)			2(18.2)				
Hollified 2007	CBT	28	USA	Adults with PTSD	40.9 (13.4)		120mins,12	7 (25)	Multiple	Unclear	Childhood or Adulthood	Community
	WL	27			43.4 (13.5)			6 (22.2)				
Jung 2013	CRIM	17	German v	CSA survivors	37.18 (10.85)	100	50 & 90mins, 2	0 (0)	Multiple	Sexual	Childhood	Community
	WL	17						0 (0)				
Keane 1989	Implosive (flooding)	11	USA	Veterans	34.7 (4.3)	0	90 minutes,14-16	1 (9.1)	Single	Non-sexual	Adulthood	Community
	WL	13			34.5 (2.1)			1(7.7)				

Study Ref	Groups included	N	Country	Participants	Age, mean (SD)	% female	Duration & N sessions available	Drops outs N (%)	Trauma exposure	Type of Trauma	Trauma onset	Treatment setting
Kip 2013	ART	US A	29	Veterans	41.0 (12.4)	20	60-75mins, 2-5	3 (10.3)	Single	Sexual or non-sexual	Adulthood	Community
	AC		28				60mins, 2	4 (14.3)				
Krakow 2001	CIT	88	USA	Sexual Assault survivors	40 (11.2) C 37 (12.7) NC	100	60-180mins , 3	22(25)	Multiple	Sexual	Childhood & Adulthood	Community
	WL	80			36 (9.3) C 31 (10.5) NC			0 (25) 2				
Krupnick 2008	IPT	32	USA	Trauma survivors	32 (10.2)	100	120 mins, 16	NR*	Multiple	Sexual & or non-sexual	Childhood & Adulthood	Community
	WL	16						NR*				
Kubany 2003	Immediate CTT-BW	19	USA	Battered women	36.4 (9.1)	100	90 mins,8-11	1 (5.3)	Multiple	Non-sexual	Adulthood	Community
	Delayed CTT-BW	18					90 mins, 8-11	4 (22.2)				
Kubany 2004	Immediate CTT-BW	63	USA	Battered women	42.2 (10.1)	100	90 mins,8-11	18 (28.6)	Multiple	Non-sexual	Adulthood	Community
	Delayed CTT-BW	62					90 mins, 8-11	22 (35.5)				
Lindauer 2005	BEP	12	Netherla nds	Trauma survivors	37.6 (10.2)	54	45-60 mins, 16	3(25)	Multiple	Non-sexual	Adulthood	Community

Study Ref	Groups included	N	Country	Participants	Age, mean (SD)	% female	Duration & N sessions available	Drops outs N (%)	Trauma exposure	Type of Trauma	Trauma onset	Treatment setting
	WL	12			40.3 (8.9)							
Marks 1998	Exposure	23	UK	Outpatients with PTSD	39 (11)	36	90 mins,10	3(13)	Single	Sexual or non-sexual	Adulthood	Community
	Cognitive	19			39 (9)		90 mins,10	1(5.3)				
	E+C	24			38 (9)		105 mins,10	5(20.8)				
	Relaxation	21			36 (10)		90 mins,10	1(4.8)				
McDonagh 2005	СВТ	29	USA	CSA survivors	39.8 (9.9)	100	90-120mins, 14	12(41)	Multiple	Sexual	Childhood	Community
	РСТ	22			39.6 (9.6)		90-120mins, 14	2(9)				
	WL	23			42.0 (9.8)			3(13)				
Monson 2006	СРТ	30	USA	Veterans	54.0 (6.3)	10	2p/w, 12	6(20)	Single	Sexual or Non-sexual	Adulthood	Community
	WL	30						4(13)				
Mueser 2008	CBT	54	USA	Severe Mental Illness patients	44.21 (10.64)	79	NR,12-16	19	Single	Sexual or non-sexual	Childhood & Adulthood	Community
	TAU	54										
Mueser 2015	СВТ	104	USA	Severe Mental Illness patients	42.96 (10.46)	72.3	NR,12- 16	37 (35.6)	Single	Sexual or non-sexual	Childhood & Adulthood	Community
	BT	97			44.52 (11.60)		NR, 3	14 (14.3)				

Study Ref	Groups included	N	Country	Participants	Age, mean (SD)	% female	Duration & N sessions available	Drops outs N (%)	Trauma exposure	Type of Trauma	Trauma onset	Treatment setting
Nijdam 2012	BEP	70	Netherla nds	Trauma survivors	37.3 (10.6)	56.4	45-60 mins,15	25 (36)	Single	Sexual or non-sexual	Adulthood	Community
	EMDR	70			38.3 (12.2)		90 mins, NR	20 (29)				
Pacella 2012	PE	40	USA	Adults with HIV	46.37 (6.30)	36.9	90-120 mins, 10	17 (42.5)	Multiple	Sexual or non-sexual	Childhood & Adulthood	Community
	WL	24						0 (0)				
Power 2002	EMDR	39	UK	Adults with PTSD	38.6 (11.8)	41.7	90 mins,10	12 (31)	Single	Sexual or non-sexual	Adulthood	Community
	E+CR	37			43.2 (11.0)		90 mins ,10	16 (43)				
	WL	29			36.5 (11.6)			5 (17)				
Resick 2002	СРТ	41	USA	Female Rape Victims	32 (9.9)	100	2 p/w 13hrs,12	11(26.8)	Multiple	Sexual & /or Non- sexual	Adulthood	Community
	PE	40					2 p/w 13hrs, 9	(27.3)				
	MA	40						(14.9)				
Scheck 1998	EMDR	30	USA	Traumatised Young Women	20.93(no SD reported)	100	90mins, 2	0 (0)	Single	Sexual	Childhood & Adulthood	Community
	AL	30					90mins, 2	1 (3.3)				
Steel 2017	CBT	30	UK	Adults with schizophrenia	42.3 (10.2)	37.7	NR ,12-16	4(13.0)	Single or Multiple	Sexual or non-sexual	Childhood & Adulthood	Community
	TAU	31						5 (16.1)				

Study Ref	Groups included	N	Country	Participants	Age, mean (SD)	% female	Duration & N sessions available	Drops outs N (%)	Trauma exposure	Type of Trauma	Trauma onset	Treatment setting
Suris 2013	СРТ	72	USA	Veterans	46.1	84.9	Unclear . 12	25(35)	Single	Sexual	Adulthood	Community
					(9.8)		,		0			5
	РСТ	57					Unclear, 10-12	10(18)				
Talbot 2014	CBT-I	29	USA	Adults with PTSD	37.1 (10.4)	68.9	Unclear	2 (6.9)	Unclear	Unclear	Unclear	Community
	WL	16			37.3(11. 0)			1 (6.3)				
ter Heide 2011	EMDR	10	Netherla nds	Asylum seekers and refugees	40.00 (9.31)	40	90 mins , 11	5 (50)	Multiple	Non-sexual	Adulthood	Community
	Stabilisation	10			43.00 (7.93)		60 mins, 11	5(50)				
ter Heide 2016	EMDR	37	Netherla nds	Refugees	43.1(10. 7)	27.8	60-90mins, 9	6 (16.7)	Multiple	Non-sexual	Adulthood	Community
	Stabilisation	37			39.8(11. 9)		60 mins , 12	8 (22.2)				
van den Berg 2015	PE	53	Netherla nds	Severe Mental Illness patients	41.2 (10.5)	54.2	90 mins, 8	13(24.5)	Single or Multiple	Sexual &/or Non-sexual	Childhood & Adulthood	Community
	EMDR	55					90 mins, 8	11(20.0)				
	WL	47										

Abbreviations: AC, Attention Control; AL, Active Listening Control; AMR, Applied Muscle Relaxation; ART, Accelerated Resolution Therapy; BEP, Brief Eclectic Psychotherapy; BPS, Borderline Personality disorder; BT, Brief Treatment; CA-CBT, Culturally Adapted Cognitive Behaviour Therapy; CBT, Cognitive Behaviour Therapy; CBT-I, Cognitive Behavioral Therapy for Insomnia; CIT, Cognitive Imagery Treatment; CPT, Cognitive Processing Therapy; CRIM, Cognitive Restructuring and Imagery Modification; CSA, Childhood Sexual Abuse; CT, Cognitive Therapy; CTT-BW, Cognitive Trauma Therapy for Battered Women; DBT, Dialectical Behavior Therapy; DBT PE, Dialectical Behavior Therapy; DET, Dialogical Exposure Therapy; DT –CBT, Delayed Treatment Cognitive Behaviour Therapy;E+C, Exposure and Cognitive; E+CR, Exposure plus Cognitive Restructuring; EMDR, Eye Movement Desensitization and Reprocessing Therapy; EXP, Experimental Treatment; EXP, Exposure

Therapy Only; Intensive CT, Intensive Cognitive Therapy; IPT, Interpersonal Psychotherapy; MA, Minimal Attention; MA WL, Minimal Attention Wait List; MCPT, Modified Cognitive Processing Therapy Intervention; MVA, Motor Vehicle Accident; PCT, Present Centred Therapy; PE, Prolonged exposure; PE-CR, Prolonged Exposure plus Cognitive Restructuring; PE-SIT, Prolonged Exposure + Stress Inoculation Training; PGT, Psychoeducational Group Therapy; PTSD, Post-traumatic Stress Disorder; RA, Repeated Assessments; REM, Rapid Eye Movement; SHB, Self-help booklet; SIT, Stress inoculation training; SMDT, Symptom –Monitoring Delayed Treatment; SMT, Self-Management Therapy; SSBT, Single Session of Behavioural Treatment; STAIRS + MPE, Skills Training in Affective and Interpersonal Regulation with modified Prolonged Exposure; TARGET, Trauma Affect Regulation: Guide for Education and Therapy; TAU, Treatment as usual; TFCBT, Trauma-Focused Cognitive-Behavioural Therapy; TMT, Trauma Management Therapy; Weekly CT, Weekly Cognitive Therapy; Weekly ST, Weekly SU, Waitlist.

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
Ahmadi 2015	EMDR REM	Emotional control subscale of the Mississippi Scale for PTSD, -5.3 (4.4) [these are the mean (SD) change scores] -6.4 (3.9) [these are the mean (SD) change	-	-	-	Interpersonal relation subscale of the Mississippi Scale for PTSD, - 4.7 (5.2)	_	Only change scores were reported (no baseline scores).	Include – although exclude in sensitivity analysis.
	Control	0.7 (2.5) [these are the mean (SD) change scores]		-		-0.08 (2.3)			
Azad marzabadi 2014	Mindfulness	-	-	-	-	Social life subscale of the WHOQOL-26, 5.5 (1.65)	-	Baseline scores indicate that the participants were on	Include
	Control	-		-		5.21 (0.97)		average at least dissatisfied in their social life.	
			1	1	1			1	1

## E. Table E.1. Participants' baseline scores on the CPTSD symptom clusters and corresponding norms

Basaglu 2007 SS	SBT				mean (SD)	(SD)	mean (SD)	interpretation	
RA	A A	-	-	-	-	WSAS, 4.1 (0.8) 4.1 (0.9)	-	Baseline scores above 4 on the WSAS suggest moderately severe or worse psychopatholo- gy (Mundt et al., 2002).	Include
Beidel 2011 TN	EXP	CAPS social and emotional subscale, 22.6 (5.3) 22.4 (3.8)	-	-	-	CAPS social and emotional subscale, 22.6 (5.3) 22.4 (3.8)	-	Using the "1, 2" rule (i.e., a frequency score of 1 and an intensity score of 2) to determine symptom severity, scores above 12 on this subscale meet the clinical threshold (Weathers, Ruscio, & Keane, 1999).	Include

Beidel 2019 TMT - - - Duration of Daily Social Interaction (outside of family interactions at home) (mins per day), 49.7 (54.3) - On averag non-clinic group age in 63.49 h	e, a Include 1 16– lgage of
EXP - 52.7 (61.9) activity per week, and activity let below 30 l indicative poor socia functionin (Hodgekir al., 2015).	els are of g s et
Bryant 2013 Support/CBT PTCI-self, 4.08 Median (SD) (1.29) among people	Include
Skills/CBT   -   4.41 (1.18)   with no trauma: 1.08 (0.76) (Foa et al., 1999)   -	
Destalla 2016 DET DET DECLastific 2.71 Madian (SD) UD C 1.24 (0.50) 1.29 (0.52) (non	Lashada NGC
Butono 2010DE1PTC1-sen, 5./1Median (SD)IIP-C, 1.34 (0.59)1.28 (0.52) (hoh-CPT(1.2) [these are the mean (SD)among people with no trauma: across both groups]1.08 (0.76) (Foa et al., 1999)	data but not DR data.
Cloitra 2002 STAID+MDE NMD 85 (15.6)	

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
	MA WL	84 (17.9)	101.6 (15.43) (non-clinical) (Cantanzaro & Mearns, 1990)	-		1.70 (0.46) (the weighted mean across these groups is 1.79)	1.28 (0.52) (non- clinical) (Brahler et al., 1999)		Include AD data but not DR data.
Difede 2007	CBT TAU	-	-	-	-	SAS-SR, 2 (0.4) 2.28 (0.44)	1.59 (0.33) (non- clinical) (Weissman et al., 1978)	-	Include
Dorrepaal 2012	EXP TAU	-	-	-	-	-	-	As all participants in this study had to meet diagnostic criteria for complex PTSD as assessed by the SIDES, this study is relevant.	Include
Duffy 2007	СТ	  -	-	-	-		  -		Include

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
	WL	-		-		SDS-social subscale, 7.7 (2.4) [these are the mean (SD) across both groups]		A score of 7 or above on this subscale indicates a marked impairment. Moreover, it has been suggested that clinicians should pay special attention to patients who score 5 or greater on this subscale (Rush et al., 2000).	
Dunn 2007	SMT PGT	SCQD, 86.14 (12.95) 79.74 (17.77)	101.2 (15.46) (non-clinical) (Mezo &	-	-	-	-	-	Include
Dunne 2012	TF-CBT		Heiby, 2004) -		-	SF-36 social functioning subscale, 43.46 (18.12)	85.66 (19.83) (among male norms aged 25- 44): 88 54	-	Include

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
	WL	-		-		45.42 (13.97)	(18.09) (among male norms aged 35-55); numerous other norms are available as well (Ware et al., 1993).		
Ehlang 2002	CT					SDS = 5.0 (2.4)		A	Inchede
Enters 2003		-	-	-	-	5D5, 5.9(2.4)		A score of 4 to 6 on this scale	Include
	RA	-		-		6.1 (1.9)		indicates a moderate impairment. Moreover, it has been suggested that clinicians should pay special attention to patients who score 5 or greater on this subscale (Rush et al., 2000).	
Ehlers 2005	СТ	-	-	-	-	SDS, 7.6 (1.9)	-		Include

Study Ref	Groups included	Name of AD assessment,	Normative AD data for	Name of NSC assessment,	Normative NSC data for	Name of DR assessment,	Normative DR data for	Other information	Decision
		(SD)	mean (SD)	(SD)	mean (SD)	(SD)	mean (SD)	interpretation	
	WL	-		-		6.7 (1.9)		A score of 7 or above on this scale indicates a marked impairment. Moreover, it has been suggested that clinicians should pay special attention to patients who score 5 or greater on this subscale (Rush et al., 2000).	
Ehlers 2014	Intensive CT	_	_	-		SDS 20.48	_	Each of the	Include
	Weekly CT Weekly ST WL	-		-	- - -	(5.55) 21.39 (5.11) 19.65 (6.97) 17.28 (7.74)		baseline mean SDS scores need to be divided by 3 so they are comparable to those of Ehlers, 2003 and 2005 above (e.g., 21.39/3 = 7.13).	
Foa 1999	PE	-	-	-	-	SAS (interview version), 3.73 (0.83)	-	Normative data for the interview	Include

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
	SIT PE-SIT WL	- -		-		3.79 (1.23) <u>4 (1.11)</u> 3.93 (1.16)		version of the SAS were not available. This version of the SAS is a 7- point scale. Assuming a 0-6 scoring method, scores of 3 or greater indicate that participants are closer to being impaired than intact.	
Foa 2005	PE PE-CR	-	-	-	-	SAS social subscale (interview version), 4 (0.9) 3.9 (1)	-	Normative data for the interview version of the SAS social	Include

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
	WL	-		-		3.9 (1.2)		subscale were not available. This version of the SAS social subscale is a 7- point scale. Assuming a 0-6 scoring method, scores of 3 or greater indicate that participants are closer to being impaired than intact.	
Forbes 2012	СРТ	-	-	-	-		-		Include

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
	TAU	-		-		Social subscale of the WHO- QOL Bref, 8.1 (2.8) [this is the mean (SD) across both groups]		Normative data for this subscale were not available. If a participant were to answer "neither satisfied or dissatisfied" on 2 of the items and "dissatisfied" on the other item they would receive a score of 8. Therefore, a score of 8.1 indicates that participants are closer to being impaired than intact.	
Ford 2011	WL	NMR, 96.9 (20)	101.6 (15.43) (non-clinical) (Cantanzaro & Mearns, 1990)	PTCI-self, 67.1 (28.3) [A mean of 67.1 is equivalent to a mean of 3.2 when scored the same way as the normative data]	Median (SD) among people with no trauma: 1.08 (0.76) (Foa et al., 1999)	RSQ secure attachment subscale, 13.5 (3.3)	15.57 (SD = 3.01) (non- clinical) (Bäackström & Holmes, 2001)	-	Include NSC data but not AD or DR data.

Study Ref	Groups included	Name of AD assessment,	Normative AD data for	Name of NSC assessment,	Normative NSC data for	Name of DR assessment,	Normative DR data for	Other information	Decision
		baseline mean (SD)	interpretation, mean (SD)	baseline mean (SD)	interpretation, mean (SD)	baseline mean (SD)	interpretation, mean (SD)	for interpretation	
	TARGET	106.1 (18.1)		51.3 (23.5) [A mean of 51.3 is equivalent to a mean of 2.44 when scored the same way as the normative data]		13.7 (3.8)			
	РСТ	103.1 (20.2)		53.7 (25.4) [A mean of 53.7 is equivalent to a mean of 2.56 when scored the same way as the normative data]		14 (3.5)			
Galovski 2012	MCPT SMDT	-	-	TRGI guilt cognitions subscale, 1.57 (0.11) [this is the least square mean (SE)] 1.62 (0.12) [this	1 (0.5) (among participants with a history of potentially traumatic CSA/CPA without any	SF-36 social functioning subscale, 42.87 (4.06) [this is the least square mean (SE)] 37.45 (4.29) [this	85.66 (19.83) (among male norms aged 25- 44); 88.54 (18.09) (among male norms aged 35-55); numerous	-	Include both NSC and DR data
				is the least square mean (SE)]	axis-I disorder; these are more severe than healthy individuals) (Rausch et al., 2016)	is the least square mean (SE)]	other norms are available as well (Ware et al., 1993).		
Ghafoori 2017	PE	-	-	-	-	SDS social	-	A score of 7 or	Include
						subscale, 7 (2.6)		above on this	menuae

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
	PCT	-		-		7.3 (2.5)		subscale indicates a marked impairment. Moreover, it has been suggested that clinicians should pay special attention to patients who score 5 or greater on this subscale (Rush et al., 2000).	
Harned 2014	DBT DBT -PE	-	-	TRGI guilt cognitions subscale, 2.4 (0.9) 2.4 (0.8)	1 (0.5) (among participants with a history of potentially traumatic CSA/CPA without any axis-I disorder;	-	-	-	Include
Hinton 2009	IT-CBT	ERS, 0.9 (0.6)	-	-	these are more severe than healthy individuals) (Rausch et al., 2016)	-	_		Include

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
	DT-CBT	0.8 (0.5)						Normative data for this scale were not available. This scale is rated on a 0-4 Likert- type scale, rating the ability to distance from affects, ranging from "not at all" to "very much so." These scores appear to indicate that participants are only edging towards somewhat being able to distance from affects.	
Hinton 2011	CA-CBT	ERS, 0.7 (0.5)	-	-	-	-	-		Include

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
	AMR	0.9 (0.4)		-		-		As above with Hinton, 2009, these scores appear to indicate that participants are only edging towards somewhat being able to distance from affects.	
									+
Hogberg 2007	EMDR	-	-	-	-	SDI, 4.5 (2.3)	1.53 (1.13)	-	Include
	WL	-		-		5.9 (4.5)	(among subjects who had experienced traumatic events but who had never developed PTSD) (Nardo et al., 2011)		
Hall: 6 ad 2007	CDT					CDC alshal active		Normating 1-t-	Tradada
Hollinea 2007	CBI	-	-	-	-	scale, $4.09(0.81)$	-	for this scale	Include

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
	WL	-		-		4 (1.02)		were not available. Participants' scores on this scale appear to be impaired as possible scores on this scale appear to range from 0 to 5, with higher scores indicating greater impairment.	
Jung 2013	CRIM WL	-	-	RSES, 22.1 (7.8) 20.6 (5.7)	49.2 (8.2) (non- clinical) (Roth et al., 2008)	-	-	This version of the RSES appears to range from 1- 60.	Include
Keane 1989	Implosive (flooding)	-	-	-	-	Social subscale of the Social Adjustment Measures, 4 (1.9)	-	Normative data for this scale were not available. The	Include

	included	assessment, baseline mean (SD)	data for interpretation, mean (SD)	assessment, baseline mean (SD)	NSC data for interpretation, mean (SD)	assessment, baseline mean (SD)	data for interpretation, mean (SD)	information for interpretation	Decision
	WL	-		-		2.6 (1.7)		weighted mean of participants' scores on this scale indicate that participants are closer to being extremely dissatisfied than extremely satisfied with their social life.	
Kip 2013	ART	-		TRGI guilt cognitions subscale, 26.7 (no SD reported) 20.2 (no SD reported)	21 (10.5) (among participants with a history of potentially traumatic CSA/CPA without any axis-I disorder; these are more severe than healthy individuals) (the mean and SD were rescaled in light of the scoring method of Kip, 2013) (Rausch et al., 2016)	Relating to others subscale of the PTGI, 11.6 (7.92; SD imputed from Nijdam, 2018) 13 (7.92; SD imputed from Nijdam, 2018)	23.04 (no SD reported) (non- clinical) (Tedeschi & Calhoun, 1996)	-	Include DR data but not NSC data.

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
Krakow 2001	CIT WL	-	-	-	-	SDS social life/leisure activities index (no baseline scores were reported)	-	Inferential statistics including effect sizes showing changes in the SDS social life/leisure activities index in the groups were reported (no baseline scores).	Include – although exclude in sensitivity analysis.
Krupnick 2008	IPT WL	-	-	-	-	IIP	-	Patients with a score of 3 or higher on any item of the IIP (indicating significant interpersonal distress) qualified for participation in	Include
Kubany 2003	Immediate CTT-BW	-	-	RSES, 13.6 (5.2)	22.62 (5.80) (non-clinical)	-	-	It is also worth noting that an	Include

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
	Delayed CTT- BW	-		12.7 (6.7)	(Sinclair et al., 2010)			inclusion criterion in this study was a score on the TRGI global guilt scale reflecting at least moderate abuse-related guilt.	
Kubany 2004	Immediate CTT-BW Delayed CTT- BW	-	-	RSES, 14.8 (5.4) 14.5 (4.5)	22.62 (5.80) (non-clinical) (Sinclair et al., 2010)	-	-	It is also worth noting that an inclusion criterion in this study was a score on the TRGI global guilt scale reflecting at least moderate abuse-related	Include
							+	guilt.	
Lindauer 2005	REP	+	+	<u> </u>	+	Relationships	<u>+</u>	At least 50% of	Include
Ellidadel 2005	DEI	-		-		Questionnaire,	-	the sample had	menude

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
	WL	-		-		50% of participants across groups had problems with relationships (binary measure, not a continuous measure; therefore no means or SDs were reported)		problems with relationships.	
Marks 1998	Exposure	-	-	-	-	WSAS, 21.5	-	Baseline scores	Include
	Cognitive E+C Relaxation		-		-	(8.9) 26.9 (8.8) 29.4 (7.9) 22.1 (9.5)		above 20 on the WSAS suggest moderately severe or worse psychopatholo- gy (Mundt et al., 2002).	
McDonagh	CBT	-	-	TSL 3 1 (0.6)	1 83 (0 34)	-	-	-	Include
2005	PCT	-		2.9 (0.5)	(non-clinical)	-			menude
	WL	-		3.2 (0.6)	(the mean and SD were rescaled in light of the scoring method of McDonagh, 2014) (Kadambi & Truscott, 2004)	-			
Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
-------------	--------------------	--	--	---	---	--	--	---	--
Monson 2006	CPT WL	ACS	-	TRGI guilt cognitions subscale	-	SAS-SR, 2.48 (0.39) 2.68 (0.54)	1.59 (0.33) (non- clinical) (Weissman et al., 1978)	-	Include (exclude AD and NSC data in sensitivity analysis, as baseline AD and NC data were not available)
Mueser 2008	CBT TAU	-	-	PTCI-self, 3.89 (1.11) 3.64 (1.14)	Median (SD) among people with no trauma:	-	-	-	Include
					1.08 (0.76) (Foa et al., 1999)				
Mueser 2015	CBT	-	-	PTCI-self, 4.10 (1.36)	Median (SD) among people with no trauma: 1.08 (0.76)	CAPS social functioning subscale, 2.35 (0.79)	-	Using the "1, 2" rule (i.e., a frequency score of 1 and an	Include NSC data but not DR data.

Study Ref	Groups included	Name of AD assessment, baseline mean	Normative AD data for interpretation,	Name of NSC assessment, baseline mean	Normative NSC data for interpretation,	Name of DR assessment, baseline mean	Normative DR data for interpretation,	Other information for	Decision
		(SD)	mean (SD)	(SD)	mean (SD)	(SD)	mean (SD)	interpretation	
	BT	-		4.15 (1.31)	(Foa et al., 1999)	2.36 (0.81)		intensity score of 2) for the CAPS social functioning subscale to determine symptom severity, scores above 3 meet the clinical threshold (Weathers et al., 1999)	
Nijdam 2012	BEP	-	-	-	-	Relating to others subscale of the PTGI, 14.38 (7.92) [this is the mean (SD) across	23.04 (no SD reported) (non- clinical) (Tedeschi & Calhoun, 1996)	-	-
	EMDR	-		-		both groups]			
<b>D</b>									
Pacella 2012	PE	-	-	PTCI-self, 3.23 (1.19)	Median (SD) among people	-	-	-	Include
	WL	-		3.04 (1.38)	with no trauma: 1.08 (0.76) (Foa et al., 1999)	-			
Power 2002	EMDR E+CR	-	-	-	-	SDS, 21.3 (5.4) 22.8 (6.3)	-	Each of the baseline mean	Include

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
	WL	-		-		23.3 (4.7)		SDS scores need to be divided by 3 so they are comparable to those of Ehlers, 2003 and 2005 above (e.g., 21.3/3 = 7.1).	
				TD CL 1 1 1 1					x 1 1
Resick 2002	СРТ	-		subscale, 2.34 (1.13)	(among participants	-	-	-	Include
	PE	-		2.53 (1.11)	with a history	-			
	МА	-		2.60 (1.03)	or potentially traumatic CSA/CPA without any axis-I disorder; these are more severe than healthy individuals) (Rausch et al., 2016)	-			
<u>Calcarla</u> 1000	EMDD			TOCO 294 57	245 75 (29 72)				T., . 1 1 .
Scheck 1998	EMDK	-		1 SCS, 284.57 (40.94)	(non-clinical)	-		-	Include
	AL	-		285.24 (38.23)	(Caplan, Henderson, Henderson, & Fleming, 2002)	-			

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
Steel 2017	CBT TAU	-	-	PTCI-self, 4.46 (1.13) [this is the mean (SD) across both groups]	Median (SD) among people with no trauma: 1.08 (0.76) (Foa et al., 1999)	-	-	-	Include
Suris 2013	CPT PCT	-	-	PTCI-self, 4.80 (1.12) 4.82 (1.25)	Median (SD) among people with no trauma: 1.08 (0.76) (Foa et al., 1999)	-	-	-	Include
Talbot 2014	CBT-I WL	-	-	-	-	WSAS, mean of >24 for both groups (as depicted in a graph)	-	Baseline scores above 20 on the WSAS suggest moderately severe or worse psychopatholo- gy (Mundt et al., 2002).	Include
ter Heide 2011	EMDR	-	-	-	-	Social subscale of the WHO- QOL Bref, 2.4 (0.86)	-	Normative data for this subscale were not available. A	Include

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
	Stabilisation	-		-		3.07 (0.49)		weighted mean of less than 3 represents being between dissatisfied (2) and neither satisfied nor dissatisfied (3; which is in the middle) on the different items of this subscale. Therefore, participants are closer to being impaired than intact.	
ter Heide 2016	EMDR	-	-	-	-	Social subscale of the WHO- QOL Bref, 2.71 (0.80)	-	Normative data for this subscale were not available. A	Include

Study Ref	Groups included	Name of AD assessment, baseline mean (SD)	Normative AD data for interpretation, mean (SD)	Name of NSC assessment, baseline mean (SD)	Normative NSC data for interpretation, mean (SD)	Name of DR assessment, baseline mean (SD)	Normative DR data for interpretation, mean (SD)	Other information for interpretation	Decision
	Stabilisation	-		-		2.55 (0.98)		weighted mean of less than 3 represents being between dissatisfied (2) and neither satisfied nor dissatisfied (3; which is in the middle) on the different items of this subscale. Therefore, participants are closer to being impaired than intact.	
van den Berg	PE	-	-	PTCI-self, 4.52	Median (SD)	-	-	-	Include
2015	EMDR WL		-	(1.22) 4.4 (1.12) 4.26 (0.96)	among people with no trauma: 1.08 (0.76) (Foa et al., 1999)				
					1999)				

Abbreviations: AC, Attention Control; ACF, Affect Control Scale; AL, Active Listening Control; AMR, Applied Muscle Relaxation; ART, Accelerated Resolution Therapy; BEP, Brief Eclectic Psychotherapy; BPS, Borderline Personality disorder; BT, Brief Treatment; CA-CBT, Culturally Adapted Cognitive Behaviour Therapy; CAPS, Clinician-Administered PTSD Scale; CBT, Cognitive Behaviour Therapy; CBT-I, Cognitive Behavioral Therapy for Insomnia; CIT, Cognitive Imagery Treatment; CPT, Cognitive Processing Therapy; CRIM, Cognitive Restructuring and Imagery Modification; CSA, Childhood Sexual Abuse; CT, Cognitive Therapy; CTT-BW, Cognitive Trauma Therapy for Battered Women; DBT, Dialectical Behavior Therapy; DBT PE, Dialectical Behavior Therapy Prolonged Exposure; DET, Dialogical Exposure Therapy; DT –CBT, Delayed Treatment Cognitive Behaviour Therapy; ERS, Emotion Regulation Scale; EXP, Experimental Treatment; EXP, Exposure Therapy Only; Intensive CT, Intensive Cognitive Therapy; IIP, Inventory of Interpersonal Problems; IIP-C, Inventory of Interpersonal Problems – Circumplex Version; IPT, Interpersonal Psychotherapy; MA, Minimal Attention; MA WL,

Minimal Attention Wait List; MCPT, Modified Cognitive Processing Therapy Intervention; MVA, Motor Vehicle Accident; NMR, General Expectancy for Negative Mood Regulation Scale; PCT, Present Centred Therapy; PE, Prolonged exposure; PE-CR, Prolonged Exposure plus Cognitive Restructuring; PE-SIT, Prolonged Exposure + Stress Inoculation Training; PGT, Psychoeducational Group Therapy; PTCI, Posttraumatic Cognitions Inventory; PTGI, Post-Traumatic Growth Inventory; PTSD, Post-traumatic Stress Disorder; RA, Repeated Assessments; REM, Rapid Eye Movement; RSES, Rosenberg Self-Esteem Scale; SAS, Social Adjustment Scale; SAS-SR, Social Adjustment Scale-Self-Report; SCQD, Self-Control Questionnaire for Depression; SDI, Social Disability Index; SDS, Sheehan Disability Scale; SF-36, Short Form-36 Health Survey; SHB, Self-help booklet; SIDES, Structured Interview for Disorders of Extreme Stress; SIT, Stress inoculation training; SMDT, Symptom –Monitoring Delayed Treatment; SMT, Self- Management Therapy; SSBT, Single Session of Behavioural Treatment; STAIRS + MPE, Skills Training in Affective and Interpersonal Regulation with modified Prolonged Exposure; TARGET, Trauma Affect Regulation: Guide for Education and Therapy; TAU, Treatment as usual; TFCBT, Trauma-Focused Cognitive-Behavioural Therapy; TMT, Trauma Management Therapy; TRGI, Trauma Related Guilt Inventory; TSCS, Tennessee Self-Concept Scale; TSI, Traumatic Stress Institute Beliefs Scale; Weekly CT, Weekly Cognitive Therapy; Weekly ST, Weekly Supportive Therapy ;WL, Waitlist; WHOQOL-26, World Health Organization Quality of Life Questionnaire; WHO-QOL Bref, The short form of the World Health Organization Quality of Life scale; WSAS, Work and Social Adjustment Scale.

# References for comparator samples or scoring guidelines referred to in Table E.1 above

- Bäckström, M., & Holmes, B. M. (2001). Measuring adult attachment: a construct validation of two self-report instruments. *Scandinavian Journal of Psychology*, 42(1), 79–86. <u>https://doi.org/10.1111/1467-9450.00216</u>
- Brahler, E., Horowitz, L. M., Kordy, H., Schumacher, J., & Strauss, B. (1999). Validation of the Inventory of Interpersonal Problems (IIP)--Results of a representative study in East and West Germany. *Psychotherapie Psychosomatik Medizinische Psychologie*, 49(11), 422–431.
- Caplan, S. M., Henderson, C. E., Henderson, J., & Fleming, D. L. (2002). Socioemotional Factors Contributing to Adjustment Among Early-Entrance College Students. *Gifted Child Quarterly*, 46(2), 124–134. <u>https://doi.org/10.1177/001698620204600205</u>
- Catarizaro, S. J., & Mearns, J. (1990). Measuring Generalized Expectancies for Negative Mood Regulation: Initial Scale Development and Implications. *Journal of Personality Assessment*, 54(3–4), 546–563. <u>https://doi.org/10.1080/00223891.1990.9674019</u>
- Foa, E. B., Tolin, D. F., Ehlers, A., Clark, D. M., & Orsillo, S. M. (1999). The Posttraumatic Cognitions Inventory (PTCI): Development and validation. *Psychological Assessment*, 11(3), 303–314. <u>https://doi.org/10.1037/1040-3590.11.3.303</u>
- Hodgekins, J., French, P., Birchwood, M., Mugford, M., Christopher, R., Marshall, M., ... Fowler, D. (2015). Comparing time use in individuals at different stages of psychosis and a non-clinical comparison group. *Schizophrenia Research*, 161(2–3), 188–193. <u>https://doi.org/10.1016/j.schres.2014.12.011</u>
- Kadambi, M. a, & Truscott, D. (2004). Vicarious Trauma Among Therapists Working with Sexual Violence, Cancer, and General Practice. *Canadian Journal of Counselling*, 38, 260–276.
- Mezo, P. G., & Heiby, E. M. (2004). A comparison of four measures of self-control skills. Assessment, 11(3), 238–250. <u>https://doi.org/10.1177/1073191104268199</u>
- Mundt, J. C., Marks, I. M., Shear, M. K., & Greist, J. M. (2002). The Work and Social Adjustment Scale: a simple measure of impairment in functioning. *The British Journal of Psychiatry*, 180(5), 461–464.
- Nardo, D., Högberg, G., Flumeri, F., Jacobsson, H., Larsson, S. A., Hällström, T., & Pagani, M. (2011). Self-rating scales assessing subjective well-being and distress correlate with rCBF in PTSD-sensitive regions. *Psychological Medicine*, 41(12), 2549–2561. <u>https://doi.org/10.1017/S0033291711000912</u>
- Rausch, S., Herzog, J., Thome, J., Lud�scher, P., M�ller-Engelmann, M., Steil, R., ... Kleindienst, N. (2016). Women with exposure to childhood interpersonal violence without psychiatric diagnoses show no signs of impairment in general functioning, quality of life and sexuality. *Borderline Personality Disorder and Emotion Dysregulation*, 3(1). <u>https://doi.org/10.1186/s40479-016-0048-y</u>
- Roth, M., Decker, O., Herzberg, P. Y., & Brähler, E. (2008). Dimensionality and norms of the Rosenberg self-esteem scale in a German general population sample. *European Journal of Psychological Assessment*, 24(3), 190–197. <u>https://doi.org/10.1027/1015-5759.24.3.190</u>

- Rush, J.A. (2000). *American Psychiatric Association: Handbook of Psychiatric Measures*. Washington, DC, American Psychiatric Press.
- Sinclair, S. J., Blais, M. A., Gansler, D. A., Sandberg, E., Bistis, K., & LoCicero, A. (2010). Psychometric properties of the Rosenberg Self-Esteem Scale: Overall and across demographic groups living within the United States. *Evaluation and the Health Professions*, 33(1), 56–80. <u>https://doi.org/10.1177/0163278709356187</u>
- Tedeschi, R. G., & Calhoun, L. G. (1996). The posttraumatic growth inventory: Measuring the positive legacy of trauma. *Journal of Traumatic Stress*, 9(3), 455–471. https://doi.org/10.1002/jts.2490090305
- Ware, J. E., Snow, K. K., Kosinski, M., & Grandek, B. (1993). SF-36 health survey : manual and interpretation guide. Boston, MA: The Health Institute
- Weathers, F. W., Ruscio, A. M., & Keane, T. M. (1999). Psychometric properties of nine scoring rules for the Clinician-Administered Posttraumatic Stress Disorder Scale. *Psychological Assessment*, 11(2), 124–133. <u>https://doi.org/10.1037/1040-3590.11.2.124</u>
- Weissman, M. M., Prusoff, B. A., Douglas Thompson, W., Harding, P. S., & Myers, J. K. (1978). Social adjustment by self-report in a community sample and in psychiatric outpatients. *Journal of Nervous and Mental Disease*, 166(5), 317–326. <u>https://doi.org/10.1097/00005053-197805000-0000</u>

# F. Risk of Bias Criteria

#### Selection Bias: random sequence generation

A judgement of unclear risk of bias was made where randomisation was referred to but described in insufficient detail to determine independent random sequence generation. There was judged to be low risk of bias where this procedure was explicitly reported.

### Selection Bias: allocation concealment

A judgement of unclear risk of bias was made where randomisation was referred to but described in insufficient detail to determine allocation concealment. There was judged to be low risk of bias where this procedure was explicitly reported.

#### Performance Bias: blinding of participants and personnel

Blinding of participants and personnel was not possible due to the nature of the interventions, as is the case with trials of psychosocial interventions in general. This resulted in a judgement of high risk of performance bias across studies.

#### Detection Bias: blinding of assessments

Detection bias was judged to be high where non-blinding of assessors was stated or where no information was given, and low if blinding was explicitly reported.

### Attrition Bias: incomplete outcome data

A judgement of high risk of attrition bias was made where data for  $\geq 25\%$  of those randomised was missing (Xia et al., 2009) or if attrition was not reported (or clearly reported) and a completer analysis was carried out. If attrition was low (<25%) and completer analysis was used risk of attrition bias was rated as low.

#### Reporting Bias: selective outcome reporting

If outcomes are pre-specified and reported a low risk of reporting bias rating was given. However, if no protocol is reported a high risk of reporting bias rating was given. If subgroup analysis is reported but not pre-specified a high risk of reporting bias rating was given.

#### **Overall Quality**

An overall quality rating for each study was also produced. Performance bias was not taken into consideration when producing this rating, as blinding of participants was not possible due to the nature of the interventions. All of the other criteria above were considered (i.e., selection bias: random sequence generation and allocation concealment, detection bias, attrition bias, and reporting bias). A high overall quality rating was given if a study received a low risk of bias rating for detection bias, at least another low risk of bias rating, and  $\leq 2$  high risk of bias ratings. A low overall quality rating was given if a study did not meet these criteria.

Study	Selection Bias:	Selection Bias:	Performance Bias:	<b>Detection Bias:</b>	Attrition Bias:	Reporting Bias:	<b>Overall Quality</b>
	random	allocation	blinding of	blinding of	incomplete	selective outcome	
	sequence	concealment	participants and	assessments	outcome data	reporting	
	generation		personnel				
Ahmadi 2015	Unclear	Unclear	High risk	High risk	High risk	High risk	Low quality
Azad marzabadi	Unclear	Unclear	High risk	High risk	Low risk	High risk	Low quality
2014							
Basoglu 2007	Low risk	Unclear	High risk	Low risk	Low risk	High risk	High quality
Beidel 2011	Unclear	Unclear	High risk	High risk	High risk	High risk	Low quality
Beidel 2019	Unclear	Unclear	High risk	High risk	High risk	High risk	Low quality
Bryant 2013	Low risk	Low risk	High risk	Low risk	High risk	High risk	High quality
Butollo 2016	Unclear	Unclear	High risk	High risk	Low risk	High risk	Low quality
Cloitre 2002	Unclear	Unclear	High risk	Low risk	Low risk	High risk	High quality
Difede 2007	Unclear	Unclear	High risk	High risk	High risk	High risk	Low quality
Dorrepaal 2012	Unclear	Low risk	High risk	Low risk	Low risk	High risk	High quality
Duffy 2007	Unclear	Unclear	High risk	High risk	Low risk	High risk	Low quality
Dunn 2007	Low risk	High risk	High risk	Low risk	High risk	High risk	High quality
Dunne 2012	Unclear	Unclear	High risk	High risk	Low risk	High risk	Low quality
Ehlers 2003	Low risk	Low risk	High risk	Low risk	Low risk	High risk	High quality
Ehlers 2005	Unclear	Unclear	High risk	Low risk	Low risk	High risk	High quality
Ehlers 2014	Low risk	Low risk	High risk	Low risk	Low risk	High risk	High quality
Foa 1999	Unclear	Unclear	High risk	Low risk	Low risk	High risk	High quality
Foa 2005	Unclear	Unclear	High risk	Low risk	High risk	High risk	Low quality
Forbes 2012	Low risk	Low risk	High risk	Low risk	Low risk	High risk	High quality
Ford 2011	Low risk	Low risk	High risk	High risk	High risk	High risk	Low quality
Galovski 2012	Low risk	Unclear	High risk	Low risk	High risk	High risk	High quality
Ghafoori 2017	Low risk	Unclear	High risk	High risk	High risk	High risk	Low quality
Harned 2014	Low risk	Unclear	High risk	Low risk	High risk	High risk	High quality
Hinton 2009	Low risk	Unclear	High risk	Low risk	Low risk	High risk	High quality
Hinton 2011	Unclear	Unclear	High risk	High risk	Low risk	High risk	Low quality
Hogberg 2007	Unclear	Low risk	High risk	High risk	Low risk	High risk	Low quality
Hollified 2007	Low risk	Low risk	High risk	High risk	High risk	High risk	Low quality
Jung 2013	Unclear	Unclear	High risk	High risk	Low risk	High risk	Low quality
Keane 1989	Unclear	Unclear	High risk	High risk	Low risk	High risk	Low quality

# G. Table F.1. Risk of Bias in Included Studies – Summary

Study	Selection Bias:	Selection Bias:	<b>Performance Bias:</b>	<b>Detection Bias:</b>	Attrition Bias:	<b>Reporting Bias:</b>	Overall Quality
	random	allocation	blinding of	blinding of	incomplete	selective outcome	
	sequence	concealment	participants and	assessments	outcome data	reporting	
	generation		personnel				
Kip 2013	Low risk	Unclear	High risk	High risk	Low risk	Low risk	Low quality
Krakow 2001	Low risk	Unclear	High risk	Low risk	Low risk	High risk	High quality
Krupnick 2008	Unclear	Unclear	High risk	High risk	High risk	High risk	Low quality
Kubany 2003	Unclear	Unclear	High risk	Low risk	Low risk	High risk	High quality
Kubany 2004	Unclear	Unclear	High risk	Low risk	High risk	High risk	Low quality
Lindauer 2005	Low risk	Unclear	High risk	Low risk	High risk	High risk	High quality
Marks 1998	Unclear	Unclear	High risk	Low risk	Low risk	High risk	High quality
McDonagh 2005	Unclear	Unclear	High risk	Low risk	Low risk	High risk	High quality
Monson 2006	Unclear	Unclear	High risk	Low risk	Low risk	High risk	High quality
Mueser 2008	Low risk	Low risk	High risk	Low risk	High risk	High risk	High quality
Mueser 2015	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	High quality
Nijdam 2012	Low risk	Low risk	High risk	Low risk	High risk	Low risk	High quality
Pacella 2012	Low risk	Unclear	High risk	Low risk	Low risk	High risk	High quality
Power 2002	Low risk	Low risk	High risk	Low risk	High risk	High risk	High quality
Resick 2002	Unclear	Unclear	High risk	High risk	High risk	High risk	Low quality
Scheck 1998	Low risk	Low risk	High risk	Low risk	High risk	High risk	High quality
Steel 2017	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	High quality
Suris 2013	Low risk	Low risk	High risk	Low risk	High risk	High risk	High quality
Talbot 2014	Low risk	Low risk	High risk	Low risk	Low risk	High risk	Low quality
ter Heide 2011	Low risk	Unclear	High risk	High risk	High risk	High risk	High quality
ter Heide 2016	Low risk	Unclear	High risk	Low risk	Low risk	Low risk	High quality
van den Berg 2015	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	High quality

11. I able 11.1. Kisk of Dias in flictuded Studies – Detaile	H.	Table H.1	. Risk of	<b>Bias</b> in	Included	Studies -	Detailed
--	----	-----------	-----------	----------------	----------	-----------	----------

Study	Selection Bias: random sequence generation	Selection Bias: allocation concealment	Performance Bias: blinding of particinants and	Detection Bias: blinding of assessments	Attrition Bias: incomplete outcome data	Reporting Bias: selective outcome reporting
	generation	conceannent	personnel		outcome auta	
Ahmadi 2015	Randomisation was referred to, but there was no more relevant information.	Randomisation was referred to, but there was no more relevant information.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	Not reported.	"The drop-out rate from the study was also high comprising over 31% of the initial participants." No follow-up data – beyond post- intervention data.	Protocol not available.
	Unclear	Unclear	High risk	High risk	High risk	High risk
Azad marzabadi 2014	Randomisation was referred to, but there was no more relevant information.	Randomisation was referred to, but there was no more relevant information.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	Not reported.	"Two participants dropped out in each group as the study continued (due to reasons like being discharged from the hospital or stopping participation in the study)."	Protocol not available.
	Unclear	Unclear	High risk	High risk	Low risk	High risk
Basoglu 2007	"A computer- generated sequence of random numbers that ensured equal cell sizes and did not lead to allocation of more than two consecutive cases to the same experimental	Not reported.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	"The assessors were blind as to the participants' experimental condition at the week 4 and week 8 assessments."	Only four non- completer cases, and ITT was used.	Protocol not available.

Study	Selection Bias: random sequence generation	Selection Bias: allocation concealment	Performance Bias: blinding of participants and personnel	Detection Bias: blinding of assessments	Attrition Bias: incomplete outcome data	Reporting Bias: selective outcome reporting
	condition was used in the randomization."					
D 11 1 2011	Low risk	Unclear	High risk	Low risk	Low risk	High risk
Beidel 2011	referred to, but there was no more relevant information.	referred to, but there was no more relevant information.	is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	relevant assessments).	participant did not complete the intervention and then another 5 did not complete the relevant post- intervention assessments. ITT was used. 10/35 = 29%. No follow-up data – beyond post- intervention data.	Protocol not available.
	Unclear	Unclear	High risk	High risk	High risk	High risk
Beidel 2019	Randomisation was referred to, but there was no more relevant information.	Randomisation was referred to, but there was no more relevant information.	"Participants were randomized to either TMT or EXP prior to initiating treatment. However, clinicians and participants were blinded to group assignment until VRET and the mid-treatment assessment were completed."	Not reported.	"The overall dropout rate was 39%, consistent with other clinical trials examining treatment for combat-related PTSD (Reger et al., 2016; Resick et al., 2015). The dropout rate was 28% for TMT and 50% for EXP, which was not significantly different ( $\chi 2 =$ 2.14, df = 91, p < 0.14)."	Protocol not available.

Study	Selection Bias:	Selection Bias:	<b>Performance Bias:</b>	<b>Detection Bias:</b>	Attrition Bias:	<b>Reporting Bias: selective</b>
-	random sequence	allocation	blinding of	blinding of	incomplete	outcome reporting
	generation	concealment	participants and	assessments	outcome data	
			personnel			
	Unclear	Unclear	High risk	High risk	High risk	High risk
Bryant 2013	"Randomization was conducted by a process of minimization stratified on gender, trauma type and Clinician Administered PTSD Scale-2 (CAPS-2; Blake et al. 1995) total score. Participants were randomly assigned according to a random numbers system administered by an individual who was independent of the study and who worked at a site that was distant from the treatment centre."	Distance randomisation – "Participants were randomly assigned according to a random numbers system administered by an individual who was independent of the study and who worked at a site that was distant from the treatment centre."	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	"Post-treatment and 6-month follow-up assessments were conducted by independent clinicians who were unaware of the treatment condition of participants. Blindness was maintained by ensuring that clinicians who conducted assessments did not have access to (a) participant notes or (b) condition allocation of participants."	"Of the participants, 51 (73%) completed treatment and 32 (46%) completed the 6-month follow- up assessment."	Protocol not available.
	Low risk	Low risk	High risk	Low risk	High risk	High risk
Butollo 2016	Randomisation was referred to, but there was no more relevant information.	Randomisation was referred to, but there was no more relevant information.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	"The IES-R [36], a self-report instrument that measures the intensity of PTSD symptoms, was our primary outcome measure. It was administered by the therapist before each session as a process	"Drop-out rates at the posttreatment assessment were 12.2% for DET (4.1% of those allocated to DET did not start treatment, 8.1% dropped out of treatment) and	Protocol not available.

Study	Selection Bias: random sequence generation	Selection Bias: allocation concealment	Performance Bias: blinding of participants and	Detection Bias: blinding of assessments	Attrition Bias: incomplete outcome data	Reporting Bias: selective outcome reporting
			personnel	measure, as well as before and after treatment and at the follow-up."	14.9% for CPT (6.0% declined treatment after allocation, 9.0% dropped out of treatment). At the 6-month follow-up, study drop-out rates were markedly higher, increasing the overall study dropout to 47.3% in the DET and 37.3% in the CPT	
	Unclear	Unclear	High risk	High risk	condition."	High risk
Cloitre 2002	Randomisation was referred to, but there was no more relevant information.	Randomisation was referred to, but there was no more relevant information.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	Clinician raters of the CAPS (PTSD measure) were blind to treatment condition at pre- and posttreatment. No reference to any blinding (e.g., re scoring) the NMR and IIP.	"Of the 58 women who entered treatment, 12 dropped out: 9 from the active treatment (29%) and 3 from the wait list (11%)." This is <25% overall.	Protocol not available.
	Unclear	Unclear	High risk	Low risk	Low risk	High risk
Difede 2007	Randomisation was referred to, but there was no more relevant information.	Randomisation was referred to, but there was no more relevant information.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	Not reported.	"Our dropout rate of 40% was higher than the typical exposure therapy study for PTSD, where dropouts are	Protocol not available.

Study	Selection Bias: random sequence generation	Selection Bias: allocation concealment	Performance Bias: blinding of participants and personnel	Detection Bias: blinding of assessments	Attrition Bias: incomplete outcome data	Reporting Bias: selective outcome reporting
					reported in the 20% to 30% percent range (Bradley et al., 2005). " Also, >30% dropped out of the allocated intervention.	
	Unclear	Unclear	High risk	High risk	High risk	High risk
Dorrepaal 2012	"The randomization was performed independently on a 1: 1 basis, stratified per site, by a methodologist not involved in the study. Condition assignments were e- mailed to the group leader, who then informed the patient without informing the researchers or assessors." No more relevant information.	"The randomization was performed independently on a 1: 1 basis, stratified per site, by a methodologist not involved in the study. Condition assignments were e- mailed to the group leader, who then informed the patient without informing the researchers or assessors."	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	"The interviews were conducted by trained independent assessors, who were blind to the treatment condition and were audiotaped for use in supervision."	Attrition was 16%.	Protocol not available.
	Unclear	Low risk	High risk	Low risk	Low risk	High risk
Duffy 2007	Randomisation was referred to, but there was no more relevant information.	Randomisation was referred to, but there was no more relevant information.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	Not reported.	21% dropped out.	Protocol not available.
	Unclear	Unclear	High risk	High risk	Low risk	High risk

Study	Selection Bias: random sequence generation	Selection Bias: allocation concealment	Performance Bias: blinding of participants and	Detection Bias: blinding of assessments	Attrition Bias: incomplete outcome data	Reporting Bias: selective outcome reporting
Dunn 2007	"The study statistician (J. Souchek [J.So.]) provided randomization numbers and group assignments from a list generated by the PLAN procedure in SAS, version 6.11. We randomized in blocks of two, in the order of participants' enrollment, to facilitate equal participant numbers in each group."	The list of random numbers provided could suggest that investigators could possibly foresee assignments.	personnel   Blinding of participants   is generally not possible   due to the nature of   psychological trials.   Nothing in this trial to   suggest otherwise.	"Interviewers were blind to participants" therapy group assignments throughout the study."	30% of those allocated to the interventions did not complete them, and 41% of those allocated to the interventions were lost to follow-up at 12 months.	Protocol not available.
	Low risk	High risk	High risk	Low risk	High risk	High risk
Dunne 2012	Randomisation was referred to, but there was no more relevant information.	Randomisation was referred to, but there was no more relevant information.	"Participants in this study were also not blinded to condition."	"Although the use of the same assessor for the diagnostic interview for all participants at all 3 time points is a methodological strength, this also meant the assessor was not blinded to the treatment condition representing a potential bias in postassessment and follow-up assessment."	"A further strength of the study was the use of the intent-to- treat sample for data analyses, despite the relatively low attrition in this study (9% at 6-mo follow-up)."	Protocol not available.

Study	Selection Bias: random sequence	Selection Bias: allocation	Performance Bias: blinding of	Detection Bias: blinding of	Attrition Bias: incomplete	Reporting Bias: selective outcome reporting
	generation	conceaiment	participants and	assessments	outcome data	
	Unclear	Unclear	High risk	High risk	Low risk	High risk
Ehlers 2003	Reference to using the random permuted blocks within strata algorithm.	Investigators enrolling participants could not possibly foresee assignments as the allocation list was kept locked in a central office and the patient's allocation was only revealed three weeks later – following the self-monitoring assessment.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	Independent assessors, blind to treatment condition, administered the CAPS. No reference to how the SDS (which is a self-report measure) was scored.	Attrition was < 25%.	Protocol not available.
	Low risk	Low risk	High risk	Low risk	Low risk	High risk
Ehlers 2005	Randomisation was referred to, but there was no more relevant information.	Randomisation was referred to, but there was no more relevant information.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	Independent assessors, blind to treatment condition, administered the CAPS. No reference to how the SDS (which is a self-report measure) was scored.	No patient dropped out.	Protocol not available.
	Unclear	Unclear	High risk	Low risk	Low risk	High risk
Ehlers 2014	"The participants were then randomly allocated to one of the four trial conditions by an independent researcher who was not involved in assessing patients using the minimization	"The participants were then randomly allocated to one of the four trial conditions by an independent researcher who was not involved in assessing patients using the minimization procedure (15) to stratify for sex and	"Participants were not blind to the nature of the treatment, but care was taken to create similarly positive expectations in each treatment group by informing them that several psychological treatments were effective in PTSD and it	"The assessments of treatment outcome were conducted by independent evaluators without knowledge of the patient's treatment condition. Patients were asked not to reveal their group	Attrition was < 25%.	Protocol not available.

Study	Selection Bias: random sequence generation	Selection Bias: allocation concealment	Performance Bias: blinding of participants and personnel	Detection Bias: blinding of assessments	Attrition Bias: incomplete outcome data	Reporting Bias: selective outcome reporting
	procedure (15) to stratify for sex and severity of PTSD symptoms. The assessors determining the suitability of a patient for inclusion were not informed about the stratification variables and algorithm."	severity of PTSD symptoms. The assessors determining the suitability of a patient for inclusion were not informed about the stratification variables and algorithm."	was unknown which worked best, and by giving a detailed rationale for the treatment condition to which the patient was allocated."	assignment to the evaluators."		
	Low risk	Low risk	High risk	Low risk	Low risk	High risk
Foa 1999	Randomisation was referred to, but there was no more relevant information.	Randomisation was referred to, but there was no more relevant information.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	Assessors were unaware of treatment assignment.	17.7 dropped out altogether; although drop-out were not spread evenly throughout the groups – 8% dropped out of the PE group, 27% of the SIT group, 27% of the PE-SIT group, 0% of the WL group. ITT was used.	Protocol not available.
	Unclear	Unclear	High risk	Low risk	Low risk	High risk
Foa 2005	Randomisation was referred to, but there was no more relevant information.	Randomisation was referred to, but there was no more relevant information.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	"Independent evaluations were conducted at pretreatment and posttreatment and 3-, 6-, and 12-month posttreatment. All	"The overall dropout rate was 32.4% and was lower for WL (3.8%) than PE/CR (40.5%) and PE (34.2%)" These	Protocol not available.

Study	Selection Bias: random sequence generation	Selection Bias: allocation concealment	Performance Bias: blinding of participants and	Detection Bias: blinding of assessments	Attrition Bias: incomplete outcome data	Reporting Bias: selective outcome reporting
	generation	conceannent	personnel	ussessments	outcome untu	
			•	evaluations were conducted by trained doctoral or master's level CTSA clinicians who were blind to study condition."	were not available for post- intervention assessment.	
	Unclear	Unclear	High risk	Low risk	High risk	High risk
Forbes 2012	A random number ordered list was used.	After full assessment by an independent clinical assessor, participants were randomised by the project manager at an independent research centre.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	Assessors were blind to allocation and treatment.	20% did not complete the post- intervention assessment and 31% did not complete the 3 month follow-up (which is a short- term follow-up.	Protocol not available.
	Low risk	Low risk	High risk	Low risk	Low risk	High risk
Ford 2011	"One hundred forty six women (ages 18– 45; M=30.7, SD=6.9) completed the screening and baseline assessment and then were randomized (by a study assessor using numbers concealed in sealed envelopes previously prepared by a different study staff member using the Excel random number generator) to WL (N=45), TARGET	"One hundred forty six women (ages $18-45$ ; M=30.7, $SD=6.9$ ) completed the screening and baseline assessment and then were randomized (by a study assessor using numbers concealed in sealed envelopes previously prepared by a different study staff member using the Excel random number generator) to WL (N=45), TARGET	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	"All interviewers were blind to the experimental condition in baseline interviews, but due to technical difficulties they were not blind to experimental condition at posttherapy or follow- up interviews."	32% of those allocated to treatment did not complete the post- intervention assessment.	Protocol not available.

Study	Selection Bias: random sequence generation	Selection Bias: allocation concealment	Performance Bias: blinding of participants and personnel	Detection Bias: blinding of assessments	Attrition Bias: incomplete outcome data	Reporting Bias: selective outcome reporting
	(N=48), or PCT (N=53). "	(N=48), or PCT (N=53). "				
	Low risk	Low risk	High risk	High risk	High risk	High risk
Galovski 2012	"If eligible, participants were randomly assigned in a 1:1 ratio using computer generated simple randomization to MCPT or to SMDT following the pre- treatment assessment."	No more relevant information other than what is in the previous column.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	"Finally, because it is technically possible to remain PTSD positive with a score of 20 on the PDS and reporting bias can exist in the therapy situation (e.g., patient wants to please therapist), a blind rater conducted the CAPS to ensure that the participant no longer met criteria for PTSD." The other relevant measures were self-report, and no reference to any blind scoring of these.	27% of those randomised did not complete an assessment at the time of the completion of the CBT intervention.	Protocol not available.
	Low risk	Unclear	High risk	Low risk	High risk	High risk
Ghafoori 2017	A pre-determined, computer-generated, randomised list was used.	No more relevant information other than what is in the previous column.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	Outcome assessors were not blind to participant assignments.	More than half of the sample terminated prematurely before completion of the treatment, and follow-up assessment time points were included.	Protocol not available.
	Low risk	Unclear	High risk	High risk	High risk	High risk

Study	Selection Bias:	Selection Bias:	<b>Performance Bias:</b>	<b>Detection Bias:</b>	Attrition Bias:	<b>Reporting Bias: selective</b>
-	random sequence	allocation	blinding of	blinding of	incomplete	outcome reporting
	generation	concealment	participants and	assessments	outcome data	
			personnel			
Harned 2014	"A minimization	No more relevant	Blinding of participants	"All assessments were	"Completion rates	Protocol not available.
	randomization	information other than	is generally not possible	conducted by	for the one year of	
	procedure was used."	what is in the previous	due to the nature of	independent clinical	treatment did not	
		column.	psychological trials.	assessors who were	differ between	
			Nothing in this trial to	blind to treatment	conditions	
			suggest otherwise.	condition."	( <i>DBT</i> =55.6%, <i>DBT</i>	
					+DBT	
					PE=58.8%)."	
	Low risk	Unclear	High risk	Low risk	High risk	High risk
Hinton 2009	Coin was tossed.	No more relevant	Blinding of participants	Outcome assessor was	"All 24 randomized	Protocol not available.
		information other than	is generally not possible	blind to treatment	patients completed	
		what is in the previous	due to the nature of	condition.	the study, and there	
		column.	psychological trials.		were no missing	
			Nothing in this trial to		data.	
	T · 1		suggest otherwise.	T • 1	T • 1	*** * * *
II. ( 2011	Low risk	Unclear	High risk	Low risk	Low risk	High risk
Hinton 2011	Randomisation was	Randomisation was	Blinding of participants	All assessments were	"There was no	Protocol not available.
	referred to, but there	referred to, but there	is generally not possible	self-report, although it	missing aata.	
	was no more relevant	was no more relevant	due to the nature of	does not clarify		
	information.	information.	psychological trials.	whether or not these		
			Nothing in this that to	were scored blind.		
	Unaloon	Unaloan	Suggest other wise.	Ligh risk	I on wish	Uigh wish
Hagharg 2007	Victear No more relevant	"The nandomization	Dlinding of participants	Not reported	"Three subjects	Drotocol not available
Thogoerg 2007	information other	was done by picking a	is generally not possible	Not reported.	dronned out after	r rotocor not available.
	than what is in the	sealed ballot in the	due to the nature of		randomization but	
	than what is in the	seared barrot in the	due to the nature of			
	next column	presence of a research	nevehological trials		hotoro	
	next column.	presence of a research	psychological trials.		before treatment/WI. One	
	next column.	presence of a research nurse who coordinated the study and followed	psychological trials. Nothing in this trial to suggest otherwise		before treatment/WL. One of them had a	
	next column.	presence of a research nurse who coordinated the study and followed the participants	psychological trials. Nothing in this trial to suggest otherwise.		bejore treatment/WL. One of them had a strong aversive	
	next column.	presence of a research nurse who coordinated the study and followed the participants through all phases "	psychological trials. Nothing in this trial to suggest otherwise.		before treatment/WL. One of them had a strong aversive reaction to the	
	next column.	presence of a research nurse who coordinated the study and followed the participants through all phases."	psychological trials. Nothing in this trial to suggest otherwise.		before treatment/WL. One of them had a strong aversive reaction to the SPECT	
	next column.	presence of a research nurse who coordinated the study and followed the participants through all phases."	psychological trials. Nothing in this trial to suggest otherwise.		before treatment/WL. One of them had a strong aversive reaction to the SPECT examination and	

Study	Selection Bias:	Selection Bias:	Performance Bias:	<b>Detection Bias:</b>	Attrition Bias:	<b>Reporting Bias: selective</b>
v	random sequence	allocation	blinding of	blinding of	incomplete	outcome reporting
	generation	concealment	participants and	assessments	outcome data	
	0		personnel			
Hollified 2007	Unclear "Before enrolling participants, 90 study ID numbers were prerandomized to study group (acupuncture, CBT or WLC) by the research	Low risk "When a participant was enrolled, the RC opened the assignment program to reveal the participant's group assignment. This allocation procedure	personnel   High risk   Blinding of participants   is generally not possible   due to the nature of   psychological trials.   Nothing in this trial to   suggest otherwise.	High risk "The research coordinator collected, entered, and helped analyze the data. Although he was aware of participant group glocation at the	the examination. Two other subjects left the study because of difficulty with finding time for the study." $5/24 =$ 21%. Low risk "Eighty-four participants were randomized, 73 began the protocol, and 61 (73% of those randomized and 84% of those	High risk Protocol not available.
	WLC) by the research coordinator (RC) using a computerized random numbers procedure without restrictions."	allocation procedure was concealed from clinicians."	High risk	group allocation at the time he collected data, he did not assist participants in completing the self- rated assessments. It is possible, yet we think quite unlikely, that he could have systematically influenced participant reports."	and 84% of those who began the protocol (acupuncture 79% vs. CBT 84% vs. WLC 88%) completed treatment or wait- list assessments. End treatment and 3-month follow-up assessments were obtained for 60 and 58 participants, respectively." 24/84 = 29%. High risk	High risk
Jung 2012	Low risk	LOW risk	Hign risk Dlinding of participants	The DDS and DSES	High risk	Hign risk Drotocol not available
Jung 2015	referred to but there	referred to but there	is generally not negricipants	(the relevant	54 participants	FIOLOCOI NOL AVAIIADIE.
	referred to, but there	referred to, but there	is generally not possible	(the relevant	were randomly	

Study	Selection Bias:	Selection Bias:	Performance Bias:	<b>Detection Bias:</b>	Attrition Bias:	<b>Reporting Bias: selective</b>
-	random sequence	allocation	blinding of	blinding of	incomplete	outcome reporting
	generation	concealment	participants and	assessments	outcome data	
	-		personnel			
	was no more relevant	was no more relevant	due to the nature of	measures) were self-	assigned to either	
	information.	information.	psychological trials.	report, but there is no	the CRIM group (n	
			Nothing in this trial to	reference to these	= 17) or WL group	
			suggest otherwise.	being blindly scored.	(n = 17). Two	
					patients in each	
					condition decided	
					against treatment	
					after randomization	
					and were defined	
					as nonstarters.	
					Further, 2 patients	
					(1 in each	
					condition) were	
					excluded from the	
					study due to	
					protocol violations,	
					specifically,	
					because they had	
					received further	
					psychological	
					treatment while	
					participating in the	
					study. No patient	
					aroppea out of	
					18%.	
	Unclear	Unclear	High risk	High risk	Low risk	High risk
Keane 1989	Randomisation was	Randomisation was	Blinding of participants	"Post-test assessments	No reference to any	"The original design of
	referred to, but there	referred to, but there	is generally not possible	for the Wait-list	missing data at	this study involved random
	was no more relevant	was no more relevant	due to the nature of	Control were	post-assessment.	assignment of subjects to
	information.	information.	psychological trials.	conducted either by		the implosive therapy
			Nothing in this trial to	the therapist who		group, a stress
			suggest otherwise.	would then treat the		management group
				patient or, in the case		wherein subjects were
				of patients to be		taught behavioral skills to

Study	Selection Bias:	Selection Bias:	Performance Bias:	<b>Detection Bias:</b>	Attrition Bias:	<b>Reporting Bias: selective</b>
· ·	random sequence	allocation	blinding of	blinding of	incomplete	outcome reporting
	generation	concealment	participants and	assessments	outcome data	
			personnel			
				treated elsewhere, by		re- duce anxiety but
				an unsystematically		exposure to traumatic
				assigned therapist		memories was limited, and
				from the four in-		the waiting list control.
				volved in the study."		Only 5 subjects completed
						the stress management
						condition, and thus, these
						data are not included in
						the present manuscript."
	Unclear	Unclear	High risk	High risk	Low risk	High risk
Kip 2013	"Eligible service	No more relevant	Blinding of participants	"Random assignment	Attrition was	"The trial was registered
	members/veterans	information other than	is generally not possible	was unblinded; hence,	<25%.	with ClinicalTrials.gov
	were randomly	what is in the previous	due to the nature of	the potential existed		<i>(NCT01559688).</i> "The
	assigned to the ART	column.	psychological trials.	for over-reporting of		relevant PTSD measure
	or AC regimen in a		Nothing in this trial to	reductions in pain		was reported as the
	1:1 ratio using a		suggest otherwise.	with the ART		primary outcome in this
	random number			intervention."		protocol.
	generator and					
	variable blocking					
	scheme (blocks of 4,					
	6, and 8)."	TT 1	TT 1 · 1			
IZ 1 2001	Low risk	Unclear	High risk	High risk		Low risk
Krakow 2001	To mask treatment	No more relevant	Blinding of participants	To limit external	Of the 168	Protocol not available.
	assignment, patients	information other than	is generally not possible	bias, blinding	randomized	
	mailed back a	what is in the previous	due to the nature of	occurred at 3 points of	participants, 96	
	postcard after intake	column.	psychological trials.	data collection: (1) at	completed 3-month	
	to complete entry into		Nothing in this trial to	intake, group	Jollow-ups by mail,	
	the protocol. The		suggest otherwise.	assignment had not	ana 99 completea	
	posicara s time and			deen establishea; (2)	ine o-month Jollow-	
	aute were logged into			ai 5-monin jouow-up,	ups in person. In	
	a computer and			questionnaires were	ioial, 114	
	entered into a			mail: and (3) at 6	inuiviauais	
	list of numbers that			mail, and (5) al 0-	1 follow up and 77	
	usi of numbers that			monin jouow-up,	1 jouow-up, ana //	

Study	Selection Bias: random sequence	Selection Bias: allocation	Performance Bias: blinding of	Detection Bias: blinding of	Attrition Bias: incomplete	Reporting Bias: selective outcome reporting
	generation	concealment	participants and personnel	assessments	outcome data	
	randomly assigned participants to treatment and control groups. All numbers and group assignments were generated at the start of the protocol."			interviewers were unaware of group status."	participants completed both follow-ups."	
	Low risk	Unclear	High risk	Low risk	Low risk	High risk
Krupnick 2008	Randomisation was referred to, but there was no more relevant information.	Randomisation was referred to, but there was no more relevant information.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	Not reported.	"At termination, we obtained assessments for only 20 (out of 32) treatment and 7 (out of 16) control participants. These figures increased to 26 treatment and 10 control participants for the 4-month follow-up interview." Therefore, attrition at termination was >25%.	Protocol not available.
	Unclear	Unclear	High risk	High risk	High risk	High risk
Kubany 2003	Randomisation was referred to, but there was no more relevant information.	Randomisation was referred to, but there was no more relevant information.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	The assessors (of the CAPS; the PTSD measure) were blind to participants' condition assignments. The RSES was also used, but this is a self-report measure and there is	"Eighteen of 19 women assigned to the Immediate CTT-BW condition completed CTT- BW. Fourteen of 18 women assigned to the Delayed CTT-	Protocol not available.

Study	Selection Bias:	Selection Bias:	Performance Bias:	<b>Detection Bias:</b>	Attrition Bias:	<b>Reporting Bias: selective</b>
-	random sequence	allocation	blinding of	blinding of	incomplete	outcome reporting
	generation	concealment	participants and	assessments	outcome data	
	-		personnel			
				no reference to this	BW condition	
				being blindly scored.	completed CTT-	
					BW. Overall, 86%	
					of the 37 women	
					who started CTT-	
					BW(n=32)	
					completed	
					treatment." It	
					appears these	
					participants also	
					completed the	
					assessment at the	
					time of the	
					completion of	
					Immediate CTT-	
					BW condition;	
					therefore there was	
					<25% attrition. Re	
					follow-ups, "three-	
					month follow-up	
					data was obtained	
					for 78% of the	
					women who	
					completed	
					Immediate CTT-	
					BW ( $n = 14$ ) and	
					for 79% of the	
					women who	
					completed Delaved	
					CTT-BW (n = )	
					11). "	
	Unclear	Unclear	High risk	Low risk	Low risk	High risk
Kubany 2004	Randomisation was	Randomisation was	Blinding of participants	The assessors (of the	Posttreatment	Protocol not available.
	referred to, but there	referred to, but there	is generally not possible	CAPS; the PTSD	assessment data	
			due to the nature of	measure) were blind to	were only available	

Study	Selection Bias:	Selection Bias:	Performance Bias:	<b>Detection Bias:</b>	Attrition Bias:	<b>Reporting Bias: selective</b>
· ·	random sequence	allocation	blinding of	blinding of	incomplete	outcome reporting
	generation	concealment	participants and	assessments	outcome data	
			personnel			
	was no more relevant	was no more relevant	psychological trials.	participants' condition	for 84 of 125	
	information.	information.	Nothing in this trial to	assignments. The	randomised	
			suggest otherwise.	RSES was also used,	participants; there	
				but this is a self-report	attrition was >25%.	
				measure and there is		
				no reference to this		
				being blindly scored.		
	Unclear	Unclear	High risk	Low risk	High risk	High risk
Lindauer 2005	"A colleague who	No more relevant	Blinding of participants	"Each patient was	"In the per-	Protocol not available.
	had done no	information other than	is generally not possible	assessed by a	protocol analysis	
	assessments used a	what is in the previous	due to the nature of	researcher (R.J.L.L. or	(patients who	
	computer program to	column.	Nothing in this trial to	E.P.M.M.), who were blind to all patients'	completed the	
	randomly assign 12 patients to each		suggest otherwise	condition "	sample sizes were 7	
	condition in a block		suggest otherwise.	conulion.	(58%) for the	
	design "				treatment group	
	ucsign.				and 11 (92%) for	
					the waitlist group"	
					Attrition = $25\%$ .	
	Low risk	Unclear	High risk	Low risk	High risk	High risk
Marks 1998	Randomisation was	Randomisation was	Blinding of participants	"Assessors were kept	"Ten subjects	Protocol not available.
	referred to, but there	referred to, but there	is generally not possible	unaware of the	(11%) dropped out	
	was no more relevant	was no more relevant	due to the nature of	treatment condition."	but they did not	
	information.	information.	psychological trials.		differ significantly	
			Nothing in this trial to		by group."	
			suggest otherwise.			
	Unclear	Unclear	High risk	Low risk	Low risk	High risk
McDonagh	Randomisation was	Randomisation was	Blinding of participants	"A separate group of	"The dropout rate	Protocol not available.
2005	referred to, but there	referred to, but there	is generally not possible	female clinicians, who	for the study was	Moreover, the following
	was no more relevant	was no more relevant	due to the nature of	were blind to	23%, with a rate of	suggests a deviation from
	information.	information.	psychological trials.	treatment condition	41% (12 of 29) for	the protocol: "When it
			Nothing in this trial to	and who had no other	CBT, 9% (2 of 22)	became clear that the
			suggest otherwise.	role in the study	for PCT, and $13\%$	dropout rate was greater
1					(3 of 23) for WL."	<i>Jor CB1, we changed the</i>

Study	Selection Bias: random sequence generation	Selection Bias: allocation concealment	Performance Bias: blinding of participants and personnel	Detection Bias: blinding of assessments	Attrition Bias: incomplete outcome data	Reporting Bias: selective outcome reporting
				conducted the four CAPS interviews."		random assignment process to increase the chance of assignment to CBT."
	Unclear	Unclear	High risk	Low risk	Low risk	High risk
Monson 2006	Randomisation was referred to, but there was no more relevant information.	"The study biostatistician provided the participants' condition assignment to the study coordinator." This does not necessarily suggest allocation concealment.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	"The independent clinician assessors were blinded to condition assignment and participants were instructed to not disclose their condition assignment to them."	"The overall drop- out rate was 16.6% (20% from CPT, 13% from the wait- list condition)."	Protocol not available.
	Unclear	Unclear	High risk	Low risk	Low risk	High risk
Mueser 2008	"Randomization was conducted at a central location in the research center by a computer based randomization program with assignments not known in advance by either clinical or research staff"	"Randomization was conducted at a central location in the research center by a computer based randomization program with assignments not known in advance by either clinical or research staff"	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	"Assessments were conducted by blinded interviewers at baseline, following the 4-6 months treatment period for the CBT program, and 3- and 6-months later."	Only 59/108 were analysed at post- treatment.	Protocol not available.
	Low risk	Low risk	High risk	Low risk	High risk	High risk
Mueser 2015	"Participants were randomised to the CBT or brief groups via a computer program operated by an off-site data manager, with no	"Participants were randomised to the CBT or brief groups via a computer program operated by an off-site data manager, with no study personnel aware	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	"All interviewers were masked to treatment assignment."	20% of randomised participants were not analysed at post-treatment.	"All study procedures were approved by the Rutgers and Dartmouth Institutional Review Boards (trial registration: clinicaltrials.gov

Study	Selection Bias: random sequence generation	Selection Bias: allocation concealment	Performance Bias: blinding of participants and personnel	Detection Bias: blinding of assessments	Attrition Bias: incomplete outcome data	Reporting Bias: selective outcome reporting
	study personnel aware of assignments in advance."	of assignments in advance."				identifier: NCT00494650)."
Nijdam 2012	Low risk "Random assignment was done on a 1:1 basis by a computer program, with a weighted maximum of subscribing four times the same treatment in a row."	Low risk "To ensure masking of assessors, one psychologist who had no other engagement in the study, had access to the computer program, kept a log file of all random assignments and assigned the patients to the therapists, "	High risk Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	Low risk "Assessments were conducted by trained, independent, masked assessors who were master's level clinical psychologists or master's level psychology students supervised by an experienced clinical psychologist."	Low risk 32% of randomised participants were lost to the first post-assessment.	Low risk "Trial registration: Dutch Trial Register, number NTR46 and ISRCTN64872147."
	Low risk	Low risk	High risk	Low risk	High risk	Low risk
Pacella 2012	"The principal investigator (DLD) generated the allocation sequence using blocked randomization (4:3 ratio of experimental:control participants)."	No more relevant information other than what is in the previous column.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	"The graduate student conducting the assessments remained blind to group membership."	"At the post- intervention assessment, 23 participants were retained in the PE group (32% drop- out rate) and 24 participants were retained in the control group (0% drop-out rate)." It also says: "Unequal numbers of participants were assigned to each group, as it was anticipated that the PE group	Protocol not available.

Study	Selection Bias: random sequence generation	Selection Bias: allocation concealment	Performance Bias: blinding of participants and personnel	Detection Bias: blinding of assessments	Attrition Bias: incomplete outcome data	Reporting Bias: selective outcome reporting
					would have a higher dropout rate."	
Power 2002	<i>Low risk</i> <i>"Randomization was</i>	Unclear "Following completion	High risk Blinding of participants	<i>Low risk</i> <i>"Assessments pre- and</i>	"A total of 105	Protocol not available.
	by means of a predetermined schedule unbeknown to the assessors, therapists or patients."	of the entire initial assessment, for those patients who met entry criteria, the blind assessor then opened a sealed envelope that informed as to which group patients were to be allocated."	is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	post-treatment were conducted by two independent assessors respectively, who were blind to treatment conditions."	patients met entry criteria and were randomized to groups as follows: 39 to EMDR, 37 to ECCR and 29 to WL. Drop-out rates between these three groups were as follows, 12 (31%) from EMDR, 16 (43%) from E+CR and five (17%) from WL."	
	Low risk	Low risk	High risk	Low risk	High risk	High risk
Resick 2002	Randomisation was referred to, but there was no more relevant information.	Randomisation was referred to, but there was no more relevant information.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	Not reported.	Attrition was > 25%.	Protocol not available.
	Unclear	Unclear	High risk	High risk	High risk	High risk
Scheck 1998	"Envelopes filled with papers labeled either EMDR or AL were shuffled and numbered 1 though 100."	"During each interview, envelopes were opened consecutively to identify which therapy was to be assigned to the participant."	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	"During the post-test assessment, a trained volunteer who was blind to group assignment administered the	29% dropped out.	Protocol not available.

Study	Selection Bias: random sequence generation	Selection Bias: allocation concealment	Performance Bias: blinding of participants and	Detection Bias: blinding of assessments	Attrition Bias: incomplete outcome data	Reporting Bias: selective outcome reporting
	8		personnel			
				standardized instru- ments".		
	Low risk	Low risk	High risk	Low risk	High risk	High risk
Steel 2017	"Block randomization was conducted independently of the research team through the OpenCDMS database specifically developed for the study."	"Block randomization was conducted independently of the research team through the OpenCDMS database specifically developed for the study."	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	"Robust procedures were adopted to minimize the risk of interviewers being able to identify the group allocation of participants"	50/61 were analysed at post- treatment.	"The trial was given ethical approval by Berkshire Research Ethics Committee (SC/09/ H0505/85) and was registered as ISTCRN 67096137."
	Low risk	Low risk	High risk	Low risk	Low risk	Low risk
Suris 2013	"For the purpose of randomization, participants were assigned sequential PIN numbers as they entered the study. Blocks of random numbers were generated for each therapist, and were allocated to either CPT or PCT using a conditional statement."	"The random number sequence was maintained on an Excel spreadsheet, and as subjects' PINs were entered into the spreadsheet, the preassigned condition was revealed."	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	Assessors were blind to treatment condition.	"Excluding data from this therapist reduced the final sample to 86 participants from the original 129."	This study has a high risk of reporting bias as lots of participants were removed from the analysis due to low treatment fidelity of a certain clinician.
	Low risk	Low risk	High risk	Low risk	High risk	High risk
Talbot 2014	"Blind assignment was determined by a computer generated random allocation schedule operated by	"Group allocation was provided to the study coordinator in opaque, sealed envelopes that were opened by the study coordinator with	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	"Clinical interviewers and the polysomnography technician were blind to participants' treatment conditions	Only 16% in total were lost to follow- up.	Protocol not available.

Study	Selection Bias: random sequence generation	Selection Bias: allocation concealment	Performance Bias: blinding of participants and personnel	Detection Bias: blinding of assessments	Attrition Bias: incomplete outcome data	Reporting Bias: selective outcome reporting
	the study statistician."	the participant following the completion of baseline measures."		during both pretreatment and posttreatment administration and scoring."		
	Low risk	Low risk	High risk	Low risk	Low risk	High risk
ter Heide 2011	Coin was tossed.	No more relevant information.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	The relevant measures were self-report and there is no reference to these being blindly scored. Re the clinician administered measure – "blindness was maintained only in 70% of SCID interviews, thus threatening the reliability of clinician rated outcomes."	10/20 (50%) dropped out.	Protocol not available.
	Low risk	Unclear	High risk	High risk	High risk	High risk
ter Heide 2016	Coin was tossed.	No more relevant information.	Blinding of participants is generally not possible due to the nature of psychological trials. Nothing in this trial to suggest otherwise.	"Interviews were administered by trained Master's students in psychology who were kept masked to treatment condition by having limited access to participant data and by asking participants not to reveal treatment content."	Attrition was <25%.	"Trial registration: NARCIS (Dutch National Academic Research and Collaborations Information System) OND1324839; ISRCTN20310201."
	Low risk	Unclear	High risk	Low risk	Low risk	Low risk

Study	Selection Bias: random sequence	Selection Bias: allocation	Performance Bias: blinding of	Detection Bias: blinding of	Attrition Bias: incomplete	<b>Reporting Bias: selective</b> outcome reporting
	generation	concealment	participants and	assessments	outcome data	
			personnel			
van den Berg	"An independent	"An independent	Blinding of participants	Assessors were blind	Attrition was	"The trial design was
2015	randomization	randomization bureau	is generally not possible	to treatment	<25%.	approved by the medical
	bureau randomized	randomized the	due to the nature of	allocation.		ethics committee of the VU
	the treatment	treatment condition	psychological trials.			University Medical Center
	condition using	using stratified	Nothing in this trial to			and was registered at
	stratified	randomization blocks	suggest otherwise.			isrctn.com
	randomization blocks	per therapist with				(ISRCTN79584912)."
	per therapist with	equal strata sizes."				
	equal strata sizes."	-				
	Low risk	Low risk	High risk	Low risk	Low risk	Low risk

# I. GRADE Assessment Criteria

We applied the following criteria for downgrading to each outcome.

# Risk of Bias

If >50% of studies had a low overall quality rating, the quality of the outcome was downgraded by 1. If >75% of studies had a low overall quality rating, the quality of the outcome was downgraded by 2.

### Imprecision

We downgraded an outcome for imprecision by 1 point if "a recommendation or clinical course of action would differ if the upper versus the lower boundary of the CI represented the truth" and/or the number of events and sample size meant the optimal information size was not reached (Guyatt et al., 2011). We downgraded by 2 points if an analysis was based on only 1-2 studies.

## Inconsistency

We downgraded an outcome for inconsistency by 1 point if the I<sup>2</sup> statistic was  $\geq 40\%$  in the context of an unclear direction of effect or  $\geq 75\%$  in the context of a clear direction of effect. We downgraded by 2 points if the I<sup>2</sup> statistic was  $\geq 75\%$  in the context of an unclear direction of effect.

# Publication Bias

We downgraded an outcome for publication bias by 1 point when, for outcomes with at least 10 studies (Higgins & Green, 2011), the funnel-plots showed asymmetry and this was not better explained by selective reporting bias or some other factor. However, if the 'trim and fill' method indicated that any publication bias was not likely to affect the overall magnitude of the effect size, we did not downgrade.

## Indirectness

Indirectness was assessed by considering the relevance of the outcome data to the construct of interest for each outcome, together with that of the study population, nature of the intervention under investigation and the control condition.
#### J. Table J.1. Other Comparisons

Studies	Outcome	Comparison (A vs B)	k included	Group A N	Group B N	Hedges' g (95% CI), p-value	Heterogeneity, I <sup>2</sup> , p-value	Quality (GRADE)
			studies Vei	rsus TAU/WL				
Krupnick 2008	PTSD + DR	IPT vs WL	1	32	16	-1.02 (-1.65, - 0.39), 0.002	-	Very low -2 RoB -2 imprecision
Azad marzabadi 2014	DR	Mindfulness vs TAU	1	14	14	-1.60 (-2.43, - 0.77), <0.001	-	Very low -2 RoB -2 imprecision
			Head-to	-head compar	risons			
Beidel 2011	PTSD, DR & AD	TMT vs exposure	1	14	16	-0.09 (-0.79, 0.61), 0.801	-	Very low -2 RoB -2 imprecision
Beidel 2019	PTSD & DR	TMT vs exposure	1	43	49	-0.05 (-0.46, 0.36), 0.815	-	Very low -2 RoB -2 imprecision
Butollo 2016	PTSD & NSC	DET vs CBT	1	66	72	0.27 (-0.07, 0.60), 0.118	-	-Very low -2 RoB -2 imprecision
Bryant 2013	PTSD & NSC	CBT + ERT vs CBT + SC	1	36	34	-0.04 (-0.51, 0.43), 0.866	-	Low -2 imprecision

Studies	Outcome	Comparison	k	Group A	Group B	Hedges' g (95%	Heterogeneity,	Quality
		(A vs B)	included	Ν	Ν	CI), p-value	I <sup>2</sup> , p-value	(GRADE)
			studies					
Harned 2014	PTSD & NSC	DBT +	1	12	6	0.51 (-0.43, 1.45),	-	Low
		Exposure vs				0.291		-2
		DBT						imprecision
ter Heide	PTSD & DR	EMDR vs	2	36	34	-0.16 (-0.61,	0%, 0.360	Low
2011,		STBT				0.29), 0.486		-2
ter Heide 2016								imprecision

Abbreviations: AD=affect dysregulation; CBT=cognitive behavioural therapy; DBT=dialectical behaviour therapy; DET=dialogical exposure therapy; DR=disturbances in relationships; EMDR=eye-movement and desensitisation and reprocessing therapy; ERT=emotion regulation training; IPT=interpersonal psychotherapy; NSC=negative self-concept; PTSD=posttraumatic stress disorder; SC=supportive counselling; STBT=stabilisation treatment; TAU=treatment as usual; TMT=trauma management therapy; WL=waiting list.

Treatment	Comparator	Outcome	g (95% CI), p-value	10% CER*	22% CER*	50% CER*
CBT	vs TAU/WL	DR	-0.66 (-0.84, -0.48), <0.001	6B (4B, 9B)	4B (3B, 6B)	4B (3B, 5B)
CBT	vs control	DR	-0.32 (-0.60, -0.03), 0.029	15B (7B, 186B)	9B (5B, 111B)	8B (4B, 84B)
CBT	vs TAU/WL	AD	-1.42 (-2.20, -0.65), <0.001	2B (1B, 6B)	2B (1B, 4B)	2B (2B, 4B)
CBT	vs control	AD	-0.82 (-2.91, 1.26), 0.440	5B (1B, 3H)	3B (1B, 2H)	3B (2B, 3H)
CBT	vs TAU/WL	NSC	-0.82 (-1.19, -0.44), <0.001	5B (3B,10B)	3B (2B, 7B)	3B (3B, 6B)
CBT	vs control	NSC	-0.24 (-0.69, 0.21), 0.295	20B (6B, 24H)	13B (4B, 15H)	11B (4B, 12H)
CBT	vs TAU/WL	PTSD	-0.90 (-1.11, -0.68), <0.001	4B (3B, 6B)	3B (2B, 4B)	3B (3B, 4B)
CBT	vs control	PTSD	-0.37 (-0.66, -0.09), 0.011	12B (6B 60B)	8B (4B, 36B)	7B (4B, 28B)
CBT	vs TAU/WL	PTSD + 1, 2 or 3	-0.81 (-1.00, -0.62), <0.001	5B (3B, 6B)	3B (3B, 5B)	3B (3B, 4B)
CBT	vs TAU/WL	PTSD + 2  or  3	-0.78 (-1.31, -0.24), 0.005	5B (2B, 20B)	4B (2B, 13B)	4B (2B, 11B)
CBT	vs TAU/WL	PTSD + 3	-0.53 (-0.96, -0.09), 0.017	8B (4B, 60B)	5B (3B, 36B)	5B (3B, 28B)
CBT	vs control	PTSD + 1, 2 or 3	-0.34 (-0.62, -0.06), 0.019	14B (6B, 91B)	9B (5B, 55B)	8B (4B, 42B)
CBT	vs control	PTSD + 2  or  3	-			
CBT	vs control	PTSD + 3	-			
Exposure	vs TAU/WL	DR	-0.59 (-1.12, -0.07), 0.028	7B (3B, 78B)	5B (2B, 47B)	5B (3B, 36B)
Exposure	vs control	DR	-0.12 (-0.60, 0.37), 0.642	44B (7B, 12H)	27B (5B, 8H)	21B (4B, 7H)
Exposure	vs TAU/WL	AD	-			
Exposure	vs control	AD	-			
Exposure	vs TAU/WL	NSC	-0.73 (-1.03, -0.43), <0.001	5B (3B, 10B)	4B (3B, 7B)	4B (3B, 6B)
Exposure	vs control	NSC	-			
Exposure	vs TAU/WL	PTSD	-1.05 (-1.52, -0.58), <0.001	3B (2B, 7B)	3B (2B, 5B)	3B (2B, 5B)
Exposure	vs control	PTSD	-0.08 (-0.47, 0.30), 0.675	68B (9B, 16H)	41B (6B 10H)	31B (6B, 8H)
Exposure	vs TAU/WL	PTSD + 1, 2 or 3	-0.86 (-1.25, -0.47), <0.001	4B (3B, 9B)	3B (2B, 6B)	3B (3B, 6B)
Exposure	vs TAU/WL	PTSD + 2  or  3	-0.56 (-0.99, -0.14), 0.009	7B (4B, 37B)	5B (3B, 23B)	5B (3B, 18B)
Exposure	vs TAU/WL	PTSD + 3	-			

#### K. Table K.1. Clinically Significant Response (number needed to treat per control group event rate)

Treatment	Comparator	Outcome	g (95% CI), p-value	10% CER*	22% CER*	50% CER*
Exposure	vs control	PTSD + 1, 2 or 3	-0.19 (-0.57, 0.20), 0.336	27B (7B, 25H)	17B (5B, 16H)	13B (5B, 13H)
Exposure	vs control	PTSD + 2  or  3	-			
Exposure	vs control	PTSD + 3	-			
EMDR	vs TAU/WL	DR	-0.76 (-1.35, -0.16), 0.012	5B (2B, 32B)	4B (2B, 20B)	4B (2B, 16B)
EMDR	vs control	DR	-0.35 (-1.01, 0.31), 0.312	13B (3B, 15H),	9B (3B, 10H)	7B (3B, 8H)
EMDR	vs TAU/WL	AD	-1.64 (-2.56, -0.72), 0.000	2B (1B, 5B)	2B (1B, 4B)	2B (2B, 4B)
EMDR	vs control	AD	0.25 (-0.57, 1.08), 0.548	20H (5B, 3H)	12H (5B, 2H)	10H (5B, 3H)
EMDR	vs TAU/WL	NSC	-0.61 (-1.04, -0.17), 0.006	7B (3B, 30B)	5B (3B, 19B)	4B (3B, 15B)
EMDR	vs control	NSC	-0.78 (-1.56, -0.01), 0.049	4B (2B, 566B)	4B (2B, 336B)	4B (2B, 251B)
EMDR	vs TAU/WL	PTSD	-1.26 (-2.01, -0.51), 0.001	3B (2B, 8B)	2B (1B, 6B)	3B (2B, 5B)
EMDR	vs control	PTSD	-0.69 (-1.35, -0.03), 0.041	6B (2B, 186B)	4B (2B, 111B)	4B (2B, 84B)
EMDR	vs TAU/WL	PTSD + 1, 2 or 3	-1.15 (-1.92, -0.37), 0.004	3B (2B, 12B)	2B (2B, 8B)	3B (2B, 7B)
EMDR	vs TAU/WL	PTSD + 2  or  3	-1.36 (-3.13, 0.42), 0.134	2B (1B, 11H)	2B (1B, 7H)	2 (2B, 6H)
EMDR	vs TAU/WL	PTSD + 3	-			
EMDR	vs control	PTSD + 1, 2 or 3	-0.52 (-0.97, -0.08), 0.020	8B (4B, 68B)	6B (3B, 41B)	5B (3B, 31B)
EMDR	vs control	PTSD + 2  or  3	-0.44 (-1.31, 0.43), 0.321	10B (2B, 10H)	7B (2B, 7H)	6B (2B, 6H)
EMDR	vs control	PTSD + 3	-			
CBT (T)	vs exposure (C)	DR	0.07 (-0.26, 0.39), 0.689	78C (19T, 12C)	47C (12T, 8C)	36C (10T, 7C)
CBT (T)	vs exposure (C)	AD	-			
CBT (T)	vs exposure (C)	NSC	-0.31 (-0.67, 0.04), 0.082	15T (6T, 139C)	10T (4T, 83C)	8T (4T, 63C)
CBT (T)	vs exposure (C)	PTSD	-0.03 (-0.23, 0.17), 0.784	186T (22T,	111T (14T,	84T (11T, 15C)
				30C)	19C)	
CBT (T)	vs exposure (C)	PTSD + 1, 2, or 3	-0.04 (-0.27, 0.19), 0.719	139T (18T,	83T (11T,	63T (9T, 13C)
				27C)	17C)	
CBT (T)	vs exposure (C)	PTSD + 2  or  3	-			
CBT (T)	vs exposure (C)	PTSD + 3	-			
CBT (T)	EMDR (C)	DR	0.28 (-0.29, 0.34), 0.338	17C (16T, 14C)	11C (11T, 9C)	9C (9T, 8C)

Treatment	Comparator	Outcome	g (95% CI), p-value	10% CER*	22% CER*	50% CER*
CBT(T)	FMDR (C)	AD				
CBT(T)	EMDR(C)	NSC	-			
CBT (T)	EMDR (C)	PTSD	0.37 (0.03, 0.71), 0.031	12C (186C, 5C)	8C (111C, 4C)	7C (84C, 4C)
CBT (T)	EMDR (C)	PTSD + 1, 2, or 3	0.31 (-0.07, 0.68), 0.111	15C (78T, 6C)	10C (47T, 4C)	8C (36T, 4C)
CBT (T)	EMDR (C)	PTSD + 2  or  3	-			
CBT (T)	EMDR (C)	PTSD + 3	-			
EMDR (T)	Exposure (C)	DR	-0.10 (-0.51, 0.31), 0.640	54T (8T, 15C)	33T (6T, 10C)	25T (5T, 8C)
EMDR (T)	Exposure (C)	AD	-			
EMDR (T)	Exposure (C)	NSC	0.16 (-0.25, 0.57), 0.444	32C (20T, 7C)	20C (12T,	16C (10T, 5C)
					47C)	
EMDR (T)	Exposure (C)	PTSD	0.10 (-0.28, 0.49), 0.604	54C (17T, 9C)	33C (11T, 6C)	25C (9T, 5C)
EMDR (T)	Exposure (C)	PTSD + 1, 2, or 3	0.06 (-0.35, 0.46), 0.789	91C (35T, 9C)	55C (9T, 6C)	42C (7T, 6C)
EMDR (T)	Exposure (C)	PTSD + 2  or  3	0.06 (-0.35, 0.46), 0.789	91C (35T, 9C)	55C (9T, 6C)	42C (7T, 6C)
EMDR (T)	Exposure (C)	PTSD + 3	-			
IPT	vs TAU/WL	PTSD + DR	-1.02 (-1.65, -0.39), 0.002	3B (2B, 12B)	3B (2B, 8B)	3B (2B, 7B)
Mindfulness	vs TAU/WL	DR	-1.60 (-2.43, -0.77), <0.001	2B (1B, 5B)	2B (1B, 4B)	2B (2B, 4B)
TMT (T)	Exposure (C)	PTSD + DR + AD	-0.09 (-0.79, 0.61), 0.801	60T (5T, 7C)	36T (3T, 5C)	28T (4T, 4C)
TMT (T)	Exposure (C)	PTSD + DR	-0.05 (-0.46, 0.36), 0.815	110T (9T, 13C)	66T (6T, 8C)	50T (6T, 7C)
DET (T)	CBT (C)	PTSD + NSC	0.27 (-0.07, 0.60), 0.118	18C (78T, 7C)	11C (47T, 5C)	9C (36T, 4C)
CBT + ERT(T)	CBT + SC(C)	PTSD + NSC	-0.04 (-0.51, 0.43), 0.866	139T (8T, 10C)	83T (6T, 7C)	63T (5T, 6C)
DBT + Exp(T)	CBT (C)	PTSD + NSC	0.51 (-0.43, 1.45), 0.291	8C (10T, 2C)	7C (7T, 2C)	5C (6T, 2C)
EMDR (T)	STBT (C)	PTSD + DR	-0.16 (-0.61, 0.29), 0.486	32T (7T, 16C)	20T (5T, 11C)	16T (4T, 9C)

Abbreviations: AD=affect dysregulation; CBT=cognitive behavioural therapy; CPTSD=complex posttraumatic stress disorder; CER=control event rate; DBT=dialectical behaviour therapy; DET=dialogical exposure therapy; DR=disturbances in relationships; DSO=disturbances in self-organisation; EMDR=eye-movement and desensitisation and reprocessing therapy; ERT=emotion regulation training; IPT=interpersonal psychotherapy; NSC=negative self-concept; PTSD=posttraumatic stress disorder; PTSD + 1, 2 or 3 CPTSD (DSO) outcomes; PTSD + 2 or 3 CPTSD (DSO) outcomes; PTSD + 2 or 3 CPTSD (DSO) outcomes;

SC=supportive counselling; STBT=stabilisation treatment; TAU=treatment as usual; TMT=trauma management therapy; WL=waiting list. \*Note: B = benefit; H = harm; T = favours T; C = favours C

#### L. Table L.1. Meta-regression Moderators (univariate)

Moderator	Coefficients (95% CI)	$\mathbf{R}^2$ or $\Delta \mathbf{R}^2$	p-value	Effects per group	Quality
(univariate)					
Random sequence	Unclear risk of bias	2%	0.358	Low (k=26)	Low
generation (low vs	-0.14 (-0.43, +0.16)			-0.64 (-0.84, -0.45)	-1 missing
unclear or high risk of				Unclear (k=26)	information
bias, k=52)				-0.78 (-0.99, -0.56)	-1 imprecise
				High (k=0)	
Allocation	Unclear risk of bias	5%	0.112	Low (k=20)	Low
concealment (low vs	-0.24 (-0.53, +0.06)			-0.57 (-0.79, -0.35)	-1 missing
unclear or high risk of				Unclear (k=32)	information
bias, k=52)				-0.80 (-0.99, -0.61)	-1 imprecise
				High (k=0)	
Detection bias (low vs	High risk of bias	1%	0.820	Low (k=35)	Moderate
unclear or high risk of	-0.04 (-0.35, +0.28)			-0.69 (-0.87, -0.52)	-1 imprecise
bias, k=52)				Unclear (k=0)	
				High $(k=17)$	
				-0.72 (-0.99, -0.47)	
Selective reporting	High risk of bias	7%	0.131	Low (k=5)	Moderate
bias (low vs unclear or	-0.35 (-0.81, +0.11)			-0.39 (-0.82, 0.04)	-1 imprecise
high risk of bias,				Unclear (k=0)	
k=52)				High $(k=47)$	
				-0.74 (-0.89, -0.59)	

Moderator	Coefficients (95% CI)	$\mathbf{R}^2$ or $\Delta \mathbf{R}^2$	p-value	Effects per group	Quality
(univariate)					
Attrition bias (low vs	High risk of bias	1%	0.524	Low (k=31)	Moderate
unclear or high risk of	+0.10(-0.20, +0.39)			-0.65 (-0.87, -0.42)	-1 imprecise
bias, k=52)				Unclear (k=0)	1
, ,				High $(k=21)$	
				-0.74 (-0.93, -0.55)	
Quality (high quality	Low quality	0%	0.754	Low quality (k=20)	Moderate
vs low quality, $k=52$ )	+0.05(-0.26, +0.35)			-0.67 (-0.91, -0.44)	-1 imprecise
				High quality (k=32)	-
				-0.72 (-0.91, -0.54)	
CPTSD symptoms	PTSD + ER	PTSD + AD	PTSD + AD	PTSD (k=3)	Moderate
(PTSD alone vs	-0.57 (-1.44, +0.32)	1%	0.215	-0.53 (-1.25, 0.18)	-1 imprecise
various CPTSD,	PTSD + NC + ID + ER	PTSD + NSC + DR	PTSD + NSC + DR	PTSD + AD (k=4)	
k=52)	-0.03 (-1.01, +0.50)	+ AD	+ AD	-1.09 (-1.66, -0.53)	
	PTSD + ID	2%	0.955	PTSD + NSC + DR + AD	
	-0.20 (-0.92, 0.53)	PTSD + DR	PTSD + DR	(k=2)	
	PTSD + NC	3%	0.593	-0.55 (-1.27, 0.17)	
	-0.11 (-0.85, 0.63)	PTSD + NSC	PTSD + NSC	PTSD + DR (k=24)	
	PTSD + NC + ID	3%	0.778	-0.72 (-0.94, -0.50)	
	-0.26 (-1.11, +0.59)	PTSD + NSC + DR	PTSD + NSC + DR	PTSD + NSC (k=15)	
		7%	0.548	-0.63 (-0.90, -0.36)	
		Overall	Overall	PTSD + NSC + DR (k=4)	
		7%	0.741	-0.78 (-1.29, -0.27)	
Comparator	Control	28%	0.001	TAU/WL (k=38)	Moderate
(TAU/WL vs control,	+0.48(+0.18,+0.77)			-0.83 (-0.99, -0.67)	-1 imprecise
k=52)				Control (k=14)	
				-0.35 (-0.60, -0.10)	

Moderator (univariate)	Coefficients (95% CI)	$\mathbf{R}^2$ or $\Delta \mathbf{R}^2$	p-value	Effects per group	Quality
Treatments (individual	EMDR	EMDR 1%	EMDR 0.736	CBT (k=33)	Moderate
CBT vs others, k=52)	-0.08 (-0.53, +0.38)	Exposure 0%	Exposure 0.834	-0.73 (-0.91, -0.54)	-1 imprecise
	Exposure	Group CBT 7%	Group CBT 0.150	EMDR (k=7)	
	+0.04(-0.53, +0.38)	Group IPT 2%	Group IPT 0.581	-0.80 (-1.23, -0.38)	
	Group CBT	Overall 10%	Overall 0.608	Exposure (k=8)	
	+0.42(-0.15, +0.99)			-0.68 (-1.06, -0.31)	
	Group IPT			Group CBT $(k=3)$	
	-0.29 (-1.34, +0.75)			-0.30(-0.86, 0.25)	
				Group IP1 ( $k=1$ )	
T1		40/	0.200	-1.02(-2.07, 0.04)	Madausta
(individual va group	Group only of in addition $\pm 0.27$ ( $0.25 \pm 0.78$ )	4%	0.309	1101V100a1 (K-48) 0.72 (0.88 0.58)	
(11101110001108 group, 12-52)	+0.27(-0.23, +0.78)			-0.75(-0.80, -0.50)	-1 imprecise
K-32)				(k-4)	
				$(\mathbf{K} - 4)$ -0.46 (-0.95, 0.03)	
Trauma onset (Adult	Child	2%	0.308	$\frac{-0.40(-0.55, 0.05)}{\text{Adult (k=37)}}$	Moderate
vs child $k=48$ )	+0.18(-0.16+0.52)	270	0.500	-0.76 (-0.93, -0.59)	-1 imprecise
vs child, k 10)	10.10 (0.10, 10.52)			Child $(k=11)$	1 impreeise
				-0.58 (-0.88, -0.29)	

Abbreviations: AD=affect dysregulation; CBT=cognitive behavioural therapy; DR=disturbances in relationships; EMDR=eye-movement and desensitisation and reprocessing therapy; k= number of included comparisons; NSC=negative self-concept; PTSD=posttraumatic stress disorder; CPTSD=complex posttraumatic stress disorder; TAU=treatment as usual; WL=waiting list.

#### M. Table M.1. Meta-regression Moderators (multivariate)

Moderator (k=48,	Coefficients (95% CI)	$\Delta \mathbf{R}^2$	p-value	Narrative summary	Quality
multivariate)					
Quality (high vs low)	Low	Low	Low	The effect for low quality	Moderate
	+0.30 (+0.00, +0.61)	1%	0.048	studies is 0.30 lower than	-1 imprecise
				high quality studies	
CPTSD symptoms	PTSD + AD	PTSD + AD	PTSD + AD	No association between	Moderate
(PTSD alone vs	-0.13 (-1.06, +0.80)	-1%	0.783	number or type of CPTSD	-1 imprecise
various CPTSD)	PTSD + NSC + DR +	PTSD + NSC + DR	PTSD + NSC + DR	symptoms reported and	
	AD	+ ER	+ ER	effect size was observed	
	+0.39 (-0.62, +1.40)	1%	0.447	(direction of effect favoured	
	PTSD + DR	PTSD + DR	PTSD + DR	smaller effects with more	
	+0.28 (-0.43, +1.00)	0%	0.437	CPTSD symptoms).	
	PTSD + NSC	PTSD + NSC	PTSD + NSC		
	+0.06 (-0.63, +0.75)	0%	0.860		
	PTSD + NSC + DR	PTSD + NSC + DR	PTSD + NSC + DR		
	+0.25 (-0.52, +1.02)	2%	0.518		
		Overall	Overall		
		2%	0.504		
Comparator	Control	Control	Control	Use of a control condition is	High
(TAU/WL vs control)	+0.69(+0.38,+1.00)	34%	< 0.0001	associated with a moderate	
				to large reduction in effect	
				size.	

Moderator (k=48,	Coefficients (95% CI)	$\Delta \mathbf{R}^2$	p-value	Narrative summary	Quality
multivariate)					
Treatments (individual	EMDR	EMDR	EMDR	No association between	Moderate
CBT vs others)	-0.25 (-0.69, +0.18)	3%	0.254	overall effect size and type	-1 imprecise
	Exposure	Exposure	Exposure	of intervention was	
	+0.07 (-0.30, +0.44)	0%	0.697	observed.	
	Group CBT	Group CBT	Group CBT		
	+0.23 (-0.36, 0.82)	7%	0.441		
	Group IPT	Group IPT	Group IPT		
	-0.70 (-1.66, 0.25)	1%	0.147		
		Overall	Overall		
		11%	0.282		
Trauma onset (adult	Child	Child	Child	Inclusion of participants	Low
vs child)	+0.35(+0.02, +0.69)	5%	0.038	with predominantly	-1 imprecise
				childhood-onset trauma is	-1
				associated with a small-	ecological
				moderate reduction in effect	bias
				size, compared to trials	
				where participants have	
				mainly adult-onset trauma	

Abbreviations: AD=affect dysregulation; CBT=cognitive behavioural therapy; DR=disturbances in relationships; EMDR=eye-movement and desensitisation and reprocessing therapy; k= number of included comparisons; NSC=negative self-concept; PTSD=posttraumatic stress disorder; CPTSD=complex posttraumatic stress disorder; TAU=treatment as usual; WL=waiting list.

### N. Forest Plots – Cognitive/imagery modification with or without exposure vs TAU/WL or non-specific control

Fig. N.1. Disturbances in relationships (DR): Cognitive/imagery modification with or without exposure vs TAU/WL



# Fig. N.2. Disturbances in relationships (DR): Cognitive/imagery modification with or without exposure vs non-specific control

Study name		Statistics for	reach stu	udy		Sample size				Hedges's g and 95%Cl			
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B						
Ehlers 2014 vs ST DR	0.508	0.224	-0.948	-0.069	0.023	61	30		—C	]—[			
Marks 1998 vs relax DR	0.331	0.275	-0.871	0.209	0.229	37	20						
Forbes 2012 vs range DR	۹.048	0.257	-0.552	0.455	0.850	30	29		-	—¢—	-		
	0.315	0.144	-0.598	-0.033	0.029	128	79		•				
								-2.00	-1.00	0.00	1.00	2.00	
									Favours A		Favours B		

Fig. N.3. Affect dysregulation (AD): Cognitive/imagery modification with or without exposure vs TAU/WL

Study name	Statistics for each study					Sample size			Hedges's g and 95%Cl			
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
linton 2009 vs WL AD	-2.443	0.529	-3.479	-1.406	0.000	12	12	-				
Cloitre 2002 vs WL AD	-1.291	0.320	-1.918	-0.664	0.000	22	24			-		
Ionson 2006 vs WL AD	-0.847	0.308	-1.451	-0.244	0.006	20	25		-(	<u></u>		
	-1.424	0.397	-2.202	-0.647	0.000	54	61					
								-4.00	-2.00	0.00	2.00	4.00
									Favours A		Favours B	

## Fig. N.4. Affect dysregulation (AD): Cognitive/imagery modification with or without exposure vs non-specific control



Fig. N.5. Negative self-concept (NSC): Cognitive/imagery modification with or without exposure vs TAU/WL

Study name		Statistics for	reach stu	udy		Samp	e size		Hedges	sgand 98	5%C	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
ubany 2003 vs WL NSC	-2.994	0.511	-3.995	-1.993	0.000	18	14		-0			
ung 2013 vs WL NSC	-0.740	0.380	-1.485	0.005	0.052	14	14		-	-0		
lonson 2006 vs WL NSC	-0.452	0.301	-1.042	0.138	0.133	20	25			-0+		
IcDonagh 2005 vs WL NS0	-0.548	0.280	-1.097	0.001	0.050	29	23			-0-		
lueser 2008 vs TAU NSC	-0.257	0.259	-0.764	0.251	0.322	32	27			-0-		
Galovski 2012 vs WL NSC	-0.935	0.257	-1.438	-0.432	0.000	34	32					
ord 2011 vs WL NSC	-1.149	0.222	-1.585	-0.714	0.000	48	45		-	-		
esick 2002 vs WL NSC	-0.888	0.212	-1.304	-0.472	0.000	62	39		-	⊡-		
ubany 2004 vs WL NSC	-0.209	0.178	-0.558	0.141	0.242	63	62			-1		
	-0.816	0.191	-1.190	-0.441	0.000	320	281		•	◆		
								-4.00	-2.00	0.00	2.00	4.00
									Favours A		Favours B	

## Fig. N.6. Negative self-concept (NSC): Cognitive/imagery modification with or without exposure vs non-specific control



#### Fig. N.7. PTSD: Cognitive/imagery modification with or without exposure vs TAU/WL

Olday Harris		Statistics for	or each s	tudy		Samp	le size		Hedg	Hedges's g and 95%	Hedges's g and 95% Cl
н	edges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B				
Kubany 2003 vs WL PTSD	-2.92	0.50	-3.90	-1.93	0.00	18	14	I			
Hinton 2009 vs WL PTSD	-1.91	0.48	-2.85	-0.97	0.00	12	12				
Enlers 2005 vs WL PTSD	-2.19	0.47	-3.11	-1.27	0.00	14	14				
Dunne 2012 vs WL PTSD	-0.92	0.42	-1.75	-0.09	0.03	12	11				
Lindauer 2005 vs WL PTSD	-0.90	0.42	-1.72	-0.09	0.03	12	12		I I		
Jung 2013 vs WL PTSD	-0.81	0.38	-1.56	-0.06	0.03	14	14				
Difede 2007 vs TAU PTSD	-0.34	0.35	-1.03	0.35	0.33	15	16				
Foa 1999 vs WL PTSD	-1.51	0.33	-2.16	-0.87	0.00	41	15				
Claitre 2002 vs WL PTSD	-1.27	0.32	-1.90	-0.65	0.00	22	24				
Power 2002 vs WL PTSD	-1.14	0.32	-1.76	-0.51	0.00	21	24				
Talbot 2014 vs WL PTSD	-0.64	0.31	-1.26	-0.03	0.04	29	16				
Hollifield 2007 vs WL PTSD	-0.68	0.29	-1.25	-0.11	0.02	25	24				
McDonach 2005 vs WL PTSD	-0.49	0.28	-1.04	0.06	0.08	29	23				
Steel 2017 vs TAU PTSD	0.16	0.28	-0.39	0.70	0.57	25	25				
Galovski 2012 vs WL PTSD	-1.50	0.28	-2.04	-0.96	0.00	34	32				
Monson 2006 vs WL PTSD	-1.13	0.28	-1.67	-0.59	0.00	30	30				
Duffy 2007 vs WL PTSD	-0.88	0.27	-1.41	-0.34	0.00	29	29				
Mueser 2008 vs TAU PTSD	-0.44	0.26	-0.95	0.07	0.09	32	27				
Basodiu 2007 vs WL PTSD	-0.44	0.26	-0.95	0.08	0.09	31	28				
Ehlers 2003 vs SH + WL PTS	D -1.08	0.25	-1.56	-0.59	0.00	28	52				
Ehlers 2014 vs WL PTSD	-1.38	0.24	-1.85	-0.90	0.00	61	30				
Dorrepaal 2012 vs TAU PTSD	-0.29	0.24	-0.75	0.17	0.22	38	33				
Foa 2005 vs WL PTSD	-0.80	0.23	-1.26	-0.34	0.00	74	26		-		
Ford 2011 vs WL PTSD	-0.96	0.22	-1.39	-0.54	0.00	48	45		-		
Residk 2002 vs WL PTSD	-1.13	0.21	-1.54	-0.73	0.00	62	47				
Kubany 2004 vs WL PTSD	-0.28	0.18	-0.63	0.07	0.12	63	62				
Krakow 2001 vs WL PTSD	-0.13	0.15	-0.43	0.17	0.39	80	88				
	-0.90	0.11	-1.11	-0.68	0.00	899	773			♠ ヿ	
								-4.00	-4.00 -2.00	-4.00 -2.00 0.00	-4.00 -2.00 0.00 2.00

## Fig. N.8. PTSD: Cognitive/imagery modification with or without exposure vs non-specific control



# Fig. N.9. PTSD plus 1, 2 or 3 CPTSD outcomes: Cognitive/imagery modification with or without exposure vs TAU/WL

Study name	5	Statistics for	each stu	dy		Sampl	e size		Hedg	es's g and 95%	a	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
bany 2003 vs WL PTSD + NSC	-2.956	0.508	-3.951	-1.961	0.000	18	14			1	1	1
nton 2009 vs WL PTSD + AD	-2.177	0.505	-3.167	-1.187	0.000	12	12					
lers 2005 vs WL PTSD + DR	-1.861	0.445	-2.732	-0.990	0.000	14	14		p	-		
nne 2012 vs WL PTSD + DR	-0.788	0.420	-1.610	0.034	0.060	12	11		<u> </u>	-0		
iede 2007 vs TAU + DR	-0.460	0.403	-1.250	0.330	0.254	7	14					
ng 2013 vs WL PTSD + NSC	-0.774	0.381	-1.520	-0.027	0.042	14	14		-	-0		
bitre 2002 vs WL PTSD + AD	-1.282	0.319	-1.908	-0.656	0.000	22	24		—C	— I		
a 1999 vs WL PTSD + DR	-1.142	0.318	-1.765	-0.518	0.000	41	15			I		
lbot 2014 vs WL PTSD + DR	-0.766	0.317	-1.387	-0.144	0.016	29	16		-	-0		
wer 2002 vs WL PTSD + DR	-0.912	0.310	-1.519	-0.305	0.003	21	24		I –			
eel 2017 vs TAU PTSD + NSC	0.076	0.303	-0.517	0.669	0.802	18	18			— <del>0</del> —		
onson 2006 vs WL PTSD + AD + DR + NSC	-0.786	0.297	-1.368	-0.204	0.008	20	25		- 1	-0		
llifield 2007 vs WL PTSD + DR	-0.625	0.288	-1.189	-0.061	0.030	25	24			-0-1		
Donagh 2005 vs WL PTSD + NSC	-0.520	0.280	-1.068	0.028	0.063	29	22			-0-		
iffy 2007 vs WL PTSD + DR	-0.820	0.278	-1.365	-0.275	0.003	29	25		- 1	-0		
dauer 2005 vs WL PTSD + DR	-0.581	0.267	-1.104	-0.058	0.029	12	12			-0-1		
alovski 2012 vs WL PTSD + NSC + DR	-1.233	0.267	-1.755	-0.710	0.000	34	32		<u>-</u> c	⊢		
Jeser 2008 vs TAU PTSD + NSC	-0.349	0.260	-0.859	0.161	0.179	32	27			-0+		
isoglu 2007 vs WL PTSD + DR	-0.290	0.259	-0.798	0.218	0.263	31	28			-0+		
lers 2003 vs SH + WL PTSD + DR	-0.939	0.244	-1.417	-0.461	0.000	28	52		· · -	<u> </u>		
vrepaal 2012 vs TAU PTSD + AD + DR + NSC	-0.338	0.238	-0.803	0.127	0.155	38	33			0+		
lers 2014 vs WL PTSD + DR	-1.144	0.235	-1.605	-0.683	0.000	61	30			<u>-  </u>		
a 2005 vs WL PTSD + DR	-0.620	0.232	-1.075	-0.165	0.008	74	25			-0-1		
rd 2011 vs WL PTSD + NSC	-1.060	0.220	-1.491	-0.629	0.000	48	45			<u> </u>		
isick 2002 vs WL PTSD + NSC	-1.011	0.210	-1.421	-0.600	0.000	62	39			Ē−		
akow 2001 vs WL PTSD + DR	-0.274	0.180	-0.626	0.079	0.129	47	48			+		
bany 2004 vs WL PTSD + NSC	-0.245	0.179	-0.595	0.105	0.170	63	62			-7+		
	-0.808	0.095	-0.995	-0.622	0.000	841	705			◆ -		
								-4.00	-2.00	0.00	2.00	4.00

# Fig. N.10. PTSD plus 2 or 3 CPTSD outcomes: Cognitive/imagery modification with or without exposure vs TAU/WL

Study name	5	Statistics for	each stu	dy		S <u>ampl</u>	e size		Hedges	s's g and 95%	a	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
nson 2006 vs WL PTSD + AD + DR + NSC	-0.786	0.297	-1.368	-0.204	0.008	20	25		-+0-	-		
lovski 2012 vs WL PTSD + NSC + DR	-1.233	0.267	-1.755	-0.710	0.000	34	32	-				
rrepaal 2012 vs TAU PTSD + AD + DR + NSC	-0.338	0.238	-0.803	0.127	0.155	38	33		—			
	-0.775	0.273	-1.310	-0.240	0.005	92	90					
								-2.00	-1.00	0.00	1.00	2.00
									Favours A		Favours B	

Fig. N.11. PTSD plus all 3 CPTSD outcomes: Cognitive/imagery modification with or without exposure vs TAU/WL

Study name	5	Statistics for	each stu	xdy		Sampl	e size		Hedges	's g and 95%	a	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
onson 2006 vs WL PTSD + AD + DR + NSC	-0.786	0.297	-1.368	-0.204	0.008	20	25		+0	-		
orrepaal 2012 vs TAU PTSD + AD + DR + NSC	-0.338	0.238	-0.803	0.127	0.155	38	33		I —			
	-0.526	0.221	-0.960	-0.093	0.017	58	58					
								-2.00	-1.00	0.00	1.00	2.00
									Favours A		Favours B	

## Fig. N.12. PTSD plus 1, 2 or 3 CPTSD outcomes: Cognitive/imagery modification with or without exposure vs non-specific control



#### O. Forest Plots – Exposure only vs TAU/WL or non-specific control

Fig. O.1. Disturbances in relationships (DR): Exposure only vs TAU/WL

Study name	5	Sta <u>tistics for</u>	eachstu	ıdy		S <u>amp</u> i	le size		Hedges	'sgand95%	6 <b>a</b>	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
Foa 1999 vs WL DR	-1.497	0.371	-2.224	-0.770	0.000	22	15		+0-	-		
Keane 1989 vs WL DR	-0.730	0.353	-1.422	-0.037	0.039	11	31		-			
Foa 2005 vs WL DR	-0.237	0.229	-0.685	0.211	0.300	78	25					
van den Berg 2015 vs WL DF	۶ -0.177	0.215	-0.599	0.244	0.410	47	39			-[]-		
	-0.594	0.270	-1.123	-0.066	0.028	158	110		-	$\bullet$		
								-4.00	-2.00	0.00	2.00	4.00
									Favours A		Favours B	

### Fig. O.2. Disturbances in relationships (DR): Exposure only vs non-specific control

Study name		Statistics for	r each stu	udy		Samp	le size		Hedges	sgand 98	5%CI	
	Hedges's g	Standard error	Lower limit	Upper İimit	p-Value	Group-A	Group-B					
Ghafoori 2017 vs PCT DR	0.116	0.248	-0.602	0.371	0.642	47	24					
	0.116	0.248	-0.602	0.371	0.642	47	24			$\blacklozenge$		
								-4.00	-2.00	0.00	2.00	4.00
									Favours A		Favours B	

### Fig. O.3. Negative self-concept (NSC): Exposure only vs TAU/WL

Study name	:	Statistics for	each stu	udy		Samp	e size		Hedges	sgand 95	% <b>a</b>	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
an den Berg 2015 vs WL NS	C -0.739	0.222	-1.174	-0.304	0.001	47	39		-		1	
esick 2002 vs WL NSC	-0.519	0.207	-0.924	-0.114	0.012	61	39					
acella 2012 vs WL NSC	-1.107	0.309	-1.712	-0.501	0.000	23	24		— C	⊢		
	-0.728	0.155	-1.031	-0.425	0.000	131	102		-	◆		
								-4.00	-2.00	0.00	2.00	4.00
									Favours A		Favours B	

### Fig. O.4. PTSD: Exposure only vs TAU/WL

Studyname	5	Statistics for	eachstu	udy		Sampi	le size		Hedges	'sgand95%	6 <b>0</b>	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
Foa 1999 vs WL PTSD	-1.915	0.392	-2.683	-1.146	0.000	23	15		—¢—			
Pacella 2012 vs WL PTSD	-2.146	0.364	-2.858	-1.434	0.000	23	24					
Keane 1989 vs WL PTSD	-0.237	0.345	-0.914	0.439	0.492	11	31					
Foa 2005 vs WL PTSD	-0.656	0.229	-1.104	-0.207	0.004	79	26					
van den Berg 2015 vs WL PTSD	-0.777	0.212	-1.191	-0.362	0.000	48	47			⊡-		
Residk 2002 vs WL PTSD	-0.862	0.201	-1.255	-0.468	0.000	62	47		-	<u>-  </u>		
	-1.048	0.241	-1.521	-0.575	0.000	246	190					
								-4.00	-2.00	0.00	2.00	4.00
									Favours A		Favours B	

#### Fig. O.5. PTSD: Exposure only vs non-specific control

Study name	5	Statistics for	each stu	udy		Samp	e size		Hedges'	sgand95%	6 <b>a</b>	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
Marks 1998 vs relax PTSD	-0.145	0.319	-0.769	0.480	0.650	20	18			-0-		
Ghafoori 2017 vs PCT PTSD	-0.044	0.248	-0.530	0.442	0.859	47	24			-0-		
	-0.082	0.196	-0.466	0.302	0.675	67	42					
								-1.00	-0.50	0.00	0.50	1.00
									Favours A		Favours B	

#### Fig. O.6. PTSD plus 1, 2 or 3 CPTSD outcomes: Exposure only vs TAU/WL

Study name	:	Statistics for	r each sti	udy		Sampl	e size		He <u>dge</u>	s's g and 95°	% <b>C</b>	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
a 1999 vs WL PTSD + DR	1.706	0.382	-2.455	-0.957	0.000	22	15			-	1	1
ine 1989 vs WL PTSD + DR	0.484	0.349	-1.168	0.200	0.165	11	16		- I -	-0+		
cella 2012 vs WL PTSD + NSC	1.627	0.337	-2.286	-0.967	0.000	23	24			-		
2005 vs WL PTSD + DR	0.447	0.229	-0.896	0.002	0.051	78	25			-0-		
den Berg 2015 vs WL PTSD + NSC + DF	₹ 0.564	0.216	-0.988	-0.140	0.009	47	39					
iick 2002 vs WL PTSD + NSC	0.690	0.204	-1.090	-0.291	0.001	61	39					
	0.861	0.200	-1.252	-0.469	0.000	242	158		•	◆		
								-4.00	-2.00	0.00	2.00	4.0
									Favours A		Favours B	

### Fig. O.7. PTSD plus 2 or 3 CPTSD outcomes: Exposure only vs TAU/WL

<u>Study name</u>	;	Statistics for	r each stu	udy		Sampl	e size		Hedg	es's g and 95	%C	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
en Berg 2015 vs WL PTSD + NSC + DR	-0.564	0.216	-0.988	-0.140	0.009	47	39					
uen beig 2013 VS WL F 13D + 143C + DK	-0.564	0.216	-0.988	-0.140	0.009	47	39			$\blacklozenge$		
								-4.00	-2.00	0.00	2.00	4.0
									Favours A		Favours B	

### Fig. O.8. PTSD plus 1, 2 or 3 CPTSD outcomes: Exposure only vs non-specific control

Study name	5	Statistics for	each stu	udy		Samp	e size		Hedge	s's gand 95°	% <b>a</b>	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
Marks 1998 vs relax PTSD + DR	-0.306	0.317	-0.927	0.315	0.334	20	18			-0+		
Ghafoori 2017 vs PCT DR	-0.116	0.248	-0.602	0.371	0.642	47	24					
	-0.188	0.195	-0.571	0.195	0.336	67	42			-		
								-4.00	-2.00	0.00	2.00	4.00
									Favours A		Favours B	

#### P. Forest Plots – EMDR vs TAU/WL or non-specific control

Fig. P.1. Disturbances in relationships (DR): EMDR vs TAU/WL



### Fig. P.2. Disturbances in relationships (DR): EMDR vs non-specific control

Study name		Statistics for	r each stu	udy		Samp	e size		Hedges's g and 95% Cl			
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
Ahmadi 2015 vs REM DR	-0.776	0.436	-1.631	0.079	0.075	11	10	- I -	+0			
Kip 2013 vs control DR	-0.075	0.279	-0.621	0.472	0.789	26	24		-	-¢-	-	
	-0.345	0.341	-1.014	0.324	0.312	37	34					
								-2.00	-1.00	0.00	1.00	2.00
									Favours A		Favours B	

### Fig. P.3. Affect dysregulation (AD): EMDR vs TAU/WL

<u>Study name</u>		Statistics for	r each stu	ıdy		Samp	le size					
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
Ahmadi 2015 vs TAU Al	1.636	0.469	-2.556	-0.717	0.000	11	12	k−□				
	-1.636	0.469	-2.556	-0.717	0.000	11	12					
								-2.00	-1.00	0.00	1.00	2.00
									FavoursA		Favours B	

### Fig. P.4. Affect dysregulation (AD): EMDR vs non-specific control

<u>Study name</u>		r each stu	udy		Samp	le size		Hedges's g and 95%Cl				
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
hmadi 2015 vs REM A	AD 0.253	0.421	-0.572	1.079	0.548	11	10		-	-+C	]	
	0.253	0.421	-0.572	1.079	0.548	11	10					
								-2.00	-1.00	0.00	1.00	2.00
									Favours A		Favours B	

### Fig. P.5. Negative self-concept (NSC): EMDR vs TAU/WL

<u>Study name</u>	:	Statistics for each study				Sample size			Hedges's g and 95%Cl				
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B						
an den Berg 2015 vs WL NS	0.607	0.223	-1.044	-0.170	0.006	44	39						
	-0.607	0.223	-1.044	-0.170	0.006	44	39						
								-2.00	-1.00	0.00	1.00	2.00	
									Favours A		Favours B		
# Fig. P.6. Negative self-concept (NSC): EMDR vs non-specific control

Study name		Statistics for	reach stu	udy		Samp	le size		Hedges's	sg and 98	5%a	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
Kip 2013 vs control NS	C .1.195	0.303	-1.789	-0.601	0.000	26	24	-				
Scheck 1998 vs AL NS	C .0.401	0.260	-0.910	0.108	0.123	30	29					
	-0.782	0.397	-1.560	-0.005	0.049	56	53					
								-2.00	-1.00	0.00	1.00	2.00
									Favours A		Favours B	

#### Fig. P.7. PTSD: EMDR vs TAU/WL



#### Fig. P.8. PTSD: EMDR vs non-specific control



## Fig. P.9. PTSD plus 1, 2 or 3 CPTSD outcomes: EMDR vs TAU/WL

Study name	:	Statistics fo	reach sti	udy		Sampl	e size		Hedges	s's g and 95%	a	
I	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
nadi 2015 vs TAU PTSD	2.325	0.529	-3.361	-1.289	0.000	11	12	-				
berg 2007 vs WL PTSD + DR	0.479	0.431	-1.323	0.365	0.266	12	9			-0+-		
er 2002 vs WL PTSD + DR	1.525	0.315	-2.142	-0.908	0.000	27	24		+	-		
den Berg 2015 vs WL PTSD + NSC + DF	R 0.513	0.215	-0.934	-0.092	0.017	44	39					
	1.146	0.395	-1.920	-0.373	0.004	94	84					
								-4.00	-2.00	0.00	2.00	40
									Favours A		Favours B	

## Fig. P.10. PTSD plus 2 or 3 CPTSD outcomes: EMDR vs TAU/WL

Study name	:	Statistics for	r each sti	udy		Sample	e size		Hedges	's g and 95'	% <b>a</b>	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
adi 2015 vs TAU PTSD	-2.325	0.529	-3.361	-1.289	0.000	11	12	-				
den Berg 2015 vs WL PTSD + NSC + DR	-0.513	0.215	-0.934	-0.092	0.017	44	39					
	-1.355	0.904	-3.126	0.417	0.134	55	51	-				
								-4.00	-2.00	0.00	2.00	4.00
									Favours A		Favours B	

# Fig. P.11. PTSD plus 1, 2 or 3 CPTSD outcomes: EMDR vs non-specific control

Study name	:	Statistics for	r each stu	udy		Sampl	e size		Hedge	s's g and 95%	a	
ŀ	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
Ahmadi 2015 vs REM PTSD	0.058	0.420	-0.764	0.880	0.890	11	10			—¢—	-	
Kip 2013 vs control PTSD + NSC + DR	0.838	0.289	-1.405	-0.271	0.004	26	24		-			
Scheck 1998 vs AL PTSD + NSC	0.556	0.265	-1.075	-0.037	0.036	30	27					
	-0.524	0.226	-0.967	-0.081	0.020	67	61			$\bullet$		
								-4.00	-2.00	0.00	200	4.0
									Favours A		Favours B	

# Fig. P.12. PTSD plus 2 or 3 CPTSD outcomes: EMDR vs non-specific control

Study name		Statistics fo	reachst	udy		Sampl	e size		Hedges	's g and 95%	a	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
Ahmadi 2015 vs REM PTSD	0.058	0.420	-0.764	0.880	0.890	11	10			—¢—	-	
Kip 2013 vs control PTSD + NSC + E	R 0.838	0.289	-1.405	-0.271	0.004	26	24			]-		
	0.442	0.445	-1.314	0.431	0.321	37	34					
								-4.00	-2.00	0.00	200	4.00
									Favours A		Favours B	

#### Q. Forest Plots – Comparison of CBT, Exposure and EMDR

Fig. Q.1. Disturbances in relationships (DR): CBT vs Exposure alone



# Fig. Q.2. Negative self-concept (NSC): CBT vs Exposure alone

Study name		Statistics for	reach stu	idy		Samp	le size		Hedges's	g and 95	5%a	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
Resick 2002 vs Exp NSC	-0.314	0.180	-0.667	0.040	0.082	62	61		-			
310K 2002 V3 Exp 1400	-0.314	0.180	-0.667	0.040	0.082	62	61					
								-2.00	-1.00	0.00	1.00	2.00
									Favours A		Favours B	

# Fig. Q.3. PTSD: CBT vs Exposure alone

Study name		Statistics for	reach stu	udy		Samp	le size		Hedges	sgand 95°	% <b>a</b>	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
Marks 1998 vs exp PTSD	0.237	0.272	-0.297	0.771	0.384	39	20					
Foa 1999 vs exp PTSD	0.180	0.258	-0.326	0.685	0.486	41	23			-+0-	_	
Resick 2002 vs Exp PTSD	-0.179	0.179	-0.529	0.172	0.318	62	62					
Foa 2005 vs exp PTSD	-0.079	0.161	-0.394	0.237	0.625	74	79			-0-		
	-0.028	0.101	-0.225	0.170	0.784	216	184			•		
								-2.00	-1.00	0.00	1.00	2.00
									FavoursA		Favours B	

# Fig. Q.4. PTSD plus 1, 2 or 3 CPTSD outcomes: CBT vs Exposure alone

Study name	5	Statistics for	eachstu	udy		Samp	le size		Hedges	's gand 95%	a	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
Marks 1998 vs Exp PTSD + DR	0.202	0.273	-0.333	0.737	0.459	37	20			-0-		
Foa 1999 vs Exp PTSD + DR	0.272	0.261	-0.240	0.784	0.297	41	22			-0-		
Resick 2002 vs Exp PTSD + NSC	-0.246	0.180	-0.598	0.106	0.171	62	61			-0-		
Foa 2005 vs Exp PTSD + DR	-0.119	0.162	-0.437	0.199	0.463	74	78			⇔		
	-0.042	0.116	-0.269	0.185	0.719	214	181			•		
								-4.00	-2.00	0.00	2.00	4.00
									Favours A		Favours B	

# Fig. Q.5. Disturbances in relationships (DR): CBT vs EMDR

Study name		Statistics for	reach stu	dy		Samp	e size		Hedges	<u>s g and 95</u> °	% <b>a</b>	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
ower 2002 vs EMDR DR	0.596	0.293	0.023	1.169	0.042	21	27					
ijdam 2012 vs EMDR DF	0.018	0.221	-0.414	0.451	0.934	38	43			—¢—	-	
	0.275	0.287	-0.287	0.838	0.338	59	70					
								-2.00	-1.00	0.00	1.00	2.0
									FavoursA		Favours B	

## Fig. Q.6. PTSD: CBT vs EMDR

Study name	5	Statistics for	eachstu	udy		Samp	e size		Hedges'	sgand 95°	% <b>a</b>	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
Power 2002 vs EMDR PTSD	0.511	0.291	-0.059	1.081	0.079	21	27			+	-0+	
Nijdam 2012 vs EVDR PTSD	0.295	0.212	-0.120	0.711	0.164	41	48			-+C	] [	
	0.370	0.171	0.034	0.706	0.031	62	75					
								-2.00	-1.00	0.00	1.00	2.00
									Favours A		Favours B	

# Fig. Q.7. PTSD plus 1, 2 or 3 CPTSD outcomes: CBT vs EMDR

Studyname	5	Statistics for	eachstu	udy		Samp	le size		Hedges	's gand 95%	6 <b>a</b>	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
Power 2002 vs EMDR PTSD + DR	0.554	0.292	-0.019	1.126	0.058	21	27				-0-+	
Nijdam 2012 vs EMDR PTSD + DF	R 0.157	0.217	-0.268	0.581	0.470	38	43				-	
	0.307	0.193	-0.071	0.684	0.111	59	70					
								-2.00	-1.00	0.00	1.00	2.00
									Favours A		Favours B	

# Fig. Q.8. Disturbances in relationships (DR): EMDR vs Exposure alone

<u>Study name</u>	:	Statistics for	eachstu	udy		Samp	esize		Hedges	s's g and 95%	.a	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
an den Berg 2015 vs Exp DF	R -0.097	0.208	-0.505	0.311	0.640	44	47			$-\Box$		
	-0.097	0.208	-0.505	0.311	0.640	44	47					
								-2.00	-1.00	0.00	1.00	2.00
									Favours A		Favours B	

# Fig. Q.9. Negative self-concept (NSC): EMDR vs Exposure alone

<u>Studyname</u>	Statistics for each study				Sample size				Hedges's g and 95%Cl					
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B							
n den Berg 2015 vs Exp NSC	0.159	0.208	-0.249	0.568	0.444	44	47				-			
	0.159	0.208	-0.249	0.568	0.444	44	47							
								-2.00	-1.00	0.00	1.00	2.00		
									Favours A		Favours B			

# Fig. Q.10. PTSD: EMDR vs Exposure alone

Study name		Statistics fo	reachstu	dy		Sampl	e size		Hedges	sgand95%	۲ <u>۵</u>	
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
van den Berg 2015 vs Exp PTSD	0.102	0.197	-0.284	0.487	0.605	55	48		-	—	]	
	0.102	0.197	-0.284	0.487	0.605	55	48		-			
								-1.00	-0.50	0.00	0.50	1.00
									Favours A		Favours B	

# Fig. Q.11. PTSD plus 1, 2 or 3 CPTSD outcomes: EMDR vs Exposure alone

<u>Study name</u>	Statistics for each study					Sample size			Hedges's g and 95%Cl			
	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
n den Berg 2015 vs Exp PTSD + NSC + I	0.055	0.204	-0.346	0.455	0.789	44	47			— <u>(</u> )—		
	J.055	0.204	-0.346	0.455	0.789	44	47			•	▶	
								-2.00	-1.00	0.00	1.00	20
									Favours A		Favours B	

# Fig. Q.12. PTSD plus 2 or 3 CPTSD outcomes: EMDR vs Exposure alone

Study name	Statistics for each study					Sample size			Hedges's g and 95%Cl			
I	Hedges's g	Standard error	Lower limit	Upper limit	p-Value	Group-A	Group-B					
den Berg 2015 vs Exp PTSD + NSC +	0.055	0.204	-0.346	0.455	0.789	44	47			— <u>(</u> )—		
	0.055	0.204	-0.346	0.455	0.789	44	47			•		
								-2.00	-1.00	0.00	1.00	20
									Favours A		Favours B	

#### **R.** Bubble Plots – Meta-regression Moderators (univariate)

Fig. R.1. Random sequence generation (low vs unclear or high risk of bias, k=52)



Regression of Hedges's g on Random sequence

Random sequence







Allocation concealment







**Detection bias** 





Regression of Hedges's g on Reporting bias

Reporting bias







Attrition bias







Quality overall

Fig. R.7. CPTSD symptoms (PTSD alone vs various CPTSD, k=52)



Regression of Hedges's g on N CPTSD symptoms

N CPTSD symptoms

## Fig. R.8. Comparator (TAU/WL vs control, k=52)



Regression of Hedges's g on Comparator

Comparator





Regression of Hedges's g on Treatment

Treatment





Regression of Hedges's g on Group only or in addition

Group only or in addition

## Fig. R.11. Trauma onset (Adult vs child, k=48)



Regression of Hedges's g on Trauma onset

Trauma onset

#### S. Bubble Plots – Meta-regression Moderators (multivariate)

#### Fig. S.1. Overall quality (high vs low)



#### Regression of Hedges's g on Quality overall





Regression of Hedges's g on N CPTSD symptoms

N CPTSD symptoms

## Fig. S.3. Comparator (TAU/WL vs control)



Regression of Hedges's g on Comparator

Comparator

#### Fig. S.4. Treatments (individual CBT vs others)



Regression of Hedges's g on Treatment

Treatment





Regression of Hedges's g on Trauma onset

Trauma onset

#### T. Funnel Plots for Meta-analyses (where publication bias is indicated)






Fig. T.2. PTSD: Cognitive/imagery modification with or without exposure vs TAU/WL

Fig. T.3. PTSD plus 1, 2 or 3 CPTSD outcomes: Cognitive/imagery modification with or without exposure vs TAU/WL



### U. PRISMA Checklist

Section/topic	#	Checklist item	Reported	
TITLE				
Title	1	Identify the report as a systematic review, meta-analysis, or both.	Yes	
ABSTRACT				
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	Yes	
INTRODUCTION				
Rationale	3	Describe the rationale for the review in the context of what is already known.	Yes	
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	Yes	
METHODS				
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	Yes	
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	Yes	
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	Yes	
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Yes	
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	Yes	
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	Yes	
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	Yes	

Section/topic	#	Checklist item	Reported
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	Yes
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	Yes
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., $I^2$ ) for each meta-analysis.	Yes

Section/topic	#	Checklist item	Reported
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	Yes
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	Yes
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	Yes
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	Yes
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	Yes
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	Yes
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	Yes
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	Yes
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	Yes
DISCUSSION			

Section/topic	#	Checklist item	Reported
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	Yes
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	Yes
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	Yes
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	N/A

#### V. References for the 51 Included Studies

- Ahmadi K, Hazrati M, Ahmadizadeh M and Noohi S (2015) REM desensitization as a new therapeutic method for post-traumatic stress disorder: a randomized controlled trial. *Acta Medica Indonesian*, **47**, 111–119.
- Azad Marzabadi E and Hashemi Zadeh SM (2014) The Effectiveness of Mindfulness Training in Improving the Quality of Life of the War Victims with Post Traumatic stress disorder (PTSD) *Iranian Journal of Psychiatry* **9**, 228–236.
- Başoğlu M, Şalcioğlu E and Livanou M (2007) A randomized controlled study of singlesession behavioural treatment of earthquake-related post-traumatic stress disorder using an earthquake simulator. *Psychological Medicine* 37, 203–213.
- Beidel DC, Frueh BC, Neer SM, Bowers CA, Trachik B, Uhde TW and Grubaugh A (2019)Trauma management therapy with virtual-reality augmented exposure therapy for combat-related PTSD: A randomized controlled trial. *Journal of Anxiety Disorders* **61**, 64–74.
- Beidel DC, Frueh BC, Uhde TW, Wong N and Mentrikoski, JM (2011) Multicomponent behavioral treatment for chronic combat-related posttraumatic stress disorder: A randomized controlled trial. *Journal of Anxiety Disorders* 25, 224–231.
- Bryant RA, Mastrodomenico J, Hopwood S, Kenny L, Cahill C, Kandris E and Taylor K (2013) Augmenting cognitive behaviour therapy for post-traumatic stress disorder with emotion tolerance training: A randomized controlled trial. *Psychological Medicine* 43, 2153–2160.
- Butollo W, Karl R, König J, and Rosner R (2016) A Randomized Controlled Clinical Trial of Dialogical Exposure Therapy versus Cognitive Processing Therapy for Adult Outpatients Suffering from PTSD after Type I Trauma in Adulthood. *Psychotherapy and Psychosomatics* 85, 16–26.

- Cloitre M, Koenen KC, Cohen LR and Han H (2002) Skills training in affective and interpersonal regulation followed by exposure: A phase-based treatment for PTSD related to childhood abuse. *Journal of Consulting and Clinical Psychology* 70, 1067– 1074.
- Difede J, Malta LS, Best S, Henn-Haase C, Metzler T, Bryant R and Marmar C (2007) A Randomized Controlled Clinical Treatment Trial for World Trade Center Attack-Related PTSD in Disaster Workers. *The Journal of Nervous and Mental Disease* 195, 861–865.
- Dorrepaal E, Thomaes K, Smit JH, Van Balkom AJLM, Veltman DJ, Hoogendoorn AW and Draijer N (2012) Stabilizing group treatment for complex posttraumatic stress disorder related to child abuse based on psychoeducation and cognitive behavioural therapy: A multisite randomized controlled trial. *Psychotherapy and Psychosomatics* 81, 217–225.
- Duffy M, Gillespie K and Clark DM (2007) Post-traumatic stress disorder in the context of terrorism and other civil conflict in Northern Ireland: randomised controlled trial. *British Journal of Medicine* 334, 1147. doi: 10.1136/bmj.39021.846852.BE.
- Dunn NJ, Rehm LP, Schillaci J, Souchek J, Mehta P, Ashton CM, Yanasak E and Hamilton JD (2007) A randomized trial of self-management and psychoeducational group therapies for comorbid chronic posttraumatic stress disorder and depressive disorder. *Journal of Traumatic Stress* 20, 221–237.
- Dunne RL, Kenardy J and Sterling M (2012) A Randomized Controlled Trial of Cognitive-behavioral Therapy for the Treatment of PTSD in the Context of Chronic Whiplash. *The Clinical Journal of Pain* 28, 755–765.

Ehlers A, Hackmann A, Grey N, Wild J, Liness S, Albert I, Deale A, Stott R and Clark DM\_(2014) A randomized controlled trial of 7-day intensive and standard weekly cognitive therapy for PTSD and emotion-focused supportive therapy. *American Journal of Psychiatry* 171, 294–304.

Ehlers A, Clark DM, Hackmann A, McManus F and Fennell M (2005) Cognitive therapy for post-traumatic stress disorder: development and evaluation. *Behaviour Research and Therapy* 43, 413–431.

- Ehlers A, Clark DM, Hackmann A, McManus F, Fennell M, Herbert C and Mayou R (2003) A randomized controlled trial of cognitive therapy, a self-help booklet, and repeated assessments as early interventions for posttraumatic stress disorder. *Archives of General Psychiatry* **60**, 1024–1032.
- Foa EB, Dancu CV, Hembree EA, Jaycox LH, Meadows EA and Street GP (1999) A comparison of exposure therapy, stress inoculation training, and their combination for reducing posttraumatic stress disorder in female assault victims. *Journal of Consulting and Clinical Psychology* 67, 194–200.

Foa EB, Hembree E, Cahill S, Rauch S, Riggs D, Feeny N and Yadin E (2005) Randomized Trial of Prolonged Exposure for Posttraumatic Stress Disorder With and Without Cognitive Restructuring: Outcome at Academic and Community Clinics. *Journal of Consulting and Clinical Psychology* 73, 953–964.

Forbes D, Lloyd D, Nixon RD, Elliott P, Varker T, Perry D, Bryant RA and Creamer M. (2012) A multisite randomized controlled effectiveness trial of cognitive processing therapy for military-related posttraumatic stress disorder. *Journal of Anxiety Disorders* 26, 442–452.

- **Ford JD, Steinberg KL and Zhang W** (2011) A randomized clinical trial comparing affect regulation and social problem-solving psychotherapies for mothers with victimization-related PTSD. *Behavior Therapy* **42**, 560-78.
- Galovski TE, Blain LM, Mott JM, Elwood L and Houle T (2012) Manualized therapy for PTSD: Flexing the structure of cognitive processing therapy. *Journal of Consulting* and Clinical Psychology 80, 968–981.
- **Ghafoori B, Hansen MC, Garibay E and Korosteleva O** (2017) Feasibility of training frontline therapists in prolonged exposure a randomized controlled pilot study of treatment of complex trauma in diverse victims of crime and violence. *Journal of Nervous and Mental Disease* **205**, 283–293.
- Harned MS, Korslund KE and Linehan MM (2014) A pilot randomized controlled trial of Dialectical Behavior Therapy with and without the Dialectical Behavior Therapy
   Prolonged Exposure protocol for suicidal and self-injuring women with borderline personality disorder and PTSD. *Behaviour Research and Therapy* 55, 7–17.
- Hinton DE, Hofmann SG, Pollack MH and Otto MW (2009) Mechanisms of efficacy of CBT for cambodian refugees with PTSD: Improvement in emotion regulation and orthostatic blood pressure response. *CNS Neuroscience and Therapeutics* 15, 255– 263.
- Hinton DE, Hofmann SG, Rivera E, Otto MW and Pollack MH (2011) Culturally adapted CBT (CA-CBT) for Latino women with treatment-resistant PTSD: A pilot study comparing CA-CBT to applied muscle relaxation. *Behaviour Research and Therapy* 49, 275–280.
- Högberg G, Pagani M, Sundin O, Soares J, Aberg-Wistedt A, Tärnell B and Hällström T (2007) On treatment with eye movement desensitization and reprocessing of

chronic post-traumatic stress disorder in public transportation workers--a randomized controlled trial. *Nordic Journal of Psychiatry* **61**, 54–61.

- Hollifield M, Sinclair-Lian N, Warner TD and Hammerschlag R (2007) Acupuncture for posttraumatic stress disorder: a randomized controlled pilot trial. *Journal of Nervous and Mental Disease* 195, 504–513.
- Jung, K and Steil R (2013) A randomized controlled trial on cognitive restructuring and imagery modification to reduce the feeling of being contaminated in adult survivors of childhood sexual abuse suffering from posttraumatic stress disorder. *Psychotherapy & Psychosomatics* 82, 213–220.
- Keane TM, Fairbank, JA, Caddell JM and Zimering, RT (1989) Implosive (flooding) therapy reduces symptoms of PTSD in Vietnam combat veterans. *Behavior Therapy* 20, 245–260.
- Kip KE, Rosenzweig L, Hernandez DF, Shuman A, Sullivan KL, Long CJ, Taylor J,
  McGhee S, Girling SA, Wittenberg T, Sahebzamani FM, Lengacher CA, Kadel
  R and Diamond DM (2013) Randomized Controlled Trial of Accelerated Resolution
  Therapy (ART) for Symptoms of Combat-Related Post-Traumatic Stress Disorder
  (PTSD) *Military Medicine* 178, 1298–1309.
- Krakow B, Hollifield M, Johnston L, Koss M, Schrader R, Warner TD, Tandberg D,
  Lauriello J, McBride L, Cutchen L, Cheng D, Emmons S, Germain A,
  Melendrez D, Sandoval D and Prince H (2001) Imagery rehearsal therapy for
  chronic nightmares in sexual assault survivors with posttraumatic stress disorder: A
  randomized controlled trial. *Journal of the American Medical Association* 286, 537–545.

- Krupnick JL, Green BL, Stockton P, Miranda J, Krause E and Mete M (2008) Group interpersonal psychotherapy for low-income women with posttraumatic stress disorder. *Psychotherapy Research* 18, 409–507.
- Kubany ES, Hill EE and Owens JA (2003) Cognitive trauma therapy for battered women with PTSD: preliminary findings. *Journal of Traumatic Stress* 16, 81–91.
- Kubany ES, Hill EE, Owens JA, Iannce-Spencer C, McCaig MA, Tremayne KJ and
   Williams PL (2004) Cognitive Trauma Therapy for Battered Women with PTSD
   (CTT-BW) Journal of Consulting and Clinical Psychology 72, 3–18.
- Lindauer RJ, Gersons BP, van Meijel EP, Blom K, Carlier IV, Vrijlandt I and Olff M (2005) Effects of brief eclectic psychotherapy in patients with posttraumatic stress disorder: Randomized clinical trial. *Journal of Traumatic Stress* 18, 205–212.
- Marks I, Lovell K and Noshirvani H (1998) Treatment of posttraumatic stress disorder by exposure and/or cognitive restructuring: a controlled study. *Archives of General* 55, 317–325.
- McDonagh A, Friedman M, McHugo G, Ford J, Sengupta A, Mueser K, Demment CC,
  Fournier D, Schnurr PP and Descamps M (2005) Randomized Trial of CognitiveBehavioral Therapy for Chronic Posttraumatic Stress Disorder in Adult Female
  Survivors of Childhood Sexual Abuse. *Journal of Consulting and Clinical Psychology* 73, 515–524.
- Monson CM, Schnurr PP, Resick PA, Friedman MJ, Young-Xu Y and Stevens SP. (2006) Cognitive processing therapy for veterans with military-related posttraumatic stress disorder. *Journal of Consulting and Clinical Psychology* 74, 898–907.
- Mueser KT, Gottlieb JD, Xie H, Lu W, Yanos PT, Rosenberg SD, Silverstein SM, Duva SM, Minsky S, Wolfe RS and McHugo GJ (2015) Evaluation of cognitive restructuring for post-traumatic stress disorder in people with severe mental illness.

British Journal of Psychiatry 206, 501–508.

Mueser KT, Rosenberg SD, Xie H, Jankowski MK, Bolton EE, Lu W, Hamblen JL,
 Rosenberg HJ, McHugo GJ and Wolfe R (2008) A Randomized Controlled Trial of
 Cognitive-Behavioral Treatment for Posttraumatic Stress Disorder in Severe Mental
 Illness. *Journal of Consulting and Clinical Psychology* 76, 259–271.

Nijdam MJ, Gersons BP, Reitsma JB, de Jongh A and Olff M (2012) Brief eclectic psychotherapy v. eye movement desensitisation and reprocessing therapy for post-traumatic stress disorder: Randomised controlled trial. *British Journal of Psychiatry* 200, 224–231.

- Pacella ML, Armelie A, Boarts J, Wagner G, Jones T, Feeny N and Delahanty DL (2012) The impact of prolonged exposure on PTSD symptoms and associated psychopathology in people living with HIV: A randomized test of concept. *AIDS and Behavior* 16, 1327–1340.
- Power K, McGoldrick T, Brown K, Buchanan R, Sharp D, Swanson V and Karatzias A (2002) A controlled comparison of Eye Movement Desensitization and Reprocessing versus exposure plus cognitive restructuring versus waiting list in the treatment of Post-traumatic Stress Disorder. *Clinical Psychology and Psychotherapy* 9, 299–318.
- Resick PA, Nishith P, Weaver TL, Astin MC and Feuer CA (2002) A comparison of cognitive-processing therapy with prolonged exposure and a waiting condition for the treatment of chronic posttraumatic stress disorder in female rape victims. *Journal of Consulting and Clinical Psychology* **70**, 867–879.
- Scheck MM, Schaeffer JA and Gillette C (1998) Brief psychological intervention with traumatized young women: The efficacy of eye movement desensitization and reprocessing. *Journal of Traumatic Stress* 11, 25–44.

Steel C, Hardy A, Smith B, Wykes T, Rose S, Enright S, Hardcastle M, Landau S, Baksh MF, Gottlieb JD, Rose D and Mueser KT (2017) Cognitive–behaviour therapy for post-traumatic stress in schizophrenia. A randomized controlled trial. *Psychological Medicine* 47, 43–51.

Surís A, Link-Malcolm J, Chard K, Ahn C and North C (2013) A randomized clinical trial of cognitive processing therapy for veterans with PTSD related to military sexual trauma. *Journal of Traumatic Stress* **26**, 28–37.

- Talbot LS, Maguen S, Metzler TJ, Schmitz M, McCaslin SE, Richards A, Perlis ML, Posner DA, Weiss B, Ruoff L, Varbel J and Neylan TC (2014) Cognitive behavioral therapy for insomnia in posttraumatic stress disorder: a randomized controlled trial. *Sleep* 37, 327–341.
- Ter Heide FJ, Mooren TM, Kleijn W, de Jongh A and Kleber RJ (2011) EMDR versus stabilisation in traumatised asylum seekers and refugees: results of a pilot study. *European Journal of Psychotraumatology* 2, 5881. doi: 10.3402/ejpt.v2i0.5881
- Ter Heide FJ, Mooren TM, van de Schoot R, de Jongh A and Kleber RJ (2016) Eye movement desensitisation and reprocessing therapy v. stabilisation as usual for refugees: Randomised controlled trial. *British Journal of Psychiatry* 209, 311–318.
- Van den Berg DP, de Bont PA, van der Vleugel BM, de Roos C, de Jongh A, Van
  Minnen A and van der Gaag M (2015) Prolonged Exposure vs Eye Movement
  Desensitization and Reprocessing vs Waiting List for Posttraumatic Stress Disorder in
  Patients With a Psychotic Disorder. *Journal of the American Medical Association Psychiatry* 72, 259-267.

#### W. References for Excluded Studies

#### Acarturk C, Konuk E, Cetinkaya M, Senay I, Sijbrandij M, Gulen B and Cuijpers P

(2016) The efficacy of eye movement desensitization and reprocessing for posttraumatic stress disorder and depression among Syrian refugees: results of a randomized controlled trial. *Psychological Medicine* **46**, 2583- 2593.

 Acierno R, Knapp R, Tuerk P, Gilmore AK, Lejuez C, Ruggiero K, Muzzy W, Egede L, Hernandez-Tejada MA and Foa EB (2017) A non-inferiority trial of Prolonged Exposure for posttraumatic stress disorder: In person versus home-based telehealth. Behaviour Research and Therapy 89, 57–65.

Akbarian F, Bajoghli H, Haghighi M, Kalak N, Holsboer-Trachsler E and Brand S (2015) The effectiveness of cognitive behavioral therapy with respect to psychological symptoms and recovering autobiographical memory in patients suffering from post-traumatic stress disorder. *Neuropsychiatric Disease and Treatment* 11, 395–404.

- Alghamdi M, Hunt N and Thomas S (2015) The effectiveness of Narrative Exposure Therapy with traumatised firefighters in Saudi Arabia: A randomized controlled study. *Behaviour Research and Therapy* **66**, 64–71.
- Arntz A, Tiesema M and Kindt M (2007) Treatment of PTSD: A comparison of imaginal exposure with and without imagery rescripting. *Journal of Behavior Therapy and Experimental Psychiatry* 38, 345–370.
- Asukai N, Saito A, Tsuruta N, Kishimoto J and Nishikawa T (2010) Efficacy of exposure therapy for Japanese patients with posttraumatic stress disorder due to mixed traumatic events: A randomized controlled study. *Journal of Traumatic Stress* 23, 744–750.

Badura-Brack A, McDermott TJ, Becker KM, Ryan TJ, Khanna MM, Pine DS, Bar-Haim Y, Heinrichs-Graham E and Wilson TW (2018) Attention training modulates resting-state neurophysiological abnormalities in posttraumatic stress disorder. *Psychiatry Research – Neuroimaging* 271, 135–141.

Bass J, Murray S, Cole G, Bolton P, Poulton C, Robinette K, Seban J, Falb K and Annan J (2016) Economic, social and mental health impacts of an economic intervention for female sexual violence survivors in Eastern Democratic Republic of Congo. *Global Mental Health* 6, doi: 10.1017/gmh.2016.13.

- Beidel DC, Stout JW, Neer SM, Frueh BC and Lejuez C (2017) An intensive outpatient treatment program for combat-related PTSD: Trauma management therapy. *Bulletin of the Menninger Clinic* **81**, 107–122.
- Belleau EL, Chin EG, Wanklyn SG, Zambrano-Vazquez L, Schumacher JA and Coffey SF (2017) Pre-treatment predictors of dropout from prolonged exposure therapy in patients with chronic posttraumatic stress disorder and comorbid substance use disorders. *Behaviour Research and Therapy* 91, 43–50.
- Betancourt TS, McBain R, Newnham EA, Akinsulure-Smith AM, Brennan RT, Weisz JR and Hansen NB (2014) A behavioral intervention for war-affected youth in Sierra Leone: A randomized controlled trial. *Journal of the American Academy of Child and Adolescent Psychiatry* 53, 1288–1297.
- Bichescu D, Neuner F, Schauer M and Elbert T (2007) Narrative exposure therapy for political imprisonment-related chronic posttraumatic stress disorder and depression. *Behaviour Research and Therapy* 45, 2212–2220.
- Blanchard EB, Hickling EJ, Devineni T, Veazey CH, Galovski TE, Mundy E, Malta LS and Buckley TC (2003) A controlled evaluation of cognitive behavioral therapy for

posttraumatic stress in motor vehicle accident survivors. *Behaviour Research and Therapy* **41**, 79–96.

- Boden MT, Kimerling R, Jacobs-Lentz J, Bowman D, Weaver C, Carney D, Walser R and Trafton JA (2012) Seeking Safety treatment for male veterans with a substance use disorder and post-traumatic stress disorder symptomatology. *Addiction* 107, 578– 586.
- Bohus M, Dyer AS, Priebe K, Krüger A, Kleindienst N, Schmahl C, Niedtfeld I and Steil R. (2013) Dialectical behaviour therapy for post-traumatic stress disorder after childhood sexual abuse in patients with and without borderline personality disorder: A randomised controlled trial. *Psychotherapy and Psychosomatics* 82, 221-233.
- Bormann JE, Hurst S and Kelly A (2013) Responses to Mantram Repetition Program from Veterans with posttraumatic stress disorder: A qualitative analysis. *Journal of Rehabilitation Research and Development* 50, 769–784.
- Bormann JE, Oman D, Walter KH and Johnson BD (2014) Mindful Attention Increases and Mediates Psychological Outcomes Following Mantram Repetition Practice in Veterans With Posttraumatic Stress Disorder. *Medical Care* **52**, S13–S18.
- Bradley RG and Follingstad DR (2003) Group therapy for incarcerated women who experienced interpersonal violence: A pilot study. *Journal of Traumatic Stress* 16, 337–340.
- Bremner JD, Mishra S, Campanella C, Shah M, Kasher N, Evans S, Fani N, Shah AJ,
  Reiff C, Davis LL, Vaccarino V and Carmody J (2017) A Pilot Study of the
  Effects of Mindfulness-Based Stress Reduction on Post-traumatic Stress Disorder
  Symptoms and Brain Response to Traumatic Reminders of Combat in Operation
  Enduring Freedom/Operation Iraqi Freedom Combat Veterans with Post-traumatic
  Stress Disorder. *Frontiers in Psychiatry* 8, 157.

Brom D, Kleber RJ and Defares PB (1989) Brief psychotherapy for posttraumatic stress disorders. *Journal of Consulting and Clinical Psychology* **57**, 607–612.

- Brunet A (2013) Randomized controlled trial of a brief dyadic cognitive-behavioral intervention designed to prevent PTSD. *European Journal of Psych traumatology* 4, doi: 10.3402/ejpt.v4i0.21572
- Bryan CJ, Clemans TA, Hernandez AM, Mintz J, Peterson AL, Yarvis JS, Resick PA and STRONG STAR Consortium (2016) Evaluating Potential Iatrogenic Suicide Risk in Trauma-Focused Group Cognitive Behavioral Therapy for the Treatment of PTSD in Active Duty Military Personnel. *Depression and Anxiety* 33, 549–557.
- Bryant RA, Moulds ML and Nixon RV (2003) Cognitive behaviour therapy of acute stress disorder: a four-year follow-up. *Behaviour Research and Therapy* **41**, 489–494.
- Bryant RA, Felmingham K, Whitford TJ, Kemp A, Hughes G, Peduto A and Williams LM (2008) Rostral anterior cingulate volume predicts treatment response to cognitive-behavioural therapy for posttraumatic stress disorder. *Journal of Psychiatry* and Neuroscience 33, 142–146.
- Buhmann CB, Nordentoft M, Ekstroem M, Carlsson J and Mortensen EL (2016) The effect of flexible cognitive-behavioural therapy and medical treatment, including antidepressants on post-traumatic stress disorder and depression in traumatised refugees: pragmatic randomised controlled clinical trial. *The British Journal of Psychiatry* 208, 252-259.
- Carlson JG, Chemtob CM, Rusnak K, Hedlund NL and Muraoka MY (1998) Eye movement desensitization and reprocessing (EDMR) treatment for combat-related posttraumatic stress disorder. *Journal of Traumatic Stress*, **11**, 3-24.
- Castillo DT, Chee CL, Nason E, Keller J, C'de Baca J, Qualls C, Fallon SK, Haaland KY, Miller MW and Keane TM (2016) Group-delivered cognitive/exposure therapy

for PTSD in women veterans: A randomized controlled trial. *Psychological Trauma: Theory, Research, Practice and Policy* **8**, 404–412.

- Catani C, Kohiladevy M, Ruf M, Schauer E, Elbert T and Neuner F (2009) Treating children traumatized by war and Tsunami: A comparison between exposure therapy and meditation-relaxation in North-East Sri Lanka. *BioMed Central Psychiatry* 9, doi: 10.1186/1471-244X-9-22
- Chard KM (2005) An evaluation of cognitive processing therapy for the treatment of posttraumatic stress disorder related to childhood sexual abuse. *Journal of Consulting and Clinical Psychology* 73, 965–971.
- Classen C, Butler LD, Koopman C, Miller E, DiMiceli S, Giese-Davis J, Fobair P, Carlson RW, Kraemer HC and Spiegel D (2001) Supportive expressive group therapy and distress in patients with metastatic breast cancer: a randomized clinical intervention trial. *Archives of General Psychiatry* 58, 494–501.
- Cloitre M, Stovall-Mc Clough KC, Nooner K, Zorbas P, Cherry S, Jackson CL Gan W and Petkova E (2010) Treatment for PTSD related to childhood abuse: A randomized controlled trial. *American Journal of Psychiatry* **167**, 915–924.
- Cloitre M, Petkova E, Wang J and Lu Lassell F (2012) An examination of the influence of a sequential treatment on the course and impact of dissociation among women with PTSD related to childhood abuse. *Depression and Anxiety* 29,709–717.
- Coffey SF, Schumacher JA, Nosen E, Littlefield AK, Henslee AM, Lappen A, Stasiewicz PR (2016)Trauma-focused exposure therapy for chronic posttraumatic stress disorder in alcohol and drug dependent patients: A randomized controlled trial. *Psychology of Addictive Behaviors* **30**, 778-790.

- Cook JM, Harb GC, Gehrman PR, Cary MS, Gamble GM, Forbes D and Ross RJ (2010) Imagery rehearsal for posttraumatic nightmares: A randomized controlled trial. *Journal of Traumatic Stress* 23, 553–563.
- **Cooper NA and Clum GA** (1989) Imaginal flooding as a supplementary treatment for PTSD in combat veterans: a controlled study. *Behavior Therapy* **20**, 381–391.
- Cooper AA, Kline AC, Graham B, Bedard-Gilligan M, Mello PG, Feeny NC and Zoellner LA (2017) Homework "Dose," Type, and Helpfulness as Predictors of Clinical Outcomes in Prolonged Exposure for PTSD. *Behavior Therapy* 48, 182–194.
- Cottraux J, Note I, Yao SN, de Mey-Guillard C, Bonasse F, Djamoussian D, Mollard E,
   Note B and Chen Y (2008) Randomized controlled comparison of cognitive behavior
   therapy with rogerian supportive therapy in chronic post-traumatic stress disorder: A
   2-year follow-up. *Psychotherapy and Psychosomatics* 77, 101–110.
- De Bont PA, van den Berg DP, van der Vleugel BM, de Roos C, de Jongh A, van der Gaag M and van Minnen AM (2016) Prolonged exposure and EMDR for PTSD v. a PTSD waiting-list condition: Effects on symptoms of psychosis, depression and social functioning in patients with chronic psychotic disorders. *Psychological Medicine* 46, 2411–2421.
- Devilly, GJ, Spence SH and Rapee RM (1998) Statistical and reliable change with eye movement desensitization and reprocessing: Treating trauma within a veteran population. *Behaviour Therapy* 29, 435–455.
- **Devilly GJ and Spence SH** (1999) The relative efficacy and treatment distress of EMDR and a cognitive-behavior trauma treatment protocol in the amelioration of posttraumatic stress disorder. *Journal of Anxiety Disorders* **13**,131–157.

- Echeburúa E, de Corral P, Zubizarreta I and Sarasua B (1997) Psychological treatment of chronic posttraumatic stress disorder in victims of sexual aggression. *Behavior Modification* 21, 433–456.
- Edmond T and Rubin A (2004) Assessing the long-term effects of EMDR: results from an 18-month follow-up study with adult female survivors of CSA. *Journal of Child Sexual Abuse* **13**, 69–86.
- Edmond T, Rubin A and Wambach KG (1999)The effectiveness of EMDR with adult female survivors of childhood sexual abuse. *Social Work Research* 23, 103–116.
- Ertl V, Pfeiffer A, Schauer E, Elbert T and Neuner F (2011) Community-implemented trauma therapy for former child soldiers in Northern Uganda: a randomized controlled trial. *Journal of the American Medical Association* **306**, 503-512.
- Fecteau G and Nicki R (1999) Cognitive behavioural treatment of post-traumatic stress disorder after motor vehicle accident. *Behavioural and Cognitive Psychotherapy* 27, 201–214.
- Feeny NC Zoellner LA and Foa EB (2002) Treatment Outcome for Chronic PTSD Among Female Assault Victims With Borderline Personality Characteristics : a Preliminary Examination. *Journal of Personality Disorders* 16, 30-40.
- Feske U(2008) Treating low-income and minority women with posttraumatic stress disorder: A pilot study comparing prolonged exposure and treatment as usual conducted by community therapists. *Journal of Interpersonal Violence*, 23, 1027–1040.
- Foa EB, Asnaani A, Rosenfield D, Zandberg LJ, Gariti P and Imms P (2017) Concurrent varenicline and prolonged exposure for patients with nicotine dependence and PTSD: A randomized controlled trial. *Journal of Consulting and Clinical Psychology* 85, 862–872.

- Foa EB, Rothbaum BO, Riggs DS and Murdock TB (1991) Treatment of posttraumatic stress disorder in rape victims: A comparison between cognitive-behavioral procedures and counselling. *Journal of Consulting and Clinical Psychology* 59, 715– 723.
- Foa EB and Rauch SA (2004) Cognitive changes during prolonged exposure versus prolonged exposure plus cognitive restructuring in female assault survivors with posttraumatic stress disorder. *Journal of Consulting and Clinical Psychology* 72, 879– 884.
- Foa EB, Yusko DA, McLean CP, Suvak MK, Bux DA Jr, Oslin D, O'Brien CP, Imms P, Riggs DS and Volpicelli J (2013) Concurrent naltrexone and prolonged exposure therapy for patients with comorbid alcohol dependence and PTSD: a randomized clinical trial. *Journal of the American Medical Association*, **310**, 488–495.
- Fredman SJ, Pukay-Martin ND, Macdonald A, Wagner AC, Vorstenbosch V and Monson CM (2016) Partner Accommodation Moderates Treatment Outcomes for Couple Therapy for Posttraumatic Stress Disorder. *Journal of Consulting & Clinical Psychology* 84, 79–87.
- Gamito P, Oliveira J, Rosa P, Morais D, Duarte N, Oliveira S and Saraiva T (2010)
   PTSD Elderly War Veterans: A Clinical Controlled Pilot Study. *Cyberpsychology, Behavior, and Social Networking*, 13, 43–48.
- Gersons BP, Carlier IV, Lamberts RD and van der Kolk BA (2000) Randomized clinical trial of brief eclectic psychotherapy for police officers with posttraumatic stress disorder. *Journal of Traumatic Stress* 13, 333-47.
- Gilboa-Schechtman E, Foa EB, Shafran N, Aderka IM, Powers MB, Rachamim L, Rosenbach L, Yadin E and Apter A (2010) Prolonged exposure versus dynamic

therapy for adolescent PTSD : a pilot randomized controlled trial. *Journal of the American Academy of Child and Adolescent Psychiatry* **49**, 1034–1042.

- Glynn SM, Eth S, Randolph ET, Foy DW, Urbaitis M, Boxer L, Paz GG, Leong GB,
  Firman G, Salk JD, Katzman JW and Crothers J (1999) A test of behavioral
  family therapy to augment exposure for combat-related posttraumatic stress disorder.
  Journal of Consulting and Clinical Psychology 67, 243–251.
- Goldstein LA, Mehling WE, Metzler TJ, Cohen BE, Barnes DE, Choucroun GJ, Silver
   A, Talbot LS, Maguen S, Hlavin JA, Chesney MA and Neylan TC (2018)
   Veterans Group Exercise: A randomized pilot trial of an Integrative Exercise program
   for veterans with posttraumatic stress. *Journal of Affective Disorders* 227, 345-352.
- Gutner CA, Gallagher MW, Baker AS, Sloan DM and Resick PA (2016) Time Course of Treatment Dropout in Cognitive-Behavioral Therapies for Posttraumatic Stress
   Disorder. *Psychological Trauma: Theory, Research, Practice and Policy* 8, 115–121.
- Hien DA, Cohen LR, Miele GM, Litt LC and Capstick C (2004) Promising Treatments for Women With Comorbid PTSD and Substance Use Disorders. *The American Journal* of Psychiatry 161, 1426-32.
- Hien DA, Lopez-Castro T, Papini S, Gorman B and Ruglass LM (2017) Emotion dysregulation moderates the effect of cognitive behavior therapy with prolonged exposure for co-occurring PTSD and substance use disorders. *Journal of Anxiety Disorders* 52, 53–61
- Hien DA, Ruglass L and Back S (2017) Concurrent treatment with prolonged exposure for co-occurring PTSD and substance use disorders: A randomized clinical trial. *Drug* and Alcohol Dependence 171, e88-e89.
- Hien DA, Wells EA, Jiang H, Suarez-Morales L, Campbell AN, Cohen LR, Miele GM, Killeen T, Brigham GS, Zhang Y, Hansen C, Hodgkins C, Hatch-Maillette M,

Brown C, Kulaga A, Kristman-Valente A, Chu M, Sage R, Robinson JA, Liu D and Nunes EV (2009) Multi-site randomized trial of behavioral interventions for women with co-occurring PTSD and substance use disorders. Journal of Consulting and Clinical Psychology 77, 607–619. <u>https://doi.org/10.1037/a0016227</u>

- Hinton DE, Chhean D, Pich V, Safren SA, Hofmann SG and Pollack MH (2005) A randomized controlled trial of cognitive-behavior therapy for Cambodian refugees with treatment-resistant PTSD and panic attacks: A cross-over design. *Journal of Traumatic Stress* 18, 617–629.
- Holliday R, Link-Malcolm J, Morris EE and Surís A (2014) Effects of Cognitive Processing Therapy on PTSD-Related Negative Cognitions in Veterans With Military Sexual Trauma. *Military Medicine* **179**, 1077–1082.
- Holliday R, Williams R, Bird J, Mullen K and Surís A (2015) The role of cognitive processing therapy in improving psychosocial functioning, health, and quality of life in veterans with military sexual trauma-related posttraumatic stress disorder. *Psychological Services* 12, 428-34.
- Ironson G, Freund B, Strauss JL and Williams J (2002) Comparison of two treatments for traumatic stress: A community-based study of EMDR and prolonged exposure. *Journal of Clinical Psychology* 58, 113–128.
- Jacob N, Neuner F, Maedl A, Schaal S and Elbert T (2014) Dissemination of psychotherapy for trauma spectrum disorders in postconflict settings: A randomized controlled trial in Rwanda. *Psychotherapy and Psychosomatics* 83, 354–363.
- Jensen JA (1994) An investigation of eye movement desensitization and reprocessing (EMD/R) as a treatment for posttraumatic stress disorder (PTSD) symptoms of Vietnam combat veterans. *Behaviour Therapy* **25**, 311–325.

- Jindani F, Turner N and Khalsa SB (2015) A yoga intervention for posttraumatic stress: a preliminary randomized control trial. *Evidence Based Complementary and Alternative Medicine* 2015, 351746. doi: 10.1155/2015/351746
- Johnson DM, Johnson NL, Perez SK, Palmieri PA and Zlotnick C (2016) Comparison of Adding Treatment of PTSD During and After Shelter Stay to Standard Care in Residents of Battered Women's Shelters: Results of a Randomized Clinical Trial. *Journal of Traumatic Stress* 29, 365–373.
- Johnson DM, Zlotnick C and Perez S (2011) Cognitive-Behavioral Treatment of PTSD in Residents of Battered Women Shelters: Results of a Randomized Clinical Trial. *Journal of Consulting and Clinical Psychology* 79, 542–551.
- Kearney DJ, McDermott K, Malte C, Martinez M and Simpson T (2013) Effects of Participation in a Mindfulness Program for Veterans With Posttraumatic Stress Disorder: A Randomized Controlled Pilot Study. *Journal of Clinical Psychology* 69, 14–27.
- Kearney DJ, Simpson TL, Malte CA, Felleman B, Martinez ME and Hunt SC (2016) Mindfulness-based Stress Reduction in Addition to Usual Care Is Associated with Improvements in Pain, Fatigue, and Cognitive Failures Among Veterans with Gulf War Illness. *American Journal of Medicine* 129, 204–214.
- Kip KE, Rosenzweig L, Hernandez DF, Shuman A, Diamond DM, Girling SA, Sullivan KL, Wittenberg T, Witt AM, Lengacher CA, Anderson B and McMillan SC (2014) Accelerated Resolution Therapy for treatment of pain secondary to symptoms of combat-related posttraumatic stress disorder. *European Journal of Psychotraumatology* 5, 10.3402/ejpt.v5.24066. doi:10.3402/ejpt.v5.24066

- König J, Onnen M, Karl R, Rosner R and Butollo W (2016) Interpersonal Subtypes and Therapy Response in Patients Treated for Posttraumatic Stress Disorder. *Clinical Psychology & Psychotherapy* 23, 97–106.
- Kruse J, Joksimovic L, Cavka M, Wöller W and Schmitz N (2009) Effects of traumafocused psychotherapy upon war refugees. *Journal of Traumatic Stress* 22, 585–592.
- Lange A, Rietdijk D, Hudcovicova M, van de Ven JP, Schrieken B and Emmelkamp PM (2003) Interapy: A controlled randomized trial of the standardized treatment of posttraumatic stress through the internet. *Journal of Consulting and Clinical Psychology* 71, 901–909.
- Lee C, Gavriel H, Drummond P, Richards J and Greenwald R (2002) Treatment of PTSD: Stress inoculation training with prolonged exposure compared to EMDR. *Journal of Clinical Psychology* 58, 1071–1089.
- Levi O, Bar-Haim Y, Kreiss Y and Fruchter E (2016) Cognitive-Behavioural Therapy and Psychodynamic Psychotherapy in the Treatment of Combat-Related Post-Traumatic Stress Disorder: A Comparative Effectiveness Study. *Clinical Psychology* & *Psychotherapy* 23, 298-307.
- Liedl A, Müller J, Morina N, Karl A, Denke C and Knaevelsrud C (2011) Physical Activity within a CBT Intervention Improves Coping with Pain in Traumatized Refugees: Results of a Randomized Controlled Design. *Pain Medicine* **12**, 234–245.
- Litz BT, Engel CC, Bryant RA and Papa A (2007) A randomized, controlled proof-ofconcept trial of an internet-based, therapist-assisted self-management treatment for posttraumatic stress disorder. *American Journal of Psychiatry* **164**, 1676–1683.
- Lovell K, Marks IM, Noshirvani H, Thrasher S, and Livanou M (2001) Do cognitive and exposure treatments improve various PTSD symptoms differently? A randomized controlled trial. *Behavioural and Cognitive Psychotherapy* **29**, 107–112.

- Macdonald A, Pukay-Martin ND, Wagner AC, Fredman SJ, Monson CM (2016)
   Cognitive-behavioral conjoint therapy for PTSD improves various PTSD symptoms and trauma-related cognitions: Results from a randomized controlled trial. *Journal of Family Psychology* 30, 157–162.
- Marcus SV, Marquis P and Sakai C (1997) Controlled study of treatment of PTSD using EMDR in an HMO setting. *Psychotherapy* 34, 307–315.
- Markowitz JC, Petkova E, Neria Y, Van Meter PE, Zhao Y, Hembree E, Lovell K,
  Biyanova T and Marshall RD (2015) Is exposure necessary? A randomized clinical trial of interpersonal psychotherapy for PTSD. *American Journal of Psychiatry* 172, 430–440.
- Markowitz JC, Neria Y, Lovell K, Van Meter PE and Petkova E (2017) History of sexual trauma moderates psychotherapy outcome for posttraumatic stress disorder. *Depression and Anxiety* 34, 692-700.
- Maxwell K, Callahan JL, Holtz P, Janis BM, Gerber MM and Connor DR (2016) Comparative study of group treatments for posttraumatic stress disorder. *Psychotherapy (Chicago, Ill.)* **53**, 433–445.
- McGovern MP, Lambert-Harris C, Xie H, Meier A, McLeman B and Saunders E (2015) A randomized controlled trial of treatments for co-occurring substance use disorders and post-traumatic stress disorder. *Addiction* **110**, 1194–1204.

McLay RN, Baird A, Webb-Murphy J, Deal W, Tran L, Anson H, Klam W and Johnston S (2017) A Randomized, Head-to-Head Study of Virtual Reality Exposure Therapy for Posttraumatic Stress Disorder. *Cyberpsychology, Behavior, and Social Networking* 20, 218–224.

- McLean CP, Su YJ and Foa EB (2014) Posttraumatic stress disorder and alcohol dependence: Does order of onset make a difference? *Journal of Anxiety Disorders* 28, 894–901.
- Meier A, McGovern MP, Lambert-Harris C, McLeman B, Franklin A, Saunders EC and Xie H (2015) Adherence and competence in two manual-guided therapies for co-occurring substance use and posttraumatic stress disorders: Clinician factors and patient outcomes. *American Journal of Drug and Alcohol Abuse* **4**1, 527–534.
- Mills KL, Teesson M, Back SE, Brady KT, Baker AL, Hopwood S, Sannibale C, Barrett EL, Merz S, Rosenfeld J and Ewer PL (2012) Integrated exposure-based therapy for co-occurring posttraumatic stress disorder and substance dependence: A randomized controlled trial. *Journal of the American Medical Association*, **308**, 690–699.
- Monson CM, Fredman SJ, Macdonald A, Pukay-Martin ND, Resick PA and Schnurr PP (2012) Effect of cognitive-behavioral couple therapy for PTSD: a randomized controlled trial. *Journal of the American Medical Association* **308**, 700-709.
- Moradi AR, Moshirpanahi S, Parhon H, Mirzaei J, Dalgleish T and Jobson L (2014) A pilot randomized controlled trial investigating the efficacy of Memory Specificity Training in improving symptoms of posttraumatic stress disorder. *Behaviour Research and Therapy* **56**, 68-74.
- Morath J, Gola H, Sommershof A, Hamuni G, Kolassa S, Catani C, Adenauer H, Ruf-Leuschner M, Schauer M, Elbert T, Groettrup M and Kolassa IT (2014) The effect of trauma-focused therapy on the altered T cell distribution in individuals with PTSD: Evidence from a randomized controlled trial. *Journal of Psychiatric Research* 54, 1–10.

# Morland LA, Mackintosh MA, Greene CJ, Rosen CS, Chard KM, Resick P and Frueh BC (2014) Cognitive processing therapy for posttraumatic stress disorder delivered to rural veterans via telemental health: a randomized noninferiority clinical trial. *Journal of Clinical Psychiatry* 75, 470–476.

- Moser JS, Cahill SP and Foa EB (2010) Evidence for poorer outcome in patients with severe negative trauma-related cognitions receiving prolonged exposure plus cognitive restructuring: implications for treatment matching in posttraumatic stress disorder. *Journal of Nervous and Mental Disease* **198**, 72–75.
- Nacasch N, Foa EB, Huppert JD, Tzur D, Fostick L, Dinstein Y, Polliack M and Zohar J. (2011) Prolonged exposure therapy for combat- and terror-related posttraumatic stress disorder: A randomized control comparison with treatment as usual. *Journal of Clinical Psychiatry* 72, 1174–1180.
- Nacasch N, Huppert JD, Su YJ, Kivity Y, Dinshtein Y, Yeh R and Foa EB (2015) Are
   60-Minute Prolonged Exposure Sessions With 20-Minute Imaginal Exposure to
   Traumatic Memories Sufficient to Successfully Treat PTSD? A Randomized
   Noninferiority Clinical Trial. *Behaviour Therapy* 46, 328–341.
- Neuner F, Kurreck S, Ruf M, Odenwald M, Elbert T and Schauer M (2010) Can asylumseekers with posttraumatic stress disorder be successfully treated? A randomized controlled pilot study. *Cognitive Behaviour Therapy* **39**, 81-91.
- Neuner F, Onyut PL, Ertl V, Odenwald M, Schauer E and Elbert T (2008) Treatment of Posttraumatic Stress Disorder by Trained Lay Counselors in an African Refugee Settlement: A Randomized Controlled Trial. *Journal of Consulting and Clinical Psychology* 76, 686–694.
- Neuner F, Schauer M, Klaschik C, Karunakara U and Elbert T (2004) A comparison of narrative exposure therapy, supportive counseling, and psychoeducation for treating

posttraumatic stress disorder in an African refugee settlement. *Journal of Consulting and Clinical Psychology* **72**, 579–587.

- Nijdam MJ, van der Meer CAI, van Zuiden M, Dashtgard P, Medema D, Qing Y,
  Zhutovsky P, Bakker A and Olff M (2018) Turning wounds into wisdom:
  Posttraumatic growth over the course of two types of trauma-focused psychotherapy
  in patients with PTSD. *Journal of Affective Disorders* 227, 424–431.
- Nosen E, Littlefield AK, Schumacher JA, Stasiewicz PR and Coffey SF (2014) Treatment of co-occurring PTSD-AUD: Effects of exposure-based and non-trauma focused psychotherapy on alcohol and trauma cue-reactivity. *Behaviour Research and Therapy* **61**, 35–42.
- Øktedalen T, Hoffart A and Langkaas TF (2015) Trauma-related shame and guilt as timevarying predictors of posttraumatic stress disorder symptoms during imagery exposure and imagery rescripting. A randomized controlled trial. *Psychotherapy Research* 25, 518–532.
- Pabst A, Schauer M, Bernhardt K, Ruf-Leuschner M, Goder R, Elbert T, Rosentraeger R, Robjant K, Aldenhoff J and Seeck-Hirschner M (2014) Evaluation of narrative exposure therapy (NET) for borderline personality disorder with comorbid posttraumatic stress disorder. *Clinical Neuropsychiatry* 11, 108–117.
- Paivio SC, Jarry JL, Chagigiorgis H, Hall I and Ralston M (2010) Efficacy of two versions of emotion-focused therapy for resolving child abuse trauma. *Psychotherapy Research* 20, 353–366.
- **Paunovic N and Ost LG (2001)** Cognitive-behavior therapy vs exposure therapy in the treatment of PTSD in refugees. *Behaviour Research and Therapy* **39**, 1183–1197.

- Peniston EG and Kulkosky PJ (1991) Alpha-theta brainwave neuro-feedback therapy for Vietnam veterans with combat-related post-traumatic stress disorder. *Medical Psychotherapy: An International Journal*, 4, 47–60.
- Polusny MA, Erbes CR, Thuras P, Moran A, Lamberty GJ, Collins RC, Rodman JL and Lim KO (2015) Mindfulness-Based Stress Reduction for Posttraumatic Stress Disorder Among Veterans. A Randomized Clinical Trial. *Journal of the American Medical Association* 314, 456-465.
- Possemato K, Bergen-Cico D, Treatman S, Allen C, Wade M and Pigeon W (2016) A Randomized Clinical Trial of Primary Care Brief Mindfulness Training for Veterans With PTSD. *Journal of Clinical Psychology* 72, 179-93.
- Pruiksma KE, Taylor DJ, Wachen JS, Mintz J, Young-McCaughan S, Peterson AL,
  Yarvis JS, Borah EV, Dondanville KA, Litz BT, Hembree EA and Resick PA
  (2016) Residual Sleep Disturbances Following PTSD Treatment in Active Duty
  Military Personnel. *Psychological Trauma: Theory, Research, Practice, and Policy* 8, 697-701.
- Reger GM, Koenen-Woods P, Zetocha K, Smolenski DJ, Holloway KM, Rothbaum BO, Difede J, Rizzo AA, Edwards-Stewart A, Skopp NA, Mishkind M, Reger MA and Gahm GA (2016) Randomized controlled trial of prolonged exposure using imaginal exposure vs. virtual reality exposure in active duty soldiers with deployment-related posttraumatic stress disorder (PTSD) *Journal of Consulting and Clinical Psychology* 84, 946–959.
- Resick PA, Galovski TE, Uhlmansiek MO, Scher CD, Clum GA and Young-Xu Y (2008) A randomized clinical trial to dismantle components of cognitive processing therapy for posttraumatic stress disorder in female victims of interpersonal violence. *Journal of Consulting and Clinical Psychology* **76**, 243–258.

**Resick PA, Nishith P and Griffin MG** (2003) How well does cognitive-behavioral therapy treat symptoms of complex PTSD? An examination of child sexual abuse survivors within a clinical trial. *CNS Spectrums* **8**, 340–355.

Resick PA, Wachen JS, Dondanville KA, Pruiksma KE, Yarvis JS, Peterson AL, Mintz J and the STRONG STAR Consortium, Borah EV, Brundige A, Hembree EA, Litz BT, Roache JD and Young-McCaughan S (2017) Effect of Group vs Individual Cognitive Processing Therapy in Active-Duty Military Seeking Treatment for Posttraumatic Stress Disorder: a Randomized Clinical Trial. *Journal of the American Medical Association Psychiatry* 74, 28–36.

Resick PA, Wachen JS, Mintz J, Young-McCaughan S, Roache JD, Borah AM, Borah EV, Dondanville KA, Hembree EA, Litz BT and Peterson AL (2015) A

randomized clinical trial of group cognitive processing therapy compared with group present-centered therapy for PTSD among active duty military personnel. *Journal of Consulting and Clinical Psychology* **83**, 1058-1068.

- Roberts NP, Roberts PA, Jones N and Bisson JI (2016) Psychological therapies for posttraumatic stress disorder and comorbid substance use disorder. *Clinical Psychology Review* 38, 25–38.
- Rothbaum BO (1997) A controlled study of eye movement desensitization and reprocessing in the treatment of posttraumatic stress disordered sexual assault victims. *Bulletin of the Menninger Clinic* **61**, 317–334.
- Rothbaum BO, Astin MC and Marsteller F (2005) Prolonged exposure versus Eye Movement Desensitization and Reprocessing (EMDR) for PTSD rape victims. *Journal of Traumatic Stress* 18, 607–616.

## Rothbaum BO, Cahill SP, Foa EB, Davidson JR, Compton J, Connor KM, Astin MC and Hahn CG (2006) Augmentation of sertraline with prolonged exposure in the treatment of posttraumatic stress disorder. *Journal of Traumatic Stress* 19, 625–638.

- Rothbaum BO, Price M, Jovanovic T, Norrholm SD, Gerardi M, Dunlop B, Davis M, Bradley B, Duncan EJ, Rizzo A and Ressler KJ (2014) A Randomized, Double-Blind Evaluation of d-Cycloserine or Alprazolam Combined With Virtual Reality Exposure Therapy for Posttraumatic Stress Disorder in Iraq and Afghanistan War Veterans. *American Journal of Psychiatry* 171, 640-648.
- Ruglass LM, Shevorykin A, Radoncic V, Smith KM, Smith PH, Galatzer-Levy IR, Papini S and Hien DA (2017) Impact of cannabis use on treatment outcomes among adults receiving cognitive-behavioral treatment for PTSD and substance use disorders. *Journal of Clinical Medicine* 6, doi.org/10.3390/jcm6020014
- Sack M, Zehl S, Otti A, Lahmann C, Henningsen P, Kruse J and Stingl M (2016) A comparison of dual attention, eye movements, and exposure only during eye movement desensitization and reprocessing for posttraumatic stress disorder: Results from a randomized clinical trial. *Psychotherapy and Psychosomatics* 85, 357–365.
- Sannibale C, Teesson M, Creamer M, Sitharthan T, Bryant RA, Sutherland K, Taylor K, Bostock-Matusko D, Visser A and Peek-O'Leary M (2013) Randomized controlled trial of cognitive behaviour therapy for comorbid post-traumatic stress disorder and alcohol use disorders. *Addiction* 108, 1397–1410.
- Sautter FJ, Glynn SM, Cretu JB, Senturk D and Vaught AS (2015) Efficacy of structured approach therapy in reducing PTSD in returning veterans: A randomized clinical trial. *Psychological Services* **12**, 199–212.

- Schaal S, Elbert T and Neuner F (2009) Narrative exposure therapy versus interpersonal psychotherapy. A pilot randomized controlled trial with Rwandan genocide orphans. Psychotherapy and Psychosomatics 78, 298–306.
- Schneier FR, Neria Y, Pavlicova M, Hembree E, Suh EJ, Amsel L and Marshall RD (2012) Combined prolonged exposure therapy and paroxetine for PTSD related to the World Trade Center attack: A randomized controlled trial. *American Journal of Psychiatry* 169, 80–88.
- Schnurr PP (2007) Cognitive behavioral therapy for posttraumatic stress disorder in women: a randomized controlled trial. *Journal of the American Medical Association* 297, 820– 830.
- Schnurr PP, Friedman MJ, Foy DW, Shea MT, Hsieh FY, Lavori PW, Glynn SM, Wattenberg M and Bernardy NC (2003) Randomized trial of trauma-focused group therapy for posttraumatic stress disorder: Results from a department of veterans affairs cooperative study. Archives of General Psychiatry 60, 481–489.
- Schnurr PP, Friedman MJ, Engel CC, Foa EB, Shea MT, Chow BK, Resick PA, Thurston V, Orsillo SM, Haug R, Turner C and Bernardy N (2007) Cognitive behavioral therapy for posttraumatic stress disorder in women - A randomized controlled trial. *Journal of the American Medical Association* 297, 820–830.
- Schnyder U, Müller J, Maercker A and Wittmann L (2011) Brief eclectic psychotherapy for PTSD: A randomized controlled trial. *Journal of Clinical Psychiatry* **72**, 564–566.
- Scott JC, Harb G, Brownlow JA, Greene J, Gur RC and Ross RJ (2017) Verbal memory functioning moderates psychotherapy treatment response for PTSD-Related nightmares. *Behaviour Research and Therapy* **91**, 24–32.
- **Shea MT, Lambert J and Reddy MK** (2013) A randomized pilot study of anger treatment for Iraq and Afghanistan veterans. *Behaviour Research and Therapy* **51**, 607–613.

- Shnaider P, Sijercic I, Wanklyn SG, Suvak MK, & Monson CM (2016)The Role of Social Support in Cognitive-Behavioral Conjoint Therapy for Posttraumatic Stress Disorder. *Behaviour Therapy* 48, 285–294.
- Sikkema KJ, Hansen NB, Kochman A, Tarakeshwar N, Neufeld S, Meade CS and Fox AM (2007) Outcomes from a group intervention for coping with HIV/AIDS and childhood sexual abuse: Reductions in traumatic stress. *AIDS and Behaviour* 11, 49– 60.
- Sin J, Spain D, Furuta M, Murrells T and Norman I (2017) Psychological interventions for post-traumatic stress disorder (PTSD) in people with severe mental illness. *Cochrane Database of Systematic Reviews* 1, CD011464. doi 10.1002/14651858.CD011464.pub2.
- Slade EP, Gottlieb JD, Lu W, Yanos PT, Rosenberg S, Silverstein SM, Minsky SK and Mueser KT (2017) Cost-Effectiveness of a PTSD Intervention Tailored for Individuals With Severe Mental Illness. *Psychiatric Services* 68, 1225-1231.
- Sloan DM, Marx BP, Bovin MJ, Feinstein BA and Gallagher MW (2012) Written exposure as an intervention for PTSD: A randomized clinical trial with motor vehicle accident survivors. *Behaviour Research and Therapy* 50, 627–635.
- Spence J, Titov N, Dear BF, Johnston L, Solley K, Lorian C, Wootton B, Zou J and Schwenke G (2011) Randomized controlled trial of Internet-delivered cognitive behavioral therapy for posttraumatic stress disorder. *Depression and Anxiety* 28, 541-550.
- Steinert C, Bumke PJ, Hollekamp RL, Larisch A, Leichsenring F, Mattheß H, Sek S, Sodemann U, Stingl M, Ret T, Vojtová H, Wöller W and Kruse J (2017) Resource activation for treating post-traumatic stress disorder, co-morbid symptoms and impaired functioning: A randomized controlled trial in Cambodia. *Psychological Medicine* 47, 553–564.

Tarrier N, Pilgrim H, Sommerfield C, Faragher B, Reynolds M, Graham E and Barrowclough C (1999) A randomized trial of cognitive therapy and imaginal exposure in the treatment of chronic posttraumatic stress disorder. *Journal of Consulting* 67, 13–18.

- Tarrier N, Sommerfield C, Pilgrim H and Humphreys L (1999) Cognitive therapy or imaginal exposure in the treatment of post-traumatic stress disorder. Twelve-month follow-up. *British Journal of Psychiatry* 175, 571–575.
- Taylor S, Thordarson DS, Maxfield L, Fedoroff IC, Lovell K and Ogrodniczuk J (2003) Comparative efficacy, speed, and adverse effects of three PTSD treatments: Exposure therapy, EMDR, and relaxation training. *Journal of Consulting and Clinical Psychology* 71, 330–338.
- van den Berg DP, de Bont PA, van der Vleugel BM, de Roos C, de Jongh A, van Minnen A and van der Gaag M (2016) Trauma-focused treatment in PTSD patients with psychosis: Symptom exacerbation, adverse events, and revictimization. *Schizophrenia Bulletin* 42, 693–702.
- van der Kolk B, Hamlin E, Gapen M, Hodgdon H, Musicaro R, Suvak M and Spinazzola J (2016) A Randomized Controlled Study of Neurofeedback for Chronic PTSD. *PLoS ONE* 11, 10.1371/journal.pone.0166752.
- van der Kolk BA, Spinazzola J, Blaustein ME, Hopper JW, Hopper EK, Korn DL and Simpson WB (2007) A randomized clinical trial of eye movement desensitization and reprocessing (EMDR), fluoxetine, and pill placebo in the treatment of posttraumatic stress disorder: treatment effects and long-term maintenance. *Journal of Clinical Psychiatry* 68, 37-46.
- van Emmerik AA, Kamphuis JH and Emmelkamp PM (2008) Treating Acute Stress Disorder and Posttraumatic Stress Disorder with Cognitive Behavioral Therapy or

Structured Writing Therapy: A Randomized Controlled Trial. *Psychotherapy and Psychosomatics* **77**, 93–100.

- Vaughan K, Armstrong MS, Gold R, O'Connor N, Jenneke W and Tarrier N (1994) A trial of eye movement desensitization compared to image habituation training and applied muscle relaxation in post-traumatic stress disorder. *Journal of Behaviour Therapy and Experimental Psychiatry* 25, 283–291.
- Wahbeh H, Goodrich E, Goy E and Oken BS (2016) Mechanistic Pathways of Mindfulness Meditation in Combat Veterans With Posttraumatic Stress Disorder. *Journal of Clinical Psychology* 72, 365–383.
- Wells A, Walton D, Lovell K and Proctor D (2015) Metacognitive therapy versus prolonged exposure in adults with chronic post-traumatic stress disorder: A parallelrandomized controlled trial. *Cognitive Therapy and Research* **39**, 70–80.
- Wilson SA, Becker LA and Tinker RH (1995) Eye movement desensitization and reprocessing (EMDR) treatment for psychologically traumatized individuals. *Journal* of Consulting and Clinical Psychology 63, 928–937.
- Wilson SA, Becker LA and Tinker RH (1997) Fifteen-month follow-up of eye movement desensitization and reprocessing (EMDR) treatment for posttraumatic stress disorder and psychological trauma. *Journal of Consulting and Clinical Psychology* 65, 1047– 1056.
- Wolff N, Huening J, Shi J, Frueh BC, Hoover DR and McHugo G (2015) Implementation and effectiveness of integrated trauma and addiction treatment for incarcerated men. *Journal of Anxiety Disorders* 30, 66–80.
- Zang Y, Hunt N and Cox T (2013) A randomised controlled pilot study: the effectiveness of narrative exposure therapy with adult survivors of the Sichuan earthquake. BMC Psychiatry 31, doi:10.1186/1471-244X-13-41
- Zang Y, Hunt N and Cox T (2014) Adapting narrative exposure therapy for Chinese earthquake survivors: a pilot randomised controlled feasibility study. *BMC Psychiatry* 14, 262. doi: 10.1186/s12888-014-0262-3.
- Zang Y, Yu J, Chazin D, Asnaani A, Zandberg LJ and Foa EB (2017) Changes in coping behavior in a randomized controlled trial of concurrent treatment for PTSD and alcohol dependence. *Behaviour Research and Therapy* 90, 9–15.
- Zlotnick C, Johnson J and Najavits LM (2009) Randomized controlled pilot study of cognitive-behavioral therapy in a sample of incarcerated women with substance use disorder and PTSD. *Behaviour Therapy* 40, 325–336.
- Zlotnick C, Shea TM, Rosen K, Simpson E, Mulrenin K, Begin A and Pearlstein T(1997) An affect-management group for women with posttraumatic stress disorder and histories of childhood sexual abuse. *Journal of Traumatic Stress* 10, 425–436.
- Zoellner LA, Foa EB and Brigidi BD (1999) Interpersonal friction and PTSD in female victims of sexual and nonsexual assault. *Journal of Traumatic Stress* **12**, 689–700.
- Zoellner LA, Telch M, Foa EB, Farach FJ, McLean CP, Gallop R, Bluett EJ, Cobb A and Gonzalez-Lima F (2017) Enhancing extinction learning in posttraumatic stress disorder with brief daily imaginal exposure and methylene blue: A randomized controlled trial. *Journal of Clinical Psychiatry* **78**, e782–e789.

#### X. Other References

Benish SG, Imel ZE and Wampold BE (2008) The relative efficacy of bona fide psychotherapies for treating post-traumatic stress disorder: A meta-analysis of direct comparisons. *Clinical Psychology Review* 28, 746–758.

Cloitre M, Garvert DW, Brewin CR, Bryant RA and Maercker A (2013) Evidence for proposed ICD-11 PTSD and complex PTSD: a latent profile analysis. *European Journal of Psychotraumatology* 4, 10.3402/ejpt.v4i0.20706. doi:10.3402/ejpt.v4i0.20706

Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J, Norris S, Falck-Ytter Y,
Glasziou P, DeBeer H, Jaeschke R, Rind D, Meerpohl J, Dahm P and
Schünemann HJ (2011) GRADE guidelines: 1. Introduction-GRADE evidence
profiles and summary of findings tables. *Journal of Clinical Epidemiology* 64, 383-94.

- Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, Savovic J, Schulz KF, Weeks L, Sterne JA; Cochrane Bias Methods Group and Cochrane
  Statistical Methods Group (2011) The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *British Medical Journal* 343, d5928.
- **Higgins JPT and Green S** (2011) Cochrane Handbook for Systematic Reviews of Interventions. The Cochrane Collaboration and John Wiley & Sons Ltd: England.
- Xia J, Adams C, Bhagat N, Bhagat V, Bhoopathi P, El-Sayeh H, Pinfold V and Takriti
   Y (2009) Losing participants before the trial ends erodes credibility of findings.
   *Psychiatric Bulletin* 33, 254–257.

#### Figure 1. PRISMA diagram



## Figure 2. Overview of studies contributing to each analysis

Study		Group A	Group B	Outcome
Ahmadi 2015		EMDR	TAU/WL	DR
Hogberg 2007		EMDR	Control	DR
Power 2002		EMDR	TAU/WL	AD
van den Berg 2015		EMDR	Control	AD
Kip 2014		EMDR	TAU/WL	NSC
Scheck 1998	A A A A A A A A A A A A A A A A A A A	EMDR	Control	NSC
Difede 2007		EMDR	TAU/WL	PTSD
Ehlers 2005		EMDR	Control	PTSD
Dunne 2012	SMACH SIL	EMDR	TAU/WL	PTSD + 1, 2 or 3
Talbot 2014	LANK LANK	EMDR	TAU/WL	PTSD + 2 or 3
Foa 1999	and the second second	EMDR	Control	PTSD + 1, 2 or 3
Monson 2006		EMDR	Control	PTSD + 2 or 3
Hollified 2007		CBT	TAU/WI	DR
Duffy 2007	- HARAMAR -	CBT	Control	DR
Galovski 2012	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	CBT	TALI/WI	AD
Basoglu 2007		CBT	Control	AD
Eblars 2003		CBT		NSC
Emers 2005		CPT	Control	NSC
Fblars 2014		CBT	тациял	PTSD
Krakow 2001		CBT	Control	PTSD
Lindever 2005		CDT	TALIANT	PTSD + 1 2 or 3
Lindauer 2005		CBI	TAU/WL TAU/WL	PTSD + 2 or 3
Marks 1998		CBI	TAU/WL	PTSD + 3
Forbes 2012		CBI	TAU/WL	PTSD + 1 2 or 3
Hinton 2009		CBI	Control	DP
Clottre 2002		Exposure alone	IAU/WL	DR
Dunn 2007		Exposure alone	Control	NSC
Hinton 2011		Exposure alone	TAU/WL	PTSD
Kubany 2003		Exposure alone	TAU/WL	PTSD
Jung 2013		Exposure alone	Control	
McDonagh 2005		Exposure alone	TAU/WL	PTSD + 2 or 3
Mueser 2008		Exposure alone	TAU/WL	PTSD + 1 2 or 2
Ford 2011		Exposure alone	Control	P13D + 1, 2013
Resick 2002		CBT	Exposure alone	DR
Kubany 2004		CBT	Exposure alone	NSC
Suris 2013		CBT	Exposure alone	PISD
Mueser 2015		CBT	Exposure alone	PTSD + 1, 2 or 3
Steel 2017		CBT	EMDR	DR
Dorrepaal 2012		CBT	EMDR	PISD
Keane 1989		CBT	EMDR	PTSD + 1, 2 or 3
Ghafoori 2017		EMDR	Exposure alone	DR
Pacella 2012		EMDR	Exposure alone	NSC
Nijdam 2012		EMDR	Exposure alone	PTSD + 1, 2 or 3
Krupnick 2008		EMDR	Exposure alone	PTSD + 2 or 3
Azad marzabadi 2014		IPT	TAU/WL	PISD + DR
Biedel 2011		Mindfulness	TAU/WL	DR
Biedel 2018		TMT	Exposure alone	PTSD, DR & AD
Butollo 2016		TMT	Exposure alone	PTSD + DR
Bryant 2013		DET	CBT	PTSD + NSC
Harned 2014		CBT + ER	CBT	PTSD + NSC
ter Heide 2011		DBT + exposure	DBT	PTSD + NSC
ter Heide 2016		EMDR	STBT	PTSD + DR

Abbreviations: AD=affect dysregulation; CBT=cognitive behavioural therapy; CPTSD=complex posttraumatic stress disorder; DBT=dialectical behaviour therapy; DET=dialogical exposure therapy; DR=disturbances in relationships; DSO=disturbances in self-organisation; EMDR=eye-movement and desensitisation and reprocessing therapy; ER=emotion regulation (training); IPT=interpersonal psychotherapy; NSC=negative self-concept; PTSD=posttraumatic stress disorder; PTSD + 1, 2 or 3 =PTSD + 1, 2 or 3 =PTSD + 3 =PTSD + all 3 CPTSD (DSO) outcomes; STBT=stabilisation treatment; TAU=treatment as usual; TMT=trauma management therapy; WL=waiting list.

Figure 3. Bubble plot of trauma onset (adult vs child) by Hedges' g, controlling for study quality, degree of CPTSD symptom severity, type of comparator, and type of treatment in multivariate meta-regression.



Regression of Hedges's g on Trauma onset

Trauma onset

Outcome	Comparator	k included studies	Treatment N	Control N	Hedges' g (95% CI), p- value	Heterogeneity, I <sup>2</sup> , p-value	Publication bias, p-value, adjusted g (95% CI), k imputed studies	Quality (GRADE)
DR	vs TAU/WL	16	485	395	-0.66 (-0.84, -0.48), <0.001	45%, 0.021	0.007, -0.39 (- 0.59, -0.20), 8	Moderate -1 publication bias
DR	vs control	3	128	79	-0.32 (-0.60, -0.03), 0.029	0%, 0.402	-	Moderate -1 imprecision
AD	vs TAU/WL	3	54	61	-1.42 (-2.20, -0.65), <0.001	71%, 0.033	-	Moderate -1 imprecision
AD	vs control	2	63	62	-0.82 (-2.91, 1.26), 0.440	94%, <0.001	-	Very low -1 inconsistency -2 imprecision
NSC	vs TAU/WL	9	320	281	-0.82 (-1.19, -0.44), <0.001	79%, <0.001	-	Moderate -1 inconsistency
NSC	vs control	4	207	163	-0.24 (-0.69, 0.21), 0.295	75%, 0.008	-	Low -1 inconsistency -1 imprecision

Table 1. Cognitive behavioural therapy with or without exposure vs TAU/WL or non-specific control

PTSD	vs TAU/WL	27	899	773	-0.90 (-1.11, -0.68), <0.001	76%, <0.001	0.002, -0.90 (- 1.11, -0.68), 0	Moderate -1 inconsistency
PTSD	vs control	9	408	323	-0.37 (-0.66, -0.09), 0.011	71%, 0.001		Moderate -1 inconsistency
PTSD + 1, 2 or 3	vs TAU/WL	27	841	705	-0.81 (-1.00, -0.62), <0.001	68%, <0.001	0.003, -0.81 (- 1.00, -0.62), 0	High
PTSD + 2 or 3	vs TAU/WL	3	92	90	-0.78 (-1.31, -0.24), 0.005	68%, 0.043		Moderate -1 imprecision
PTSD + 3	vs TAU/WL	2	58	58	-0.53 (-0.96, -0.09), 0.017	28%, 0.239		Low -2 imprecision
PTSD + 1, 2 or 3	vs control	9	398	314	-0.34 (-0.62, -0.06), 0.019	68%, 0.001		Low -1 imprecision -1 inconsistency
PTSD + 2  or  3	vs control	0	-	-	-	-		
PTSD + 3	vs control	0	-	-	-	-		

Abbreviations: AD=affect dysregulation; CPTSD=complex posttraumatic stress disorder; DR=disturbances in relationships; DSO=disturbances in self-organisation; NSC=negative self-concept; PTSD=posttraumatic stress disorder; PTSD + 1, 2 or 3=PTSD + 1, 2 or 3 CPTSD (DSO) outcomes; PTSD + 2 or 3=PTSD + 2 or 3 CPTSD (DSO) outcomes; PTSD + 3=PTSD + all 3 CPTSD (DSO) outcomes; TAU=treatment as usual; WL=waiting list.

Outcome	Comparator	k included studies	Treatment N	Control N	Hedges' g (95% CI), p- value	Heterogeneity, I <sup>2</sup> , p-value	Publication bias, p-value, adjusted g (95% CI), k imputed studies	Quality (GRADE)
DR	vs TAU/WL	4	158	110	-0.59 (-1.12, -0.07), 0.028	73%, 0.011	-	Moderate -1 imprecision
DR	vs control	1	47	24	-0.12 (-0.60, 0.37), 0.642	-	-	Very low -2 RoB -2 imprecision
AD	vs TAU/WL	0	-	-	-	-	-	-
AD	vs control	0	-	-	-	-	-	-
NSC	vs TAU/WL	3	131	102	-0.73 (-1.03, -0.43), <0.001	21%, 0.283	-	Moderate -1 imprecision
NSC	vs control	0	-	-	-	-	-	-
PTSD	vs TAU/WL	6	246	190	-1.05 (-1.52, -0.58), <0.001	79%, <0.001	-	Low -2 imprecision
PTSD	vs control	2	67	42	-0.08 (-0.47, 0.30), 0.675	0%, 0.803	-	Low -2 imprecision

# Table 2. Exposure only vs TAU/WL or non-specific control

Outcome	Comparator	k included studies	Treatment N	Control N	Hedges' g (95% CI), p- value	Heterogeneity, I <sup>2</sup> , p-value	Publication bias, p-value, adjusted g (95% CI), k imputed studies	Quality (GRADE)
PTSD + 1, 2 or 3	vs TAU/WL	6	242	158	-0.86 (-1.25, -0.47), <0.001	69%, 0.006	-	High
PTSD + 2 or 3	vs TAU/WL	1	47	39	-0.56 (-0.99, -0.14), 0.009	-	-	Low -2 imprecision
PTSD + 3	vs TAU/WL	0	-	-	-	-	-	-
$\frac{\text{PTSD} + 1, 2 \text{ or}}{3}$	vs control	2	67	42	-0.19 (-0.57, 0.20), 0.336	0%, 0.636	-	Low -2 imprecision
PTSD + 2  or  3	vs control	0	-	-	-	-	-	-
PTSD + 3	vs control	-	-	-	-	-	-	-

Abbreviations: AD=affect dysregulation; CPTSD=complex posttraumatic stress disorder; DSO=disturbances in self-organisation; DR=disturbances in relationships; NSC=negative self-concept; PTSD=posttraumatic stress disorder; PTSD + 1, 2 or 3=PTSD + 1, 2 or 3 CPTSD (DSO) outcomes; PTSD + 2 or 3=PTSD + 2 or 3 CPTSD (DSO) outcomes; PTSD + 3=PTSD + all 3 CPTSD (DSO) outcomes; TAU=treatment as usual; WL=waiting list.

Ta	ble 3.	EMDR	vs	TAU/WL	or non-	-specific	control
----	--------	------	----	--------	---------	-----------	---------

Outcome	Comparator	k included studies	Treatment N	Control N	Hedges' g (95% CI), p- value	Heterogeneity, I <sup>2</sup> , p-value	Publication bias, p-value, adjusted g (95% CI), k imputed studies	Quality (GRADE)
DR	vs TAU/WL	4	94	84	-0.76 (-1.35, - 0.16), 0.012	70%, 0.019	-	Moderate -1 imprecision
DR	vs control	2	37	34	-0.35 (-1.01, 0.31), 0.312	46%, 0.174	-	Very low -2 RoB -2 imprecision -1 inconsistency
AD	vs TAU/WL	1	11	12	-1.64 (-2.56, - 0.72), 0.000	-	-	Very low -2 RoB -2 imprecision
AD	vs control	1	11	10	0.25 (-0.57, 1.08), 0.548	-	-	Very low -2 RoB -2 imprecision
NSC	vs TAU/WL	1	44	39	-0.61 (-1.04, - 0.17), 0.006	-	-	Low -2 imprecision
NSC	vs control	2	56	53	-0.78 (-1.56, - 0.01), 0.049	75%, 0.047	-	Very low -1 inconsistency

Outcome	Comparator	k included studies	Treatment N	Control N	Hedges' g (95% CI), p- value	Heterogeneity, I <sup>2</sup> , p-value	Publication bias, p-value, adjusted g (95% CI), k imputed studies	Quality (GRADE)
								-2 imprecision
PTSD	vs TAU/WL	4	105	92	-1.26 (-2.01, - 0.51), 0.001	79%, 0.002	-	Low -1 inconsistency -1 imprecision
PTSD	vs control	3	70	65	-0.69 (-1.35, - 0.03), 0.041	70%, 0.035	-	Very low -1 RoB -1 inconsistency -1 imprecision
PTSD + 1, 2 or 3	vs TAU/WL	4	94	84	-1.15 (-1.92, - 0.37), 0.004	81%, 0.002	-	Low -1 inconsistency -1 imprecision
PTSD + 2  or  3	vs TAU/WL	2	55	51	-1.36 (-3.13, 0.42), 0.134	90%, 0.001	-	Very low -1 inconsistency -2 imprecision
PISD + 3	VS IAU/WL	U	-	-	-	-	-	-

Outcome	Comparator	k included studies	Treatment N	Control N	Hedges' g (95% CI), p- value	Heterogeneity, I <sup>2</sup> , p-value	Publication bias, p-value, adjusted g (95% CI), k imputed studies	Quality (GRADE)
PTSD + 1, 2 or 3	vs control	3	67	61	-0.52 (-0.97, - 0.08), 0.020	35%, 0.213	-	Low -1 RoB -1 imprecision
PTSD + 2  or  3	vs control	2	37	34	-0.44 (-1.31, 0.43), 0.321	68%, 0.079	-	Very low -2 RoB -1 inconsistency -2 imprecision
PTSD + 3	vs control	0	-	-	-	-	-	-

Abbreviations: AD=affect dysregulation; CPTSD=complex posttraumatic stress disorder; DR=disturbances in relationships; DSO=disturbances in self-organisation; EMDR=eye-movement and desensitisation and reprocessing therapy; NSC=negative self-concept; PTSD=posttraumatic stress disorder; PTSD + 1, 2 or 3=PTSD + 1, 2 or 3 CPTSD (DSO) outcomes; PTSD + 2 or 3=PTSD + 2 or 3 CPTSD (DSO) outcomes; PTSD + 3=PTSD + all 3 CPTSD (DSO) outcomes; TAU=treatment as usual; WL=waiting list.

## Table 4. Comparison of CBT, Exposure and EMDR

Outcome	Comparison (A vs B)	k included studies	Group A N	Group B N	Hedges' g (95% CI), p- value	Heterogeneity, I <sup>2</sup> , p-value	Publication bias, p-value, adjusted g (95% CI), k imputed studies	Quality (GRADE)
DR	CBT vs exposure alone	3	152	120	0.07 (-0.26, 0.39), 0.689	38%, 0.200	-	Moderate -1 imprecision
AD	CBT vs exposure alone	0	-	-	-	-	-	-
NSC	CBT vs exposure alone	1	62	61	-0.31 (-0.67, 0.04), 0.082	-	-	Very low -2 RoB -2 imprecision
PTSD	CBT vs exposure alone	4	216	184	-0.03 (-0.23, 0.17), 0.784	0%, 0.493	-	Moderate -1 imprecision
PTSD + 1, 2 or 3	CBT vs exposure alone	4	214	181	-0.04 (-0.27, 0.19), 0.719	20%, 0.291	-	Moderate -1 imprecision
PTSD + 2  or  3	CBT vs exposure alone	0	-	-	-	-	-	-
PTSD + 3	CBT vs exposure alone	0	-	-	-	-	-	-

Outcome	Comparison (A vs B)	k included studies	Group A N	Group B N	Hedges' g (95% CI), p- value	Heterogeneity, I <sup>2</sup> , p-value	Publication bias, p-value, adjusted g (95% CI), k imputed studies	Quality (GRADE)
DR	CBT vs EMDR	2	59	70	0.28 (-0.29, 0.34), 0.338	60%, 0.115	-	Very low -2 imprecision -1 inconsistency
AD	CBT vs EMDR	0	-	-	-	-	-	-
NSC	CBT vs EMDR	0	-	-	-	-	-	-
PTSD	CBT vs EMDR	2	62	75	0.37 (0.03, 0.71), 0.031	0%, 0.548	-	Low -2 imprecision
PTSD + 1, 2 or 3	CBT vs EMDR	2	59	70	0.31 (-0.07, 0.68), 0.111	16%, 0.275	-	Low -2 imprecision
PTSD + 2  or  3	CBT vs EMDR	0	-	-	-	-	-	-
PTSD + 3	CBT vs EMDR	0	-	-	-	-	-	-
DR	EMDR vs exposure alone	1	44	47	-0.10 (-0.51, 0.31), 0.640	-	-	Low -2 imprecision
AD	EMDR vs exposure alone	0	-	-	-	-	-	-

Outcome	Comparison (A vs B)	k included studies	Group A N	Group B N	Hedges' g (95% CI), p- value	Heterogeneity, I <sup>2</sup> , p-value	Publication bias, p-value, adjusted g (95% CI), k imputed studies	Quality (GRADE)
NSC	EMDR vs exposure alone	1	44	47	0.16 (-0.25, 0.57), 0.444	-	-	Low -2 imprecision
PTSD	EMDR vs exposure alone	1	55	48	0.10 (-0.28, 0.49), 0.604	-	-	Low -2 imprecision
PTSD + 1, 2 or 3	EMDR vs exposure alone	1	44	47	0.06 (-0.35, 0.46), 0.789	-	-	Low -2 imprecision
PTSD + 2 or 3	EMDR vs exposure alone	1	44	47	0.06 (-0.35, 0.46), 0.789	-	-	Low -2 imprecision
PTSD + 3	EMDR vs exposure alone	0	-	-	-	-	-	-

Abbreviations: AD=affect dysregulation; CPTSD=complex posttraumatic stress disorder; DR=disturbances in relationships; DSO=disturbances in self-organisation; EMDR=eye-movement and desensitisation and reprocessing therapy; NSC=negative self-concept; PTSD=posttraumatic stress disorder; PTSD + 1, 2 or 3=PTSD + 1, 2 or 3 CPTSD (DSO) outcomes; PTSD + 2 or 3=PTSD + 2 or 3 CPTSD (DSO) outcomes; PTSD + 3=PTSD + all 3 CPTSD (DSO) outcomes.