

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/120170/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Imms, Phoebe, Clemente, Adam, Cook, Mark, D'Souza, Wendyl, Wilson, Peter H., Jones, Derek K. and Caeyenberghs, Karen 2019. The structural connectome in traumatic brain injury: A meta-analysis of graph metrics. *Neuroscience and Biobehavioral Reviews* 99 , pp. 128-137. 10.1016/j.neubiorev.2019.01.002

Publishers page: <http://dx.doi.org/10.1016/j.neubiorev.2019.01.002>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



## Abstract

IMMS, P., A. Clemente, M. Cook, W. D'Souza, P. H. Wilson, D. K., Jones & K.

Caeyenberghs. Insights into traumatic brain injury from graph theory: A meta-analysis.

NEUROSCI BIOBEHAV REV 95(1) XXX-XXX, 2018. - Although recent structural

connectivity studies of traumatic brain injury (TBI) have used graph theory to evaluate

alterations in global integration and functional segregation, pooled analysis is needed to

examine the robust patterns of change in graph metrics across studies. Following a systematic

search, 15 studies met the inclusion criteria for review. Of these, ten studies were included in

a random-effects meta-analysis of global graph metrics, and subgroup analyses examined the

confounding effects of severity and time since injury. The meta-analysis revealed

significantly higher values of normalised clustering coefficient ( $g=1.445$ ,  $CI=[0.512, 2.378]$ ,

$p=0.002$ ) and longer characteristic path length ( $g=0.514$ ,  $CI=[0.190, 0.838]$ ,  $p=0.002$ ) in TBI

patients compared with healthy controls. Our findings suggest that the TBI structural network

has shifted away from the balanced small-world network towards a regular lattice. Therefore,

these graph metrics may be useful markers of neurocognitive dysfunction in TBI. We

conclude that the pattern of change revealed by our analysis should be used to guide

hypothesis-driven research into the role of graph metrics as diagnostic and prognostic

biomarkers.

**Key words:** Traumatic Brain Injury; graph theory; graph metrics; structural connectomics;

network analysis; diffusion MRI; biomarkers; meta-analysis; systematic search; narrative

review.