

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/122962/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

de Graaf, Barend H and Dewitte, Walter 2019. Fertilisation and cell cycle in angiosperms. *Annual Plant Reviews Online* 2019 (2)
10.1002/9781119312994.apr0641

Publishers page: <https://doi.org/10.1002/9781119312994.apr0641>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Fertilisation and Cell Cycle in Angiosperms

Barend H. J. de Graaf and Walter Dewitte

*SCHOOL OF BIOSCIENCES, CARDIFF UNIVERSITY,
CARDIFF, UK*

1 Introduction

Angiosperms, the large groups of seed plants better known as flowering plants, are the most successful land plants on our planet today, representing an estimate of 95% of all terrestrial vegetation (Simonin and Roddy, 2018). Their seeds germinate only during favourable growth conditions culminating in the development of a new generation of plants, which can show variable periods of vegetative growth depending on the species, habitat, and growth conditions. The transition of many of these angiosperm plants into the flowering phase is triggered by seasonal and environmental factors and involves a switch from vegetative growth into developing reproductive structures. Upon fertilisation, the embryo encapsulated in a seed ensures the survival of the species. The seeds of some flowering plant species can be stored for very long periods of time and retain their viability even deep in the soil. Disruption of the substrate by ploughing or digging changes the micro environment and disrupts seed dormancy (Hilhorst et al., 2010; Shen-Miller et al., 1995). A large variation in shape, nectar availability, volatile production, and colours exists among flowers. Hence most flowers are very attractive to the human eye. In Angiosperms successful reproduction depends on pollen grains, often carried by wind and insect pollinators, that land on the stigmatic surface and fertilise the ovules. During the relatively rapid development of flowers, male and female reproductive organs, anthers and pistils, respectively, are formed at the centre of the flower. Within anthers and pistils, sporophytic cells are recruited to generate the new germ line from which male and female gametophytes develop. The egg cell is specified inside the developing embryo sac, which is the female gametophyte of a plant. Two sperm cells (SCs) develop inside a single vegetative cell (VC) surrounded by several durable layers, forming and protecting the male gametophyte, referred to as the pollen grain. Once flowers are fully mature and ready for reproduction, they open and the pollen grains are released from the anthers.

In the 320 000 different flowering plant species, many different mechanisms have evolved that ensure outcrossing to maximise genetic variability in the offspring (Lughadha et al., 2016; Fujii et al., 2016; Hiscock, 2011).

After pollen lands on a stigma, a series of complex recognition events results in rejection of all foreign pollen by as yet unknown mechanisms. In addition, approximately half of all angiosperm species display a genetically determined mechanism, which prevents self-fertilisation called self-incompatibility (SI). Pollen rejection can vary between different SI mechanisms, either by preventing pollen germination immediately after pollen landing by blocking water uptake from the stigma, or by triggering calcium signalling mediated programmed cell death in incompatible pollen, or by the activity of pistil produced S-RNases after translocation into the growing pollen tubes (Allen et al., 2010; Fujii et al., 2016).

All ‘own’ and compatible pollen grains rapidly hydrate, followed by pollen grain germination and the outgrowth of a pollen tube, driven by the tip growth of the VC. The pollen tube accurately delivers two SCs into one embryo sac deep enclosed in the ovule inside the pistil. In flowering plants, upon pollen tube arrival and SC release, a unique double fertilisation takes place in the female embryo sac. One SC fuses with the egg cell, generating the diploid zygote that will form a new sporophyte (embryo), whereas the second sperm merges with the diploid large central cell (CC) and results in the formation of the triploid endosperm precursor, triggering seed development. The endosperm functions as a nutrient source for the developing plant embryo, either during seed development and maturation before release from the mother plant, or during the early stages of plant development after seed dispersal and germination (de Graaf et al., 2001; Rousseau et al., 2008; Mizuta and Higashiyama, 2018).

Here, we survey the critical processes for plant reproduction and explore this in the context of cell cycle regulation. The coordination of cell cycle is required to produce the gametes, synchronise the cell cycle phase for fertilisation as well as the creation of an embryo from a zygote.

2 Fertilisation

2.1 Gametophyte Development

The developmental pathways of both male and female gametophytes producing both sex cell types for plant reproduction are highly conserved throughout angiosperms. Nevertheless, there are considerable differences in the timing of developmental steps, this can lead to a different number of cells in the pollen grain. In some species, the last mitotic division has not been executed and we see bicellular pollen, instead of tricellular. The total numbers of cells in the female gametophyte depends on the species as well (Yadegari and Drews, 2004).

The final cell number in most male gametophytes (pollen) appears to be rigidly conserved in all higher plants, formed by a well-defined sequence of divisions. Meiotic rounds of division (I and II) of a single diploid microsporocyte (microspore mother cell) gives rise to a tetrad of four haploid microspores embedded in callose (β 1,3 glucan). The tapetum cell layer in the anther, surrounding the developing microspores, is responsible for the degradation of callose, releasing free microspores. These microspores undergo a first round of mitotic and asymmetric division that results in the formation of a larger VC and a smaller generative cell (GC). At this point of development, the GC becomes invaginated by the VC giving rise to a cell within a cell generating a bicellular pollen grain. The anther tapetum and developing pollen are both responsible for the production and deposition of two extracellular pollen layers, the intine composed mainly of cellulose-pectic compounds and an exine layer – composed of sporopollenin, which besides its species-specific structural properties to optimise pollen transfer, is composed of a biochemically decay-resistant substance protecting the pollen once released from the anther upon flower opening (Borg and Twell, 2010; Twell, 2011; Gómez et al., 2015).

2.2 Attraction to the Ovule

Among angiosperms, there are considerable structural differences in the size of reproductive organs, which also confers large differences in distance between stigma surface of the pistil, the landing platform for pollen, and the ovules inside the pistil. As such, the time for pollen tubes to reach an ovule depends on the species. For example, it takes approximately five hours for the fastest growing vegetative pollen tube cell of *Arabidopsis* to grow approximately 3 mm inside the pistil to deliver the SCs and it takes an additional four to five hours to fertilise every ovule (Jean-Emmanuel et al., 2002). In maize, pollen tubes need to grow up to 25 cm in length for approximately 24 h, growing at an astonishing rate of almost 1.0 cm h^{-1} (Dresselhaus et al., 2016; Zhou et al., 2017). In rice, it only takes 15–60 min after pollen tube germination to reach an ovule and two to five hours to

complete fertilisation (Lan et al., 2004; Dong and Yang, 1989). Moreover, pollen tube growth is affected by the structural, physical, and nutritious properties of the different pistil tissues. A switch in tube growth speed is clearly observed in bicellular pollen species, such as *Nicotiana tabacum* (see below), which was suggested to be correlated with the staging of the second mitotic division of the GC into two SCs while growing through the style transmitting tissue (TT) of the pistil. In Arabidopsis, however, a species with tricellular pollen upon anthesis, the tubes grow slower through the TT of the short style and the long septum in the centre of the ovary as compared to the initial stigma growth (Lennon et al., 1998; Cheung et al., 2010). Pollen tubes show directional growth with high precision due to structural and biochemical clues that are provided by the different pistil tissues, i.e. stigma, style, ovary, and ultimately, the embryo sacs inside the ovules.

During pollen–pistil interaction, specific members of a large family of membrane localised receptor-like kinases (RLKs) are involved in the guidance of the expanding pollen tube tip through the different pistil tissues. Both pollen tube (ANX1/2, BUPS1/2 and LIPS1/2) and ovule-specific RLKs (STRUBBELIG) are responsible for sensing directional clues for micropyle entrance into an ovule and for the communication between pollen tube tip and the different types of embryo sac cells such as the synergid cells with their filiform apparatus (FERONIA), the eggs and CC, but also the integument cell layers surrounding, protecting and communicating with the embryo sac during ovule development (Muschiatti and Wengier 2018; Wang et al., 2016).

‘Defensin-like’ proteins, i.e. LUREs and EA1 produced by synergids and egg cell respectively (see below), are responsible for a species-specific close-range attraction and guidance of pollen tubes into the micropyle of the ovule (Márton et al., 2012; Okuda et al., 2013; Takeuchi and Higashiyama, 2012). Remarkably, the interaction between growing pollen tubes and female pistil tissues during the progamic phase of plant reproduction is essential to prime the pollen tubes to respond to the LURE protein attractants and to release the SCs inside the embryo sac (Okuda et al., 2013; Leydon and Johnson, 2013). Interestingly, an additional style TT produced component that triggers the competence of pollen tubes to respond to these LURE proteins is not a protein or peptide but is shown to be an arabinogalactan polysaccharide (AMOR) (Mizukami et al., 2016; Okuda et al., 2013). One of the two synergid cells at the entrance of the embryo sac collapses upon pollen tube arrival at this site, at least in Arabidopsis (Jean-Emmanuel et al., 2002) concomitantly triggering the arrest and rupture of the pollen tube tip, releasing its content including both SCs (Leydon and Johnson, 2013; Leydon et al., 2013).

Curiously, an important structural characteristic of all different types of cells within an embryo sac (Yadegari and Drews, 2004) but also of the pollen tube arriving at one of the synergid cells and as well as of the two released SCs is the development of a cell wall. This may create a potential structural barrier between cells and thus prevent direct membrane–membrane interaction and animal-like cell communication and fusion event(s) (Wang et al., 2010; Mori et al., 2014). Signalling between these cells and the communication by making use of membrane-localised receptors, such as the FERONIA receptor, GPI-anchored LORELEI protein or the SC-specific MDIS1-MIK receptor kinases, GCS1/HAP2, and GEX2 (Mori et al., 2005, 2014; Escobar-Restrepo et al., 2007; Igawa et al., 2013; Liu et al., 2016; Dresselhaus and Snell, 2014; Wang et al., 2016; Berger et al., 2008; von Besser et al., 2006), can only occur based on the release (exocytosis) of highly diffusible compounds, e.g. Ca^{2+} and signalling proteins such as the group of small secreted cysteine-rich globular proteins (CRPs), e.g. the LURE proteins (Sprunck et al., 2012; Dresselhaus et al., 2016; Mizuta and Higashiyama, 2018; Márton et al., 2012). However, it has been demonstrated that in *N. tabacum*, high molecular weight and glycosylated arabinogalactan-like proline-rich proteins from the pistil TT can traverse pollen tubes walls *in vivo* ending up against the pollen tube membrane (de Graaf et al., 2000). Cell wall modifying enzymes such as pectinases, glucanases, and expansins may play a role in changing the structural properties of these cell walls facilitating protein movement across plant cell walls, supporting cell–cell communication (de Graaf et al., 2000). The CRPs and other defensin-related small peptides, including the LURE proteins, and especially the short RALF (*r*apid *a*lkalization *f*actor) peptides (see above), may be able to traverse cell walls more easily, quickly, and specifically (Takeuchi and Higashiyama, 2012; Márton et al., 2012).

In both bicellular and tricellular pollen species, the GC and the two SCs are spatially, structurally, and temporally connected with the vegetative tube cell nucleus (VN), defined as a male germ unit (MGU). The GC and SCs are surrounded by a continuous vegetative membrane which has originated from the microspore membrane during pollen mitosis I – during pollen development in the anther locule – when the GC becomes ‘endocytosed’ from the vegetative pollen cell after asymmetric division (see above). An intercellular matrix or ‘periplasm-like’ region, rich in pectins and arabinogalactan proteins (AGPs), separates both SCs from the vegetative tube cell. One end of the SCs forms a characteristic hook structure and creates an intimate contact between the VN of the tube cell and either the GC or SCs and is suggested to play a role in communication (Jiang et al., 2015; Grant-Downton et al., 2013; Slotkin et al., 2009).

However, plasmodesmata-like cytoplasmic connections between both cell types appear to be absent (van Aelst and van Went, 1992; Cresti et al., 1987; Southworth and Kwiatkowski, 1996). After pollination and pollen tube germination, the MGU becomes translocated into the growing tube where it migrates with and at a specific distance from the growing pollen tube apex. Importantly, once pollen tubes grow inside the female pistil, the interaction between the male gametophyte and maternal tissues triggers different gene expression in the pollen tube, compared to *in vitro* growing pollen tube cultures. This indicated that pistil pollen tube communication is essential for directed pollen tube growth, and indeed pollen tubes respond to the LURE protein attractants produced by the female gametophyte (Lin et al., 2014; Qin et al., 2009). In addition, maturation of the SCs in the tricellular pollen of Arabidopsis seems to be completed only by the time pollen tubes reach the embryo sacs (Friedman, 2006). An important question that needs to be resolved is whether SCs, when transported through a growing pollen tube – are indeed actively involved in pollen tube guidance to an ovule and fertilisation (Mori et al., 2005). When the SC expressed HAP2/GCS1 protein is absent, these pollen tubes are disrupted in their ability to find the micropyle of the ovules, and those that do arrive fail to initiate fertilisation (von Besser et al., 2006). Recently, however, Glöckle et al. (2018) demonstrated that mutant pollen – in which the second mitotic division of the GC into two SCs is blocked – are still capable of producing a pollen tube, and find their way to an ovule *in vivo*. These results suggest that SCs are not important for the function of the pollen VC, which is the formation, polarised and directional growth of the pollen tube (Glöckle et al., 2018; Zhao et al., 2012). How communication is established between all the different cell types needs to be resolved.

2.3 Double Fertilisation

Recently, it has been demonstrated that besides reactive oxygen species (ROS), RALF peptides specifically expressed in the pollen tubes and in the ovule are involved in the actual disruption of the pollen tube tip membrane, at least in tissue culture (Ge et al., 2017; Mecchia et al., 2017; Duan et al., 2014). Remarkably, reducing the level of these RALF peptides does not result in a ‘nonbursting’ pollen tube phenotype in plants which suggests that other pollen and or ovule components are involved in the sperm release from pollen tubes. Following the release of the SC pair in the embryo sac, one of the synergids degenerates. Initially, both SCs move in close proximity showing a Brownian-like movement inside the embryo sac, up to the point that they separate, and move in different directions: one travels to the egg cell, whereas the second one find its way to the CC to

fuse and form the new zygote (sporophyte) and endosperm, respectively (Zhang et al., 1999; Huang and Russell, 1994; Kawashima et al., 2014), such events have now been captured by live imaging (Hamamura et al., 2011). The SCs have a protoplast-like morphology (Jean-Emmanuel et al., 2002), decorated with myosin, and are proposed to be guided precisely to the egg and CC by two actin coronas for double fertilisation to take place (Huang and Russell, 1994; Zhang et al., 1999; Kawashima et al., 2014; Ohnishi et al., 2014). At this stage, the vegetative membrane surrounding the two SCs and the pectic/AGP matrix must be removed to be able to make the contact between sperm membrane and the actin coronas and to achieve fusion with the egg and CC ultimately, with a lag time of eight minutes between sperm release and actual fusion (Hamamura et al., 2011; Denninger et al., 2014). Interestingly, in rice, the chalazal end of the egg cell does not present a proper cell wall which is suggested to be the site of SC fusion whereas formation of a cell wall early after sperm-egg fusion in *Arabidopsis* and maize is suggested to be important to prevent polyspermy (Kranz et al., 1995; Okamoto, 2010; Tekleyohans et al., 2017).

When exactly SCs are fully mature and capable to fuse with the egg and CC inside the embryo sac is currently not known. *In vitro* egg-sperm fusion experiments, using tissue cultured tobacco pollen tubes for the isolation of the SCs and egg cell protoplasts, suggests that the sperm is 'more or less' fully mature and capable of fusing with the egg cells immediately after division (Jean-Emmanuel et al., 2002; Khalequzzaman and Haq, 2005; Kranz et al., 2008). However, in tissue culture fertilisation assays, it has become evident that pollen tube attraction by ovules, the release of SCs and the fusion between sperm and egg cell is much more efficient when ovules, sperm, and egg cells are isolated from pre-pollinated pistils. Recently, it has been suggested that the small cysteine-rich EC1 protein, secreted by the egg cell, is important for rapid SC 'activation' to fuse with the egg cell more efficiently *in vivo* (Dresselhaus et al., 2016). Although it has been clearly demonstrated that pollen tubes need the interaction and communication with the different pistil tissues and the embryo sac containing the egg cell, respectively, it is still debated whether within a single male gametophyte (pollen), the VC or VN communicates directly with one or both SCs (Zhang et al., 2017a; McCue et al., 2011; Slotkin et al., 2009; Grant-Downton et al., 2013; Jiang et al., 2015).

The diploid CC inside the embryo sac fuses with one of the two SCs and this fusion product, the endosperm initial, develops a triploid endosperm, which functions as a temporary nutrient source for the actual developing zygote into the new embryo and seedling ultimately. The nutrient storage capacity and composition of endosperm tissue varies greatly between different species (Esau, 1962). Remarkably, in

gymnosperms, producing naked seeds, although two SCs are being delivered by a single pollen tube, only a single fertilisation event takes place due to the absence of a CC. The second sperm degenerates during further seed development.

2.4 Preventing Polyubuy

In Arabidopsis, upon pollen tube arrival and fertilisation, ovules also seem to become reprogrammed and either stop producing the short and – most probably – long distance guidance clues, or perhaps start producing repellent-like products preventing additional pollen tubes to grow towards and into an ‘unavailable’ ovule (Maruyama et al., 2015; Völz et al., 2013). For example, the production of LURE, a pollen tube attractant is turned off upon fertilisation (Palanivelu and Preuss, 2006; Maruyama et al., 2013). This process prevents the attraction of excess pollen tubes and the degeneration and fusion of the remaining synergid with the CC is key in this process. In Arabidopsis, this remaining synergid, the second synergid nuclei, degenerates after fertilisation, and this cell fuses with the CC in response to ethylene produced by the newly formed zygote (Maruyama et al., 2015; Völz et al., 2013). FIS-PRC2 is a chromatin-modifying complex involved in gene silencing via trimethylation of lysine 27 on histone H3 (H3K27me) (Kohler et al., 2012), and the activity of this complex is required for mitosis associated degradation of the synergid's nucleus (Maruyama et al., 2015). The fusion of the persistent synergid with the CC results in the dilution of the synergid produced attractants. JAGGER, an AGP is required for the degeneration of the remaining synergid (Pereira et al., 2016a, b). Interestingly, in *Oryza sativa* (rice, a monocotyledon), the degeneration of one of the synergids starts just before anthesis, thus even before pollination and this seems to be developmentally regulated rather than by an approaching pollen tube (Dong and Yang, 1989). However, in Arabidopsis, mutants that do allow more than one pollen tube to enter an ovule, the second set of SCs released were not able to fertilise the already fertilised egg or CC (Scott et al., 2008).

3 Cell Division and Fertilisation

The formation of a gametophyte involves three basic steps, the recruitment of diploid germ line cells in floral tissues, meiosis to generate haploid cells and one or more rounds of mitosis to generate the cells that will populate the gametophyte, including the gametes. During the mitotic

cell cycle, the duplication of genetic material during the S phase and segregation of the two copies in two different cells during M phase is temporally separated by two gap phases (G), G₁ and G₂ during which nutrient status and developmental signals can be integrated and DNA integrity control can be performed. The meiotic cell cycle is a modified cell cycle with a single S-phase but with two rounds of chromosome segregation generating haploid daughter cells.

In all eukaryotes, progression through both the meiotic and mitotic cell cycle is driven by the action of cyclin (CYC)/cyclin-dependent kinases (CDKs) complexes (De Veylder et al., 2007). These kinases phosphorylate key actors in cell cycle progression and directly or indirectly provide transcriptional and post-translational control over the core cell cycle machinery. The CDK/CYC kinases act in feedforward networks, wherein controlled destruction of cell cycle factors results in irreversible switches that provide direction to cell cycle progression (Zhao et al., 2012). To this end, cell cycle regulators are degraded by the proteasomes upon tagging by ubiquitin, implicating the action of ubiquitin ligases in cell cycle regulation. SCF (Skp, Cullin, F-box containing) complexes ubiquitinate factors involved in both G₁/S and G₂/M transitions, and the APC/C (anaphase-promoting complex/cyclosome) controls the metaphase–anaphase transition during mitosis (Genschik et al., 2014).

In higher plants, the G₁/S transition of the mitotic cell cycle is driven by the action of CDKA/CYC kinase complexes, whereas progression through G₂/M involves the action of both CYC/CDKA and CYC/CDKB kinases (Polyn et al., 2015). The activity of these CDK kinases is post-translationally controlled by phosphorylation. Activating phosphorylations that stabilise the T loop of the CDK are mediated by CDK-activating kinases (CAKs), while WEE1 kinases reduce activity by phosphorylating residues in the ATP-binding pocket. Interaction with inhibitors, of which the Kip-related proteins (KRPs)/interactor-inhibitor of cyclin-dependant kinase (ICK) and SIAMESE related (SIMS) are the best documented, and the control of CYCLIN levels by proteolysis are important negative feedback mechanisms. A key target of G₁/S CYCD/CDK and CYCA/CDK kinases is the retinoblastoma-related protein. Hyperphosphorylation of retinoblastoma related protein (RBR) by these kinases reduces its potential to interact with transcription factors and factors involved in chromatin structure and cell cycle progression. The action of phosphatases at the end of M-phase strips RBR of phosphate groups resulting in its reactivation. Proteolysis of the KRP inhibitors stimulated by the action of CYC/CDK via inactivation of RBR provides a positive feedback resulting in irreversible switch-like behaviour of the G₁/S cell cycle transition (Zhao et al., 2012). However, G₁/S kinases also regulate levels of RBR (Dewitte et al., 2003),

and E2Fa-RBR complexes suppresses E2F targets genes involved in cell differentiation (Magyar et al., 2012). This indicates that both RBR bound and free E2F fractions can have a role in suppressing differentiation and activating cell division. Further research into the role and stoichiometry of these fractions is required to elucidate this mechanism.

3.1 Gamete Formation and Cell Division

In preparation for sexual reproduction, male and female haploid gametes are produced in hermaphrodite flowers containing both male and female reproductive organs, anthers and pistils, or in male or female flowers that are formed on the same plant or on separate individuals, termed monoecious and dioecious plants respectively (Barrett and Hough, 2013). During flower development after the recruitment of a germ cell in the floral organs of the sporophyte, a series of meiotic and mitotic divisions, in combination with cell differentiation and programmed cell death, results in the formation of mature dry pollen grains and embryo sacs starting from diploid microsporocytes and megasporocytes, respectively.

Cyclins have been shown to be involved in meiosis, the correct execution of meiosis in the male microspore mother cell requires the action of six CYCAs, one CYCB type cyclin and the SOLODANCERS (SDS), related cyclins (Bulankova et al., 2013), which are activated in different stages of the process. CYCA1;2 is required for timely entry and prevents premature exit from meiosis after the first division. Its mutant alleles are known as *TARDY ASYNCHRONOUS MEIOSIS* (d'Erfurth et al., 2010). CDKA;1-CYCA1;2 phosphorylates and destabilises THREE DIVISION MUTANT1/MALE STERILE5 (TDM1) until meiosis I is successfully performed. TDM1 is required for meiotic exit by stimulating the APC/C activity at the end of meiosis II (Cifuentes et al., 2016). OSD1 seems to be also involved in controlling levels of APC activity, and *osd1 tam1* mutants even fail to enter into meiosis I (d'Erfurth et al., 2010). Several cyclins play a role in male meiosis. CYCA2s play an important role for the segregation of meiotic chromosomes. CYCB3;1 is localised to the meiotic spindles, and together with SDS represses the premature onset of cell wall formation (Bulankova et al., 2013). SDS is required for chromosome pairing. Meiotic exit requires the activation of TDM1 and the nonsense-mediated RNA decay factor encoding SMG7 genes (Bulankova et al., 2010). Male meiosis is sensitive to elevated temperatures, and the CDKG1-CYCLINL complex is involved in RNA splicing and is critical for chromosome pairing and recombination and the robustness of the meiotic process at ambient temperatures (Huang et al., 2013; Zheng et al., 2014b).

Secondly, after meiosis, several rounds of mitosis provide the appropriate cell number in the gametophyte. The male gametophyte contains three cells at the time of fertilisation. The pollen grain contains the SCs enveloped in the VC. This acquisition of new daughter cells fates involves RBR action as *rbr* mutant VCs retain aspects of the microspore whilst reiterating these unequal divisions (Chen et al., 2009). A second mitotic division of the GC generates the two SCs. Whilst this process is strictly conserved among all flowering plants, the timing of these divisions with respect to pollination and fertilisation seems to be variable in different species (Friedman, 1999). Degradation of KRP inhibitors by the SKP1-Cullin1-F-box protein (SCF) mediated by the FBL17 F-box protein is required to trigger the symmetrical division of the GC in order to produce two SCs (Kim et al., 2008), and CDKA;1 activity is required for their timely division (Nowack et al., 2006; Iwakawa et al., 2006; Aw et al., 2010). Cell cycle arrest of the VC requires the action of RBR, as well as suppression of CDKA;1 activity by the KRP inhibitors (KRP6 and KRP7 in Arabidopsis). SAMBA-mediated APC/C activity destabilisation of CYCA2;3 is required for progression through mitosis I (Eloy et al., 2012). The activation of CDKs by the combined action of the CDKD CAK kinases has proved to be essential for pollen mitosis I and II (Takatsuka et al., 2015).

The coordination of the final mitotic cell division and fate adoption during SC formation is intriguing. In gymnosperm male gametophytes, we also see a mitotic cycle that produces two SCs, even if this is unnecessary as only a single sperm is required for fertilisation, suggesting that cell cycle progression is not only necessary to generate the correct cell number but also a requirement for cell fate acquisition. However, cell division and sperm fate acquisition can be uncoupled in mutants impaired in the chromatin assembly factor 1 pathways in which single fertile SCs are formed (Chen et al., 2008), suggesting that defects in histone deposition might overcome the requirement of normal cell cycle progression to adopt a new cell fate. This in combination with the observation that in mutants of ARID1, which physically interacts with histone deacetylase, histone acetylation expands which is normally a feature of the SC, also expands into the VCs chromatin (Zheng et al., 2014a), suggest that regulation of histone deposition and activity is key in the process of SC specification. The coordination of fate acquisition and cell division during sperm formation requires the action of DUO1, a germline-specific Myb protein and a positive regulator of CYCB1;1 amongst other targets. DUO1 is itself a target of the ARID1 transcription factor. Both *duo1* and *arid1* mutant alleles display a high proportion of blocked divisions in the GCs (Durberry et al., 2005; Brownfield et al., 2009). The expression of a subset of DUO1

targets requires the action of DUO3, a homeobox-related factor (Brownfield et al., 2009), which acts independently of *CYCB1;1*, but has a wider role in sporophytic cell division and embryo patterning.

In order to ensure an ovule produces a single seed, it is essential that only a single cell inside the female ovule primordium enters the meiotic programme. Several key factors in this process have been discovered. Entry into meiosis requires the action of SWITCH/AMEIOTIC 1 (Mercier et al., 2001; Pawlowski et al., 2009). Furthermore, RBR plays a key dual role in this and suppresses WUSCHEL, a factor controlling stem cell identity involved in meiocytes specification, and limits the number of mitotic divisions to trigger meiosis in a single cell (Zhao et al., 2017). During evolution, the development of the female gametophyte has evolved from a four nucleate, four-celled gametophyte created by two mitotic divisions of the megaspore into seven-celled, eight nuclei-containing structures by nuclear migration followed by the duplication of modules (Friedman and Ryerson, 2009). Additional variations in the pattern of division led to a wide flexibility of the cell (and nuclei) number in the female gametophyte (Schmid et al., 2015). The plant retinoblastoma-related protein is a key factor to limit the divisions in the female gametophyte (Ebel et al., 2004), and local elevated expression of a D-type cyclin can prevent the arrest of the CC (Sornay et al., 2015). The current working hypothesis for cell fate acquisition in the female gametophyte is that this is mediated through an auxin gradient in combination with accurate nuclear migration and positioning (Sprunck and Gross-Hardt, 2011).

Maintenance of DNA integrity is essential in the context of gamete production and integral to cell cycle progression. Hence, it is worth noting that cell cycle machinery and DNA repair are closely linked. E2F activity is suppressed by the conserved DNA damage protein SNI, and they act in a negative feedback loop, linking the G₁/S checkpoint with DNA repair (Magyar et al., 2012). E2Fa-RBR complexes are recruited to sites of DNA damage and interact with AtBRCA1, a conserved DNA repair-associated protein. Furthermore, RBR is required to recruit RAD51 to sites of DNA damage, suggesting a central role for RBR in complexes involved in DNA repair (Horvath et al., 2017; Biedermann et al., 2017). Specific CDK activity is essential for DNA repair to overcome mitotic arrest, *CYCB1;1* is rapidly induced upon DNA damage, the attraction of RBR and RAD51 to damaged loci is dependent on *cycb1;1*/CDKB kinase action and RAD51 is a substrate of *CYCB1*/CDKB1 kinases (Weimer et al., 2016; Schnittger and de Veylder, 2018). The WEE1 kinase arrests the cell cycle by phosphorylation of CDK interfering with ATP binding, and its action is essential to block the cell cycle in case of replication defects (De Schutter et al., 2007).

3.2 Coordination of Cell Cycle for Fertilisation

Fertilisation is the result of plasmogamy followed by karyogamy, the fusion of cells followed by the fusion of nuclei. In angiosperms, double fertilisation requires two mature SCs to physically interact and fuse with the egg cell and CC to form the zygote and endosperm precursors (see above).

Angiosperms can be divided into two groups based on pollen morphology: one group, comprising approximately 70% of the total number of species, releases pollen at anthesis at the bicellular stage after pollen dehydration and anther locule degeneration. In the second group of angiosperms ($\pm 30\%$), pollen will continue their development following a second mitotic step of the GC into two SC before pollen dehydration and anthesis, which is called tricellular pollen (McCue et al., 2011).

Friedman and colleagues established that there are three options for karyogamy in seed plants (Carmichael and Friedman, 1995). G_1 karyogamy, whereby the zygote undergoes DNA replication, or S-phase where karyogamy is initiated in G_1 but nuclei pass through S-phase before completing fusion, or G_2 karyogamy where nuclei fuse after S-phase and once completed enter M-phase. The sperm nuclei are guided towards the egg cell by interaction with the actin cytoskeleton, and once in close proximity, the egg cell chromatin moves into the sperm nucleus (Ohnishi et al., 2014). The synchrony of gametes required for karyogamy is a challenge from the point of view of cell cycle regulation. Male SCs have to reach the correct cell cycle phase, and the egg cell has to remain quiescent until karyogamy. It implies that core cell cycle regulators are involved that blocks gamete cells either in G_1 or in G_2 , especially in the waiting or quiescent egg cell, and that in both the male and female gametophyte key regulatory pathways ensure this coordination. Progress in gamete isolation (Englhart et al., 2017; Schoft et al., 2015; Santos et al., 2017) and developments in transcriptomics and proteomics (Zhang et al., 2017b) are anticipated to put us in a position to identify the key core cell cycle regulators in this process. Although our knowledge is still fragmented due to our limits of detection, the emerging data on sperm (Table 1) and egg cells are in line with our knowledge about the cell cycle regulators involved in the key cell cycle transitions. Rice gametes fuse in G_1 , and transcripts for key factors such as KRPs, CDKB1;1, WEE1, and KRP2 were detected by RT PCR (Sukawa and Okamoto, 2018). An array-based analysis revealed significant levels of transcripts encoding for CDKB1;1, CYCB1-2, and KRP1 whereas proteomics detected only CDKA;1. This suggests that rice

SCs, while still in G₁ express a number of cell cycle genes, which have been implicated in DNA repair and G₂/M transition. In the egg cell, on the other hand, mainly transcripts encoding for KRP1 and CDKA;1 were detected (Sukawa and Okamoto, 2018). One can speculate that DNA repair mechanism are activated in SCs given that pollen are more exposed to environmental factors compared to the female gametophyte deep inside the ovary, alternatively, the paternal genome could be the main source of G₂/M factor encoding transcripts.

Table 1 Transcriptomics and Proteomics in Male Gametes

	G ₁	S	G ₂
Rice, single-cell proteomics (Abiko et al., 2013a)		PCNA, CDKA;1	
Rice sperm, transcriptomics (Abiko et al., 2013b)		KRP1 CDKB1;1 CYCB1-2	
Arabidopsis pollen (Grobei et al., 2009) (Weimer et al., 2018) (Zhao et al., 2012)		CDKA;1, CDKB2;1, CDKB2;2, CYCA2;1, CYCA3;1, CYCB1;3, CYCB2;4, CYCD2;1, CYCD4;1, RBR CYCD7;1-GFP CDKA;1-YFP	
Arabidopsis sperm RNA (Borges et al., 2008)		CDKA;1 (—), CDKB1;2(++), CDKB2;2 (++) , CDKB2;1 (—), RBR (++) , CYCB3;1 (++) , CYCA1s (+), CYCA3 (—), CYCD2;1 (—), RBR (++) , CYCD3;1 & D3;3 s (—), CYCD3;2 (+), CYCD4;1 (+)	
Lilly (Zhao et al., 2013)		CYCD1;1 (+), CYCD5;1 (—), CYCD2; 1(—), KRP1 (—), KRP2 (—), KRP3 (—), CDC6, CDC25/ACR2, CDC48	SC enriched in SCF complex-mediated proteolysis components and CDKB1;1
Tobacco pollen transcriptomics (Hafidh et al., 2012)			CDKA;1 CYCB1;1 CDKB2;2 KRP3

Identified proteins or transcripts in species with pollen described to be in G₁, S or G₂.
Blue. transcripts: purple. native proteins: green. translational fluorescent protein fusions.

[ntbz001.eps]

In Arabidopsis pollen CYCLIND, CDKA and CDKB transcripts and the proteins they encode were detected. Whilst some KRP transcripts were detected their levels were relatively low, suggesting that if they are involved in the S-phase arrest of the SCs, the SCs are preloaded with KRP proteins after the last division. The localisation of CYCD7;1 in the SCs is intriguing given that CYCD7;1-GFP accumulates in the precursor of stomatal guard cells, an unrelated cell type (Weimer et al., 2018).

3.3 Fertilisation and Activation of Cell Division in the Zygote

Fertilisation of both the egg and the CC forms the zygote (2n) and endosperm precursor, which is the starting point for development of the new sporophyte and the endosperm inside the developing seed. The origin of the double fertilisation mechanism and the gain of fitness associated with this development is an intriguing problem in developmental biology. The requirement for the contribution of paternal gene expression to sustain proliferation of the endosperm is ensured by genomic imprinting. The need for the paternal contribution can be bypassed in mutants in the FIS-class genes (Nowack et al., 2007).

Sperm and CC plasmogamy is sufficient to trigger the first divisions of the endosperm precursor, evidenced by the pollination with *cdka;1* mutant sperm which prevents karyogamy of the homodiploid CC and the sperm nucleus but does trigger the first rounds of mitosis (Aw et al., 2010). This suggests that the contribution of the second sperm for the fertilisation of the CC is more involved in the timing of the division, rather than a strict requirement for the division mechanism. In combination with the autonomous endosperm development in some mutant backgrounds, this observation supports the hypothesis of a gametophytic origin of the endosperm (Strassburger, 1900; Nowack et al., 2007), and that the fertilisation by the second sperm, in combination with imprinting provides a mechanism to synchronise the proliferation of the endosperm with the formation of the zygote.

Upon the division of the egg cell, the two daughters not only differ in size but also in developmental fate, hence the first division of the zygote is asymmetric and differential, generating a smaller terminal cell and a larger basal cell. This division establishes the apical–basal axis of the future embryo. Proliferation of the apical cell will develop all the embryonic structures apart from the quiescent centre and root cap. Proliferation of the basal cell will form the quiescent centre and root cap and the suspensor (ten Hove et al., 2015). *De novo* transcription of cell-type specific homeobox genes in both cell types guarantees the polarised development of the embryo proper and suspensor cells (Ueda et al., 2011; Armenta-Medina et al., 2017). This first asymmetric division of the zygote requires extensive reorganisation of the cytoskeleton (Kimata et al., 2016). Upon fertilisation, microtubules form a transverse ring to drive apical cell growth. Actin filaments reorganise to position the nucleus in the apical end of the zygote before the asymmetric division takes place. Furthermore, the anaphase-promoting complex ubiquitin ligase has to be active to target CYCB1;1 for destruction for proper formation of the new cell wall (Guo et al., 2016).

From a cell cycle perspective, this poses several questions. How does successful karyogamy triggers cell cycle progression, and how is an asymmetric division established?

One of the key players during the whole process of sexual reproduction in plants, as in animals and humans, is calcium (Fan et al., 2008; Edel et al., 2017). This universal signalling component is important in many – if not all – intracellular plant cell signalling events. Calcium as second messenger functions either at the ion level or by highly cell type-specific changes in free cytoplasmic concentrations or changes in stored calcium in various intracellular compartments. These so-called ‘calcium signatures’ can be triggered by intracellular and extracellular stimuli (Edel et al., 2017). Pollen development in the anther requires Ca influx from the surrounding sporophytic tissue, and upon landing on the stigma, pollen tube growth through the pistil tissues towards and into the ovule is highly dependent on calcium. Upon entering the micropyle, the entrance site of the ovule near the egg and synergid cells, the growing pollen tube first encounters one of the two synergid cells (SY). The physical interaction between growing pollen tube tip and this SY cell initiates a calcium response of continuous oscillations in the SY, leading to programmed cell death (PCD) and collapse of only this particular SY cell and the release of stored calcium into the cavity where the pollen tube resides (Denninger et al., 2014). This localised increase of free calcium may coincide with slowing down pollen tube growth (Holdaway-Clarke et al., 2003; Hill et al., 2012).

A distinctive short calcium spike response is observed in the egg cell upon SC release whereas upon the arrival of the sperm at the egg cell, which shows a distinctive deformation, a second transient rise in intracellular calcium is detected in the egg cell upon fertilisation, but only after successful karyogamy (Denninger et al., 2014). This second, longer lasting rise in calcium trigger *de novo* cell wall synthesis of the fertilised egg and prevents the unlikely event of polyspermy. Once fertilisation has occurred, the production of the LURE protein, which is the pollen tube attractant, is turned off (Palanivelu and Preuss, 2006; Maruyama et al., 2013).

The second calcium signature, a transient broader spike, detected upon successful karyogamy (Denninger et al., 2014; Palanivelu and Preuss, 2006), besides preventing polyspermy, may also be the intrinsic autonomous and decisive signal in the fertilised egg cell to re-start the cell cycle, and thus early embryogenesis. Importantly, immediately after karyogamy, the first division of the single-celled zygote is asymmetric, producing a small apical cell that will develop further into the embryo and new sporophyte ultimately, whereas a much larger basal cell is responsible for the formation of the suspensor (Yadegari and Drews, 2004). It is tempting to propose a role for various Ca²⁺/calmodulin-dependent protein kinases in this process. Also, of particular interest in this process is the

ZAR1 receptor kinase. *Zygotic arrest 1* was identified as a mutant in which the asymmetric division and the daughter cell fate acquisition was perturbed. *ZAR1* encodes a receptor-like kinase/Pelle kinase (Gish and Clark, 2011) that interacts with calmodulin and the heterotrimeric G protein G β . In *zar1* the basal cell is short and the apical cell has basal cell fate (Yu et al., 2016).

De novo transcription of cell-type specific homeobox genes and miRNA biosynthesis in both cell types guarantees the polarised development of the embryo proper and suspensor cells (Ueda et al., 2011; Armenta-Medina et al., 2017).

4 Concluding Remarks

The complexity of the interactions required for the coordination of fertilisation in seed plants is emerging and advanced live imaging is revealing new features. In this process, the control of the cell cycle is vital to generate the gametes, instigate cell cycle synchrony for gamete fusion and trigger cell division upon successful germination. The identification of cell cycle factors involved has started, and links with coordinating pathways discovered. This research has the potential to provide us with new breeding strategies to enable fertilisation and thus seed setting under challenging conditions.

Related Articles

Sexual Reproduction: From Sexual Differentiation to Fertilization

Embryogenesis

The Genetic and Molecular Control of Ovule Development

The Developmental Biology of Pollen

Plant Embryogenesis—the Cellular Design of a Plant

Endosperm Development

Pollination Signals and Flower Senescence

Re-Staging Plant Mitosis

The Plant Cyclins

CDK Inhibitors

The UPS: An Engine that Drives the Cell Cycle
CDK Phosphorylation
E2F–DP Transcription Factors
Function of the Retinoblastoma-Related Protein in Plants
Cell Cycle Control During Leaf Development
Cell Cycle Control and Fruit Development
Cell Cycle and Endosperm Development
Hormonal Regulation of Cell Cycle Progression and Its Role in Development
Cell Cycle and Environmental Stresses
Calcium Signals and Their Regulation
Signaling by Protein Phosphorylation in Cell Division

REFERENCES

- Abiko, M., Furuta, K., Yamauchi, Y. et al. (2013a). Identification of proteins enriched in rice egg or sperm cells by single-cell proteomics. *PLoS One* **8**: e69578.
- Abiko, M., Maeda, H., Tamura, K. et al. (2013b). Gene expression profiles in rice gametes and zygotes: identification of gamete-enriched genes and up- or down-regulated genes in zygotes after fertilization. *Journal of Experimental Botany* **64**: 1927–1940.
- van Aelst, A.C. and van Went, J.L. (1992). Ultrastructural immunolocalization of pectins and glycoproteins in *Arabidopsis thaliana* pollen grains. *Protoplasma* **168**: 14–19.
- Allen, A.M., Lexer, C., and Hiscock, S.J. (2010). Characterisation of sunflower-21 (SF21) genes expressed in pollen and pistil of *Senecio squalidus* (Asteraceae) and their relationship with other members of the SF21 gene family. *Sexual Plant Reproduction* **23**: 173–186.
- Armenta-Medina, A., Lepe-Soltero, D., Xiang, D. et al. (2017). *Arabidopsis thaliana* miRNAs promote embryo pattern formation beginning in the zygote. *Developmental Biology* **431**: 145–151.
- Aw, S.J., Hamamura, Y., Chen, Z. et al. (2010). Sperm entry is sufficient to trigger division of the central cell but the paternal genome is required for endosperm development in *Arabidopsis*. *Development* **137**: 2683–2690.

- Barrett, S.C. and Hough, J. (2013). Sexual dimorphism in flowering plants. *Journal of Experimental Botany* **64**: 67–82.
- Berger, F., Hamamura, Y., Ingouff, M., and Higashiyama, T. (2008). Double fertilization – caught in the act. *Trends in Plant Science* **13**: 437–443.
- von Besser, K., Frank, A.C., Johnson, M.A., and Preuss, D. (2006). Arabidopsis HAP2 (GCS1) is a sperm-specific gene required for pollen tube guidance and fertilization. *Development* **133**: 4761–4769.
- Biedermann, S., Harashima, H., Chen, P. et al. (2017). The retinoblastoma homolog RBR1 mediates localization of the repair protein RAD51 to DNA lesions in Arabidopsis. *The EMBO Journal* **36**: 1279–1297.
- Borg, M. and Twell, D. (2010). Life after meiosis: patterning the angiosperm male gametophyte. *Biochemical Society Transactions* **38** (2): 577–582.
- Borges, F., Gomes, G., Gardner, R. et al. (2008). Comparative transcriptomics of Arabidopsis sperm cells. *Plant Physiology* **148**: 1168–1181.
- Brownfield, L., Hafidh, S., Borg, M. et al. (2009). A plant germline-specific integrator of sperm specification and cell cycle progression. *PLoS Genetics* **5**: e1000430.
- Bulankova, P., Riehs-Kearnan, N., Nowack, M.K. et al. (2010). Meiotic progression in Arabidopsis is governed by complex regulatory interactions between SMG7, TDM1, and the meiosis I-specific cyclin TAM. *Plant Cell* **22**: 3791–3803.
- Bulankova, P., Akimcheva, S., Fellner, N., and Riha, K. (2013). Identification of Arabidopsis meiotic cyclins reveals functional diversification among plant cyclin genes. *PLoS Genetics* **9**: e1003508.
- Carmichael, J.S. and Friedman, W.E. (1995). Double fertilization in gnetum gnemon: the relationship between the cell cycle and sexual reproduction. *Plant Cell* **7**: 1975–1988.
- Chen, Z., Tan, J.L., Ingouff, M. et al. (2008). Chromatin assembly factor 1 regulates the cell cycle but not cell fate during male gametogenesis in *Arabidopsis thaliana*. *Development* **135**: 65–73.
- Chen, Z., Hafidh, S., Poh, S.H. et al. (2009). Proliferation and cell fate establishment during Arabidopsis male gametogenesis depends on the Retinoblastoma protein. *Proceedings of the National Academy of Sciences of the United States of America* **106**: 7257–7262.

- Cheung, A.Y., Boavida, L.C., Aggarwal, M. et al. (2010). The pollen tube journey in the pistil and imaging the in vivo process by two-photon microscopy. *Journal of Experimental Botany* **61**: 1907–1915.
- Cifuentes, M., Jolivet, S., Cromer, L. et al. (2016). TDM1 regulation determines the number of meiotic divisions. *PLoS Genetics* **12**: e1005856.
- Cresti, M., Lancelle, S.A., and Hepler, P.K. (1987). Structure of the generative cell wall complex after freeze substitution in pollen tubes of *Nicotiana* and *Impatiens*. *Journal of Cell Science* **88**: 373–378.
- Denninger, P., Bleckmann, A., Lausser, A. et al. (2014). Male–female communication triggers calcium signatures during fertilization in Arabidopsis. *Nature Communications* **5**: 4645.
- D'erfurth, I., Cromer, L., Jolivet, S. et al. (2010). The cyclin-A CYCA1;2/TAM is required for the meiosis I to meiosis II transition and cooperates with OSD1 for the prophase to first meiotic division transition. *PLoS Genetics* **6**: e1000989.
- Dewitte, W., Riou-Khamlichi, C., Scofield, S. et al. (2003). Altered cell cycle distribution, hyperplasia, and inhibited differentiation in Arabidopsis caused by the D-type cyclin CYCD3. *The Plant Cell* **15**: 79–92.
- Dong, J. and Yang, H.-Y. (1989). An ultrastructural study embryo sac in *Oryza sativa* L. *Acta Botanica Sinica* **31**: 81–88.
- Dresselhaus, T. and Snell, W.J. (2014). Fertilization: a sticky sperm protein in plants. *Current Biology* **24**: R164–R166.
- Dresselhaus, T., Sprunck, S., and Wessel, G.M. (2016). Fertilization mechanisms in flowering plants. *Current Biology* **26**: R125–R139.
- Duan, Q., Kita, D., Johnson, E.A. et al. (2014). Reactive oxygen species mediate pollen tube rupture to release sperm for fertilization in Arabidopsis. *Nature Communications* **5**: 3129.
- Durberry, A., Vizir, I., and Twell, D. (2005). Male germ line development in Arabidopsis. duo pollen mutants reveal gametophytic regulators of generative cell cycle progression. *Plant Physiology* **137**: 297–307.
- Ebel, C., Mariconti, L., and Gruissem, W. (2004). Plant retinoblastoma homologues control nuclear proliferation in the female gametophyte. *Nature* **429**: 776–780.
- Edel, K.H., Marchadier, E., Brownlee, C. et al. (2017). The evolution of calcium-based signalling in plants. *Current Biology* **27**: R667–R679.

- Eloy, N.B., Gonzalez, N., van Leene, J. et al. (2012). SAMBA, a plant-specific anaphase-promoting complex/cyclosome regulator is involved in early development and A-type cyclin stabilization. *Proceedings of the National Academy of Sciences of the United States of America* **109**: 13853–13858.
- Englhart, M., Soljic, L., and Sprunck, S. (2017). Manual isolation of living cells from the *Arabidopsis thaliana* female gametophyte by micromanipulation. *Methods in Molecular Biology* **1669**: 221–234.
- Esau, K. (1962). *Anatomy of Seed Plants*. Wiley.
- Escobar-Restrepo, J.-M., Huck, N., Kessler, S. et al. (2007). The FERONIA receptor-like kinase mediates male-female interactions during pollen tube reception. *Science* **317**: 656–660.
- Fan, Y.-F., Jiang, L., Gong, H.-Q., and Liu, C.-M. (2008). Sexual reproduction in higher plants I: fertilization and the initiation of zygotic program. *Journal of Integrative Plant Biology* **50**: 860–867.
- Friedman, W.E. (1999). Expression of the cell cycle in sperm of *Arabidopsis*: implications for understanding patterns of gametogenesis and fertilization in plants and other eukaryotes. *Development* **126**: 1065–1075.
- Friedman, W.E. (2006). Embryological evidence for developmental lability during early angiosperm evolution. *Nature* **441**: 337–340.
- Friedman, W.E. and Ryerson, K.C. (2009). Reconstructing the ancestral female gametophyte of angiosperms: insights from *Amborella* and other ancient lineages of flowering plants. *American Journal of Botany* **96**: 129–143.
- Fujii, S., Kubo, K.-I., and Takayama, S. (2016). Non-self- and self-recognition models in plant self-incompatibility. *Nature Plants* **2**: 16130.
- Ge, Z., Bergonci, T., Zhao, Y. et al. (2017). *Arabidopsis* pollen tube integrity and sperm release are regulated by RALF-mediated signaling. *Science* **358**: 1596–1600.
- Genschik, P., Marrocco, K., Bach, L. et al. (2014). Selective protein degradation: a rheostat to modulate cell-cycle phase transitions. *Journal of Experimental Botany* **65**: 2603–2615.
- Gish, L.A. and Clark, S.E. (2011). The RLK/Pelle family of kinases. *The Plant Journal* **66**: 117–127.
- Glöckle, B., Urban, W.J., Nagahara, S. et al. (2018). Pollen differentiation as well as pollen tube guidance and discharge

- are independent of the presence of gametes. *Development* **145**: pii: dev152645.
- Gómez, J.F., Talle, B., and Wilson, Z.A. (2015). Anther and pollen development: a conserved developmental pathway. *Journal of Integrative Plant Biology* **57**: 876–891.
- de Graaf, B.H.J., Knuiman, B., Bosch, M.J.M., and Mariani, C. (2000). The class III pistil-specific extensin-like proteins of *Nicotiana tabacum* show arabinogalactan-protein-like characteristics and are non-specifically translocated through pollen tube walls in vivo. In: *Cell and Developmental Biology of Arabinogalactan-Proteins* (eds E.A. Nothnagel, A. Bacic and A.E. Clarke). Boston, MA: Springer US.
- de Graaf, B.H.J., Derksen, J.W.M., and Mariani, C. (2001). Pollen and pistil in the progamic phase. *Sexual Plant Reproduction* **14**: 41–55.
- Grant-Downton, R., Kourmpetli, S., Hafidh, S. et al. (2013). Artificial microRNAs reveal cell-specific differences in small RNA activity in pollen. *Current Biology* **23**: R599–R601.
- Grobei, M.A., Qeli, E., Brunner, E. et al. (2009). Deterministic protein inference for shotgun proteomics data provides new insights into *Arabidopsis* pollen development and function. *Genome Research* **19**: 1786–1800.
- Guo, L., Jiang, L., Zhang, Y. et al. (2016). The anaphase-promoting complex initiates zygote division in *Arabidopsis* through degradation of cyclin B1. *Plant Journal* **86** (2): 161–174. doi: 10.1111/tpj.13158.
- Hafidh, S., Breznenova, K., and Honys, D. (2012). De novo post-pollen mitosis II tobacco pollen tube transcriptome. *Plant Signaling & Behavior* **7**: 918–921.
- Hamamura, Y., Saito, C., Awai, C. et al. (2011). Live-cell imaging reveals the dynamics of two sperm cells during double fertilization in *Arabidopsis thaliana*. *Current Biology* **21**: 497–502.
- Hilhorst, H.W.M., Finch-Savage, W.E., Buitink, J. et al. (2010). Dormancy in plant seeds. In: *Dormancy and Resistance in Harsh Environments* (eds E. Lubzens, J. Cerda and M. Clark). Berlin, Heidelberg: Springer Berlin Heidelberg.
- Hill, A., Shachar-Hill, B., Skepper, J. et al. (2012). An osmotic model of the growing pollen tube. *PLoS One* **7**: e36585.
- Hiscock, S.J. (2011). Sexual plant reproduction. *Annals of Botany* **108**: 585–587.
- Holdaway-Clarke, T.L., Weddle, N.M., Kim, S. et al. (2003). Effect of extracellular calcium, pH and borate on growth oscillations in

- Lilium formosanum* pollen tubes. *Journal of Experimental Botany* **54**: 65–72.
- Horvath, B.M., Kourova, H., Nagy, S. et al. (2017). Arabidopsis RETINOBLASTOMA RELATED directly regulates DNA damage responses through functions beyond cell cycle control. *The EMBO Journal* **36**: 1261–1278.
- Huang, B.-Q. and Russell, S.D. (1994). Fertilization in *Nicotiana tabacum*: cytoskeletal modifications in the embryo sac during synergid degeneration. *Planta* **194**: 200–214.
- Huang, X.Y., Niu, J., Sun, M.X. et al. (2013). CYCLIN-DEPENDENT KINASE G1 is associated with the spliceosome to regulate CALLOSE SYNTHASE5 splicing and pollen wall formation in Arabidopsis. *Plant Cell* **25**: 637–648.
- Igawa, T., Yanagawa, Y., Miyagishima, S.-Y., and Mori, T. (2013). Analysis of gamete membrane dynamics during double fertilization of Arabidopsis. *Journal of Plant Research* **126**: 387–394.
- Iwakawa, H., Shinmyo, A., and Sekine, M. (2006). Arabidopsis CDKA;1, a cdc2 homologue, controls proliferation of generative cells in male gametogenesis. *The Plant Journal* **45**: 819–831.
- Jean-Emmanuel, F., Nicolas, R., Philippe, F., and Christian, D. (2002). Fertilization in *Arabidopsis thaliana* wild type: developmental stages and time course. *The Plant Journal* **30**: 481–488.
- Jiang, H., Yi, J., Boavida, L.C. et al. (2015). Intercellular communication in *Arabidopsis thaliana* pollen discovered via *AHG3* transcript movement from the vegetative cell to sperm. *Proceedings of the National Academy of Sciences of the United States of America* **112** (43): 13378–13383.
- Kawashima, T., Maruyama, D., Shagirov, M. et al. (2014). Dynamic F-actin movement is essential for fertilization in *Arabidopsis thaliana*. *eLife* **3**: e04501.
- Khalequzzaman, M. and Haq, N. (2005). Isolation and in vitro fusion of egg and sperm cells in *Oryza sativa*. *Plant Physiology and Biochemistry* **43**: 69–75.
- Kim, H.J., Oh, S.A., Brownfield, L. et al. (2008). Control of plant germline proliferation by SCF(FBL17) degradation of cell cycle inhibitors. *Nature* **455**: 1134–1137.
- Kimata, Y., Higaki, T., Kawashima, T. et al. (2016). Cytoskeleton dynamics control the first asymmetric cell division in *Arabidopsis* zygote. *Proceedings of the National Academy of*

Sciences of the United States of America **113** (49): 14157–14162.

- Kohler, C., Wolff, P., and Spillane, C. (2012). Epigenetic mechanisms underlying genomic imprinting in plants. *Annual Review of Plant Biology* **63**: 331–352.
- Kranz, E., Wiegen, P., and Lörz, H. (1995). Early cytological events after induction of cell division in egg cells and zygote development following in vitro fertilization with angiosperm gametes. *The Plant Journal* **8**: 9–23.
- Kranz, E., Hoshino, Y., and Okamoto, T. (2008). In vitro fertilization with isolated higher plant gametes. In: *Plant Embryogenesis* (eds M.F. Suárez and P.V. Bozhkov). Totowa, NJ: Humana Press.
- Lan, L., Chen, W., Lai, Y. et al. (2004). Monitoring of gene expression profiles and isolation of candidate genes involved in pollination and fertilization in rice (*Oryza sativa* L.) with a 10K cDNA microarray. *Plant Molecular Biology* **54**: 471–487.
- Lennon, K.A., Roy, S., Hepler, P.K., and Lord, E.M. (1998). The structure of the transmitting tissue of *Arabidopsis thaliana* (L.) and the path of pollen tube growth. *Sexual Plant Reproduction* **11**: 49–59.
- Leydon, A.R. and Johnson, M.A. (2013). Preparing for explosion: pollen tubes are capacitated by the pistil. *Molecular Reproduction and Development* **80**: 427–427.
- Leydon, A.R., Beale, K.M., Woroniecka, K. et al. (2013). Three MYB transcription factors control pollen tube differentiation required for sperm release. *Current Biology: CB* **23**: 1209–1214.
- Lin, S.-Y., Chen, P.-W., Chuang, M.-H. et al. (2014). Profiling of transcriptomes of in vivo-grown pollen tubes reveals genes with roles in micropylar guidance during pollination in *Arabidopsis*. *The Plant Cell Online* **26**: 602–618.
- Liu, X., Castro, C., Wang, Y. et al. (2016). The role of LORELEI in pollen tube reception at the interface of the synergid cell and pollen tube requires the modified eight-cysteine motif and the receptor-like kinase FERONIA. *The Plant Cell* **28**: 1035–1052.
- Lughadha, E.N., Govaerts, R., Belyaeva, I. et al. (2016). Counting counts: revised estimates of numbers of accepted species of flowering plants, seed plants, vascular plants and land plants with a review of other recent estimates. *Phytotaxa* **272**: 82.
- Magyar, Z., Horváth, B., Khan, S. et al. (2012). *Arabidopsis* E2FA stimulates proliferation and endocycle separately through RBR-bound and RBR-free complexes. *The EMBO Journal* **31**: 1480–1493.

- Márton, M.L., Fastner, A., Uebler, S., and Dresselhaus, T. (2012). Overcoming hybridization barriers by the secretion of the maize pollen tube attractant ZmEA1 from *Arabidopsis* ovules. *Current Biology* **22**: 1194–1198.
- Maruyama, D., Hamamura, Y., Takeuchi, H. et al. (2013). Independent control by each female gamete prevents the attraction of multiple pollen tubes. *Developmental Cell* **25**: 317–323.
- Maruyama, D., Volz, R., Takeuchi, H. et al. (2015). Rapid elimination of the persistent synergid through a cell fusion mechanism. *Cell* **161**: 907–918.
- Mccue, A.D., Cresti, M., Feijó, J.A., and Slotkin, R.K. (2011). Cytoplasmic connection of sperm cells to the pollen vegetative cell nucleus: potential roles of the male germ unit revisited. *Journal of Experimental Botany* **62**: 1621–1631.
- Mecchia, M.A., Santos-Fernandez, G., Duss, N.N. et al. (2017). RALF4/19 peptides interact with LRX proteins to control pollen tube growth in *Arabidopsis*. *Science* **358**: 1600–1603.
- Mercier, R., Vezon, D., Bullier, E. et al. (2001). SWITCH1 (SWI1): a novel protein required for the establishment of sister chromatid cohesion and for bivalent formation at meiosis. *Genes & Development* **15**: 1859–1871.
- Mizukami, A.G., Inatsugi, R. et al. (2016). The AMOR Arabinogalactan sugar chain induces pollen-tube competency to respond to ovular guidance. *Current Biology* **26**: 1091–1097.
- Mizuta, Y. and Higashiyama, T. (2018). Chemical signaling for pollen tube guidance at a glance. *Journal of Cell Science* **131**.
- Mori, T., Kuroiwa, H., Higashiyama, T., and Kuroiwa, T. (2005). GENERATIVE CELL SPECIFIC 1 is essential for angiosperm fertilization. *Nature Cell Biology* **8**: 64.
- Mori, T., Igawa, T., Tamiya, G. et al. (2014). Gamete attachment requires GEX2 for successful fertilization in *Arabidopsis*. *Current Biology* **24**: 170–175.
- Muschietti, J.P., and Diego, L.W. (2018). How many receptor-like kinases are required to operate a pollen tube. *Current Opinion in Plant Biology* **41**: 73–82.
- ten Hove, C.A., Lu, K.J., and Weijers, D. (2015). Building a plant: cell fate specification in the early *Arabidopsis* embryo. *Development* **142**: 420–430.
- Nowack, M.K., Grini, P.E., Jakoby, M.J. et al. (2006). A positive signal from the fertilization of the egg cell sets off endosperm

- proliferation in angiosperm embryogenesis. *Nature Genetics* **38**: 63–67.
- Nowack, M.K., Shirzadi, R., Dissmeyer, N. et al. (2007). Bypassing genomic imprinting allows seed development. *Nature* **447**: 312–315.
- Ohnishi, Y., Hoshino, R., and Okamoto, T. (2014). Dynamics of male and female chromatin during karyogamy in rice zygotes. *Plant Physiology* **165**: 1533–1543.
- Okamoto, T. (2010). Gamete fusion site on the egg cell and autonomous establishment of cell polarity in the zygote. *Plant Signaling & Behavior* **5**: 1464–1467.
- Okuda, S., Suzuki, T., Kanaoka, M.M. et al. (2013). Acquisition of LURE-binding activity at the pollen tube tip of *Torenia fournieri*. *Molecular Plant* **6**: 1074–1090.
- Palanivelu, R. and Preuss, D. (2006). Distinct short-range ovule signals attract or repel *Arabidopsis thaliana* pollen tubes in vitro. *BMC Plant Biology* **6**: 7–7.
- Pawlowski, W.P., Wang, C.J., Golubovskaya, I.N. et al. (2009). Maize AMELOTIC1 is essential for multiple early meiotic processes and likely required for the initiation of meiosis. *Proceedings of the National Academy of Sciences of the United States of America* **106**: 3603–3608.
- Pereira, A.M., Lopes, A.L., and Coimbra, S. (2016a). JAGGER, an AGP essential for persistent synergid degeneration and polytubey block in *Arabidopsis*. *Plant Signaling & Behavior* **11**: e1209616.
- Pereira, A.M., Nobre, M.S., Pinto, S.C. et al. (2016b). "Love Is Strong, and You're so Sweet": JAGGER is essential for persistent synergid degeneration and polytubey block in *Arabidopsis thaliana*. *Molecular Plant* **9**: 601–614.
- Polyn, S., Willems, A., and de Veylder, L. (2015). Cell cycle entry, maintenance, and exit during plant development. *Current Opinion in Plant Biology* **23**: 1–7.
- Qin, Y., Leydon, A.R., Manziello, A. et al. (2009). Penetration of the stigma and style elicits a novel transcriptome in pollen tubes, pointing to genes critical for growth in a pistil. *PLoS Genetics* **5**: e1000621–e1000621.
- Rousseau, D.-D., Schevin, P., Ferrier, J. et al. (2008). Long-distance pollen transport from North America to Greenland in spring. *Journal of Geophysical Research* **113**: G02013. doi: 10.1029/2007JG000456.
- Santos, M.R., Bispo, C., and Becker, J.D. (2017). Isolation of *Arabidopsis* pollen, sperm cells, and vegetative nuclei by

- fluorescence-activated cell sorting (FACS). *Methods in Molecular Biology* **1669**: 193–210.
- Schmid, M.W., Schmidt, A., and Grossniklaus, U. (2015). The female gametophyte: an emerging model for cell type-specific systems biology in plant development. *Frontiers in Plant Science* **6**: 907.
- Schnittger, A. and de Veylder, L. (2018). The dual face of cyclin. *Trends in Plant Sciences* **23** (6): 475–478.
- Schoft, V.K., Chumak, N., Bindics, J. et al. (2015). SYBR green-activated sorting of *Arabidopsis* pollen nuclei based on different DNA/RNA content. *Plant Reproduction* **28**: 61–72.
- de Schutter, K., Joubès, J., Cools, T. et al. (2007). *Arabidopsis* WEE1 kinase controls cell cycle arrest in response to activation of the DNA integrity checkpoint. *The Plant Cell* **19**: 211–225.
- Scott, R.J., Armstrong, S.J., Doughty, J., and Spielman, M. (2008). Double fertilization in *Arabidopsis thaliana* involves a polyspermy block on the egg but not the central cell. *Molecular Plant* **1**: 611–619.
- Shen-Miller, J., Mudgett, M.B., Schopf, J.W. et al. (1995). Exceptional seed longevity and robust growth: ancient Sacred Lotus from China. *American Journal of Botany* **82**: 1367–1380.
- Simonin, K.A. and Roddy, A.B. (2018). Genome downsizing, physiological novelty, and the global dominance of flowering plants. *PLoS Biology* **16** (1): e2003706.
- Slotkin, R.K., Vaughn, M., Borges, F. et al. (2009). Epigenetic reprogramming and small RNA silencing of transposable elements in pollen. *Cell* **136**: 461–472.
- Sornay, E., Forzani, C., Forero-Vargas, M. et al. (2015). Activation of CYCD7;1 in the central cell and early endosperm overcomes cell-cycle arrest in the *Arabidopsis* female gametophyte, and promotes early endosperm and embryo development. *The Plant Journal* **84**: 41–55.
- Southworth, D. and Kwiatkowski, S. (1996). Arabinogalactan proteins at the cell surface of *Brassica* sperm and *Lilium* sperm and generative cells. *Sexual Plant Reproduction* **9**: 269–272.
- Sprunck, S. and Gross-Hardt, R. (2011). Nuclear behavior, cell polarity, and cell specification in the female gametophyte. *Sexual Plant Reproduction* **24**: 123–136.
- Sprunck, S., Rademacher, S., Vogler, F. et al. (2012). Egg cell-secreted EC1 triggers sperm cell activation during double fertilization. *Science* **338**: 1093–1097.

- Strassburger, E. (1900). *Lehrbuch der Botanik für Hochschulen*. Jena: Verlag von Gustav Fischer.
- Sukawa, Y. and Okamoto, T. (2018). Cell cycle in egg cell and its progression during zygotic development in rice. *Plant Reproduction* **31**: 107–116.
- Takatsuka, H., Umeda-Hara, C., and Umeda, M. (2015). Cyclin-dependent kinase-activating kinases CDKD;1 and CDKD;3 are essential for preserving mitotic activity in *Arabidopsis thaliana*. *The Plant Journal* **82**: 1004–1017.
- Takeuchi, H. and Higashiyama, T. (2012). A species-specific cluster of defensin-like genes encodes diffusible pollen tube attractants in *Arabidopsis*. *PLoS Biology* **10**: e1001449.
- Tekleyohans, D.G., Mao, Y., Kägi, C. et al. (2017). Polyspermy barriers: a plant perspective. *Current Opinion in Plant Biology* **35**: 131–137.
- Twell, D. (2011). Male gametogenesis and germline specification in flowering plants. *Sexual Plant Reproduction* **24**: 149–160.
- Ueda, M., Zhang, Z., and Laux, T. (2011). Transcriptional activation of arabidopsis axis patterning genes WOX8/9 links zygote polarity to embryo development. *Developmental Cell* **20**: 264–270.
- de Veylder, L., Beeckman, T., and Inze, D. (2007). The ins and outs of the plant cell cycle. *Nature Reviews. Molecular Cell Biology* **8**: 655–665.
- Völz, R., Heydlauff, J., Ripper, D. et al. (2013). Ethylene signaling is required for synergid degeneration and the establishment of a pollen tube block. *Developmental Cell* **25**: 310–316.
- Wang, D.-Y., Zhang, Q., Liu, Y. et al. (2010). The levels of male gametic mitochondrial DNA are highly regulated in angiosperms with regard to mitochondrial inheritance. *The Plant Cell* **22**: 2402–2416.
- Wang, J.-G., Feng, C., Liu, H.-H. et al. (2016). HAPLESS13-mediated trafficking of STRUBBELIG is critical for ovule development in arabidopsis. *PLoS Genetics* **12** (8): e1006269.
- Wang, T., Liang, L., Xue, Y. et al. (2016). A receptor heteromer mediates the male perception of female attractants in plants. *Nature* **531**: 241.
- Weimer, A.K., Biedermann, S., Harashima, H. et al. (2016). The plant-specific CDKB1-CYCB1 complex mediates homologous recombination repair in *Arabidopsis*. *The EMBO Journal* **35**: 2068–2086.
- Weimer, A.K., Matos, J.L., Sharma, N. et al. (2018). Lineage- and stage-specific expressed CYCD7;1 coordinates the single

symmetric division that creates stomatal guard cells. *Development* **145**.

- Yadegari, R. and Drews, G.N. (2004). Female gametophyte development. *The Plant Cell* **16**: S133–S141.
- Yu, T.Y., Shi, D.Q., Jia, P.F. et al. (2016). The Arabidopsis receptor kinase ZAR1 is required for zygote asymmetric division and its daughter cell fate. *PLoS Genetics* **12**: e1005933.
- Zhang, Z., Tian, H.Q., and Russell, S.D. (1999). Localization of myosin on sperm-cell-associated membranes of tobacco (*Nicotiana tabacum* L.). *Protoplasma* **208**: 123–128.
- Zhang, J., Huang, Q., Zhong, S. et al. (2017a). Sperm cells are passive cargo of the pollen tube in plant fertilization. *Nature Plants* **3**: 17079.
- Zhang, Z., Hu, M., Feng, X. et al. (2017b). Proteomes and phosphoproteomes of anther and pollen: availability and progress. *Proteomics* **17**: 1600458. doi: 10.1002/pmic.201600458.
- Zhao, X., Harashima, H., Dissmeyer, N. et al. (2012). A general G1/S-phase cell-cycle control module in the flowering plant *Arabidopsis thaliana*. *PLoS Genetics* **8**: e1002847.
- Zhao, X., Yang, N., and Wang, T. (2013). Comparative proteomic analysis of generative and sperm cells reveals molecular characteristics associated with sperm development and function specialization. *Journal of Proteome Research* **12**: 5058–5071.
- Zhao, X., Bramsiepe, J., van Durme, M. et al. (2017). RETINOBLASTOMA RELATED1 mediates germline entry in Arabidopsis. *Science* **356**.
- Zheng, B., He, H., Zheng, Y. et al. (2014a). An ARID domain-containing protein within nuclear bodies is required for sperm cell formation in *Arabidopsis thaliana*. *PLoS Genetics* **10**: e1004421.
- Zheng, T., Nibau, C., Phillips, D.W. et al. (2014b). CDKG1 protein kinase is essential for synapsis and male meiosis at high ambient temperature in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences of the United States of America* **111**: 2182–2187.
- Zhou, L., Juranic, M., and Dresselhaus, T. (2017). Germline development and fertilization mechanisms in maize. *Molecular Plant* **10** (3): 389–401.