

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/123902/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Jack, Alison A., Nordli, Henriette R., Powell, Lydia C. , Farnell, Damian J. J. , Pukstad, Brita, Rye, Philip D., Thomas, David W., Chinga-Carrasco, Gary and Hill, Katja E. 2019. Cellulose nanofibril formulations incorporating a low molecular weight alginate oligosaccharide modify bacterial biofilm development. *Biomacromolecules* 20 (8) , pp. 2953-2961. 10.1021/acs.biomac.9b00522

Publishers page: <http://dx.doi.org/10.1021/acs.biomac.9b00522>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.






Figure S1. Effect of CNF dispersions and dried materials (aerogels and films) on growth of *S. aureus* 1061A: Growth curves of *S. aureus* in T3 CNF dispersions (0.2 %) in (a) MH broth, (b) Phosphate buffered saline (PBS), (c) sterile distilled water (n=3). (d) Log₁₀ reduction values of planktonic *S. aureus* 1061A incubated with CNF formulations (A0.2, A0.4, A0.6 or F0.2) for 24 h (n=3). [†]≥3 log₁₀ CFU/ml indicates bacteriocidal activity. (e) Images of the dried CNF materials: aerogels (A0.2, A0.4 and A0.6) and film (F0.2).




Figure S2. COMSTAT image analysis of LIVE/DEAD® staining of 24 h single and dual species biofilms grown on (a, b) 0.2% CNF films (Control F0.2, OligoG-coated F0.2G and PEG400 biocomposite F0.2P). (c) 0.2% CNF aerogels (Control A0.2 and Oligo-bio-nanocomposite A0.2G) showing (a) mean biofilm thickness. (b, c) roughness coefficients of LIVE (green) and DEAD (red) cells (*significantly different compared to the equivalent no OligoG control; n=10 images each from n=3 replicates; P<0.05).