

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/123966/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Furby, Hannah , Warnert, Esther, Marley, Christopher, Bailey, Damian and Wise, Richard 2020. Cardiorespiratory fitness is associated with increased middle cerebral arterial compliance and decreased cerebral blood flow in young healthy adults: a pulsed ASL MRI study. *Journal of Cerebral Blood Flow and Metabolism* 40 (9) , pp. 1879-1889. 10.1177/0271678X19865449

Publishers page: <https://doi.org/10.1177/0271678X19865449>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies. See <http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Original Article

Cardiorespiratory fitness is associated with increased middle cerebral arterial compliance and decreased cerebral blood flow in young healthy adults: a pulsed ASL MRI study

Hannah V Furby^{1,4}, Esther AH Warnert^{1,3}, Christopher J Marley², Damian M Bailey², Richard G Wise¹

¹Cardiff University Brain Research Imaging Centre (CUBRIC), School of Psychology, Cardiff University, Cardiff, UK

²Neurovascular Research Laboratory, Faculty of Life Sciences and Education, University of South Wales, Pontypridd, UK.

³Department of Radiology and Nuclear Medicine, Erasmus MC, Rotterdam, the Netherlands

⁴UCL Queen Square Institute of Neurology, University College London, London, UK

Corresponding Author:

Professor RG Wise, CUBRIC, School of Psychology, Cardiff University, Maindy Road, Cardiff CF24 4HQ, UK.

E-mail: wiserg@cardiff.ac.uk

HVF received support from the Neuroscience and Mental Health Research Institute Scholarship from Cardiff University. RGW and DMB acknowledge the support of the Higher Education Funding Council for Wales. DMB acknowledges financial support from the JPR Williams Trust. DMB is supported by a Royal Society Wolfson Research Fellowship (#WM170007)

Abstract = 175

Total Words = 5,157 (max. 6,000)

Figures = 3

Tables=1

Abstract

Cardiorespiratory fitness is thought to have beneficial effects on systemic vascular health, in part, by decreasing arterial stiffness. However, in the absence of non-invasive methods, it remains unknown whether this effect extends to the cerebrovasculature. The present study uses a novel pulsed arterial spin labelling (pASL) technique to explore the relationship between cardiorespiratory fitness and arterial compliance of the middle cerebral arteries (MCAC). Other markers of cerebrovascular health, including resting cerebral blood flow (CBF) and cerebrovascular reactivity to CO₂ (CVR_{CO2}) were also investigated. Eleven healthy males aged 21±2 years with varying levels of cardiorespiratory fitness (maximal oxygen uptake ($\dot{V}O_{2MAX}$) 38-76 ml/min/kg) underwent MRI scanning at 3 Tesla. Higher $\dot{V}O_{2MAX}$ was associated with greater MCAC ($R^2=0.64$, $p<0.01$) and lower resting grey matter CBF ($R^2=0.75$, $p<0.01$). However, $\dot{V}O_{2MAX}$ was not predictive of global grey matter BOLD-based CVR ($R^2=0.47$, $p=0.17$) or CBF-based CVR ($R^2=0.19$, $p=0.21$). The current experiment builds upon the established benefits of exercise on arterial compliance in the systemic vasculature, by showing that increased cardiorespiratory fitness is associated with greater cerebral arterial compliance in early adulthood.

Keywords: arterial compliance; ASL; CBF; CVR; fitness

Introduction

Physical exercise is well known for its cardiovascular benefits¹, yet the challenge remains of identifying how exercise is beneficial to the brain. Although studies using ultrasound methods have reported increases in resting cerebral blood flow velocity² and cerebrovascular reactivity³ associated with cardiorespiratory fitness, these methods often lack spatial specificity, reliability and consistency across individuals⁴. More recently, advances in arterial spin labelling (ASL) magnetic resonance imaging (MRI) have started to offer non-invasive measures of cerebrovascular function with enhanced spatial sensitivity for quantifying individual differences in cerebral dynamics compared to ultrasound methods⁵.

Cerebral arterial compliance (AC) permits the arteries and arterioles to buffer pressure pulsations that arise from the heart, smoothing blood flow to the capillaries. Cerebrovascular reactivity (CVR) refers to dilation or constriction of vessels to control cerebral blood flow (CBF), relying on complex signalling processes. In the healthy brain, compliance and reactivity work together to regulate local blood flow, protect against fluctuations in blood pressure and preserve autoregulation⁶. Formation of arterial plaques or vessel stiffening, which occur naturally in ageing and disease⁷, can disrupt these vascular mechanisms thereby putting the downstream microvasculature at risk; a potential contributor to small vessel disease⁸ and cognitive decline⁹.

Ultrasound imaging with simultaneous arterial applanation tonometry of the arterial waveform, has shown that central arterial stiffness is reduced in those who exercise regularly^{10,11}. Due to a limited ability to assess diameters of intracranial arteries, however, ultrasound techniques are currently restricted to providing

information about blood velocity and not volume or flow. One ultrasound method, transcranial Doppler (TCD) sonography, is only able to inform us about compliance of a distal vascular bed and not the local stiffness profile of the larger cerebral vessels themselves¹². Optical imaging methods have also demonstrated a relationship between cardiorespiratory fitness and cerebral AC, as well as a regional correspondence with age and cognitive function¹³. This method has the advantage, over TCD, of extracting cerebral pulsatile waveform measurements from the arteries over which they are placed¹⁴ yet despite this spatial advantage, the limited penetration of optical imaging precludes examination of the deeper vasculature.

Due to the added spatial resolution, MRI methods therefore allow more precise quantification of the *local* arterial wall properties rather than those distal to the site of measurement. ASL MRI is primarily used to map CBF. However, using novel ASL methods that measure changes in arterial blood volume (aBV) within cerebral arteries throughout the cardiac cycle^{15,16}, it is possible to estimate AC in the major cerebral arteries.

The present study examined the association between cardiorespiratory fitness ($\dot{V}O_{2MAX}$) and middle cerebral AC in a cohort of healthy young males. Evidence, although indirect, of the systemic vasculature has demonstrated a greater vessel compliance in fitter young adults¹⁷ and greater AC in fronto-parietal regions has been associated with increased fitness using optical imaging methods across ages¹³. We hypothesised that greater cardiorespiratory fitness would predict higher AC using the ASL MRI method¹⁸ and that those with higher aerobic fitness ($\dot{V}O_{2MAX}$) would show elevated baseline CBF, as has been seen previously in ASL studies of children¹⁹, older

adults²⁰ and patients²¹. Using a breath-hold stimulus, we also probed the relationship between $\dot{V}O_{2MAX}$ and MRI measures of both BOLD- and CBF-based CVR²².

Materials and Methods

Participants

Eleven healthy males, aged 21 ± 2 years old, provided informed consent under ethical approval from the University of South Wales and Cardiff University School of Psychology Ethics Committees. All experiments were performed in accordance with the guidelines stated in the Cardiff University Research Framework (version 4.0, 2010). We specifically chose to exclude females as oestrogen levels (during the menstrual cycle, menopause, and hormone replacement therapy) have been associated with intracranial vasodilatation and increased CBF²³. In order to recruit a wide fitness range, participants who engaged in >150 minutes per week of self-reported moderate-to-vigorous intensity recreational aerobic activity were recruited from running and cycling clubs, while general University wide advertisement and word of mouth was used to recruit more sedentary participants. Clubs that involved higher impact sports e.g. rugby, were excluded in this study to avoid complications that may arise from a history of concussion.

Subjects underwent a detailed clinical examination that included 12-lead functional diagnostic exercise electrocardiography (ECG) and were excluded if they showed signs of, or reported, any cardiovascular, cerebrovascular or respiratory disease. Participants were also screened by self-report for any neurological or psychiatric illnesses, regular smoking or prescribed medication. Individual differences in haematocrit (Hct) were assessed by sampling capillary blood from the middle

finger. Samples were centrifuged for 10-min via ultracentrifugation and a micro-haematocrit reader (Hawksley and Sons Ltd, Sussex, England) used to quantify Hct. Three samples were acquired and mean Hct reported.

Study Design

All participants took part in two separate testing sessions. Participants first underwent cardiorespiratory fitness testing at the University of South Wales and were then followed up for a second visit at Cardiff University Brain Research Imaging Centre, where they underwent 3T MRI. Prior to each visit, participants were asked to refrain from drinking caffeinated drinks, taking any recreational drugs or engaging in any exhaustive exercise that may elevate heart rate and subsequently confound CBF measurements.

Cardiorespiratory Fitness Testing

The $\dot{V}O_{2\text{MAX}}$ test is a test of maximal oxygen uptake and is an established ²⁴ of cardiorespiratory fitness, where $\dot{V}O_{2\text{MAX}}$ refers to the highest rate at which oxygen can be taken up and consumed by the body during intense exercise.

Online respiratory gas analysis (Medgraphics, MA, USA) was performed during an incremental cycling exercise test to volitional exhaustion on an electronically braked, semi-recumbent cycle ergometer (Lode Corival, Cranlea & Company, UK) for the specific determination of ventilation, $\dot{V}O_2$ and $\dot{V}CO_2$. The test began with 2 minutes of rest, followed by 5 minutes of unloaded pedalling (0W) and increased by 5W every

$\dot{V}O_{2\text{MAX}}$ predicts MCA compliance and baseline CBF

HV Furby *et al.*

10s thereafter. Participants were required to maintain a cadence of ~70 revolutions per minute (RPM). Maximum exertion and corresponding $\dot{V}O_{2\text{MAX}}$ was confirmed when at least two of the following established criteria were met: 1) Failure to increase $\dot{V}O_2$ with increasing exercise load 2) a respiratory exchange ratio (RER; the ratio between $\dot{V}CO_2$ and $\dot{V}O_2$ during cycling) of >1.15, or 3) a heart rate within 10 beats of an age-predicted maximum (*i.e.* 220 – age in years) (e.g. Barnes *et al.*, 2013).

MRI data acquisition

All scanning was carried out using a 3T GE HDx scanner (GE Healthcare, Milwaukee, WI, USA) equipped with an 8-channel receive-only head coil. All participants underwent whole-brain T₁-weighted structural scans (3D FSPGR, 1x1x1 mm³ voxels, TI= 450ms, TR =7.8ms, TE = 3 ms) for registration purposes.

Middle Cerebral Artery Compliance and Grey Matter CBF

A multi-inversion time (MTI) pulsed ASL acquisition was performed at rest. A Proximal Inversion with Control of Off-Resonance Effects (PICORE) ASL sequence was used to improve the profile of the labelling slice. A QUIPSS II (quantitative imaging of perfusion using a single subtraction) cut-off was also applied at 700ms²⁶ to reduce the sensitivity of the arterial transit time. Ten inversion times (TI's) were acquired, whereby short (TI's = 250, 350, 450, 550, 650ms) medium (TI's = 750, 850) and long TI's (TI's = 1,000, 1,500, 2000ms) were acquired as separate scans in which the label (width=200mm) was applied 10mm below the most proximal slice. Images were acquired with similar parameters to those described elsewhere¹⁸ using a spiral readout single shot gradient

echo sequence (TE=2.7ms) with the following acquisition parameters: a variable repetition time (1,000ms to 3,400ms), eight control–tag pairs per TI, 12 slices, slice gap=1mm, voxel size=3x3x7mm³. Total acquisition time was ~18 minutes. For quantification of perfusion, a (M_0) calibration scan was acquired without labelling in which the same acquisition parameters were applied as above, but with a long TR.

Cerebrovascular reactivity

A breath-hold paradigm was carried out as described elsewhere²⁷. Participants were instructed to complete five end-expiration breath-holds (15s each) interleaved with 30s periods of paced breathing at a rate of 12 breaths per minute²⁸. After each breath-hold the subject was cued to exhale first to obtain a measure of peak end-tidal CO₂. Total scan duration was approximately four minutes during which quantitative arterial spin labelling (pASL) and BOLD-weighted images were acquired with a single-shot PICORE QUIPSS II (Wong *et al.*, 1998) pulse sequence (TR=2.2 s, TI1=700ms, TI2=1500ms, 20-cm tag width, and a 1-cm gap between the distal end of the tag and the most proximal imaging slice) with a dual-echo gradient echo (GRE) readout²⁹ and spiral acquisition of k-space (TE1=2.7ms, TE2= 29ms, flip angle=90°, field of view (FOV)=22 cm, 64x64 matrix). Twelve slices of 7mm thickness were imaged, with an inter-slice gap of 1mm.

Physiological Monitoring

Throughout scanning, the cardiac pulse was recorded using a finger plethysmograph and a pneumatic belt just below the ribcage was used to measure the respiratory cycle. Expired gas content was monitored continuously via a nasal cannula whereby

end-tidal O₂ and CO₂ data were recorded using a rapidly responding gas analyser (AEI Technologies, PA, USA) to provide representative measures of arterial partial pressures of both gases at the prevailing barometric pressure. Brachial artery blood pressure (BP) was measured at three time-points across the scan session using an MRI-compatible BP cuff (OMRON, Tokyo, Japan).

MRI Data analysis

Physiological Noise Correction. Physiological noise correction was carried out on the raw data using a modified RETROICOR pipeline³⁰. For the raw CBF data, the 1st and 2nd harmonics of the cardiac and respiratory cycles (and the interaction term) were calculated, as well as variance related to end-tidal CO₂, end-tidal O₂, heart rate, and respiration volume per time (RVT; Birn *et al.*, 2009) using a general linear model framework and subsequently regressed from the raw CBF signal. For the MCAC data, only respiratory noise correction was performed.

MCAC Quantification. Arterial compliance measurements were carried out using the methods described by¹⁸ (equation 1). Arterial blood volume (aBV) within the bilateral middle cerebral arteries (MCA) was assessed in systole and diastole. Brachial artery blood pressure cuff recordings were averaged over three time points to calculate average systolic and diastolic BP for each subject. Only data from short TI's (250-850ms) were necessary for deriving aBV to ensure that signal being measured was originating from the arteries rather than the tissue. To determine systole and diastole, the cardiac cycle was divided into 5 phases using the finger plethysmography trace. The short TI images were retrospectively organized into the 5 cardiac phases and an

arterial input function was fitted voxel-wise for each of the 5 phases. The cardiac phases with the maximum and minimum blood volumes, averaged over both MCAs, were used as systole and diastole, respectively.

Equation (1) was used to calculate AC (%/mmHg). Voxel-wise differences between aBV_{Sys} and aBV_{Dia} were calculated, normalised for the aBV in diastole to produce AC values of percentage change in aBV /mmHg (%/mm Hg). Masks of the bilateral MCA were obtained at the level of the M1 segments, branching from the circle of Willis, by thresholding the aBV images ($aBV > 0.1$ % of the voxel) and masking out the anterior and posterior arteries.

$$AC = \frac{aBV_{Sys} - aBV_{Dia}}{aBV_{Dia} * (BP_{Sys} - BP_{Dia})} * 100\% \quad (\text{Equation 1})$$

Grey Matter CBF Quantification. The full MTI time series was used for quantification of resting CBF. Signal within the ventricles ($M_{0\text{CSF}}$) was used to estimate $M_{0,\text{blood}}$ ³² and subsequently modelled to calculate whole-brain perfusion maps based on the entire MTI dataset using FSL BASIL toolbox (FMRIB Software Library, Oxford, UK). Due to the inherently low SNR in ASL imaging, an ROI approach was chosen, *a priori*, in favour of a voxel-wise CBF analysis. Grey matter ROIs were computed by performing whole brain automated segmentation of the T_1 -weighted structural image using the FSL FAST toolbox (Zhang *et al.*, 2001). Segmented grey matter masks were spatially down-sampled into functional space, and binarised to produce an individual grey matter

specific mask for each subject. Whole-brain GM masks were applied to CBF maps to produce a median GM CBF estimate.

CVR Quantification. Simultaneously acquired CBF and BOLD time-series images were corrected for head motion with MCFLIRT³⁴, brain-extracted³⁵ and spatially smoothed with a Gaussian kernel of 6 mm using SUSAN³⁶. BOLD images were calculated from the second echo data using interpolated surround averaging of the tag and control images to yield a BOLD weighted time-series, as described previously³⁷. The first echo data were used to calculate a subtraction time-series³⁸ from which CBF was quantified using the standard single-compartment CBF model²⁶. BOLD and CBF time-series data were converted to percentage change in the signal relative to the baseline (mean) of the time-series to produce a $\% \Delta \text{BOLD}$ and $\% \Delta \text{CBF}$ time-series respectively. Signal was averaged across whole-brain grey matter. A regression analysis was performed to measure $\% \Delta \text{BOLD}$ and $\% \Delta \text{CBF}$ per mmHg change in absolute end-tidal CO_2 with a 3rd order polynomial included to remove slow signal drift. Temporal lag-fitting (time-shift steps of 0.1s) was also carried out, to account for the delay between end-tidal CO_2 increase in response to breath-holding and the subsequent blood flow response²⁷. CVR was thus defined as the beta-weight from the regression model, where BOLD and CBF were measured in units of $\% \text{BOLD}/\text{mmHg}$ or $\% \text{CBF}/\text{mmHg}$ respectively.

Statistical Analysis

Pearson's correlation was used to assess the relationship between cardiorespiratory fitness ($\dot{V}O_{2\text{MAX}}$) and physiological measures (Table 1). Linear regression was used to assess the predictive effect of cardiorespiratory fitness on MRI metrics across whole

$\dot{V}O_{2\text{MAX}}$ predicts MCA compliance and baseline CBF

HV Furby *et al.*

brain grey matter and their relationship with heart rate and Hct. Pearson correlation coefficients were used to assess the relationships between MCAC, CVR and CBF. Bootstrapped confidence intervals (95%) were computed unless otherwise stated, whereby analyses were bootstrapped to 1000 samples. Bias corrected and accelerated confidence intervals (CIs) are reported. Analysis was performed in SPSS version 20 (IBM).

Results

Physiological Measures

$\dot{V}O_{2MAX}$ ranged from 38 to 76ml/min/kg (57.3 ± 12.7 ml/kg/min). Body mass was 76.6 ± 8.3 kg, systolic BP was 122 ± 4 mm/Hg and diastolic BP was 72 ± 9 mm/Hg. $\dot{V}O_{2MAX}$ was positively associated with time-to-exhaustion ($R^2 = 0.59$, $F(1, 9) = 12.74$, $p < 0.01$, 95% CI [3.02, 13.49]) and a visual inspection suggested greater endurance (time to exhaustion) in those recruited from cycling and running clubs compared to other participants (figure 1). $\dot{V}O_{2MAX}$ was not associated with any other physiological metric (Table 1).

Resting heart rate was not predictive of MCAC ($R^2 = 1.00$, $F(1, 7) = 0.75$, $\beta = -0.006$, $p = 0.42$, 95% CI [-0.02, 0.01]), global GM CBF ($R^2 = 0.001$, $F(1, 6) = 0.006$, $\beta = 0.23$, $p = 0.94$, 95% CI [-0.70, 0.74]) or CVR measures (BOLD $R^2 = 0.01$, $F(1, 8) = 0.10$, $p = 0.76$, 95% CI [-0.001, 0.009]); CBF $R^2 = 0.06$, $F(1, 8) = 0.53$, $p = 0.49$, 95% CI [-0.18, 0.09]). Hct was not correlated with any of the MRI measures (MCAC ($r(7) = -0.15$, $p = 0.71$); global GM CBF ($r(6) = 0.07$, $p = 0.87$); CBF CVR ($r(8) = 0.41$, $p = 0.24$); BOLD CVR ($r(8) = 0.35$, $p = 0.33$)).

MCAC

Nine participants contributed to the MCAC analysis as two were removed due to severe head movement, observed during visual inspection of MR images, that could not be rectified by volume removal. Averaged over all participants, we calculated bilateral MCAC to be 0.41 ± 0.16 %/mmHg. Linear regression revealed a significant

relationship between $\dot{V}O_{2MAX}$ and MCAC, whereby fitter individuals showed the hypothesised greater compliance of the middle cerebral arteries (figure 2a) and this relationship was significant ($R^2 = 0.64$, $\beta=0.01$, $F(1,8) = 12.6$, $p=0.009$, 95% CI [0.003, 0.018]). These values indicate that arterial compliance in the MCA increased by 0.01%/mmHg for each ml/min/kg increase in $\dot{V}O_{2MAX}$.

Retrospective synchronisation of images across the cardiac cycle was inspected to ensure that there was not a bias between the number of tag and control images for a particular TI or cardiac phase. A repeated-measures ANOVA revealed that the number of tag and control images did not differ significantly between TI ($F(6,48) = 1.3$, $p=0.30$, n.s.) or cardiac phase ($F(1,8) = 0.7$, $p=0.43$, n.s.), nor was there an interaction between the number of images within each cardiac phase at each TI ($F(6,48) = 0.54$, $p=0.78$, n.s.). On average, for a single TI there were 6 tag and 7 control images in diastole, and 6 tag and 6 control images in systole.

Grey Matter CBF

Whole-brain GM averaged CBF values ranged from 53.8 to 73.1 ml/100g/min (59.4 ± 6.7). Eight participants contributed to baseline CBF analysis (3 were excluded due to severe head motion). Linear regression revealed a significant inverse relationship between $\dot{V}O_{2MAX}$ and resting whole-brain GM CBF ($R^2 = 0.75$, $\beta=-0.47$, $F(1,7) = 18.3$, $p= 0.005$, 95% CI [-0.73, -0.20]; figure 2b). Whole-brain grey matter CBF decreased 0.47 ml/100g/min for each ml/min/kg increase in $\dot{V}O_{2MAX}$. An inverse relationship was

observed within each cortical ROI, however, none of these was significant (p -values > 0.05).

CVR

Cerebrovascular reactivity data was excluded for one participant because the subject was unable to breathe through his nose, so that 10 subjects contributed to the CVR analysis. BOLD data demonstrated better signal-to-noise (SNR) than CBF measurements in response to breath-holding (see figure 3a). However, CBF CVR was positively correlated with the BOLD CVR measurements across whole-brain GM ($R^2 = 0.52$, $F(1,9) = 8.8$, $p = 0.01$, 95% CI [0.034, 0.078; see figure 3b]. Both measurements showed a decline in CVR with increasing $\dot{V}O_{2MAX}$ within whole-brain grey matter (figure 3c). However, linear regression did not find this to be significant for either the BOLD ($R^2 = 0.47$, $\beta = -0.004$, $F(1,9) = 2.27$, $p = 0.17$, 95% CI [-0.009, 0.002] or CBF ($R^2 = 0.19$, $\beta = -0.05$, $F(1,9) = 1.88$, $p = 0.21$, 95% CI [-0.15, -0.04] CVR).

Relationship between MCAC, whole-brain CBF and CVR

MCAC was not correlated with either BOLD CVR ($r(7) = -0.20$, $p = 0.68$, 95% CI [-0.89, 0.44]), CBF CVR ($r(7) = -0.14$, $p = 0.78$, 95% CI [-0.54, 0.39]), nor with resting grey matter perfusion ($r(7) = -0.45$, $p = 0.30$, 95% CI [-0.73, -0.33]). There was, however, a significant correlation between resting grey matter perfusion with both CBF CVR ($r(7) = 0.76$, $p < 0.05$, 95% CI [0.04, 0.95] and BOLD CVR ($r(7) = 0.78$, $p < 0.05$, 95% CI [-0.35, 0.97]).

Discussion

Whilst the benefits of physical activity on cognition and mental health are well recognised, the physiological mechanisms by which exercise exerts its beneficial effects on the brain remain poorly understood. In this study, we demonstrate that ASL MRI is a useful tool for understanding how fitness may influence vascular function, in particular, cerebral arterial compliance.

MCAC

The present study utilised a novel, non-invasive, measure of MCAC based on pASL MRI to demonstrate the link between cardiorespiratory fitness and cerebral arterial compliance in a sample of young males spanning a range of VO_{2MAX} values from fair (36.5-42.2 ml/kg/min) to higher (52.4 ml/kg/min) for adults aged 20-29 years³⁹. We showed that MCAC was higher in individuals with higher $\dot{V}O_{2MAX}$, a finding consistent with non-MRI methods in other major arteries throughout the body^{10,11,13}. The present study is the first to use MRI to measure fitness related changes in MCAC and provides promising evidence towards the cerebrovascular benefits of physical activity, as well as insight into the potential mechanisms at play.

It has been suggested that the ability of cerebral arteries to dampen changes in pulse pressure may prevent downstream tissue damage where vessels are vulnerable to deterioration⁸. Higher MCAC, as measured here, can be thought to reflect healthy, more 'elastic' vessel walls than those with lower MCAC, a possible marker of better cerebrovascular health in those with high cardiorespiratory capacity. Damage to the microvasculature has been associated with poorer memory, processing

speed and executive function^{13,40,41}. ‘Training’ the vessel, through increasing AC could give rise to some of the cognitive benefits that have been reported as a result of exercise, by preventing age-related arterial stiffening and reducing a down-stream deleterious effect of pulsatile flow on the microvasculature within the tissue bed.

Our MRI results corroborate indirect evidence from the ultrasound literature that shows an increase in extracranial compliance with cardiorespiratory fitness⁴². Validation of this link using our ASL methods lends support for future interventional exercise studies, where the mechanisms underpinning MCAC can be explored in different ages, and with different types and modes of exercise. Resistance training has been found previously to reduce AC, or have no effect on, carotid artery compliance whereas aerobic training leads to increased AC⁴³. Similarly, high intensity interval training (HIIT) differs from continuous moderate intensity exercise on measures of arterial stiffness⁴⁴⁻⁴⁶. It has been proposed that a moderate or higher load of training may be required to influence endothelial function in healthy people⁴⁷ where repeated shear stress stimulation is required to drive adaptation⁴⁸ and arterial remodelling of endothelial and vascular smooth muscle cells that are located within the medial layer of the arterial wall⁴⁹ and regulate vascular function⁶. Although our participants were recruited from cycling and running clubs to ensure a broad range of cardiorespiratory fitness, the volume, intensity duration and mode of training was not controlled for. Further research into the effects of specific types of exercise on AC in the brain using this novel MRI method is warranted to elucidate these potentially variable effects.

Our cross-sectional design explored $\dot{V}O_{2MAX}$ as a surrogate measure of physical fitness, but with this method, we are unable to elucidate the temporal dynamics of

arterial remodelling or its causal linkage to $\dot{V}O_{2MAX}$. $\dot{V}O_{2MAX}$ can decrease surprisingly quickly in the absence of any aerobic training and it would be of interest to measure the immediate effects of detraining on MCAC in a longitudinal design. A previous study using aortic pulse wave velocity (PWV) as a measure of arterial distensibility, showed increased arterial distensibility after 8 weeks of cycling, that returned to baseline after just 4 weeks of detraining⁵⁰.

Average MCA compliance across participants in the present study was 0.41% \pm 0.16% per mmHg which is consistent, albeit slightly lower than reported previously in a sample of 5 participants (right MCAC = 0.57% \pm 0.20%; left MCAC = 0.50% \pm 0.30% per mmHg)¹⁸. The current findings demonstrate variation in cerebral AC in the MCA, however using MRI it is also possible to investigate the posterior and anterior cerebral arteries^{15,16,18}. Unfortunately, due to the scan duration and sample size used in this study, SNR was too low to assess compliance in these smaller arteries.

Grey Matter CBF

This is the first study to assess resting cerebral blood flow using ASL in a cohort of young adults in the moderate-to-high fitness range. We report a reduction in resting CBF with increased fitness levels, a finding which contrasts with a handful of studies from the ultrasound literature, whereby fitness has been positively associated with cerebral blood velocity^{2,3,20} and flow in children¹⁹, older adults²⁰ and patients with coronary artery disease²¹.

Across adulthood, age decreases cerebral metabolic rates of oxygen (CMRO₂) and glucose by \sim 5% per decade, and reduced metabolic rate is coupled with lower CBF^{51,52}. It has been proposed that exercise could ameliorate age-related cognitive

decline^{53,54} by enhancing vasodilatory signalling via nitrous oxide synthase activity, promoting endothelial repair mechanisms and angiogenesis to effectively meet the demands of the metabolising cerebral tissue^{55,56}. It is widely assumed, but less well proven, that these mechanisms lead to a net increase in resting CBF in the healthy adult brain following exercise. In this study, we find that CBF is *lower* in young males with higher cardiorespiratory fitness.

Interpretation should be made cautiously given the modest size of the present study, however there are a number of possible mechanisms that could drive the negative association between cardiorespiratory fitness and CBF. These include reduction of arteriolar luminal diameter, changes in capillary density and an alteration of tissue oxygen extraction. The former seems unlikely, since exercise has been shown to decrease the intima media thickness (IMT) of the arterial wall, thereby increasing lumen diameter and allowing for an increase in blood flow through the artery (Sandrock et al., 2008). However, an increase in lumen diameter is not a consistent observation in young adults (Popovic et al., 2011) and has not been explored in the cerebral arteries, making it worthy of further investigation. It is unlikely that lower CBF in fitter subjects is due to a reduction of capillary density, since a number of preclinical studies have provided evidence of increased vessel density in the rodent brain following exercise⁵⁷⁻⁵⁹. It is possible that such an increase in vessel surface area with increased capillary number could *reduce* the demand for CBF⁵⁹, where shorter diffusion distances mean nutrient extraction is facilitated. This raises the possibility that fitter individuals have more efficient gas exchange from the capillary bed, permitting a reduction in the amount of flow needed to meet metabolic oxygen demand.

It has been shown that a reduction in CBF seen *during* exercise was accompanied by an increase in oxygen extraction, resulting in a maintained cerebral metabolic rate of oxygen consumption ($CMRO_2$)⁶⁰. Future research could use calibrated fMRI measures of oxygen extraction and $CMRO_2$ ^{61–64} in highly fit individuals, to assess whether efficiency of nutrient supply via the cerebral microvasculature can explain the inverse relationship between $\dot{V}O_{2MAX}$ and CBF. Hct contributes to an individual's O_2 carrying capacity, and alongside CBF and the arterial oxyhaemoglobin saturation, dictates cerebral oxygen delivery⁶⁵. We explored the relationship between Hct and CBF and, Hct and $\dot{V}O_{2MAX}$, but neither was significant in this sample.

Cerebrovascular Reactivity

To date, studies that have investigated the relationship between CVR and fitness have relied upon either ultrasound methods^{2,3} or BOLD measurements^{22,66} which have found opposing results. Since the BOLD signal does not represent blood *flow*, BOLD CVR alone is not sufficient for understanding the mechanisms at play⁶⁷. The current study used pulsed ASL methods that allowed simultaneous measurement of BOLD and CBF, to assess whether differences in BOLD previously reported are likely to be due to a change in blood flow. In line with the BOLD MRI literature, whole-brain grey matter CVR showed an inverse trend with $\dot{V}O_{2MAX}$, for both BOLD and CBF metrics, although neither effect was statistically significant.

Within the healthy brain, an increase in arterial CO_2 is expected to produce a rapid vasodilatory response, yielding an elevation in CBF. This vascular reactivity is thought to be an adaptive physiological response, such that a decline in reactivity

could be considered maladaptive. Nonetheless this, and previous studies, have found a negative trend whereby CVR is lower in fitter subjects. For example, BOLD CVR was found to decrease in a study of elderly masters athletes with increased $\dot{V}O_{2MAX}$ in response to a 5% CO₂ hypercapnic challenge²². A separate study found reductions in frontal BOLD CVR with increased $\dot{V}O_{2MAX}$ despite fitter subjects performing better at a frontal executive cognitive task⁶⁶. Together with our results, it seems that the link between cardiorespiratory fitness and cerebrovascular health may be more complex than previously suggested. One proposed explanation is that chronic elevations in venous CO₂ during prolonged periods of exercise over years of training may lead to desensitisation of the vasodilatory mechanisms such as the bioavailability of nitric oxide, that mediate the reactivity of the blood vessels and regulate blood flow⁴⁷. This same mechanism may also explain the negative relationship between fitness and resting CBF observed in our study.

Hct levels are associated with variation in task-based BOLD estimates⁶⁵. We did not observe a significant relationship between Hct levels and BOLD-based or CBF-based CVR in this study. However, we exercise caution when interpreting MRI measures in light of Hct since, blood and MRI measures were acquired on separate days.

It is possible that the breath-hold paradigm used here may not have been sensitive enough to detect a significant difference in CVR in this sample. Targeted gas challenges tend to provide a more robust measure of CVR, as CO₂ is directly manipulated²⁸ and comparable levels of hypercapnia can be achieved between subjects. However, breath-hold offers greater experimental convenience. It has been

previously shown that breath-holds are a reliable measure of BOLD CVR, even when breath-holding is poor²⁷. SNR is inherently lower in CBF than BOLD data. Nevertheless, we observed a significant relationship between BOLD and CBF CVR measures.

Unlike the multi inversion time arterial spin labelling scheme used for estimating baseline CBF, an inherent limitation of the PASL single inversion time approach used for measuring CVR is that it assumes all the labelled blood has flowed into the imaging slice. This acquisition scheme was chosen for time efficiency and because we were interested in the dual-echo (i.e. CBF and BOLD) readout. However, it is possible that bias in CVR estimates may be introduced where differing amounts of the labelled bolus arrive in the imaged slice during normo- and hyper-capnia.

Limitations

Care should be taken when generalizing these findings since the cohort used here was small. Our study design specifically recruited those across a range of moderate-high $\dot{V}O_{2MAX}$ to exacerbate any association with vascular MRI parameters. Due to the correlational design of this study, cause and effect cannot easily be determined and a randomized clinical trial (RCT) involving a specified mode, intensity, frequency and duration of exercise would address this issue.

An inherent limitation of using $\dot{V}O_{2MAX}$ testing, is that performance may be biased by mode of exercise (e.g. treadmill vs. cycle ergometer). We did not control for the amount of cycle training engaged in by each of our participants prior to testing, however, those recruited from cycling clubs did not differ convincingly from those recruited from running clubs. Future studies should take this bias into consideration.

We did not address potential genetic and other environmental factors that could mediate the relationship between fitness and vascular health. However, emerging evidence suggests that the process of arterial stiffening may have a genetic component⁶⁸ that may be relevant when looking at individual differences in response to exercise.

Conclusions and Future Research

In conclusion, $\dot{V}O_{2MAX}$ was found to be associated with several cerebrovascular parameters, including an elevation in MCAC and a decline in resting CBF. This is the first time an association has been reported between cardiorespiratory fitness and AC *within* the brain using this novel MRI technique¹⁸. The relationship between fitness and MCA compliance in this group of healthy young males, provides promising clues towards the influence of exercise on cerebrovascular health early in life, before cognitive decline becomes evident, and sheds light on the possible mechanisms by which exercise impacts cerebrovascular health.

ACKNOWLEDGEMENTS

The authors would like to thank Dr Alan Stone and Peter Hobden for their imaging assistance and Dr Catherine Foster for her contribution to the conceptual development.

AUTHOR CONTRIBUTION STATEMENT

HF performed recruitment, MRI imaging, analysis and write up. HF, CM, DB and RW were involved in study concept and design. EW and HF played a major role in MRI data

acquisition, data processing; data analysis and interpretation. CM and DB were involved in subject recruitment and responsible for acquisition of VO_{2MAX} , blood and anthropomorphic data. All authors contributed to interpretation and critical revision of manuscript.

DISCLOSURE/CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

1. Nocon M, Hiemann T, Müller-Riemenschneider F, et al. Association of physical activity with all-cause and cardiovascular mortality: a systematic review and meta-analysis. *Eur J Cardiovasc Prev Rehabil*; 15: 239–246.
2. Ainslie PN, Cotter JD, George KP, et al. Elevation in cerebral blood flow velocity with aerobic fitness throughout healthy human ageing. *J Physiol*; 586: 4005–10.
3. Bailey DM, Marley CJ, Brugniaux J V, et al. Elevated aerobic fitness sustained throughout the adult lifespan is associated with improved cerebral hemodynamics. *Stroke*; 44: 3235–8.
4. Willie CK, Colino FL, Bailey DM, et al. Utility of transcranial Doppler ultrasound for the integrative assessment of cerebrovascular function. *J Neurosci Methods*; 196: 221–37.
5. Tymko MM, Ainslie PN, Smith KJ. Evaluating the methods used for measuring cerebral blood flow at rest and during exercise in humans. *Eur J Appl Physiol*; 0: 1–12.
6. Peterson EC, Wang Z, Britz G. Regulation of cerebral blood flow. *Int J Vasc Med* 2011; 2011.
7. Lacolley P, Regnault V, Segers P, et al. Vascular Smooth Muscle Cells and Arterial Stiffening: Relevance in Development, Aging, and Disease. *Physiol Rev*; 97: 1555–1617.
8. Poels MMF, Zaccari K, Verwoert GC, et al. Arterial stiffness and cerebral small vessel disease: The rotterdam scan study. *Stroke*; 43: 2637–2642.
9. Pase MP, Himali JJ, Mitchell GF, et al. Association of Aortic Stiffness With Cognition and Brain Aging in Young and Middle-Aged Adults. *Hypertension*; 67: 513–519.
10. Tanaka H, Dinunno F a, Monahan KD, et al. Aging, habitual exercise, and dynamic arterial compliance. *Circulation*; 102: 1270–1275.
11. Tabara Y, Yuasa T, Oshiumi a, et al. Effect of acute and long-term aerobic exercise on arterial stiffness in the elderly. *HypertensRes*; 30: 895–902.

12. Carrera E, Kim DJ, Castellani G, et al. Effect of Hyper- and Hypocapnia on Cerebral Arterial Compliance in Normal Subjects. *J Neuroimaging*; 21: 121–125.
13. Tan CH, Low KA, Kong T, et al. Mapping cerebral pulse pressure and arterial compliance over the adult lifespan with optical imaging. *PLoS One*; 12: 1–23.
14. Fabiani M, Low KA, Tan C, et al. Taking the pulse of aging : Mapping pulse pressure and elasticity in cerebral arteries with optical methods. 51: 1072–1088.
15. Warnert EAH, Hart EC, Hall JE, et al. The major cerebral arteries proximal to the Circle of Willis contribute to cerebrovascular resistance in humans. *J Cereb blood flow Metab*; 1–12.
16. Warnert EAH, Hart EC, Hall JE, et al. The major cerebral arteries proximal to the Circle of Willis contribute to cerebrovascular resistance in humans. *J Cereb Blood Flow Metab*; 0271678X15617952.
17. Edwards NM, Daniels SR, Claytor RP, et al. Physical activity is independently associated with multiple measures of arterial stiffness in adolescents and young adults. *Metabolism*; 61: 869–872.
18. Warnert EA, Murphy K, Hall JE, et al. Noninvasive assessment of arterial compliance of human cerebral arteries with short inversion time arterial spin labeling. *J Cereb Blood Flow Metab*; 35: 461–468.
19. Chaddock-Heyman L, Erickson KI, Chappell MA, et al. Developmental Cognitive Neuroscience Aerobic fitness is associated with greater hippocampal cerebral blood flow in children. *Dev Cogn Neurosci*; 20: 52–58.
20. Zimmerman B, Sutton BP, Low K a., et al. Cardiorespiratory fitness mediates the effects of aging on cerebral blood flow. *Front Aging Neurosci*; 6: 1–13.
21. MacIntosh BJ, Swardfager W, Crane DE, et al. Cardiopulmonary fitness correlates with regional cerebral grey matter perfusion and density in men with coronary artery disease. *PLoS One*; 9: e91251.
22. Thomas B, Yezhuvath U, Tseng B, et al. Life-long aerobic exercise preserved baseline cerebral blood flow but reduced vascular reactivity to CO₂. *J Magn Reson Imaging*; 38: 1177–83.
23. Nevo O, Soustiel JF, Thaler I. Cerebral blood flow is increased during controlled ovarian stimulation. *Am J Physiol Heart Circ Physiol*; 293: H3265–H3269.
24. Ross R, Blair SN, Arena R, et al. *Importance of Assessing Cardiorespiratory Fitness in Clinical Practice : A Case for Fitness as a Clinical*. 2016.
25. Barnes JN, Taylor JL, Kluck BN, et al. Cerebrovascular reactivity is associated with maximal aerobic capacity in healthy older adults. *J Appl Physiol*; 114: 1383–7.
26. Wong EC, Buxton RB, Frank LR. Quantitative imaging of perfusion using a single subtraction (QUIPSS and QUIPSS II). *Magn Reson Med*; 39: 702–8.
27. Bright MG, Murphy K. Reliable quantification of BOLD fMRI cerebrovascular reactivity despite poor breath-hold performance. *Neuroimage*; 83: 559–68.
28. Tancredi FB, Hoge RD. Comparison of cerebral vascular reactivity measures obtained using breath-holding and CO₂ inhalation. *J Cereb Blood Flow Metab*; 33: 1066–74.
29. Liu JZ, Dai TH, Sahgal V, et al. Nonlinear cortical modulation of muscle fatigue:

- a functional MRI study. *Brain Res*; 957: 320–9.
30. Glover GH, Li TQ, Ress D. Image-based method for retrospective correction of physiological motion effects in fMRI: RETROICOR. *Magn Reson Med*; 44: 162–7.
 31. Birn RM, Murphy K, Handwerker D a, et al. fMRI in the presence of task-correlated breathing variations. *Neuroimage*; 47: 1092–104.
 32. Lu H, Clingman C, Golay X, et al. Determining the longitudinal relaxation time (T1) of blood at 3.0 Tesla. *Magn Reson Med*; 52: 679–82.
 33. Zhang Y, Brady M, Smith S. Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE Trans Med Imag*; 20: 45–57.
 34. Jenkinson M, Bannister P, Brady M, et al. Improved Optimization for the Robust and Accurate Linear Registration and Motion Correction of Brain Images. *Neuroimage*; 17: 825–841.
 35. Smith SM. Fast robust automated brain extraction. *Hum Brain Mapp*; 17: 143–55.
 36. Smith SM, Brady J. SUSAN - a new approach to low level image processing. 1997.
 37. Liu TT, Wong EC. A signal processing model for arterial spin labeling functional MRI. *Neuroimage*; 24: 207–15.
 38. Murphy K, Harris AD, Diukova A, et al. Pulsed arterial spin labeling perfusion imaging at 3 T: estimating the number of subjects required in common designs of clinical trials. *Magn Reson Imaging*; 29: 1382–9.
 39. Heyward VH. *Advance Fitness Assessment & Exercise Prescription*. 3rd ed. Dallas TX, 1998.
 40. Davenport MH, Hogan DB, Eskes G a, et al. Cerebrovascular reserve: the link between fitness and cognitive function? *Exerc Sport Sci Rev*; 40: 153–8.
 41. Mitchell GF, Van Buchem M a., Sigurdsson S, et al. Arterial stiffness, pressure and flow pulsatility and brain structure and function: The Age, Gene/Environment Susceptibility-Reykjavik Study. *Brain*; 134: 3398–3407.
 42. Tanaka H, DeSouza C a, Seals DR. Absence of age-related increase in central arterial stiffness in physically active women. *Arterioscler Thromb Vasc Biol*; 18: 127–132.
 43. Miyachi M, Donato AJ, Yamamoto K, et al. Greater age-related reductions in central arterial compliance in resistance-trained men. *Hypertension*; 41: 130–135.
 44. Kim H, Hwang C, Yoo J, et al. All-Extremity Exercise Training Improves Arterial Stiffness in Older Adults. *Med Sci Sports Exerc*; 49: 1404–1411.
 45. Guimarães GV, Ciolac EG, Carvalho VO, et al. Effects of continuous vs. interval exercise training on blood pressure and arterial stiffness in treated hypertension. *Hypertens Res*; 33: 627–632.
 46. Ciolac EG, Bocchi E a, Bortolotto L a, et al. Effects of high-intensity aerobic interval training vs. moderate exercise on hemodynamic, metabolic and neuro-humoral abnormalities of young normotensive women at high familial risk for hypertension. *Hypertens Res*; 33: 836–843.
 47. Green DJ, Maiorana A, O'Driscoll G, et al. Effect of exercise training on endothelium-derived nitric oxide function in humans. *J Physiol*; 561: 1–25.

48. Green DJ, Bilsborough W, Naylor LH, et al. Comparison of forearm blood flow responses to incremental handgrip and cycle ergometer exercise: relative contribution of nitric oxide. *J Physiol*; 562: 617–628.
49. Newcomer SC, Thijssen DHJ, Green DJ. Effects of exercise on endothelium and endothelium/smooth muscle cross talk: role of exercise-induced hemodynamics. *J Appl Physiol*; 111: 311–320.
50. Kakiyama T, Sugawara JUN, Murakami H, et al. Effects of Short-Term Endurance Training on Aortic Distensibility in Young Males. *Med Sci Sports Exerc*; 37: 267–271.
51. Leenders KL, Perani D, Lammertsma AA, et al. CEREBRAL BLOOD FLOW , BLOOD VOLUME AND OXYGEN UTILIZATION NORMAL VALUES AND EFFECT OF AGE. *Brain*; 113: 27–47.
52. Petit-Taboue MC, Landeau B, Desson JF, et al. Effects of healthy aging on regional cerebral metabolic rates of glucose: Assessment with positron emission tomography and statistical parametric mapping. *Ann Neurol*; 40: M48–M48.
53. Tarumi T, Gonzales MM, Fallow B, et al. Central artery stiffness, neuropsychological function, and cerebral perfusion in sedentary and endurance-trained middle-aged adults. *J Hypertens*; 31: 2400–9.
54. Smith AM, Spiegler KM, Sauce B, et al. Voluntary aerobic exercise increases the cognitive enhancing effects of working memory training. *Behav Brain Res*; 256: 626–35.
55. Gertz K, Priller J, Kronenberg G, et al. Physical activity improves long-term stroke outcome via endothelial nitric oxide synthase-dependent augmentation of neovascularization and cerebral blood flow. *Circ Res*; 99: 1132–40.
56. Bolduc V, Thorin-Trescases N, Thorin E. Endothelium-dependent control of cerebrovascular functions through age: exercise for healthy cerebrovascular aging. *AJP Hear Circ Physiol*; 305: H620–H633.
57. Swain R., Harris a. ., Wiener E., et al. Prolonged exercise induces angiogenesis and increases cerebral blood volume in primary motor cortex of the rat. *Neuroscience*; 117: 1037–1046.
58. van Praag H, Shubert T, Zhao C, et al. Exercise enhances learning and hippocampal neurogenesis in aged mice. *J Neurosci*; 25: 8680–5.
59. Al-Jarrah M, Jamous M, Al Zailaey K, et al. Endurance exercise training promotes angiogenesis in the brain of chronic/progressive mouse model of parkinson's disease. *NeuroRehabilitation*; 26: 369–373.
60. Trangmar SJ, Chiesa ST, Stock CG, et al. Dehydration affects cerebral blood flow but not its metabolic rate for oxygen during maximal exercise in trained humans. *J Physiol*; 592: 3143–60.
61. Wise RG, Harris AD, Stone AJ, et al. Measurement of OEF and absolute CMRO₂: MRI-based methods using interleaved and combined hypercapnia and hyperoxia. *Neuroimage*; 83: 135–47.
62. Merola A, Murphy K, Stone AJ, et al. Measurement of oxygen extraction fraction (OEF): an optimised BOLD signal model for use with hypercapnic and hyperoxic calibration. *Neuroimage*; 5–6.
63. Germuska M, Merola A, Murphy K, et al. A new framework for dual calibrated

- estimation of OEF R2v1. 2016.
64. Merola A, Germuska MA, Murphy K, et al. Assessing the repeatability of absolute CMRO₂, OEF and haemodynamic measurements from calibrated fMRI. *Neuroimage*; 173: 113–126.
 65. Xu F, Li W, Liu P, et al. Accounting for the Role of Hematocrit in Between-Subject Variations of MRI-Derived Baseline Cerebral Hemodynamic Parameters and Functional BOLD Responses. *Hum Brain Mapp*; 39: 344–353.
 66. Gauthier CJ, Lefort M, Mekary S, et al. Hearts and minds: linking vascular rigidity and aerobic fitness with cognitive aging. *Neurobiol Aging*; 36: 304–314.
 67. Buxton RB, Uludağ K, Dubowitz DJ, et al. Modeling the hemodynamic response to brain activation. *Neuroimage*; 23 Suppl 1: S220-33.
 68. Lacolley P, Challande P, Osborne-Pellegrin M, et al. Genetics and pathophysiology of arterial stiffness. *Cardiovasc Res*; 81: 637–648.

FIGURES

Figure 1. Those recruited from cycling (circles) and running (triangles) clubs appeared to have higher VO_{2MAX} and longer time to exhaustion than community controls.




Figure 2. Increased VO_{2MAX} is associated with (a) increased arterial compliance within the bilateral middle cerebral arteries (MCA)($p < 0.01$) (b) decreased GM CBF at rest ($p < 0.01$) (c) decreased GM CBF CVR ($p = 0.21$, n.s.).




Figure 3. (a) BOLD and (b) CBF responses to the breath-hold task within a grey matter mask. Coloured lines represent individual subject data; black lines reflect the average response across participants. BOLD time-series showed better signal-to-noise than CBF. (c) BOLD CVR ($\% \Delta \text{BOLD} / \text{mmHg PET CO}_2$) and CBF CVR ($\% \Delta \text{CBF} / \text{mmHg PET CO}_2$) were significantly correlated.




Table 1. Anthropomorphic measures for all subjects.

Subject	VO_{2MAX} (ml/kg/min)	Age (years)	Height (cm)	Weight (Kg)	BMI (kg/m ²)	Total Body Fat (%)	Haematocrit (%)	Systolic Blood Pressure (mmHg)	Diastolic Blood Pressure (mmHg)	Resting Heart Rate (bpm)	Time to exhaustion (secs)
1	56.3	23	1.87	85.9	25	10.7	48	128	70	62	810
2	75.8	20	1.76	62.7	20	12.5	46	126	84	65	690
3	58.5	21	1.76	65.5	21	10.4	46	116	60	71	680
4	64	19	1.84	74.7	22	8.8	50	118	74	54	830
5	63.1	22	1.82	77.1	23	9.1	45	120	74	55	830
6	67.3	23	1.84	75.6	22	10.6	44	124	72	60	810
7	61.5	21	1.79	69.5	22	14.9	48	124	50	54	650
8	42.5	20	1.88	67.6	19	10	45	124	78	77	500
9	40.8	19	1.67	64.3	23	18.6	42	126	76	79	480
10	37.8	20	1.75	85.6	28.0	18.8	48	122	76	54	530
11	45.2	19	1.76	84.6	27.3	16.2	48	118	78	72	620
Mean (±SD)	56 (12)	21 (2)	1.79 (0.06)	74 (9)	23 (3)	13 (4)	46(2)	122 (4)	72 (9)	64 (10)	675 (133)
Correlation (Pearson's r)		0.13	0.29	-0.29	-0.49	-0.61	0.11	0.02	-0.14	-0.44	-0.77
p-value (α=0.05)		n.s	n.s	n.s	n.s	n.s	n.s	n.s	n.s	n.s	<0.01