

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/125181/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Brydges, Nichola M, Best, Caroline and Thomas, Kerrie ORCID:
<https://orcid.org/0000-0003-3355-9583> 2020. Female HPA axis displays heightened sensitivity to pre-pubertal stress. *Stress* 23 (2) , pp. 190-200.
10.1080/10253890.2019.1658738 file

Publishers page: <http://dx.doi.org/10.1080/10253890.2019.1658738>
< <http://dx.doi.org/10.1080/10253890.2019.1658738> >

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1 **Female HPA axis displays heightened sensitivity to pre-pubertal stress**

2 **Running title: Pre-pubertal stress and adult HPA axis**

3 Nichola M. Brydges^{1*}, Caroline Best¹ & Kerrie Thomas^{1,2}.

- 4
- 5 1. Neuroscience and Mental Health Research Institute, Cardiff University, Hadyn Ellis Building,
6 Maindy Road, Cardiff, CF24 4HQ, UK. 2. School of Biosciences, Cardiff University, Museum
7 Avenue, Cardiff, CF10 3AX, UK.
- 8
- 9
- 10
- 11
- 12
- 13

14 *Corresponding author. Tel: +44 (0)29 208 8339. E-mail: brydgesn@cardiff.ac.uk.

15

16

17

18

19

20

21

22 **Abstract**

23 Early life stress (ELS) is a risk factor in the development of psychiatric disorders. The underlying
24 biological mechanisms governing this phenomenon are not fully understood, but dysregulation of
25 stress responses is likely to play a key role. Males and females differ in their propensity to develop
26 psychiatric disorders, with far higher rates of anxiety, major depressive disorder, affective disorders
27 and post-traumatic stress disorder found in women. We hypothesised that sex differences in response
28 to ELS may play a crucial role in differential vulnerability between the sexes. To test this, we evaluated
29 the consequences of pre-pubertal stress (PPS) on the HPA axis in adult female and male Lister Hooded
30 rats. PPS animals were exposed to swim, restraint and elevated platform stress on postnatal days 25-
31 27, controls remained in their home cage. Once adult, animals were either a) sacrificed directly and
32 brains collected or b) sacrificed 20 minutes or 1 week after a social test and trunk blood collected. In
33 the female hippocampal formation, PPS increased expression of *FKBP5* and *AVPR1a*. In the female
34 prefrontal cortex, PPS resulted in increased glucocorticoid receptor expression, increased
35 glucocorticoid:mineralocorticoid (*GR:MR*) receptor expression ratio and decreased *AVPR1a*
36 expression. Females exposed to PPS did not show the normal rise in blood corticosterone levels
37 following a social interaction test. In contrast, PPS did not alter the expression of oxytocin or oxytocin
38 receptors, and no effects of PPS were seen in males. However, striking sex differences were found.
39 Females had higher oxytocin receptor expression in the prefrontal cortex and *AVPR1a* and oxytocin
40 expression in the hypothalamus, whereas males demonstrated higher expression of *GR*, *MR*, *GR:MR*,
41 *FKBP5* and oxytocin receptor in the hypothalamus. These results demonstrate heightened reactivity
42 of the female HPA axis to PPS and may help explain why in humans females display an increased
43 susceptibility to certain stress-related psychopathologies.

44

45

46

47 Lay Summary

48 Women are at greater risk of developing several psychiatric illnesses. Using a rodent model, we show
49 that the female stress system is more reactive to the lasting effects of early life stress. This heightened
50 reactivity of the female stress response may help explain why women are at a greater risk of
51 developing psychiatric disorders.

52 **Keywords:** pre-pubertal stress, HPA axis, sex differences, GR, MR, FKBP5

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68 **Introduction**

69 Adverse experiences early in life are linked with an increased risk of developing psychiatric disorders
70 later in life(Heim & Nemeroff, 2001; Juruena, Baes, Menezes, & Graeff, 2015; Teicher & Samson, 2016;
71 Teicher, Samson, Anderson, & Ohashi, 2016). Dysregulation of the stress response is a potential
72 mechanism through which early life stress (ELS) increases vulnerability to illness. Prolonged or
73 excessive stress may lead to a maladaptive stress response, and when experienced early in life could
74 also alter brain development, increasing vulnerability to psychiatric disorders. Stress results in several
75 adaptive physiological and behavioural responses, a major mediator of this is the hypothalamic-
76 pituitary-adrenal (HPA) axis.

77 Both psychological and physical stressors result in the release of corticotrophin releasing
78 hormone (CRH) and arginine vasopressin (AVP) from the paraventricular nucleus (PVN) of the
79 hypothalamus. These neuropeptides act on the pituitary, stimulating the release of
80 adrenocorticotrophic hormone (ACTH) which in turn causes the release of glucocorticoid stress
81 hormones (corticosterone in rodents, cortisol in humans (CORT)) from the adrenal cortex(de Kloet,
82 Joels, & Holsboer, 2005). Glucocorticoids cross the blood brain barrier and bind to corticosteroid
83 receptors (CR: glucocorticoid (GR) and mineralocorticoid (MR) receptors) distributed throughout the
84 brain. Feedback mechanisms then ensure the response is terminated in a healthy system. In contrast
85 to AVP, the closely related neuropeptide oxytocin (OXT) inhibits the activity of the HPA axis(Neumann
86 & Landgraf, 2019). HPA axis dysfunction is prevalent in psychiatric illness, for example HPA axis
87 hyperactivity is often found in major depression and bipolar disorder, and increased or decreased HPA
88 axis activity may be a direct consequence of ELS(Juruena, Cleare, & Young 2018; Murri et al., 2016;
89 Zorn et al., 2017).

90 Long-term effects of ELS on the HPA axis differ between the sexes. Early trauma is associated
91 with a more severely blunted cortisol response to social stress in women, and fewer stressful events
92 early in life are required to trigger liability to PTSD in women. Conversely, lower levels of recent stress

93 are capable of provoking major depression in men than women (Bunea, Szentagotai-Tatar, & Miu,
94 2017; McLaughlin, Conron, Koenen, & Gilman, 2010). Furthermore, women display 2-3 times higher
95 rates of anxiety, affective disorders, major depressive disorder and post-traumatic stress disorder
96 (PTSD)(Christiansen & Hansen, 2015; Kessler et al., 2003; Kessler, Chiu, Demler, Merikangas, &
97 Walters, 2005; Kessler, McGonagle, Swartz, Blazer, & Nelson, 1993; Remes, Brayne, van der Linde, &
98 Lafortune, 2016). These differences are not purely attributable to sex-specific life experiences; studies
99 controlling for stressful life events and sex-specific risk factors still find higher prevalence in
100 women(Tiwari & Gonzalez, 2018). Sex differences in HPA axis function may underlie this. Basal
101 secretion of CORT from the adrenal gland is higher in females than males, and this is attributed to sex-
102 differences in gonadal hormones, with estrogen sensitising and testosterone dampening the HPA-
103 axis(Heck & Handa, 2019; Seale et al., 2004).

104 Animal studies demonstrate that ELS has profound implications for later HPA axis function.
105 Prenatal and early post-natal stressors alter basal and stress-induced corticosterone release from the
106 adrenal glands, brain corticosteroid receptor (*GR* and *MR*) expression, as well as expression of AVP
107 and OXT in a timing and sometimes sex-specific manner(Llorente et al., 2011; Lupien, McEwen,
108 Gunnar, & Heim, 2009; Neumann & Landgraf, 2019; Schroeder, Notaras, Du, & Hill, 2018; Tobon,
109 Newport, & Nemeroff, 2018). However, despite well-established sex differences in the HPA axis,
110 comparatively few preclinical studies include male and female animals. Compared to the prenatal and
111 post-natal periods, less is known about the effects of stress experienced in the post-weaning, pre-
112 pubertal phase (PPS), a time-point suggested as more akin to human childhood(Brydges, 2016). The
113 limbic system and prefrontal cortex are undergoing maturation during this period, areas which are
114 crucial for cognition and emotion and are extremely stress reactive due to high densities of CR,
115 particularly in the hippocampal formation(Herman, 1993).

116 The present study investigated the effects of PPS on long-term neurochemical and molecular
117 alterations in the adult HPA axis in male and female animals by measuring the brain regional

118 expression of CR (*GR* and *MR*), *AVP*, *OXT* and their receptors (AVP receptor 1a (*AVPR1a*) and oxytocin
119 receptor (*OXTR*)) and *FKBP5*. *FKBP5* encodes the FK506 binding protein 51 co-chaperone protein of
120 the GR complex, and is extremely responsive to stress(Wochnik et al., 2005). When *FKBP5* is bound to
121 the GR complex, *CORT* binds with lower affinity and nuclear translocation of the receptor is less
122 efficient, decreasing negative feedback regulation of the HPA axis(Wochnik et al., 2005). There is
123 evidence that genetic modifications in *FKBP5* interact with childhood, but not adulthood stress to
124 increase risk for several psychiatric disorders(Matossin, Halldorsdottir, & Binder, 2018). We also
125 measured plasma corticosterone following a social test in adult rats as a behavioural measure of
126 altered HPA axis function. Altered social function is a core component of several adult psychiatric
127 illnesses and ELS has been shown to impact on social behaviour and functioning in both animal and
128 human studies(Nicol, Pope, Romaniuk, & Hall, 2015; Palmier-Claus et al., 2016; Sandi & Haller, 2015).
129 Furthermore, early life trauma is associated with blunted cortisol responses to social stress in humans,
130 particularly in women(Bunea et al., 2017). For this reason, we elected to focus on corticosterone
131 rather than other components of the HPA axis, such as ACTH. Further studies are need to determine
132 whether ACTH reflects the sex differences we observed in corticosterone.

133 We hypothesised that PPS would alter the expression of *GR*, *MR*, *GR:MR* ratio, *AVPR1a*, *AVP*
134 and *OXT* in the rodent brain, and the direction of change would be region and receptor/neuropeptide
135 specific. Given their higher vulnerability to stress-related psychiatric illnesses, we hypothesised
136 dysregulation resulting from PPS would be more pronounced in females. We also hypothesised that
137 corticosterone responses to social stress would be blunted in both males and females, but more
138 exaggerated in females, as early life trauma is often associated with a more pronounced blunting of
139 the corticosterone response in females(Bunea et al., 2017; McLaughlin et al., 2010).

140

141 **Methods**

142 *Animals.* Male and female Lister Hooded rats were bred at Cardiff University from 16 adult pairs
143 (Charles River). Females were primiparous. Litters ranged between 11 and 18 animals, with an average
144 of 14.5, and an average sex-ratio of 6.2 males to 8.3 females. All litters were used and weaning from
145 the birth dams took place on postnatal day (PND) 21, and offspring were housed in groups of 2-4 in
146 same litter, same sex cages (32cm x 50cm x 21cm) lined with wood shavings. Light was maintained on
147 12:12 hour light/dark cycle, a wooden stick, nesting material and cardboard tube were provided for
148 enrichment and food and water provided *ad libitum*. All experiments were approved by Cardiff
149 University's Animal Welfare and Ethical Review Body and adhered to the UK Home Office Animals
150 (Scientific Procedures) Act 1986 and European regulations on animal experimentation.

151 *Pre-pubertal stress (PPS).* Half of the offspring (8 litters) were pseudo-randomly allocated to a PPS
152 protocol (Jacobson-Pick & Richter-Levin, 2010) on PND 25-27 such that litters and sexes were equally
153 distributed between treatment groups (PPS/control, male/female). PPS took place in a designated
154 room separate to the holding room, with regular room lighting. On PND 25, animals were placed into
155 an opaque swim tank (25cm high, 34cm diameter) filled with 6L of 25±1°C water for a 10 minute swim
156 stress. On PND26 the rats were restrained in plastic restraint tubes (15cm length 5cm diameter) for
157 3x30 minute sessions (separated by 30 minute breaks in the home cage) and lastly on PND27 they
158 were exposed to elevated platforms (15x15cm, 115cm high) for 3x30 minute sessions (separated by
159 60 minute breaks in the home cage). **Animals were observed by the experimenter during all stress**
160 **procedures, and males and females reacted in a similar manner to each stressor.** Following PPS,
161 animals were left undisturbed until adulthood aside from weekly cage cleaning. Control animals were
162 left undisturbed from weaning until adulthood, aside from weekly cage cleaning.

163 *RT-qPCR.* Forty rats (male: 12 control, 10 PPS; female: 8 control, 10 PPS) were sacrificed at PND 60-70
164 using a rising concentration of CO₂. Brains were removed, dissected and stored at -80°C until analysis.
165 Total cell RNA was extracted from hippocampal formation, prefrontal cortex and hypothalamus using
166 the Qiagen RNeasy Kit (Qiagen, Manchester, UK) and DNase treated in accordance with the supplied

167 protocols. RNA was used to create cDNA for analysis using RNA to cDNA Easy Premix (Clontech
168 Laboratories, France), heated at 42°C for 75 minutes, followed by 80°C for 15 minutes. Sample was
169 then diluted 1:15 in nuclease-free water. 96-well plates were loaded, each well containing a total of
170 15µl reaction mixture (1.9µl sterile RNAase free water, 0.3µl 10µM forward primer, 0.3µl 10µM
171 reverse primer, 7.5µl SensiMix (Bioline) and 5µl cDNA). *Gapdh* and *Hprt1* primers (Sigma) were used
172 as housekeeping controls and all results were normalised from these values. After loading, plates were
173 spun down at 3,000 rpm for approximately 10-20 seconds before being transferred to Real-Time PCR
174 instrument (Applied Biosystems®) and run for 45 cycles (95°C for 20s, 60°C for 20s, 72°C for 20s). The
175 expression of *GR*, *MR*, *AVP*, *OXT*, *AVPR1a*, *OXTR* and *FKBP5* was measured (see Table 1 for primers).

176 *Social test.* Sixty-two animals (females: 22 control, 18 PPS; male: 12 control, 10 PPS) were given a social
177 test in same-sex pairs in adulthood (PND 60-67). Three hours before testing animals were single
178 housed in the holding room, and one hour before testing transferred to the testing room. All animals
179 were given an intraperitoneal injection of a vehicle (15%DMSO, 2% Tween 80 in 0.9% saline) 30
180 minutes before testing as part of a design to measure the effect of PPS on social behaviour directly,
181 an experiment which included a drug treated group. PPS had a significant effect on social interaction,
182 and this behavioural data is reported elsewhere (*Brydges et al. under review*). Animals were weighed
183 on the day of testing and placed in weight-matched pairs (weight difference did not exceed 20g) into
184 a clear acrylic arena (65cmx65cmx40cm high) on the floor in the middle of a dimly lit room (45lux) for
185 15 minutes. Animal pairs were from the same group, control or PPS, but different litters so were
186 strangers to each other.

187

188 *Corticosterone ELISA.* One animal from each pair was sacrificed 20 minutes after the social test to
189 investigate corticosterone responses to social interaction, the other sacrificed one week later for
190 baseline analysis. Animals were decapitated and trunk blood was collected using EDTA microvette
191 collection tubes (Sarstedt, Germany). Blood was spun at 1500 x g for 10 minutes, plasma was removed

192 and stored at -20°C until analysis. Corticosterone was analysed by ELISA, according to the
193 manufacturer's instructions (Abcam, UK, ab108821). The sensitivity of this ELISA is 0.28ng/ml and the
194 intra-assay coefficient of variation is 5.3%. Samples were run in triplicate on several plates,
195 counterbalancing between groups.

196

197 **Data analysis**

198 JMP statistical software (SAS Institute, Cary, NC, USA) was used to run generalised linear models. For
199 mRNA analysis, group (control/PPS), sex and group*sex were fitted as factors and mRNA expression
200 (normalised to *GAPDH* & *Hprt1*) as response. For corticosterone, group (control/PPS), time of sacrifice
201 (baseline vs 20 mins post social testing), sex and all two and three way interactions were fitted as
202 factors, corticosterone level as response. For all models, litter was nested within group and fitted as
203 a random factor to account for the use of multiple animals per litter. Data were checked for normality
204 and homogeneity of variance. Post-hoc t-tests were used when significant interactions were found.
205 The most relevant statistics are reported below, please see Table 2 for a full statistical summary.

206

207 **Results**

208 *mRNA* – Hippocampal formation. *FKBP5* (group*sex: $F_{1,29.85}=4.33$, $p=0.04$, Fig. 1a) and *AVPR1a*
209 (group*sex: $F_{1,26.64}=4.47$, $p=0.04$, Fig 1b) expression was significantly higher in the female hippocampal
210 formation following PPS, whereas *GR* (group: $F_{1,7.63}=0.01$, $p=0.92$), *MR* (group: $F_{1,9.44}=0.86$, $p=0.38$),
211 *GR:MR* (group: $F_{1,8.06}=0.88$, $p=0.38$), *AVP* (group: $F_{1,10.93}=0.62$, $p=0.45$), *OXT* (group: $F_{1,9.56}=0.26$, $p=0.62$)
212 and *OXTR* (group: $F_{1,9.84}=0.02$, $p=0.9$) were unchanged in males and females.

213 PFC. In the female PFC, PPS resulted in significantly higher *GR* expression (group*sex: $F_{1,24.94}=7.17$,
214 $p=0.01$, Fig. 2a), a higher *GR:MR* ratio (group*sex: $F_{1,25.37}=4.97$, $p=0.03$, Fig. 2b) and reduced *AVPR1a*
215 expression (group*sex: $F_{1,23.17}=6.94$, $p=0.01$, Fig. 2c) when compared to control females. *MR* (group:

216 $F_{1,8.81}=0.14$, $p=0.72$), *AVP* (group: $F_{1,9.77}=0.87$, $p=0.37$), *OXT* (group: $F_{1,10.34}=0.17$, $p=0.68$), *OXTR* (group:
217 $F_{1,10.39}=0.16$, $p=0.7$) and *FKBP5* (group: $F_{1,9.23}=1.43$, $p=0.26$) were unchanged in male and female PFC
218 following PPS, but *OXTR* expression was higher in females than males (sex: $F_{1,27.9}=28.65$, $p<0.0001$, Fig.
219 2d).

220 Hypothalamus. In the hypothalamus, *GR* (sex: $F_{1,30.7}=13.68$, $p<0.001$), *MR* (sex: $F_{1,25.52}=63.73$,
221 $p<0.0001$), *GR:MR* (sex: $F_{1,28.33}=8.93$, $p<0.01$), *FKBP5* (sex: $F_{1,26.23}=43.7$, $p<0.0001$) and *OXTR* (sex:
222 $F_{1,29.24}=118.81$, $p<0.0001$) were lower in females than males regardless of treatment (Fig 3a-e),
223 whereas *AVPR1a* (sex: $F_{1,25.55}=27.06$, $p<0.0001$) and *OXT* (sex: $F_{1,26.76}=42.06$, $p<0.0001$, Fig. 3f-g) were
224 higher in females. There was no effect of PPS on *GR* (group: $F_{1,10.79}=0.08$, $p=0.78$), *MR* (group: $F_{1,9}=0.53$,
225 $p=0.49$), *GR:MR* (group: $F_{1,11.03}=0.15$, $p=0.71$), *AVP* (group: $F_{1,7.69}=0.86$, $p=0.38$), *AVPR1a* (group:
226 $F_{1,10.42}=0.04$, $p=0.84$), *OXT* (group: $F_{1,10.33}=0.4$, $p=0.54$), *OXTR* (group: $F_{1,10.84}=0.43$, $p=0.53$) or *FKBP5*
227 (group: $F_{1,10.64}=0.63$, $p=0.44$), expression. See Table 3 for summary of regional gene expression
228 changes.

229 *Corticosterone*. There was no effect of PPS on baseline expression of plasma corticosterone, but 20
230 minutes after a social interaction test PPS blunted the normal corticosterone rise in females
231 (group*sex*time of sacrifice: $F_{7,37.77}=3.53$, $p=0.0052$, Fig. 4., Table 3).

232

233 **Discussion**

234 PPS altered the expression of receptors and neuropeptides involved in HPA axis function in the
235 hippocampal formation and PFC, but not hypothalamus of adult females, whereas in males the
236 expression of major HPA axis components were unaffected in all brain regions studied. In the female
237 prefrontal cortex, PPS increased *GR* and *GR:MR* receptor ratio and reduced *AVPR1a* expression. In the
238 female hippocampal formation, PPS increased expression of *FKBP5* and *AVPR1a*. In the periphery,
239 females exposed to PPS did not show the normal rise in blood corticosterone levels following a social

240 interaction test. We also found sex differences in baseline gene expression, particularly in the
241 hypothalamus.

242 GR and MR are nuclear receptors/transcription factors which mediate the actions of
243 glucocorticoid stress hormones, playing a key role in the stress response and also regulation of brain
244 development and neuronal plasticity(Liston & Gan, 2011). These corticosteroid receptors (CR) act
245 through delayed, long-lasting transcription-dependent mechanisms, but also exert more rapid effects
246 which dampen the activated HPA axis in a negative feedback, transcription-independent
247 manner(Gjerstad, Lightman, & Spiga, 2018; Tasker & Herman, 2011). Distributed throughout the brain,
248 CR expression is highest in the limbic system(Herman, 1993). We found that PPS had no effect on *GR*
249 or *MR* expression in male or female hippocampal formation. In contrast, stressors applied at earlier
250 time points (e.g. prenatal stress and maternal separation) generally decrease hippocampal CR
251 expression in males and females (although precise effects can vary depending on nature of the stress)
252 (Aisa, Tordera, Lasheras, Del Rio, & Ramirez, 2008; Brunton & Russell, 2010; Kapoor, Dunn, Kostaki,
253 Andrews, & Matthews, 2006; Levitt, Lindsay, Holmes, & Seckl, 1996; Maccari et al., 1995; Plotsky &
254 Meaney, 1993; van Bodegom, Homberg, & Henckens, 2017; Welberg, Seckl, & Holmes, 2001).
255 Stressors at later time points produce different effects, with chronic variable stress in adolescence
256 decreasing *GR* in the male hippocampus, and increasing/decreasing *MR* in male/female hippocampus
257 respectively(Isgor, Kabbaj, Akil, & Watson, 2004; Llorente et al., 2011). One study using mice found
258 that PPS increased *MR* and decreased *GR:MR* ratio in the hippocampus of adult male and female
259 animals(Brydges et al., 2014). Although PPS did not alter hippocampal CR expression directly in the
260 present study, we did find evidence of altered CR activity following PPS in females through increased
261 expression of *FKBP5*. *FKBP5* is a co-chaperone of heat shock protein 90 (hsp90) which regulates GR
262 sensitivity. Activation of GR leads to increased expression of *FKBP5*, creating an ultrashort negative
263 feedback loop which inhibits GR signalling(Wochnik et al., 2005; Zannas, Wiechmann, Gassen, &
264 Binder, 2016). Therefore, increased *FKBP5* likely indicates increased GR activity. Indeed, increased
265 expression of *FKBP5* in the limbic system (hippocampus and amygdala) is associated with increased

266 stress responsiveness (anxiety) and decreased stress coping behaviours, whereas experimental
267 reduction of *FKBP5* has opposite effects(Touma et al., 2011; Zannas et al., 2016).

268 The brain undergoes significant development postnatally, and the PFC is one of the last brain
269 regions to mature, undergoing synaptic remodelling in childhood and adolescence(Barfield & Gourley,
270 2018). Therefore pre-pubertal and adolescent stress may be particularly detrimental for the PFC, yet
271 little is known of the effects of **early life stress (ELS)** on CR expression in this region(Patel, Katz, Karssen,
272 & Lyons, 2008). In agreement with a previous study, we found that PPS did not impact *GR* or *MR*
273 expression in the male PFC(Fuentes, Carrasco, Armario, & Nadal, 2014). However, PPS did increase *GR*
274 and *GR:MR* ratio in the female PFC. Stress at an earlier timepoint, between PND 7-14, increased *GR*
275 expression in the PFC of female and male rats, again highlighting the importance of timing and
276 sex(Alteba, Korem, & Akirav, 2016). We found no evidence of altered *FKBP5* in the PFC following PPS,
277 but prenatal stress and maternal separation in rats decreases *FKBP5* expression in the male PFC with
278 no effects in the hippocampus, whereas chronic unpredictable stress in adolescence increases *FKBP5*
279 in the male hippocampus, PFC and amygdala(Szymanska et al., 2009; van der Doelen et al., 2014; Xu
280 et al., 2017; Xu et al., 2019). Overall our results suggest that PPS alters CR function in the adult
281 hippocampus and PFC, and this effect is specific to females.

282 PPS did not impact baseline corticosterone in males or females in agreement with the majority
283 of previous rodent research(Fuentes et al., 2014; Grigoryan, Ardi, Albrecht, Richter-Levin, & Segal,
284 2015; Jacobson-Pick & Richter-Levin, 2010). In humans, studies have found increased, decreased and
285 no change in basal cortisol following ELS(Agorastos, Pervanidou, Chrousos, & Baker, 2019; Lupien et
286 al., 2009). Differences are likely attributable to variation in the nature and timing of stress as well as
287 genetics, factors which are rarely considered in human studies. In the present study, exposure to social
288 interaction with a stranger resulted in elevated corticosterone in the plasma of control females and
289 all males, but this response was blunted in females with experience of PPS. Note that the increases in
290 corticosterone are not an acute response to systemic vehicle administration but are a result of the

291 social interaction, since all animals, both baseline and socially experienced rats, received vehicle
292 injections. Furthermore, the injection occurred sixty-five minutes before sacrifice, so any acute
293 corticosterone rise resulting from this would no longer be detectable. In contrast to the results
294 presented here, mild prenatal stress results in *heightened* corticosterone response to restraint stress
295 in adult females(Aisa et al., 2008). Interestingly, more prolonged prenatal stress is necessary to induce
296 the same effects in males(Gobinath, Mahmoud, & Galea, 2015). Maternal separation elevates or
297 blunts male and female corticosterone responses to restraint stress, depending on the
298 study(Desbonnet, Garrett, Daly, McDermott, & Dinan, 2008; Lehmann, Russig, Feldon, & Pryce, 2002;
299 Roman, Gustafsson, Berg, & Nylander, 2006), and chronic variable adolescent stress between PND 45-
300 58/37-49 blunted corticosterone responses to a stressor in adult females but not males(Bourke &
301 Neigh, 2011; Wulsin, Wick-Carlson, Packard, Morano, & Herman, 2016). This again suggests that the
302 female HPA axis is more sensitive to ELS, although specific outcomes are mediated by exact timing of
303 stress and adult testing paradigm (e.g. social vs restraint). An adaptive stress response is characterised
304 by a rapid **corticosterone or cortisol** (CORT) increase, followed by a progressive decline. Excessive or
305 repeated activation of the HPA axis and release of CORT can lead to blunted CORT secretion in
306 response to acute stress(Kinlein, Wilson, & Karatsoreos, 2015). A healthy CORT response is necessary
307 for appropriate behaviour and survival, therefore a blunted CORT response to acute stress may be
308 considered a maladaptive phenotype.

309 PPS increased expression of *AVPR1a* in the female hippocampal formation and decreased it in
310 the PFC. No changes were observed in the hypothalamus. OXT and AVP are closely related
311 neuropeptides that exert opposite effects on the HPA axis. Stress results in the release of
312 hypothalamic AVP, which stimulates the release of adrenocorticotrophic hormone from the pituitary
313 and eventual production of CORT(de Kloet et al., 2005). In contrast, OXT dampens the HPA
314 axis(Neumann & Landgraf, 2019). Both AVP and OXT exert effects on behaviour through OXTR and
315 AVPR1a/AVPR1b situated in the brain(Song & Albers, 2018). Effects of prenatal stress on AVP/OXT
316 systems are mixed, with some studies finding decreased OXT/AVP expression in the male

317 hypothalamus, others no changes in males or females (Desbonnet et al., 2008; Lee, Brady, Shapiro,
318 Dorsa, & Koenig, 2007; Schmidt et al., 2018). Poor maternal care in rodents decreases OXT and OXTR
319 expression centrally (hypothalamus and amygdala) and peripherally (blood plasma) in female
320 animals (Francis, Young, Meaney, & Insel, 2002; Tobon, Jeffrey, & Nemeroff, 2018), whereas maternal
321 separation increases hypothalamic AVP and alters OXT expression and OXT/AVP receptor binding in
322 an age and sex-specific manner (Lukas, Bredewold, Neumann, & Veenema, 2010; Murgatroyd et al.,
323 2009; Veenema, Bredewold, & Neumann, 2007; Veenema & Neumann, 2009). Our previous work
324 found PPS increased protein levels of AVP in the supraoptic (but not paraventricular) nucleus of the
325 hypothalamus and blood plasma in male and female rats (*Brydges et al. in Review*). In the present
326 study, the hypothalamus was analysed as a whole, it is possible differences may have been found if
327 the supraoptic and paraventricular nuclei had been analysed separately. Alternatively, PPS may alter
328 translation rather than transcription of AVP in this region. Considering their opposing effects on
329 behaviour (AVP exerts anxiogenic and depressive-like effects, whereas OXT is an endogenous
330 anxiolytic), the balance of AVP and OXT in the brain is thought crucial for appropriate emotional
331 behaviours (Mak, Broussard, Vacy, & Broadbear, 2012; Neumann & Landgraf, 2012). In the present
332 study, we find altered *AVPR1a* expression in the female limbic system in the absence of altered
333 OXT/OXTR expression. This indicates a dysregulated HPA axis which may predispose towards anxiety
334 or depressive phenotypes following stress (Lesse, Rether, Groger, Braun, & Bock, 2017; Neumann &
335 Slattery, 2016; Nowacka-Chmielewska, Kasprowska-Liskiewicz, Barski, Obuchowicz, & Malecki, 2017).

336 PPS increased *AVPR1a* expression in the female hippocampal formation, yet decreased
337 expression in the PFC. Bi-directional projections exist between the hippocampus and hypothalamus
338 (production site of AVP): the hippocampus is a target of AVP and is capable of decreasing AVP
339 expression in the hypothalamus (Nettles, Pesold, & Goldman, 2000; Zhang & Hernandez, 2013). PPS
340 leads to increased AVP (*Brydges et al. under review*), therefore increased *AVPR1a* expression in the
341 hippocampal formation may be a compensatory mechanism, enhancing the sensitivity to and
342 subsequent inhibitory effects of the hippocampus on hypothalamic AVP secretion. The PFC is also

343 thought to exert inhibitory effects over the hypothalamus, but direct connections between these two
344 structures are lacking, and it is hypothesised that the PFC may act via other structures to exert this
345 influence(Spencer, Buller, & Day, 2005). Whether the decreased *AVPR1a* expression following PPS in
346 this region is due to adaptation or pathology remains to be elucidated.

347 Striking sex differences were seen regardless of PPS. *GR*, *MR*, *GR:MR*, *FKBP5* and *OXTR*
348 expression were significantly higher in male than female hypothalamus, whereas *AVPR1a* and *OXT*
349 showed the opposite pattern. *OXTR* expression was higher in female PFC. These findings are consistent
350 with previous studies finding *AVPR1a* expression is higher in female vs male rodents, and *GR*, *MR* and
351 *OXTR* expression higher in the male hypothalamus (although species, age and region studied can all
352 affect direction of difference)(Albers, 2015; Bale & Dorsa, 1995; Dumais, Bredewold, Mayer, &
353 Veenema, 2013; Smith et al., 2017; Turner, 1990). One study investigating binding in 35 different
354 rodent brains regions similarly found sex differences in *OXTR* and *AVPR1a* expression
355 (increased/decreased depending on region)(Smith et al., 2017), but less is known about *FKBP5*. These
356 sex differences may confer a natural heightened reactivity to stress in females which may underlie the
357 greater vulnerability of the female HPA axis to ELS.

358 The balance between MR and GR functioning is thought crucial for appropriate HPA axis
359 function, and dysregulation and imbalance between CR is suggested as a candidate mechanism
360 underlying psychiatric disorders such as major depression, a disorder which has been repeatedly
361 associated with hyperactive HPA axis function(de Kloet et al., 2005; Juruena et al., 2015; Oitzl,
362 Champagne, van der Veen, & de Kloet, 2010). Polymorphisms associated with enhanced expression of
363 *FKBP5* following GR activation are overrepresented in major depression, bipolar and PTSD(Binder,
364 2009; Matosin et al., 2018). *FKBP5* is implicated in a number psychiatric disorders, particularly in
365 combination with early life stress(Wang, Shelton, & Dwivedi, 2018). In humans, there is an interaction
366 between *FKBP5* (FK506 binding protein 5) variability and childhood trauma on psychosis, paranoia,
367 social stress appraisal and prefrontal cortex function(Harms et al., 2017; Misiak et al., 2018; Wang et

368 al., 2018). Our results suggest that PPS plays a role in altered FKBP5 functioning in females, a key
369 regulator of the HPA axis. Also in agreement with our findings, the human literature shows early
370 trauma is associated with blunted CORT responses to social stimuli, particularly in women.

371

372 **Conclusions**

373 We found the adult female HPA axis was sensitive to PPS, with changes seen throughout the system.
374 This highlights the pre-pubertal phase as a particularly sensitive time for re-programming of the
375 female HPA axis by stress. In contrast, the male HPA axis was unaffected. This sex-specific vulnerability
376 may underlie the greater propensity for women to develop psychiatric disorders including depression,
377 anxiety and PTSD, disorders which are frequently associated with HPA axis dysregulation. Although
378 we found greater effects in females in the present study, males are not immune to the effects of PPS.
379 For example, in previous studies we found that PPS significantly impaired hippocampal-dependent
380 behaviour and hippocampal neurogenesis in males but not females, and social behaviour is equally
381 affected in both sexes(Brydges et al., 2018, *Brydges et al. in review*). Furthermore, others have found
382 several behavioural and neurobiological effects of PPS in male animals(Albrecht et al., 2017; Brydges,
383 2016). This suggests males and females differ in their responses to PPS, potentially resulting in sex-
384 specific vulnerabilities to certain disorders. This strengthens the argument for including both sexes in
385 preclinical and clinical studies.

386

387 **Figure Legends**

388 **Figure 1. Hippocampal formation.** PPS increased expression of a) FKBP5 and b) AVPR1a in the female
389 hippocampal formation. Con=control, PPS=pre-pubertal stress, F=female, M=male. Male: 12 control,
390 10 PPS; female: 8 control, 10 PPS. *=p<0.05. Error bars represent 1 S.E. and bars joined by a line and
391 asterisk are significantly different to one another.

392 **Figure 2. PFC.** PPS increased a) GR and b) GR:MR ratio and decreased c) AVPR1a in the female PFC.
393 OXTR expression was higher in female than male PFC. Con=control, PPS=pre-pubertal stress,
394 F=female, M=male. Male: 12 control, 10 PPS; female: 8 control, 10 PPS. *=p<0.05, **p<0.01,
395 ***p<0.0001. Error bars represent 1 S.E. and bars joined by a line and asterisk are significantly
396 different to one another.

397 **Figure 3. Hypothalamus.** a) GR, b) MR, c) GR:MR, d) FKBP5 and e) OXTR were higher in male than
398 female hypothalamus, whereas f) AVPR1a and g) OXT were higher in female hypothalamus.
399 Con=control, PPS=pre-pubertal stress, F=female, M=male. Male: 12 control, 10 PPS; female: 8 control,
400 10 PPS. **p<0.01, ***p<0.0001. Error bars represent 1 S.E. and bars joined by a line and asterisk are
401 significantly different to one another.

402 **Figure 4. Corticosterone.** Social interaction significantly elevated corticosterone above baseline in
403 control animals and PPS males. This response was blunted in PPS females. Con=control, PPS=pre-
404 pubertal stress, F=female, M=male. Females: 22 control, 18 PPS; male: 12 control, 10 PPS *=p<0.05.
405 Error bars represent 1 S.E. and bars joined by a line and asterisk are significantly different to one
406 another.

407

408 **Acknowledgements**

409 We wish to acknowledge support from the Cardiff University Neuroscience and Mental Health
410 Research Institute and The Jane Hodge Foundation who provided NB with fellowship funding during
411 this research, as well as The Waterloo Foundation who provided grant funding for preliminary work
412 (grant number 918-1875).

413

414 **Declaration of interest**

415 The authors declare no competing interest.

- 417 Agorastos, A., Pervanidou, P., Chrousos, G. P., & Baker, D. G. (2019). Developmental Trajectories of
418 Early Life Stress and Trauma: A Narrative Review on Neurobiological Aspects Beyond Stress
419 System Dysregulation. *Frontiers in Psychiatry*, *10*. doi:10.3389/fpsy.2019.00118
- 420 Aisa, B., Tordera, R., Lasheras, B., Del Rio, J., & Ramirez, M. J. (2008). Effects of maternal separation
421 on hypothalamic-pituitary-adrenal responses, cognition and vulnerability to stress in adult
422 female rats. *Neuroscience*, *154*(4), 1218-1226. doi:10.1016/j.neuroscience.2008.05.011
- 423 Albers, H. E. (2015). Species, sex and individual differences in the vasotocin/vasopressin system:
424 Relationship to neurochemical signaling in the social behavior neural network. *Frontiers in*
425 *Neuroendocrinology*, *36*, 49-71. doi:10.1016/j.yfrne.2014.07.001
- 426 Albrecht, A., Muller, I., Ardi, Z., Caliskan, G., Gruber, D., Ivens, S., . . . Richter-Levin, G. (2017).
427 Neurobiological consequences of juvenile stress: A GABAergic perspective on risk and
428 resilience. *Neuroscience and Biobehavioral Reviews*, *74*, 21-43.
429 doi:10.1016/j.neubiorev.2017.01.005
- 430 Alteba, S., Korem, N., & Akirav, I. (2016). Cannabinoids reverse the effects of early stress on
431 neurocognitive performance in adulthood. *Learning & Memory*, *23*(7), 349-358.
432 doi:10.1101/lm.041608.116
- 433 Binder, E. B. (2009). The role of FKBP5, a co-chaperone of the glucocorticoid receptor in the
434 pathogenesis and therapy of affective and anxiety disorders. In (Vol. 34, pp. S186-195).
435 Psychoneuroendocrinology.
- 436 Bale, T. L., & Dorsa, D. M. (1995). Sex-differences in and effects of estrogen on oxytocin receptor
437 messenger-ribonucleic-acid expression in the ventromedial hypothalamus. *S. Endocrinology*,
438 *136*(1), 27-32. doi:10.1210/en.136.1.27
- 439 Barfield, E. T., & Gourley, S. L. (2018). Prefrontal cortical trkB, glucocorticoids, and their interactions
440 in stress and developmental contexts. *Neuroscience and Biobehavioral Reviews*, *95*, 535-558.
441 doi:10.1016/j.neubiorev.2018.10.015
- 442 Bourke, C. H., & Neigh, G. N. (2011). Behavioral effects of chronic adolescent stress are sustained and
443 sexually dimorphic. *Hormones and Behavior*, *60*(1), 112-120.
444 doi:10.1016/j.yhbeh.2011.03.011
- 445 Brunton, P. J., & Russell, J. A. (2010). Prenatal Social Stress in the Rat Programmes Neuroendocrine
446 and Behavioural Responses to Stress in the Adult Offspring: Sex-Specific Effects. *Journal of*
447 *Neuroendocrinology*, *22*(4), 258-271. doi:10.1111/j.1365-2826.2010.01969.x
- 448 Brydges, N. M. (2016). Pre-pubertal stress and brain development in rodents. *Current Opinion in*
449 *Behavioral Sciences*, *7*, 8-14. doi:10.1016/j.cobeha.2015.08.003
- 450 Brydges, N. M., Jin, R. W., Seckl, J., Holmes, M. C., Drake, A. J., & Hall, J. (2014). Juvenile stress enhances
451 anxiety and alters corticosteroid receptor expression in adulthood. *Brain and Behavior*, *4*(1),
452 4-13. doi:10.1002/brb3.182
- 453 Brydges, N. M., Moon, A., Rule, L., Watkin, H., Thomas, K. L., & Hall, J. (2018). Sex specific effects of
454 pre-pubertal stress on hippocampal neurogenesis and behaviour. *Translational Psychiatry*, *8*.
455 doi:10.1038/s41398-018-0322-4
- 456 *Brydges et al. in review.* Brydges, N.M., Hall, J., Best, C., Rule, L, Watkin, H., Drake, A.J., Lewis, C.,
457 Thomas, K.L. & Hall, J. Early life stress impairs social function through AVP-dependent
458 mechanisms.
- 459 Bunea, I. M., Szentagotai-Tatar, A., & Miu, A. C. (2017). Early-life adversity and cortisol response to
460 social stress: a meta-analysis. *Translational Psychiatry*, *7*, 8. doi:10.1038/s41398-017-0032-3
- 461 Christiansen, D. M., & Hansen, M. (2015). Accounting for sex differences in PTSD: A multi-variable
462 mediation model. *European Journal of Psychotraumatology*, *6*, 10. doi:10.3402/ejpt.v6.26068
- 463 de Kloet, E. R., Joels, M., & Holsboer, F. (2005). Stress and the brain: From adaptation to disease.
464 *Nature Reviews Neuroscience*, *6*(6), 463-475. doi:10.1038/nrn1683

465 Desbonnet, L., Garrett, L., Daly, E., McDermott, K. W., & Dinan, T. G. (2008). Sexually dimorphic effects
466 of maternal separation stress on corticotrophin-releasing factor and vasopressin systems in
467 the adult rat brain. *International Journal of Developmental Neuroscience*, 26(3-4), 259-268.
468 doi:10.1016/j.ijdevneu.2008.02.004

469 Dumais, K. M., Bredewold, R., Mayer, T. E., & Veenema, A. H. (2013). Sex differences in oxytocin
470 receptor binding in forebrain regions: Correlations with social interest in brain region- and
471 sex- specific ways. *Hormones and Behavior*, 64(4), 693-701. doi:10.1016/j.yhbeh.2013.08.012

472 Juruena, M.F., Cleare, A.J., & Young, A.H. (2018). The role of early life stress in HPA axis and depression.
473 In (pp. 71-80). *Understanding Depression*: Springer.

474 Francis, D. D., Young, L. J., Meaney, M. J., & Insel, T. R. (2002). Naturally occurring differences in
475 maternal care are associated with the expression of oxytocin and vasopressin (V1a) receptors:
476 Gender differences. *Journal of Neuroendocrinology*, 14(5), 349-353. doi:10.1046/j.0007-
477 1331.2002.00776.x

478 Fuentes, S., Carrasco, J., Armario, A., & Nadal, R. (2014). Behavioral and neuroendocrine consequences
479 of juvenile stress combined with adult immobilization in male rats. *Hormones and Behavior*,
480 66(3), 475-486. doi:10.1016/j.yhbeh.2014.07.003

481 Gjerstad, J. K., Lightman, S. L., & Spiga, F. (2018). Role of glucocorticoid negative feedback in the
482 regulation of HPA axis pulsatility. *Stress-the International Journal on the Biology of Stress*,
483 21(5), 403-416. doi:10.1080/10253890.2018.1470238

484 Gobinath, A. R., Mahmoud, R., & Galea, L. A. M. (2015). Influence of sex and stress exposure across
485 the lifespan on endophenotypes of depression: focus on behavior glucocorticoids, and
486 hippocampus. *Frontiers in Neuroscience*, 8. doi:10.3389/fnins.2014.00420

487 Grigoryan, G., Ardi, Z., Albrecht, A., Richter-Levin, G., & Segal, M. (2015). Juvenile stress alters LTP in
488 ventral hippocampal slices: Involvement of noradrenergic mechanisms. *Behavioural Brain
489 Research*, 278, 559-562. doi:10.1016/j.bbr.2014.09.047

490 Harms, M. B., Birn, R., Provencal, N., Wiechmann, T., Binder, E. B., Giakas, S. W., . . . Pollak, S. D. (2017).
491 Early life stress, FK506 binding protein 5 gene (FKBP5) methylation, and inhibition-related
492 prefrontal function: A prospective longitudinal study. *Development and Psychopathology*,
493 29(5), 1895-1903. doi:10.1017/s095457941700147x

494 Heck, A. L., & Handa, R. J. (2019). Sex differences in the hypothalamic-pituitary-adrenal axis' response
495 to stress: an important role for gonadal hormones. *Neuropsychopharmacology*, 44(1), 45-58.
496 doi:10.1038/s41386-018-0167-9

497 Heim, C., & Nemeroff, C. B. (2001). The role of childhood trauma in the neurobiology of mood and
498 anxiety disorders: Preclinical and clinical studies. *Biological Psychiatry*, 49(12), 1023-1039.
499 doi:10.1016/s0006-3223(01)01157-x

500 Herman, J. P. (1993). Regulation of adrenocorticosteroid receptor messenger-RNA expression in the
501 central nervous system. *Cellular and Molecular Neurobiology*, 13(4), 349-372.
502 doi:10.1007/bf00711577

503 Isgor, C., Kabbaj, M., Akil, H., & Watson, S. J. (2004). Delayed effects of chronic variable stress during
504 peripubertal-juvenile period on hippocampal morphology and on cognitive and stress axis
505 functions in rats. *Hippocampus*, 14(5), 636-648. doi:10.1002/hipo.10207

506 Jacobson-Pick, S., & Richter-Levin, G. (2010). Differential impact of juvenile stress and corticosterone
507 in juvenility and in adulthood, in male and female rats. *Behavioural Brain Research*, 214(2),
508 268-276. doi:10.1016/j.bbr.2010.05.036

509 Juruena, M. F., Baes, C. V., Menezes, I. C., & Graeff, F. G. (2015). Early Life Stress in Depressive Patients:
510 Role of Glucocorticoid and Mineralocorticoid Receptors and of Hypothalamic-Pituitary-
511 Adrenal Axis Activity. *Current Pharmaceutical Design*, 21(11), 1369-1378.
512 doi:10.2174/1381612821666150105125500

513 Kapoor, A., Dunn, E., Kostaki, A., Andrews, M. H., & Matthews, S. G. (2006). Fetal programming of
514 hypothalamo-pituitary-adrenal function: prenatal stress and glucocorticoids. *Journal of
515 Physiology-London*, 572(1), 31-44. doi:10.1113/jphysiol.2006.105254

516 Kessler, R. C., Berglund, P., Demler, O., Jin, R., Koretz, D., Merikangas, K. R., . . . Wang, P. S. (2003). The
517 epidemiology of major depressive disorder - Results from the National Comorbidity Survey
518 Replication (NCS-R). *Jama-Journal of the American Medical Association*, 289(23), 3095-3105.
519 doi:10.1001/jama.289.23.3095

520 Kessler, R. C., Chiu, W. T., Demler, O., Merikangas, K. R., & Walters, E. E. (2005). Prevalence, severity,
521 and comorbidity of 12-month DSM-IV disorders in the national comorbidity survey replication.
522 (vol 62, pg 617, 2005). *Archives of General Psychiatry*, 62(7), 709-709.

523 Kessler, R. C., McGonagle, K. A., Swartz, M., Blazer, D. G., & Nelson, C. B. (1993). Sex and depression
524 in the national comorbidity survey .1. lifetime prevalence, chronicity and recurrence.. *Journal*
525 *of Affective Disorders*, 29(2-3), 85-96. doi:10.1016/0165-0327(93)90026-g

526 Kinlein, S. A., Wilson, C. D., & Karatsoreos, I. N. (2015). Dysregulated hypothalamic-pituitary-adrenal
527 axis function contributes to altered endocrine and neurobehavioral responses to acute stress.
528 *Frontiers in Psychiatry*, 6. doi:10.3389/fpsy.2015.00031

529 Tobon, A., Jeffrey, N., & Nemeroff, C.B. (2018). The role of oxytocin in early life adversity and later
530 psychopathology: a review of preclinical and clinical studies. In (Vol. 5, pp. 401-415). *Current*
531 *Treatment Options in Psychiatry*.

532 Lee, P. R., Brady, D. L., Shapiro, R. A., Dorsa, D. M., & Koenig, J. I. (2007). Prenatal stress generates
533 deficits in rat social behavior: Reversal by oxytocin. *Brain Research*, 1156, 152-167.
534 doi:10.1016/j.brainres.2007.04.042

535 Lehmann, J., Russig, H., Feldon, J., & Pryce, C. R. (2002). Effect of a single maternal separation at
536 different pup ages on the corticosterone stress response in adult and aged rats. *Pharmacology*
537 *Biochemistry and Behavior*, 73(1), 141-145. doi:10.1016/s0091-3057(02)00788-8

538 Lesse, A., Rether, K., Groger, N., Braun, K., & Bock, J. (2017). Chronic Postnatal Stress Induces
539 Depressive-like Behavior in Male Mice and Programs second-Hit Stress-Induced Gene
540 Expression Patterns of OxtR and AvpR1a in Adulthood. *Molecular Neurobiology*, 54(6), 4813-
541 4819. doi:10.1007/s12035-016-0043-8

542 Levitt, N. S., Lindsay, R. S., Holmes, M. C., & Seckl, J. R. (1996). Dexamethasone in the last week of
543 pregnancy attenuates hippocampal glucocorticoid receptor gene expression and elevates
544 blood pressure in the adult offspring in the rat. *Neuroendocrinology*, 64(6), 412-418.
545 doi:10.1159/000127146

546 Liston, C., & Gan, W. B. (2011). Glucocorticoids are critical regulators of dendritic spine development
547 and plasticity in vivo. *Proceedings of the National Academy of Sciences of the United States of*
548 *America*, 108(38), 16074-16079. doi:10.1073/pnas.1110444108

549 Llorente, R., Miguel-Blanco, C., Aisa, B., Lachize, S., Borcel, E., Meijer, O. C., . . . Viveros, M. P. (2011).
550 Long Term Sex-Dependent Psychoneuroendocrine Effects of Maternal Deprivation and
551 Juvenile Unpredictable Stress in Rats. *Journal of Neuroendocrinology*, 23(4), 329-344.
552 doi:10.1111/j.1365-2826.2011.02109.x

553 Lukas, M., Bredewold, R., Neumann, I. D., & Veenema, A. H. (2010). Maternal separation interferes
554 with developmental changes in brain vasopressin and oxytocin receptor binding in male rats.
555 *Neuropharmacology*, 58(1), 78-87. doi:10.1016/j.neuropharm.2009.06.020

556 Lupien, S. J., McEwen, B. S., Gunnar, M. R., & Heim, C. (2009). Effects of stress throughout the lifespan
557 on the brain, behaviour and cognition. *Nature Reviews Neuroscience*, 10(6), 434-445.
558 doi:10.1038/nrn2639

559 Maccari, S., Piazza, P. V., Kabbaj, M., Barbazanges, A., Simon, H., & Lemoal, M. (1995). Adoption
560 reverses the long-term impairment in glucocorticoid feedback induced by prenatal stress.
561 *Journal of Neuroscience*, 15(1), 110-116.

562 Mak, P., Broussard, C., Vacy, K., & Broadbear, J. H. (2012). Modulation of anxiety behavior in the
563 elevated plus maze using peptidic oxytocin and vasopressin receptor ligands in the rat. *Journal*
564 *of Psychopharmacology*, 26(4), 532-542. doi:10.1177/0269881111416687

565 Matosin, N., Halldorsdottir, T., & Binder, E. B. (2018). Understanding the Molecular Mechanisms
566 Underpinning Gene by Environment Interactions in Psychiatric Disorders: The FKBP5 Model.
567 *Biological Psychiatry*, *83*(10), 821-830. doi:10.1016/j.biopsych.2018.01.021

568 McLaughlin, K. A., Conron, K. J., Koenen, K. C., & Gilman, S. E. (2010). Childhood adversity, adult
569 stressful life events, and risk of past-year psychiatric disorder: a test of the stress sensitization
570 hypothesis in a population-based sample of adults. *Psychological Medicine*, *40*(10), 1647-
571 1658. doi:10.1017/s0033291709992121

572 Misiak, B., Stramecki, F., Gaweda, L., Prochwicz, K., Sasiadek, M. M., Moustafa, A. A., & Frydecka, D.
573 (2018). Interactions Between Variation in Candidate Genes and Environmental Factors in the
574 Etiology of Schizophrenia and Bipolar Disorder: a Systematic Review. *Molecular Neurobiology*,
575 *55*(6), 5075-5100. doi:10.1007/s12035-017-0708-y

576 Murgatroyd, C., Patchev, A. V., Wu, Y., Micale, V., Bockmuhl, Y., Fischer, D., . . . Spengler, D. (2009).
577 Dynamic DNA methylation programs persistent adverse effects of early-life stress. *Nature*
578 *Neuroscience*, *12*(12), 1559-U1108. doi:10.1038/nn.2436

579 Murri, M. B., Prestia, D., Mondelli, V., Pariante, C., Patti, S., Olivieri, B., . . . Amore, M. (2016). The HPA
580 axis in bipolar disorder: Systematic review and meta-analysis. *Psychoneuroendocrinology*, *63*,
581 327-342. doi:10.1016/j.psyneuen.2015.10.014

582 Nettles, K. W., Pesold, C., & Goldman, M. B. (2000). Influence of the ventral hippocampal formation
583 on plasma vasopressin, hypothalamic-pituitary-adrenal axis, and behavioral responses to
584 novel acoustic stress. *Brain Research*, *858*(1), 181-190. doi:10.1016/s0006-8993(99)02281-7

585 Neumann, I. D., & Landgraf, R. (2012). Balance of brain oxytocin and vasopressin: implications for
586 anxiety, depression, and social behaviors. *Trends in Neurosciences*, *35*(11), 649-659.
587 doi:10.1016/j.tins.2012.08.004

588 Neumann, I. D., & Landgraf, R. (2019). Tracking oxytocin functions in the rodent brain during the last
589 30 years: From push-pull perfusion to chemogenetic silencing. *Journal of Neuroendocrinology*,
590 *31*(3). doi:10.1111/jne.12695

591 Neumann, I. D., & Slattery, D. A. (2016). Oxytocin in General Anxiety and Social Fear: A Translational
592 Approach. *Biological Psychiatry*, *79*(3), 213-221. doi:10.1016/j.biopsych.2015.06.004

593 Nicol, K., Pope, M., Romaniuk, L., & Hall, J. (2015). Childhood trauma, midbrain activation and
594 psychotic symptoms in borderline personality disorder. *Translational Psychiatry*, *5*.
595 doi:10.1038/tp.2015.53

596 Nowacka-Chmielewska, M. M., Kasprowska-Liskiewicz, D., Barski, J. J., Obuchowicz, E., & Malecki, A.
597 (2017). The behavioral and molecular evaluation of effects of social instability stress as a
598 model of stress-related disorders in adult female rats. *Stress-the International Journal on the*
599 *Biology of Stress*, *20*(6), 549-561. doi:10.1080/10253890.2017.1376185

600 Oitzl, M. S., Champagne, D. L., van der Veen, R., & de Kloet, E. R. (2010). Brain development under
601 stress: Hypotheses of glucocorticoid actions revisited. *Neuroscience and Biobehavioral*
602 *Reviews*, *34*(6), 853-866. doi:10.1016/j.neubiorev.2009.07.006

603 Palmier-Claus, J., Berry, K., Darrell-Berry, H., Emsley, R., Parker, S., Drake, R., & Bucci, S. (2016).
604 Childhood adversity and social functioning in psychosis: Exploring clinical and cognitive
605 mediators. *Psychiatry Research*, *238*, 25-32. doi:10.1016/j.psychres.2016.02.004

606 Patel, P. D., Katz, M., Karssen, A. M., & Lyons, D. M. (2008). Stress-induced changes in corticosteroid
607 receptor expression in primate hippocampus and prefrontal cortex.
608 *Psychoneuroendocrinology*, *33*(3), 360-367. doi:10.1016/j.psyneuen.2007.12.003

609 Plotsky, P. M., & Meaney, M. J. (1993). Early postnatal experience alters hypothalamic corticotropin-
610 releasing factor (CRF) messenger-RNA, median eminence CRF content and stress-induced
611 release in adult rats. *Molecular Brain Research*, *18*(3), 195-200. doi:10.1016/0169-
612 328x(93)90189-v

613 Remes, O., Brayne, C., van der Linde, R., & Lafortune, L. (2016). A systematic review of reviews on the
614 prevalence of anxiety disorders in adult populations. *Brain and Behavior*, *6*(7).
615 doi:10.1002/brb3.497

616 Roman, E., Gustafsson, L., Berg, M., & Nylander, I. (2006). Behavioral profiles and stress-induced
617 corticosteroid secretion in male Wistar rats subjected to short and prolonged periods of
618 maternal separation. *Hormones and Behavior*, *50*(5), 736-747.
619 doi:10.1016/j.yhbeh.2006.06.016

620 Sandi, C., & Haller, J. (2015). Stress and the social brain: behavioural effects and neurobiological
621 mechanisms. *Nature Reviews Neuroscience*, *16*(5), 290-304. doi:10.1038/nrn3918

622 Schmidt, M., Braun, K., Brandwein, C., Rossetti, A. C., Ciurana, S. G., Riva, M. A., . . . Groger, N. (2018).
623 Maternal stress during pregnancy induces depressive-like behavior only in female offspring
624 and correlates to their hippocampal Avp and Oxt receptor expression. *Behavioural Brain
625 Research*, *353*, 1-10. doi:10.1016/j.bbr.2018.06.027

626 Schroeder, A., Notaras, M., Du, X., & Hill, R. A. (2018). On the Developmental Timing of Stress:
627 Delineating Sex-Specific Effects of Stress across Development on Adult Behavior. *Brain
628 Sciences*, *8*(7). doi:10.3390/brainsci8070121

629 Seale, J. V., Wood, S. A., Atkinson, H. C., Bate, E., Lightman, S. L., Ingram, C. D., . . . Harbuz, M. S. (2004).
630 Gonadectomy reverses the sexually dimorphic patterns of circadian and stress-induced
631 hypothalamic-pituitary-adrenal axis activity in male and female rats. *Journal of
632 Neuroendocrinology*, *16*(6), 516-524. doi:10.1111/j.1365-2826.2004.01195.x

633 Smith, C. J. W., Poehlmann, M. L., Li, S., Ratnaseelan, A. M., Bredewold, R., & Veenema, A. H. (2017).
634 Age and sex differences in oxytocin and vasopressin V1a receptor binding densities in the rat
635 brain: focus on the social decision-making network. *Brain Structure & Function*, *222*(2), 981-
636 1006. doi:10.1007/s00429-016-1260-7

637 Song, Z. M., & Albers, H. E. (2018). Cross-talk among oxytocin and arginine-vasopressin receptors:
638 Relevance for basic and clinical studies of the brain and periphery. *Frontiers in
639 Neuroendocrinology*, *51*, 14-24. doi:10.1016/j.yfrne.2017.10.004

640 Spencer, S. J., Buller, K. M., & Day, T. A. (2005). Medial prefrontal cortex control of the Paraventricular
641 hypothalamic nucleus response to psychological stress: Possible role of the bed nucleus of the
642 stria terminalis. *Journal of Comparative Neurology*, *481*(4), 363-376. doi:10.1002/cne.20376

643 Szymanska, M., Budziszewska, B., Jaworska-Feil, L., Basta-Kaim, A., Kubera, M., Leskiewicz, M., . . .
644 Lason, W. (2009). The effect of antidepressant drugs on the HPA axis activity, glucocorticoid
645 receptor level and FKBP51 concentration in prenatally stressed rats.
646 *Psychoneuroendocrinology*, *34*(6), 822-832. doi:10.1016/j.psyneuen.2008.12.012

647 Tasker, J. G., & Herman, J. P. (2011). Mechanisms of rapid glucocorticoid feedback inhibition of the
648 hypothalamic--pituitary--adrenal axis. *Stress-the International Journal on the Biology of Stress*,
649 *14*(4), 398-406. doi:10.3109/10253890.2011.586446

650 Teicher, M. H., & Samson, J. A. (2016). Annual Research Review: Enduring neurobiological effects of
651 childhood abuse and neglect. *Journal of Child Psychology and Psychiatry*, *57*(3), 241-266.
652 doi:10.1111/jcpp.12507

653 Teicher, M. H., Samson, J. A., Anderson, C. M., & Ohashi, K. (2016). The effects of childhood
654 maltreatment on brain structure, function and connectivity. *Nature Reviews Neuroscience*,
655 *17*(10), 652. doi:10.1038/nrn.2016.111

656 Tiwari, A., & Gonzalez, A. (2018). Biological alterations affecting risk of adult psychopathology
657 following childhood trauma: A review of sex differences. *Clinical Psychology Review*, *66*, 69-
658 79. doi:10.1016/j.cpr.2018.01.006

659 Tobon, A. L., Newport, D. J., & Nemeroff, C. B. (2018). The role of oxytocin in early life adversity and
660 later psychopathology: a review of preclinical and clinical studies. In (Vol. 5, pp. 401-415).
661 Current Treatment Options in Psychiatry.

662 Touma, C., Gassen, N. C., Herrmann, L., Cheung-Flynn, J., Bull, D. R., Ionescu, I. A., . . . Rein, T. (2011).
663 FK506 Binding Protein 5 Shapes Stress Responsiveness: Modulation of Neuroendocrine
664 Reactivity and Coping Behavior. *Biological Psychiatry*, *70*(10), 928-936.
665 doi:10.1016/j.biopsych.2011.07.023

666 Turner, B. B. (1990). Sex difference in glucocorticoid binding in rat pituitary is estrogen dependent.
667 *Life Sciences*, 46(19), 1399-1406. doi:10.1016/0024-3205(90)90340-w

668 van Bodegom, M., Homberg, J. R., & Henckens, M. (2017). Modulation of the Hypothalamic-Pituitary-
669 Adrenal Axis by Early Life Stress Exposure. *Frontiers in Cellular Neuroscience*, 11, 33.
670 doi:10.3389/fncel.2017.00087

671 van der Doelen, R. H. A., Calabrese, F., Guidotti, G., Geenen, B., Riva, M. A., Kozicz, T., & Homberg, J.
672 R. (2014). Early life stress and serotonin transporter gene variation interact to affect the
673 transcription of the glucocorticoid and mineralocorticoid receptors, and the co-chaperone
674 FKBP5, in the adult rat brain. *Frontiers in Behavioral Neuroscience*, 8.
675 doi:10.3389/fnbeh.2014.00355

676 Veenema, A. H., Bredewold, R., & Neumann, I. D. (2007). Opposite effects of maternal separation on
677 intermale and maternal aggression in C57BL/6 mice: Link to hypothalamic vasopressin and
678 oxytocin immunoreactivity. *Psychoneuroendocrinology*, 32(5), 437-450.
679 doi:10.1016/j.psyneuen.2007.02.008

680 Veenema, A. H., & Neumann, I. D. (2009). Maternal separation enhances offensive play-fighting, basal
681 corticosterone and hypothalamic vasopressin mRNA expression in juvenile male rats.
682 *Psychoneuroendocrinology*, 34(3), 463-467. doi:10.1016/j.psyneuen.2008.10.017

683 Wang, Q. Z., Shelton, R. C., & Dwivedi, Y. (2018). Interaction between early-life stress and FKBP5 gene
684 variants in major depressive disorder and post-traumatic stress disorder: A systematic review
685 and meta-analysis. *Journal of Affective Disorders*, 225, 422-428.
686 doi:10.1016/j.jad.2017.08.066

687 Welberg, L. A. M., Seckl, J. R., & Holmes, M. C. (2001). Prenatal glucocorticoid programming of brain
688 corticosteroid receptors and corticotrophin-releasing hormone: Possible implications for
689 behaviour. *Neuroscience*, 104(1), 71-79. doi:10.1016/s0306-4522(01)00065-3

690 Wochnik, G. M., Ruegg, J., Abel, G. A., Schmidt, U., Holsboer, F., & Rein, T. (2005). FK506-binding
691 proteins 51 and 52 differentially regulate dynein interaction and nuclear translocation of the
692 glucocorticoid receptor in mammalian cells. *Journal of Biological Chemistry*, 280(6), 4609-
693 4616. doi:10.1074/jbc.M407498200

694 Wulsin, A. C., Wick-Carlson, D., Packard, B. A., Morano, R., & Herman, J. P. (2016). Adolescent chronic
695 stress causes hypothalamo-pituitary-adrenocortical hypo-responsiveness and depression-like
696 behavior in adult female rats. In (Vol. 65, pp. 109-117). *Psychoneuroendocrinology*.

697 Xu, J. J., Wang, R., Liu, Y., Liu, D. X., Jiang, H., & Pan, F. (2017). FKBP5 and specific microRNAs via
698 glucocorticoid receptor in the basolateral amygdala involved in the susceptibility to depressive
699 disorder in early adolescent stressed rats. *Journal of Psychiatric Research*, 95, 102-113.
700 doi:10.1016/j.jpsychires.2017.08.010

701 Xu, J. J., Wang, R., Liu, Y., Wang, W., Liu, D. X., Jiang, H., & Pan, F. (2019). Short- and long-term
702 alterations of FKBP5-GR and specific microRNAs in the prefrontal cortex and hippocampus of
703 male rats induced by adolescent stress contribute to depression susceptibility.
704 *Psychoneuroendocrinology*, 101, 204-215. doi:10.1016/j.psyneuen.2018.11.008

705 Zannas, A. S., Wiechmann, T., Gassen, N. C., & Binder, E. B. (2016). Gene-Stress-Epigenetic Regulation
706 of FKBP5: Clinical and Translational Implications. *Neuropsychopharmacology*, 41(1), 261-274.
707 doi:10.1038/npp.2015.235

708 Zhang, L., & Hernandez, V. S. (2013). Synaptic innervation to rat hippocampus by vasopressin-immuno-
709 positive fibres from the hypothalamic supraoptic and paraventricular nuclei. *Neuroscience*,
710 228, 139-162. doi:10.1016/j.neuroscience.2012.10.010

711 Zorn, J. V., Schur, R. R., Boks, M. P., Kahn, R. S., Joels, M., & Vinkers, C. H. (2017). Cortisol stress
712 reactivity across psychiatric disorders: A systematic review and meta-analysis.
713 *Psychoneuroendocrinology*, 77, 25-36. doi:10.1016/j.psyneuen.2016.11.036